

Subarachnoid haemorrhage

[O] Evidence review for imaging strategies for follow-up

NICE guideline <number>

Evidence review underpinning

February 2021

Draft for consultation

*Developed by the National Guideline Centre,
hosted by the Royal College of Physicians*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and, where appropriate, their carer or guardian.

Local commissioners and providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to Notice of rights.

ISBN

[add for final publication version only, delete this text for consultation version]

Contents

1	Imaging strategies for follow-up	5
1.1	Review question: What is the clinical and cost effectiveness of different imaging strategies for follow-up of adults with confirmed aneurysmal subarachnoid haemorrhage?	5
1.2	Introduction	5
1.3	PICO table	5
1.4	Clinical evidence	6
1.4.1	Included studies	6
1.4.2	Excluded studies	6
1.4.3	Summary of clinical studies included in the evidence review	7
1.4.4	Quality assessment of clinical studies included in the evidence review	7
1.5	Economic evidence	8
1.5.1	Included studies	8
1.5.2	Excluded studies	8
1.5.3	Unit costs	8
1.6	The committee's discussion of the evidence	8
1.6.1	Interpreting the evidence	8
1.6.2	Cost effectiveness and resource use	9
1.6.3	Other factors the committee took into account	10
	Appendices	21
	Appendix A: Review protocols	21
	Appendix B: Literature search strategies	28
	B.1 Clinical search literature search strategy	28
	B.2 Health Economics literature search strategy	33
	Appendix C: Clinical evidence selection	37
	Appendix D: Clinical evidence tables	38
	Appendix E: Forest plots	39
	Appendix F: GRADE tables	40
	Appendix G: Health economic evidence selection	41
	Appendix H: Health economic evidence tables	43
	Appendix I: Excluded studies	44
	I.1 Excluded clinical studies	44
	I.2 Excluded health economic studies	47

1 ¹ Imaging strategies for follow-up

² Evidence review underpinning recommendation 1.4.3 in the NICE guideline.

1.1 ³ Review question: What is the clinical and cost ⁴ effectiveness of different imaging strategies for follow-up ⁵ of adults with confirmed aneurysmal subarachnoid ⁶ haemorrhage?

1.2 ⁷ Introduction

⁸ Prior to the introduction of aneurysm coiling, long-term imaging follow-up for clipped brain
⁹ aneurysms was uncommon. Aneurysm clipping was known to achieve complete aneurysm
¹⁰ occlusion, as the aneurysm clip pinched the endothelial surfaces of the aneurysm wall
¹¹ together (the direct contact of biological surfaces resulting in healing across the aneurysm
¹² neck akin to scar formation).

¹³ In contemporary practice, imaging follow-up is not generally required after aneurysm clipping,
¹⁴ but catheter angiography is arranged if there is concern about adequacy of clip placement, if
¹⁵ there is a residual aneurysm after clipping, or if there is another known unruptured aneurysm
¹⁶ that may require future treatment.

¹⁷ Technology that enabled endovascular treatment of brain aneurysms at a population level
¹⁸ was developed in the early 1990s and evaluated primarily by the ISAT trial. Histological and
¹⁹ angiographic evaluation of the results of early coiling treatments showed that aneurysm neck
²⁰ healing after treatment with coils was frequently incomplete, as the extent that reconstruction
²¹ of the arterial wall occurs over a metal scaffold varies depending on a number of factors.
²² Knowledge that aneurysm treatment with coiling may not result in complete aneurysm
²³ occlusion (when compared with aneurysm clipping) resulted in the evolution of imaging
²⁴ strategies to evaluate the immediate and longer-term results of endovascular aneurysm
²⁵ treatment.

²⁶ In some cases where it is thought that follow-up imaging will not change patient
²⁷ management, either due to a low likelihood of further events in the patient's estimated life-
²⁸ time, or due to a very poor clinical outcome, patients are discharged from imaging
²⁹ surveillance.

³⁰ This review was carried out to assess the clinical and cost-effectiveness of different follow-up
³¹ imaging strategies for adults with confirmed aneurysmal SAH.

1.3 ³² PICO table

³³ For full details see the review protocol in Appendix A:.

³⁴ **Table 1: PICO characteristics of review question**

Population	Adults (16 and older) with a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm.
Interventions	Follow up imaging strategy at: <ul style="list-style-type: none">• <1 year<ul style="list-style-type: none">○ 6-monthly follow-up○ Yearly follow-up• 1-2 years<ul style="list-style-type: none">○ 6-monthly follow-up

	<ul style="list-style-type: none">○ Yearly follow-up○ Follow-up at >yearly intervals● >2-5 years<ul style="list-style-type: none">○ 6-monthly follow-up○ Yearly follow-up○ Follow-up at >yearly intervals
Comparisons	<ul style="list-style-type: none">● To each other● No imaging follow up
Outcomes	<p>Critical:</p> <ul style="list-style-type: none">● Mortality● Health and social-related quality of life (any validated score)● Degree of disability or dependence in daily activities, (e.g. Modified Rankin Scale and patient-reported outcome measures)● Complications of investigation (e.g. stroke, vascular injury) <p>Important:</p> <ul style="list-style-type: none">● Subsequent subarachnoid haemorrhage● Return to daily activity (e.g. work)● Need for retreatment● Length of hospital stay (if rehospitalised)
Study design	<ul style="list-style-type: none">● Randomised controlled trials (RCTs), systematic reviews of RCTs.● If insufficient RCT evidence is available, non-randomised studies will be considered if they adjust for key confounders (age), starting with prospective cohort studies.

1

1.4 2 Clinical evidence

1.4.1 3 Included studies

4 No relevant clinical studies comparing follow up imaging strategies were identified.

5 See also the study selection flow chart in Appendix C:.

1.4.2 6 Excluded studies

7 See the excluded studies list in Appendix I:.

8

1.4.3 1 Summary of clinical studies included in the evidence review

2 No evidence was identified for this review.

1.4.4 3 Quality assessment of clinical studies included in the evidence review

4 No evidence was identified for this review.

5

6

1.5 1 Economic evidence

1.5.1 2 Included studies

3 No health economic studies were included.

1.5.2 4 Excluded studies

5 No relevant health economic studies were excluded due to assessment of limited
6 applicability or methodological limitations.

7 See also the health economic study selection flow chart in Appendix G:

1.5.3 8 Unit costs

9 Relevant unit costs are provided below to aid consideration of cost effectiveness.

10 **Table 2: UK costs of imaging modalities for follow-up**

Drug	Description	Average cost
Computerised Tomography Angiography	Computerised Tomography Scan of One Area, with Post-Contrast Only, 19 years and over [NHS Reference Cost code: RD21A]	£101
Magnetic Resonance Angiography	Magnetic Resonance Imaging Scan of One Area, with Pre and Post-Contrast, 19 years and over [NHS Reference Cost code: RD03Z]	£190
Digital Subtraction Angiography	Percutaneous Transluminal Arteriography, of Intracranial or Extracranial Blood Vessel (day case) [NHS Reference Cost code: YA11Z]	£1,448

11 *Source: NHS Reference Cost 2018/19⁸³*

1.6 12 The committee's discussion of the evidence

1.6.1 13 Interpreting the evidence

1.6.1.1 14 The outcomes that matter most

15 The committee agreed that the intention of follow-up imaging is to monitor any non-culprit
16 aneurysms, assess the status of previous interventions, and to help detect and manage any
17 treatment complications. The committee considered critical outcomes for decision making to
18 be mortality, health and social related quality of life, degree of disability or dependence in
19 daily activities (e.g. Modified Rankin scale and patient reported outcome measures), and
20 complications of investigation. The committee also considered subsequent subarachnoid
21 haemorrhage, return to daily activity, complications of intervention, need for re-treatment,
22 and length of hospital stay (if rehospitalised) to be important outcomes.

1.6.1.2 23 The quality of the evidence

24 No evidence was identified. As such, the committee used consensus to provide clinical
25 recommendation to consider neuroimaging as part of a follow up strategy for some people
26 depending on individual clinical need and risk factors.

27 Given the lack of evidence the committee discussed making a research recommendation on
28 the optimal frequency, duration and indications for follow-up. This is a difficult area to study
29 as decisions to carry out neuroimaging during follow-up of a patient are based on multiple

1 factors, such as perceived effectiveness of the initial treatment, estimated risk of aneurysm
2 recurrence, actual aneurysm recurrence, patient age, the presence and estimated risk of
3 associated aneurysms and comorbidities.

4 The committee agreed that an important first point is estimation of risk of subarachnoid
5 haemorrhage and the research recommendation that they made on the development and
6 evaluation of risk stratification tools to estimate the risk of subsequent aneurysmal
7 subarachnoid haemorrhage within the evidence review in this guideline on the risk of
8 subsequent SAH would inform indications for follow-up imaging (see evidence review N,
9 Appendix H). They considered therefore that a separate research recommendation around
10 follow up imaging was not appropriate at this time.

1.6.1.31 Benefits and harms

12 The committee agreed that there are benefits to follow-up imaging: assessment of the
13 operative result and detection of aneurysm recurrence, identification of complications of SAH
14 or treatment, and evaluation and surveillance of non-culprit, or de-novo aneurysms. Some of
15 these are necessary for early patient management and the benefits will generally outweigh
16 any harms. Following endovascular treatment, imaging follow-up is usually required to
17 confirm that operative treatment has been effective and that on-going risks from the treated
18 aneurysm are negligible and or remain low for the duration of follow-up. The main long-term
19 harms from imaging relate to effects of ionising radiation as well as use of resources. The
20 committee noted however that patients with treated aSAH are at risk of both aneurysm
21 recurrence and formation of new aneurysm(s). The committee considered the evidence
22 presented in evidence review N on the risk of subsequent SAH showing that the risk of
23 rebleed or subsequent aSAH in people who receive endovascular or neurosurgical
24 intervention is generally around 0.5% per annum, although this incidence rate varies
25 between individual patients. These data also showed that the overall risk of subsequent SAH
26 beyond 1-year post-ictus is closer to 0.1% per annum. People who have had aSAH can also
27 benefit from reassurance of follow up imaging to help manage any anxiety and management
28 of continuing symptoms such as headache.

29 The lack of evidence on both the clinical and cost effectiveness of follow-up strategies
30 prevented the committee from making a strong recommendation. The committee therefore
31 decided to make a consensus recommendation to consider neuroimaging as part of a follow
32 up strategy for some people depending on individual clinical need and risk factors.

33 Individual clinical factors, and type and outcome of any neurointervention or neurosurgery
34 may influence this risk and should inform any follow-up strategy.

35 The committee agreed that follow-up imaging can be done by MRA, which is the preferred
36 modality as it avoids exposure to ionising radiation. CT angiography can also be used but
37 involves ionising radiation and CT images may be degraded by artefact from aneurysm metal
38 coils. Catheter angiography (DSA) also requires ionising radiation, is an invasive procedure
39 associated with a small risk, and associated with additional cost compared with non-invasive
40 alternatives (MRA, CTA). DSA is reserved for patients in whom MRA and CTA are
41 contraindicated or not tolerated. In current practice the majority of patients are followed up
42 with non-invasive MRA.

43 The committee therefore recommended that the choice of imaging follow-up technique will be
44 at clinician discretion but should take account of the treatment method, presence of non-
45 culprit aneurysm(s), perceived risk of further bleeding, risks of planned investigations and
46 any subsequent interventions and patient preference.

1.6.27 Cost effectiveness and resource use

48 No published economic evaluations were identified or included in this review. Unit costs of
49 different imaging modalities were presented to the committee, but the lack of clinical

- 1 evidence precluded assessment of the cost effectiveness of different follow up strategies.
- 2 The committee made a consensus recommendation, which will not change current clinical
- 3 practice or result in a resource impact to the NHS.

1.6.3 4 Other factors the committee took into account

- 5 The committee noted that current standard practice is to offer a follow-up appointment to
- 6 patients at 6 months and at 2 years. There is no nationally agreed standard surveillance
- 7 interval, and strategies will be individualised. Imaging strategies should take into account the
- 8 accuracy and any associated risk of the test, as well as acceptability to the patient. The
- 9 approach to imaging follow-up should weigh the impact of uncertainty on a patient's well-
- 10 being and be responsive to patient wishes.
- 11

1 References

- 2 1. Aboukais R, Zairi F, Bourgeois P, Boustia F, Leclerc X, Lejeune JP. Pericallosal
3 aneurysm: a difficult challenge for microsurgery and endovascular treatment. *Neuro-
4 Chirurgie*. 2015; 61(4):244-249
- 5 2. Adeeb N, Griessenauer CJ, Foreman PM, Moore JM, Motiei-Langroudi R, Chua MH
6 et al. Comparison of stent-assisted coil embolization and the pipeline embolization
7 device for endovascular treatment of ophthalmic segment aneurysms: a multicenter
8 cohort study. *World Neurosurgery*. 2017; 105:206-212
- 9 3. Ahmed SU, Mocco J, Zhang X, Kelly M, Doshi A, Nael K et al. MRA versus DSA for
10 the follow-up imaging of intracranial aneurysms treated using endovascular
11 techniques: a meta-analysis. *Journal of Neurointerventional Surgery*. 2019;
12 11(10):1009-1014
- 13 4. Anzalone N, De Filippis C, Scomazzoni F, Calori G, Iadanza A, Simionato F et al.
14 Longitudinal follow up of coiled intracranial aneurysms: the impact of contrast
15 enhanced MRA in comparison to 3DTOF MRA at 3T. *Neurovascular Imaging*. 2015;
16 1:11
- 17 5. Arrese I, Sarabia R, Pintado R, Delgado-Rodriguez M. Flow-diverter devices for
18 intracranial aneurysms: systematic review and meta-analysis. *Neurosurgery*. 2013;
19 73(2):193-199
- 20 6. Arthur AS, Molyneux A, Coon AL, Saatci I, Szikora I, Baltacioglu F et al. The safety
21 and effectiveness of the Woven EndoBridge (WEB) system for the treatment of wide-
22 necked bifurcation aneurysms: final 12-month results of the pivotal WEB Intracranial
23 Therapy (WEB-IT) study. *Journal of NeuroInterventional Surgery*. 2019; 11(9):924-
24 930
- 25 7. Atasoy D, Kandasamy N, Hart J, Lynch J, Yang SH, Walsh D et al. Outcome study of
26 the pipeline embolization device with shield technology in unruptured aneurysms
27 (PEDSU). *American Journal of Neuroradiology*. 2019; 40(12):2094-2101
- 28 8. Aydin K, Stracke CP, Barbuoglu M, Yamac E, Berdikhojayev M, Sencer S et al.
29 Long-term outcomes of wide-necked intracranial bifurcation aneurysms treated with
30 T-stent-assisted coiling. *Journal of Neurosurgery*. 2019;
31 <https://dx.doi.org/10.3171/2019.9.JNS191733>
- 32 9. Aydin Y, Cavusoglu H, Kahyaoglu O, Musluman AM, Yilmaz A, Turkmenoglu ON et
33 al. Clip ligation of unruptured intracranial aneurysms: a prospective midterm outcome
34 study. *Acta Neurochirurgica*. 2012; 154(7):1135-1144
- 35 10. Bakker NA, Groen RJ, Foumani M, Uyttenboogaart M, Eshghi OS, Metzemaekers JD
36 et al. Repeat digital subtraction angiography after a negative baseline assessment in
37 nonperimesencephalic subarachnoid hemorrhage: a pooled data meta-analysis.
38 *Journal of Neurosurgery*. 2014; 120(1):99-103
- 39 11. Bendok BR, Rahme RJ. Complex shaped detachable platinum coil system for the
40 treatment of cerebral aneurysms: the Codman TruFill DCS and TruFill DCS Orbit
41 Detachable Coil System COMPLEX Registry final results. *Journal of
42 Neurointerventional Surgery*. 2013; 5(1):54-61
- 43 12. Bor AS, Rinkel GJ, van Norden J, Wermer MJ. Long-term, serial screening for
44 intracranial aneurysms in individuals with a family history of aneurysmal subarachnoid
45 haemorrhage: a cohort study. *Lancet Neurology*. 2014; 13(4):385-392

- 1 13. Bousset L, Rayz V, McCulloch C, Martin A, Acevedo-Bolton G, Lawton M et al.
2 Aneurysm growth occurs at region of low wall shear stress: patient-specific
3 correlation of hemodynamics and growth in a longitudinal study. *Stroke*. 2008;
4 39(11):2997-3002
- 5 14. Bracard S, Lebedinsky A, Anxionnat R, Neto JM, Audibert G, Long Y et al.
6 Endovascular treatment of Hunt and Hess grade IV and V aneurysms. *American*
7 *Journal of Neuroradiology*. 2002; 23(6):953-957
- 8 15. Bruneau M, Rynkowski M, Smida-Rynkowska K, Brotchi J, De Witte O, Lubicz B.
9 Long-term follow-up survey reveals a high yield, up to 30% of patients presenting
10 newly detected aneurysms more than 10 years after ruptured intracranial aneurysms
11 clipping. *Neurosurgical Review*. 2011; 34(4):485-496
- 12 16. CBS Office of Statistics Netherlands. National life tables. 2000. Available from:
13 <https://www.cbs.nl/> Last accessed: 17/10/2019.
- 14 17. Chalouhi N, Bovenzi CD, Thakkar V, Dressler J, Jabbour P, Starke RM et al. Long-
15 term catheter angiography after aneurysm coil therapy: results of 209 patients and
16 predictors of delayed recurrence and retreatment. *Journal of Neurosurgery*. 2014;
17 121(5):1102-1106
- 18 18. Chen X, Sun Z, Shi L, Xu L, Yu J, Fang B et al. Endovascular management of
19 ruptured distal posterior inferior cerebellar artery aneurysms A retrospective cohort
20 study. *Medicine*. 2018; 97(49):e13300
- 21 19. Cho YD, Jeon JP, Rhim JK, Park JJ, Yoo RE, Kang HS et al. Progressive thrombosis
22 of small saccular aneurysms filled with contrast immediately after coil embolization:
23 analysis of related factors and long-term follow-up. *Neuroradiology*. 2015; 57(6):615-
24 623
- 25 20. Cho YD, Kim KM, Lee WJ, Sohn CH, Kang HS, Kim JE et al. Time-of-flight magnetic
26 resonance angiography for follow-up of coil embolization with enterprise stent for
27 intracranial aneurysm: usefulness of source images. *Korean Journal of Radiology*.
28 2014; 15(1):161-168
- 29 21. Cloft HJ, Joseph GJ, Dion JE. Risk of cerebral angiography in patients with
30 subarachnoid hemorrhage, cerebral aneurysm, and arteriovenous malformation: a
31 meta-analysis. *Stroke*. 1999; 30(2):317-320
- 32 22. Cognard C, Januel AC. Remnants and recurrences after the use of the WEB
33 intrasaccular device in large-neck bifurcation aneurysms. *Neurosurgery*. 2015;
34 76(5):522-530;discussion 530
- 35 23. Crawley F, Clifton A, Brown MM. Should we screen for familial intracranial aneurysm?
36 *Stroke*. 1999; 30(2):312-316
- 37 24. David CA, Vishteh AG, Spetzler RF, Lemole M, Lawton MT, Partovi S. Late
38 angiographic follow-up review of surgically treated aneurysms. *Journal of*
39 *Neurosurgery*. 1999; 91(3):396-401
- 40 25. De Letter JA, Moll FL, Welten RJ, Eikelboom BC, Ackerstaff RG, Vermeulen FE et al.
41 Benefits of carotid patching: a prospective randomized study with long-term follow-up.
42 *Annals of Vascular Surgery*. 1994; 8(1):54-58
- 43 26. Delgado Almandoz JE, Jagadeesan BD, Refai D, Moran CJ, Cross DT, 3rd, Chicoine
44 MR et al. Diagnostic yield of repeat catheter angiography in patients with catheter and
45 computed tomography angiography negative subarachnoid hemorrhage.
46 *Neurosurgery*. 2012; 70(5):1135-1142

- 1 27. Disney L, Weir B, Grace M. Factors influencing the outcome of aneurysm rupture in
2 poor grade patients: a prospective series. *Neurosurgery*. 1988; 23(1):1-9
- 3 28. Dorhout Mees SM, Van Den Bergh WM, Algra A, Rinkel GJE. Antiplatelet therapy for
4 aneurysmal subarachnoid haemorrhage. *Cochrane Database of Systematic Reviews*
5 2007, Issue 4. Art. No.: CD006184. DOI:
6 <http://dx.doi.org/10.1002/14651858.CD006184.pub2>.
- 7 29. Flores BC, White JA, Hunt Batjer H, Samson DS. The 25th anniversary of the
8 retrograde suction decompression technique (Dallas technique) for the surgical
9 management of paraclinoid aneurysms: historical background, systematic review, and
10 pooled analysis of the literature. *Journal of Neurosurgery*. 2019; 130(3):902-916
- 11 30. Fountas KN, Kassam M, MacHinis TG, Dimopoulos VG, Robinson IJS, Ajjan M et al.
12 C-reactive protein might predict outcome in aneurysmal subarachnoid haemorrhage.
13 *Cerebral Vasospasm: New Strategies in Research and Treatment*. 2008; *Acta*
14 *Neurochirurgica, Supplementum*.(104):377-381
- 15 31. Fujimura M, Joo JY, Kim JS, Hatta M, Yokoyama Y, Tominaga T. Preventive effect of
16 clazosentan against cerebral vasospasm after clipping surgery for aneurysmal
17 subarachnoid hemorrhage in Japanese and Korean patients. *Cerebrovascular*
18 *Diseases*. 2017; 44(1-2):59-67
- 19 32. Gallas S, Januel AC, Pasco A, Drouineau J, Gabrillargues J, Gaston A et al. Long-
20 term follow-up of 1036 cerebral aneurysms treated by bare coils: a multicentric cohort
21 treated between 1998 and 2003. *American Journal of Neuroradiology*. 2009;
22 30(10):1986-1992
- 23 33. Gallas S, Pasco A, Cottier JP, Gabrillargues J, Drouineau J, Cognard C et al. A
24 multicenter study of 705 ruptured intracranial aneurysms treated with Guglielmi
25 detachable coils. *American Journal of Neuroradiology*. 2005; 26(7):1723-1731
- 26 34. Garg K, Sinha S, Kale SS, Chandra PS, Suri A, Singh MM et al. Role of simvastatin
27 in prevention of vasospasm and improving functional outcome after aneurysmal sub-
28 arachnoid hemorrhage: a prospective, randomized, double-blind, placebo-controlled
29 pilot trial. *British Journal of Neurosurgery*. 2013; 27(2):181-186
- 30 35. Gauvrit JY, Leclerc X, Pernodet M, Lubicz B, Lejeune JP, Leys D et al. Intracranial
31 aneurysms treated with Guglielmi detachable coils: usefulness of 6-month imaging
32 follow-up with contrast-enhanced MR angiography. *American Journal of*
33 *Neuroradiology*. 2005; 26(3):515-521
- 34 36. Geng B, Wu X, Brackett A, Malhotra A. Meta-analysis of recent literature on utility of
35 follow-up imaging in isolated perimesencephalic hemorrhage. *Clinical Neurology and*
36 *Neurosurgery*. 2019; 180:111-116
- 37 37. Germano A, Priola S, Angileri FF, Conti A, La Torre D, Cardali S et al. Long-term
38 follow-up of ruptured intracranial aneurysms treated by microsurgical wrapping with
39 autologous muscle. *Neurosurgical Review*. 2013; 36(1):123-131; discussion 132
- 40 38. Geyik S, Yavuz K, Yurttutan N, Saatci I, Cekirge HS. Stent-assisted coiling in
41 endovascular treatment of 500 consecutive cerebral aneurysms with long-term follow-
42 up. *American Journal of Neuroradiology*. 2013; 34(11):2157-2162
- 43 39. Ghogawala Z, Amin-Hanjani S, Curran J, Ciarleglio M, Berenstein A, Stabile L et al.
44 The effect of carotid endarterectomy on cerebral blood flow and cognitive function.
45 *Journal of Stroke and Cerebrovascular Diseases*. 2013; 22(7):1029-1037

- 1 40. Gibbs GF, Huston J, 3rd, Qian Q, Kubly V, Harris PC, Brown RD, Jr. et al. Follow-up
2 of intracranial aneurysms in autosomal-dominant polycystic kidney disease. *Kidney*
3 *International*. 2004; 65(5):1621-1627
- 4 41. Goksu E, Korkmaz E, Akyuz M, Ozgur O, Sindel T, Tuncer R. The analysis of long-
5 term follow-up screening in patients with surgically treated intracranial aneurysms.
6 *Turkish Neurosurgery*. 2015; 25(3):404-409
- 7 42. Groden C, Eckert B, Ries T, Probst EN, Kucinski T, Zeumer H. Angiographic follow-
8 up of vertebrobasilar artery aneurysms treated with detachable coils. *Neuroradiology*.
9 2003; 45(7):435-440
- 10 43. Hai J, Deng DF, Chen ZQ, Pan QG. Endovascular embolization of small ruptured
11 intracranial aneurysms using a biplane angiographic system with three-dimensional
12 rotational digital subtraction angiography. *Journal of Clinical Neuroscience*. 2009;
13 16(8):1028-1033
- 14 44. Hashimoto H, Iida JI, Hironaka Y, Okada M, Sakaki T. Use of spiral computerized
15 tomography angiography in patients with subarachnoid hemorrhage in whom
16 subtraction angiography did not reveal cerebral aneurysms. *Journal of Neurosurgery*.
17 2000; 92(2):278-283
- 18 45. Hassan AE, Zacharatos H, Rodriguez GJ, Suri MF, Tariq N, Vazquez G et al. Long-
19 term clinical and angiographic outcomes in patients with spontaneous cervico-cranial
20 arterial dissections treated with stent placement. *Journal of Neuroimaging*. 2012;
21 22(4):384-393
- 22 46. Heller R, Calnan DR, Lanfranchi M, Madan N, Malek AM. Incomplete stent apposition
23 in Enterprise stent-mediated coiling of aneurysms: persistence over time and risk of
24 delayed ischemic events. *Journal of Neurosurgery*. 2013; 118(5):1014-1022
- 25 47. Helthuis JHG, Bhat S, Van Doormaal TPC, Kumar RK, Van Der Zwan A. Proximal
26 and distal occlusion of complex cerebral aneurysms-implications of flow modeling by
27 fluid-structure interaction analysis. *Operative Neurosurgery*. 2018; 15(2):217-230
- 28 48. Hendryk S, Jarzab B, Josko J. Increase of the IL-1 beta and IL-6 levels in CSF in
29 patients with vasospasm following aneurysmal SAH. *Neuroendocrinology Letters*.
30 2004; 25(1-2):141-147
- 31 49. Hijdra A, Braakman R, van Gijn J, Vermeulen M, van Crevel H. Aneurysmal
32 subarachnoid hemorrhage. Complications and outcome in a hospital population.
33 *Stroke*. 1987; 18(6):1061-1067
- 34 50. Hop JW, Rinkel GJ, Algra A, van Gijn J. Case-fatality rates and functional outcome
35 after subarachnoid hemorrhage: a systematic review. *Stroke*. 1997; 28(3):660-664
- 36 51. Hosono M, Machida K, Matsui T, Honda N, Takahashi T, Dei S et al. Non-invasive
37 quantitative monitoring of cerebral blood flow in aneurysmal subarachnoid
38 haemorrhage with 99mTc-ECD. *Nuclear Medicine Communications*. 2002; 23(1):5-11
- 39 52. Hussain SI, Lynch JR, Wolfe T, Fitzsimmons BF, Zaidat OO. Stent-assisted parent
40 artery occlusion of giant cerebrovascular aneurysms to avoid mass effect. *Journal of*
41 *Neuroimaging*. 2009; 19(4):370-374
- 42 53. Ji W, Kang H, Liu A, Li Y, Feng X, Qian Z et al. Stent-assisted coiling of very small
43 wide-necked intracranial aneurysms: complications, anatomical results and clinical
44 outcomes. *Neurologia i Neurochirurgia Polska*. 2016; 50(6):410-417

- 1 54. Juvela S, Porras M, Heiskanen O. Natural history of unruptured intracranial
2 aneurysms: a long-term follow-up study. *Journal of Neurosurgery*. 1993; 79(2):174-
3 182
- 4 55. Juvela S, Poussa K, Porras M. Factors affecting formation and growth of intracranial
5 aneurysms: a long-term follow-up study. *Stroke*. 2001; 32(2):485-491
- 6 56. Kalra VB, Wu X, Matouk CC, Malhotra A. Use of follow-up imaging in isolated
7 perimesencephalic subarachnoid hemorrhage: a meta-analysis. *Stroke*. 2015;
8 46(2):401-406
- 9 57. Kannath SK, Mohimen A, Raman KT, Abraham M, Nair S, Rajan JE. Single centre
10 experience of flow diverter treatment of complex intracranial aneurysms from South
11 India: intermediate and long-term outcomes. *Neurology India*. 2019; 67(3):797-802
- 12 58. Kao HL, Hung CS, Li HY, Yeh CF, Huang CC, Chen YH et al. Long-term outcomes
13 after endovascular recanalization in patients with chronic carotid artery occlusion.
14 *American Journal of Cardiology*. 2018; 122(10):1779-1783
- 15 59. Kapapa T, Woischneck D, Tjahjadi M. Long-term health-related quality of life after
16 spontaneous nontraumatic subarachnoid hemorrhage: self and proxy reports in a 10-
17 year period. *World Neurosurgery*. 2014; 81(1):105-109
- 18 60. Kasner SE, Hankins LL, Bratina P, Morgenstern LB. Magnetic resonance
19 angiography demonstrates vascular healing of carotid and vertebral artery
20 dissections. *Stroke*. 1997; 28(10):1993-1997
- 21 61. Kim CH, Kim YH, Sung SK, Son DW, Song GS, Lee SW. Clinical safety and
22 effectiveness of stent-assisted coil embolization with neuroform atlas stent in
23 intracranial aneurysm. *Journal of Korean Neurosurgical Society*. 2019; 63(1):80-88
- 24 62. Kim P, Jang SJ. Management of recurrent cerebral aneurysm after surgical clipping:
25 clinical article. *Journal of Korean Neurosurgical Society*. 2018; 61(2):212-218
- 26 63. Kim ST, Baek JW, Lee WH, Lee KS, Kwon WH, Pyo S et al. Causes of early
27 rebleeding after coil embolization of ruptured cerebral aneurysms. *Clinical Neurology
28 and Neurosurgery*. 2018; 174:108-116
- 29 64. King JT, Jr., Berlin JA, Flamm ES. Morbidity and mortality from elective surgery for
30 asymptomatic, unruptured, intracranial aneurysms: a meta-analysis. *Journal of
31 Neurosurgery*. 1994; 81(6):837-842
- 32 65. Kwee TC, Kwee RM. MR angiography in the follow-up of intracranial aneurysms
33 treated with Guglielmi detachable coils: systematic review and meta-analysis.
34 *Neuroradiology*. 2007; 49(9):703-713
- 35 66. Lindgren A, Vergouwen M, van der Schaaf I, Algra A, Wermer M, Clarke M et al.
36 Endovascular coiling versus neurosurgical clipping for people with aneurysmal
37 subarachnoid haemorrhage. *Cochrane Database of Systematic Reviews* 2018, Issue
38 8. Art. No.: CD003085. DOI: 10.1002/14651858.CD003085.pub3.
- 39 67. Lindvall P, Borota L, Birgander R, Jonasson P, Ridderheim PA. Long-term follow-up
40 of intracranial aneurysms treated with endovascular coiling: experience from one
41 institution. *Vascular and Endovascular Surgery*. 2012; 46(4):325-328
- 42 68. Liu JP, Ye ZN, Lv SY, Zhuang Z, Zhang XS, Zhang X et al. The rise of soluble
43 platelet-derived growth factor receptor beta in CSF early after subarachnoid
44 hemorrhage correlates with cerebral vasospasm. *Neurological Sciences*. 2018;
45 39(6):1105-1111

- 1 69. Lizza BD, Kosteva A, Maas MB, Rosenberg NF, Liotta E, Guth J et al. Preadmission
2 statin use does not improve functional outcomes or prevent delayed ischemic events
3 in patients with spontaneous subarachnoid hemorrhage. *Pharmacotherapy*. 2014;
4 34(8):811-817
- 5 70. Lublinsky S, Major S, Kola V, Horst V, Santos E, Platz J et al. Early blood-brain
6 barrier dysfunction predicts neurological outcome following aneurysmal subarachnoid
7 hemorrhage. *EBioMedicine*. 2019; 43:460-472
- 8 71. Ma C, Zhou W, Yan Z, Qu M, Bu X. Toll-like receptor 4 (TLR4) is associated with
9 cerebral vasospasm and delayed cerebral ischemia in aneurysmal subarachnoid
10 hemorrhage. *Neurologia Medico-Chirurgica*. 2015; 55(12):878-884
- 11 72. Maimaitili A, Maimaitili M, Rexidan A, Lu J, Ajimu K, Cheng X et al. Pituitary hormone
12 level changes and hyponatremia in aneurysmal subarachnoid hemorrhage.
13 *Experimental and Therapeutic Medicine*. 2013; 5:1657-1662
- 14 73. Marquardt G, Niebauer T, Schick U, Lorenz R. Long term follow up after
15 perimesencephalic subarachnoid haemorrhage. *Journal of Neurology, Neurosurgery
16 and Psychiatry*. 2000; 69(1):127-130
- 17 74. Mooney MA, Brigeman S, Bohl MA, Simon ED, Sheehy JP, Chang SW et al. Analysis
18 of overlapping surgery in patients undergoing microsurgical aneurysm clipping: acute
19 and long-term outcomes from the Barrow Ruptured Aneurysm Trial. *Journal of
20 Neurosurgery*. 2018; 129(3):711-717
- 21 75. Morais R, Mine B, Bruyere PJ, Naeije G, Lubicz B. Endovascular treatment of
22 intracranial aneurysms with the p64 flow diverter stent: mid-term results in 35 patients
23 with 41 intracranial aneurysms. *Neuroradiology*. 2017; 59(3):263-269
- 24 76. Moritz JL, Vendrell JF, Hoa D, Menjot N, Costalat V, Brunel H et al. Mid-term clinical
25 and angiographic results of cerebral aneurysms treated with matrix2() coils. *Journal of
26 Neuroradiology*. 2012; 39(5):326-331
- 27 77. Mortimer AM, Marsh H, Klimczak K, Joshi D, Barton H, Nelson RJ et al. Is long-term
28 follow-up of adequately coil-occluded ruptured cerebral aneurysms always
29 necessary? A single-center study of recurrences after endovascular treatment.
30 *Journal of Neurointerventional Surgery*. 2015; 7(5):373-379
- 31 78. Murakami T, Nishida T, Asai K, Kadono Y, Nakamura H, Fujinaka T et al. Long-term
32 results and follow-up examinations after endovascular embolization for unruptured
33 cerebral aneurysms. *American Journal of Neuroradiology*. 2019; 40(7):1191-1196
- 34 79. Murias Quintana E, Gil Garcia A, Vega Valdes P, Cuellar H, Meilan Martinez A, Saiz
35 Ayala A et al. Anatomical results, rebleeding and factors that affect the degree of
36 occlusion in ruptured cerebral aneurysms after endovascular therapy. *Journal of
37 Neurointerventional Surgery*. 2015; 7(12):892-897
- 38 80. Nagano A, Yamada Y, Miyake H, Domen K, Koyama T. Increased resting energy
39 expenditure after endovascular coiling for subarachnoid hemorrhage. *Journal of
40 Stroke and Cerebrovascular Diseases*. 2016; 25(4):813-818
- 41 81. Nakagawa I, Yokoyama S, Omoto K, Takeshima Y, Matsuda R, Nishimura F et al.
42 Omega-3 fatty acids ethyl esters suppress cerebral vasospasm and improve clinical
43 outcome following aneurysmal subarachnoid hemorrhage. *World Neurosurgery*.
44 2017; 99:457-464
- 45 82. National Institute for Health and Care Excellence. Developing NICE guidelines: the
46 manual [updated October 2018]. London. National Institute for Health and Care

- 1 Excellence, 2014. Available from:
2 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20overview>
- 3 83. NHS England and NHS Improvement. National cost collection for the NHS 2018-19.
4 2019. Available from: <https://improvement.nhs.uk/resources/national-cost-collection/>
5 Last accessed: 01/04/2020.
- 6 84. Niiro M, Shimosuru T, Nakamura K, Kadota K, Kuratsu JI. Long-term follow-up study
7 of patients with cavernous sinus aneurysm treated by proximal occlusion. *Neurologia*
8 *Medico-Chirurgica*. 2000; 40(2):88-97
- 9 85. Nossek E, Chalif DJ, Chakraborty S, Lombardo K, Black KS, Setton A. Concurrent
10 use of the pipeline embolization device and coils for intracranial aneurysms:
11 technique, safety, and efficacy. *Journal of Neurosurgery*. 2015; 122(4):904-911
- 12 86. Ocal O, Peker A, Balci S, Arat A. Placement of a stent within a flow diverter improves
13 aneurysm occlusion rates. *American Journal of Neuroradiology*. 2019; 40(11):1932-
14 1938
- 15 87. Okada T, Ishikawa T, Moroi J, Suzuki A. Timing of retreatment for patients with
16 previously coiled or clipped intracranial aneurysms: analysis of 156 patients with
17 multiple treatments. *Surgical Neurology International*. 2016; 7(Suppl 2):S40-48
- 18 88. Peschillo S, Caporlingua A, Resta MC, Paul Peluso JP, Burdi N, Sourour N et al.
19 Endovascular treatment of large and giant carotid aneurysms with flow-diverter stents
20 alone or in combination with coils: a multicenter experience and long-term follow-up.
21 *Operative Neurosurgery*. 2017; 13(4):492-502
- 22 89. Petridis AK, Cornelius JF, Kamp MA, Falahati S, Fischer I, Steiger HJ. Level of
23 headaches after surgical aneurysm clipping decreases significantly faster compared
24 to endovascular coiled patients. *Clinics and Practice*. 2017; 7(2):56-59
- 25 90. Piano M, Valvassori L, Lozupone E, Pero G, Quilici L, Boccardi E. FRED Italian
26 Registry: a multicenter experience with the flow re-direction endoluminal device for
27 intracranial aneurysms. *Journal of Neurosurgery*. 2020; 133(1):174-181
- 28 91. Pierot L, Spelle L, Molyneux A, Byrne J. Clinical and anatomical follow-up in patients
29 with aneurysms treated with the WEB device: 1-year follow-up report in the
30 cumulated population of 2 prospective, multicenter series (WEBCAST and French
31 Observatory). *Neurosurgery*. 2015; 78(1):133-139
- 32 92. Piske RL, Kanashiro LH, Paschoal E, Agner C, Lima SS, Aguiar PH. Evaluation of
33 Onyx HD-500 embolic system in the treatment of 84 wide-neck intracranial
34 aneurysms. *Neurosurgery*. 2009; 64(5):E865-875; discussion E875
- 35 93. Poon TK, Ho WS, Pang KY, Wong CK. Comparison of computerized tomography
36 angiography and digital subtraction angiography in ruptured cerebral aneurysm
37 surgery. *Surgical Practice*. 2006; 10(1):8-13
- 38 94. Potter CA, Fink KR, Ginn AL, Haynor DR. Perimesencephalic hemorrhage: yield of
39 single versus multiple dsa examinations-a single-center study and meta-analysis.
40 *Radiology*. 2016; 281(3):858-864
- 41 95. Proust F, Gerardin E, Derrey S, Lesveque S, Ramos S, Langlois O et al.
42 Interdisciplinary treatment of ruptured cerebral aneurysms in elderly patients. *Journal*
43 *of Neurosurgery*. 2010; 112(6):1200-1207
- 44 96. Qin F, Li Z, Fang X, Zhao X, Liu J, Wu D et al. Therapeutic effect of enterprise stent-
45 assisted embolization for very small ruptured intracranial aneurysms. *Medicine*. 2017;
46 96(34):e7832

- 1 97. Raffi L, Simonetti L, Cenni P, Agati R, Bacci A, Leonardi M. Follow up study of
2 embolized brain aneurysms. *Rivista di Neuroradiologia*. 2003; 16(4):595-608
- 3 98. Raschi M, Mut F, Byrne G, Putman CM, Tateshima S, Vinuela F et al. CFD and PIV
4 analysis of hemodynamics in a growing intracranial aneurysm. *International Journal
5 for Numerical Methods in Biomedical Engineering*. 2012; 28(2):214-228
- 6 99. Rautio R, Rahi M, Katila A, Rinne J. Single-center experience with six-month follow-
7 up of FRED Jr flow diverters for intracranial aneurysms in small arteries. *Acta
8 Radiologica*. 2019; 60(7):917-924
- 9 100. Ruppert V, Erz K, Burklein D, Traitl M, Steckmeier B, Stelzer W et al. Double tube
10 stent-grafts for infrarenal aortic aneurysm: a new concept. *Journal of Endovascular
11 Therapy*. 2007; 14(2):144-149
- 12 101. Saatci I, Cekirge HS, Ciceri EF, Mawad ME, Pamuk AG, Besim A. CT and MR
13 imaging findings and their implications in the follow-up of patients with intracranial
14 aneurysms treated with endosaccular occlusion with onyx. *American Journal of
15 Neuroradiology*. 2003; 24(4):567-578
- 16 102. Serafin Z, Strzesniewski P, Lasek W, Beuth W. Methods and time schedule for follow-
17 up of intracranial aneurysms treated with endovascular embolization: a systematic
18 review. *Neurologia i Neurochirurgia Polska*. 2011; 45(5):421-430
- 19 103. Slater LA, Soufan C, Holt M, Chong W. Effect of flow diversion with silk on aneurysm
20 size: a single center experience. *Interventional Neuroradiology*. 2015; 21(1):12-18
- 21 104. Slosberg PS. Unexpected results in long-term medically treated ruptured intracranial
22 aneurysm including data on 14 patients followed more than 30 years each. *Acta
23 Neurochirurgica*. 1997; 139(8):697-705
- 24 105. Sprengers ME, van Rooij WJ, Sluzewski M, Rinkel GJ, Velthuis BK, de Kort GA et al.
25 MR angiography follow-up 5 years after coiling: frequency of new aneurysms and
26 enlargement of untreated aneurysms. *American Journal of Neuroradiology*. 2009;
27 30(2):303-307
- 28 106. Sun H, Ma J, Liu Y, Lan Z, You C. Diagnosing residual or recurrent cerebral
29 aneurysms after clipping by computed tomographic angiography: meta-analysis.
30 *Neurology India*. 2013; 61(1):51-55
- 31 107. Tailor J, Goetz P, Chandrashekar H, Stephen T, Schiariti M, Grieve J et al. Stability of
32 ruptured intracranial aneurysms treated with detachable coils: is delayed follow-up
33 angiography warranted? *British Journal of Neurosurgery*. 2010; 24(4):405-409
- 34 108. Tao C, Fan C, Hu X, Ma J, Ma L, Li H et al. The effect of fenestration of the lamina
35 terminalis on the incidence of shunt-dependent hydrocephalus after aneurysmal
36 subarachnoid hemorrhage (FISH) Study protocol for a randomized controlled trial.
37 *Medicine*. 2016; 95(52):e5727
- 38 109. Thaker NG, Turner JD, Cobb WS, Hussain I, Janjua N, He W et al. Computed
39 tomographic angiography versus digital subtraction angiography for the postoperative
40 detection of residual aneurysms: a single-institution series and meta-analysis. *Journal
41 of Neurointerventional Surgery*. 2012; 4(3):219-225
- 42 110. Tsutsumi K, Ueki K, Morita A, Usui M, Kirino T. Risk of aneurysm recurrence in
43 patients with clipped cerebral aneurysms: results of long-term follow-up angiography.
44 *Stroke*. 2001; 32(5):1191-1194

- 1 111. Tsutsumi K, Ueki K, Usui M, Kwak S, Kirino T. Risk of recurrent subarachnoid
2 hemorrhage after complete obliteration of cerebral aneurysms. *Stroke*. 1998;
3 29(12):2511-2513
- 4 112. van Amerongen MJ, Boogaarts HD, de Vries J, Verbeek AL, Meijer FJ, Prokop M et
5 al. MRA versus DSA for follow-up of coiled intracranial aneurysms: a meta-analysis.
6 *American Journal of Neuroradiology*. 2014; 35(9):1655-1661
- 7 113. van Eijck M, Bechan RS, Sluzewski M, Peluso JP, Roks G, van Rooij WJ. Clinical
8 and imaging follow-up of patients with coiled basilar tip aneurysms up to 20 years.
9 *American Journal of Neuroradiology*. 2015; 36(11):2108-2113
- 10 114. Vendrell JF, Menjot N, Costalat V, Hoa D, Moritz J, Brunel H et al. Endovascular
11 treatment of 174 middle cerebral artery aneurysms: clinical outcome and radiologic
12 results at long-term follow-up. *Radiology*. 2009; 253(1):191-198
- 13 115. Wang CB, Shi WW, Zhang GX, Lu HC, Ma J. Flow diverter treatment of posterior
14 circulation aneurysms. A meta-analysis. *Neuroradiology*. 2016; 58(4):391-400
- 15 116. Wang T, Zhang JH, Qin X. Non-aneurysm subarachnoid hemorrhage in young adults.
16 *Acta Neurochirurgica - Supplement*. 2011; 110(Pt 1):209-213
- 17 117. Waqas M, Vakharia K, Gong AD, Rai HH, Wack A, Fayyaz N et al. One and done?
18 The effect of number of pipeline embolization devices on aneurysm treatment
19 outcomes. *Interventional Neuroradiology*. 2020; 26(2):147-155
- 20 118. Wardlaw JM, White PM. The detection and management of unruptured intracranial
21 aneurysms. *Brain*. 2000; 123(Pt 2):205-221
- 22 119. Weidauer S, Lanfermann H, Raabe A, Zanella F, Seifert V, Beck J. Impairment of
23 cerebral perfusion and infarct patterns attributable to vasospasm after aneurysmal
24 subarachnoid hemorrhage: a prospective MRI and DSA study. *Stroke*. 2007;
25 38(6):1831-1836
- 26 120. Weng HH, Jao SY, Yang CY, Tsai YH. Meta-analysis on diagnostic accuracy of MR
27 angiography in the follow-up of residual intracranial aneurysms treated with guglielmi
28 detachable coils. *Interventional Neuroradiology*. 2008; 14 (Suppl 2):53-63
- 29 121. Wenz H, Ehrlich G, Wenz R, Al Mahdi MM, Scharf J, Groden C et al. MR angiography
30 follow-up 10 years after cryptogenic nonperimesencephalic subarachnoid
31 hemorrhage. *PloS One*. 2015; 10(2):e0117925
- 32 122. Wermer MJ, Buskens E, van der Schaaf IC, Bossuyt PM, Rinkel GJ. Yield of
33 screening for new aneurysms after treatment for subarachnoid hemorrhage.
34 *Neurology*. 2004; 62(3):369-375
- 35 123. Wermer MJ, van der Schaaf IC, Velthuis BK, Algra A, Buskens E, Rinkel GJ et al.
36 Follow-up screening after subarachnoid haemorrhage: frequency and determinants of
37 new aneurysms and enlargement of existing aneurysms. *Brain*. 2005; 128(Pt
38 10):2421-2429
- 39 124. Wermer MJ, van der Schaaf IC, Velthuis BK, Majoie CB, Albrecht KW, Rinkel GJ.
40 Yield of short-term follow-up CT/MR angiography for small aneurysms detected at
41 screening. *Stroke*. 2006; 37(2):414-418
- 42 125. Wisniewski K, Tomasik B, Bobeff EJ, Stefanczyk L, Hupalo M, Jaskolski DJ.
43 Predictors for ophthalmic segment aneurysms recanalization after coiling and flow
44 diverter embolization in 6- and 12-month follow-up. *Journal of Clinical Neuroscience*.
45 2019; 68:151-157

- 1 126. Woo SW, Kim JH, Kang HI, Kim DR, Moon BG, Kim JS. High-dose simvastatin is
2 effective in preventing cerebral vasospasm after aneurysmal subarachnoid
3 hemorrhage: a prospective cohort study in Korean patients. *Journal of Korean*
4 *Neurosurgical Society*. 2015; 58(4):328-333
- 5 127. Xia Y, Ju Y, Chen J, You C. Hemorrhagic stroke and cerebral paragonimiasis. *Stroke*.
6 2014; 45(11):3420-3422
- 7 128. Yan Y, Zhu D, Tang H, Huang Q. Safety and efficacy of flow diverter treatment for
8 aneurysm in small cerebral vessels: a systematic review and meta-analysis. *World*
9 *Neurosurgery*. 2018; 115:54-64
- 10 129. Yu SC, Chan MS, Boet R, Wong JK, Lam JM, Poon WS. Intracranial aneurysms
11 treated with Guglielmi detachable coils: midterm clinical and radiological outcome in
12 97 consecutive Chinese patients in Hong Kong. *American Journal of Neuroradiology*.
13 2004; 25(2):307-313
- 14 130. Yu SC, Wong GK, Wong JK, Poon WS. Endovascular coiling versus neurosurgical
15 clipping for ruptured intracranial aneurysms: significant benefits in clinical outcome
16 and reduced consumption of hospital resources in Hong Kong Chinese patients.
17 *Hong Kong Medical Journal*. 2007; 13(4):271-278
- 18 131. Zenteno MA, Santos-Franco JA, Freitas-Modenesi JM, Gomez C, Murillo-Bonilla L,
19 Aburto-Murrieta Y et al. Use of the sole stenting technique for the management of
20 aneurysms in the posterior circulation in a prospective series of 20 patients. *Journal*
21 *of Neurosurgery*. 2008; 108(6):1104-1118
- 22 132. Zijlstra IA, Verbaan D, Majoie CB, Vandertop P, van den Berg R. Coiling and clipping
23 of middle cerebral artery aneurysms: a systematic review on clinical and imaging
24 outcome. *Journal of Neurointerventional Surgery*. 2016; 8(1):24-29
- 25
- 26

1 Appendices

2 Appendix A: Review protocols

3 Table 3: Review protocol: Imaging strategies for follow-up

ID	Field	Content
0.	PROSPERO registration number	CRD42019153672
1.	Review title	What is the clinical and cost effectiveness of different imaging strategies for follow-up of adults with confirmed aneurysmal subarachnoid haemorrhage?
2.	Review question	What is the clinical and cost effectiveness of different imaging strategies for follow-up of adults with confirmed aneurysmal subarachnoid haemorrhage?
3.	Objective	To determine the clinical and cost effectiveness of imaging strategies for subarachnoid haemorrhage.
4.	Searches	<p>The following databases will be searched:</p> <ul style="list-style-type: none"> • Cochrane Central Register of Controlled Trials (CENTRAL) • Cochrane Database of Systematic Reviews (CDSR) • Embase • MEDLINE <p>Searches will be restricted by:</p> <ul style="list-style-type: none"> • English language studies <p>The searches may be re-run 6 weeks before the final committee meeting and further studies retrieved for inclusion if relevant.</p> <p>The full search strategies will be published in the final review.</p>
5.	Condition or domain being studied	Aneurysmal subarachnoid haemorrhage
6.	Population	<p>Inclusion: Adults (16 and older) with a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm.</p> <p>Exclusion:</p> <ul style="list-style-type: none"> • Adults with subarachnoid haemorrhage caused by head injury, ischaemic stroke or an arteriovenous malformation. • Children and young people aged 15 years and younger. <p>Strata:</p> <ul style="list-style-type: none"> • If received intervention

		<ul style="list-style-type: none"> ○ Clip following aSAH ○ Coil following aSAH ● Treated conservatively (no clipping/coiling)
7.	Intervention/Exposure/Test	<p>Follow up imaging strategy at:</p> <ul style="list-style-type: none"> ● <1 year <ul style="list-style-type: none"> ○ 6-monthly follow-up ○ Yearly follow-up ● 1-2 years <ul style="list-style-type: none"> ○ 6-monthly follow-up ○ Yearly follow-up ○ Follow-up at >yearly intervals ● >2-5 years <ul style="list-style-type: none"> ○ 6-monthly follow-up ○ Yearly follow-up ○ Follow-up at >yearly intervals <p>Follow up intervention to include:</p> <ul style="list-style-type: none"> ● MR Angiography ● DSA ● CT angiography
8.	Comparator/Reference standard/Confounding factors	<p>Comparators:</p> <ul style="list-style-type: none"> ● To each other ● No imaging follow up
9.	Types of study to be included	<ul style="list-style-type: none"> ● Randomised controlled trials (RCTs), systematic reviews of RCTs. ● If insufficient RCT evidence is available, non-randomised studies will be considered if they adjust for key confounders (age), starting with prospective cohort studies.
10.	Other exclusion criteria	<p>Exclusions:</p> <ul style="list-style-type: none"> ● Non- English language studies ● Conference abstracts will be excluded as it is expected there will be sufficient full text published studies available.
11.	Context	<p>Review focused on the efficacy of imaging strategies at follow up, once the person with aSAH has been discharged.</p>
12.	Primary outcomes (critical outcomes)	<ul style="list-style-type: none"> ● Mortality ● Health and social-related quality of life (any validated score) ● Degree of disability or dependence in daily activities, (e.g. Modified Rankin Scale and patient-reported outcome measures) ● Complications of investigation (e.g. stroke, vascular injury) <p>Outcomes will be captured at or after the point of imaging follow-up. Outcomes will therefore be grouped at <1 year, 1-2 years and >2-5 years</p>

13.	Secondary outcomes (important outcomes)	<ul style="list-style-type: none"> • Subsequent subarachnoid haemorrhage • Return to daily activity (e.g. work) • Need for retreatment • Length of hospital stay (if rehospitalised) <p>Outcomes will be captured at or after the point of imaging follow-up. Outcomes will therefore be grouped at <1 year, 1-2 years and >2-5 years.</p>
14.	Data extraction (selection and coding)	<p>EndNote will be used for reference management, sifting, citations and bibliographies. All references identified by the searches and from other sources will be screened for inclusion. 10% of the abstracts will be reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer. The full text of potentially eligible studies will be retrieved and will be assessed in line with the criteria outlined above.</p> <p>EviBASE will be used for data extraction.</p>
15.	Risk of bias (quality) assessment	<p>Risk of bias will be assessed using the appropriate checklist as described in Developing NICE guidelines: the manual.</p> <p>For Intervention reviews</p> <ul style="list-style-type: none"> • Systematic reviews: Risk of Bias in Systematic Reviews (ROBIS) • Randomised Controlled Trial: Cochrane RoB (2.0) • Non randomised study, including cohort studies: Cochrane ROBINS-I <p>10% of all evidence reviews are quality assured by a senior research fellow. This includes checking:</p> <ul style="list-style-type: none"> • papers were included /excluded appropriately • a sample of the data extractions • correct methods are used to synthesise data • a sample of the risk of bias assessments <p>Disagreements between the review authors over the risk of bias in particular studies will be resolved by discussion, with involvement of a third review author where necessary.</p>
16.	Strategy for data synthesis	<ul style="list-style-type: none"> • Pairwise meta-analyses will be performed using Cochrane Review Manager (RevMan5). • GRADEpro will be used to assess the quality of evidence for each outcome, taking into account individual study quality and the meta-analysis results. The 4 main quality elements (risk of bias, indirectness, inconsistency and imprecision) will be appraised for each

		<p>outcome. Publication bias is tested for when there are more than 5 studies for an outcome.</p> <ul style="list-style-type: none"> • The risk of bias across all available evidence was evaluated for each outcome using an adaptation of the 'Grading of Recommendations Assessment, Development and Evaluation (GRADE) toolbox' developed by the international GRADE working group http://www.gradeworkinggroup.org/ • Where meta-analysis is not possible, data will be presented and quality assessed individually per outcome. • Heterogeneity between the studies in effect measures will be assessed using the I² statistic and visually inspected. An I² value greater than 50% will be considered indicative of substantial heterogeneity. Sensitivity analyses will be conducted based on pre-specified subgroups using stratified meta-analysis to explore the heterogeneity in effect estimates. If this does not explain the heterogeneity, the results will be reported in full. 		
17.	Analysis of sub-groups	<p>Strata:</p> <ul style="list-style-type: none"> • Received intervention <ul style="list-style-type: none"> ○ Clip following aSAH ○ coil following aSAH • Treated conservatively (no clipping/coiling) <p>Subgroups (if heterogeneity):</p> <ul style="list-style-type: none"> • Monitoring technique <ul style="list-style-type: none"> ○ MR Angiography ○ DSA ○ CT angiography 		
18.	Type and method of review	<input checked="" type="checkbox"/>	Intervention	
		<input type="checkbox"/>	Diagnostic	
		<input type="checkbox"/>	Prognostic	
		<input type="checkbox"/>	Qualitative	
		<input type="checkbox"/>	Epidemiologic	
		<input type="checkbox"/>	Service Delivery	
		<input type="checkbox"/>	Other (please specify)	
19.	Language	English		
20.	Country	England		
21.	Anticipated or actual start date			
22.	Anticipated completion date	3 February 2021		
23.		Review stage	Started	Completed

	Stage of review at time of this submission	Preliminary searches	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Piloting of the study selection process	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Formal screening of search results against eligibility criteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Data extraction	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Risk of bias (quality) assessment	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Data analysis	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
24.	Named contact	<p>5a. Named contact National Guideline Centre</p> <p>5b Named contact e-mail SAH@nice.org.uk</p> <p>5e Organisational affiliation of the review National Institute for Health and Care Excellence (NICE) and the National Guideline Centre</p>		
25.	Review team members	<p>From the National Guideline Centre:</p> <ul style="list-style-type: none"> • Ms Gill Ritchie • Mr Ben Mayer • Mr Audrius Stonkus • Mr Vimal Bedia • Ms Emma Cowles • Ms Jill Cobb • Ms Amelia Unsworth 		
26.	Funding sources/sponsor	<p>This systematic review is being completed by the National Guideline Centre which receives funding from NICE.</p>		
27.	Conflicts of interest	<p>All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's</p>		

		declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline.	
28.	Collaborators	Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of Developing NICE guidelines: the manual . Members of the guideline committee are available on the NICE website.	
29.	Other registration details		
30.	Reference/URL for published protocol		
31.	Dissemination plans	<p>NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as:</p> <ul style="list-style-type: none"> • notifying registered stakeholders of publication • publicising the guideline through NICE's newsletter and alerts • issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE. 	
32.	Keywords	Subarachnoid haemorrhage; imaging; follow-up	
33.	Details of existing review of same topic by same authors	None	
34.	Current review status	<input type="checkbox"/>	Ongoing
		<input type="checkbox"/>	Completed but not published
		<input type="checkbox"/>	Completed and published
		<input type="checkbox"/>	Completed, published and being updated
		<input type="checkbox"/>	Discontinued
35.	Additional information		
36.	Details of final publication	www.nice.org.uk	

1
2

1 **Table 4: Health economic review protocol**

Review question	All questions where health economic evidence applicable
Objectives	To identify health economic studies relevant to any of the review questions.
Search criteria	<ul style="list-style-type: none"> • Populations, interventions and comparators must be as specified in the clinical review protocol above. • Studies must be of a relevant health economic study design (cost–utility analysis, cost-effectiveness analysis, cost–benefit analysis, cost–consequences analysis, comparative cost analysis). • Studies must not be a letter, editorial or commentary, or a review of health economic evaluations. (Recent reviews will be ordered although not reviewed. The bibliographies will be checked for relevant studies, which will then be ordered.) • Unpublished reports will not be considered unless submitted as part of a call for evidence. • Studies must be in English.
Search strategy	A health economic study search will be undertaken using population-specific terms and a health economic study filter.
Review strategy	<p>Studies not meeting any of the search criteria above will be excluded. Studies published before 2003, abstract-only studies and studies from non-OECD countries or the USA will also be excluded.</p> <p>Each remaining study will be assessed for applicability and methodological limitations using the NICE economic evaluation checklist which can be found in appendix H of Developing NICE guidelines: the manual.⁸²</p> <p>Inclusion and exclusion criteria</p> <ul style="list-style-type: none"> • If a study is rated as both ‘Directly applicable’ and with ‘Minor limitations’ then it will be included in the guideline. A health economic evidence table will be completed and it will be included in the health economic evidence profile. • If a study is rated as either ‘Not applicable’ or with ‘Very serious limitations’ then it will usually be excluded from the guideline. If it is excluded then a health economic evidence table will not be completed and it will not be included in the health economic evidence profile. • If a study is rated as ‘Partially applicable’, with ‘Potentially serious limitations’ or both then there is discretion over whether it should be included. <p>Where there is discretion</p> <p>The health economist will decide based on the relative applicability and quality of the available evidence for that question, in discussion with the guideline committee if required. The ultimate aim is to include health economic studies that are helpful for decision-making in the context of the guideline and the current NHS setting. If several studies are considered of sufficiently high applicability and methodological quality that they could all be included, then the health economist, in discussion with the committee if required, may decide to include only the most applicable studies and to selectively exclude the remaining studies. All studies excluded based on applicability or methodological limitations will be listed with explanation in the excluded health economic studies appendix below.</p> <p>The health economist will be guided by the following hierarchies.</p> <p><i>Setting:</i></p> <ul style="list-style-type: none"> • UK NHS (most applicable). • OECD countries with predominantly public health insurance systems (for example, France, Germany, Sweden). • OECD countries with predominantly private health insurance systems (for example, Switzerland).

- Studies set in non-OECD countries or in the USA will be excluded before being assessed for applicability and methodological limitations.
- Health economic study type:*
- Cost–utility analysis (most applicable).
 - Other type of full economic evaluation (cost–benefit analysis, cost-effectiveness analysis, cost–consequences analysis).
 - Comparative cost analysis.
 - Non-comparative cost analyses including cost-of-illness studies will be excluded before being assessed for applicability and methodological limitations.
- Year of analysis:*
- The more recent the study, the more applicable it will be.
 - Studies published in 2003 or later but that depend on unit costs and resource data entirely or predominantly from before 2003 will be rated as ‘Not applicable’.
 - Studies published before 2003 will be excluded before being assessed for applicability and methodological limitations.
- Quality and relevance of effectiveness data used in the health economic analysis:*
- The more closely the clinical effectiveness data used in the health economic analysis match with the outcomes of the studies included in the clinical review the more useful the analysis will be for decision-making in the guideline.

1

2 Appendix B: Literature search strategies

3 This literature search strategy was used for the following review;

- 4 • What is the clinical and cost effectiveness of different imaging strategies for follow-up
5 of adults with confirmed aneurysmal subarachnoid haemorrhage?

6 The literature searches for this review are detailed below and complied with the methodology
7 outlined in Developing NICE guidelines: the manual⁸² For more information, please see the
8 Methods Report published as part of the accompanying documents for this guideline.

9 B.1 Clinical search literature search strategy

10 Searches were constructed using a PICO framework where population (P) terms were
11 combined with Intervention (I) and in some cases Comparison (C) terms. Outcomes (O) are
12 rarely used in search strategies for interventions as these concepts may not be well
13 described in title, abstract or indexes and therefore difficult to retrieve. Search filters were
14 applied to the search where appropriate.

15 **Table 5: Database date parameters and filters used**

Database	Dates searched	Search filter used
Medline (OVID)	1946 – 26 June 2020	Exclusions Randomised controlled trials Systematic review studies Observational studies
Embase (OVID)	1974 – 26 June 2020	Exclusions Randomised controlled trials Systematic review studies Observational studies
The Cochrane Library (Wiley)	Cochrane Reviews to 2020 Issue 6 of 12	None

Database	Dates searched	Search filter used
	CENTRAL to 2020 Issue 6 of 12	

1 Medline (Ovid) search terms

1.	exp Subarachnoid Hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp Intracranial Aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.
15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16
18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23
25.	6 not 24
26.	(exp child/ or exp pediatrics/ or exp infant/) not (exp adolescent/ or exp adult/ or exp middle age/ or exp aged/)
27.	25 not 26
28.	limit 27 to English language
29.	Epidemiologic studies/
30.	Observational study/
31.	exp Cohort studies/
32.	(cohort adj (study or studies or analys* or data)).ti,ab.
33.	((follow up or observational or uncontrolled or non randomi#ed or epidemiologic*) adj (study or studies or data)).ti,ab.
34.	((longitudinal or retrospective or prospective or cross sectional) and (study or studies or review or analys* or cohort* or data)).ti,ab.
35.	Controlled Before-After Studies/
36.	Historically Controlled Study/
37.	Interrupted Time Series Analysis/

38.	(before adj2 after adj2 (study or studies or data)).ti,ab.
39.	or/29-38
40.	exp case control study/
41.	case control*.ti,ab.
42.	or/40-41
43.	39 or 42
44.	Cross-sectional studies/
45.	(cross sectional and (study or studies or review or analys* or cohort* or data)).ti,ab.
46.	or/44-45
47.	39 or 46
48.	39 or 42 or 46
49.	Meta-Analysis/
50.	exp Meta-Analysis as Topic/
51.	(meta analy* or metanaly* or metaanaly* or meta regression).ti,ab.
52.	((systematic* or evidence*) adj3 (review* or overview*)).ti,ab.
53.	(reference list* or bibliograph* or hand search* or manual search* or relevant journals).ab.
54.	(search strategy or search criteria or systematic search or study selection or data extraction).ab.
55.	(search* adj4 literature).ab.
56.	(medline or pubmed or cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or science citation index or bids or cancerlit).ab.
57.	cochrane.jw.
58.	((multiple treatment* or indirect or mixed) adj2 comparison*).ti,ab.
59.	or/49-57
60.	randomized controlled trial.pt.
61.	controlled clinical trial.pt.
62.	randomi#ed.ti,ab.
63.	placebo.ab.
64.	randomly.ti,ab.
65.	Clinical Trials as topic.sh.
66.	trial.ti.
67.	or/60-66
68.	Magnetic Resonance Angiography/ or Angiography, Digital Subtraction/ or Computed Tomography Angiography/
69.	((magnetic resonance or digital subtraction or computed tomograph*) adj3 angiograph*).ti,ab.
70.	((MR or DS or CT) adj3 (angiograph* or angiogram*)).ti,ab.
71.	(MRA or DSA or CTA).ti,ab.
72.	or/68-71
73.	28 and 72 and (48 or 59 or 67)

1 Embase (Ovid) search terms

1.	*subarachnoid hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.

4.	exp intracranial aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	Case report/ or Case study/
11.	(letter or comment*).ti.
12.	or/7-11
13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	Nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental animal/
19.	Animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	(exp child/ or exp pediatrics/) not (exp adult/ or exp adolescent/)
25.	23 not 24
26.	limit 25 to English language
27.	Clinical study/
28.	Observational study/
29.	family study/
30.	longitudinal study/
31.	retrospective study/
32.	prospective study/
33.	cohort analysis/
34.	follow-up/
35.	cohort*.ti,ab.
36.	34 and 35
37.	(cohort adj (study or studies or analys* or data)).ti,ab.
38.	((follow up or observational or uncontrolled or non randomi#ed or epidemiologic*) adj (study or studies or data)).ti,ab.
39.	((longitudinal or retrospective or prospective or cross sectional) and (study or studies or review or analys* or cohort* or data)).ti,ab.
40.	(before adj2 after adj2 (study or studies or data)).ti,ab.
41.	or/27-33,36-40
42.	exp case control study/
43.	case control*.ti,ab.
44.	or/42-43
45.	41 or 44

46.	cross-sectional study/
47.	(cross sectional and (study or studies or review or analys* or cohort* or data)).ti,ab.
48.	or/46-47
49.	41 or 48
50.	41 or 44 or 48
51.	random*.ti,ab.
52.	factorial*.ti,ab.
53.	(crossover* or cross over*).ti,ab.
54.	((doubl* or singl*) adj blind*).ti,ab.
55.	(assign* or allocat* or volunteer* or placebo*).ti,ab.
56.	crossover procedure/
57.	single blind procedure/
58.	randomized controlled trial/
59.	double blind procedure/
60.	or/51-59
61.	systematic review/
62.	meta-analysis/
63.	(meta analy* or metanaly* or metaanaly* or meta regression).ti,ab.
64.	((systematic or evidence) adj3 (review* or overview*)).ti,ab.
65.	(reference list* or bibliograph* or hand search* or manual search* or relevant journals).ab.
66.	(search strategy or search criteria or systematic search or study selection or data extraction).ab.
67.	(search* adj4 literature).ab.
68.	(medline or pubmed or cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or science citation index or bids or cancerlit).ab.
69.	((pool* or combined) adj2 (data or trials or studies or results)).ab.
70.	cochrane.jw.
71.	((multiple treatment* or indirect or mixed) adj2 comparison*).ti,ab.
72.	or/61-70
73.	magnetic resonance angiography/ or computed tomographic angiography/ or digital subtraction angiography/
74.	((magnetic resonance or digital subtraction or computed tomograph*) adj3 angiograph*).ti,ab.
75.	((MR or DS or CT) adj3 (angiograph* or angiogram*)).ti,ab.
76.	(MRA or DSA or CTA).ti,ab.
77.	or/73-76
78.	26 and 77 and (50 or 60 or 72)

1 Cochrane Library (Wiley) search terms

#1.	MeSH descriptor: [Subarachnoid Hemorrhage] explode all trees
#2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) near/3 (hemorrhag* or haemorrhag* or bleed* or blood*)):ti,ab
#3.	(SAH or aSAH):ti,ab
#4.	MeSH descriptor: [Intracranial Aneurysm] explode all trees
#5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) near/3 (aneurysm* or aneurism* or hematoma* or haematoma*)):ti,ab

#6.	(or #1-#5)
#7.	MeSH descriptor: [Magnetic Resonance Angiography] this term only
#8.	MeSH descriptor: [Angiography, Digital Subtraction] this term only
#9.	MeSH descriptor: [Computed Tomography Angiography] this term only
#10.	((magnetic resonance or digital subtraction or computed tomograph*) near/3 angiograph*):ti,ab
#11.	((MR or DS or CT) near/3 (angiograph* or angiogram*)):ti,ab
#12.	(MRA or DSA or CTA):ti,ab
#13.	(or #7-#12)
#14.	#6 and #13

B.2.1 Health Economics literature search strategy

2 Health economic evidence was identified by conducting a broad search relating to
3 subarachnoid haemorrhage population in NHS Economic Evaluation Database (NHS EED –
4 this ceased to be updated after March 2015) and the Health Technology Assessment
5 database (HTA) with no date restrictions. NHS EED and HTA databases are hosted by the
6 Centre for Research and Dissemination (CRD). Additional searches were run on Medline and
7 Embase.

8 **Table 6: Database date parameters and filters used**

Database	Dates searched	Search filter used
Medline	2003 – 23 June 2020	Exclusions Health economics studies
Embase	2003 – 23 June 2020	Exclusions Health economics studies
Centre for Research and Dissemination (CRD)	HTA - Inception – 23 June 2020 NHSEED - Inception to March 2015	None

9 Medline (Ovid) search terms

1.	exp Subarachnoid Hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)):ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp Intracranial Aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)):ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.

15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16
18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23
25.	6 not 24
26.	limit 25 to English language
27.	Economics/
28.	Value of life/
29.	exp "Costs and Cost Analysis"/
30.	exp Economics, Hospital/
31.	exp Economics, Medical/
32.	Economics, Nursing/
33.	Economics, Pharmaceutical/
34.	exp "Fees and Charges"/
35.	exp Budgets/
36.	budget*.ti,ab.
37.	cost*.ti.
38.	(economic* or pharmaco?economic*).ti.
39.	(price* or pricing*).ti,ab.
40.	(cost* adj2 (effective* or utilit* or benefit* or minimi* or unit* or estimat* or variable*)).ab.
41.	(financ* or fee or fees).ti,ab.
42.	(value adj2 (money or monetary)).ti,ab.
43.	or/27-42
44.	26 and 43

1 Embase (Ovid) search terms

1.	subarachnoid hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp intracranial aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	case report/ or case study/

11.	(letter or comment*).ti.
12.	or/7-11
13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental Animal/
19.	animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	limit 23 to English language
25.	health economics/
26.	exp economic evaluation/
27.	exp health care cost/
28.	exp fee/
29.	budget/
30.	funding/
31.	budget*.ti,ab.
32.	cost*.ti.
33.	(economic* or pharmaco?economic*).ti.
34.	(price* or pricing*).ti,ab.
35.	(cost* adj2 (effective* or utilit* or benefit* or minimi* or unit* or estimat* or variable*)).ab.
36.	(financ* or fee or fees).ti,ab.
37.	(value adj2 (money or monetary)).ti,ab.
38.	or/25-37
39.	24 and 38

1 NHS EED and HTA (CRD) search terms


#1.	MeSH DESCRIPTOR Subarachnoid Hemorrhage EXPLODE ALL TREES
#2.	MeSH DESCRIPTOR Intracranial Hemorrhages EXPLODE ALL TREES
#3.	((((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)))
#4.	((SAH or aSAH))
#5.	#1 OR #2 OR #3 OR #4
#6.	MeSH DESCRIPTOR Aneurysm EXPLODE ALL TREES
#7.	((aneurysm* or hematoma* or haematoma*))
#8.	#6 OR #7
#9.	MeSH DESCRIPTOR Intracranial Aneurysm EXPLODE ALL TREES
#10.	((((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (aneurysm* or hematoma* or haematoma*)))
#11.	#9 OR #10
#12.	MeSH DESCRIPTOR Aneurysm, ruptured

#13.	(((ruptur* or weak* or brain or trauma*) adj3 (aneurysm* or hematoma* or haematoma*)))
#14.	#12 OR #13
#15.	(#5 or #8 or #11 or #14)

1

1 Appendix C: Clinical evidence selection

Figure 1: Flow chart of clinical study selection for the review of imaging strategies for follow-up



2

3

1 **Appendix D: Clinical evidence tables**

2 No studies were included

3

1 **Appendix E: Forest plots**

2 No studies were included

3

4

5

6

7

1 **Appendix F: GRADE tables**

2 No studies were included

3

4


5

6

7

1 **Appendix G: Health economic evidence** 2 **selection**

Figure 2: Flow chart of health economic study selection for the guideline



* Non-relevant population, intervention, comparison, design or setting; non-English language

1 **Appendix H: Health economic evidence tables**

2 None.

3

4

1

2 Appendix I: Excluded studies

I.1.3 Excluded clinical studies

4 Table 7: Studies excluded from the clinical review

Study	Exclusion reason
Aboukais 2015 ¹	Not in English
Adeeb 2017 ²	Inappropriate population – ophthalmic segment aneurysm
Ahmed 2019 ³	Systematic review - references checked
Anzalone 2015 ⁴	Inappropriate study design – non comparative study
Arrese 2013 ⁵	Systematic review - references checked
Arthur 2019 ⁶	Inappropriate study design – non comparative study
Atasoy 2019 ⁷	Not review population - patients with unruptured aneurysms
Aydin 2012 ⁹	Inappropriate study design – non comparative study
Aydin 2019 ⁸	Not review population - patients with wide-necked complex bifurcation aneurysms (99 out of 102 unruptured)
Bakker 2014 ¹⁰	Systematic review - references checked
Bendok 2013 ¹¹	Inappropriate study design – no relevant outcomes
Bor 2014 ¹²	Inappropriate population – relatives of people with SAH
Boussel 2008 ¹³	Inappropriate study design - no relevant outcomes
Bracard 2002 ¹⁴	Inappropriate comparison – single cohort studied at multiple time points
Bruneau 2011 ¹⁵	Inappropriate intervention – angiogram 10 years post intervention (no previous angiogram)
CBS Office of Statistics Netherlands 2000 ¹⁶	Paper not available
Chalouhi 2014 ¹⁷	Inappropriate study design - no relevant outcomes
Chen 2018 ¹⁸	Inappropriate comparison/No relevant outcomes – single cohort studied at multiple time points
Cho 2014 ²⁰	Inappropriate study design - no relevant outcomes
Cho 2015 ¹⁹	Inappropriate study design - no relevant outcomes
Cloft 1999 ²¹	Inappropriate study design - no relevant outcomes
Cognard 2015 ²²	Inappropriate study design / inappropriate population - no relevant outcomes/ all forms of aneurysm
Crawley 1999 ²³	Inappropriate study design – theoretical models
David 1999 ²⁴	Inappropriate study design – non comparative study
De Letter 1994 ²⁵	Inappropriate comparison/ Inappropriate population – patching after carotid endarterectomy
Delgado Almandoz 2012 ²⁶	Inappropriate study design - no relevant outcomes
Disney 1988 ²⁷	Inappropriate study design – case series
Dorhout Mees 2007 ²⁸	Systematic review - references checked
Flores 2019 ²⁹	Systematic review - references checked
Fountas 2008 ³⁰	Inappropriate comparison – no angiographic follow up
Fujimura 2017 ³¹	Inappropriate study design/inappropriate comparison – medication comparison

Study	Exclusion reason
Gallas 2005 ³³	Inappropriate study design – non comparative study
Gallas 2009 ³²	Inappropriate study design – non comparative study
Garg 2013 ³⁴	Inappropriate intervention – no angiographic follow up
Gauvrit 2005 ³⁵	Inappropriate study design - no relevant outcomes
Geng 2019 ³⁶	Systematic review - references checked
Germano 2013 ³⁷	Inappropriate comparison – MRA at follow up
Geyik 2013 ³⁸	Inappropriate study design - no relevant outcomes
Ghogawala 2013 ³⁹	Inappropriate population – carotid artery stenosis
Gibbs 2004 ⁴⁰	Inappropriate population – people with polycystic kidney disease and SAH
Goksu 2015 ⁴¹	Inappropriate study design – non comparative study
Groden 2003 ⁴²	Inappropriate study design – non comparative study
Hai 2009 ⁴³	Inappropriate study design - no relevant outcomes
Hashimoto 2000 ⁴⁴	Inappropriate study design / no relevant outcomes – non comparative study / no angiographic follow up results
Hassan 2012 ⁴⁵	Inappropriate population – arterial dissection
Heller 2013 ⁴⁶	Inappropriate study design - no relevant outcomes
Helthuis 2018 ⁴⁷	Inappropriate study design / no relevant outcomes – case series
Hendryk 2004 ⁴⁸	Inappropriate intervention – no angiographic follow up
Hijdra 1987 ⁴⁹	Inappropriate study design - no relevant outcomes
Hop 1997 ⁵⁰	Systematic review - references checked
Hosono 2002 ⁵¹	Inappropriate comparison - single photon emission CT (SPECT performed at 1 and 7 days after SAH)
Hussain 2009 ⁵²	Inappropriate comparison- results of last follow up
Ji 2016 ⁵³	Inappropriate study design – non comparative study
Juvela 1993 ⁵⁴	Inappropriate population – unruptured aneurysms
Juvela 2001 ⁵⁵	Inappropriate study design – non comparative study
Kalra 2015 ⁵⁶	Systematic review - references checked
Kannath 2019 ⁵⁷	Inappropriate study design - non comparative study/case series
Kao 2018 ⁵⁸	Not review population - patients with chronic carotid artery occlusion
Kapapa 2014 ⁵⁹	Inappropriate study design – surveys and questionnaires
Kasner 1997 ⁶⁰	Inappropriate population – arterial dissection
Kim 2018 ⁶³	Inappropriate study design – non comparative study
Kim 2018 ⁶²	Inappropriate study design – non comparative study
Kim 2019 ⁶¹	Not review population - patients with unruptured aneurysms
King 1994 ⁶⁴	Systematic review - references checked
Kwee 2007 ⁶⁵	Systematic review - references checked
Lindgren 2018 ⁶⁶	Systematic review - references checked
Lindvall 2012 ⁶⁷	Inappropriate study design – non comparative study
Liu 2018 ⁶⁸	Inappropriate study design - no relevant outcomes
Lizza 2014 ⁶⁹	Inappropriate comparison – stroke or delayed cerebral infarction
Lublinsky 2019 ⁷⁰	Inappropriate study design - no relevant outcomes
Ma 2015 ⁷¹	Inappropriate intervention – flow cytometry
Maimaitili 2013 ⁷²	Inappropriate study design - no relevant outcomes
Marquardt 2000 ⁷³	Inappropriate study design - no relevant outcomes

Study	Exclusion reason
Mooney 2018 ⁷⁴	Inappropriate study design/inappropriate comparison – no angiographic follow up
Morais 2017 ⁷⁵	Inappropriate comparison – single cohort studied at multiple time points
Moritz 2012 ⁷⁶	Inappropriate study design - no relevant outcomes
Mortimer 2015 ⁷⁷	Inappropriate study design - no relevant outcomes
Murakami 2019 ⁷⁸	Not review population - patients with unruptured aneurysms
Murias Quintana 2015 ⁷⁹	Inappropriate study design – non comparative study
Nagano 2016 ⁸⁰	inappropriate comparison – no angiographic follow up
Nakagawa 2017 ⁸¹	Inappropriate comparison – DSA up to 10 days post SAH
Niuro 2000 ⁸⁴	Inappropriate population – all types of aneurysm
Nosseck 2015 ⁸⁵	Inappropriate study design - no relevant outcomes
Ocal 2019 ⁸⁶	Inappropriate comparison - surpass flow diverter only versus stent + flow diverter
Okada 2016 ⁸⁷	Inappropriate study design no relevant outcomes
Peschillo 2017 ⁸⁸	Inappropriate study design – case series
Petridis 2017 ⁸⁹	Inappropriate study design - no relevant outcomes
Piano 2020 ⁹⁰	Inappropriate study design – non comparative study
Pierot 2015 ⁹¹	Systematic review - references checked
Piske 2009 ⁹²	Inappropriate comparison – outcomes compared by aneurysm size
Poon 2006 ⁹³	Systematic review - references checked
Potter 2016 ⁹⁴	Systematic review - references checked
Proust 2010 ⁹⁵	Inappropriate comparison – outcomes 6 months post-surgery
Qin 2017 ⁹⁶	Inappropriate study design – non comparative study
Raffi 2003 ⁹⁷	Inappropriate study design - No relevant outcomes
Raschi 2012 ⁹⁸	Inappropriate study design – modelling
Rautio 2019 ⁹⁹	Inappropriate study design – non-comparative study
Ruppert 2007 ¹⁰⁰	Inappropriate study design - no relevant outcomes
Saatci 2003 ¹⁰¹	Inappropriate study design - non-comparative study
Serafin 2011 ¹⁰²	Systematic review - references checked
Slater 2015 ¹⁰³	Inappropriate study design - no relevant outcomes
Slosberg 1997 ¹⁰⁴	Inappropriate comparison – overall results
Sprengers 2009 ¹⁰⁵	Inappropriate study design – non comparative study
Sun 2013 ¹⁰⁶	Systematic review - references checked
Taylor, 2010 ¹⁰⁷	Inappropriate study design - non-comparative,
Tao 2016 ¹⁰⁸	Inappropriate study design – study protocol
Thaker 2012 ¹⁰⁹	Inappropriate study design – no comparison group
Tsutsumi 2001 ¹¹⁰	Inappropriate study design – non comparative study
Tsutsumi 1998 ¹¹¹	Inappropriate study design – non comparative study
van Amerongen 2014 ¹¹²	Systematic review - references checked
van Eijck 2015 ¹¹³	Inappropriate study design - no relevant outcomes
Vendrell 2009 ¹¹⁴	Inappropriate comparison – single cohort studied at multiple time points
Wang 2011 ¹¹⁶	Inappropriate study design – no angiographic follow up
Wang 2016 ¹¹⁵	Systematic review - references checked
Waqas 2020 ¹¹⁷	No relevant outcomes

Study	Exclusion reason
Wardlaw 2000 ¹¹⁸	Inappropriate population – unruptured aneurysms
Weidauer 2007 ¹¹⁹	Inappropriate study design - no relevant outcomes
Weng 2008 ¹²⁰	Systematic review - references checked
Wenz 2015 ¹²¹	Inappropriate study design – non comparative study
Wermer 2005 ¹²³	Inappropriate study design – non comparative study
Wermer 2006 ¹²⁴	Inappropriate study design – non comparative study
Wermer 2004 ¹²²	Inappropriate study design- Markov economical model
Wisniewski 2019 ¹²⁵	No relevant outcomes
Woo 2015 ¹²⁶	Inappropriate comparison – no angiographic follow up
Xia 2014 ¹²⁷	Inappropriate population - Pargonimiasis
Yan 2018 ¹²⁸	Systematic review - references checked
Yu 2004 ¹²⁹	Inappropriate comparison – patients with SAH and mass effect / all patients followed up
Yu 2007 ¹³⁰	Inappropriate study design - no relevant outcomes
Zenteno 2008 ¹³¹	Inappropriate study design – case series
Zijlstra 2016 ¹³²	Systematic review - references checked

1

I.2.2 Excluded health economic studies

3 Published health economic studies that met the inclusion criteria (relevant population,
4 comparators, economic study design, published 2003 or later and not from non-OECD
5 country or USA) but that were excluded following appraisal of applicability and
6 methodological quality are listed below. See the health economic protocol for more details.

7 **Table 8: Studies excluded from the health economic review**

Reference	Reason for exclusion
None.	

8