

Tenotomy of horizontal eye muscles for nystagmus (with reattachment at their original insertions)

Interventional procedures guidance

Published: 27 May 2009

www.nice.org.uk/guidance/ipg299

1 Guidance

- 1.1 The evidence on tenotomy of horizontal eye muscles for nystagmus (with reattachment at their original insertions) raises no major safety concerns, but current evidence on its efficacy is inadequate in quantity. Therefore, this procedure should only be used with special arrangements for clinical governance, consent, and audit or research.
- 1.2 Clinicians wishing to undertake tenotomy of horizontal eye muscles for nystagmus (with reattachment at their original insertions) should take the following actions.
 - Inform the clinical governance leads in their Trusts.

- Ensure that patients and their carers understand the uncertainty about the procedure's safety and efficacy and provide them with clear written information. In addition, the use of NICE's [information for patients](#) ('Understanding NICE guidance') is recommended.
 - Audit and review clinical outcomes of all patients having tenotomy of horizontal eye muscles for nystagmus (with reattachment at their original insertions) (see section 3.1).
- 1.3 Patient selection and follow-up should take place in specialist units with experience in the management of ocular motility disorders.
- 1.4 NICE encourages further collaborative data collection, including information on visual acuity and quality of life, and may review the procedure on publication of further evidence.

2 The procedure

2.1 Indications and current treatments

- 2.1.1 Nystagmus is an involuntary oscillatory movement of the eyes, usually from side to side, but sometimes the eyes move up and down or in a circular motion. Most people with nystagmus have impaired vision.
- 2.1.2 There are several types of nystagmus but there is no definitive classification system to describe them. Nystagmus may be present at birth, caused by defects in the eye or the visual pathway from the eye to the brain. It can occur in a wide range of childhood eye disorders and may be found in children with multiple disabilities. Nystagmus can also develop later in life as a symptom of a variety of conditions, including stroke, multiple sclerosis or head injury.
- 2.1.3 There is currently no curative treatment for nystagmus. Spectacles or contact lenses may be worn to improve visual acuity but they do not correct the nystagmus.

2.2 Outline of the procedure

- 2.2.1 Tenotomy for nystagmus is carried out with the patient under general anaesthesia, and involves division of the attachments of the two horizontal rectus muscles (lateral and medial) of each eye. A limbal incision is made in the conjunctiva and each muscle is detached from the sclera. The muscles are then reattached at their original places of insertion.
- 2.2.2 The aims of the procedure are to reduce the frequency and amplitude of nystagmus (how often and how far the eyes oscillate) and to improve visual acuity. The exact mechanism by which this procedure might improve nystagmus is unknown.

Sections 2.3 and 2.4 describe efficacy and safety outcomes which were available in the published literature and which the Committee considered as part of the evidence about this procedure. For more detailed information on the evidence, see the [overview](#).

2.3 Efficacy

- 2.3.1 In a case series of ten adults treated with tenotomy, nine patients had increased and one patient had decreased expanded Nystagmus Acuity Function (NAFX) scores at 1 year (this score is an objective measure used to predict visual acuity on the basis of the patient's foveation period, defined as their ability to fix an image on the fovea, the most visually precise part of the retina). A case series of nine patients with infantile nystagmus reported that eight patients had increased NAFX scores (mean 60%) at 1 year. The NAFX score remained unchanged in one patient at 1 year. All nine patients had reduced nystagmus amplitude (mean 33%) and increased foveation period (mean 104%) at 1 year. In a case series of five patients with infantile nystagmus, two were assessed for eye movement. The NAFX scores improved by 8% in one child at 1 year and by 36% in the other child at 6 months.
- 2.3.2 Two case series of five children and ten adults reported improvements in best-corrected visual acuity of at least five letters on the Early Treatment

Diabetic Retinopathy Study chart (corresponding to a one-line improvement on the Snellen chart) in four and five patients, respectively (assessed at 6 weeks and 12 months, respectively). The case series of nine patients reported that three patients had an improvement in visual acuity of at least one line, three had an improvement of a few letters and three had no change (method of assessment and follow-up was not described).

- 2.3.3 In a case series of five patients, all patients had a reduced time to target acquisition (measured by infrared reflection or high-speed digital video) 1 year after the procedure (precise reduction not stated).
- 2.3.4 The Specialist Advisers considered efficacy outcomes to include best-corrected binocular visual acuity under varying gaze angles, null point width, stereoacuity, ocular movement recordings, nystagmus, visual function in day-to-day life, quality of life, cosmesis and head posture.

2.4 Safety

- 2.4.1 No adverse events were reported in the literature.
- 2.4.2 The Specialist Advisers considered theoretical adverse events to include damage to the retina or perforation of the globe, infection and misalignment of muscles causing redness, swelling, diplopia, induced strabismus and possible loss of vision. Revision surgery may be needed to correct these adverse outcomes. One Specialist Adviser stated that development of a conjunctival cyst had been reported in the literature.

3 Further information

- 3.1 This guidance requires that clinicians undertaking the procedure make special arrangements for audit. NICE has identified relevant audit criteria and has developed [audit support](#) (which is for use at local discretion).

Information for patients

NICE has produced [information on this procedure for patients and carers](#) ('Understanding

NICE guidance'). It explains the nature of the procedure and the guidance issued by NICE, and has been written with patient consent in mind.

A large print version is also available.

4 About this guidance

NICE interventional procedure guidance makes recommendations on the safety and efficacy of the procedure. It does not cover whether or not the NHS should fund a procedure. Funding decisions are taken by local NHS bodies after considering the clinical effectiveness of the procedure and whether it represents value for money for the NHS. It is for healthcare professionals and people using the NHS in England, Wales, Scotland and Northern Ireland, and is endorsed by Healthcare Improvement Scotland for implementation by NHSScotland.

This guidance was developed using the NICE [interventional procedure guidance](#) process.

We have produced a [summary of this guidance for patients and carers](#). Tools to help you put the guidance into practice and information about the evidence it is based on are also [available](#).

Changes since publication

7 January 2012: minor maintenance.

Your responsibility

This guidance represents the views of NICE and was arrived at after careful consideration of the available evidence. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of healthcare professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have regard to promoting equality of opportunity. Nothing in this

guidance should be interpreted in a way which would be inconsistent with compliance with those duties.

Copyright

© National Institute for Health and Clinical Excellence 2009. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

Contact NICE

National Institute for Health and Clinical Excellence
Level 1A, City Tower, Piccadilly Plaza, Manchester M1 4BT

www.nice.org.uk
nice@nice.org.uk
0845 033 7780

Endorsing organisation

This guidance has been endorsed by [Healthcare Improvement Scotland](#).

Accreditation

