

## Abdominal aortic aneurysm: diagnosis and management

Evidence review N: Signs, symptoms and risk factors predicting ruptured or symptomatic unruptured aneurysms before arrival at the hospital, and in non-specialist hospital settings

*NICE guideline <number>*

*Evidence reviews*

*May 2018*

*Draft for Consultation*

*Commissioned by the National Institute  
for Health and Care Excellence*



## **Disclaimer**

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

## **Copyright**

© NICE [2018]. All rights reserved. Subject to [Notice of rights](#).

ISBN:

## Contents

Predicting ruptured or symptomatic aneurysms before arrival at the hospital, and in non-specialist hospital settings .....	5
Review question .....	5
Introduction .....	5
PICO 5	
Methods and process .....	6
Clinical evidence .....	6
Summary of clinical studies included in the evidence review .....	6
Quality assessment of clinical studies included in the evidence review .....	7
Economic evidence .....	7
Evidence statements .....	7
Recommendations .....	7
Rationale and impact.....	7
The committee’s discussion of the evidence.....	8
<b>Appendices .....</b>	<b>10</b>
Appendix A – Review protocols .....	10
Review protocol for risk factors associated with aneurysm growth or rupture .....	10
Appendix B – Literature search strategies .....	12
Clinical search literature search strategy .....	12
Health Economics literature search strategy .....	13
Appendix C – Clinical evidence study selection .....	16
Appendix D – Economic evidence study selection .....	17
Appendix E – Excluded studies .....	18
Clinical studies .....	18
Economic studies .....	18
Appendix F – Glossary .....	19

# 1 Predicting ruptured or symptomatic 2 aneurysms before arrival at the 3 hospital, and in non-specialist hospital 4 settings

## 5 Review question

6 Which signs, symptoms, risk factors (or combinations of these) and diagnostic  
7 assessment tools are most accurate in indicating the presence of a ruptured or  
8 symptomatic unruptured abdominal aortic aneurysm i) before arrival at the hospital  
9 and ii) in a non-specialist hospital setting?

## 10 Introduction

11 This review question aims to determine which signs, symptoms and risk factors:

- 12 • predict the presence of 'emergency' (that is, ruptured or symptomatic  
13 unruptured) abdominal aortic aneurysms
- 14 • would lead to immediate transfer to a specialist vascular unit (i.e. bypassing  
15 non-specialist A&E)
- 16 • should initiate definitive imaging in the emergency department (i.e. sufficient  
17 suspicion to initiate imaging, but not so high that a patient should be  
18 transferred to a vascular unit without imaging)

## 19 PICO

### 20 Table 1: Inclusion criteria

Parameter	Inclusion criteria
Population	<ul style="list-style-type: none"> <li>• People with a suspected ruptured or symptomatic unruptured abdominal aortic aneurysm</li> </ul>
Index test / factors of interest	<ul style="list-style-type: none"> <li>• Pulsatile abdominal mass</li> <li>• Abdominal, back or groin pain</li> <li>• Sweating</li> <li>• Clamminess</li> <li>• Shock</li> <li>• Altered mental/cognitive status</li> <li>• Loss of consciousness</li> <li>• Collapse</li> <li>• Paralysis</li> <li>• Dizziness</li> <li>• Nausea</li> <li>• Vomiting</li> <li>• Hypotension</li> <li>• Fast pulse / tachycardia</li> <li>• Shortness of breath</li> <li>• Cyanosis</li> </ul>

Parameter	Inclusion criteria
	<ul style="list-style-type: none"> <li>• Urinary symptoms</li> <li>• Colic</li> <li>• Peripheral ischemia / absent peripheral pulses</li> <li>• Temperature</li> <li>• Capillary refill time</li> <li>• Ultrasound ('fast scan')</li> </ul>
Endpoints	<ul style="list-style-type: none"> <li>• Surgical or CT confirmation of rupture</li> <li>• CT confirmation of AAA accompanied by symptoms</li> </ul>

## 21 Methods and process

22 This evidence review was developed using the methods and process described in  
23 [Developing NICE guidelines: the manual](#). Methods specific to this review question  
24 are described in the review protocol in Appendix A.

25 Declarations of interest were recorded according to NICE's 2014 conflicts of interest  
26 policy.

27 A broad search was used to identify all studies that examine the diagnosis,  
28 surveillance or monitoring of abdominal aortic aneurysms (AAA). This was a 'bulk'  
29 search covering multiple review questions. The database was sifted to identify all  
30 studies that met the criteria detailed in Table 1. The relevant review protocol can be  
31 found in Appendix A.

32 Studies were considered for inclusion if they were were cross-sectional studies or  
33 systematic reviews (of cross-sectional studies) exploring signs, symptoms or risk  
34 factors indicative of AAA rupture outside a regional vascular service setting.

35 Studies were excluded if they:

- 36 • were not in English
- 37 • were not full reports of the study (for example, published only as an abstract)
- 38 • were not peer-reviewed.

## 39 Clinical evidence

### 40 Included studies

41 From an initial database of 16,274 abstracts, 2 were identified as being potentially  
42 relevant. Following full-text review of these articles, no studies were included.

43 An update search was conducted in December 2017, to identify any relevant studies  
44 published during guideline development. The search found 2,180 abstracts; all of  
45 which were not considered relevant to this review question. As a result no additional  
46 studies were included.

### 47 Excluded studies

48 The list of papers excluded at full-text review, with reasons, is given in Appendix E.

## 49 Summary of clinical studies included in the evidence review

50 No studies were included following full text review.

## 51 **Quality assessment of clinical studies included in the evidence review**

52 No studies were included following full text review.

## 53 **Economic evidence**

### 54 **Included studies**

55 A literature search was conducted jointly for all review questions by applying  
56 standard health economic filters to a clinical search for AAA. This search returned a  
57 total of 5,173 citations. Following review of all titles and abstracts, no studies were  
58 identified as being potentially relevant to the review question. No full texts were  
59 retrieved, and no studies were included as economic evidence.

60 An update search was conducted in December 2017, to identify any relevant health  
61 economic analyses published during guideline development. The search found 814  
62 abstracts; all of which were not considered relevant to this review question. As a  
63 result no additional studies were included.

### 64 **Excluded studies**

65 No studies were retrieved for full-text review.

## 66 **Evidence statements**

67 No evidence was identified for this review question.

## 68 **Recommendations**

69 N1. Think about the possibility of ruptured AAA in people with new abdominal and/or  
70 back pain, cardiovascular collapse, or loss of consciousness. Be aware that ruptured  
71 AAA is particularly likely if they also have any of the following risk factors:

- 72 • an existing diagnosis of AAA
- 73 • age over 60
- 74 • they smoke or used to smoke
- 75 • history of hypertension.

76 N2. Be aware that AAAs are more likely to rupture in women than men.

77 N3. Immediately discuss all people with suspected ruptured AAA with a regional  
78 vascular service.

## 79 **Rationale and impact**

### 80 **Why the committee made the recommendations**

81 Based on their own experience, the committee highlighted the most important signs  
82 and symptoms of ruptured AAAs, because:

- 83 • non-specialists commonly misdiagnose them
- 84 • reducing misdiagnosis should increase the chance of survival
- 85 • urgent discussion of a suspected ruptured AAA with a regional vascular service  
86 will improve the chances of appropriate treatment and survival.

**87 Impact of the recommendations on practice**

88 There is variation in awareness of AAAs among non-specialists. Implementing the  
89 recommendations should reduce this variation and increase the chance of ruptured  
90 AAAs being diagnosed earlier.

**91 The committee's discussion of the evidence****92 Interpreting the evidence****93 *The outcomes that matter most***

94 The committee considered that the outcomes that matter most in non-specialist  
95 settings were symptoms and signs that, when considered in combination with risk  
96 factors, could be used as a strong indication of the presence of a ruptured aneurysm.

**97 *The quality of the evidence***

98 The committee noted that there are currently no national criteria for predicting the  
99 presence of ruptured AAAs and the use of criteria varied according to local  
100 arrangements. In the absence of any evidence from prospective observational  
101 studies, the committee agreed that consensus recommendations were needed to  
102 ensure that clinicians in non-specialist settings are clear when a ruptured AAA should  
103 form part of a differential diagnosis.

104 Following discussion of the range of symptoms and risk factors presented in the  
105 review protocol the committee agreed that it would not be feasible to provide a  
106 comprehensive list of factors indicative of ruptured AAAs in non-specialist care  
107 settings. The committee also agreed that individual symptoms and signs are not  
108 usually used in isolation to predict the presence of ruptured aneurysms. As a result,  
109 the committee discussed key symptoms and signs that, when considered in  
110 combination with risk factors, would give the strongest indication of a ruptured  
111 aneurysm.

112 After considering the symptoms, signs and risk factors outlined in the review protocol,  
113 the committee agreed that a change in body temperature was not sensitive enough  
114 to predict the presence of a ruptured AAA in non-specialist care settings. Conversely,  
115 new abdominal and/or back pain, cardiovascular collapse and loss of consciousness  
116 were considered important symptoms indicative of a ruptured aneurysm. It was noted  
117 that the reliability of the aforementioned symptoms would be strengthened when  
118 observed in the presence of one or more significant risk factors.

119 In relation to risk factors, the committee agreed that people with a previously  
120 diagnosed asymptomatic AAA who present with any of the above-mentioned  
121 symptoms were most likely to have a ruptured aneurysm. Age was considered to be  
122 the second most important risk factor. The committee discussed various age  
123 thresholds, noting that the risk of AAA gradually increases with age but there is no  
124 published evidence of a specific age at which the risk of aneurysm rupture increases  
125 markedly. The committee agreed that ruptured AAAs were exceptionally rare in  
126 people younger than 60. Furthermore, they were in agreement that, in their  
127 experience, people over 60 were more likely to present at an A&E department with a  
128 ruptured AAA than people younger than 60. It was noted that the guideline  
129 recommendation aligns with the National AAA Screening Programme which does not  
130 offer screening to men until they reach 65.

131 **Benefits and harms**

132 The committee felt that defining a set of symptoms, signs and risk factors that are  
133 particularly indicative of a ruptured aneurysm, would highlight when clinicians in non-  
134 specialist settings would know when AAA should form part of a differential diagnosis.  
135 This would increase the chances of aneurysms being detected and therefore  
136 increase the likelihood of patient survival.

137 **Cost effectiveness and resource use**

138 The committee felt that the recommendations are unlikely to impact on costs and  
139 resource use, as they were informative in nature.

140 **Other factors the committee took into account**

141 Evidence identified from literature searches for review question 3 (risk factors for  
142 aneurysm growth or rupture in people with known AAA) established that women had  
143 a higher likelihood of aneurysm rupture than men. The committee agreed that  
144 including sex in the list of risk factors may remove focus from the risk of rupture in  
145 men. As a result, it was agreed that a separate recommendation should be made for  
146 women.

147 The committee discussed whether it was necessary to make a recommendation  
148 about imaging of suspected ruptures in non-specialist care settings. The committee  
149 noted that, when they reviewed evidence relating to imaging techniques for  
150 diagnosing unruptured and ruptured AAA, they recommended that an immediate  
151 bedside aortic ultrasound should be offered to people with suspected symptomatic or  
152 ruptured AAA. As a result, they did not think that repeating this recommendation was  
153 necessary. Instead, the committee opted to outline the importance of immediately  
154 discussing suspected ruptures with a vascular surgeon in order obtain specialist  
155 advice and facilitate prompt treatment.  
156

## 157 Appendices

### 158 Appendix A – Review protocols

#### 159 Review protocol for risk factors associated with aneurysm growth or 160 rupture

Review question 15	Which signs, symptoms, risk factors (or combinations of these) and diagnostic assessment tools are most accurate in indicating the presence of a ruptured or symptomatic unruptured abdominal aortic aneurysm i) before arrival at the hospital and ii) in a non-specialist hospital setting?
Objectives	<p>To determine which signs, symptoms and risk factors predict the presence of 'emergency' (that is, ruptured or symptomatic unruptured) abdominal aortic aneurysms</p> <p>To determine which signs, symptoms and risk factors would lead to immediate transfer to a regional vascular service (i.e. bypassing non-specialist A&amp;E)</p> <p>To determine which signs, symptoms and risk factors should initiate definitive imaging in the emergency department (i.e. sufficient suspicion to initiate imaging, but not so high that a patient should be hurried on to vascular unit without imaging)</p>
Type of review	Diagnostic
Language	English
Study design	<p>Systematic reviews of study designs listed below</p> <p>Cross-sectional studies</p>
Status	<p>Published papers only (full text)</p> <p>No date restrictions</p>
Population	People with a suspected ruptured or symptomatic unruptured abdominal aortic aneurysm
Index test / factors of interest	<p>Pulsatile abdominal mass</p> <p>Abdominal, back or groin pain</p> <p>Sweating</p> <p>Clamminess</p> <p>Shock</p> <p>Altered mental/cognitive status</p> <p>Loss of consciousness</p> <p>Collapse</p> <p>Paralysis</p> <p>Dizziness</p> <p>Nausea</p> <p>Vomiting</p> <p>Hypotension</p> <p>Fast pulse / tachycardia</p> <p>Shortness of breath</p> <p>Cyanosis</p> <p>Urinary symptoms</p> <p>Colic</p> <p>Peripheral ischemia / absent peripheral pulses</p> <p>Temperature</p> <p>Capillary refill time</p> <p>Ultrasound ('fast scan')</p>
Endpoint	Surgical or CT confirmation of rupture

<b>Review question 15</b>	<b>Which signs, symptoms, risk factors (or combinations of these) and diagnostic assessment tools are most accurate in indicating the presence of a ruptured or symptomatic unruptured abdominal aortic aneurysm i) before arrival at the hospital and ii) in a non-specialist hospital setting?</b>
	CT confirmation of AAA accompanied by symptoms
Other criteria for inclusion / exclusion of studies	Exclusion: Non-English language Abstract/non-published (i only)
Baseline characteristics to be extracted in evidence tables	Age Sex Size of aneurysm Comorbidities BMI
Search strategies	See Appendix B
Review strategies	Appropriate NICE Methodology Checklists, depending on study designs, will be used as a guide to appraise the quality of individual studies. 20% will be appraised by a second reviewer. Data on all included studies will be extracted into evidence tables. Where statistically possible, a meta-analytic approach will be used to give an overall summary effect. All key findings from evidence will be presented in GRADE profiles and further summarised in evidence statements.
Key papers	None identified

## Appendix B – Literature search strategies

### Clinical search literature search strategy

#### Main searches

Bibliographic databases searched for the guideline

- Cumulative Index to Nursing and Allied Health Literature - CINAHL (EBSCO)
- Cochrane Database of Systematic Reviews – CDSR (Wiley)
- Cochrane Central Register of Controlled Trials – CENTRAL (Wiley)
- Database of Abstracts of Reviews of Effects – DARE (Wiley)
- Health Technology Assessment Database – HTA (Wiley)
- EMBASE (Ovid)
- MEDLINE (Ovid)
- MEDLINE Epub Ahead of Print (Ovid)
- MEDLINE In-Process (Ovid)

#### Identification of evidence for review questions

The searches were conducted between November 2015 and October 2017 for 31 review questions (RQ). In collaboration with Cochrane, the evidence for several review questions was identified by an update of an existing Cochrane review. Review questions in this category are indicated below. Where review questions had a broader scope, supplement searches were undertaken by NICE.

Searches were re-run in December 2017.

Where appropriate, study design filters (either designed in-house or by McMaster) were used to limit the retrieval to, for example, randomised controlled trials. Details of the study design filters used can be found in section 4.

#### Search strategy review question 15

**Medline Strategy, searched 29th September 2016**

**Database: 1946 to September Week 3 2016**

**Search Strategy:**

- 1 Aortic Aneurysm, Abdominal/
- 2 Aortic Rupture/
- 3 (aneurysm\* adj4 (abdom\* or thoracoabdom\* or thoraco-abdom\* or aort\* or spontan\* or juxtarenal\* or juxta-renal\* or juxta renal\* or paraarenal\* or para-renal\* or para renal\* or suprarenal\* or supra renal\* or supra-renal\* or short neck\* or short-neck\* or shortneck\* or visceral aortic segment\*).tw.
- 4 or/1-3
- 5 prognosis.sh.
- 6 diagnosed.tw.
- 7 cohort.mp.
- 8 predictor:.tw.
- 9 death.tw.

**Medline Strategy, searched 29th September 2016**

**Database: 1946 to September Week 3 2016**

**Search Strategy:**

10 exp models, statistical/  
11 or/5-10  
12 (sensitiv: or predictive value:).mp. or accurac:.tw.  
13 11 or 12  
14 "signs and symptoms"/  
15 ((sign or signs) adj5 symptom\*).tw.  
16 Risk Factors/  
17 factor\*.tw.  
18 predict\*.tw.  
19 or/14-18  
20 13 or 19  
21 4 and 20  
22 animals/ not humans/  
23 21 not 22 (12444)  
24 limit 23 to english language

## Health Economics literature search strategy

### Sources searched to identify economic evaluations

- NHS Economic Evaluation Database – NHS EED (Wiley) last updated Dec 2014
- Health Technology Assessment Database – HTA (Wiley) last updated Oct 2016
- Embase (Ovid)
- MEDLINE (Ovid)
- MEDLINE In-Process (Ovid)

Search filters to retrieve economic evaluations and quality of life papers were appended to the population and intervention terms to identify relevant evidence. Searches were not undertaken for qualitative RQs. For social care topic questions additional terms were added. Searches were re-run in September 2017 where the filters were added to the population terms.

### Health economics search strategy

**Medline Strategy**

Economic evaluations

1 Economics/  
2 exp "Costs and Cost Analysis"/  
3 Economics, Dental/  
4 exp Economics, Hospital/  
5 exp Economics, Medical/  
6 Economics, Nursing/  
7 Economics, Pharmaceutical/  
8 Budgets/  
9 exp Models, Economic/  
10 Markov Chains/

## Medline Strategy

- 11 Monte Carlo Method/
- 12 Decision Trees/
- 13 econom\*.tw.
- 14 cba.tw.
- 15 cea.tw.
- 16 cua.tw.
- 17 markov\*.tw.
- 18 (monte adj carlo).tw.
- 19 (decision adj3 (tree\* or analys\*)).tw.
- 20 (cost or costs or costing\* or costly or costed).tw.
- 21 (price\* or pricing\*).tw.
- 22 budget\*.tw.
- 23 expenditure\*.tw.
- 24 (value adj3 (money or monetary)).tw.
- 25 (pharmacoeconomic\* or (pharmac adj economic\*)).tw.
- 26 or/1-25


### Quality of life

- 1 "Quality of Life"/
- 2 quality of life.tw.
- 3 "Value of Life"/
- 4 Quality-Adjusted Life Years/
- 5 quality adjusted life.tw.
- 6 (qaly\* or qald\* or qale\* or qtime\*).tw.
- 7 disability adjusted life.tw.
- 8 daly\*.tw.
- 9 Health Status Indicators/
- 10 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).tw.
- 11 (sf6 or sf 6 or short form 6 or shortform 6 or sf six or sfsix or shortform six or short form six).tw.
- 12 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).tw.
- 13 (sf16 or sf 16 or short form 16 or shortform 16 or sf sixteen or sfsixteen or shortform sixteen or short form sixteen).tw.
- 14 (sf20 or sf 20 or short form 20 or shortform 20 or sf twenty or sftwenty or shortform twenty or short form twenty).tw.
- 15 (euroqol or euro qol or eq5d or eq 5d).tw.
- 16 (qol or hql or hqol or hrqol).tw.
- 17 (hye or hyes).tw.
- 18 health\* year\* equivalent\*.tw.
- 19 utilit\*.tw.
- 20 (hui or hui1 or hui2 or hui3).tw.
- 21 disutili\*.tw.
- 22 rosser.tw.
- 23 quality of wellbeing.tw.
- 24 quality of well-being.tw.
- 25 qwb.tw.


### Medline Strategy

- 26 willingness to pay.tw.
- 27 standard gamble\*.tw.
- 28 time trade off.tw.
- 29 time tradeoff.tw.
- 30 tto.tw.
- 31 or/1-30

## Appendix C – Clinical evidence study selection



## Appendix D – Economic evidence study selection



## Appendix E – Excluded studies

### Clinical studies

No.	Study	Reason for exclusion
1	Iino Misako, Kuribayashi Sachio, Imakita Satoshi, Takamiya Makoto, Matsuo Hiroshi, Ookita Yutaka, Ando Motomi, and Ueda Hatsue (2002) Sensitivity and specificity of CT in the diagnosis of inflammatory abdominal aortic aneurysms. Journal of computer assisted tomography 26(6), 1006-12	The study explored the diagnostic accuracy of CT scans for detecting inflamed abdominal aortic aneurysms. It was not specified which setting (specialist versus non-specialist) the scans were performed.
2	Maybury Rubie Sue, Chang David C, and Freischlag Julie A (2011) Rural hospitals face a higher burden of ruptured abdominal aortic aneurysm and are more likely to transfer patients for emergent repair. Journal of the American College of Surgeons 212(6), 1061-7	Case control evaluating medical records of people with a diagnosis of intact AAA who underwent surgical repair or people with a diagnosis of ruptured AAA upon presentation to a healthcare facility.

### Economic studies

No full text papers were retrieved. All studies were excluded at review of titles and abstracts.

## Appendix F – Glossary

### Abdominal Aortic Aneurysm (AAA)

A localised bulge in the abdominal aorta (the major blood vessel that supplies blood to the lower half of the body including the abdomen, pelvis and lower limbs) caused by weakening of the aortic wall. It is defined as an aortic diameter greater than 3 cm or a diameter more than 50% larger than the normal width of a healthy aorta. The clinical relevance of AAA is that the condition may lead to a life threatening rupture of the affected artery. Abdominal aortic aneurysms are generally characterised by their shape, size and cause:

- **Infrarenal AAA:** an aneurysm located in the lower segment of the abdominal aorta below the kidneys.
- **Juxtarenal AAA:** a type of infrarenal aneurysm that extends to, and sometimes, includes the lower margin of renal artery origins.
- **Suprarenal AAA:** an aneurysm involving the aorta below the diaphragm and above the renal arteries involving some or all of the visceral aortic segment and hence the origins of the renal, superior mesenteric, and celiac arteries, it may extend down to the aortic bifurcation.

### Abdominal compartment syndrome

Abdominal compartment syndrome occurs when the pressure within the abdominal cavity increases above 20 mm Hg (intra-abdominal hypertension). In the context of a ruptured AAA this is due to the mass effect of a volume of blood within or behind the abdominal cavity. The increased abdominal pressure reduces blood flow to abdominal organs and impairs pulmonary, cardiovascular, renal, and gastro-intestinal function. This can cause multiple organ dysfunction and eventually lead to death.

### Cardiopulmonary exercise testing

Cardiopulmonary Exercise Testing (CPET, sometimes also called CPX testing) is a non-invasive approach used to assess how the body performs before and during exercise. During CPET, the patient performs exercise on a stationary bicycle while breathing through a mouthpiece. Each breath is measured to assess the performance of the lungs and cardiovascular system. A heart tracing device (Electrocardiogram) will also record the hearts electrical activity before, during and after exercise.

### Device migration

Migration can occur after device implantation when there is any movement or displacement of a stent-graft from its original position relative to the aorta or renal arteries. The risk of migration increases with time and can result in the loss of device fixation. Device migration may not need further treatment but should be monitored as it can lead to complications such as aneurysm rupture or endoleak.

## Endoleak

An endoleak is the persistence of blood flow outside an endovascular stent - graft but within the aneurysm sac in which the graft is placed.

- Type I – Perigraft (at the proximal or distal seal zones): This form of endoleak is caused by blood flowing into the aneurysm because of an incomplete or ineffective seal at either end of an endograft. The blood flow creates pressure within the sac and significantly increases the risk of sac enlargement and rupture. As a result, Type I endoleaks typically require urgent attention.
- Type II – Retrograde or collateral (mesenteric, lumbar, renal accessory): These endoleaks are the most common type of endoleak. They occur when blood bleeds into the sac from small side branches of the aorta. They are generally considered benign because they are usually at low pressure and tend to resolve spontaneously over time without any need for intervention. Treatment of the endoleak is indicated if the aneurysm sac continues to expand.
- Type III – Midgraft (fabric tear, graft dislocation, graft disintegration): These endoleaks occur when blood flows into the aneurysm sac through defects in the endograft (such as graft fractures, misaligned graft joints and holes in the graft fabric). Similarly to Type I endoleak, a Type III endoleak results in systemic blood pressure within the aneurysm sac that increases the risk of rupture. Therefore, Type III endoleaks typically require urgent attention.
- Type IV– Graft porosity: These endoleaks often occur soon after AAA repair and are associated with the porosity of certain graft materials. They are caused by blood flowing through the graft fabric into the aneurysm sac. They do not usually require treatment and tend to resolve within a few days of graft placement.
- Type V – Endotension: A Type V endoleak is a phenomenon in which there is continued sac expansion without radiographic evidence of a leak site. It is a poorly understood abnormality. One theory that it is caused by pulsation of the graft wall, with transmission of the pulse wave through the aneurysm sac to the native aneurysm wall. Alternatively it may be due to intermittent leaks which are not apparent at imaging. It can be difficult to identify and treat any cause.

## Endovascular aneurysm repair

Endovascular aneurysm repair (EVAR) is a technique that involves placing a stent –graft prosthesis within an aneurysm. The stent-graft is inserted through a small incision in the femoral artery in the groin, then delivered to the site of the aneurysm using catheters and guidewires and placed in position under X-ray guidance.

- Conventional EVAR refers to placement of an endovascular stent graft in an AAA where the anatomy of the aneurysm is such that the ‘instructions for use’ of that particular device are adhered to. Instructions for use define tolerances for AAA anatomy that the device manufacturer considers appropriate for that device. Common limitations on AAA anatomy are infrarenal neck length (usually >10mm), diameter (usually ≤30mm) and neck angle relative to the main body of the AAA
- Complex EVAR refers to a number of endovascular strategies that have been developed to address the challenges of aortic proximal neck fixation associated with complicated aneurysm anatomies like those seen in juxtarenal and suprarenal AAAs.

These strategies include using conventional infrarenal aortic stent grafts outside their 'instructions for use', using physician-modified endografts, utilisation of customised fenestrated endografts, and employing snorkel or chimney approaches with parallel covered stents.

### **Goal directed therapy**

Goal directed therapy refers to a method of fluid administration that relies on minimally invasive cardiac output monitoring to tailor fluid administration to a maximal cardiac output or other reliable markers of cardiac function such as stroke volume variation or pulse pressure variation.

### **Post processing technique**

For the purpose of this review, a post-processing technique refers to a software package that is used to augment imaging obtained from CT scans, (which are conventionally presented as axial images), to provide additional 2- or 3-dimensional imaging and data relating to an aneurysm's, size, position and anatomy.

### **Permissive hypotension**

Permissive hypotension (also known as hypotensive resuscitation and restrictive volume resuscitation) is a method of fluid administration commonly used in people with haemorrhage after trauma. The basic principle of the technique is to maintain haemostasis (the stopping of blood flow) by keeping a person's blood pressure within a lower than normal range. In theory, a lower blood pressure means that blood loss will be slower, and more easily controlled by the pressure of internal self-tamponade and clot formation.

### **Remote ischemic preconditioning**

Remote ischemic preconditioning is a procedure that aims to reduce damage (ischaemic injury) that may occur from a restriction in the blood supply to tissues during surgery. The technique aims to trigger the body's natural protective functions. It is sometimes performed before surgery and involves repeated, temporary cessation of blood flow to a limb to create ischemia (lack of oxygen and glucose) in the tissue. In theory, this "conditioning" activates physiological pathways that render the heart muscle resistant to subsequent prolonged periods of ischaemia.

### **Tranexamic acid**

Tranexamic acid is an antifibrinolytic agent (medication that promotes blood clotting) that can be used to prevent, stop or reduce unwanted bleeding. It is often used to reduce the need for blood transfusion in adults having surgery, in trauma and in massive obstetric haemorrhage.