

Coil embolisation of ruptured intracranial aneurysms

1 Guidance

- 1.1 Current evidence on the safety and efficacy of coil embolisation of ruptured intracranial aneurysms appears adequate to support use of the procedure, provided that normal arrangements are in place for consent, audit and clinical governance.
- 1.2 The procedure should only be performed in specialist units with expertise in the endovascular treatment of intracranial aneurysms. Clear arrangements should be in place for the involvement of different clinical disciplines in treatment and follow-up.
- 1.3 Patients with subarachnoid haemorrhage should have rapid access to appropriate specialist care.

2 The procedure

2.1 Indications

- 2.1.1 Intracranial aneurysms are small balloon-like dilated portions of blood vessels that may occasionally rupture, causing haemorrhage, stroke or death. Usually the cause is unknown but people with genetic causes of weak blood vessels are more likely to develop aneurysms. Rupture of intracranial aneurysms causes subarachnoid haemorrhage and has a poor prognosis. About 30% of people die within 24 hours and a further 25–30% die within 4 weeks.
- 2.1.2 The alternative treatment for ruptured intracranial aneurysm involves open surgery to clip the aneurysm inside the skull.

2.2 Outline of the procedure

- 2.2.1 The coil technique involves approaching the aneurysm from inside the diseased blood vessel, thereby avoiding the need to open the skull. A thin tube containing the coil on a guidewire is inserted into a large artery, usually in the groin, and passed up into the skull under radiological guidance. The coil is placed inside the aneurysm and detached from the guidewire. Once in position, it causes clotting and stops blood from entering the aneurysm. Multiple coils may be inserted into the aneurysm through the same tube until the aneurysm is filled with coils.

2.3 Efficacy

- 2.3.1 Use of the procedure was supported by a high-quality randomised controlled trial. In the trial, 'dependency' reflected a moderate to severe disability as defined by the modified Rankin score. The trial showed a 7% absolute risk reduction in dependency or death for patients treated with coils compared with patients treated by surgical clipping. For more details, refer to the Sources of evidence.
- 2.3.2 The Specialist Advisors stated that, in the short term, the procedure is more effective clinically than surgical clipping. However, the long-term durability of coil embolisation has not been established.

Interventional Procedure Guidance 106

This guidance is written in the following context:

This guidance represents the view of the Institute which was arrived at after careful consideration of the available evidence. Health professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of health professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

2.4 Safety

- 2.4.1 Complications associated with the procedure included perforation of the aneurysm, intracranial haematoma and re-bleeding. In a case series of 403 patients, aneurysm perforation was observed in 11 patients (3%) and cerebral clot embolisation in 10 patients (2%). Coil migration occurred in two patients (0.5%). For more details, refer to the Sources of evidence.
- 2.4.2 The Specialist Advisors considered this procedure to be safer than surgical clipping. They stated that procedural mortality and stroke were the main adverse events. They also stated that there is a small risk of re-bleeding, and that this should be monitored over the long term.

2.5 Other comments

- 2.5.1 Currently, evidence on the procedure's long-term results is limited to a mean follow-up of 3.7 years.

3 Further information

- 3.1 This guidance relates to ruptured intracranial aneurysms. The Institute has also published separate guidance on the use of coil embolisation for unruptured intracranial aneurysms (www.nice.org.uk/IPG105guidance).
- 3.2 In 2002, the Royal College of Physicians published guidelines on the management of stroke, including subarachnoid haemorrhage (www.rcplondon.ac.uk).

Andrew Dillon
Chief Executive
January 2005

Information for the public

NICE has produced information describing its guidance on this procedure for patients, carers and those with a wider interest in healthcare. It explains the nature of the procedure and the decision made, and has been written with patient consent in mind. This information is available, in English and Welsh, from www.nice.org.uk/IPG106publicinfo

Sources of evidence

The evidence considered by the Interventional Procedures Advisory Committee is described in the following document.

Interventional procedure overview of endovascular treatment of intracranial aneurysms, December 2002.

Available from: www.nice.org.uk/IP038overview

Ordering information

Copies of this guidance can be obtained from the NHS Response Line by telephoning 0870 1555 455 and quoting reference number N0799. *Information for the public* can be obtained by quoting reference number N0800 for the English version and N0801 for a version in English and Welsh.

The distribution list for this guidance is available on the NICE website at www.nice.org.uk/IPG106distributionlist

Published by the National Institute for Clinical Excellence, January 2005 ISBN: 1-84257-858-8

© National Institute for Clinical Excellence, January 2005. All rights reserved. This material may be freely reproduced for educational and not for profit purposes within the NHS. No reproduction by or for commercial organisations is permitted without the express written permission of the Institute.

National Institute for Clinical Excellence

MidCity Place, 71 High Holborn, London WC1V 6NA, website: www.nice.org.uk

N0799 1P 20k Jan 05 (ABA)