

**New indicators to be added to the NICE
indicator menu for
clinical commissioning groups**

1st August 2017

Summary tables for the NICE menu

This paper provides the latest set of indicators NICE has published for inclusion on the NICE indicator menu for CCGs.

For information the full NICE indicator menu for CCGs is also provided in table 2.

NHS England and other organisations can use the NICE menu to identify suitable indicators for national and local indicator frameworks. The indicators have been tested for use at CCG level; they could also be used to support measurement across the sustainability and transformation plan footprints.

The paper consists of the following sections:

- newly developed indicators for inclusion in the NICE menu (table 1)
- the full NICE indicator menu of CCG indicators (table 2)

The full NICE indicator menu and the associated supporting documentation will be available via the [NICE website](#) following publication on 1st August 2017.

Table 1: New CCG indicators recommended for the NICE indicator menu

10 weeks antenatal booking appointment			
Indicator ID	Indicator wording	Evidence source	Why is this indicator important to measure?
CCG81	Proportion of pregnant women accessing antenatal care who are seen for booking by 10 weeks and 0 days	NICE guideline on antenatal care 1.1.1.1 NICE QS22 – antenatal care – statement 1	Pregnant women accessing antenatal care should be seen early in their pregnancy in order to determine if there is a need for additional care. NICE guidance recommends 3 audit targets during gestation: 10 weeks, 12 weeks + 6 days (13 weeks) and 20 weeks of pregnancy

Antenatal and postnatal mental health			
Indicator ID	Indicator wording	Evidence source	Why is this indicator important to measure?
CCG82	The proportion of pregnant women who were asked about their mental health at their first booking appointment	NICE guideline on Antenatal and postnatal mental health . 1.5.4 and 1.5.8 NICE QS115 – antenatal and postnatal mental health – statement 4	The first booking appointment allows healthcare professionals to discuss emotional wellbeing with women and identify potential mental health problems. This will help health professionals provide appropriate support. The management of mental health problems during pregnancy and the postnatal period differs from at other times because of the nature of this life stage and the potential impact of any difficulties and treatments on the woman and the baby

Contents

Atrial fibrillation	5
Cancer	5
Chronic obstructive pulmonary disorder	5
Dementia	6
Depression and anxiety	6
Diabetes	6
Embolism and thrombosis	9
Gynaecological conditions	10
Heart failure	10
Hip fracture	10
Pregnancy and neonates	11
Smoking	13
Stroke and ischaemic attack	13
Surgical care	14

Atrial fibrillation

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG55	Proportion of patients with atrial fibrillation on anticoagulation admitted to hospital for stroke	Overarching outcome measure	Atrial fibrillation QS93	NICE menu
CCG56	Proportion of patients with atrial fibrillation not on anticoagulation admitted to hospital for stroke	Outcome measure (statement 1)	Atrial fibrillation QS93: statement 1	NICE menu

Cancer

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG01	The proportion of invasive cases of cancer diagnosed via an emergency route	Overarching outcome measure		In CCG OIS
CCG02	The proportion of new cases of cancer with a stage of diagnosis recorded	Overarching outcome measure		In 2016/17 CCG IAF In CCG OIS
CCG03	The proportion of people with cancer who have the cancer detected at stage 1 or 2	Overarching outcome measure		In CCG OIS
CCG04	Breast cancer mortality rates	Overarching outcome measure		In CCG OIS
CCG06	The proportion of lung cancer cases for which a stage at diagnosis is recorded	Overarching outcome measure		In CCG OIS

Chronic obstructive pulmonary disorder

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG08	The proportion of people with COPD and MRC Dyspnoea Scale 3 and above referred to a pulmonary rehabilitation programme	NICE guideline CG101 : rec 1.2.8.2	COPD in adults QS10: statement 4	In CCG OIS

Dementia

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG18	Proportion of people with dementia prescribed anti-psychotic medication	NICE guideline CG42 recs 1.7.2.2, 1.7.2.3	Dementia: support in health and social care QS1: Statement 7	In CCG OIS

Depression and anxiety

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG41	The proportion of people of all ages with mental health problems who clinically recover following talking therapies	Overarching outcome measure	Anxiety disorders QS53	In CCG OIS
CCG54	Proportion of people treated by IAPT for anxiety disorders who return to full function	Overarching outcome measure	Anxiety disorders QS53	NICE menu

Diabetes

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG12	The proportion of adults with a diagnosis of diabetes who have a recording of MI, stroke and end stage kidney disease.	Overarching outcome measure		In CCG OIS
CCG13	The proportion of adults with diabetes who have received all nine basic care processes performed (see guidance for the details of the process)	NICE guideline NG28 recs 1.8–1.14 NICE guideline NG17 recs 1.2.5, 1.13, 1.15		In CCG OIS

CCG14	The proportion of adults with diabetes referred to a structured education programme within 12 months of diagnosis	NICE guideline NG28 rec 1.1.1 NICE guideline NG17 recs 1.3.1, 1.3.8 NICE guideline NG19 rec 1.3.13	Diabetes in adults QS6: statements 2 and 3	In 2016/17 CCG IAF In CCG OIS
CCG15	The proportion of adults with diabetes who have an emergency admissions for diabetic ketoacidosis.	Overarching outcome measure		In CCG OIS
CCG16	Proportion of adults with diabetes with a HES recording of National Diabetes Audit complications	Overarching outcome measure		In CCG OIS
CCG58	Admission rates for people with diabetes due to complications associated with diabetes	NICE guideline NG28 NICE guideline NG17		NICE menu
CCG59	Admissions rates due to angina in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 rec 1.13.7		NICE menu
CCG60	Admission rates due to myocardial infarction in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 rec 1.13.7		NICE menu
CCG61	Admission rates due to heart failure in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 rec 1.13.7		NICE menu

CCG62	Admission rates due to stroke in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 rec 1.13.7		NICE menu
CCG63	Admissions rates for renal replacement therapy in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 rec 1.3.13		NICE menu
CCG64	Admission rates due to diabetic ketoacidosis in people with diabetes	NICE guideline NG28 rec 1.5.2 NICE guideline NG17 recs 1.15.10 - 1.15.18		NICE menu
CCG65	Admission rates due to lower limb amputations in people with diabetes	NICE guideline NG19		NICE menu
CCG66	Proportion of children and young people with diabetes who receive the following individual care processes in the past 12 months: <ul style="list-style-type: none"> • Glycated Haemoglobin A1c (HbA1c) monitoring • Body Mass Index (BMI) • Blood pressure • Urinary Albumin • Eye screening • Foot examination • Smoking • Screening for thyroid disease • Psychological assessment 	NICE guideline NG18 : recs 1.2.4, 1.2.12-14, 1.2.45, 1.2.71, 1.2.95, 1.2.98, 1.2.110, 1.3.3, 1.3.8-10, 1.3.20, 1.3.28, 1.3.34, 1.3.37, 1.3.43 NICE guideline NG19 : recs 1.3.2, 1.3.11 NICE guideline NG20 : rec 1.1.1		NICE menu
CCG67	Proportion of children and young people with diabetes who have had their glycated haemoglobin A1c (HbA1c) monitored in the previous 12 months	NICE guideline NG18 : recs 1.2.71, 1.3.28		NICE menu

CCG68	Proportion of children and young people with diabetes who have had their Body Mass Index (BMI) recorded	NICE guideline NG18 : recs 1.2.45, 1.3.20		NICE menu
CCG69	Proportion of young people aged 12-18 years with diabetes who have had their blood pressure recorded in the previous 12 months	NICE guideline NG18 : recs 1.2.110, 1.3.43		NICE menu
CCG70	Proportion of young people aged 12-18 years with diabetes who have had their urinary albumin recorded in the previous 12 months	NICE guideline NG18 : recs 1.2.110, 1.3.43		NICE menu
CCG72	Proportion of young people aged 12-18 years with diabetes who have a record of eye screening in the previous 12 months	NICE guideline NG18 : recs 1.2.4, 1.3.3		NICE menu
CCG73	Proportion of children and young people with diabetes who have a record of a foot examination in the previous 12 months	NICE guideline NG19 : recs 1.3.2, 1.3.11		NICE menu
CCG74	Proportion of young people age 12 -18 who have had their smoking status recorded in the previous 12 months	NICE guideline NG18 : recs 1.2.12-14, 1.3.8-10		NICE menu
CCG75	Proportion of children and young people with type 1 diabetes who have been screened for thyroid disease	NICE guideline NG18 : rec 1.2.110		NICE menu
CCG76	Proportion of children and young people with diabetes who have received a psychological assessment in the previous 12 months.	NICE guideline NG18 recs 1.2.95, 1.2.98, 1.3.34, 1.3.37	Diabetes in children and young people QS125: statement 6	NICE menu

Embolism and thrombosis

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG80	Mortality rates directly associated with VTE	Overarching outcome measure.	Venous thromboembolism in adults: diagnosis and management QS29	NICE menu

Gynaecological conditions

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG77	Rates of hysterectomy	NICE guideline CG44 recs 1.7.3, 1.7.5, 1.7.6	Heavy menstrual bleeding QS47: statement 6	NICE menu
CCG78	Rates of endometrial ablation	NICE guideline CG44 recs 1.6.4, 1.6.5	Heavy menstrual bleeding QS47: statement 5	NICE menu

Heart failure

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG10	The proportion of people with coronary heart disease referred for cardiac rehabilitation	NICE guideline CG108 rec 1.3.1.1	Chronic heart failure in adults QS9: statement 6	In CCG OIS
CCG11	All cause mortality - 12 months following admission to hospital for heart failure	Outcome measure	Chronic heart failure in adults QS9	In CCG OIS

Hip fracture

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG20	The proportion of people with hip fracture, who receive a formal hip fracture programme from admission	NICE guideline CG124 : recs 1.8.1, 1.8.5, 1.8.6	Hip fracture in adults QS16: statement 1	In CCG OIS
CCG21	The proportion of people with hip fracture, who receive surgery on the day of, or the day after, admission	NICE guideline CG124 : recs 1.2.1, 1.2.2	Hip fracture in adults QS16: statement 2	In CCG OIS

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG22	The proportion of people with hip fracture, who receive a multifactorial risk assessment of future falls risk	NICE guideline CG161 : recs 1.1.2.1, 1.1.2.2	Falls in older people QS86: statements 2 and 4	In CCG OIS
CCG23	The proportion of people in the national hip fracture database who have received all nine care processes performed (see guidance for the details of the process)	NICE guideline CG124 : rec 1.8.1		In CCG OIS
CCG24	The rate of people admitted with a primary diagnosis of hip fracture per 100,000 CCG population	Outcome measure		In CCG OIS

Pregnancy and neonates

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG30	Proportion of women who receive antenatal assessments by 13 weeks of pregnancy	NICE guideline CG62 : rec 1.1.1.1	Antenatal care QS22: statement 1	In CCG OIS
CCG31	Proportion of pregnant women who were smokers during pregnancy	NICE guideline PH26 : rec 1	Antenatal care QS22: statement 5	NICE menu
CCG32	Proportion of pregnant women who were smokers at the time of delivery	NICE guideline PH26 : rec 1	Antenatal care QS22: statement 5	In 2016/17 CCG IAF In CCG OIS
CCG33	Proportion of successful births where breast feeding is initiated	NICE guideline CG62 : recs 1.1.1.1, 1.1.1.6 NICE guideline PH11 : recs 1, 7	Postnatal care QS37: statement 5	NICE menu

CCG34	The proportion of pregnancies resulting in a neonatal or still birth	Overarching outcome measure	Antenatal care QS22 Caesarean section QS32 Multiple pregnancy: twin and triplet pregnancies QS46	In 2016/17 CCG IAF In CCG OIS
CCG35	The proportion of full term births where the child has a low birth weight	Overarching outcome measure	Antenatal care QS22 Caesarean section QS32 Maternal and child nutrition QS98	In CCG OIS
CCG36	The proportion of births resulting in a neonatal unit admission	Overarching outcome measure	Antenatal care QS22 Caesarean section QS32 Multiple pregnancy: twin and triplet pregnancies QS46	In CCG OIS
CCG37	The proportion of pregnant women having a planned caesarean section who have the procedure carried out at or after 39 weeks 0 days	NICE guideline CG132 : rec 1.4.1.1	Caesarean section QS32: statement 5	In CCG OIS
CCG38	Rates of exclusive or partial breastfeeding at 6-8 weeks after the birth	NICE guideline CG37 : rec 1.3.3 NICE guideline PH11 : rec 7	Postnatal care QS37: statement 5	In CCG OIS
CCG81	Proportion of pregnant women accessing antenatal care who are seen for booking by 10 weeks and 0 days	NICE guideline on antenatal care 1.1.1.1	Antenatal care QS22: statement 1	New NICE menu indicator

CCG82	The proportion of pregnant women who were asked about their mental health at their first booking appointment	NICE guideline on Antenatal and postnatal mental health . 1.5.4 and 1.5.8	Antenatal and postnatal mental health QS115 – statement 4	New NICE menu indicator
-------	--	---	---	-------------------------

Smoking

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG44	The proportion of people with severe mental illness who are recorded as current smokers	NICE guideline PH48 : rec 2		In CCG OIS

Stroke and ischaemic attack

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG45	Mortality within 30 days of hospital admission for stroke	Overarching outcome measure.		In CCG OIS
CCG46	Proportion of people who receive psychological support for mood, behaviour or cognitive disturbance 6 months after an admission to hospital with a stroke	NICE guideline CG162 : rec 1.1.3	Stroke in adults QS2: statement 3	In CCG OIS
CCG47	Proportion of people with stroke who receive joint health and social care plans on discharge from hospital	NICE guideline CG162 : rec 1.11.15	Stroke in adults QS2: statements 4 and 7	In CCG OIS
CCG48	Proportion of people who had a stroke who are reviewed within 6 months of being discharged from hospital	NICE guideline CG162 : rec 1.11.15	Stroke in adults QS2: statement 7	In CCG OIS

CCG49	The proportion of people who had a stroke that are supported by a stroke skilled early supported discharge (ESD) team	NICE guideline CG162 : rec 1.1.8	Stroke in adults QS2: statement 4	In CCG OIS
CCG50	Proportion of people who have had an acute stroke who receive thrombolysis for stroke	NICE guideline CG68 : rec. 1.4.1.1		In CCG OIS
CCG51	Proportion of people who have had or are having a stroke who are admitted to an acute stroke unit within 4 hours of arrival to hospital	NICE guideline CG68 : rec 1.3.1.1	Stroke in adults QS2: statement 1	In CCG OIS
CCG52	Proportion of patients who have had an acute stroke who spend 90% or more of their stay on a stroke unit	NICE guideline CG68 : rec 1.3.1.1	Stroke in adults QS2: statement 1	In CCG OIS
CCG53	The proportion of people who have had an acute stroke whose swallowing is screened by a specially trained healthcare professional within 4 hours of admission to hospital	RCP Stroke guideline 5th edition (2016) 3.10.1E		NICE menu

Surgical care

NICE menu ID	Indicator	Evidence base	NICE quality standard	Current status
CCG79	Readmission rates for surgical site infections within 30 days of discharge from surgery	Outcome measure	Surgical site infection QS49: statement 7	NICE menu