

Cystic Fibrosis

Diagnosis and management

NICE Guideline

Methods, evidence and recommendations

04 May 2017

Draft for Consultation

*Developed by the National Guideline Alliance, hosted
by the Royal College of Obstetricians and
Gynaecologists*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the Welsh Government, Scottish Government, and Northern Ireland Executive. All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© National Institute for Health and Care Excellence 2017. All rights reserved.

Contents

1	Introduction	14
2	Guideline summary	15
2.1	Guideline Committee membership, NGA staff and acknowledgements	15
2.2	Other versions of the guideline	16
2.3	Schedule for updating the guideline.....	16
3	Development of this guideline	17
3.1	What is a NICE Guideline?	17
3.2	Remit.....	17
3.3	Who developed this guideline?	18
3.4	What this guideline covers.....	18
3.4.1	Groups that will be covered.....	18
3.4.2	Key clinical issues that will be covered.....	18
3.5	What this guideline does not cover	19
3.5.1	Clinical issues that will not be covered	19
3.6	Relationship between the guideline and other NICE guidance.....	19
3.6.1	Related NICE guidance.....	19
4	Guideline development methodology	20
4.1	Developing the review questions and outcomes	20
4.2	Searching for evidence.....	28
4.2.1	Health economic literature search.....	29
4.3	Reviewing and synthesising research evidence	30
4.3.1	Systematic review process.....	30
4.3.2	Inclusion/exclusion criteria	31
4.3.3	Type of studies.....	31
4.3.4	Types of data and methods for synthesis	32
4.3.5	Appraising the quality of the evidence by outcomes	38
4.3.6	Evidence statements.....	50
4.3.7	Evidence of cost effectiveness	50
4.4	Developing recommendations	51
4.4.1	Guideline recommendations.....	51
4.4.2	Research recommendations	52
4.5	Validation process	52
4.6	Updating the guideline.....	52
4.7	Disclaimer	52
4.8	Funding	52
5	Diagnosis of cystic fibrosis	53
5.1	Introduction	53
5.2	Description of clinical evidence.....	53

5.2.1	Respiratory symptoms	54
5.2.2	Faltering growth	54
5.2.3	Symptoms of malabsorption.....	54
5.2.4	Azoospermia.....	54
5.2.5	Acute pancreatitis	54
5.2.6	Meconium ileus	54
5.3	Summary of included studies and results	54
5.4	Clinical evidence profile.....	60
5.5	Economic evidence	60
5.6	Evidence statements	60
5.6.1	Respiratory symptoms	60
5.6.2	Faltering growth	61
5.6.3	Symptoms of malabsorption.....	61
5.6.4	Azoospermia.....	61
5.6.5	Acute pancreatitis	61
5.6.6	Meconium ileus.....	61
5.6.7	Economic evidence statements.....	61
5.7	Evidence to recommendations	62
5.7.1	Consideration of clinical benefits and harms	62
	These Committee noted that these recommendations are consistent with the NHS Service Specifications for cystic fibrosis.	64
5.7.2	Consideration of economic benefits and harms.....	64
5.7.3	Quality of evidence	64
5.7.4	Other considerations.....	64
5.7.5	Key conclusions	65
5.8	Recommendations	65
6	Information and Support.....	67
6.1	Introduction	67
6.2	Description of clinical evidence.....	67
6.3	Summary of included studies.....	68
6.4	Clinical evidence profile.....	83
6.4.1	Theme maps.....	83
6.4.2	Clinical evidence profile: information needs for people with cystic fibrosis and their parents or carers	85
6.4.3	Clinical evidence profile: support needs for people with cystic fibrosis and their parents or carers	104
6.5	Economic evidence	124
6.6	Evidence statements	124
6.6.1	Information needs for people with cystic fibrosis and their parents or carers.....	124
6.6.2	Support needs for people with cystic fibrosis and their parents or carers	125

6.6.3	Economic evidence statements.....	126
6.7	Evidence to recommendations	126
6.7.1	Relative value placed on the themes considered	126
6.7.2	Consideration of clinical benefits and harms	126
6.7.3	Consideration of economic benefits and harms.....	127
6.7.4	Quality of evidence	128
6.7.5	Other considerations.....	128
6.7.6	Key conclusions	128
6.8	Recommendations	129
7	Service delivery	131
7.1	Service configuration.....	131
7.1.1	Introduction	131
7.1.2	Description of clinical evidence	131
7.1.3	Summary of included studies	133
7.1.4	Clinical evidence profile	140
7.1.5	Economic evidence.....	154
7.1.6	Evidence statements.....	155
7.1.7	Evidence to recommendations	164
7.1.8	Recommendations	170
7.2	Multidisciplinary team	171
7.2.1	Introduction	171
7.2.2	Description of the clinical evidence	171
7.2.3	Summary of included studies	172
7.2.4	Clinical evidence profile	172
7.2.5	Economic evidence.....	172
7.2.6	Evidence statements.....	173
7.2.7	Evidence to recommendations	173
7.2.8	Recommendations	176
7.3	Transition to adult services.....	177
7.3.1	Introduction	177
7.3.2	Description of clinical evidence	178
7.3.3	Summary of included studies	179
7.3.4	Clinical evidence	182
7.3.5	Clinical evidence profile	184
7.3.6	Economic evidence.....	202
7.3.7	Evidence statements.....	202
7.3.8	Evidence to recommendations	203
7.3.9	Recommendations	206
8	Complications of cystic fibrosis.....	208
8.1	Introduction	208

8.2	Malnutrition.....	208
8.2.1	Description of clinical evidence: malnutrition	208
8.2.2	Summary of included studies and results: malnutrition.....	209
8.2.3	Evidence statements: malnutrition.....	212
8.3	Impaired growth.....	213
8.3.1	Description of clinical evidence: impaired growth	213
8.3.2	Summary of included studies and results: impaired growth.....	213
8.3.3	Evidence statements: impaired growth.....	219
8.4	Cystic fibrosis related renal disease	221
8.4.1	Description of clinical evidence: cystic fibrosis related renal disease.....	221
8.4.2	Summary of included studies and results: cystic fibrosis related renal disease	221
8.4.3	Evidence statements: cystic fibrosis related renal disease	223
8.5	Delayed puberty	224
8.5.1	Description of clinical evidence: delayed puberty	224
8.5.2	Summary of included studies and results: delayed puberty.....	224
8.5.3	Evidence statements: delayed puberty.....	225
8.6	Abdominal pain	225
8.6.1	Description of clinical evidence: abdominal pain	225
8.6.2	Summary of included studies and results: abdominal pain	225
8.6.3	Evidence statements: abdominal pain.....	225
8.7	Cystic Fibrosis related diabetes.....	225
8.7.1	Description of clinical evidence: cystic fibrosis related diabetes	225
8.7.2	Summary of included studies and results: cystic fibrosis related diabetes	226
8.7.3	Evidence statements: cystic fibrosis related diabetes.....	228
8.8	Upper airways disease	228
8.8.1	Description of clinical evidence: upper airways disease	228
8.8.2	Summary of included studies and results: upper airways disease	229
8.8.3	Evidence statements: Upper airways disease	229
8.9	Cystic fibrosis related musculoskeletal disorders.....	229
8.9.1	Description of clinical evidence: cystic fibrosis related musculoskeletal disorders	229
8.9.2	Summary of included studies and results: Cystic fibrosis related musculoskeletal disorders.....	230
8.9.3	Evidence statements: cystic fibrosis related musculoskeletal disorders ..	231
8.10	Urinary stress incontinence	231
8.10.1	Description of clinical evidence: urinary stress incontinence	231
8.10.2	Summary of included studies and results: urinary stress incontinence ...	231
8.10.3	Evidence statements: urinary stress incontinence.....	231
8.11	Reduced bone mineral density	232

8.11.1	Description of clinical evidence: reduced bone mineral density	232
8.11.2	Summary of included studies and results: reduced bone mineral density	232
8.11.3	Evidence statements: reduced bone mineral density	234
8.12	Cystic fibrosis related liver disease.....	234
8.12.1	Description of clinical evidence: cystic fibrosis related liver disease	234
8.12.2	Summary of included studies and results: cystic fibrosis related liver disease	235
8.12.3	Evidence statements: cystic fibrosis related liver disease	235
8.13	Infertility.....	236
8.13.1	Description of clinical evidence: infertility	236
8.13.2	Summary of included studies and results: infertility	236
8.13.3	Evidence statements: infertility.....	236
8.14	Distal Intestinal Obstruction Syndrome (DIOS).....	236
8.14.1	Description of clinical evidence: Distal Intestinal Obstruction Syndrome (DIOS)	236
8.14.2	Summary of included studies and results: Distal Intestinal Obstruction Syndrome (DIOS)	237
8.14.3	Evidence statements: Distal Intestinal Obstruction Syndrome (DIOS)....	238
8.15	Meconium ileus	239
8.15.1	Description of clinical evidence: meconium ileus.....	239
8.15.2	Summary of included studies and results: meconium ileus	239
8.15.3	Evidence statements: meconium ileus	239
8.16	Economic evidence	240
8.17	Evidence to recommendations	240
8.17.1	Relative value placed on the outcomes considered.....	240
8.17.2	Consideration of clinical benefits and harms	240
8.17.3	Consideration of economic benefits and harms.....	244
8.17.4	Quality of evidence	244
8.17.5	Other considerations.....	244
8.17.6	Key conclusions	244
8.18	Recommendations	245
9	Pulmonary monitoring, assessment and management	246
9.1	Pulmonary monitoring	246
9.1.1	Introduction	246
9.1.2	Description of clinical evidence	246
9.1.3	Summary of included studies	248
9.1.4	Clinical evidence profiles.....	249
9.1.5	Economic evidence.....	252
9.1.6	Evidence statements.....	253
9.1.7	Evidence to recommendations	255

9.1.8	Recommendations	260
9.1.9	Research recommendations	262
9.2	Airway clearance techniques	263
9.2.1	Introduction	263
9.2.2	Description of clinical evidence	263
9.2.3	Summary of included studies	265
9.2.4	Clinical evidence profile	272
9.2.5	Economic evidence	281
9.2.6	Evidence statements.....	281
9.2.7	Evidence to recommendations	288
9.2.8	Recommendations	291
9.2.9	Research recommendations	292
9.3	Mucoactive agents	294
9.3.1	Introduction	294
9.3.2	Description of clinical evidence	294
9.3.3	Summary of included studies	296
9.3.4	Clinical evidence profile	304
9.3.5	Economic evidence	322
9.3.6	Evidence statements.....	322
9.3.7	Evidence to recommendations	331
9.3.8	Recommendations	334
9.3.9	Research recommendations	335
9.4	Pulmonary Infection.....	336
9.4.1	Prophylaxis	337
9.4.2	Acute	351
9.4.3	Chronic	387
9.4.4	Recommendations	456
9.5	Immunomodulatory agents	461
9.5.1	Introduction	461
9.5.2	Description of clinical evidence	461
9.5.3	Summary of included studies	462
9.5.4	Clinical evidence profile	466
9.5.5	Economic evidence	481
9.5.6	Evidence statements.....	482
9.5.7	Evidence to recommendations	486
9.5.8	Recommendations	489
10	Other monitoring, assessment and management	490
10.1	Nutritional Interventions.....	490
10.1.1	Introduction	490
10.1.2	Description of clinical evidence	490

10.1.3	Summary of included studies	491
10.1.4	Clinical evidence profile	498
10.1.5	Economic evidence	520
10.1.6	Evidence statements	520
10.1.7	Evidence to recommendations	529
10.1.8	Recommendations	533
10.2	Exocrine pancreatic insufficiency	533
10.2.1	Introduction	533
10.2.2	Description of clinical evidence	534
10.2.3	Summary of included studies	535
10.2.4	Clinical evidence profile	537
10.2.5	Economic evidence	543
10.2.6	Evidence statements	544
10.2.7	Evidence to recommendations	547
10.2.8	Recommendations	550
10.3	Distal ileal obstruction syndrome	550
10.3.1	Introduction	550
10.3.2	Description of clinical evidence	550
10.3.3	Summary of included studies	551
10.3.4	Clinical evidence profile	551
10.3.5	Economic evidence	551
10.3.6	Evidence statements	551
10.3.7	Evidence to recommendations	551
10.3.8	Recommendations	554
10.4	Monitoring for liver disease	555
10.4.1	Introduction	555
10.4.2	Description of clinical evidence	556
10.4.3	Summary of included studies	557
10.4.4	Clinical evidence profile	565
10.4.5	Economic evidence	565
10.4.6	Evidence statements	565
10.4.7	Evidence to recommendations	577
10.4.8	Recommendations	581
10.5	Ursodeoxycholic acid	581
10.5.1	Introduction	581
10.5.2	Description of clinical evidence	581
10.5.3	Summary of included studies	582
10.5.4	Clinical evidence profile	583
10.5.5	Economic evidence	585
10.5.6	Evidence statements	585

10.5.7	Evidence to recommendations	586
10.5.8	Recommendations	589
10.5.9	Research recommendations	589
10.6	Monitoring for cystic fibrosis related diabetes	590
10.6.1	Introduction	590
10.6.2	Description of clinical evidence	590
10.6.3	Summary of included studies	591
10.6.4	Clinical evidence profile	591
10.6.5	Economic evidence	591
10.6.6	Evidence statements.....	591
10.6.7	Evidence to recommendations	591
10.6.8	Recommendations	594
10.6.9	Research recommendations	594
10.7	Monitoring for low bone mineral density	596
10.7.1	Introduction	596
10.7.2	Description of clinical evidence	596
10.7.3	Summary of included studies	597
10.7.4	Clinical evidence profile	599
10.7.5	Economic evidence	602
10.7.6	Evidence statements.....	603
10.7.7	Evidence to recommendations	605
10.7.8	Recommendations	607
10.8	Exercise	608
10.8.1	Introduction	608
10.8.2	Description of clinical evidence	608
10.8.3	Summary of included studies	610
10.8.4	Clinical evidence profile	618
10.8.5	Economic evidence	642
10.8.6	Evidence statements.....	642
10.8.7	Evidence to recommendations	654
10.8.8	Recommendations	657
10.9	Psychological assessment	657
10.9.1	Introduction	657
10.9.2	Description of clinical evidence	658
10.9.3	Summary of included studies and results	661
10.9.4	Clinical evidence profile	664
10.9.5	Economic evidence	664
10.9.6	Evidence statements.....	664
10.9.7	Evidence to recommendations	666
10.9.8	Recommendations	669

10.9.9 Research recommendations	670
11 Prevention of cross infection	672
11.1 Introduction	672
11.2 Description of clinical evidence.....	672
11.3 Summary of included studies.....	674
11.4 Clinical evidence profile.....	683
11.4.1 Outpatient care	683
11.4.2 Inpatient care	686
11.4.3 Combined inpatient and outpatient care.....	687
11.5 Economic evidence	692
11.6 Evidence statements	693
11.6.1 Outpatient care	693
11.6.2 Inpatient care	696
11.6.3 Combined inpatient and outpatient care.....	697
11.6.4 Economic evidence statements.....	702
11.7 Evidence to recommendations	703
11.7.1 Relative value placed on the outcomes considered.....	703
11.7.2 Consideration of clinical benefits and harms	703
11.7.3 Consideration of economic benefits and harms.....	705
11.7.4 Quality of evidence	707
11.7.5 Other considerations	708
11.7.6 Key conclusions	708
11.8 Recommendations	708
12 References.....	710
13 Glossary and abbreviations.....	749

1

1 Introduction

Cystic fibrosis is a multi-system genetic disorder affecting the lungs, pancreas, liver and intestine. It can have a significant impact on life expectancy and quality of life. The current median age of those who have died is 28 years and the median predicted survival is 45.1 years.

Diagnosis is primarily made during newborn screening. The median age at diagnosis is 3 months, and 1 in every 2500 babies born in the UK has cystic fibrosis. More than 57% of people on the UK cystic fibrosis registry are aged over 16 years.

Many different mutations are responsible for cystic fibrosis. The UK registry shows that 90.8% of people with cystic fibrosis have one known genotype. However 8.9% of people have at least one unknown genotype.

Lung function is often reduced in cystic fibrosis. The typical measure of lung function is forced expiratory volume in 1 second (FEV₁). A FEV₁ of 50% and above will enable people to live relatively normal lives and is associated with fewer difficulties in completing activities of daily living. A FEV₁ above 85% indicates normal or near-normal lung function.

Lung infections are a cause of significant morbidity in cystic fibrosis. Chronic infection (for example with *S aureus* and *Pseudomonas aeruginosa*) may need long-term use of antibiotics.

There is variation across the country in the multidisciplinary team structures used, the arrangements services make for providing care, and in the resources available to support services. Particular problems may arise with smaller shared-care clinic arrangements. In some centres, both inpatient and outpatient facilities are limited. For example, there may be problems in arranging admission to single rooms with en-suite facilities. If adequate protocols are not in place, then there is a risk of cross-infection.

By making robust recommendations based on the available evidence and best practice in cystic fibrosis care, this guideline will help improve care for this highly complex condition.

2 Guideline summary

2.1 Guideline Committee membership, NGA staff and acknowledgements

Table 1: Guideline Committee Members

Name	Role
Mandy Bryon	Senior Consultant Clinical Psychologist Head of Psychological Services, Great Ormond Street Hospital for Children NHS Trust
Janis Bloomer	Paediatric Nurse Specialist (Children and young people) Cystic Fibrosis, Great North Children's Hospital Royal Victoria Infirmary
Sarah Collins	Cystic Fibrosis Specialist Dietician, Nutrition & Dietetic Department, Royal Brompton Hospital
Alexander Darlington	Lay Member
Iolo Doull	Consultant Respiratory Paediatrician, Children's Hospital for Wales, Cardiff
Elaine Edwards	Advanced Physiotherapist, Sheffield Children's NHS Foundation Trust
Zoe Elliott	Lay Member
Andrew Jones	Consultant and Honorary Reader in Respiratory Medicine and Cystic Fibrosis, North West Lung Centre, University Hospitals South Manchester NHS Foundation Trust
David Lacy	General Paediatrician Wirral University Teaching Hospital NHS Foundation Trust
Nichola MacDuff	Adult Specialist Nurse, Advanced Clinical Nurse Specialist & Lead Nurse, Black Country Adult CF Centre, Royal Wolverhampton NHS Trust.
Helen McCabe	Principal Paediatric Dietitian - Royal Victoria Infirmary, Newcastle upon Tyne Hospitals NHS Foundation Trust.
Helen Parrott	Physiotherapist, Clinical Specialty Lead, Adult CF Therapies Royal Brompton Hospital
Sarah Pople	Pharmacist, Leicester Royal Infirmary Children's Hospital / De Montfort University
Keith Thompson	Senior Respiratory Pharmacist, Royal Brompton and Harefield NHS Foundation Trust
Martin Walshaw (Chair)	Consultant Physician in General and Chest Medicine at Royal Liverpool and Broadgreen University Hospitals NHS Trust, and The Liverpool Heart and Chest Hospital NHS Foundation Trust
Co-opted members	
Stuart Elborn	Dean, School of Medicine, Dentistry and Biomedical Sciences, Queen's University, Belfast
Stephen Holden	Consultant Medical Microbiologist Nottingham University Hospitals NHS Trust

Table 2: NGA Staff

Name	Role
Omnia Abdulrazeg	Research Fellow (until January 2016)
Stephanie Arnold	Information Scientist (from May 2016)
Alexander Bates	Guideline Lead (November 2016 – January 2017)
Zosia Beckles	Information Scientist (until March 2016)

Name	Role
Bishal Bhandari	Assistant Systematic Reviewer (September 2016 – November 2016)
Lisa Boardman	Guideline Lead (from January 2017)
Me'leshah Brown	Project Administrator
Shona Burman-Roy	Senior Systematic Reviewer (March 2015 – May 2016)
Rami Cosulich	Systematic Reviewer (from March 2016)
Katherine Cullen	Senior Health Economist (until December 2014)
Vanessa Delgado-Nunes	Guideline Lead (until November 2016)
Annabel Flint	Project Manager (until September 2016)
Paul Jacklin	Senior Health Economist (January 2015 – July 2015)
Gemma Marцениuk	Health Economist (from July 2015)
Maija Kallioinen	Systematic Reviewer (August 2016 – September 2016)
Stephen Murphy	Clinical Advisor
Fionnuala O'Brien	Project Manager (from September 2016)
Hugo Pedder	Statistician
Gemma Villanueva	Senior Systematic Reviewer (from March 2015)

1 Acknowledgements

2 Additional support was received from Ebenezer Ademoisoye (Health Economist), Alex Bates
3 (Senior Health Economist), Maija Kallioinen (Systematic Reviewer), Abigail Moore (Assistant
4 Systematic Reviewer), Ferruccio Pelone (Systematic Reviewer) and Natasha Pillai
5 (Systematic Reviewer), Tim Reeves (Information Scientist).

6 2.2 Other versions of the guideline

7 NICE produces a number of versions of this guideline:

- 8 • The 'short guideline' lists the recommendations, context and recommendations for
9 research.
- 10 • NICE Pathways brings together all connected NICE guidance.

11 2.3 Schedule for updating the guideline

12 For the most up-to-date information about guideline reviews, please see the latest version of
13 the NICE guidelines manual available from the NICE website.

3 Development of this guideline

3.1 What is a NICE Guideline?

NICE guidelines are recommendations for the care of individuals in specific clinical conditions or circumstances within the NHS – from prevention and self-care through primary and secondary care to more specialised services. We base our NICE guidelines on the best available research evidence, with the aim of improving the quality of healthcare. We use predetermined and systematic methods to identify and evaluate the evidence relating to specific review questions.

NICE guidelines can:

- Provide recommendations for the treatment and care of people by healthcare professionals.
- Be used to develop standards to assess the clinical practice of individual healthcare professionals.
- Be used in the education and training of healthcare professionals.
- Help patients to make informed decisions.
- Improve communication between patients and healthcare professionals.

While guidelines assist the practice of healthcare professionals, they do not replace their knowledge and skills.

We produce our guidelines using the following steps:

- The guideline topic is referred to NICE from the Department of Health.
- Stakeholders register an interest in the guideline and are consulted throughout the development process.
- The scope is prepared by the NGA.
- The NGA establishes a guideline committee.
- A draft guideline is produced after the Committee members assess the available evidence and makes recommendations.
- There is a consultation on the draft guideline.
- The final guideline is produced.

The NGA and NICE produce a number of versions of this guideline.

- The 'full guideline' contains all the recommendations, together with details of the methods used and the underpinning evidence.
- The 'short guideline' lists the recommendations, context and recommendations for research.
- NICE Pathways brings together all connected NICE guidance.

3.2 Remit

NICE received the remit for this guideline from the Department of Health. It commissioned the NGA to produce the guideline.

The remit for this guideline is to develop a NICE guideline on the diagnosis and management of Cystic Fibrosis.

1 **3.3 Who developed this guideline?**

2 A multidisciplinary Guideline Committee comprising healthcare professionals and
3 researchers as well as lay members developed this guideline (see the list of group members
4 and acknowledgements).

5 NICE funds the NGA and thus supported the development of this guideline. The Guideline
6 Committee was convened by the NGA and chaired by Dr Martin Walshaw in accordance with
7 guidance from NICE.

8 The group met every 4 to 6 weeks during the development of the guideline. At the start of the
9 guideline development process all group members declared interests including
10 consultancies, fee-paid work, shareholdings, fellowships and support from the healthcare
11 industry. At all subsequent group meetings, members declared arising conflicts of interest.

12 Members were either required to withdraw completely or for part of the discussion if their
13 declared interest necessitated it appropriate to do so. The details of declared interests and
14 the actions taken are shown in Appendix C.

15 Staff from the NGA provided methodological support and guidance for the development
16 process. The team working on the guideline included a guideline lead, a project manager,
17 systematic reviewers, health economists, a statistician and information scientists. They
18 undertook systematic searches of the literature, appraised the evidence, conducted meta-
19 analysis and cost-effectiveness analysis where appropriate and drafted the guideline in
20 collaboration with the group.

21 **3.4 What this guideline covers**

22 **3.4.1 Groups that will be covered**

23 The guideline covers the following groups.

- 24 • Infants, children, young people and adults with cystic fibrosis, including those who have
25 non-classic cystic fibrosis and those who have had an organ transplant.

26 **3.4.2 Key clinical issues that will be covered**

27 The following clinical issues will be covered in this guideline:

- 28 • The clinical manifestations of cystic fibrosis at the time of diagnosis in infants, children,
29 young people and adults.
- 30 • The complications of cystic fibrosis.
- 31 • Management of chest disease:
- 32 ○ Routine monitoring of lung disease, including microbiological surveillance, radiological
33 imaging and pulmonary function testing.
 - 34 • Antimicrobial management in cystic fibrosis to:
 - 35 ○ prevent bacterial colonisation
 - 36 ○ treat acute pulmonary infection
 - 37 ○ treat chronic pulmonary infection, including clinical exacerbations and colonisation.
 - 38 • Immunomodulatory management in chest disease.
 - 39 • Management with mucoactive or mucolytic agents.
 - 40 • Chest physiotherapy.
 - 41 • The role of exercise in maintaining health.
 - 42 • Management of nutrition.

- 1 • Management of exocrine pancreatic insufficiency.
- 2 • Management of distal intestinal obstruction syndrome.
- 3 • Surveillance for cystic-fibrosis-related diabetes.
- 4 • Surveillance for cystic-fibrosis-related liver disease and prevention of progression.
- 5 • Surveillance for reduced bone mineral density.
- 6 • Recognising psychological and behavioural problems.
- 7 • Models for delivery of care and multidisciplinary teams.
- 8 • Provision of information and support for infants, children, young people, adults and their
- 9 carers

10 Note that guideline recommendations will normally fall within licensed indications.
11 Exceptionally, and only if clearly supported by evidence, the use outside a licensed indication
12 may be recommended. This guideline will assume that prescribers will use a drug's summary
13 of product characteristics to inform decisions made with individual patients.

14 For further details please refer to the scope in Appendix A and review questions in Appendix
15 D.

16 **3.5 What this guideline does not cover**

17 **3.5.1 Clinical issues that will not be covered**

18 This guideline does not cover:

- 19 • Specialist management of cystic-fibrosis-related diabetes.
- 20 • Specialist management of cystic-fibrosis-related fertility and pregnancy problems.
- 21 • Specialist management of cystic-fibrosis-related liver disease.
- 22 • Specialist management of cystic-fibrosis-related bone disease.
- 23 • Specialist management of cystic-fibrosis-related ear, nose and throat (ENT) disorders.
- 24 • Specialist management of cystic-fibrosis-related renal disease.
- 25 • Surgical management of gastrointestinal complications.
- 26 • Referral for, and management of, transplantation.
- 27 • Management of specific psychological conditions.
- 28 • Management of specific behavioural problems.

29 **3.6 Relationship between the guideline and other NICE** 30 **guidance**

31 **3.6.1 Related NICE guidance**

- 32 • Gastro-oesophageal reflux disease (2015) NICE guideline NG1
- 33 • Dyspepsia and gastro- oesophageal reflux disease (2014) NICE guideline CG184
- 34 • Infection: prevention and control of healthcare-associated infections in primary and
35 community care (2012) NICE guideline CG139
- 36 • Constipation in children and young people (2010) NICE guideline CG99
- 37 • Depression in adults with a chronic physical health problem (2009) NICE guideline CG91
- 38 • Living-donor lung transplantation for end-stage lung disease (2006) NICE interventional
39 procedure guidance 170

4 Guideline development methodology

This chapter describes the methods used to review the evidence and generate the recommendations presented in subsequent chapters. This guidance was developed in accordance with the methods outlined in the NICE guidelines manual 2012 (PMG 6) for the scoping phase, and the NICE guidelines manual 2014 (PMG 20) from the development phase.

Declarations of interest were recorded according to the 2014 NICE conflicts of interest policy.

4.1 Developing the review questions and outcomes

The review questions were drafted by the NGA, and refined and validated by the guideline Committee. The questions were based on the key areas identified in the guideline scope (See Appendix A).

A total of 29 questions were identified (See Table 3).

The review questions were based on the following frameworks:

- intervention reviews – using population, intervention, comparator and outcome (PICO framework)
- reviews of diagnostic test accuracy – using population, diagnostic test (index tests), reference standard and target condition
- prognostic reviews – using population, presence or absence of a risk factor, and outcome
- prevalence reviews – using population and outcome (prevalence of target condition)
- qualitative reviews – using population, area of interest and themes of interest

These frameworks guided the literature searching process, critical appraisal and synthesis of evidence and facilitated the development of recommendations by the Guideline Committee.

Full literature searches, critical appraisals and evidence reviews were completed for all review questions.

Table 3: Description of review questions

Chapter or section number	Type of review	Review questions	Outcomes
5	Clinical prediction	In infants, children, young people and adults (including those that have undergone newborn screening) when should cystic fibrosis be suspected?	<ul style="list-style-type: none"> • Risk of cystic fibrosis (risk ratios and odds ratios) • Sensitivity • Specificity • Positive predictive value • Negative predictive values • Positive likelihood ratios • Negative likelihood ratios
6	Qualitative	What information and support should be given to children, young people and adults with cystic fibrosis?	<p>Themes will be identified from the literature, but expected themes are, for example:</p> <ul style="list-style-type: none"> • Timing of when support is given • Regular formal assessments tailored to needs • Psychosocial support

Chapter or section number	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Information about named individual for point of contact. • Information at the time of diagnosis. • Clear and accurate information about cystic fibrosis • Social media, apps and technology. • Discussion about planning management of cystic fibrosis. • Information for access to resources for managing co-morbidities. • Information on pregnancy and fertility. • Education and healthcare at school.
7.1	Intervention	What is the effectiveness of different models of care (for example, specialist centre, shared care [delivered by a Network CF Clinic which is part of an agreed designated network with a Specialist CF Centre], community, telehealth and/or home care for people with cystic fibrosis?	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) ◦ Lung clearance index (LCI) • Time to next pulmonary exacerbation • Mortality • Nutritional status • Quality of life (measured with CF-QOL or CFQR) • Patient and carer satisfaction • Frequency of cross-infections (<i>Pseudomonas aeruginosa</i>, <i>Burkholderia cepacia</i>) • Staff experience • Adherence to treatment
7.2	Intervention	What is the clinical and cost-effectiveness of multidisciplinary teams of various compositions?	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) ◦ Lung clearance index (LCI) • Time to next pulmonary exacerbation • Mortality • Nutritional status • Quality of life (measured with CF-QOL or CFQR) • Patient and carer satisfaction • Frequency of cross-infections (<i>P. aeruginosa</i>, <i>B. cepacia</i>) • Staff experience • Adherence to treatment
7.3	Qualitative	What parts of the transition from children's to adult services are most important for young people with cystic fibrosis and their family	<p>Themes will be identified from the literature, but expected themes are, for example:</p> <ul style="list-style-type: none"> • Transition clinic

Chapter or section number	Type of review	Review questions	Outcomes
		members and carers?	<ul style="list-style-type: none"> • Transition lead (consultant/social worker) preparation of plan of transition for individual and family/carers, multidisciplinary structured approach. • Involvement of young people and family/carer in planning, implementing and reviewing transition. • Key transition therapist as part of paediatric and adult service. • Communication/co-ordination between paediatric and adult services. • Timing of transition.
8	Prevalence	What are the non-lower-respiratory complications of cystic fibrosis in infants, children, young people and adults?	<p>Prevalence of complications of cystic fibrosis.</p> <ul style="list-style-type: none"> • Malnutrition • Impaired growth • Cystic fibrosis related renal disease • Delayed puberty • Distal intestinal obstruction syndrome (DIOS) • Abdominal pain • Cystic fibrosis related diabetes • Upper airways disease • Cystic fibrosis related musculoskeletal disorders • Urinary stress incontinence • Reduced bone mineral density • Cystic fibrosis related liver disease • Infertility • Meconium ileus
10.8	Intervention	What is the effectiveness of programmes of exercise in the management of cystic fibrosis?	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Forced vital capacity (FVC) ○ Lung clearance index (LCI) • VO₂ • Time to next exacerbation • Body composition • Quality of life (measured with CF-QOL or CFQR) • Preference for training programme • Adverse event
10.1	Intervention	What is the clinical and cost effectiveness of nutritional interventions in people with cystic fibrosis?	<ul style="list-style-type: none"> • Body composition • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Pulmonary exacerbations • Changes to body composition

Chapter or section number	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> detected by anthropometric measure Quality of life (measured with CF-QOL or CFQR) Satisfaction Adverse effects
10.2	Intervention	In people with cystic fibrosis, what is the most effective regimen of enzyme replacement therapy in the treatment of exocrine pancreatic insufficiency?	<ul style="list-style-type: none"> Nutritional status Reduction of steatorrhea and faecal fat (coefficient of fat absorption and faecal fat excretion specific outcomes and others) Resolution of symptoms of malabsorption Quality of life (measured with CF-QOL or CFQR) Satisfaction Drug related side effects/ adverse events
10.3	Intervention	What are the effective strategies for treatment and secondary prevention of distal ileal obstruction syndrome?	<p>Primary treatment of DIOS:</p> <ul style="list-style-type: none"> Reduction in clinical manifestations including Adverse events from treatment Patient satisfaction Duration of hospital stay Treatment failure (need for surgery) Adverse events from surgery <p>Secondary prevention of DIOS:</p> <ul style="list-style-type: none"> Reduction in clinical manifestations including Adverse events from treatment Patient satisfaction Recurrence of DIOS Admission to hospital
10.4	Diagnostic	What is the diagnostic accuracy of tests to detect or strategies to detect early and late cystic fibrosis liver disease?	<p>For the diagnostic accuracy question:</p> <ul style="list-style-type: none"> Positive likelihood ratios/ Negative likelihood ratios (LR+/ LR-) Sensitivity/ Specificity Area under the curve (AUC) <p>For the following target conditions:</p> <ul style="list-style-type: none"> Liver disease Cirrhosis Portal hypertension Oesophageal varices
10.4	Prognostic	What is the diagnostic and prognostic value of different strategies to detect cystic fibrosis liver disease and predict progression (including	<ul style="list-style-type: none"> Adjusted odds ratios (adjORs) Adjusted hazard ratios (adjHRs) For the identification of: Liver disease

Chapter or section number	Type of review	Review questions	Outcomes
		progression to cirrhosis and portal hypertension with (out) oesophageal varices)?	<ul style="list-style-type: none"> • Cirrhosis • Portal hypertension
10.5	Intervention	What is the effectiveness of ursodeoxycholic acid for preventing the development or progression of liver disease in people with cystic fibrosis?	<ul style="list-style-type: none"> • Change of hepatocellular enzymes or bilirubin level • Liver failure • Liver transplantation • Liver related mortality • Development of portal hypertension indicated by an: <ul style="list-style-type: none"> ○ Enlarged spleen (increased by at least 15%) ○ Development of varices ○ Ultrasound evidence of portal hypertension ○ No development of liver disease • Quality of life (measured with CF-QOL or CFQR)
10.6	Prognostic	What is the most effective strategy to monitor for the onset of CF-related diabetes (CFRD)?	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Forced vital capacity (FVC) ○ Lung clearance index (LCI) • Pulmonary exacerbation • Body mass index (BMI) • Adverse events • Patient acceptability/ satisfaction (with insulin therapy)
10.7	Prognostic	What is the most effective strategy to monitor for the identification of reduced bone mineral density in people with cystic fibrosis?	<ul style="list-style-type: none"> • Change in body bone mineral density • Number of fractures • Quality of life (measured with CF-QOL or CFQR)
10.9	Clinical prediction	What strategies are effective at identifying people with cystic fibrosis for the presence of a psychological and/or behavioural problem?	<ul style="list-style-type: none"> • Sensitivity • Specificity • Positive likelihood ratio (LR+) • Negative likelihood ratio (LR-) • AUROC • Reliability and validity
9.1	Intervention	<p>What is the value of the following investigative strategies in monitoring the onset of pulmonary disease in people with cystic fibrosis without clinical signs or symptoms of lung disease?</p> <ul style="list-style-type: none"> • Non-invasive microbiological investigation: induced sputum samples, cough swab, throat swab, and nasopharyngeal aspiration 	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Lung clearance index (LCI) • Quality of life (measured with CF-QOL or CFQR) • Nutritional parameters • Time to chronic infection with <i>P. aeruginosa</i> or <i>Staphylococcus aureus</i> • Clearance of the organism from the cultures

Chapter or section number	Type of review	Review questions	Outcomes
		<ul style="list-style-type: none"> • Invasive microbiological investigation: bronchoalveolar lavage • Lung physiological function tests: ardiopulmonary exercise testing, Spirometry and Lung Clearance Index • Imaging techniques: chest x-ray and CT scan 	
9.1	Intervention	<p>What is the value of the following investigative strategies in monitoring evolving pulmonary disease in people with established lung disease?</p> <ul style="list-style-type: none"> • Non-invasive microbiological investigation: induced sputum samples, cough swab, throat swab, and nasopharangeal aspiration • Invasive microbiological investigation: bronchoalveolar lavage • Lung physiological function tests: cardiopulmonary exercise testing, Spirometry and lung clearance index • Imaging techniques: chest X-ray and CT scan <p>Evolving pulmonary disease defined as decline in lung function (based on FEV₁), increased exacerbations and/or infections, and CT changes.</p>	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Lung clearance index (LCI) • VO₂ max • Nutritional parameters • Time to next exacerbation • Time to chronic infection using any recognized definition with <i>P aeruginosa</i> or with <i>S aureus</i> • Mortality • Quality of life (measured with CF-QOL or CFQR)
9.1	Intervention	<p>What is the added value of imaging and invasive microbiological testing in addition to non-invasive microbiological testing and lung function tests in monitoring the response to treatment following an acute exacerbation?</p> <p>Definition of stablished lung disease: clinical signs and symptoms and/or radiological signs of lung disease.</p>	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Lung clearance index (LCI) • Oxygen saturation • High-resolution computed tomography (CT) appearances or Chest X-ray appearances • Nutritional parameters • Time to next exacerbation • Clearance of the organism from the cultures • Inflammatory markers • Quality of life (measured with CF-QOL or CFQR)
9.2	Intervention	<p>What is the effectiveness of airway clearance techniques in people with cystic fibrosis?</p>	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ○ Forced expiratory volume (FEV₁) ○ Forced vital capacity (FVC)

Chapter or section number	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Oxygen saturation • Expectorated secretions (mucus, sputum, phlegm) • Sputum volume • Patient preference • Pulmonary exacerbations • Hospitalisations, change in frequency • Quality of Life (measured with CF-QOL or CFQR)
9.3	Intervention	What is the effectiveness of mucoactive or mucolytic agents, including dornase alpha, nebulised sodium chloride (isotonic and hypertonic) and mannitol in people with cystic fibrosis?	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume(FEV₁) • Inflammatory markers (Serum or Sputum) • Time to pulmonary exacerbations • Need for intravenous antibiotics for pulmonary exacerbation • Quality of life (measured with CF-QOL or CFQR) • Adverse events
9.4.1	Intervention	What is the effectiveness of long-term antimicrobial prophylaxis to prevent pulmonary bacterial colonisation with <i>S aureus</i> in people with cystic fibrosis?	<ul style="list-style-type: none"> • Time to identification of the pathogen (<i>S aureus</i>) in sputum culture • Number of positive pathogen cultures (<i>S aureus</i>) identified during study period • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume(FEV₁) ◦ Lung clearance index (LCI) • Time to next pulmonary exacerbation • Quality of life (measured with CF-QOL or CFQR) • Adherence to treatment • Adverse events • Emergence of resistant organisms
9.4.2	Intervention	What is the effectiveness of antimicrobial treatment for acute pulmonary infection or those with an exacerbation in children and adults with cystic fibrosis?	<ul style="list-style-type: none"> • For pulmonary exacerbation: • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) ◦ Lung clearance index (LCI) • Eradication of specific pathogen • Resolution of infection/exacerbation or measure of treatment failure (e.g. need for additional antibiotics) • Duration of the acute episode • Quality of life (measured with CF-QOL or CFQR) • Mortality

Chapter or section number	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Adverse events • For acute infection: • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) ◦ Lung clearance index (LCI) • Eradication of specific pathogen • Time to next pulmonary exacerbation • Resolution of infection/exacerbation or measure of treatment failure • Quality of life (measured with CF-QOL or CFQR) • Adverse events
9.4.3	Intervention	<p>What is the effectiveness of antimicrobial regimens in suppressing chronic pulmonary infection in children and adults with cystic fibrosis with any of the following pathogens:</p> <ul style="list-style-type: none"> • <i>Pseudomonas aeruginosa</i> • <i>B cepacia complex</i> • <i>S aureus</i> • <i>Aspergillus fumigatus</i> 	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) • Time to next pulmonary exacerbation • Number of patients with at least 1 pulmonary exacerbation • Eradication of the specified organism from sputum/airway cultures • Nutritional status • Quality of life (measured with CF-QOL or CFQR) • Adverse events • Emergence of resistant organisms/antibiotic resistance
9.5	Intervention	<p>What is the effectiveness of immunomodulatory agents in the management of lung disease?</p>	<ul style="list-style-type: none"> • Lung function <ul style="list-style-type: none"> ◦ Forced expiratory volume (FEV₁) • Nutritional status • Time to next pulmonary exacerbation • Quality of life (measured with CF-QOL or CFQR) • Adverse effects • Mortality
11	Intervention	<p>What is the effectiveness of cohorting on the basis of pathogen status versus not cohorting on the basis of pathogen status in reducing transmission of CF pathogens?</p>	<ul style="list-style-type: none"> • Incidence of patients infected with transmissible pathogens • Prevalence of patients infected with transmissible pathogens • Quality of life (measured with CF-QOL or CFQR)
11	Intervention	<p>What is the effectiveness of different models of segregating patient's in reducing transmission of CF pathogens?</p>	<ul style="list-style-type: none"> • Emotional function including anxiety and depression • Carer satisfaction • Patient satisfaction

Chapter or section number	Type of review	Review questions	Outcomes
11	Intervention	What is the effectiveness of individual protective equipment in reducing transmission of CF pathogens?	<ul style="list-style-type: none"> • Staff experience • Staff and patient compliance
11	Intervention	What is the effectiveness of the combination of cohorting, segregating and protective equipment in reducing transmission of CF pathogens?	

1 4.2 Searching for evidence

2 Systematic literature searches were undertaken to identify all published clinical evidence
3 relevant to the review questions.

4 Databases were searched using relevant medical subject headings, free-text terms and
5 study type filters where appropriate. Studies published in languages other than English were
6 not reviewed. Where possible, searches were restricted to retrieve only articles published in
7 English. All searches were conducted in MEDLINE, Embase and The Cochrane Library. The
8 following searches were updated in January 2017.

- 9
- 10 • Monitoring for liver disease
 - 11 • Airway clearance
 - 12 • Monitoring pulmonary disease
 - 13 • Mucoactive agents
 - 14 • Immunomodulatory agents
 - 15 • Antimicrobials: prophylaxis
 - 16 • Antimicrobials: acute
 - 17 • Antimicrobials: chronic
 - 18 • Nutrition interventions
 - 19 • Exercise
 - 20 • Service configuration: cross-infection
 - 21 • Service configuration: MDT
 - 22 • Service configuration: models of care

22 We prioritised the list below for re-runs based on the following criteria:

- 23 • Topics where network meta-analyses (NMAs) and HE modelling work have been
24 conducted
- 25 • Topics with significant evidence movement where it is likely that new evidence will have
26 been published
- 27 • Topics that are part of the service delivery component of the guideline
- 28 • Topics with empty reviews (i.e. MDT)
- 29 • Topics that have been covered earlier in guideline development which may now be at
30 greater risk of being out of date.

31 Any studies added to the databases after this date (even those published prior to this date)
32 were not included unless specifically stated in the text.

1 Search strategies were quality assured by cross-checking reference lists of highly relevant
2 papers, analysing search strategies in other systematic reviews and asking the group
3 members to highlight any additional studies. The questions, the study types applied, the
4 databases searched and the years covered can be found in Appendix E.

5 The titles and abstracts of records retrieved by the searches were inspected for relevance,
6 with potentially significant publications obtained in full text. These were assessed against the
7 inclusion criteria.

8 During the scoping stage, a search was conducted for guidelines and reports on websites of
9 organisations relevant to the topic. Searching for grey literature or unpublished literature was
10 not undertaken. Searches for electronic, ahead-of-print publications were not routinely
11 undertaken unless indicated by the Guideline Committee. All references suggested by
12 stakeholders at the scoping consultation were initially considered.

13 4.2.1 Health economic literature search

14 A global search of economic evidence relating to all treatments for cystic fibrosis was
15 undertaken in April 2015 and re-ran in January 2017. The following databases were
16 searched:

- 17 • MEDLINE (Ovid);
- 18 • EMBASE (Ovid);
- 19 • Cochrane Central Register of Controlled Trials (CCTR);
- 20 • HTA database (HTA);
- 21 • NHS Economic Evaluations Database (NHS EED).

22 Further to the database searches, the Committee was contacted with a request for details of
23 relevant published and unpublished studies of which they may have knowledge; reference
24 lists of key identified studies were also reviewed for any potentially relevant studies. Finally,
25 the NICE website was searched for any recently published guidance relating to cystic fibrosis
26 that had not been already identified via the database searches.

27 The search strategy for existing economic evaluations combined terms capturing the target
28 condition (cystic fibrosis) and, for searches undertaken in MEDLINE, EMBASE and CCTR,
29 terms to capture economic evaluations. No restrictions on language or setting were applied
30 to any of the searches, but a standard exclusions filter was applied (letters, animals, etc).
31 Conference abstracts were considered for inclusion from 1st January 2014, as high-quality
32 studies reported in abstract form before 2014 were expected to have been published in a
33 peer-reviewed journal. Full details of the search strategies are presented in Appendix E.

34 The titles and abstracts of papers identified through the searches were independently
35 assessed for inclusion using pre-defined eligibility criteria defined in Table 4.

36 **Table 4: Inclusion and exclusion criteria for the systematic reviews of economic**
37 **evaluations**

Inclusion criteria
intervention or comparators according to the scope
study population according to the scope
full economic evaluations (cost-utility, cost-effectiveness, cost-benefit or cost-consequence analyses) that assess both the costs and outcomes associated with the interventions of interest
Exclusion criteria
abstracts with insufficient methodological details
cost-of-illness type studies
conference papers pre January 2014

1 Once the screening of titles and abstracts was complete, full versions of the selected papers
2 were acquired for assessment. The Preferred Reporting Items for Systematic Reviews and
3 Meta-Analyses (PRISMA) for this search on economic evaluations is presented in Appendix
4 F.

5 The quality of evidence was assessed using the economic evaluations checklist as specified
6 in the NICE guidelines manual. Quality assessments of included studies and data extraction
7 tables are provided in Appendix L and M, respectively.

8 **4.3 Reviewing and synthesising research evidence**


9 **4.3.1 Systematic review process**

- 10 • The evidence was reviewed following these steps (See Figure 1):
- 11 • Potentially relevant studies were identified for each review question from the relevant
12 search results by reviewing titles and abstracts. Full papers were then obtained.
- 13 • Full papers were reviewed against pre-specified inclusion and exclusion criteria in the
14 review protocols (in Appendix D).
- 15 • Key information was extracted on the study's methods, according to the factors specified
16 in the protocols and results. These were presented in summary tables (in each review
17 chapter) and evidence tables (in Appendix G)
- 18 • Relevant studies were critically appraised using the appropriate checklist as specified in
19 the NICE guidelines manual (NICE 2014).
- 20 • Summaries of evidence were generated by outcome (included in the relevant review
21 chapters) and were presented in committee meetings (details of how the evidence was
22 appraised is described in Section 4.3.5 below):
- 23 ○ Randomised studies: meta-analysis was carried out where appropriate and results
24 were reported in GRADE profiles (for intervention reviews).
 - 25 ○ Observational studies: data were presented individually by study in GRADE profiles.
 - 26 ○ Prognostic studies: data were presented individually by study, usually in terms of the
27 relative effect as reported by the authors.
 - 28 ○ Diagnostic studies: data were presented individually by study as measures of
29 diagnostic test accuracy (sensitivity and specificity, positive and negative likelihood
30 ratios) and were presented in modified GRADE profiles.
 - 31 ○ Prevalence studies: data were presented as measures of prevalence during a period of
32 time (proportions with their 95% confidence intervals); the decision if meta-analysis
33 could be conducted was based on the consideration of the heterogeneity of the studies.
 - 34 ○ Qualitative studies: each study was summarised by theme and meta-synthesis was
35 carried out where appropriate to identify an overarching framework of themes and
36 subthemes. These were then presented in GRADE-CERQual (Lewin 2015) profiles,
37 where CERQual stands for Confidence in the Evidence from Reviews of Qualitative
38 research.

39 For quality assurance of study identification, either whole study selections or a sample of the
40 study selection results were double checked by a second reviewer. Searches related to the
41 NMA were also double sifted.

42 A sample of all evidence tables, including a sample of evidence tables related to the NMA
43 were. All drafts of reviews were checked by a second reviewer. Any discrepancies were
44 resolved by discussion between the 2 reviewers.

Figure 1: Step-by-step review of evidence in the guideline



1 **4.3.2 Inclusion/exclusion criteria**

2 The inclusion and exclusion of studies was based on the review protocols, which can be
3 found in appendix D. Excluded studies by review question (with the reasons for their
4 exclusion) are listed in appendix H. In addition, the Committee was consulted about any
5 uncertainty regarding inclusion or exclusion.

6 **4.3.3 Type of studies**

7 Systematic reviews (SRs) with meta-analyses were considered the highest quality evidence
8 to be selected for inclusion.

9 For most intervention reviews in this guideline, parallel randomised controlled trials (RCTs)
10 were prioritised because they are considered the most robust type of study design that could
11 produce an unbiased estimate of the intervention effects. Crossover RCTs were appropriate
12 for some of the interventional questions. If there was limited evidence from RCTs,
13 observational studies were included.

14 For diagnostic reviews, cross-sectional, retrospective or prospective observational studies
15 were considered for inclusion. Where evidence was limited, case-control studies were also
16 considered for inclusion.

17 For clinical prediction and prognostic reviews, prospective and retrospective cohort studies
18 were included.

1 For prevalence reviews, the Committee prioritised the CF UK registry. Where no evidence
2 was available or the Committee agreed the CF registry data provided limited for a particular
3 complication, cross-sectional studies and prospective cohort studies (national registries were
4 preferred) were also included.

5 For qualitative reviews, studies using focus groups, or structured or semi-structured
6 interviews were considered for inclusion. Survey data or other types of questionnaires were
7 only included if they provided analysis from open-ended questions, but not if they reported
8 descriptive quantitative data only.

9 Literature reviews, posters, letters, editorials, comment articles, unpublished studies and
10 studies not in English were excluded. Conference abstracts were only considered for
11 inclusion in the absence of full published studies.

12 **4.3.3.1 Incorporation of technology appraisals**

13 Two NICE Technology Appraisals were identified as relevant for this guideline (TA276 and
14 TA266). It was agreed that the recommendations were still current, and therefore they were
15 lifted unchanged from the original guidance.

16 It was also decided that full systematic reviews were to be conducted based on the review
17 protocols agreed with the Committee, as this would allow to include studies that were not
18 included in the Technology Appraisals.

19 **4.3.4 Types of data and methods for synthesis**

20 **4.3.4.1 Data synthesis for intervention studies**

21 **4.3.4.1.1 Pairwise meta-analysis**

22 Meta-analysis was conducted whenever it could be robustly performed, to combine the
23 results of studies for each review question using Cochrane Review Manager (RevMan5)
24 software.

25 The generic inverse variance option in RevMan5 was used where any studies reporting
26 solely the summary treatment effect and 95% confidence interval (95% CI) or standard error
27 could be included.

28 Fixed-effect (Mantel–Haenszel) techniques were used in the first instance to calculate risk
29 ratios (relative risk) for binary outcomes, such as rate of adverse events or rate of people
30 with symptom improvements (Mantel 1959).

31 For continuous outcomes, measures of central tendency (mean) and variation (standard
32 deviation) are required for meta-analysis. However, in cases where standard deviations were
33 not reported per intervention group, the standard error (SE) for the mean difference was
34 calculated from other reported statistics (p-values or 95% CIs): meta-analysis was then
35 undertaken for the mean difference and SE using the generic inverse variance method in
36 RevMan5

37 When the only evidence was based on studies summarising results by presenting medians
38 (and interquartile ranges) or only p values were given, this information was assessed in
39 terms of the study's sample size and was included in the GRADE tables without calculating
40 the relative or absolute effects. Consequently, aspects of quality assessment, such as
41 imprecision of effect, could not be assessed for evidence of this type. However, the limited
42 reporting of this outcome was classified as a risk of bias in study limitations.

1 Stratified analyses were predefined for some review questions at the protocol stage when the
2 Committee identified that these strata are different in terms of biological and clinical
3 characteristics and the interventions were expected to have a different effect.

4 Statistical heterogeneity was assessed by visually examining the forest plots (please see
5 Appendix I) and by considering the chi-squared test for significance at $p < 0.1$ or an I-squared
6 inconsistency statistic (with an I-squared value of more than 50% indicating considerable
7 heterogeneity). Where considerable heterogeneity was present, predefined subgroup
8 analyses were performed.

9 Assessments of potential differences in effect between subgroups were based on the chi-
10 squared tests for heterogeneity statistics between subgroups. If no sensitivity analysis was
11 found to completely resolve statistical heterogeneity, then a random-effects (DerSimonian
12 and Laird) model was employed to provide a more conservative estimate of the effect –
13 (DerSimonian 1986).

14 Where data from observational studies were included, the Committee decided that the
15 results for each outcome should be presented separately for each study and meta-analysis
16 was not conducted.

174.3.4.1.2 **Network Meta-Analysis (NMA)**

18 In some circumstances, the results of conventional pairwise meta-analyses of direct
19 evidence does not help assess which intervention is most effective. The challenge of
20 interpretation may arise for two main reasons:

- 21 • Relative treatment efficacies based on separate individual pairwise comparisons across
22 multiple treatments are difficult to assess.
- 23 • Direct RCT comparison between treatments of clinical interest are not available in
24 published literature.

25 To overcome these issues, NMA can be performed. Advantages of performing this type of
26 analysis are:

- 27 • It allows the synthesis of data from direct and indirect comparisons without breaking
28 randomisation, to produce measures of treatment effect and ranking of different
29 interventions. If treatment A has never been compared against treatment B head to head,
30 but these two interventions have been compared to a common comparator, then an
31 indirect treatment comparison can use the relative effects of the two treatments versus the
32 common comparator. This is also the case whenever there is a path linking two
33 treatments through a set of common comparators. All the randomised evidence is
34 considered within the same model.
- 35 • For every intervention in a connected network, a relative effect estimate (with its 95%
36 credible intervals (95% CrI) can be estimated versus any other intervention. These
37 estimates provide a useful clinical summary of the results and facilitate the formation of
38 recommendations based on all of the best available evidence, whilst appropriately
39 accounting for uncertainty. Furthermore, these estimates will be used to parameterise
40 treatment effectiveness in the de novo cost-effectiveness modelling.

41 There are 3 key assumptions behind an NMA: similarity, transitivity and consistency.

42 Consistency is the assumption that the direct estimates are equal to the indirect estimates
43 (i.e. that the relative effect of A *versus* C is equal to the relative effect of A *versus* B
44 *versus* C).

45 Similarity across trials is the critical rationale for the consistency assumption to be valid as,
46 by ensuring the clinical characteristics of the trials are similar, we ensure consistency in the
47 data analysis.

1 More specifically, randomisation holds only within individual trials, not across the trials.
2 Therefore, if the trials differ in terms of patient characteristics, measurement and/or definition
3 of outcome, length of follow-up across the direct comparisons, the similarity assumption is
4 violated and this can bias the analysis. Potential sources of heterogeneity arising from trials
5 and attempts made to identify and account for heterogeneity are:

- 6 • Different duration of treatment or study follow-up:
 - 7 ○ The impact of study follow-up on treatment efficacy was assessed to identify if
 - 8 statistically there were differences between long and short duration studies.
 - 9 ○ Where differences were found, or where the Committee believed there were likely to be
 - 10 clinical differences, NMAs were conducted separately for short and long duration
 - 11 studies.
- 12 • Different dosages of pharmacological treatments:
 - 13 ○ These typically showed little variation and were within the dose ranges specified by the
 - 14 British National Formulary (BNF).

15 Transitivity is the assumption that an intervention (A) will have the same efficacy in a study
16 comparing A *versus* B as it will in a study comparing A *versus* C. Another way of looking at it,
17 in terms of the study participants, is that we assume that it is equally likely that any patient in
18 the network could have been given any of the treatments in the network and would have
19 responded to the treatments in the same way (depending on how efficacious the treatments
20 are). This assumption is closely related to similarity in that if participants in a study
21 comparing A *versus* B are not the same as those in a study comparing A *versus* C.

22 As it is the case for ordinary pairwise meta-analysis, NMA may be conducted using either
23 fixed or random effects models. A fixed effects model typically assumes that there is no
24 variation in relative effects across trials for a particular pairwise comparison and any
25 observed differences are solely due to chance. For a random effects model, it is assumed
26 that the relative effects are different in each trial but that they are from a single common
27 distribution. The variance reflecting heterogeneity is often assumed to be constant across
28 trials.

29 For continuous outcomes (e.g. forced expiratory volume), where SEs could not be calculated
30 from the data, we imputed absolute values for them from other studies that reported
31 measures of uncertainty/variance, using the method of Stevens (2011).

32 In a Bayesian analysis, for each parameter the evidence distribution is weighted by a
33 distribution of prior beliefs. Markov Chain Monte Carlo (MCMC) algorithm was used to
34 generate a sequence of samples from a joint posterior distribution of 2 or more random
35 variables and is particularly well adapted to sampling the treatment effects (known as
36 posterior distribution) of a Bayesian network. A non-informative prior distribution was used to
37 maximise the weighting given to the data and to generate the posterior distribution for each
38 log odds ratio (OR), log rate ratio or mean difference (MD) of interest in the networks. We
39 used the median of the distribution as our point estimate and the centiles provided the 95%
40 Credible Intervals (CrI).

41 Non-informative priors were used which were normally distributed with a mean of 0 and
42 standard deviation of 100. However, for dichotomous data, where there was sparse data
43 within the network, we investigated whether the use of informative priors generated from
44 empirical data would give a more stable between-study variance (Turner 2012; Appendix L).

45 For the analyses, a series of 40,000 (100,000 for the multivariate NMA) burn-in simulations
46 were run to allow the posterior distributions to convergence and then a further 100,000
47 simulations were run to produce the outputs. Convergence was assessed by examining the
48 history, autocorrelation and Brooks-Gelman-Rubin plots.

49 Goodness-of-fit of the model was also estimated by using the posterior mean of the sum of
50 the deviance contributions for each item by calculating the residual deviance and deviance

1 information criteria (DIC). If the residual deviance was close to the number of unconstrained
2 data points (the number of trial arms in the analysis) then the model was explaining the data
3 at a satisfactory level. The choice of a fixed or random effects model can be made by
4 comparing their goodness-of-fit to the data.

5 Incoherence in NMA between direct and indirect evidence can be assessed in closed
6 treatment loops within the network. These closed treatment loops are regions within a
7 network where direct evidence is available on at least 3 different treatments that form a
8 closed “circuit” of treatment comparisons (for example A *versus* B, B *versus* C, C *versus* A).
9 If closed treatment loops existed then discrepancies between direct and indirect evidence
10 was assessed for each loop using node-splitting (van Valkenhoef 2016).

11 The outputs of the NMA were:

- 12 • Treatment specific log ORs, log rate ratios, and MDs with their 95% CrI were generated
13 for every possible pairs of comparisons by combining direct and indirect evidence in each
14 network.
- 15 • The probability that each treatment is ranked as the best treatment, based on the
16 proportion of Markov chain iterations in which the treatment effect for an intervention is
17 ranked best, 2nd best and so forth. This was calculated by taking the treatment effect of
18 each intervention compared to placebo and counting the proportion of simulations of the
19 Markov chain in which each intervention had the highest treatment effect.
- 20 • The ranking of treatments (presented as median rank and its 95% CrI).

21 One of the main advantages of the Bayesian approach is that the method leads to a decision
22 framework that supports decision making. The Bayesian approach also allows the probability
23 that each intervention is best for achieving a particular outcome, as well as its ranking, to be
24 calculated.

25 We adapted a model templates for continuous and dichotomous data available from NICE
26 Decision Support UNIT (DSU) technical support document number 2. This model accounts
27 for the within-study correlation between treatment effects induced by multi-arm trials.

28 NMA was considered particularly important for the 2 review questions where it was used
29 because it allows use of indirect evidence to make comparisons between treatments that
30 have not been compared in head-to-head RCTs. The networks in those 2 reviews were
31 mostly compared to a single comparator (namely placebo) and therefore NMA allows us to
32 estimate relative effects between all active treatments. NMA also allows all treatments to be
33 compared to a single comparator, which is useful for health economic analysis that takes a
34 fully incremental approach to determine the most cost-effective treatment out of all
35 treatments under consideration. The primary motivation behind NMA for the chosen review
36 question was that health economic analysis was prioritised for both of those review
37 questions.

38 **4.3.4.2 Data synthesis for diagnostic test accuracy reviews**

39 There are a number of diagnostic test accuracy measures. Sensitivity, specificity, positive
40 and negative likelihood ratios were used as outcomes for diagnostic reviews in this guideline.
41 These diagnostic accuracy parameters (with 95% CI) were obtained from the studies or
42 calculated by the technical team using data from the studies (See Table 5).

43 Sensitivity and specificity are measures of the ability of a test to correctly classify a person as
44 having a disorder or not having a disorder. When Sensitivity is high, a Negative test result
45 rules out the target disorder. When Specificity is high, a Positive test result rules in the target
46 disorder. An ideal test would be both highly sensitive and highly specific, but this is frequently
47 not possible and typically there is a trade-off.

The following definitions were used when summarising the levels of sensitivity or specificity for the Committee:

- High: 90% and above
- Moderate: 75% to 89%
- Low: 74% or below

Positive and negative likelihood ratios are measures of the association between a test result and the target disorder. A positive likelihood ratio greater than one indicates a positive test is associated with having the disorder, whilst a negative likelihood ratio less than one indicates a negative test is associated with not having the disorder.

The following definitions were used when summarising the likelihood ratios for the Committee:

- Very useful test: LR+ higher than 10; LR- lower than 0.1
- Moderately useful test: LR+ 5 to 10; LR- 0.1 to 0.2
- Not a useful test: LR+ lower than 5; LR- higher than 0.2

Table 5: '2 x 2' table for calculation of diagnostic accuracy parameters

	Reference standard positive	Reference standard negative	Total
Index test result positive	True positive (TP)	False positive (FP)	TP+FP (Total number of subjects with positive result in screening tool)
Index test result negative	False negative (FN)	True negative (TN)	FN+TN (Total number of subjects with negative results in screening tool)
Total	TP+FN (Total number of subjects with diagnosis)	FP+TN (Total number of subjects without diagnosis)	TP+FP+FN+Tn=N (Total number of subjects in study)
<p>Note: $Sensitivity = TP / (TP + FN)$ $Specificity = TN / (TN + FP)$ $Positive\ likelihood\ ratio = sensitivity / (1 - specificity)$ $Negative\ likelihood\ ratio = (1 - sensitivity) / specificity$</p>			

4.3.4.3 Data synthesis for clinical prediction and prognostic reviews

For the investigation of pre-specified prognostic/predictive factors, odds ratios (ORs), risk ratios (RRs) or hazard ratios (HRs), with their 95% confidence intervals (95% CIs) were extracted from the papers when reported. As prognostic factors are not attributes that can be randomised (either for practical or ethical reasons), evidence came from observational studies. For this type of review, we looked for studies that took into account possible key confounders as reported in multivariable analyses. The reported measures were therefore adjusted to take into account other characteristics that might explain some of the association between the prognostic factor and the outcome. Studies did this in a pre-specified manner using stratification, or used statistical methods that included confounding variables (such as multivariable logistic regression). Results from these analyses would then indicate which characteristics are most likely to be independent prognostic factors rather than confounding factors only spuriously related to the outcome.

1 4.3.4.4 Data synthesis for prevalence reviews

2 Study results were presented according to the PRISMA guidelines (Preferred Reporting
3 Items for Systematic Reviews and Meta-analyses). Risk factors that were assessed in a
4 multi-variated regression analysis model with adjustment for important confounders were
5 reported. To assist with the ease of interpretation, only results from studies where outcomes
6 were assessed dichotomously were included and reported. Prevalence estimates
7 (proportions) with their 95% confidence intervals were reported or calculated where sufficient
8 data were available.

9 Studies were categorised according to type of outcome and where data were available,
10 results were reported by subgroups pre-specified in the review protocol.

11 4.3.4.5 Data synthesis for qualitative reviews

12 Where possible, a meta-synthesis was conducted to combine qualitative study results. The
13 main aim of the synthesis of qualitative data was to produce a description of the topics that
14 may influence the experience of person with cystic fibrosis, those people important to them
15 and healthcare professionals involved in their care, rather than build new theories or
16 reconceptualise the topic under review. Whenever studies identified a qualitative theme, this
17 was extracted and the main characteristics were summarised. When all themes were
18 extracted from studies, common concepts were categorised and tabulated. This included
19 information on how many studies had contributed to an identified overarching theme.

20 In qualitative synthesis, a theme being reported by different studies more often than other
21 themes does not necessarily mean that it would be more important than those other themes.
22 The aim of qualitative research is to identify new perspectives on a particular topic. Study
23 type and population in qualitative research can differ widely, meaning that themes identified
24 by just one or a few studies can provide important new information for a given topic.
25 Therefore, for the purpose of the qualitative reviews in this guideline, we did not add further
26 studies when they reported the same themes that had already been identified from the same
27 perspectives because the emphasis was on conceptual robustness rather than the
28 quantitative completeness of evidence. This has implications for the types and numbers of
29 studies that are included in the qualitative reviews. Study inclusion continued until no new
30 relevant data could be found regarding a topic that would add to or refute it, a concept
31 referred to in the literature as 'theoretical saturation' (Dixon-Woods 2005).

32 The most relevant evidence in this respect would originate from studies set in the target
33 context of the UK NHS setting. Themes from individual studies were then integrated into a
34 wider context and, when possible, overarching categories of themes with sub-themes were
35 identified. Themes were derived from data presented in individual studies based directly on
36 quotes from interviewees. When themes were extracted, theme names derived from the
37 studies that provided it were used. The names of overarching themes, however, were named
38 by the systematic reviewers.

39 Emerging themes were then placed into a thematic map that presents the relationship
40 between themes and subthemes. The purpose of the map was to show relationships
41 between overarching themes and their subthemes. The mapping part of the review was
42 drafted by a member of the technical team, but the final framework of themes was further
43 shaped and, when necessary, re-classified through discussion with at least one other
44 member of the technical team. The Committee could then draw conclusions from each theme
45 in each setting or country and how they may help in forming recommendations.

1 4.3.5 Appraising the quality of the evidence by outcomes

2 4.3.5.1 GRADE methodology

3 For intervention reviews, the evidence for outcomes from the included RCTs and
4 observational studies were evaluated and presented using GRADE, which was developed by
5 the international GRADE working group. Modified GRADE assessments were also carried
6 out for accuracy measures in diagnostic reviews. For the appraisal of the quality of the
7 evidence from qualitative reviews an adapted GRADE-CERQual (Lewin 2015) approach was
8 used, where CERQual stands for Confidence in the Evidence from Reviews of Qualitative
9 research.

10 The software developed by the GRADE working group (GRADEpro) was used to assess the
11 quality of each outcome, taking into account individual study quality factors and the meta-
12 analysis results. The clinical/economic evidence profile tables include details of the quality
13 assessment and pooled outcome data, where appropriate, an absolute measure of
14 intervention effect and the summary of quality of evidence for that outcome. In this table, the
15 columns for intervention and control indicate summary measures of effect and measures of
16 dispersion (such as mean and standard deviation or median and interquartile range) for
17 continuous outcomes and frequency of events (n/N: the sum across studies of the number of
18 patients with events divided by sum of the number of completers) for binary outcomes.
19 Reporting or publication bias was only taken into consideration in the quality assessment and
20 included in the clinical evidence profile tables if it was apparent.

21 The selection of outcomes for each review question was decided when each review protocol
22 was discussed with the Committee. However, given the nature of most of the review
23 questions included in this guideline (driven by short- or long-term outcomes), the
24 categorisation of outcomes as critical and important did not follow the standard GRADE
25 approach. The outcomes selected for a review question were critical for decision-making in a
26 specific context.

27 The evidence for each outcome in interventional reviews was examined separately for the
28 quality elements listed and defined in Table 6.

29 **Table 6: Description of quality elements in GRADE for intervention studies**

Quality element	Description
Risk of bias (study limitations)	Limitations in the study design and implementation may bias the estimates of the treatment effect. High risk of bias for the majority of the evidence decreases confidence in the estimate of the effect.
Inconsistency	Inconsistency refers to an unexplained heterogeneity of results or findings.
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question, or recommendation made, such that the effect estimate is changed. This is also related to applicability or generalisability of findings.
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of the effect. Imprecision results if the confidence interval includes the clinically important threshold. For qualitative research this can relate to the sufficiency of data within each theme.
Publication bias	Publication bias is a systematic underestimate or an overestimate of the underlying beneficial or harmful effect due to the selective publication of studies.

1 The GRADE toolbox is designed only for RCTs and observational studies, but we adapted
2 the quality assessment elements and outcome presentation for diagnostic accuracy and
3 qualitative studies, subject to data availability.

4 For example, for diagnostic accuracy studies, the GRADE tables were modified to include
5 the most appropriate measures of diagnostic accuracy (sensitivity and specificity) (See Table
6 7). For prognostic factors, an adapted GRADE approach was conducted. This looked at the
7 body of the evidence for each risk factor across studies for 1 outcome (See Table 8).

8 **Table 7: Description of the elements in GRADE and how they are used to assess the**
9 **quality for diagnostic accuracy reviews**

Quality element	Description
Risk of bias ('Study limitations')	Limitations in the study design and implementation may bias the estimates of the diagnostic accuracy. High risk of bias for the majority of the evidence decreases confidence in the estimate of the effect. Diagnostic accuracy studies are not usually randomised and therefore would not be downgraded for study design from the outset and start as high level evidence.
Inconsistency	Inconsistency refers to an unexplained heterogeneity of test accuracy measures such as sensitivity and specificity between studies.
Indirectness	Indirectness refers to differences in study population, differences in index tests across studies, reference standards and outcomes between the available evidence and the review question.
Imprecision	Results are considered imprecise when studies include relatively few patients and the probability to be diagnosed correctly in this group is low. Accuracy measures would therefore have wide confidence intervals around the estimate of the effect.

10 **Table 8: Description of the elements in GRADE and how they are used to assess the**
11 **quality for prognostic reviews**

Quality element	Description
Risk of bias ('Study limitations')	Limitations in the study design and implementation may bias the estimates and the interpretation of the effect of the prognostic risk factor. High risk of bias for the majority of the evidence decreases confidence in the estimate of the effect. Prognostic studies are not usually randomised and therefore would not be downgraded for study design from the outset and start as high level evidence.
Inconsistency	Inconsistency refers to an unexplained heterogeneity between studies looking at the same sign or symptom, resulting in wide variability between ORs, RRs or HRs, with little or no overlap in confidence intervals.
Indirectness	Indirectness refers to any departure from the review protocol, for instance differences in study population or risk factor, that may affect how results can be generalised from the reviewed evidence.
Imprecision	Results are considered imprecise when studies include relatively few patients and also when the number of patients is too low for a multivariable analysis (as a rule of thumb a number of 10 participants per variable). This was assessed by looking at the confidence interval and where it lies in relation to the point estimate of the study.

12 For qualitative studies an adapted GRADE-CERQual (Lewin 2015) approach was used,
13 where CERQual stands for confidence in the evidence from reviews of qualitative research.
14 This looked at the quality of evidence by theme. These themes may have originated from an
15 individual study or may have been identified through a number of individual themes or
16 components of themes across a number of included studies (See Table 9).

1 **Table 9: Description of the elements in the adapted GRADE-CERQual approach used**
2 **to assess qualitative evidence by theme**

Quality element	Description
Risk of bias ('Study limitations')	Limitations in the study design and implementation may bias the interpretation of the qualitative themes that are identified. High risk of bias for the majority of the evidence decreases confidence in the estimate of the effect. Qualitative studies are not usually randomised and therefore would not be downgraded for study design from the outset and start as high level evidence.
Coherence of findings	The extent to which different individual themes or components of themes from studies fit into a wider network of overarching themes. For example, many components (relationship and rapport, clinical experience, information provision) can contribute to an overarching theme of healthcare professional factors in shared decision-making. Even though each individual study may not mention each factor, the overall theme is coherent.
Applicability (or relevance) of evidence	The extent to which the evidence supporting the review finding is applicable to the context specified in the review question. In the case of this guideline, qualitative evidence from the UK was prioritised over and above data from other contexts.
Theme saturation / sufficiency	Theme saturation or sufficiency refers to a similar concept in qualitative research. This refers to whether a theoretical point of theme saturation was achieved, at which point no further citations or observations would provide more insight or suggest a different interpretation of this theme. Individual studies that may have contributed to a theme or subtheme may have been conducted in a manner that by design would have not reached theoretical saturation on an individual study level.

3 The main criteria considered in the rating of these elements are discussed below (see
4 section 4.3.5.2). Footnotes were used to describe reasons for grading a quality element as
5 having serious or very serious problems. The ratings for each component were summed to
6 obtain an overall assessment for each outcome.

7 The main criteria considered in the rating of these elements are discussed below. Footnotes
8 beneath GRADE tables were used to describe reasons for grading a quality element as
9 having serious or very serious limitations. The ratings for each component were summed to
10 obtain an overall assessment for each outcome (See Table 12).

11 **4.3.5.2 Grading the quality of clinical evidence**

12 After results were pooled using data synthesis methods, the overall quality of evidence for
13 each outcome was considered. The following procedure was adopted when using the
14 GRADE approach:

- 15 • An initial quality rating was assigned, based on the study design. RCTs start as 'High' in
16 intervention reviews and observational studies as 'Low'. In diagnostic, prognostic and
17 qualitative reviews, evidence from non-randomised studies start as 'High'.
- 18 • The rating was then downgraded for the specified criteria: risk of bias (study limitations);
19 inconsistency; indirectness; imprecision; and publication bias. These criteria are detailed
20 below. Evidence from observational studies (which had not previously been downgraded)
21 was upgraded if there was a large magnitude of effect or a dose-response gradient, and if
22 all plausible confounding would reduce a demonstrated effect, or suggest a spurious
23 effect when results showed no effect.

24 Each quality element considered to have 'serious' or 'very serious' issues was rated down by
25 1 or 2 points respectively. Value based judgements for relevant interpretation of the levels of
26 quality elements were informed by discussion with the Committee for each review to balance
27 consistency of approach across the guideline and clinical relevance within each review (see

Table 10). The downgraded/upgraded ratings were then summed and the overall quality rating was revised, taking into account the relative contributions from the individual studies within a meta-analysis, where performed. For example, RCTs start as high and the overall quality becomes moderate, low or very low if 1, 2 or 3 points are deducted respectively. The reasons or criteria used for downgrading were specified in the footnotes.

For qualitative reviews, each quality element considered to have 'minor or 'serious' limitations was rated down by 1 or 2 points respectively. A quality assessment of 'Unclear' was added to the list of possible GRADE-CERQual levels. Together with the Committee, it was decided that in qualitative reviews 1 'Unclear' rating did not mean an automatic downgrade of the evidence for this theme. However, 2 'Unclear' ratings were downgraded by 1. Footnotes were not used for the CERQual tables (See Table 11).

Table 10: Levels of quality elements in GRADE for intervention, diagnostic and prognostic reviews

Level	Description
None	There are no serious issues with the evidence.
Serious	The issues are serious enough to downgrade the outcome evidence by 1 level.
Very serious	The issues are serious enough to downgrade the outcome evidence by 2 levels.

Table 11: Levels of quality elements in GRADE for qualitative reviews

Level	Description
No limitations	There are no serious issues with the evidence.
Minor limitations	The issues are serious enough to downgrade the outcome evidence by 1 level.
Serious limitations	The issues are serious enough to downgrade the outcome evidence by 2 levels.
Unclear	There is not enough information available to assess the domain.

Table 12: Overall quality of outcome evidence in GRADE

Level	Description
High	Further research is very unlikely to change our confidence in the estimate of effect.
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
Very low	Any estimate of effect is very uncertain.

The details of the criteria used for each of the main quality elements are discussed further in sections 4.3.5.2.1 to 4.3.5.3.4 below.

4.3.5.3 Risk of bias/ methodological limitations

Intervention studies

For intervention studies, the Cochrane Risk of Bias tool was used for randomised control trials (see appendix H in the NICE guidelines manual 2014).

Bias can be defined as anything that causes a consistent deviation from the truth. Bias can be perceived as a systematic error. The risk of bias for a given study and outcome is associated with the risk of over or underestimation of the true effect. Sources of bias in randomised controlled trials are listed in Table 13).

1 A study with a poor methodological design does not automatically imply high risk of bias; the
2 bias is considered individually for each outcome and it is assessed whether this poor design
3 will impact on the estimation of the intervention effect.

4 **Table 13: Summary of Cochrane risk of bias tool**

Risk of bias	Explanation
Allocation concealment	Those enrolling patients are aware of the group to which the next enrolled patient will be allocated (this is a major problem in 'pseudo' or 'quasi' randomised trials with allocation by, for example, day of week, birth date, chart number).
Lack of blinding	Patient, caregivers, those recording outcomes, those adjudicating outcomes or data analysts are aware of the arm to which patients are allocated.
Incomplete accounting of patients and outcome events	Missing data not accounted for and failure of the investigators to adhere to the intention to treat principle when indicated.
Selective outcome reporting	Reporting of some outcomes and not others on the basis of the results.
Other risks of bias	For example: <ul style="list-style-type: none"> • stopping early for benefit observed in randomised trials, in particular in the absence of adequate stopping rules • use of unvalidated patient-reported outcomes • recruitment bias in cluster randomised trials.

5 For observational studies, quality was assessed using the Newcastle-Ottawa Scale (Wells
6 2008) (see appendix H in the NICE guidelines manual 2014).

7 The risk of bias was derived by assessing the risk of bias across 3 domains – selection,
8 comparability and outcome. Studies are given a rating depending on how they perform on
9 each of the domains. More details about the quality assessment items for observational
10 studies are shown in Table 14.

11 **Table 14: Summary of Newcastle and Ottawa scale**

Risk of bias category	Quality assessment item
Selection	Representativeness of the cohort
	Selection of the non-exposed cohort
	Ascertainment of exposure
	Demonstration that the outcome of interest was not present at the start of the study
Comparability	Comparability of cohorts on the basis of the design or analysis
Outcome	Assessment of outcome
	Was follow-up long enough for outcomes to occur
	Adequacy of follow-up of cohorts

12 **Diagnostic studies**

13 For diagnostic accuracy studies, the Quality Assessment of Diagnostic Accuracy Studies
14 version 2 (QUADAS- 2) checklist was used ([http://www.bristol.ac.uk/social-community-
15 medicine/projects/quadas/quadas-2/](http://www.bristol.ac.uk/social-community-medicine/projects/quadas/quadas-2/)) (see appendix H in the NICE guidelines manual 2014).

16 Risk of bias and applicability in primary diagnostic accuracy studies in QUADAS- 2 consists
17 of 4 domains:

- 18 • patient selection

- 1 • index test
- 2 • reference standard
- 3 • flow and timing.

4 More details about the quality assessment of diagnostic studies are shown in Table 15.

5 **Table 15: Summary of QUADAS-2**

Domain	Patient Selection	Index text	Reference standard	Flow and timing
Description	Describe methods of patient selection: Describe included patients (prior testing, presentation, intended use of index test and setting):	Describe the index test and how it was conducted and interpreted:	Describe the reference standard and how it was conducted and interpreted:	Describe any patients who did not receive the index test(s) and/or reference standard or who were excluded from the 2x2 table (refer to flow diagram): Describe the time interval and any interventions between index test(s) and reference standard:
Signalling questions (yes/no/unclear)	Was a consecutive or random sample of patients enrolled?	Were the index test results interpreted without knowledge of the results of the reference standard?	Is the reference standard likely to correctly classify the target condition?	Was there an appropriate interval between index test(s) and reference standard?
	Was a case-control design avoided?	If a threshold was used, was it pre-specified?	Were the reference standard results interpreted without knowledge of the results of the index test?	Did all patients receive a reference standard?
	Did the study avoid inappropriate exclusions?			Did all patients receive the same reference standard?
				Were all patients included in the analysis?
Risk of bias: (high/low/unclear)	Could the selection of patients have introduced bias?	Could the conduct or interpretation of the index test have introduced bias?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Could the patient flow have introduced bias?
Concerns regarding applicability: (high/low/unclear)	Are there concerns that the included patients do not match the review question?	Are there concerns that the index test, its conduct, or interpretation differ from the review	Are there concerns that the target condition as defined by the reference standard does not match the	

Domain	Patient Selection	Index text	Reference standard	Flow and timing
		question?	review question?	

1 **Prevalence studies**

2 For prevalence studies, the quality was assessed using the checklist created by The Joanna
3 Briggs Institute (The Joanna Briggs Institute, 2014; Munn et al., 2014) (see appendix H in the
4 NICE guidelines manual 2014).

5 The quality was assessed based on answering ‘yes’, ‘no’, ‘unclear’, or “not applicable” to the
6 following questions:

- 7 • Was the sample representative of the target population?
- 8 • Were the study participants recruited in an appropriate way?
- 9 • Was the sample size adequate?
- 10 • Were the study subjects and setting described in detail?
- 11 • Is the data analysis conducted with sufficient coverage of the identified sample?
- 12 • Were objective, standard criteria used for measurement of the condition?
- 13 • Was the condition measured reliably?
- 14 • Was there appropriate statistical analysis?
- 15 • Are all important confounding factors/ subgroups/differences identified and accounted for?
- 16 • Were subpopulations identified using objective criteria?

17 The assessment of the overall quality of the evidence was based on the reviewer’s judgment
18 considering the answers to the questions above. For example, if there were several “no” and
19 “unclear” answers, the quality of the evidence was considered to be low, or if there were
20 some “unclear” answers the quality of the evidence was considered to be moderate.

21 **Prognostic studies**

22 For prognostic studies, the quality was assessed using the checklist created by Hayden et al.
23 (2013) (see appendix H in the NICE guidelines manual 2014).

24 This risk of bias for each risk factor across studies was derived by assessing the risk of bias
25 across 6 domains for each study: study participation, study attrition, prognostic factor
26 measurement, outcome measurement, study confounding, and statistical analysis and
27 reporting, with the last 4 domains being assessed for each outcome. More details about the
28 quality assessment for prognostic studies are shown in Table 16.

29 The assessment of the overall quality of the evidence was based on the reviewer’s judgment
30 considering the assessment of all the 6 domains, for example, if there was a high risk of bias
31 in any domain, the evidence was considered to be of low quality and if there was low risk of
32 bias in all domains, the evidence was considered to be of high quality.

33 **Table 16: Assessment of risk of bias for prognostic studies based on Hayden et al.**
34 **(2013)**

Risk of bias	Explanation
Study participation	Assessment of whether or not there was adequate participation in the study by eligible individuals; if the population and sample were described; if the recruitment and sampling were described and considered appropriate; if inclusion and exclusion criteria were adequately described.
Study attrition	Assessment of whether there was an adequate follow-up rate for study participants; reasons for losses to follow-up were described; the individuals

Risk of bias	Explanation
	lost to follow-up were adequately described; assessment was done whether the ones lost to follow-up differed from the ones who completed the follow-up.
Prognostic factor measurement	Assessment of whether or not a clear description of the prognostic (risk) factor is provided; the method of assessing or measuring the prognostic factor is valid and reliable; and is the same for every participant.
Outcome measurement	Assessment of whether or not a clear definition of the outcome was provided; the measurement or assessment of outcome is valid and reliable; the method and setting of outcome measurement is the same for every participant.
Study confounding	Assessment of whether or not important confounders were adequately measured, described and adjusted for in the analyses.
Statistical analysis and reporting	Assessment of whether or not there is sufficient presentation of data to assess the adequacy of the analytical strategy; the statistical model is adequate; the reporting of results is adequate, clear and not selective.

1 Qualitative studies

2 For qualitative studies, quality was assessed using a checklist for qualitative studies (as
3 suggested in Appendix H in the NICE guidelines manual 2014). This was based on the
4 Critical Appraisal Skills Programme (CASP) checklist for qualitative studies. The quality
5 rating for risk of bias (low, high and unclear) was derived by assessing the risk of bias across
6 6 domains.

7 The evidence was then assessed by theme using GRADE-CERQual across studies as
8 described above and labelled (no limitations, minor limitations, major limitations and unclear),
9 see Table 17.

10 **Table 17: Summary of CASP tool for qualitative studies**

Risk of bias	Explanation
Aim and appropriateness of qualitative evidence.	This refers to an assessment of whether the aims and relevance of the study were clearly described and whether qualitative research methods were appropriate for investigating the research question.
Rigour in study design or validity of theoretical approach	This domain assesses whether the study approach has been clearly described and is based on a theoretical framework (for example ethnography or grounded theory). This does not necessarily mean that the framework has to be explicitly stated, but that at least a detailed description is provided which makes it transparent and reproducible.
Sample selection	The background, the procedure and reasons for the chosen method of selecting participants should be stated. It should also be assessed whether there was a relationship between the researcher and the informant and if so, how this may have influenced the findings that were described.
Data collection	Consideration was given to how well the method of data collection (in-depth interviews, semi-structured interviews, focus groups or observations) was described, whether details were provided and how the data were collected (who conducted the interviews, how long did they last and where did they take place).
Data analysis	For this criterion it is assessed whether sufficient detail is provided about the analytical process and whether it is in accordance with the theoretical approach. For instance, if a thematic analysis was used, it is assessed whether there was a clear description of how the theme was arrived at. Data saturation is also part of this section. This refers to whether a theoretical point of theme saturation was achieved at which point no further citations or observations would provide more insight or suggest a different interpretation of this theme. This could be explicitly stated, or it may be clear from the citations presented that it may have been possible to find more themes.

Risk of bias	Explanation
Results	In relation to this section the reasoning about the results are important, for instance whether a theoretical proposal or framework is provided rather than being restricted to citations / presentation of data.

1 4.3.5.4 Inconsistency / coherence of findings

2 Inconsistency refers to unexplained heterogeneity of results. When estimates of treatment
3 effect measures vary widely across studies (that is, there is heterogeneity or variability in
4 results between studies), this suggests true differences in underlying effects.

5 Heterogeneity in meta-analyses was evaluated. If present, sensitivity and subgroup analyses
6 were performed as pre-specified in the protocols (Appendix D).

7 When heterogeneity existed (chi-squared probability less than 0.1, I-squared inconsistency
8 statistic of greater than 50%, or from visually examining forest plots), but no plausible
9 explanation (for example duration of intervention or different follow-up periods) could be
10 found, the quality of the evidence was downgraded in GRADE by 1 or 2 levels, depending on
11 the extent of inconsistency in the results. When outcomes are derived from a single trial,
12 inconsistency is not an issue for downgrading the quality of evidence. However, 'no
13 inconsistency' is nevertheless used to describe this quality assessment in the GRADE
14 profiles as this is the default option in the GRADEpro software used.

15 For qualitative research, a similar concept to inconsistency is coherence, which refers to the
16 way findings within themes are described and whether they make sense. This concept was
17 used in the quality assessment across studies for individual themes. This does not mean that
18 contradictory data was downgraded automatically, but that it was highlighted and presented,
19 and that reasoning was provided. As long as the themes, or components of themes, from
20 individual studies fit into a theoretical framework, they do not necessarily have to have the
21 same perspective. It should, however, be possible to explain these by differences in context
22 (for example, the views of healthcare professionals might not be the same as those of family
23 members, but they could contribute to the same overarching theme). Coherence was graded
24 across studies with the following labels: coherent, incoherent or unclear.

25 4.3.5.5 Indirectness / applicability or relevance of findings

26 For quantitative reviews, directness refers to the extent to which the populations,
27 intervention, comparisons and outcome measures are similar to those defined in the
28 inclusion criteria for the reviews. Indirectness is important when these differences are
29 expected to contribute to a difference in effect size, or may affect the balance of harms and
30 benefits considered for an intervention.

31 Relevance of findings in qualitative research is the equivalent of indirectness for quantitative
32 outcomes and refers to how closely the aims and context of the studies contributing to a
33 theme reflect the objectives outlined in the review protocol of the guideline question.


34 4.3.5.6 Imprecision / theme saturation or sufficiency

35 For quantitative reviews, imprecision in guidelines concerns whether the uncertainty
36 (confidence interval) around the effect estimate means that it is not clear whether there is a
37 clinically important difference between interventions or not (that is, whether the evidence
38 would clearly support one recommendation or appear to be consistent with several different
39 types of recommendations). Therefore, imprecision differs from the other aspects of evidence
40 quality because it is not really concerned with whether the point estimate is accurate or
41 correct (has internal or external validity). Instead, it is concerned with the uncertainty about
42 what the point estimate actually is. This uncertainty is reflected in the width of the confidence
43 interval.

1 The 95% confidence interval (95% CI) is defined as the range of values within which the
2 population value will fall on 95% of repeated samples, were this procedure to be repeated.
3 The larger the trial, the smaller the 95% CI and the more certain the effect estimate.

4 Imprecision in the evidence reviews was assessed by considering whether the width of the
5 95% CI of the effect estimate was relevant to decision-making, taking each outcome in
6 isolation. This is explained in Figure 2, which considers a positive outcome for the
7 comparison of treatment A versus treatment B. Three decision-making zones can be
8 identified, bounded by the thresholds for clinical importance (minimal important difference,
9 MID) for benefit and for harm. The MID for harm for a positive outcome means the threshold
10 at which drug A is less effective than drug B by an amount that is clinically important to
11 patients (favours B).

Figure 2: Illustration of precise, imprecise and very imprecise evidence based on the confidence interval of outcomes in forest plots



12 When the confidence interval of the effect estimate is wholly contained in 1 of the 3 zones
13 (for example clinically important benefit), we are not uncertain about the size and direction of
14 effect (whether there is a clinically important benefit, or the effect is not clinically important, or
15 there is a clinically important harm), so there is no imprecision.

16 When a wide confidence interval lies partly in each of 2 zones, it is uncertain in which zone
17 the true value of effect estimate lies and therefore there is uncertainty over which decision to
18 make (based on this outcome alone). The confidence interval is consistent with 2 possible
19 decisions and so this is considered to be imprecise in the GRADE analysis and the evidence
20 is downgraded by 1 level ('serious imprecision').

21 If the confidence interval of the effect estimate crosses into 3 zones, this is considered to be
22 very imprecise evidence because the confidence interval is consistent with 3 possible clinical
23 decisions and there is therefore a considerable lack of confidence in the results. The
24 evidence is therefore downgraded by 2 levels in the GRADE analysis ('very serious
25 imprecision').

26 Implicitly, assessing whether the confidence interval is in, or partially in, a clinically important
27 zone, requires the Committee to estimate an MID or to say whether they would make
28 different decisions for the 2 confidence limits.

29 Minimally Important Differences

30 The literature was searched for established minimally important differences (MIDs) for the
31 selected outcomes in the evidence reviews, such as symptom measurement tools. MIDs
32 specific to each review are reported in the review protocol (Appendix D), though the following
33 MIDs were used consistently throughout the guideline:

- 1 • For CF-QOL a published MID was used (European Medicines Agency 2012) which was a
2 difference of 5 points.
- 3 • For time to next exacerbation the Committee agreed that any change was considered to
4 be clinically important, so the MID threshold was one on a ratio scale.
- 5 • For adverse events that led to discontinuation of treatment the Committee agreed that any
6 change was considered to be clinically important, so the MID threshold was one on a ratio
7 scale.
- 8 • For mortality the Committee agreed that any change was considered to be clinically
9 important, so the MID threshold was one on a ratio scale.

10 Finally, if no published or acceptable MIDs were identified, the Committee considered
11 whether it was clinically acceptable to use the GRADE default MID to assess imprecision.
12 For binary outcomes clinically important thresholds for a risk ratio of 0.8 and 1.25
13 respectively were used (due to the statistical distribution of this measure this means that this
14 is not a symmetrical interval). This default MID was used for all the binary outcomes in the
15 interventions' evidence reviews as a starting point and decisions on clinical importance were
16 then considered based on the absolute risk difference. For continuous outcomes GRADE
17 default MIDs were half of the median standard deviation of the control group.

18 The same principle was used for prognostic factors, for example using the default MID as a
19 starting point for the Committee discussion, to assess whether the size of the outcome effect
20 would be large enough to be meaningful in clinical practice.

21 In diagnostic accuracy measures, it was first considered whether sensitivity, specificity,
22 positive likelihood ratios or negative likelihood ratios would be given more weight in the
23 decision-making process. If one measure was given more importance than the other, then
24 imprecision was rated on this statistical measure using the following MID thresholds:

- 25 • Sensitivity and specificity
 - 26 ○ High: >90%
 - 27 ○ Moderate: 75-90%
 - 28 ○ Low: <75%
- 29 • Positive likelihood ratio:
 - 30 ○ Very useful test: >10
 - 31 ○ Moderately useful test: 5-10
 - 32 ○ Not a useful test: <5
- 33 • Negative likelihood ratio:
 - 34 ○ Very useful test: <0.1
 - 35 ○ Moderately useful test: 0.1 to 0.2
 - 36 ○ Not a useful test: >0.2

37 Theme saturation or sufficiency refers to a similar concept in qualitative research. This refers
38 to whether a theoretical point of theme saturation was achieved, at which point no further
39 citations or observations would provide more insight or suggest a different interpretation of
40 this theme. As already highlighted in a previous section on qualitative reviewing methods, it
41 is not equivalent to the number of studies contributing to a theme, but rather to the depth of
42 data and whether sufficient quotes or observations were provided that could underpin these
43 findings.

44 **4.3.5.7 NMA quality appraisal**

45 The use of GRADE to assess the quality of studies addressing a particular review question
46 for pairwise comparisons of interventions is relatively established. However, the use of
47 GRADE to assess the quality of evidence across a NMA is still a developing methodology.

1 While most criteria for pairwise meta-analyses still apply, it is important to adapt some of the
2 criteria to take into consideration additional factors, such as how each 'link' or pairwise
3 comparison within the network applies to the others. As a result, we used the following
4 adapted GRADE approach for appraising the quality of NMA.

5 **Table 18: Rationale for downgrading quality of evidence in NMAs**

GRADE criteria	Example reasons for downgrading quality
Risk of bias	Risk of bias was assessed in accordance with GRADE, as specified in 'The guidelines manual (2012)'. This includes limitations in the design or execution of the study, including concealment of allocation, blinding, loss to follow up (these can reduce the quality rating).
Inconsistency	Evidence of any inconsistency between the direct and indirect estimates of effect was assessed using the residual deviance, deviance information criterion and the statistic tau; outcome was downgraded if tau > 0.5
Indirectness	The extent to which the available evidence fails to address the specific review question (this can reduce the quality rating). This may be in relation to the setting, population, outcomes, interventions or study designs used in the evidence base. Evidence was only downgraded if this was likely to have an impact on the overall rankings (that is, within smaller networks where there is a lack of evidence or within larger networks in large trials which show large reductions in outcomes).
Imprecision	This is considered to be present when there is uncertainty around the estimate of effect, and reflects the confidence in, or 'credibility' of, the estimate of effect. It is assessed based on the overall distribution of the rankings, such that evidence was downgraded if no interventions had rank credible intervals $\leq 33\%$ of total distribution of comparators.

6 **4.3.5.8 Assessing clinical significance**

7 **Intervention and prognostic reviews**

8 The Committee assessed the evidence by outcome. To facilitate this, where possible, binary
9 outcomes were converted into absolute risk differences (ARDs) using GRADEpro software:
10 the median control group risk across studies was used to calculate the ARD and its 95% CI
11 from the pooled risk ratio. For continuous outcomes, the mean difference between the
12 intervention and control arm of the trail was calculated. This was then assessed in relation to
13 the default MID (0.5 times the median control group standard deviation).

14 The clinical significance of a treatment effect or prognostic factor was evaluated as a
15 combination of the minimally / clinically important difference (MID) thresholds and statistical
16 significance / the null hypothesis value (zero for continuous outcomes and 1 for RRs, ORs
17 and HRs):

- 18 • If the point estimate for a treatment effect / prognostic factor exceeded the MID and the
19 95% CI did not include the null hypothesis value then the result was considered to be
20 "clinically significant"
- 21 • If the point estimate for a treatment effect / prognostic factor did not exceed the MID then
22 the result was not considered to be "clinically significant"

23 **Diagnostic reviews**

24 The clinical usefulness of a test for diagnosis was determined based on either sensitivity,
25 specificity, positive likelihood ratio or negative likelihood ratio, depending on what the
26 Committee believed was the most important – correctly identifying if a patient had the target
27 disorder (ruling in) or correctly identifying if a patient did not have the target disorder (ruling
28 out).

1 The value of the point estimate within the different MID thresholds for sensitivity, specificity,
2 positive likelihood ratio or negative likelihood ratio were used to determine clinical
3 usefulness.

4 **Qualitative reviews**

5 For themes stemming from qualitative findings, clinical significance was decided upon by the
6 Committee taking into account the generalisability of the context from which the theme was
7 derived and whether it was convincing enough to support or warrant a change in current
8 practice, as well as the evidence quality.

9 **4.3.6 Evidence statements**

10 Evidence statements are summary statements that are presented after the GRADE profiles,
11 summarising the key features of the clinical evidence presented. The wording of the
12 evidence statements reflects the certainty or uncertainty in the estimate of effect. The
13 evidence statements are presented by outcome or theme and encompass the following key
14 features of the evidence:

- 15 • the quality of the evidence (GRADE rating)
- 16 • the number of studies and the number of participants for a particular outcome
- 17 • a brief description of the participants
- 18 • the clinical significance of the effect and an indication of its direction (for example, if a
19 treatment is clinically significant [beneficial or harmful] compared with another, or whether
20 there is no clinically significant difference between the tested treatments).

21 **4.3.7 Evidence of cost effectiveness**

22 The aims of the health economic input to the guideline were to inform the Guideline
23 Committee of potential economic issues related to the diagnosis and management of cystic
24 fibrosis to ensure that recommendations represented a cost-effective use of healthcare
25 resources. Health economic evaluations aim to integrate data on healthcare benefits (ideally
26 in terms of quality-adjusted life-years (QALYs) with the costs of different care options. In
27 addition, the health economic input aimed to identify areas of high resource impact;
28 recommendations which – while nevertheless cost-effect – might have a large impact on
29 CCG or Trust finances and so need special attention.

30 **4.3.7.1 Undertaking new health economic analysis**

31 As well as reviewing the published economic literature, as described above, new economic
32 analysis was undertaken by the Health Economist in selected areas. The following priority
33 areas for de novo economic analysis were agreed by the Committee after formation of the
34 review questions and consideration of the available health economic evidence:

- 35 • immunomodulatory agents in the management of lung disease,
- 36 • antimicrobial regimens in suppressing chronic pulmonary disease,
- 37 • configuration of services to minimise the risk of cross-infection.

38 A costing tool was also developed for the review question relating to models of care, where
39 little clinical evidence was uncovered. It was thought that the Committee may wish to make
40 recommendations that would lead to a high resource impact, although current practice was
41 recommended.

42 The methods and results of de novo economic analyses are reported in Appendix K. When
43 new economic analysis was not prioritised, the Committee made a qualitative judgement
44 regarding cost effectiveness by considering expected differences in resource and cost use
45 between options, alongside clinical effectiveness evidence identified from the clinical

1 evidence review. Cost descriptions used to aid considerations of cost effectiveness are also
2 reported in Appendix K.

3 **4.3.7.2 Cost effectiveness criteria**

4 NICE's report Social value judgements: principles for the development of NICE guidance
5 sets out the principles that Committees should consider when judging whether an
6 intervention offers good value for money. In general, an intervention was considered to be
7 cost effective if either of the following criteria applied (given that the estimate was considered
8 plausible):

- 9 • the intervention dominated other relevant strategies (that is, it was both less costly in
10 terms of resource use and more clinically effective compared with all the other relevant
11 alternative strategies), or;
- 12 • the intervention cost less than £20,000 per QALY gained compared with the next best
13 strategy, or;
- 14 • the intervention provided clinically significant benefits at an acceptable additional cost
15 when compared with the next best strategy.

16 The Committee's considerations of cost-effectiveness are discussed explicitly in the
17 'Consideration of economic benefits and harms' section of the relevant sections.

18 **4.4 Developing recommendations**

19 **4.4.1 Guideline recommendations**

20 Over the course of the guideline development process, the Committee was presented with:

- 21 • evidence tables of the clinical and economic evidence reviewed from the literature: all
22 evidence tables are in Appendix H and economic evidence tables are in Appendix L
- 23 • summary of clinical and economic evidence and quality assessment (as presented in
24 Chapters 5 to 11)
- 25 • forest plots (Appendix I)
- 26 • a description of the methods and results of the cost-effectiveness analysis undertaken for
27 the guideline (Appendix K).

28 Recommendations were drafted on the basis of the group's interpretation of the available
29 evidence, taking into account the balance of benefits, harms and costs between different
30 courses of action. This was either done formally, in an economic model, or informally. Firstly,
31 the net benefit over harm (clinical effectiveness) was considered, focusing on the critical
32 outcomes, although most of the reviews in the guideline were outcome driven. When this
33 was done informally, the group took into account the clinical benefits and harms when one
34 intervention was compared with another. The assessment of net benefit was moderated by
35 the importance placed on the outcomes (the group's values and preferences) and the
36 confidence the group had in the evidence (evidence quality). Secondly, the group assessed
37 whether the net benefit justified any differences in costs.

38 When clinical and economic evidence was of poor quality, conflicting or absent, the group
39 drafted recommendations based on their expert opinion. The considerations for making
40 consensus-based recommendations include the balance between potential harms and
41 benefits, the economic costs or implications compared with the economic benefits, current
42 practices, recommendations made in other relevant guidelines, patient preferences and
43 equality issues. The group also considered whether the uncertainty was sufficient to justify
44 delaying making a recommendation to await further research, taking into account the
45 potential harm of failing to make a clear recommendation.

1 The wording of recommendations was agreed by the group and focused on the following
2 factors:

- 3 • the actions healthcare professionals need to take,
- 4 • the information readers of the guideline need to know,
- 5 • the strength of the recommendation (for example the word 'offer' was used for strong
6 recommendations and 'consider' for weak recommendations),
- 7 • the involvement of patients (and their carers if needed) in decisions about treatment and
8 care,
- 9 • consistency with NICE's standard advice on recommendations about drugs, waiting times
10 and ineffective intervention.

11 The main considerations specific to each recommendation are outlined in the
12 'Recommendations and link to evidence' sections within each chapter.

13 **4.4.2 Research recommendations**

14 When areas were identified for which good evidence was lacking, the Committee considered
15 making recommendations for future research. Decisions about inclusion were based on
16 factors such as:

- 17 • the importance to patients or the population,
- 18 • national priorities,
- 19 • potential impact on the NHS and future NICE guidance,
- 20 • ethical and technical feasibility.

21 **4.5 Validation process**

22 This guidance is subject to a 6-week public consultation and feedback as part of the quality
23 assurance and peer review of the document. All comments received from registered
24 stakeholders are responded to in turn and posted on the NICE website at publication.

25 **4.6 Updating the guideline**

26 Following publication, and in accordance with the NICE guidelines manual, NICE will
27 undertake a review of whether the evidence base has progressed significantly to alter the
28 guideline recommendations and warrant an update.

29 **4.7 Disclaimer**

30 Healthcare providers need to use clinical judgement, knowledge and expertise when
31 deciding whether it is appropriate to apply guidelines. The recommendations cited here are a
32 guide and may not be appropriate for use in all situations. The decision to adopt any of the
33 recommendations cited here must be made by practitioners in light of individual patient
34 circumstances, the wishes of the patient, clinical expertise and resources.

35 The NGA disclaims any responsibility for damages arising out of the use or non-use of these
36 guidelines and the literature used in support of these guidelines.

37 **4.8 Funding**

38 The NGA was commissioned by the National Institute for Health and Care Excellence (NICE)
39 to undertake the work on this guideline.

40

5 Diagnosis of cystic fibrosis

Review question: In infants, children, young people and adults (including those that have undergone new-born screening) when should cystic fibrosis be suspected?

5.1 Introduction

Today, most people with cystic fibrosis will be diagnosed at birth as part of the national Newborn Screening Programme using the blood spot immunoreactive trypsin test. Screening was introduced UK-wide in 2006 and so there remains a cohort of young people and adults with cystic fibrosis who have not been screened and not yet have been diagnosed though a clinical assessment. Although highly successful, the Programme is not able to screen for all cystic fibrosis associated genetic variants and so some infants will inevitably remain undiagnosed. It should be noted that over 2,000 mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene have been identified. Many of these variants are rare and have not been shown to lead to clinical disease and therefore their clinical significance can be unclear. Of these 2,000 variants, around 200 have been shown to lead to cystic fibrosis disease although only common disease-associated variants are tested for as part of the Newborn Screening Programme.

Additionally, infants may not undergo newborn screening due to parent's wishes or if newborn screening is not carried out in the country of birth. People with atypical manifestations of cystic fibrosis can reach adulthood undiagnosed and therefore untreated when clinical intervention would be beneficial.

Given the potential for people with cystic fibrosis to remain undiagnosed at all stages in life we aim with this review question to determine what lists of symptoms may indicate a possibility of cystic fibrosis and so warrant further investigation.

5.2 Description of clinical evidence

The aim of this review was to support health care professionals in identifying cystic fibrosis even in people who have been through new-born screening.

In this review, our index test were clinical symptoms and signs, including:

- Respiratory features (including recurrent infection, chest x-ray evidence of chronic disease)
- Faltering growth
- Symptoms of malabsorption
- Azoospermia
- Acute pancreatitis
- Meconium ileus (in infants).

The Committee agreed DIOS is known to be a symptom unique to cystic fibrosis, and therefore there was no need to include it in the review. We looked for prospective or retrospective cohort studies to identify diagnostic or prognostic factors, but no relevant studies were found. Therefore we looked for observational studies that reported the prevalence of cystic fibrosis among people presenting with one of the symptoms of interest.

As no diagnostic or prognostic data were finally included in the review, a GRADE approach (as specified in the protocol) was no longer deemed appropriate. Therefore for this review the quality appraisal of the evidence has been conducted by study, and not by outcome.

For full details of the protocol see Appendix D.

1 **5.2.1 Respiratory symptoms**

2 Four observational studies were identified, 2 prospective (Ooi 2012, Seer 1997) and 2
3 retrospective (Hubert 2014, Grimaldi 2015).

4 Sample sizes ranged from 72 to 601, and the studies were conducted in Canada (Ooi 2012,
5 Seer 1997) and France (Hubert 2014, Grimaldi 2015).

6 **5.2.2 Faltering growth**

7 No studies were identified.

8 **5.2.3 Symptoms of malabsorption**

9 No studies were identified.

10 **5.2.4 Azoospermia**

11 One observational study was identified (Ooi 2012).

12 Sample size was 92 and it was conducted in Canada.

13 **5.2.5 Acute pancreatitis**

14 Two observational studies were identified (Lucidi 2011, Ooi 2012).

15 Sample sizes ranged from 44 to 78, and the studies were conducted in Canada (Ooi 2012)
16 and Italy (Lucidi 2011).

17 **5.2.6 Meconium ileus**

18 No studies were identified.

19 A summary of the included studies is presented in Table 19. See also study selection flow
20 chart in Appendix F, excluded studies list in Appendix H, and study evidence tables in
21 Appendix G.

22 **5.3 Summary of included studies and results**

23 A summary of the studies that were included in this review and the results is presented in
24 Table 19.

25

1 Table 19: Summary of included studies and results

Study	Clinical symptoms & signs and reference test	Population	Findings	Comments	Quality
Respiratory symptoms					
Hubert 2004 (France) Retrospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> • bronchiectasis • <i>Bronchiectasis defined as chronic mucopurulent sputum production and recurrent lower respiratory tract infection, were confirmed by high-resolution CT.</i> <p>Reference test:</p> <ul style="list-style-type: none"> • sweat chloride test (ST) • <i>Thresholds:</i> • <i>Diagnosis of CF: >60 mmol/l.</i> • <i>Suggestive, but not diagnostic of CF: 40 to 60 mmol/l.</i> 	<p>N=601</p> <p>Adults referred for diffuse bronchiectasis from 1992 to 2001</p> <p>Mean age (range): 31 years (18 to 56)</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> • Confirmed CF diagnosis: n=37; 6.16% • Borderline CF diagnosis: n=9; 1.50% • Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> • It is unknown whether these patients underwent newborn screening, but seems unlikely as newborn screening was implemented in France in 2002 (Grimaldi 2015) • Two sweat test were performed for each patient. • Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> • Very low
Grimaldi 2015 (France) Retrospective, observational, multicentre study	<p>Symptoms:</p> <ul style="list-style-type: none"> • asthma, • chronic cough, • lower airway infections, • bronchiectasis • <i>Definition for symptoms not provided.</i> <p>Reference test:</p> <ul style="list-style-type: none"> • sweat chloride test (ST) • <i>Thresholds for patients >6 months:</i> • <i>Positive ST: ≥60 mmol/l.</i> • <i>Intermediate ST: 40 to 59 mmol/l.</i> • <i>Negative ST: ≤39 mmol/l.</i> • <i>Thresholds for infants up to 6 months:</i> • <i>Positive ST: ≥60 mmol/l.</i> 	<p>N=502</p> <p>Infants and children who had a negative CF newborn screening presenting respiratory symptoms</p> <p>Mean age ± SD (range): 36±28 months (1 month to 10 years)</p> <ul style="list-style-type: none"> • Asthma: n=358 • Chronic cough: n=263 • Lower airway infections: n=212 	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> • In children with asthma: n=1; 0.3% • In children with chronic cough: n=4; 1.5% • In children with lower airway infections: n=4; 1.8% • In children with bronchiectasis: n=2; 5.7% • Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> • All children had a negative CF newborn screening. • Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> • Very low

Study	Clinical symptoms & signs and reference test	Population	Findings	Comments	Quality
	<ul style="list-style-type: none"> Intermediate ST: 29 to 59 mmol/l. Negative ST: ≤29 mmol/l. 	<ul style="list-style-type: none"> Bronchiectasis: n=35 			
Ooi 2012 (Canada) Prospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> Idiopathic chronic sinopulmonary disease <p><i>Idiopathic sinopulmonary disease defined as recurrent or chronic sinusitis (including sinusoidal pain, nasal discharge, postnasal drip), nasal polyps, recurrent or chronic bronchitis, recurrent pneumonia and/or bronchiectasis for at least 6 months. All enrolled subjects with sinopulmonary disease had three or more of these symptoms.</i></p> <p>Reference test:</p> <ul style="list-style-type: none"> sweat chloride test (ST) cystic fibrosis transmembrane conductance regulator gene (CFTR) <p><i>ST thresholds for the diagnosis of CF according to European consensus recommendations.</i></p> <p><i>Extensive CFTR genotyping according to European diagnostic process (interpretation of results from genotyping was based on the number of mutations identified).</i></p>	<p>N=72</p> <p>Undiagnosed individuals with single organ manifestations of CF were enrolled between 1994 and 2008</p> <p>Mean age ± SD (range): 38.5±15.9 (9.9 to 66.7) years</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> Classic CF: n=14; 19.4% CFTR dysfunction: n=3; 4.2% Inconclusive: n=1; 1.4% Unlikely diagnosis: n=54; 75.0% Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> It is unknown whether these patients underwent newborn screening, Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> Very low
Seear 1997 (Canada) Prospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> Chronic cough No definition given. But just children with a history of >3 months of productive cough, of unknown cause, were included <p>Reference test:</p>	<p>N=81</p> <p>Children with a history of >3 months of productive cough</p> <p>Age: not reported</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> Diagnosis of CF: n=1; 1.23% Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> It is unknown whether these patients underwent newborn screening, Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> Very low

Study	Clinical symptoms & signs and reference test	Population	Findings	Comments	Quality
	<ul style="list-style-type: none"> sweat chloride test (ST) <i>Thresholds for diagnosis of CF not reported.</i> 				
Faltering growth					
No studies were identified.					
Symptoms of malabsorption					
No studies were identified.					
Azoospermia					
Ooi 2012 (Canada) Prospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> Infertility due to obstructive azoospermia <i>A diagnosis of obstructive azoospermia (congenital unilateral or bilateral absence of vas deferens) was confirmed by physical examination, transrectal ultrasound and evidence of azoospermia on two separate occasions.</i> <p>Reference test:</p> <ul style="list-style-type: none"> sweat chloride test (ST) cystic fibrosis transmembrane conductance regulator gene (CFTR) <i>ST thresholds for the diagnosis of CF according to European consensus recommendations.</i> <i>Extensive CFTR genotyping according to European diagnostic process (interpretation of results from genotyping was based on the number of mutations identified).</i> 	<p>N=92</p> <p>Undiagnosed individuals with single organ manifestations of CF were enrolled between 1994 and 2008</p> <p>Mean age ± SD (range): 34.8±5.3 (25.4 to 56.6) years</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> Classic CF: n=19; 20.7% CFTR dysfunction: n=21; 22.8% Inconclusive: n=9; 9.8% Unlikely diagnosis: n=43; 46.7% Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> It is unknown whether these patients underwent newborn screening, Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> Very low
Acute pancreatitis					

Study	Clinical symptoms & signs and reference test	Population	Findings	Comments	Quality
Lucidi 2011 (Italy) Retrospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> acute recurrent pancreatitis <i>Acute recurrent pancreatitis defined as 2 or more separate documented episodes of acute pancreatitis with serum amylase and/ or lipase levels at least 3 times the upper reference limit</i> <p>Reference tests:</p> <ul style="list-style-type: none"> sweat chloride test (ST), genetic test (CFTR mutation) <p><i>ST thresholds for diagnosis of CF not reported.</i></p> <p><i>CFTR genotyping diagnostic process not reported.</i></p>	<p>N=78</p> <p>Paediatric population affected by acute recurrent pancreatitis between 2003 and 2008</p> <p>Mean age \pm SD (range): 8.8\pm5.1 years (4 months to 18 years)</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> Diagnosis of CF with ST: n=1; 1.3% Borderline diagnosis of CF with ST: n=7; 9% Diagnosis of CF by the detection of 2 CF-causing mutations: n=1 (percentage cannot be calculated as the study does not report for how many people there was data available) CFTR mutation: 39.6% (data available for n=53) Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> 42.3% (n=33) patients had a positive family history of chronic pancreatitis/ cystic fibrosis and/ or positive genetic testing and/ or altered sweat test. However it is unknown whether all these patients underwent newborn screening. Critical confounders not taken into consideration. 	Very low
Ooi 2012 (Canada) Prospective, observational study	<p>Symptom:</p> <ul style="list-style-type: none"> Idiopathic recurrent, acute or chronic pancreatitis <p><i>A diagnosis of idiopathic recurrent acute pancreatitis was accepted following at least two episodes of abdominal pain associated with raised serum amylase and/or lipase (more than two times the upper limit of the reference range), and/or imaging evidence of acute pancreatitis such as pancreatic oedema, haemorrhage or necrosis. Patients with chronic pancreatitis had chronic pain in association with pancreatic</i></p>	<p>N=44</p> <p>Undiagnosed individuals with single organ manifestations of CF were enrolled between 1994 and 2008</p> <p>Mean age \pm SD (range): 24.3\pm13.2 (7.9 to 59.9) years</p>	<p>Clinical diagnosis of CF</p> <ul style="list-style-type: none"> Classic CF: n=2; 4.5% CFTR dysfunction: n=6; 13.6% Inconclusive: n=1; 2.3% Unlikely diagnosis: n=35; 79.6% Sensitivity, specificity, LR+, LR-, PPV and NPV not reported and not calculable. 	<ul style="list-style-type: none"> It is unknown whether these patients underwent newborn screening, Critical confounders not taken into consideration. 	<ul style="list-style-type: none"> Very low

Study	Clinical symptoms & signs and reference test	Population	Findings	Comments	Quality
	<p><i>calcifications and/or characteristic ductal changes.</i></p> <p>Reference test:</p> <ul style="list-style-type: none"> • sweat chloride test (ST) • cystic fibrosis transmembrane conductance regulator gene (CFTR) • <i>ST thresholds for the diagnosis of CF according to European consensus recommendations.</i> • <i>Extensive CFTR genotyping according to European diagnostic process (interpretation of results from genotyping was based on the number of mutations identified).</i> 				
Meconium ileus					
No studies were identified					

1 Abbreviations: CF: cystic fibrosis; CFTR: CF transmembrane conductance regulator; LR: likelihood ratio; NPV: negative predictive value; PPV: positive predictive value; SDS: standardized by age and gender; ST: sweat test

3

4

1 5.4 Clinical evidence profile

2 See summary of results in Table 19.

3 5.5 Economic evidence

4 This review question is not relevant for economic analysis because it does not involve a
5 decision between alternative courses of action.

6 No economic evaluations to identify the clinical manifestations suggestive of cystic fibrosis
7 were identified in the literature search conducted for this guideline. Full details of the search
8 and economic article selection flow chart can be found in Appendix E and F, respectively.

9 5.6 Evidence statements

10 5.6.1 Respiratory symptoms

11 Very low quality evidence from 1 prospective observational study showed that among 81
12 children with a history >3 months of productive cough 1.23% (n=1) had a diagnosis of cystic
13 fibrosis (by means of sweat test, thresholds not reported). The study does not report whether
14 these children had undergone newborn screening.

15 Very low quality evidence from 1 retrospective observational study showed that among 502
16 infants and children (age range: 1 month to 10 years) who had a negative cystic fibrosis
17 newborn screening presenting respiratory symptoms:

- 18 • 0.3% (n=1; N=358) of the children presenting with asthma had a diagnosis of cystic
19 fibrosis (sweat test ≥ 60 mmol/l)
- 20 • 1.5% (n=4; N=263) of the children presenting with chronic cough had a diagnosis of cystic
21 fibrosis (sweat test ≥ 60 mmol/l)
- 22 • 1.8% (n=4; N=212) of the children presenting with lower airway infections had a diagnosis
23 of cystic fibrosis (sweat test ≥ 60 mmol/l)
- 24 • 5.7% (n=2; N=35) of the children presenting with bronchiectasis had a diagnosis of cystic
25 fibrosis (sweat test ≥ 60 mmol/l).

26 Very low quality evidence from 1 retrospective observational study showed that among 601
27 adults (mean age: 31 years) referred for diffuse bronchiectasis:

- 28 • 6.16% (n=37) had confirmed diagnosis of cystic fibrosis (sweat test >60 mmol/l)
- 29 • 1.50% (n=9) had borderline diagnosis of cystic fibrosis (sweat test 40 to 60 mmol/l).

30 The study does not report whether these people had undergone newborn screening, but
31 seems unlikely as newborn screening was implemented in France in 2003.

32 Very low quality evidence from 1 prospective observational study showed that among 72
33 children, young people and adults (mean age; range: 34.8 years; 9.9 to 66.7 years) with
34 idiopathic chronic sinopulmonary disease:

- 35 • 19.4% (n=14) had a diagnosis of classic cystic fibrosis based on sweat test (European
36 Consensus Recommendations)
- 37 • 4.2% (n=3) had a CFTR abnormality (based on extensive CFTR genotyping, and
38 according to European Guidelines). This is also known as non-classic or atypical cystic
39 fibrosis
- 40 • 1.4% (n=1) had an inconclusive diagnosis.

41 The study does not report whether these people had undergone newborn screening.

1 **5.6.2 Faltering growth**

2 No evidence was found for this sign.

3 **5.6.3 Symptoms of malabsorption**

4 No evidence was found for this symptom.

5 **5.6.4 Azoospermia**

6 Very low quality evidence from 1 prospective observational study showed that among 92
7 adult men (mean age; range: 34.8 years; 25.4 to 56.6 years) with infertility due to obstructive
8 azoospermia:

- 9 • 20.7% (n=19) had a diagnosis of classic cystic fibrosis based on sweat test (European
10 Consensus Recommendations)
- 11 • 22.8% (n=21) had a CFTR abnormality
- 12 • 9.8% (n=9) had an inconclusive diagnosis.

13 The study does not report whether these men had undergone newborn screening.

14 **5.6.5 Acute pancreatitis**

15 Very low quality from 1 retrospective observational study with 78 infants, children and young
16 people (age range: 4 months to 18 years) affected by acute pancreatitis:

- 17 • 1.3% (n=1) had a diagnosis of cystic fibrosis based on sweat test (thresholds not
18 reported)
- 19 • 9% (n=7) had a borderline diagnosis of cystic fibrosis based on sweat test (thresholds not
20 reported);
- 21 • 39.6% had a single CFTR mutation on genetic testing
- 22 • n=1 had a diagnosis of cystic fibrosis based on the detection of 2 cystic fibrosis-causing
23 mutations (the study included 78 children and young people, but it is unknown for how
24 many of them genetic testing was available)

25 The study does not report whether these people had undergone newborn screening.

26 Very low quality evidence from 1 prospective observational study showed that among 44
27 children, young people and adults (mean age; range: 24.3 years; 7.9 to 59.9 years) with
28 idiopathic recurrent, acute or chronic pancreatitis:

- 29 • 4.5% (n=2) had a diagnosis of classic cystic fibrosis based on sweat test (European
30 Consensus Recommendations)
- 31 • 13.6% (n=6) had a CFTR abnormality
- 32 • 2.3% (n=1) had an inconclusive diagnosis.

33 The study does not report whether these people had undergone newborn screening.

34 **5.6.6 Meconium ileus**

35 No evidence was found for symptoms in infants.

36 **5.6.7 Economic evidence statements**

37 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

1 **5.7 Evidence to recommendations**

2 The aim of this review was to support health care professionals in identifying cystic fibrosis
3 even in people who have been through new-born screening.

4 The Committee chose clinical diagnosis of cystic fibrosis as a critical outcome for this
5 evidence review. Sensitivity, specificity, positive predictive value, negative predictive value,
6 positive likelihood ratio and negative likelihood ratio were rated as important outcomes.

7 **5.7.1 Consideration of clinical benefits and harms**

8 The Committee agreed that a definition of cystic fibrosis was needed in order to emphasise
9 that cystic fibrosis is a syndrome rather than a disease.

10 There are a number of investigations (such as sweat test or genetic test) that can be done in
11 the event that cystic fibrosis is suspected, but there is no gold standard as such. These tests
12 are useful to confirm cystic fibrosis, but they cannot completely exclude cystic fibrosis.
13 Ultimately, the diagnosis can be made based on the clinical signs and symptoms alone, even
14 if tests results are negative. They discussed the emotional implications of being diagnosed
15 with cystic fibrosis with a negative sweat test and cystic fibrosis gene mutations that have
16 been conclusively shown to be disease causing.

17 The Committee discussed the limitations of the available evidence. None of the studies
18 reported diagnostic accuracy data. They agreed the usefulness of prevalence data reported
19 in the studies was very limited as studies did not adequately define the population, and they
20 analysis did not control for confounders. Due to this they concluded it was not very useful in
21 informing their recommendation and these were based on their clinical experience.

22 The Committee discussed the relevance of each sign and symptom included in the evidence
23 review. The Committee agreed that infants (children under 1 year of age), children and
24 young people and adults, required separate recommendations depending on the pertinence
25 of the symptoms for each subgroup.

- 26 • The Committee noted that the presence of meconium ileus in infants was considered a
27 highly suggestive sign of cystic fibrosis that should lead to further investigation. This is
28 because meconium ileus is a unique feature of cystic fibrosis.
- 29 • Likewise, the suspected diagnosis of DIOS in children, young people and adults (which is
30 the equivalent to meconium ileus in infants) is also a well-known factor suggestive of
31 cystic fibrosis that should also lead to further investigations of cystic fibrosis.
- 32 • In relation to respiratory symptoms, the Committee emphasized that one single respiratory
33 event should not necessarily lead to further investigation. The Committee considered
34 recurrent lower respiratory tract infections, chronic lung disease or chest X-ray with
35 persistent changes as reasons for referral in infants. Similarly, chronic sinus disease and
36 chronic wet or productive cough should also be considered as reasons for referral in
37 children, young people and adults. They noted that children do not produce sputum and
38 they agreed to use the term wet cough. In adults, they agreed it was more appropriate to
39 use productive sputum instead of wet cough, as adults are more aware of having sputum.
40 Finally they noted the presence of chronic or repeated chest infection regardless of
41 species may raise the possibility of cystic fibrosis. They agreed not to specify pathogens
42 causing chest infection as people with cystic fibrosis may become colonised or chronically
43 infected with many different opportunistic infections, and so highlighting any individual
44 species is likely to confuse diagnosis.
- 45 • The Committee discussed that people with cystic fibrosis often show signs of
46 bronchiectasis due to recurrent inflammation and infection. Additionally, people with cystic
47 fibrosis may present with asthma-like symptoms, such as wheezing, coughing, chest
48 tightness and shortness of breath, due to inflammation and infection of the airways.

- 1 • The Committee acknowledged that cystic fibrosis is known to cause pancreatitis.
2 Approximately 10-15% of people with cystic fibrosis are exocrine pancreatic sufficient and
3 so do not show malabsorption symptoms or diabetes mellitus, however these patients do
4 show a high incidence of attacks of pancreatitis.
- 5 • The Committee agreed that cystic fibrosis can also be suspected if there are signs of
6 faltering growth in infants and pre-school children or undernutrition in older children,
7 young people and adults. They noted that in young people, undernutrition can lead to
8 delayed puberty, but they agreed not to include it as a sign where cystic fibrosis should be
9 considered as there are also other reasons that can lead to delayed puberty. They also
10 noted that in adults with cystic fibrosis undernutrition normally goes in association with
11 other symptoms, such as pancreatitis and malabsorption.
- 12 • The Committee stressed that malabsorption should be separated from undernutrition as
13 malabsorption, such as steatorrhea, may be indicative of pancreatic insufficiency, a
14 common complication of cystic fibrosis (as discussed above), whereas undernutrition has
15 a variety of non-cystic fibrosis causes.
- 16 • The Committee highlighted that azoospermia showed an important association with a
17 diagnosis of cystic fibrosis. They noted that this was consistent with previous findings, as
18 it is estimated 99% of men with cystic fibrosis are infertile. They also noted that some men
19 can produce sperm, but they are still infertile. They discussed whether this symptom
20 applied to young people too, but they agreed that generally will not be apparent to people
21 under 18 years. It was also discussed that women may present with sub-infertility too, but
22 they agreed that this is rather unusual, and that clinically it is not possible to use the term
23 sub-infertility.

24 The Committee also discussed about other signs and symptoms not included in the review
25 protocol were also discussed, such as congenital intestinal atresia, DIOS, rectal prolapse,
26 pseudo-Bartter syndrome or diabetes mellitus.

- 27 • The Committee noted congenital intestinal atresia is a rare condition that leads to the
28 complete occlusion of the intestinal lumen in neonates and it has been associated with the
29 presence of cystic fibrosis.
- 30 • As highlighted in the protocol, DIOS is considered a symptom of cystic fibrosis. Therefore,
31 the Committee agreed that the suspected diagnosis of DIOS in children, young people
32 and adults (which is the equivalent to meconium ileus in infants) should also lead to
33 further investigations of cystic fibrosis.
- 34 • They noted that rectal prolapse can also be a sign that health care professionals should
35 be aware of. However, this sign does not present on its own, and it is associated with
36 pancreatitis.
- 37 • Cystic fibrosis may present as Pseudo-Bartter Syndrome (dehydration and salt depletion
38 due to dysregulation of salt homeostasis in cystic fibrosis), but this is rare in the UK and
39 more common in warmer climates.
- 40 • They also discussed whether the presence of diabetes mellitus in young people could be
41 a sign of cystic fibrosis. However, they agreed not to include it in the recommendations, as
42 cystic fibrosis-related diabetes mellitus is diagnosed in the presence of other symptoms
43 related to cystic fibrosis.

44 The Committee agreed that a sweat test should be recommended in infants, children and
45 young people. If the test was positive or borderline, the person should be referred for further
46 investigations at a specialist cystic fibrosis centre.

47 Furthermore, the Committee was concerned about carrying out genetic testing in a child
48 carriers to determine their carrier status (i.e. the child is heterozygous), as they cannot give
49 consent. Thus, they agreed that genetic testing should only be considered when sweat tests
50 results are uncertain. In adults, however, sweat test is less reliable, and cystic fibrosis gene
51 testing is preferred.

1 With regards to gene testing, the Committee noted that over 2000 variants in the CFTR have
2 been identified but at present only around 10% of these have been linked to the development
3 of cystic fibrosis. At present genetic tests only return results for the most common variants
4 associated with disease. Clinicians should be aware this means results from genetic tests
5 cannot rule out a diagnosis of cystic fibrosis.

6 The Committee discussed how the results of genetic testing results related to the obligation
7 of clinicians to refer people to a specialist cystic fibrosis centre. They agreed that a gene
8 testing revealing 1 or 2 cystic fibrosis mutations was a reason for referral. It was also decided
9 that due to the possibility that negative results do not entirely rule out cystic fibrosis, a referral
10 should be based on the clinicians own judgement in light of gene test results and apparent
11 symptoms (as discussed above).

12 These Committee noted that these recommendations are consistent with the NHS Service
13 Specifications for cystic fibrosis.

14 **5.7.2 Consideration of economic benefits and harms**

15 The Committee advised that their recommendation to offer a sweat test or gene test follows
16 current clinical practice to identify the clinical manifestations of cystic fibrosis and its
17 complications. Those tests would subsequently inform the patient's management to improve
18 their health-related quality of life. As a result, the Committee agreed their recommendations
19 promoted a cost-effective use of resources.

20 Moreover, knowing what clinical manifestations suggest a diagnosis of cystic fibrosis and
21 knowing the complications of cystic fibrosis, may lead to better identification and thus more
22 timely management and has therefore, indirectly, potentially important resource implications.
23 Therefore, it was important those manifestations and complications were included in the
24 Committee's recommendations.

25 **5.7.3 Quality of evidence**

26 Prospective and retrospective observational studies were included in the review. The quality
27 of evidence as assessed per individual studies was very low. The main sources of bias in the
28 studies were:

- 29 • Selection bias: It was noted that most studies do not indicate whether children had
30 undergone newborn screening, although Committee members were able to assume that
31 in some of the studies it was likely the participants were no assessed based on the date of
32 the study and the country (Hubert 2004 - France).
- 33 • Prognostic factor of interest or outcome of interest not defined: It was noted that some
34 studies did not adequately define the symptom evaluated. Other studies did not provide
35 details about how sweat test was conducted, or the thresholds used for diagnosis.
- 36 • Lack of control of potential confounders: Studies did not indicated either if the people
37 included in the studies presented other sign or symptoms suggestive of cystic fibrosis,
38 which was considered a very serious limitation.

39 Due to all these limitations, the Committee considered that the usefulness of prevalence data
40 reported in the studies is very limited.

41 None of the studies reported diagnostic accuracy data, or enough data to calculate
42 diagnostic outcomes.

43 **5.7.4 Other considerations**

44 No equality issues were identified by the Committee for this review question.

1 The Committee agreed there was no need to prioritise a research recommendation for this
2 topic. They noted that a universal screening programme is in place in the UK since 2006. In
3 addition, they agreed their clinical experience and expertise was sufficient to draft
4 recommendations regarding the clinical features that should lead to investigation of cystic
5 fibrosis. t.

6 **5.7.5 Key conclusions**

7 The Committee concluded that cystic fibrosis is a clinical syndrome that is diagnosed based
8 on clinical presentation and this diagnosis can be confirmed by sweat test and a genetic test.
9 In infants, children, young people and adults that have not been previously diagnosed with
10 cystic fibrosis, including those who had a negative newborn screening test, cystic fibrosis can
11 be suspected if one or more of the following signs or symptoms are present: meconium ileus
12 or DIOS, respiratory symptoms, pancreatitis, faltering growth or malnutrition, symptoms of
13 malabsorption, rectal prolapse or pseudo-Bartter syndrome.

14 **5.8 Recommendations**

15 **1. Be aware that cystic fibrosis can be diagnosed based on:**

- 16 • clinical manifestations, supported by sweat or gene test results for
17 confirmation or
- 18 • clinical manifestations alone, for people with symptoms who have
19 normal sweat or gene test results or
- 20 • positive test results in people with no symptoms, for example infant
21 screening (blood spot immunoreactive trypsin test) followed by sweat
22 and gene tests for confirmation.

23 **2. Assess for cystic fibrosis and, when clinically appropriate, perform a sweat test 24 (for children and young people) or a cystic fibrosis gene test (for adults) in people 25 with any of the following:**

- 26 • congenital intestinal atresia
- 27 • meconium ileus
- 28 • symptoms and signs that suggest distal intestinal obstruction syndrome
- 29 • faltering growth (in infants and young children)
- 30 • undernutrition
- 31 • recurrent and chronic pulmonary disease, such as:
 - 32 ○ recurrent lower respiratory tract infections
 - 33 ○ clinical or radiological lung disease (in particular bronchiectasis)
 - 34 ○ persistent chest X-ray changes
 - 35 ○ chronic wet or productive cough
- 36 • chronic sinus disease
- 37 • obstructive azoospermia (in young people and adults)
- 38 • acute or chronic pancreatitis
- 39 • malabsorption
- 40 • rectal prolapse (in children)
- 41 • pseudo-Bartter syndrome.

42 **3. Refer people with suspected cystic fibrosis to a specialist cystic fibrosis centre if:**

- 43 • they have a positive or equivocal sweat test result

- 1
 - 2
 - 3
- their assessment suggests they have cystic fibrosis but their test results are normal
 - gene testing reveals 1 or 2 cystic fibrosis mutations.

6 Information and Support

Review question: What information and support should be given to children, young people and adults with cystic fibrosis?

6.1 Introduction

People with cystic fibrosis and their families and carers often report that the information and support available to them at different stages of diagnosis and management is inconsistent. While the information needed differs between people, it is widely acknowledged that they need information and support at all stages of life with cystic fibrosis. Having the right information and support is an important factor in the effective management of cystic fibrosis, for example in order to achieve optimal health and quality of life.

Access to the right information and support can help to reduce anxiety and increase empowerment and confidence in managing symptoms and providing effective care for someone with cystic fibrosis. Equally, not having access to the right information and support can hinder people to make informed decisions about treatment and management.

Due to the variations in provision of information and support, the Committee agreed it was important to consider what information and support people with cystic fibrosis and their family and carers need.

The Committee considered not only the type of information and support available, but also the format in which it was delivered. Providing too much information in an inaccessible format may be equally detrimental as not enough quality information. They also considered some of the concerns people might have about access to reliable sources of information or support. An example of this is the increasing use of online forums or online communities where people who have cystic fibrosis, their carers or family members may look for answers to questions to supplement information gaps from their health care providers. The Committee did not want to negate the applicability of such avenues of support, but they did want to consider how they may be useful as an adjunct to other sources of information rather than as replacement.

6.2 Description of clinical evidence

The aim of this review was to identify the information and support that should be provided to people with cystic fibrosis and their parents and carers.

Studies were considered for inclusion if they collected data using qualitative methods (such as semi-structured interviews, focus groups and surveys with open-ended questions) in which the authors analysed the data qualitatively (including thematic analysis, framework thematic analysis or content analysis). Survey studies restricted to reporting descriptive data that was analysed quantitatively were excluded.

Given the nature of qualitative reviews, findings or themes were summarised from the literature. They were not restricted to those identified as likely themes by the Committee at protocol stage.

For full details see review protocol in Appendix D.

There were 31 studies included in this review.

All the studies were qualitative studies except 1 which used mixed methods (Hilliard 2014) and 2 studies based on questionnaires that included open questions (Fair 2000 and Widerman 2003). All qualitative studies included semi-structured interviews except 2 which

used in-depth interviews (D’Auria 2000, Johannesson 1998), 1 study that used unstructured interviews (Jessup and Parkinson 2010), 1 study that used an online ethnographical approach, which involved observing, downloading and analysing discussion group postings (Kirk 2016). There were 3 studies which used focus groups in addition to semi-structured interviews (Braithwaite 2011, MacDonald and Greggans 2010, Tipping 2010).

The size of the studies ranged from 4 to 203 participants. There were 2 studies which included parents of infants with cystic fibrosis (Tluczek 2006, Tluczek 2009), 1 study included parents of infants and children with cystic fibrosis (Jessup 2016), 4 studies included parents of children with cystic fibrosis (Coates 2007, Filigno 2012, Grob 2008, Whyte 1995), 5 studies included parents of children and young people with cystic fibrosis (Grossoehme 2014, Hodgkinson and Lester 2002, Hummelinck and Pollock 2006, Lang 2005, Miller 2009), 1 study included children with cystic fibrosis and their parents (Angst and Deatrck 1996), 1 study included infants, children, young people and adults with cystic fibrosis and their families (Roehrer 2013), 1 study included children, young people and adults with cystic fibrosis and their parents (Jessup and Parkinson 2010), 1 study included young people with cystic fibrosis and their parents (Kirk 2016), 7 studies included adults with cystic fibrosis (Braithwaite 2011, Hilliard 2014, Johannesson 1998, Kazmerski 2016, Widerman 2002, Widerman 2003, Widerman 2004), 2 studies included children and young people with cystic fibrosis (Barker 2012, Beresford 2002), 2 studies included young people and adults with cystic fibrosis (D’Auria 2000, Fair 2000), 1 study included pediatric nurses for cystic fibrosis (Bagnasco 2013), 1 study included physicians that cared for either children with cystic fibrosis, adults with cystic fibrosis or both children and adults with cystic fibrosis (Dellon 2012), 1 study included children and young people with cystic fibrosis, their parents, befrienders, play therapists and education liaison personnel (MacDonald and Greggans 2010), 1 study included parents or children and young people with cystic fibrosis and physiotherapists (Tipping 2010).

There were 9 studies conducted in the UK (Beresford 2002, Coates 2007, Fair 2000, Hodgkinson and Lester 2002, Hummelinck and Pollock 2006, Kirk 2016, Lang 2005, MacDonald and Greggans 2010, Whyte 1995), 13 in the USA (Angst and Deatrck 1996, Barker 2012, D’Auria 2000, Dellon 2012, Filigno 2012, Grob 2008, Grossoehme 2014, Hilliard 2014, Kazmerski 2016, Tluczek 2006, Tluczek 2009, Widerman 2002, Widerman 2004), 5 in Australia (Braithwaite 2011, Jessup 2016, Jessup and Parkinson 2010, Roehrer 2013, Tipping 2010), 1 in Italy (Bagnasco 2013), 1 in Sweden (Johannesson 1998), 1 in Canada (Miller 2009), 1 in multiple countries (Widerman 2003).

A summary of the included studies is presented in Table 159. See study selection flow chart in Appendix F, study evidence tables in Appendix G and list of excluded studies in Appendix H. For presentation of findings, a theme map was generated according to the themes emerged from studies (Figure 1). Due to the qualitative nature of these studies, evidence is summarised in GRADE-CERQual tables within the evidence report. Therefore no separate Appendix is provided for this.

6.3 Summary of included studies

A summary of the studies that were included in this review are presented in Table 20.

Table 20: Summary of included studies

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
Angst and Deatrck (1996) USA	Semi-structured interviews (secondary analysis of two	N=28 mixed population N=20 children with CF and both parents of each	To describe how children with chronic illness and their parents are	<ul style="list-style-type: none"> • Overall quality: low • Sample selection was not reported. • Description of data collection method was

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
	data sets)	child Age of children: range 7 to 11 years (median 9 years).	involved in health care decisions.	vaguely described. <ul style="list-style-type: none"> The analytical process was reported vaguely. Description of emerging and overarching themes was reported, but saturation of data was not reported.
Bagnasco et al. (2013) Italy	Semi-structured interviews	N= 12 paediatric nurses N=7 paediatric nurses for CF	To explore how nurses perceived autonomy in parents, adolescents, and children related to the management of chronic disease.	<ul style="list-style-type: none"> Overall quality: low Mixed sample with Neuro muscular and CF unit. No clear differentiation with overlap between samples. No information on structure of interview or whether topic guide reported. Description of how "themes" were arrived at was discussed but information was not sufficient to conclude if data collection process was robust. No information on data saturation and full exploration of theme. The analytical process was described with description of themes and categories. Researchers' role and potential influences in the analytical process not critically reviewed. Consistency between the researchers not reported.
Barker et al. (2012) USA	Semi-structured interviews	N= 24 children and young people with CF Age range: 11 to 18 years; mean age: 15.7	To explore the role of family and friends in supporting cystic fibrosis disease management during adolescence.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. Data collection relied on the semi-structured interviews. Process for semi structured interview was clearly reported but topic guide was not reported. The analytical process was described, with the use of predefined template analysis from the literature. No

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				<p>description of how "themes" were arrived at</p> <ul style="list-style-type: none"> • The researchers' roles and potential influences in the analytical process not critically reviewed • Multiple researchers but consistency between them not reported
Beresford et al. (2002) UK	Semi-structured interviews	N= 63 children and young people with chronic condition (n=11 with CF) Age: 11 to 16 years	To explore the experiences of chronically ill adolescents in communicating with health professionals, including the identification of factors which hinder or facilitate their use of health professionals as an information source.	<ul style="list-style-type: none"> • Overall quality: low • Sample selection was clearly reported. • Unclear about topic guides and limited information about group meetings. • The analytical process was not described in detail, no description of how "themes" were arrived at; researchers did not critically review their own roles in the process. • The study was not clear about the number of researchers involved in data collection or interviews.
Braithwaite et al. (2011) Australia	Focus groups (with staff) and semi-structured interviews (with people with CF and with family members)	N= 42 <ul style="list-style-type: none"> • n=12 people with CF • n=10 family members of people with CF who had died • n=20 staff • Age range of people with CF: 26 to 53 years; mean age: 35 years • 	To explore the unmet needs and key issues for people with CF, their families and the staff providing their care while awaiting organ transplantation.	<p>Overall quality: high</p> <ul style="list-style-type: none"> • Sample selection was clearly reported. • Structure of interview and topic guide decided by the experts within the hospital • The analytical process was described in detail. • Study conducted by lone researcher and may lack some of the formal research vigour.
Coates et al. (2007) UK	Semi-structured interviews	N=8 mothers of children with CF Children's age: 3 to 7	To supplement existing research to gain insight into mothers' experiences of informing relatives about CF and to look at patterns of	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported • Structure of interview and topic guide reported. Description of how "themes" were arrived at was discussed. Data

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
			communication within these families.	<p>saturation and full exploration of theme not clear.</p> <ul style="list-style-type: none"> The analytical process was described with description of themes and categories. Whether sufficient data were gathered to fully explore the themes is not clear. No critical review of the researchers' role in the process. Discrepancies between the researchers were addressed by the senior researcher with oversight.
D'Auria et al. (2000) USA	In-depth interviews	N=15 young people and adults with CF Age of respondents with CF: between 17 - 22 years with mean age of 19 years.	To explore the influence of peer relationships on adjustment to CF.	<p>Overall quality: low</p> <ul style="list-style-type: none"> Sample selection was not clearly reported. Structure of interview and topic guide not reported. No discussion on whether saturation has been reached for any of the themes reported. Description of how "themes" were arrived at was discussed. No critical review of the researchers' role in the process Study conducted by multiple researchers but the level of consistency between them not reported
Dellon et al. (2012) USA	Follow-up semi-structured interviews after surveys	N=26 physicians at two major CF centres Physicians worked with either children, or adults, or both children and adults.	To give an account of the physician perspective on communication with patients about the use of non-invasive and invasive mechanical ventilation for respiratory failure.	<p>Overall quality: low</p> <ul style="list-style-type: none"> Sample selection was process was not clearly reported. The relationship between the researcher and the participants was not reported. Description of data collection method was vaguely described and the setting of the interview was unclear. The analytical process was reported vaguely. Description of emerging

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				<p>themes and data saturation was not reported.</p> <ul style="list-style-type: none"> Results were presented clearly (e.g., citation/data and the researchers' own input distinguished)
Fair et al. (2000) UK	Survey (postal questionnaire with open-ended questions)	N=136 young people and adults with CF Age: ≥16 years	<p>To determine:</p> <ul style="list-style-type: none"> attitudes about fertility and pregnancy among subjects with CF satisfaction with communication on this issue from health care professionals 	<p>Overall quality: Low</p> <ul style="list-style-type: none"> Sample selection was not clearly reported. The relationship between the researcher and the participants was not reported. Limited information on the development of questionnaire. Study design limits the exploration of themes or development and eliciting further information. The analytical process of interpreting open ended question was not reported although use of specific qualitative software (NUD*ist) reported. No information on data saturation or identification of specific themes Results were presented clearly (e.g., citation/data and the researchers' own input distinguished).
Filigno et al. (2012) USA	Semi-structured interviews	N=8 parents of children with CF Mean age of children at the time of interview=8.2 years (SD 0.8).	<p>To:</p> <ul style="list-style-type: none"> better understand how families used the strategies taught in a behaviour-nutrition intervention identify the challenges with CF management families experienced during the 	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. Data collection method was described The analytical process was reported. Description of emerging and overarching themes, and saturation was reported. Findings or results: <ul style="list-style-type: none"> Results were presented, however, quotations/citations

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
			developmental transition from toddlerhood to school-age, particularly nutrition.	from respondents were not reported clearly
Grob (2008) USA	Semi-structured interviews	N=35 parents of children with CF (33 mothers and 2 fathers) (16 parents received a newborn screening diagnosis; 4 received a prenatal diagnosis; 4 received a diagnosis within a few days or weeks of birth for an asymptomatic child; 11 received a later diagnosis for a symptomatic child (between 2 months and 7 years). Age of children at the time of the interviews: Not reported	To examine how the expansion of mandatory genetic testing at birth structures, for parents, specific kinds of disjuncture between their child's medically defined disorder and the manifest illness.	<ul style="list-style-type: none"> • Overall quality: Low • Sample selection was not reported. • Data collection method was not described clearly. • The analytical process was reported, but vague in description. • Description of emerging and overarching themes was reported, but saturation of data was not reported.
Grossoehme (2014) USA	Semi-structured interviews	N=25 parents of children and young people with CF Age of children and young people: < 13 years	To describe parent experiences developing and utilizing CF care routines.	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported. • Structure of interview reported. Description of how "themes" were arrived at was discussed in depth. • Data saturation and full exploration of theme reported. • The analytical process was described with description of themes and categories and use of specific software for processing. • Researchers' role and potential influences in the analytical process not critically reviewed. • Study conducted by multiple researchers but the level of consistency

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				between the researchers not reported.
Hilliard et al. (2014) USA	Mixed-methods study including semi-structured interviews	N=16 adults with CF Age: 21 to 48 years	To involve individuals with CF in guiding the development of future mHealth apps for adherence promotion.	<ul style="list-style-type: none"> • Overall quality: high • Sample selection was clearly reported. • The analytical process was described in detail. • Description of how "themes" were arrived at; saturation of data and exploring all the themes in detail was reported
Hodgkinson and Lester (2002) UK	Semi-structured interviews	N=17 mothers of children and young people with CF Age range of people with CF: 2-13 years	To explore the current stresses and coping strategies used by mothers and to identify roles and strategies that nursing professionals could extend or adopt to support them and families of children with CF.	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported. • Data collection method was appropriately described, including saturation of themes • The analytical process was reported, but not in detail. Saturation of themes was described in the data collection section
Hummelinck and Pollock (2006) UK	Semi-structured interviews	N=27 parents from 20 families with a child with a chronic condition (n=4 children and young people with CF) Age range of children and young people: 0-16	To explore the complexity of parents' information needs and how current information provision is evaluated.	<ul style="list-style-type: none"> • Overall quality: low • Sample selection was clearly reported. The relationship between the researcher and the respondents clearly reported • Data collection method was not described in detail and cross referenced to other study for detail information • The analytical process was clearly reported. Unclear if saturation of data was achieved. Development of theme was described. No report on transcribing interview, validation or use of qualitative software for processing of information • Results were presented clearly and the findings

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				discussed in detail (e.g., citation/data and the researchers' own input distinguished)
Jessup et al. (2016) Australia	Semi-structured interviews	N=10 parents from 7 families of infants and children with CF Age of infants and children: 1 to 8 years	To explore the education needs of 10 parents following their infant's diagnosis with CF via newborn screening.	<p>Overall quality: moderate</p> <ul style="list-style-type: none"> • Sample selection was clearly reported. • Structure of interview and topic guide reported. Description of how "themes" were arrived at was discussed. • Data saturation determining further sample recruitment was reported. • The analytical process was described with description of themes and categories. • Researchers' role and potential influences in the analytical process not critically reviewed • Study conducted by multiple researchers but the level of consistency not reported.
Jessup and Parkinson (2010) Australia	Un-structured interviews	N=8 families with a son/daughter with CF (n=7 people with CF, n=17 parents either as couples or individually) Age of people with CF: range: 2-21, average: 10.5	To explore the experiences of living with CF.	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported. • The authors clearly explained and justified clearly how the data were collected. However, the authors did not discuss data saturation. • There was an in-depth description of the analysis process. It is clear how the themes were derived from the data. However, there was no critical review of the researcher's role in the process. • A colleague with expertise in phenomenology challenged perceived anomalies in the analysis. However, the total number of analysts

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				involved was unclear. There was no respondent validation due to concerns about the research burden on the CF population.
Johannesson et al. (1998) Sweden	In-depth interviews	N=14 women with CF Age: 22 to 34 years	To investigate psychosocial issues concerning puberty and motherhood among CF adult females, to see how they had obtained and conceived information on these matters and how they would like information to be given.	<ul style="list-style-type: none"> • Overall quality: low • Sample selection was clearly reported. The relationship between the researcher and the respondents was clearly reported. • Description of data collection method was vaguely described • The analytical process was not clearly reported. • Description of how emerging and overarching themes were reached was not reported, saturation of data was not reported. • Results were presented clearly supported with quotes and findings discussed in depth (e.g., citation/data and the researchers' own input distinguished)
Kazmerski et al. (2016) USA	Semi-structured interviews	N=22 women with CF Age: 18 to 30 years N=16 CF program directors	To explore the attitudes, preferences, and experiences of patients with CF and CF providers toward sexual and reproductive health care for young women with CF.	<p>Overall quality: moderate</p> <ul style="list-style-type: none"> • Sample selection was clearly reported. • Structure of interview and topic guide reported. • Description of how "themes" were arrived at was discussed. Data saturation and full exploration of theme reported. • The analytical process was described with description of themes and categories. • Researchers' role and potential influences in the analytical process not critically reviewed. • Study conducted by multiple researchers but the level of consistency between the

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				<p>researchers not reported.</p> <ul style="list-style-type: none"> Findings cannot be generalised to all CF care providers (for example those who are not directors).
Kirk et al. (2016) UK	Online ethnographical approach, involving observing, downloading and analysing discussion group postings	N= 182 participants on the parent's discussion group (103 discussion threads) and 97 participants on the young people's discussion group (48 discussion threads). Age of participants: not reported	To explore how online peer support is used by young people and parents to support self-care in relation to CF.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. Whether the sample is genuine representative of the CF population is unclear and it was based on a single online discussion forum. Data collection relied on the discussion thread. Description of how "themes" were arrived at was unclear or whether data saturation reached. The analytical process was described but was inadequate in description. Researchers' role and potential influences in the analytical process not critically reviewed.
Lang et al. (2005) UK	Semi-structured interviews	N=8 parents of children and young people with CF Age of children and young people: 3 to 16 years	To explore the experiences of parents of children with CF who had been asked to consider lung transplantation as a treatment choice and subsequently referred to one of two national lung transplant centres. Especially to explore their views on how the flow of information should be managed, and how the process of initial introduction by	<p>Overall quality: Moderate</p> <ul style="list-style-type: none"> Sample selection was clearly reported. The relationship between the researcher and the participants was not reported. Description of data collection method was vaguely described although validated qualitative research tool was used. The analytical process was reported vaguely. Description of emerging and overarching themes was reported, but saturation of data was not reported. Coding and category identification by two independent researchers suggest

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
			the referring centre could be improved.	<ul style="list-style-type: none"> reliability of findings Findings/results: Results were presented clearly. Discussion of the finding was limited and cross reference to citation/data and the researchers' own input was not adequately presented
MacDonald and Greggans (2010) UK	Semi-structured interviews and focus groups	N=17 participants of which n=10 were children or young people with CF or their parents, n=3 were befrienders, n=2 play therapists, n=2 education liaison personnel Age of children and young people with CF: 8 to 18 years	To evaluate the impact of a community youth befriending programme on a group of young people with CF and their carers.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. Structure of interview and topic guide reported. Description of how "themes" were arrived at was discussed. Data saturation and full exploration of theme not reported. The analytical process was described with description of themes and categories. Researchers' role and potential influences in the analytical process not critically reviewed. Study conducted by multiple researchers but the level of consistency between the researchers and rigour of the process reported.
Miller et al. (2009) Canada	Semi-structured interviews	N=47 parents of children and young people with complex chronic health conditions (n=7 with CF) Age of participants with CF not reported.	To examine how the experiences and perceptions of parents of children with complex chronic health conditions fit with the perspectives of academic and service providers on continuity of care. To identify the salient factors in the experience of continuity in this population.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. Structure of interview process reported but no information on use of topic guide. Description of how "themes" were arrived at was discussed. Data saturation and full exploration of theme reported. The analytical process was described with description of themes and categories and use of specific technology.

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				<p>No critical review of the researchers' role in the process.</p> <ul style="list-style-type: none"> • Interview conducted by single researcher but the level of consistency and accuracy not reported.
Roehrer et al. (2013) Australia	Semi-structured interviews	<p>N= 15 people with CF and their families</p> <p>Age of people with CF: 19 months to 52 years</p>	To provide an overview of the evaluation of a pilot trial of an information system conceptualised to develop to assist people with CF and their families, to enhance their skills and communication in relation to self-management for their condition.	<ul style="list-style-type: none"> • Overall quality: low • Sample selection was clearly reported. • Data collection method was described. • The analytical process was reported. Description of emerging and overarching themes was reported, but saturation of data was not reported. • Results were not presented clearly (e.g., citation/data and the researchers' own input distinguished).
Tipping (2010) Australia	Focus group of physiotherapists and semi-structured interviews with parents	<p>N=11</p> <ul style="list-style-type: none"> • n=5 physiotherapists • n=6 parents of children or young people with CF <p>Age of children: 2 to 16</p>	To identify factors that impair the delivery and retention of physiotherapy education for parents of children with CF and factors that impair effective physiotherapy treatment in the home environment.	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported. • Data collection was clearly reported, including the number of interviews, data saturation and the use of an interview map and digital recording. • The analysis process was described in detail, including details on how categories, sub-categories and themes were developed. However, there was no mention of contradictory data and researchers did not critically examine their own role, potential bias and influence during analysis and data selection. • Data analysis was completed independently by two researchers to enhance credibility. Moreover,

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				participants were asked to review the emerging themes and comment on their accuracy.
Tluczek (2006) USA	Semi-structured interviews	N= 33 families of infants with abnormal newborn results (8 with CF confirmed after sweat test) Age of infants: 1.5 to 6 months	<ul style="list-style-type: none"> To understand parents' perceptions about genetic counselling received while awaiting their infant's sweat test results, identify conditions that may affect the quality of their experience, develop a model for genetic counselling under the conditions of NBS using CF as a prototype for NBS programs using gene technologies. 	<ul style="list-style-type: none"> Overall quality: moderate Sample selection including process of recruitment was clearly reported. Structure of interview and topic guide not reported. Description of how "themes" were arrived at was discussed. Data saturation and full exploration of theme reported. The analytical process was described with description of themes and categories. Researchers' role and potential influences in the analytical process not critically reviewed.
Tluczek (2009) USA	Semi-structured interviews	N= 193 parents of 100 infants (100 interviews) with abnormal newborn screening results for CF (n=16 diagnosed with CF) Age of infants: <6 months	To examine how, when and from whom parents were informed about NBS, as well as to obtain their perspectives about how to improve parent education about NBS.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was reported. The relationship between the researchers was reported. Description of data collection method was reported. The analytical process was reported. Description of emerging and overarching themes was reported, but saturation of data was not reported.
Whyte et al. (1995) UK	Semi-structured interviews	N=16 families (n=4 with CF) Age of children: 4 years	To increase understanding of the needs of families caring for children with chronic illness; to investigate the continuity,	<ul style="list-style-type: none"> Overall quality: low Sample selection was clearly reported. Description of data collection method was not clearly described The analytical process was reported.

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
			effectiveness and acceptability of care from the parents' perspectives.	Description of methodology of emerging and overarching themes was not clearly reported, and saturation of data was not reported.
Widerman (2002) USA	Semi-structured interviews	N= 36 adults who received a diagnosis of CF at age 20 years or older	To determine the extent to which the needs and issues attendant to the adult CF interview are addressed by existing bad-news and paediatric CF recommendations and, if necessary, to develop and present recommendations to supplement them.	<ul style="list-style-type: none"> • Overall quality: moderate • Sample selection was clearly reported. • Data collection method was appropriately described • The analytical process was reported. Authors' interpretation was checked independently. Unclear if saturation of themes was achieved
Widerman (2003) International	Survey (survey instrument included open-ended questions)	N=130 adults diagnosed with CF after age 18	To address evidence gap about the actual and self-perceived knowledge of people diagnosed with CF as adults in order to inform the development of educational materials for this sub-population and to guide caregivers in educating them.	<ul style="list-style-type: none"> • Overall quality: low • Research method was not the most appropriate for answering the research question. Qualitative semi structured interview would have been better. • Sample selection was unclear. The need for international sample solicited through email was not justified. • Data collection relied on the postal questionnaire with open ended question. Survey instrument described. There was no theme or topic guide or report of data saturation. • The analytical process was poorly described and involved grouping information based on frequency. No further analysis of the qualitative information.

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				<ul style="list-style-type: none"> • Researchers' role and potential influences in the analytical process not critically reviewed. • Was sponsored by pharmaceutical company.
Widerman (2004) USA	Semi structured interview	N=36 young adults with CF	To explore the experience of receiving a diagnosis of cystic fibrosis after age 20.	<p>Overall quality: Moderate</p> <ul style="list-style-type: none"> • Sample selection: Sample selection was clearly reported. The relationship between the researcher and the respondents clearly reported • Data collection method was appropriately described • The analytical process was reported but was unclear on use of analytical software. Authors' interpretation was checked independently. • Unclear if saturation of themes was achieved • Results were presented clearly (e.g., citation/data and the researchers' own input distinguished).

Abbreviations: CF-cystic fibrosis; NBS-newborn screening; SD- standard deviation; UK-United Kingdom; USA- United States of America.

1
2
3

6.4.1 Clinical evidence profile

6.4.1.2 Theme maps

3 The theme maps are presented in Figure 3 and Figure 4.

Figure 3: Theme map: information needs for people with cystic fibrosis and their parents or carers




Figure 4: Theme map: support needs for people with cystic fibrosis and their parents or carers



1 The clinical evidence (GRADE-CERQual) for the information and support question is presented in Table 21 to Table 30.

6.4.21 Clinical evidence profile: information needs for people with cystic fibrosis and their parents or carers

2 Table 21: Summary of clinical evidence (GRADE-CERQual): Theme 1. Information at diagnosis

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: new-born screening					
5 studies (Grob 2008, Jessup 2010, Tipping 2010, Tluczek 2006, Whyte 1994)	5 studies using interviews	<p>5 studies conducted in Australia, the UK, the USA explored the information needs of the families of infants with abnormal newborn screening or children with cystic fibrosis.</p> <p>Parents reported they were relieved when they received the diagnosis of cystic fibrosis: <i>"..when we got it I was totally relieved. Even though he cystic fibrosis and I knew what it was and I knew the outcome of it, it was a relief, because I knew I was gonna be treated correctly. I knew..that I wasn't crazy, that I wasn't looking for something to be wrong with him, you know?" (mother of a child aged 7 years with delayed diagnosis of cystic fibrosis)</i></p> <p>The issue of delayed diagnosis was also discussed. Parents were concerned about their observations and suggestions being dismissed by the health care professional regarding their child's health: <i>"At the doctor's office, I would cry every time because he wasn't gaining [weight]. I think they kind of looked at me like this hysterical first-time mother, and the doctor whom I kept going to see kept saying 'Oh he'll kick in, some babies take a while to kick in.' That was really hard, being so powerless..." (mother of child who had delayed diagnosis)</i></p> <p>Some parents discussed about the information needs they had before or after receiving the results of the test: <i>"I want all the information. That way I know what I'm waiting to hear about. So it's not this black box that I don't know anything about and when I find out.. I think it was good that we had the information ahead of time. And, in fact, we were even seeking</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>it out.”</p> <p><i>"I'm the kind of person, I'm really proactive, so if I find out about a problem or an issue I want to dive into it and figure out what's the best way to do this, or what should we do? So I want all the information I can get.. I don't just want to be clueless and think, 'Oh she'll be fine, she'll beat the odds.' I want to know the dirty truth. I want to know what these people [with CF] go through so that I know how I can prepare myself and how I can prepare Alexandra". (parent of child with cystic fibrosis diagnosis)</i></p> <p>Another mother reported she did not know of cystic fibrosis until she received a positive new-born screen result for cystic fibrosis:</p> <p><i>"[Margo was] chunky," she recalls. "She was over nine pounds at birth, so I mean there was no indicators, you know, I mean visually, you know looking at her there wasn't anything to think there was anything wrong with her.." (mother of child of 2.5 years diagnosed at birth).</i></p> <p>Following diagnosis, parents sought information on future implication of cystic fibrosis:</p> <p><i>"I think probably for his future. What is it going to mean for (him) when he gets married and has his own kids? How is that (being a CF carrier) going to play a role? How do we tell him about all this and explain it to him so that he understands? And I think more of how is it going to affect him then us, now? How in the future is it going to affect him?" (mother of carrier infant)</i></p> <p>Information is seen as helpful to find reassurance. Parents experienced difficulties immediately after diagnosis as they felt confused and frightened when presented with information because they could not find answers to specific questions or resolve anxieties that confronted them:</p> <p><i>"I felt I was standing outside, watching it all going on, I did not even know...the IV's he was having straight away and I did not even know what that was all about. And they were just</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>pumping all these drugs into him. It was just one injection after the other in, pretty scary actually. You are left on your own in a room, you know. Sitting there, waiting for the next lot of IV's which of course I did not know when that would be. I did not know how many he had to have. It was difficult. " (Mother of a child with cystic fibrosis age 7 years)</i></p> <p>Some parents reported they received too much information at the time of diagnosis, and found this overwhelming. They felt they were "snowed under" by initial information "blah, blah, blah" (parent).</p> <p>Another parent also said "I think firstly in that first week we felt information overload... to be told like especially with the physiotherapy...that you've got to do this every single day for your child's life it's just overwhelming."</p> <p>Whereas some parents felt they had to fight to get the information they wanted:</p> <p><i>"We wanted to know all the details, and there would be things where we would ask the question and they would hedge as if to say, "We really don't like to tell everybody all those details to start with." Because we were both biology trained, we just wanted the absolute details . . . it seemed like getting blood out of a stone." (couple of parents)</i></p> <p><i>"not just the stuff they want the parents to hear" (one parent who read whatever she could and said that she could recognize the erroneous facts she received from health care professionals)</i></p> <p><i>"I didn't know anything about it and I got sent home with a little bit of information but it was like a week later before I had my first clinic. I was there all by myself and calling like half of [city name] it felt like trying to find somebody that...was home and that could talk...They told me it was genetics but I didn't understand what genetics truly meant either so I was like well what did I do and you know what I can have done different.." (mother of a one month old infant diagnosed with cystic</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>fibrosis).</p> <p>Or that they have to look the information themselves: <i>"I got the initial diagnosis over the phone.... they don't tell you much. It's like: "We think she's got cystic fibrosis and we want you to come in and talk about, and I can see you in two or three days' time'. And then it's like: 'Oh my God,' and 'Help, Google!'... (parent)</i></p> <p>Parents appreciate honesty, and did not want professionals to withhold information: <i>"One problem I have with some doctors is that they talk down to you and don't explain things thoroughly". (Parent of a child with cystic fibrosis diagnosis) (parent)</i></p> <p>However, they also appreciate information that is simple, and "not scary": <i>"We had one doctor. and we walked into the room and she sat down and we sat down and she said, 'Having cystic fibrosis is not a good prognosis.' And I sort of thought 'I don't really need to hear this. I'm well aware of what it does.' I didn't really think that was very thoughtful to say to someone while holding their new baby.. This one doctor. could be quite callous. And not really think about how you might be feeling as a parent..[What I needed was] the basics for the moment. You can find out everything else as you go along. It's not necessary to know everything right from the start". (mother of a child with cystic fibrosis diagnosis) (mother)</i></p> <p>And timely, at a rhythm that is comfortable for them: <i>"They had some hand-outs and things...as far as treatment and dietary concerns, you know," "But "there was just too much at that time to absorb, so we [would] look at it a little bit [at a time].." (mother of a child with cystic fibrosis diagnosis)</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 2: adult diagnosis					
2 studies (Wideman 2002, Wideman 2003)	1 study using questionnaires and 1 study using interviews	<p>2 studies conducted in USA and Europe with 166 adults with cystic fibrosis explored the experience of receiving a diagnosis of cystic fibrosis as adults.</p> <p>Some adults with suspected of cystic fibrosis were told by their physicians not to be concerned as they would probably test negative. Participants would have appreciated more clarity in this regard, as they were left confused and questioning whether they actually had cystic fibrosis:</p> <p><i>"[My doctor] decided to have me tested for CF...He said, 'Don't worry, you're too old'. It can't be, blah, blah, blah, blah. And it came back positive. So they did it again. And it was positive again" (46 year old woman)</i></p> <p><i>"OK you tested positive for CF, but we wanted to make sure you really have the illness. So, therefore we are going to forward you another clinic for confirmation" (Participant comment)</i></p> <p>Participants highlighted the need for information following diagnosis.</p> <p><i>"What do I do now?" or "how do I care for myself?"</i></p> <p>However, they noted information was non-specific and not relevant to them:</p> <p><i>"I wanted to know everything" (participant comment)</i></p> <p><i>"I got a booklet from the CF Foundation listing the median age of survival as 21. I was diagnosed at 24!" (woman diagnosed as adult)</i></p> <p><i>"I immediately went to the library the next day and looked up CF. And, everything said you were going to die by the time you were 16. And here I was 40". (man diagnosed as adult)</i></p> <p><i>"None (of the materials) addressed social, economic, psychological, or political issues and obstacles." (man diagnosed as adult)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>"We were given two books, but only one little paragraph really applied to me." (man diagnosed as adult)</p> <p>"I want to know about adult stuff. I want to know what to look for in symptoms, hints to better activities now, not to think I am going to die soon so often". (male participant)</p> <p>Some participants questioned why they were diagnosed with a paediatric condition, why they had not experienced symptoms, and how many others are diagnosed as adults. (author comment)</p> <p>Participants said that health care professionals should offer more information and spend more time with them:</p> <p>"Doctors need to have more information available. I had 15 min counsel with my diagnosis and that was it. I would call with a question. They would answer, but otherwise I have been on my own for information. It's scary to have a bomb dropped on you and then it's like here, deal with it." (female, 31 years)</p> <p>"My physicians gave me no information. I actually supplied them with articles and I have continued my self-education once I realized that physicians often do not keep up on the literature." (young woman with cystic fibrosis)</p> <p>They also found difficult to make questions at the time of diagnosis, as they knew nothing about the condition:</p> <p>"Because how can you ask questions about something you know nothing about? First you assimilate the disease, then you question it" (female participant)</p> <p>"I was confused. I didn't know what to ask" (male participant)</p> <p>In particular, newly diagnosed adults would like to know about general aspects, such as treatment available and impact of cystic fibrosis in their lives.</p> <p>For example they wanted to know how cystic fibrosis would change their lives (whether they could or should have children), and what to expect in the future. (author comments)</p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		They also wanted to know what treatments would be effective, and signs to look out for as indicators of their health status, and if their symptoms could be controlled (authors comments). They also wanted to know how long they could expect to live and whether being diagnosed as an adult was associated with longer life expectancy. (author comment)			

1 Table 22: Summary of clinical evidence (GRADE-CERQual): Theme 2. Information about treatment

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: trust in healthcare professionals					
1 study (Hodgkinson 2002)	1 study using interviews	In 1 study conducted in the UK with mother of children with cystic fibrosis, mothers reported that the primary care team's unfamiliarity with cystic fibrosis drugs, which contributed to distrust of primary care advice: <i>"The GP says-'what do you normally have?' and it's sort of, well it would be nice if they could tell me what they think". (mother)</i>	Limitation of evidence	Minor limitations	VERY LOW QUALITY
			Coherence of findings	Unclear	
			Applicability of evidence	Unclear	
			Saturation	Not saturated	
Sub-theme 2: more information					
2 studies (Angst 1996, Lang 2005)	2 studies using interviews	In 1 study conducted in the USA with children with cystic fibrosis and their parents, most families reported they were not given information about alternative care (for example, home vs hospital antibiotic therapy, and different means to enhance their child's nutritional status). (author comment) In another study conducted in the UK with families of children with cystic fibrosis, parents agreed that a gradual and informal process of discussing cystic fibrosis and treatment options prior to crisis-point was recommended as a means of preparing and supporting families more effectively. (author comment)	Limitation of evidence	Major limitations	VERY LOW QUALITY
			Coherence of findings	Unclear	
			Applicability of evidence	Unclear	
			Saturation	Not saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 3: physiotherapy					
2 studies (Jessup 2016, Tipping 2010)	2 studies using interviews	<p>In 1 study conducted in Australia with families of infants with cystic fibrosis, several parents referred the usefulness of the practical instructions given by the physiotherapist, which represented something they could do:</p> <p><i>“You know what was terrific was the little physio card that she made up for us...So when we’re home going ‘What do we do next?’ we could go back to the card and check.” (parent)</i></p> <p>However, in another study conducted in Australia with parents of young children with cystic fibrosis and physiotherapist, parents reported the information given was really outdated:</p> <p><i>“They gave us a video as well, really outdated ... I didn’t think it was the greatest video but anyway it was a bit old fashioned.” (parent)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY

1 Table 23: Summary of clinical evidence (GRADE-CERQual): Theme 3. Information about management of complications

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: lung transplant					
1 study (Lang 2005)	1 study using interviews	<p>One study conducted in the UK with 10 families of children explored the role of information in relation to lung transplantation.</p> <p>Most of the parents thought information helped them to prepare and face the reality of the situation:</p> <p><i>“Distressing but facing reality”</i></p> <p><i>“I wanted to be told everything (eg. Assessment criteria, procedures, complications and outcomes, including statistics, quality of life, drug-side effects and long-term prognosis”.</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Unclear</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>"Information makes you more aware and prepared"</i> <i>"I would have liked more information"</i></p> <p>However, other parents were reluctant to receive information: <i>"I didn't want information"</i> <i>"I didn't really want to deal with it"</i> <i>"Having the information is depressing"</i></p>			
Sub-theme 2: infertility					
3 studies (Fair 2000, Johannesson 1998, Kazmerski 2016)		<p>In 3 studies conducted in Sweden, the USA, and the UK with adults with cystic fibrosis and centre directors discussed the role of information in the context of sexual health and fertility.</p> <p>People with cystic fibrosis and care providers said they felt uncomfortable discussing these issues, as they are too intimate. Solutions to overcome this barrier were proposed: <i>"Sometimes women are afraid to speak up and they keep these things [SRH issues] personal to them... and might feel uncomfortable."</i> (woman with cystic fibrosis) <i>"I think the number 1 [reason] is that a lot of the younger women are ...embarrassed, especially because I'm a middle aged man, they're just a little embarrassed to bring it up."</i> "You do what you're comfortable with. I'm not good at fielding questions about sexuality, so I probably don't bring it up as often as I should." (male care provider) <i>"Sometimes, if people were to feel uncomfortable... maybe be given a pamphlet. Or some papers that have Web sites that you can, you know, go on... or maybe there would be an online thing where you can actually ask questions, kinda like be anonymous because maybe some people are embarrassed."</i> (woman with cystic fibrosis) Providers: "...a concise booklet that was [...] very accurate [with] all the different [SRH] subjects at a comprehensive level.</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>Because some [patients] probably don't necessarily want to talk about it in the clinic, but ... we could provide accurate information to them that they could access at their own convenience." (care provider)</i></p> <p>Timing of these discussions was also identified as important: <i>"[SRH] was brought up in school when I was in 4th or 5th grade, so I was probably 9...I think between 8 and 10, depending on if puberty is starting, I think you should be informed." (person with cystic fibrosis)</i></p> <p><i>"Honestly, for me, the easiest thing would be to just start [SRH discussions] young and have it be an expectation. We walk in here and know that people are going to talk to us about bowel movements, that's just part of what we know is gonna be asked. So, if you start [SRH discussions] at a young age, I think it just becomes part of the routine and it doesn't become as uncomfortable as it would be." (young person with cystic fibrosis)</i></p> <p>Women described satisfactory and unsatisfactory discussions with doctors and nurses: <i>"I felt no one would help me at least try and come to a decision. I had so little information. I was constantly told the risks were too high and now it's too late and I feel there's a huge gap in my life" (F, 33 years, FEV₁ 62% predicted).</i></p> <p>They also indicated they want more information about health consequences of pregnancy: <i>"(explain that) you will feel ill during and maybe for years after giving birth and when you have your baby, there's almost no time to be unwell yourself which can cause problems" (F, 23 years, FEV₁ 44% predicted).</i></p> <p><i>"Information about general health during pregnancy and risks about the actual birth" (F, 16 years, FEV₁ 50% predicted).</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>“To let them know that 14 days of IVS will be administered at home after the birth of the baby” (F, 29 years, FEV₁ 60% predicted).</i></p> <p>Men suggested they wanted to know “the facts: <i>“Simple cans and can’ts, facts ... to the point, no ‘maybe you can but ... etc.’” (M, 20 years)</i></p> <p><i>“I feel that there should be discussions and literature handed out in CF Clinic at an earlier stage e.g. not later than 16” (M, 20 years, FEV₁ 65% predicted).</i></p> <p><i>“Facts as they stand with hope via assisted fertility information” (M, 28 years, FEV₁ 22% predicted).</i></p> <p><i>“Make sure he knows about it early so he can learn to accept it easier” (M, 27 years, FEV₁ 40% predicted).</i></p> <p>The authors of 1 study noted that most men did not seem to be aware of the relatively low success rate of assisted fertility treatment. Of 30 comments from men, 9 were on the positive chance that they would be able to have children through assisted fertility and no man commented on the low success rate of fertility programmes. (author comment)</p> <p>In particular they noted that younger men aged < 20 years were much less likely than women or older men to make any comment on what information they wanted.</p> <p>Older men with good lung function seemed most likely to be distressed by their infertility:</p> <p><i>“I would like more information on how other people are handling the fact that they cannot have children” (M, 35 years, FEV₁ 84% predicted).</i></p>			

1 Table 24: Summary of clinical evidence (GRADE-CERQual): Theme 4. Provision of information

Study information	Description of theme or finding	Quality assessment
-------------------	---------------------------------	--------------------

Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: accessible information					
2 studies (Dellon 2012, Hilliard 2014)	1 study using surveys and 1 study using interviews	<p>In 1 study conducted in the USA with adults with CF, participants expressed the need for an accessible resource for general information about cystic fibrosis: <i>“Sometimes I’ll [wonder when] something happens health-related to me, ‘Is that normal for everyone or...is that happening to me because I have CF?’ And it’s hard to find particular sources where I can find that out.” (Age 35, Female)</i></p> <p>In another study conducted in the UK with health care professionals (pulmonologists), participants also highlighted the need to ensure access to same information for all patients and families: <i>“It would be helpful to formalize the structure. We do a lot of this stuff in a pretty informal ad-hoc fashion.” (health care professional)</i></p>	Limitation of evidence	Minor limitations	LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	
Sub-theme 2: adapted information/ relevant to the person					
5 studies (Beresford 2003, Hilliard 2014, Jessup 2016, Tluczek 2006, Widerman 2004)	5 studies using interviews	<p>In 1 study conducted in the UK with young people with cystic fibrosis, 2 studies conducted in the USA with adults with cystic fibrosis and 2 studies conducted in Australia and the USA with parents of children with cystic fibrosis reported on the importance of giving adapted information.</p> <p>Participants highlighted that information has to be relevant to the person: <i>“We just got a booklet that was produced in America and it wasn’t relevant here, and there was so much ambiguity.” (parent)</i> <i>“Sometimes I’ll [wonder when] something happens health-related to me, ‘Is that normal for everyone or...is that happening to me because I have CF?’ And it’s hard to find particular sources where I can find that out.” (Age 35, Female)</i></p> <p>Many participants who were diagnosed with cystic fibrosis as adults said they were given information addressed to a</p>	Limitation of evidence	Minor limitations	MODERATE QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>paediatric audience, and that few adult materials were available:</p> <p><i>"It seems there is not enough information for me to research on my own out there. I want to know about adult stuff" (man recently diagnosed with cystic fibrosis)</i></p> <p>A woman wanted directions on how to do chest percussions; she was given a booklet with illustrations of an infant. (author comment)</p> <p>Young people with cystic fibrosis also noted that healthcare professionals should communicate adequately, as sometimes they do not feel part of the conversation:</p> <p><i>"I go to see him, but not sure why 'cos mum talks about things."</i></p> <p><i>"He doesn't talk at my level. He ignores me and talks to my mum."</i></p> <p>The use of simple language was also important to parents:</p> <p><i>"We like to know that he does or he does not have this, instead of using all these big, giant words that make your head swim. It is confusing because one (test) could be a good thing that he's negative, another one could be a bad thing that he's negative."</i> (mother)</p> <p>Participants also discussed about relevance of information at diagnosis. Please see Table 1.</p>			
Sub-theme 3: right amount					
3 studies (Braithwaite 2011, Hummelinck 2006, Jessup)	3 studies using interviews	There were 3 studies conducted in Australia and the UK with parents of children with cystic fibrosis which explored parents' views on information needs, and reported on adequacy of information provided.	Limitation of evidence Coherence of findings Applicability of	Minor limitations Coherent Applicable	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
2016)		<p>Family members of children with cystic fibrosis sometimes felt that they were given too much information without giving them time to process it.</p> <p><i>“You get a bit overwhelmed by the information ... when you think about it later you think, Oh, what did they say?” (parent)</i></p> <p>Similarly, the authors of another study noted that parents of children with cystic fibrosis (that strictly required multidisciplinary care input and secondary care management) reported receiving an <i>‘information overload’</i>, particularly at the time of diagnosis. (author comment)</p> <p>This presents a challenge when people may have differing requirements.</p> <p>For one parent: <i>“We were just trying to get everything we could”</i>;</p> <p>Whereas for another: <i>“I think at that point for us it was probably all we needed”</i></p> <p>As 1 parent reflected, in reality there is no easy way: <i>“You’re given probably more than you want but it’s what you need. I don’t think there’s anything that you could stop or take away from that process. I don’t think there is any right way to do it... It’s just a process that you have to go through.”</i></p> <p>Participants also discussed about adequacy of information at diagnosis. Please see Table 1.</p>	evidence Saturation	Saturated	
Sub-theme 4: ongoing and timely					
5 studies (Braithwaite 2011, Dellon 2012, Jessup 2016, Lang 2005, Widerman 2004)	4 studies using interviews and 1 study using surveys	Two studies conducted in Australia and the UK with parents of children with cystic fibrosis, 1 study conducted in Australia with children and young people with cystic fibrosis and their parents and 1 study conducted in the USA with adults with cystic fibrosis identified receiving ongoing information as important.	Limitation of evidence Coherence of findings Applicability of evidence	Minor limitations Coherent Applicable	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>In 1 study, parents wanted information at different times, either as soon as possible or when the treatment became an option for their child. Therefore, it seems beneficial to have information available for parents to access as and when they wish. (author comment)</p> <p>For example, some parents felt they needed developmentally relevant information delivered at intervals during their child's first year:</p> <p><i>“So I think It’d be better to have a booklet day one up to may be six months pre food and then another booklet that’s relevant for that timeSome more of these fact sheets....and use those as a thing to go through with parents over time, rather than overload with information all at once.” (parent of an infant with cystic fibrosis)</i></p> <p><i>“... Just the practical day to day stuff would be great ... maybe when the baby is about 6 months, it would be great to have some day to day tips about CF: things to avoid, things to do.”</i></p> <p><i>“You’re not ready to hear what’ll happen in three years or things like that. You’re more ready to hear that stuff once you’ve found your platform of confidence and comfort.” (parent)</i></p> <p>Families also sought timely information to help them prepare. Participants showed their frustration, as they were not given information at time of diagnosis. (author comment)</p> <p><i>“I would have liked more information when (patient)'s health was better so I wasn’t in shock.” (parent)</i></p> <p>In another study conducted in the USA with a group of pulmonologists, they suggestive to take a proactive rather than a reactive approach to information giving:</p> <p><i>“Maybe if we can find a way of bringing this up at earlier points in the disease it wouldn’t become such a heavy weight on the patient. It would be helpful to be able to say, ‘This is not</i></p>	Saturation	Saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<i>something we are doing uniquely for you, this is just part of what we do.” (healthcare professional)</i>			
Sub-theme 5: content					
2 studies (Jessup 2016, Widerman 2004)	2 studies using interviews	<p>One study conducted in the USA with adults with cystic fibrosis and 1 study conducted in Australia with parents of children with cystic fibrosis participants discussed about the information they need.</p> <p>Many participants reported they needed general information, rather than illness-specific.</p> <p>Participants were not particularly interested in biomedical descriptions of cystic fibrosis, or even in instructions on self-care. (author comment)</p> <p>A woman recently diagnosed with cystic fibrosis lamented her education involved <i>"technical things"</i> and <i>"nothing about what life would be like"</i>.</p> <p>Another man said: <i>"We need more on everyday stuff"</i></p> <p>Several people referred to the practical instructions given by the physiotherapist, which represented something they could do:</p> <p><i>"You know what was terrific was the little physio card that she made up for us...So when we're home going 'What do we do next?' we could go back to the card and check."</i></p> <p>Some participants complained the information was unnecessarily "frightening or upsetting":</p> <p><i>"We received a sheet ... with genetics with some basic pointers about cystic fibrosis... but the very first line on it was the life expectancy for CF... and the number ... it's really confronting to have that as one of the first pieces of information."</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>“The first thing you see when you open their website is that 30% of teenagers died from CF ... To see that figure is like ‘Oh my God.’” and one father explained: ‘This is what we got at the time ... All we could see, there’s a little kid in jail: ‘Just one cell mutation can trap you for life’.... It’s just awful.”</i></p>			
Sub-theme 6: format					
3 studies (Jessup 2016, Hilliard 2014, Lang 2005)	3 studies using interviews	<p>In 2 studies conducted in Australia and the UK with families of children with cystic fibrosis, participants commented on the format, with recommendations from simple fact sheets, to brochures, booklets, DVDs, a map, personal accounts, specific transplant group meetings and counselling. They needed developmentally relevant information delivered at intervals during their child's first year. (author comment)</p> <p><i>“So I think It’d be better to have a booklet day one up to may be six months pre food and then another booklet that’s relevant for that timeSome more of these fact sheets....and use those as a thing to go through with parents over time, rather than overload with information all at once.”</i></p> <p>One study conducted in the USA with adults with cystic fibrosis explored the use of online information and mobile applications. Although some participants noted the benefits of using mobile applications as a way of contacting with healthcare professionals, others reported that cystic fibrosis providers may not be responsive if contacted via telephone or email between visits, and would prefer to see the provider in person:</p> <p><i>“I’m not always in a place where I can call them, so if I can just shoot a text...that would be convenient...If they want me to do something out of the ordinary...I want to [ask], ‘How exactly did you want me to do this?’ ” (Age 23, Male)</i></p> <p><i>“In the everyday world [electronic communication] just seems to be replacing talk and conversation and you know,</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Unclear</p> <p>Unclear</p>	<p>LOW QUALITY</p>

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>communicating that way, I don't want that to happen [with my doctors]."</i> (Age 35, Female)</p> <p>Similarly, some participants reported that they would not need another channel of communication with their family: <i>"I would use other more direct messages."</i> (Age 28, Male) <i>"I would probably continue to communicate with them the way I already do."</i> (Age 33, Female)</p>			
Sub-theme 7: privacy					
1 study (Beresford 2003)	1 study using interviews	<p>In 1 study conducted in the UK with young people with cystic fibrosis, some participants felt the privacy was important to communicate freely. <i>"You don't tell the doctor anything because you don't want them [student doctors] to hear."</i> (young person with cystic fibrosis)</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	LOW QUALITY

1 Table 25: Summary of clinical evidence (GRADE-CERQual): Theme 5. Perceived benefits of receiving information

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: empowerment and control					
4 studies (Braithwaite 2011, Hummelinck 2006, Jessup 2016, Lang 2005)	4 studies using interviews	<p>Four studies conducted in Australia and the UK with parents of children with cystic fibrosis and people with cystic fibrosis explored the reasons why parents want to receive information.</p> <p>Participants thought that they required information so that they could ask relevant questions when required: <i>"I need to ask more questions but sometimes I don't even know what to ask"</i> (adult with cystic fibrosis)</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>Parents wished for more information to feel involved in management of their child's illness and to be able to understand decisions being made. Understanding what was happening helped some parents to cope with the illness and re-establish a sense of control:</p> <p><i>"I want everything [all there is to know about cystic fibrosis] now. Because then when something arises, you can go [snaps his fingers] 'Right, I recognise that, we have got to do this' or 'I know what that is, we do not need to panic'. We know, we are in control. I think we are in control anyway, but when he's not well...It might not be nice, knowing what might be, but it's better to know. At least you are in control that way". (Father of a child with cystic fibrosis age years)</i></p> <p>Some parents also saw information as crucial to prepare. (author comment)</p> <p>In addition, parents of children with cystic fibrosis reported that information gives them reassurance and empowerment:</p> <p><i>"It would have been a bit more empowering if I'd more information ... Because you kind of want parents to become experts." (parent)</i></p> <p>One parent also mentioned that the unknown exacerbated fear:</p> <p><i>"I think the biggest thing was not knowing what might happen, which gives you a bit of a panic." (parent)</i></p>			
Sub-theme 2: hope					
2 studies (Jessup 2016, Wideman 2002)	2 studies using interviews	<p>In 1 study conducted in Australia with 10 parents of infants with cystic fibrosis, parents saw information as a source of hope:</p> <p><i>"So that's the most important thing, that optimism is really important for parents."</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p>	<p>Minor limitations</p> <p>Unclear</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>This was particularly so because it: "...moves you out of a 'victim' state to a 'move on with it' state."</p> <p>As one mother declared: "I just wanted to get on with bringing up my baby".</p> <p>Parents sought information that enabled them to assign their child's position on the 'severity spectrum' of cystic fibrosis disease, explained by this father: "I want to kind of put the severity of CF on a spectrum, and then ask: 'Where does my child fit? Is she worse or better off compared to someone else?'"</p> <p>However, in spite of the best intentions and sources of current information, prognosis evolves and eludes: "I know obviously they can't tell you for various reasons because they don't know themselves, but that's one question that I was asking myself a lot ... because we still don't know."</p> <p>Similarly, in another study conducted in the USA with adults newly diagnosed with cystic fibrosis, participants reported that they reacted in a positive manner when their questions were answered, and given information about cystic fibrosis, as well as positive messages to promote hope. (author comment)</p>	<p>Applicability of evidence</p> <p>Saturation</p>	<p>Applicable</p> <p>Not saturated</p>	

6.4.31 Clinical evidence profile: support needs for people with cystic fibrosis and their parents or carers

2 Table 26: Summary of clinical evidence (GRADE-CERQual): Theme 1. Support at specific stages in the course of the condition

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: new-born screening and diagnosis of cystic fibrosis					
1 study (Grob 2008, Whyte)	2 studies using interviews and 1	Three studies conducted in the UK, USA and Europe with parents of children diagnosed with cystic fibrosis reported on	Limitation of evidence	Major limitations	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
1994, Widerman 2003)	study using a questionnaire	<p>the experience following diagnosis.</p> <p>Parents felt overwhelmed and distress when they found out about the diagnosis and following discharge from hospital: <i>"My husband and I were completely traumatised at first and only kept going on adrenalin for the first year. Now I grieve for the baby I wanted and didn't get. However, we still love Jane" (mother of infant diagnosed at birth)</i> <i>"we were told that she might die when she was in her teens-I shall always remember the doctor saying that, and the effect it had on us" (mother of an infant diagnosed at 15 months age, who had symptoms of loose stools)</i> <i>"..It just seemed like everything changed.. It was like there is so much more now to taking care of her, and are we really fit to do that?.. [I]t was just so overwhelming. I mean the first time we went to the clinic they were like well, you have to do this and this. And we met with nutritionists, respiratory therapists and pulmonologists and social workers and you know it was just all so overwhelming, all this stuff we were going to have to do. I remember leaving there thinking "how am I going to do all this stuff in one day?" ". (parent of infant diagnosed with cystic fibrosis, after discharge from hospital)</i></p> <p>Mothers reported that they seek support from professionals upon receiving a positive screening test for cystic fibrosis: <i>"Well, I would say in the very beginning [when we got the NBS diagnosis]. me and my husband, we were both kind of like what do you [doctors] want to do, what do you think we should do, what do you think we should do, what do you think is best? You guys are the doctors, you know" (mother of infant diagnosed with cystic fibrosis).</i></p> <p>Mothers reported that they contacted the health care professional for advice, expertise, reassurance and instructions about caring for their infant:</p>	Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>"Once I found out that she possibly could have the cystic fibrosis, I called so many times in the middle of the night. I'm like 'Oh my god, she's breathing really heavy, I don't know if this is right..' There was just a lot of follow-up that came from the hospital that helped.' (mother of one month old infant)</p> <p>One mother found it difficult to approach the health care professional as they regarded her as being overly needy: <i>"What was hard I think in the beginning [was] being new as a parent for one and not knowing what was normal for children. and then dealing with the disease, the health care.. The people in health care were somewhat hard to deal with... because I would call a lot because I didn't know, because I was so scared, because there was such a fear..I would call the nurse a lot and say 'I don't know if this normal or not, this doesn't seem right.'</i> (mother of new born infant diagnosed with cystic fibrosis).</p> <p>With regards to diagnosis of cystic fibrosis as adults, 1 study conducted across different centres in the USA and Europe explored the experience of people who receive the diagnosis of cystic fibrosis as adults. A 51-year-old man diagnosed 2 years previously said, <i>"The concern for my emotional health by the medical professionals was almost non-existent."</i> A 37-year-old male simply wrote, <i>'It seems like care is lacking.'</i></p>			
Sub-theme 2: support through end of life					
1 study (Braithwaite 2011)	1 study using interviews and focus groups	<p>One study conducted in Australia with 42 participants, including people with cystic fibrosis, their parents and the healthcare professionals looking after them noted that family members needed support to address the imminent death of their children: <i>"We had spoken about death and his wishesI could just</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p>	<p>No limitations</p> <p>Coherent</p> <p>Applicable</p>	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>focus on (patient), say the things I needed to say... have no regrets... prepare myself for the worst...which I think helped me to accept" (parent)</i></p> <p>Likewise, people with cystic fibrosis felt vulnerable and thought they needed psychological support: <i>"I would need some psychological support ... I worry about my family and how they will cope... knowing there is counselling is a comfort to me." (person with cystic fibrosis)</i></p>	Saturation	Not saturated	

1 Table 27: Summary of clinical evidence (GRADE-CERQual): Theme 2. Support with treatment

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: continuity of care					
2 studies (Beresford 2003, Miller 2009)	2 studies using interviews	<p>One study conducted in Canada that included 7 parents of children with cystic fibrosis and another study conducted in the UK with young people with cystic fibrosis noted the importance of continuity of care.</p> <p>They reported lack of coordination between different organisations. <i>"To me, it was like you were cut off from life. You turn six, that's it. You're gone. When they do it from zero to six, they coordinated. They stayed on top of it, they tell you what they need. As soon as they get into the school system ... I'm not even sure who coordinates it then." (parent)</i> <i>"It would be better just to have one doctor so we could move on to different parts instead of getting the same questions again and again." (young person with cystic fibrosis)</i></p> <p>Parents believed that knowledge of the child, according to</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>parents, developed through relationships with a consistent set of service providers, both in and outside of medical settings.</p> <p><i>"You need to see the regular faces, because they're the ones you feel at least know your child best," the mother said. "They know the history," the father added, "so you feel they have the whole story." (parent)</i></p> <p><i>"It's nice when relationships do develop, you know. Kate knows the nurses [in the cystic fibrosis clinic] and she likes them, and ... she's not scared when she goes down there. Those faces are familiar to her, and if she is sick, it's not scary, it's not somebody she doesn't know." (parent)</i></p>			
Sub-theme 2: active involvement from health care professionals					
1 study (Hodgkinson 2002)	1 study using interviews	<p>In 1 study conducted in the UK with 17 mothers of children with cystic fibrosis, some participants reported that there was a need for GPs to become more involved in their children's treatment as their role was limited:</p> <p><i>"Well basically he [GP] just writes prescriptions for us...he hasn't played a big part in the cystic fibrosis part of it". (mother)</i></p> <p>Many mothers turned to nurse specialists for support and advice, and was critical in interpreting information and aiding understanding of treatment and compliance:</p> <p><i>"If we do have any worries or concerns [we] just contact the hospital straight away...we never bother going up to the doctors, we always just contact S, she's one of the nurses". (mother)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Unclear</p> <p>Unclear</p> <p>Not saturated</p>	VERY LOW QUALITY
Sub-theme 3: decision making					
2 studies (Angst 1996, Dellon 2012)	1 survey using interviews and 1 study using a survey	<p>Two studies conducted in the USA, 1 with 34 healthcare professionals (pulmonologists) and another with 20 children with cystic fibrosis and their parents, reported on the importance of supporting parents in relation to decision</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p>	<p>Major limitations</p> <p>Coherent</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>making.</p> <p>Most parents viewed themselves as involved in the decision making process (discussion with the healthcare provider, decisions at home regarding enzymes and respiratory therapies). (author comment)</p> <p>However, some parents did not see themselves as having much room to make decisions. They noted decisions were based on recommendations made by health care professionals. (author comment)</p> <p><i>"We pretty much get the plan from [the doctor] and then we just implement it...I just do what he tells me, basically. I think if I had something that was a nagging concern, I certainly know he would listen and respond to those concerns, but to date I've just not had any...I figure when he wants to change the programme, he'll tell me and we'll just do it" (mother of child with cystic fibrosis).</i></p> <p>Parents viewed the outcome of decisions about their child's health as potentially very serious. They identified the outcome of making the wrong decision as illness progression and even death, which is why they considered the healthcare professional's recommendations seriously. (author comment)</p> <p>With regards to children, some parents reported they did not involve children in decision making:</p> <p><i>"It was presented as a need for him to get back to approximately where he was on the growth curve. And if he does, then he avoids the tube. If he doesn't, then he gets the tube...We don't have that much that's negotiable...I don't see that there's two equivalent paths of therapy that are offered. Generally there's only one. Therefore, there's no need for discussion." (father)</i></p> <p><i>"I don't think he really should have much choice. I think we should just tell him. He certainly has as much right to ask</i></p>	<p>Applicability of evidence</p> <p>Saturation</p>	<p>Applicable</p> <p>Saturated</p>	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>questions and get answers as I do, but I want him to know that it's very important to do what we're told in this case. Not to be a creative thinker." (mother's comment)</i></p> <p>Other parents did not previously think about involving their children in decision making as they waited for a cue from the healthcare providers as to when it was appropriate to involve their children:</p> <p><i>"I guess I never thought to ask [child]...I guess as he's gotten older, there's no reason not to ask his opinion". (mother)</i></p> <p><i>"As a parent, I guess I need to push or just be told it's OK to do this now. You know that this is the stage that the child can handle it. You know, because when [the health care professional] tells you it's OK, it's a lot easier than you making that decision". (father)</i></p> <p>Children did not see themselves as involved in planning or decision making either:</p> <p><i>"When I go to clinic, he doesn't usually talk to me...when I loose weight, he yells at my mom for it". (girl with cystic fibrosis)</i></p> <p>Most children liked being uninvolved; however, many children wanted greater involvement:</p> <p><i>"Sometimes they want me to take more medicine, and I don't even know what the medicine is, and if I stop taking other medicines. And so I have to ask my parents and they have to ask. If they at least told me, I think I would feel a little better about why I'm taking this medicine..I think I'd feel more comfortable if I got to talk to them". (girl's comment)</i></p> <p>HCPs suggested that educational and decision support tools were seen as useful prompt in-depth discussions about treatment options and preferences:</p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<i>"Having tools to facilitate the discussion would help. We may misread the kind of information people want. And if you are trying to explain treatments, I have no materials to give people about things like bilevel pressure ventilation." (HCP)</i>			
Sub-theme 4: online support					
1 study (Kirk 2016)	1 study using information from online forums	<p>1 study conducted in the UK explored how online peer support is used by young people and parents (n=279) to support self-care in relation to cystic fibrosis.</p> <p>They noted the group sought advice and support about how to manage different treatments and therapies.</p> <p><i>"Have you tried going swimming together just the girls in yr family make exercise fun and then when she has done it go and have lunch together or buy her a treat, That has worked with my son 13." (parent)</i></p> <p><i>"We used to use straws – blowing cotton wool balls across the kitchen table – actually our physio who has retired showed us this one – it was great fun when we all joined in as a family. "</i> (parent)</p>	Limitation of evidence	Minor limitations	LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Unclear	
			Saturation	Not saturated	
Sub-theme 5: physiotherapy					
2 studies (Tipping 2010, Whyte 1994)	2 studies using interviews	<p>2 studies conducted in the UK and Australia with families of children with cystic fibrosis and physiotherapists noted parents found helpful learning about physiotherapy in hospital and they considered the hospital as a primary source of professional support, for problems with their child adhering to physiotherapy:</p> <p><i>"we do it three times a day mostly-on different parts. We do front and back in the mornings, sides at lunch time and tops in the evening. We learnt that in hospital, it was helpful" (parent of a child with cystic fibrosis)</i></p> <p><i>"the new physiotherapist came out to get the message over. 'If</i></p>	Limitation of evidence	Major limitations	LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>you won't let mum do it, someone else has to come"</i> (mother of a child with cystic fibrosis)</p> <p>Similarly, they also reported that the visit of the physiotherapist to the school was helpful: <i>"the physiotherapist went to the school and told them all about CF, and that helped the teachers to understand Rachel's problems"</i> (mother of a child with cystic fibrosis, at school)</p> <p>Mothers reported that continued support from the physiotherapist was helpful in crisis situations: <i>"I wasn't coping and I went to the hospital before my appointment was due and I just broke down in tears. The physiotherapist was very good, and said it wasn't a unique situation, and that I was coping well. I was happy that I had spoken to someone about it-it helped"</i> (mother of a child with cystic fibrosis) <i>"I asked to see the physio at the time... it was just impossible to try and keep someone of that age still for 20 min to half an hour to finish the treatment... he started telling us about doing some bubble games"</i> (parent of a young child with cystic fibrosis)</p>			
Sub-theme 6: nutrition					
1 study (Filigno 2012)	1 study using interviews	<p>In 1 study conducted in the USA with 8 parents of children with cystic fibrosis highlighted the difficulties parents experience in relation to nutrition.</p> <p>Parents recalled that learning how to deliver both positive consequences (praises and rewards) and negative consequences (removal of privileges) to manage mealtime behaviour was helpful.</p> <p>They also reported intense desperation to get their child to eat, including preparing meals for the child so that the child would eat.</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Unclear</p> <p>Unclear</p> <p>Not saturated</p>	VERY LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>Parents found that an ongoing challenge was general behavioural non-compliance including refusal to eat, take enzymes, and complete a fecal fat test.</p> <p>Parents found challenges with transfer of treatment responsibility from them to their child for certain aspects of cystic fibrosis management.</p> <p>Parents/families reported that managing transition to school was difficult as parents were not able to monitor their child's nutrition during the school day, and found that they were compensating their child's food intake at home (dinner). Parents also reported that there was a negative impact of missing school due to hospitalisation and illness. Parents struggled with partnering with schools to ensure that their children received appropriate accommodations.</p>			
Sub-theme 7: attitude from parents and friends					
1 study (Barker 2012)	1 study using interviews	<p>In 1 study conducted in the USA with 24 children and young people with cystic fibrosis participants perceived treatment-related behaviours as non-supportive, although they were reluctant to rate family members or friends as being 'unsupportive' even when annoyed by them.</p> <p>One young person stated, <i>"Their intentions are good but the way they pursue it isn't that wonderful. I'd rather them tell me to do it instead of them yelling at me to do it. I mean I'm a person, too, I forget things."</i></p> <p>Young adults reported becoming annoyed when reminders were given after the treatment was completed or when the adolescent has a plan to complete the treatment.</p> <p><i>"I get annoyed 'cause sometimes [mom] reminds me and I</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Unclear</p> <p>Unclear</p> <p>Saturated</p>	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>already did them. When talking about support from a close friend," another participant said, "It starts to get a little nagging at times, he's like 'You gotta do it, you gotta do it.' And I'm like, 'I know, I have a set time for this. I'll do 'em, don't worry!"</i></p> <p>They also found reminders annoying when the reminder interrupted other activities.</p> <p><i>"Well, like they'll tell me to do stuff. And if I'm talking on the phone or hanging out with my friends, then I don't want to do it and it gets on my nerves." Another participant said, "Well sometimes like when I want to watch a show or something, she tells me to do my treatment, so I have to stop the activity and I go do it – that gets annoying."</i></p> <p>One adolescent said, <i>"cause sometimes she'll say it and it'll really get to me and I'll be like, 'Don't tell me what to do' or 'I'll do whatever I want', you know, 'I can take care of myself'. So it's not that she's saying anything differently, it's just the way I'm perceiving it that day."</i></p> <p>However they recognized the need for persistent reminders and their benefits even when they are annoyed by them.</p> <p>One participant reported, <i>"[Mom] keeps telling me to do it whether I want to or not, she knows that it's going to help me so it's pretty supportive."</i></p> <p>While another stated, <i>"[Mom] usually tells me to do [airway clearance] daily 'cause sometimes I don't like doing it so she usually has to tell me or else I won't do it."</i></p> <p>Similarly, one adolescent talked about reminders from her friend, She pretty much says, <i>"Hey ok, if we're going to go out, you know, just like, let's get your meds done.' She wouldn't say, 'Ok you have to do your meds now' she'd say, 'So let's get your meds done just before we go or whatever so we don't have to do it later.' She'll present it in the way that it's not like something I have to do."</i> She rated her friends' reminders as very supportive because they were as encouraging and not as demanding the treatment be completed.</p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall

1 Table 28: Summary of clinical evidence (GRADE-CERQual): Theme 3. Management of the complications of cystic fibrosis

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: infertility					
2 studies (Fair 2000, Kazmerski 2016)	1 study using interviews and 1 study using a questionnaire	<p>2 studies conducted in the UK and the USA with young people and adults with cystic fibrosis and HCPs discussed about the support needed regarding fertility issues.</p> <p>Participants reported they want the emotional impact of infertility to be recognised by their health professionals: <i>“I do feel this can be a very emotional issue” (M, 23 years, FEV₁ 45% predicted).</i> <i>“Give some hope of being able to father and try and make them not feel a failure if they can’t father children” (M, 31 years, FEV₁ 96% predicted).</i> <i>“It is terrifying for men not to be able to father a child” (M, 34 years, FEV₁ 84% predicted).</i></p> <p>Both professionals and people with cystic fibrosis reported this issue is often neglected: <i>“It’s always just like you don’t talk about it [SRH], it’s one of those things that’s left to the side, it’s [...] like they [CF providers] feel it’s not as important as everything else, but sometimes it is. I mean, it [SRH] wasn’t life or death threatening, but it could’ve changed my life a lot.” (person with cystic fibrosis)</i> <i>We’ve been so focused on nutrition and liver disease and lung disease and diabetes, but now that [...] quality of life continues to improve, this will be a big issue, a more important issue for everyone.” (HCP)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	VERY LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall

1 Table 29: Summary of clinical evidence (GRADE-CERQual): Theme 4. Psychological and emotional support

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: from healthcare professionals					
6 studies (Braithwaite 2011, Hodgkinton 2002, Jessup 2016, Kazmerski 2016, Lang 2005, Tipping 2010)	6 studies using interviews	<p>4 studies conducted in Austria and the UK with parents of children with cystic fibrosis, and another study conducted in Australia with parents of children with cystic fibrosis and healthcare professionals commented on the perceived emotional support from healthcare professionals</p> <p>Many parents felt supportive by the staff, and they found this to be important and encouraging, and also helpful to achieve a good relationship with staff:</p> <p>For example one parent commented about their physiotherapist: <i>"I wasn't coping very well... the physio was very good. I don't know why, but for some reason I think the physio part of Cystic Fibrosis, not only is it huge because of what it does, I just think it's huge in terms of support. We just asked them so many questions ... I've placed my baggage on them and they've taken that really well."</i> (parent of a toddler with cystic fibrosis)</p> <p>Another mother commented: <i>"He will always, without fail, give you praise...and he'll say Wonderful specimen mother. Well done, keep it up, wont you, you're doing marvellous', and it's what you need"</i>.</p> <p>Having a good relationship with the clinician introducing and discussing was seen as essential. Parents felt the process would also be less formal and less distressing if the cystic fibrosis nurse, who knew the family and who had perhaps been</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>previously emotionally supportive, was central to this. (author's comment)</p> <p>Family members said they needed timely reassurances that their children are looked after: <i>"Even though there is no new information we still want to hear from the team ...otherwise you know you're not being abandoned but you feel abandoned". (parent)</i></p> <p>They also expect this support to continue throughout their illness: <i>"The team has rescued me a number of times now and I hope they can just keep doing that until transplant" (parent)</i></p> <p>However, some parents raised that they were overwhelmed by the number of different people involved with their care and not being empathetic: <i>"All I can remember is five people in a room watching me cry, feeling like a real goose." (parent)</i> <i>"I felt quite teary. It was really, really draining ...I remember people asking me questions...and I didn't know who to ask, and I just felt like I was surviving." (parent)</i> <i>"We came in on the three days. It was sort-of like people came and saw you and then you waited for the next lot of people to come through, and then you waited some more...It was such a nightmare... It was the actual doing, it was terrible." (mother)</i></p>			
Sub-theme 2: from family and carers					
1 study (Widerman 2004)	1 study using interviews	<p>In 1 study conducted in the USA with 16 people diagnosed with cystic fibrosis as adults, moderate or seriously ill participants admitted feeling self-pity and wanting sympathy, particularly from family members and caregivers. <i>"Once in a while, I'd like someone to feel sorry for me" (woman diagnosed with cystic fibrosis as adult)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p>	VERY LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		Participants said they envied and resented the concern of the public for cystic fibrosis "poster children". But because cystic fibrosis is not outwardly apparent in most adults, participants felt their families, friends and coworkers underestimated its impact (authors' comment).	Saturation	Not saturated	
Sub-theme 3: online support					
1 study (Kirk 2016)	1 study using information from online forums	<p>1 study conducted in the UK explored how online peer support is used by young people and parents (n=279).</p> <p>The results suggested that online peer support groups are useful for managing emotions, as group discussion provided an outlet for parents and young adults to express their emotions:</p> <p><i>"Hey, You are certainly not alone! I think everyone with CF has felt like tha sometimes. I know for a fact I hve felt like why do I bother but I tend to do it when i'm well bcoz i cant see any differece when i take my tablet sna if i miss them but I've learnt now that i have to do my nebs and stuff"</i></p> <p>Some parents tried to validate their identity and to justify that they were good parents</p> <p><i>"my daughter is 15 now but i remember like it was yesterday going through the stuff u r now and to be honest u sound like a great mum and my only advice is a mum knows best just listen to your heart and u wont go far wrong."</i></p> <p>Although the groups were a place where negative emotions could be expressed, it appeared that there were boundaries to this. Indeed, the online group was not always seen as being an appropriate place to discuss certain experiences and feelings:</p> <p><i>"This is a short post. I ashamed to say I often feel the same (perhapes manage it a little better though). I dont want to open up on this subject on here though. Feel free to email me on</i></p>	Limitation of evidence	Minor limitations	LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>XXX" (Parent)</p> <p>Participants advised parents to take an assertive stance and question medical decision making.</p> <p><i>"I just wanted to say that I think your attitude towards your son's care is fantastic. I know a lot of parents struggle to stand up to/question medical staff (including my mum when I was younger) so it's great that you have already managed to gain the confidence to do it when your son is still at such a young age."</i></p>			

1 Table 30: Summary of clinical evidence (GRADE-CERQual): Theme 5. Social support

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: support groups					
2 studies (D'Auria 2000, Whyte 1994)	2 studies using interviews	<p>In 1 study conducted with 15 young people and adults with cystic fibrosis, participants commented on the usefulness of attending support groups with other people with cystic fibrosis.</p> <p><i>"There were 16 cystic fibrosis patients on the floor that holds maybe 30. We all went out to dinner. That's the kind of thing that balances out even though you miss school and the occasional homecoming dance. You have at least something to balance out and just say, Yeah, I missed that, but I've made a lot of good friends here, too."</i></p> <p>Similarly, in another study conducted in the UK with 4 families of children with cystic fibrosis some parents found that joining support groups was helpful, and parents considered volunteering to work with families who were newly-diagnosed with cystic fibrosis:</p> <p><i>"I would like to work with newly-diagnosed families. I needed</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Not applicable</p> <p>Not saturated</p>	VERY LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<i>much more help with the emotional side of things during the whole of the first year" (mother of a child with cystic fibrosis)</i>			
Sub-theme 2: support from family and friends					
4 studies (Hodgkinson 2002, MacDonald 2010, Tipping 2010, Whyte 1984)	3 studies using interviews and 1 study using both interviews and focus groups	<p>2 studies conducted in the UK with parents of children with cystic fibrosis and 1 study conducted in Australia with parents of young children with cystic fibrosis and HCPs explored the roles of families and friends in providing social support.</p> <p>Mothers found support from various sources including family, partner, friends, cystic fibrosis liaison nurses and their primary health care team. (authors comment)</p> <p>Family was perceived as supportive by some people: "At the time we also had a bit of support unit with my mum and sister in law there and they were taught [physiotherapy techniques] as well for back up and a bit of emotional and moral support for me." (parent) But it was described as unsupportive by others: "My husband who is an angel, he is fantastic, he also wasn't 100% support[ive] in that manner, he left most of it (physiotherapy) for me" (parent)</p> <p>Friends were identified as an important source of support because they could talk about other things than cystic fibrosis: "If I visit my friend who lives round the corner, we don't sit there talking about S, you know, we can have a gossip and just things like that. CF isn't the centre of the conversation all the time, which like, it shouldn't have to be" (mother)</p> <p>A mother of twins with cystic fibrosis found that the help she received from the community support scheme was not helpful: "The help I got was not the same as help from the family. if only my mum had been around...we do miss a granny figure"</p>	Limitation of evidence Coherence of findings Applicability of evidence Saturation	Major limitations Coherent Applicable Unclear	LOW QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>One study conducted in the UK with children and young people with cystic fibrosis, their parents and the healthcare professionals looking after them, explored the benefits of a befriending programme.</p> <p>Befriending was seen as helpful by both parents and young adults. For example a young adult of 15 years was happy in the company of befriender: <i>"it's what I expected, going out having a wee bit of a laugh and when I come back my dad says I'm always happier than when I left home."</i></p> <p>In particular, young adults understood that having a befriender took the pressure of parents:</p> <p><i>"when they first asked me if I wanted a befriender, I just wanted to go through it myself, saves my mum and dad having to do all that stuff."</i> (young person with cystic fibrosis)</p> <p>Parents also recognise that their children might share their emotion with befriender when they sometime struggle to share with them:</p> <p><i>"... this is one of the reasons that the befriender ... plays a role in it ... that builds up a sort of friendship with (son) ... that if he's got any fears like that, hopefully he'll speak tae the befriender."</i> (parent of a befriender)</p> <p>Some criticism of befriending was around the continuity, as befrienders were mostly young people who were in transition between education and employment and continuity was thus difficult:</p> <p>For example 1 young adult was unhappy about the lack of continuity <i>"Don't know what happened to the first one.....I thought it was me.... I could never get in touch with her."</i></p> <p>Similarly, a parent of a 16 year old befriender said <i>"They'd be better if the lassies were a wee bit older, ken they're away on holiday, I dunno what age, she can only be in her 20s,</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>changing jobs, its months since we've seen her."</i></p> <p>In relation to this, a parent shared the same concern <i>"I feel let down and (son) has been let down because he was getting close to her."</i></p> <p>Another difficulty raised is that befrienders also faced challenges in building and boundaries of the relationship: <i>"...It can take a while to get to that stage, I'm now totally comfortable with (child), we can talk about anything..."</i> <i>"He doesn't understand why he can't come to my house, or I can't bring my child along and neither do I."</i></p> <p>Befrienders were also concerned about their lack of knowledge when discussing with parent or young adult about cystic fibrosis. <i>"I would like more training about CF, the parents talk to you in jargon you don't understand – what's IV's?"</i></p>			
Sub-theme 3: peer support/ virtual support					
2 studies (Hilliard 2014, Kirk 2016)	1 study using interviews and 1 study using information from online forums	<p>One study conducted in the UK with 279 young people with cystic fibrosis and their parents explored the use of an online forum as means of social support.</p> <p>While parents mainly used the site to seek specific advice or emotional support (as seen before), young people used it as a social networking site. <i>"hey Tina welcome back! how did you get on with the exams? i'm good thanx, back on the IV's in June. how are you?" (young person with cystic fibrosis)</i> <i>"aww i'm gd. on my lv's atm. halfway there! yay! a week is too long though. i feel like pulling the needle out! glad to hear you're feeling good. i think i need a bronch to! havent had one in ages and sometimes you can just tell you need one! now is one of these times! (but don't tell the doctor!" (young person with cystic fibrosis)</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Unclear</p>	MODERATE QUALITY

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p>Another study conducted in the USA with 15 adults with cystic fibrosis using a mobile application showed that some people saw this as a novel opportunity to network, given prohibitions on face-to face contact.</p> <p><i>“Because people with CF can’t be in the same room as each other, ...being able to see someone else with CF is much more profound than just exchanging emails with some anonymous person.” (Age 28, Male)</i></p> <p><i>“I think that support for the family and friends is important...for people who have CF...talking to significant others of people who have CF.” (Age 28, Male)</i></p> <p>Others could feel discouraged or guilty seeing others “<i>doing better or worse than you</i>”:</p> <p><i>“I don’t like hearing about CF people that aren’t doing well. I have a hard time distancing myself from it. It’s hard having to filter through all this sadness to get kind of connected with someone.” (Age 26, Female)</i></p> <p><i>“I didn’t really like it only because some people had it worse than me and if it kind of brought me down because I felt like this is where I’m heading and I just didn’t like that...So I don’t know if I’d really want to talk to any other people with CF, I don’t want to like be depressed.” (Age 32, Male)</i></p>			

- 1
- 2
- 3

1 **6.5 Economic evidence**

2 No economic evaluations related to information and support in cystic fibrosis were identified
3 in the literature search conducted for this guideline. Full details of the search and economic
4 article selection flow chart can be found in Appendix E and F, respectively.

5 This review question is not relevant for economic analysis because it does not involve a
6 decision between alternative courses of action.

7 **6.6 Evidence statements**

8 **6.6.1 Information needs for people with cystic fibrosis and their parents or carers**

9 **Information at diagnosis**

10 Low to moderate quality evidence from 7 qualitative studies conducted with parents of
11 children with abnormal newborn screening results, children with cystic fibrosis or adults
12 recently diagnosed with cystic fibrosis discussed the information needs prior to and following
13 diagnosis. Both parents and adults wanted to know information about the condition and the
14 future implications of cystic fibrosis. However, some discrepancies were found between
15 those parents who felt they received too much information and those who felt they had to
16 fight to get the information they needed, or to look for information themselves. In balance
17 parents appreciate timely information, at a pace that it is convenient for them. Adults were
18 particularly concerned about not being given information specifically related to them.

19 **Information about treatment**

20 Very low to low quality evidence from 5 qualitative studies conducted with children with cystic
21 fibrosis and parents of children with cystic fibrosis noted the importance of trusting healthcare
22 professionals in relation to the information received about the treatment. They wanted to
23 know about the available treatment options and about alternative care. Parents particularly
24 welcomed practical instructions in relation to the physiotherapy treatment.

25 **Information about management of complications**

26 Low quality evidence from 4 qualitative studies conducted with families of children with cystic
27 fibrosis, adults with cystic fibrosis and healthcare professionals looked at specific
28 complications related to cystic fibrosis. With regards to lung transplant, most parents found
29 information helped them to face the situation, although some parents found it difficult to deal
30 with it. Adults reported they were concerned in relation to fertility issues, but it seemed they
31 sometimes found it difficult and uncomfortable to discuss their worries and questions with the
32 healthcare professionals. To overcome this barrier, some participants suggested this topic
33 should be discussed earlier and these discussions should be part of the routine. Women in
34 particular reported that they would like more information about the consequences of
35 pregnancy and the risks of the actual birth.

36 **Provision of information**

37 Low to moderate quality evidence from 10 qualitative studies with young people and adults
38 with cystic fibrosis, parents of children with cystic fibrosis and healthcare professionals
39 discussed aspects relevant to the provision of information. People valued receiving relevant
40 information for them, accessibility to information resources, the use of simple language and
41 privacy. In relation to the content, people wanted to get general information that is practical
42 rather than technical information. They noted that information can sometimes be upsetting.

1 The amount of information needed was discussed. People have different needs, those who
2 required lots of information and those who preferred information to be delivered at intervals.
3 Ultimately, people seemed to prefer timely information that helps them to prepare. With
4 regards to the format, people commented on different possibilities, such as written
5 information, specific groups and DVDs. Many people felt reluctant to use online resources or
6 mobile applications, as a way of getting information or communicating with healthcare
7 professionals or the family.

8 **Perceived benefits of receiving information**

9 Low to moderate quality evidence from 5 qualitative studies conducted with parents of
10 children with cystic fibrosis and with people with cystic fibrosis reported on the benefits of
11 receiving information. Most participants reported that information helped them to understand
12 the condition and gave them a sense of control. It helped them to prepare for treatments and
13 to ask meaningful questions when needed. Information was seen as reassuring. Both adults
14 with cystic fibrosis and parents of children with cystic fibrosis saw information as a source of
15 hope.

16 **6.6.2 Support needs for people with cystic fibrosis and their parents or carers**

17 **Support at specific stages**

18 Low to moderate quality evidence from 2 qualitative studies conducted with people with
19 cystic fibrosis, parents of children with cystic fibrosis, and healthcare professionals reported
20 on the support needed at important stages during the condition, including newborn
21 screening, diagnosis and end of life care. Parents reported feeling overwhelmed after the
22 diagnosis and wanting to contact healthcare professionals for advice, expertise and
23 reassurance, although some felt difficult to approach professionals. Similarly, people with
24 cystic fibrosis and their parents indicated they needed support to address the end of life care.

25 **Support with treatment**

26 Very low to low quality evidence from 10 qualitative studies conducted with children and
27 young people with cystic fibrosis, parents of children and healthcare professionals explored
28 the support needed in relation to treatment. Parents and young people with cystic fibrosis
29 noted the lack of continuity of care and emphasised that having a consistent set of providers
30 would be beneficial. Parents would also appreciate GPs to be more involved in the care of
31 their child. Parents seek help in relation to treatment either from professionals or other
32 parents. They particularly noted the need for help with practical aspects in relation to nutrition
33 and physiotherapy. Finally, the role people should take in decision making regarding
34 treatment was discussed by several participants. Parents viewed themselves as being
35 involved in the process, although they acknowledged their decisions were based on the
36 recommendations of healthcare professionals because of the seriousness of the condition.
37 Children felt they were not very involved in decision making and they noted they would like to
38 have a more active role. In relation to this, parents said they did not want their children to be
39 involved, or that they were waiting for the right time.

40 **Management of complications**

41 Very low quality evidence from 2 qualitative studies conducted with young people with cystic
42 fibrosis, parents of children with cystic fibrosis and healthcare professionals explored the
43 issue of fertility and the support needed in this regard. They all agreed the issue of fertility is
44 often neglected as it is not considered life-threatening. Young people in particular highlighted
45 they need healthcare professionals to recognise the emotional impact of infertility.

1 **Psychological and emotional support**

2 Very to moderate quality evidence from 5 qualitative studies conducted with young people
3 with cystic fibrosis, people diagnosed with cystic fibrosis as adults, parents of children with
4 cystic fibrosis and healthcare professionals explored the importance of emotional and
5 psychological support. Parents highlighted they needed timely reassurance and continuous
6 support from staff. Feeling emotionally supported by staff was very encouraging for them and
7 helped build a good relationship with them. Some parents had experienced lack of empathy
8 and this made them feel misunderstood. Similarly, people that had been diagnosed as adults
9 felt the need for emotional support and sympathy from their relatives and friends. The use of
10 online groups was described as helpful for managing emotions.

11 **Social support**

12 Very low to moderate quality evidence from 7 qualitative studies conducted with young
13 people and adults with cystic fibrosis, parent of children with cystic fibrosis and healthcare
14 professionals explored the benefits of social support. Young people with cystic fibrosis and
15 parents of children with cystic fibrosis described attending support groups as useful.
16 Furthermore, the use of virtual support, either online support or mobile applications, was also
17 found to be very useful to meet people with the same condition. This is particularly useful
18 given the recommendations to avoid face to face contact and risk of cross infection.
19 However, some felt it could be discouraging seeing other people doing better or worse than
20 them.

21 Families and friends were seen as supportive and unsupportive. A befriending programme
22 was described as helpful by both young people with cystic fibrosis and their parents as it took
23 the pressure off parents and allowed for young people to share their emotions. However they
24 were concerned about the lack of continuity.

25 **6.6.3 Economic evidence statements**

26 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

27 **6.7 Evidence to recommendations**

28 **6.7.1 Relative value placed on the themes considered**

29 The aim of this review was to identify the information and support that should be provided to
30 people with cystic fibrosis and their parents and carers. Although there were many themes in
31 the literature, the Committee identified those that they thought would be important when the
32 protocol was developed. They agreed that the following themes would provide useful
33 perspectives: provision and timing of psychosocial support, regular individualised
34 assessment, support groups, support at school, information at the time of diagnosis, social
35 media, care planning and social services and benefits.

36 **6.7.2 Consideration of clinical benefits and harms**

37 While acknowledging the limitations of the evidence, the Committee agreed that the themes
38 and sub-themes identified in the literature were useful and relevant.

39 The Committee discussed the evidence regarding information and support needs around the
40 time of diagnosis. The literature reported that people with cystic fibrosis and their parents
41 want to receive information following diagnosis, mostly regarding information about the
42 condition and the future implications. In addition, the Committee emphasised that receiving
43 the diagnosis of cystic fibrosis is a stressful event and it is important that information is
44 delivered by a specialist in cystic fibrosis.

1 As highlighted in the evidence, the Committee acknowledged the requirement for honest
2 dialogue between people with cystic fibrosis, their parents or carers and health care
3 professionals. They agreed information should be relevant and adequate for the person, but
4 not overwhelming to the receivers. They also agreed general information should be readily
5 available, but the detail people want may differ and should be based on individual's needs.

6 The Committee discussed it is important to offer ongoing and staggered information, as
7 reported in the evidence. They agreed that the person or their family may require information
8 at specific times, for example before and after diagnosis, or for specific topics, such as
9 treatment for cystic fibrosis-related complications (liver disease, cystic fibrosis diabetes, lung
10 transplant or fertility issues). They suggested to have an honest and open dialogue on the
11 symptoms of cystic fibrosis and possible complications. They also agreed that information on
12 coping with anxiety from cystic fibrosis is important.

13 The Committee noted that all people with cystic fibrosis have a pathway of care, which
14 describes the care they are receiving and that they should be able to have access to it.

15 The Committee noted it is important to deliver information through different formats and the
16 use of plain English for the information to be useful, as suggested in the literature, taking into
17 account individual variances in literacy and comprehension levels.

18 The Committee underlined that psychological support is important for people with cystic
19 fibrosis and their families or carers. The literature highlights the time of diagnosis and when
20 approaching end of life as key periods when psychological support is needed. They
21 highlighted additional support may be needed during times of change, such as transition from
22 education to work, living independently or having a family (the latter due to fertility
23 complication and issues which may arise in pregnancy for expectant mothers with cystic
24 fibrosis).

25 The Committee discussed the use of online support, including social media, as reported in
26 the literature. They expressed potential concerns that social media is unregulated and so
27 information may be inaccurate. However, they highlighted that monitoring and regulating
28 online discussion forums is far beyond the remit of healthcare professionals. Based on this,
29 they agreed it is important to advise people with cystic fibrosis and their parents to be aware
30 of the potential issues when they use internet to look up for health information. However,
31 information and support available from trustworthy sources of information, such as the NHS,
32 CF Trust (and its international counterparts) could be signposted to. They also agreed it is
33 important to give the person with cystic fibrosis and the families and opportunity to discuss
34 what they have read.

35 Some reports in the literature suggest that support groups can be perceived as beneficial by
36 some people with cystic fibrosis. However, the Committee strongly recommends these
37 groups do not take place face to face in order to prevent cross-infection which can lead to
38 further complications. Information about online support groups would be beneficial.
39 Facilitating access to parent or carer only face to face social support groups can be
40 considered if there is a demand. But with these the risk to their children meeting socially
41 must be emphasised.

42 **6.7.3 Consideration of economic benefits and harms**

43 This review question is not relevant for economic analysis because it does not involve a
44 decision between alternative courses of action. Even so, there are considerations for the
45 resources and costs providing information and support may entail.

46 Whilst there are aspects of providing information and support which have opportunity costs,
47 such as the staff time and some of the different communication formats that may be useful in
48 this population, these are typically small and would ordinarily be considered within the
49 provision of standard services and care. Good communication is recognised as important

1 generally within healthcare provision and that patient care can suffer as a result of poor or
2 ineffective communication. The Committee considered that, overall, their recommendations
3 regarding information and support would have a minimal resource impact and that they
4 would promote the cost-effective use of NHS resources.

5 **6.7.4 Quality of evidence**

6 Moderate to very low-quality evidence was presented in this review as assessed by GRADE-
7 CERQual. Below the main reasons leading to downgrading the evidence are described.

8 The methodological limitation of the studies, especially the inadequate reporting on
9 processing and analysis of the data. Some of the studies reported were descriptive when
10 thematic analysis would have been more appropriate and informative. The studies where
11 thematic analysis was done, the authors did not always report in detail how findings or
12 themes were derived or emerged from the data in their research.

13 Another methodological limitation was the inadequate information on the researcher's role in
14 sample recruitment, data collection or data analysis. Few studies clearly reported the
15 relationship between researchers, interviewers and the respondents, whether the
16 researchers had a pre-understanding about the topic or the possible influence of that in data
17 collection and analytical process. Lack of verification of findings was not reported either in
18 any of the studies.

19 Downgrading of evidence was also related to data saturation, applicability and incoherence.
20 Some of the studies did not report whether saturation was achieved in terms of data
21 collection or data analysis. It was difficult to ascertain from the information reported in those
22 studies whether all possible views had been explored. Some other studies were done in an
23 isolated group of population in other countries and were not applicable to UK population.
24 Incoherence was also a factor in downgrading the quality of evidence in few of the studies.

25 **6.7.5 Other considerations**

26 No equality issues were identified by the Committee for this review question.

27 The recommendations related to this evidence review were based on the evidence and the
28 Committee's clinical experience. In addition, the Committee acknowledged some of the
29 principles about communication and information in the NICE guideline on "patient experience
30 in adult NHS services: improving the experience of care for people using adult NHS services"
31 (NICE CG138) could also be applied here. Therefore they agreed to draft a recommendation
32 which directed readers to this guideline. The Committee concluded that a research
33 recommendation was not needed as the guidance provided in the guideline combined with
34 an individual healthcare professionals experience is sufficient to provide the necessary
35 support and information. Given the limitations in research funding it was determined that this
36 should not be a priority for further research.

37 **6.7.6 Key conclusions**

38 The Committee concluded that receiving a diagnosis of cystic fibrosis is a stressful event. It
39 is important to ensure that information and support is provided by healthcare professionals
40 with expertise in cystic fibrosis. Information should be tailored to the individual needs and
41 given on an ongoing and staggered basis. They agreed that information and support may
42 also be needed regarding new treatments, management of comorbidities and fertility.

1 **6.8 Recommendations**

- 2 **4. Provide people who are newly diagnosed with cystic fibrosis and their family**
3 **members or carers (as appropriate) with opportunities to discuss their concerns.**
- 4 **5. Information and support should be provided by healthcare professionals with**
5 **expertise in cystic fibrosis.**
- 6 **6. Provide people with suspected or diagnosed cystic fibrosis and their family**
7 **members or carers (as appropriate) with relevant information in clear English and**
8 **opportunities for discussion on topics such as:**
- 9 • their diagnosis
 - 10 • monitoring of their condition
 - 11 • management choices for their condition
 - 12 • possible or existing complications or comorbidities
 - 13 • implications for independent living.
- 14 **7. Provide people with cystic fibrosis and their family members or carers (as**
15 **appropriate) with information about their care pathway (for example, by providing**
16 **them with copies of correspondence).**
- 17 **8. Give information to people with cystic fibrosis and to family members or carers in**
18 **ways that are individually appropriate. Avoid jargon and use formats that they**
19 **prefer, for example:**
- 20 • face-to-face discussions
 - 21 • written information (such as leaflets)
 - 22 • any digital media and reliable internet sources that are available.
- 23 **9. When appropriate, provide people with cystic fibrosis and their family members or**
24 **carers opportunities for discussion with relevant expert professionals on:**
- 25 • available resources and support, such as local support and advocacy
26 groups, including those aimed specifically at children, young people,
27 adults, parents and carers
 - 28 • implications of the condition for school and education
 - 29 • career planning
 - 30 • transition to adult care
 - 31 • fertility and contraception
 - 32 • pregnancy and parenting
 - 33 • organ transplantation.
- 34 **10. Provide people with cystic fibrosis with information about how to contact other**
35 **people with cystic fibrosis without risking cross-infection (see Prevention of cross**
36 **infection), for example, by directing them to online support groups.**
- 37 **11. For more information on communication, providing information and shared**
38 **decision-making in adult NHS services, see the NICE guideline on patient**
39 **experience in adult NHS services.**

- 1 **12. Be aware that people with cystic fibrosis and their family members or carers may**
2 **need emotional and psychological support (see psychological assessment), in**
3 **particular:**
- 4 • at diagnosis
 - 5 • at times of transition (for example, when moving from education to work
 - 6 or when living independently for the first time)
 - 7 • during pregnancy
 - 8 • to deal with complications, such as infertility
 - 9 • when approaching the end of life.

7 Service delivery

7.1 Service configuration

Review question: What is the effectiveness of different models of care (for example, specialist centre, shared care [delivered by a Network CF Clinic which is part of an agreed designated network with a Specialist CF Centre], community, telehealth and/or home care) for people with cystic fibrosis?

7.1.1 Introduction

There is a wide range of ways in which care for people with cystic fibrosis is organised and delivered.

In the UK, care for adults with cystic fibrosis is primarily concentrated in specialist cystic fibrosis centres.

Children with cystic fibrosis may be cared for solely by a specialist cystic fibrosis centre but there are some circumstances where they may also be seen in conjunction with a team in a local hospital as part of a paediatric network (shared-care clinic). Such an arrangement reduces disruption to schooling and improves the family's local support.

In addition there are other models of care and service which people with cystic fibrosis may utilise including telemedicine, clinical review and treatment at home, outreach care and community care by primary health care professionals.

Each of these models provide a different patient experience and may be associated with different outcomes and cost effectiveness. Additionally, limitations in a patient's socio-economic background and local geography have significant impacts on how care can be arranged and delivered.

7.1.2 Description of clinical evidence

The aim of this review was to assess the effectiveness of different models of care (for example specialist centre, shared care, community care, telehealth and home care) for the care of people with cystic fibrosis.

For the purpose of this review, the different models of care were defined as follows:

- Specialist centre:
 - normally with >100 patients and minimum of >50 patients
 - commissioned by NHS England to provide care for people with cystic fibrosis
 - will have the core MDT available.
- Shared care (delivered by a network clinic):
 - local hospital looking after a small number of children with cystic fibrosis
 - has input from a specialist centre at least twice a year, which is responsible for the standard of care
 - may not have the core MDT and may not be full time.
- Community care:
 - community nurses of the region who look after people with cystic fibrosis and will administer treatments such as home intravenous (IV) antibiotics
 - normally delivered by health visitors and school nurses.
 - might include palliative services.

- 1 • Home care:
- 2 ○ care that would normally be given at hospital which is given at home
- 3 ○ delivered by cystic fibrosis specialist (e.g. nurse, dietician, physiotherapist).
- 4 • Outreach:
- 5 ○ specialist centre team conduct a clinic in a local hospital.
- 6 • Telemedicine:
- 7 ○ involves videoconferencing patients and remote monitoring.

8 As per protocol, studies that assessed the effectiveness of different models of care were
9 eligible for inclusion in this review if they were randomised controlled trials or comparative
10 cohort studies that were conducted in Western countries. Studies based on registry and audit
11 data from the UK were also eligible for inclusion. Conference abstracts of RCTs were
12 considered if RCTs were unavailable. Prospective studies were prioritized over retrospective
13 studies, therefore retrospective studies were excluded if there were prospective studies
14 which covered the same intervention and outcomes. Evidence from questionnaires
15 conducted within cross sectional studies was included (for example, for quality of life, or
16 patient satisfaction).

17 For full details see review protocol in Appendix D.

18 Two Cochrane systematic reviews were included:

- 19 • Balaguer 2015 evaluated the effectiveness of IV antibiotic therapy administered at home
20 compared to IV antibiotic therapy administered in hospital, 1 RCT was included from this
21 review (Wolter 1997).
- 22 • Goldbeck 2014 evaluated the effectiveness of psychological interventions compared to
23 standard care, 1 RCT was included from this review (Wilkinson 2008)
- 24 • In addition, 4 prospective cohort studies (Donati 1987, Esmond 2006, Riethmueller 2002,
25 van Koolwijk 2002), 3 retrospective cohort studies (Bosworth 1997, Thomas 2008,
26 Finkelstein 1992) and 2 cross sectional studies (Thomas 2006, Walters 1994) were
27 included.

28 Where possible, data and risk of bias assessment were taken directly from the Cochrane
29 systematic reviews. Individual studies were retrieved for completeness and accuracy, and
30 were also checked for additional outcomes of interest.

31 The size of the studies ranged from 6 to 746 participants. Three studies included adults only
32 (Esmond 2006, Wilkinson 2008, Wolter 1997), 1 study included people aged ≥ 15 years
33 (Walters 1994), 1 study included people aged ≥ 12 years (Donati 1987), 5 studies included
34 children, young people and adults (Bosworth 1997, Finkelstein 1992, Riethmueller 2002,
35 Thomas 2006, Thomas 2008), 1 study included children and young people (Van Koolwijk
36 2002).

37 Three studies were conducted in the UK (Esmond 2006, Walters 1994, Wilkinson 2008), 3 in
38 the USA (Bosworth 1997, Donati 1987, Finkelstein 1992), 3 in Australia (Thomas 2006,
39 Thomas 2008, Wolter 1997), 1 in Germany (Riethmueller 2002), 1 in the Netherlands (van
40 Koolwijk 2002).

41 Four studies (Bosworth 1997, Donati 1987, Esmond 2006, Wolter 1997) compared the
42 effectiveness of IV antibiotics for treatment of pulmonary exacerbations administered at
43 home and in the hospital. One study (Riethmueller 2002) compared the effectiveness of IV
44 antibiotics for people with a chronic infection administered at home and in the hospital. Three
45 studies compared centre care to shared care (Thomas 2006, Thomas 2008, Van Koolwijk
46 2002). Two studies (Thomas 2008, Van Koolwijk 2002) compared centre care to local care
47 (below UK standards). One study (Thomas 2008) compared shared care above UK
48 standards to shared care at UK standards. One study (Walters 1994) compared centre care
49 to usual care (by non-cystic fibrosis specialists). One study (Finkelstein 1992) compared a

1 telemedicine intervention (daily recording of physical measurements and symptoms) to usual
2 care. One study (Wilkinson 2008) compared a telemedicine intervention (weekly
3 videoconferences for clinical assessment and psychological support) to usual care.

4 A summary of the included studies is presented in Table 159. See study selection flow chart
5 in Appendix F, study evidence tables in Appendix G, list of excluded studies in Appendix H,
6 forest plots in Appendix I, and full GRADE profiles in Appendix J.

7 7.1.3 Summary of included studies

8 A summary of the studies that were included in this review are presented in Table 31.

9 **Table 31: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
Cochrane reviews				
Balaguer 2015 Cochrane SR	Comparison 1: home-based care (IV antibiotic therapy administered at home) versus hospital-based care (IV antibiotic therapy administered in hospital) (Wolter 1997)	Children, young people and adults with CF	<ul style="list-style-type: none"> • Change in FEV₁ (% predicted) • Weight gain (kg) • Not reported: • Mortality • Patient and carer satisfaction • Lung Index Clearance • Time to next pulmonary exacerbation • Quality of life • Frequency of cross-infections • Staff experience • Adherence to treatment 	
Goldbeck 2014 Cochrane SR	Comparison 1: Psychological intervention versus usual care (Wilkinson 2008)	Children, young people and adults with CF	<ul style="list-style-type: none"> • Change in quality of life • Not reported: • FEV₁ % predicted • Mortality • Patient and carer satisfaction • Lung Index Clearance • Time to next pulmonary exacerbation • Nutritional status • Frequency of cross-infections • Staff experience • Adherence to treatment 	

Study	Intervention/ Comparison	Population	Outcomes	Comments
Primary studies included in the Cochrane SR				
Wilkinson 2008 (UK) RCT	<p>Intervention: telemedicine</p> <ul style="list-style-type: none"> weekly videoconferences from home for a clinical assessment, psychological support and the opportunity for discussion with any member of the multidisciplinary team <p>Control: Standard care</p>	<p>N=7 adults with CF on a transplantation list (16 recruited, 11 completed the baseline assessment and 7 completed the study)</p> <ul style="list-style-type: none"> Telemedicine: n=4 Control arm: n=3 <p>Age range of people who were randomized (median): 21 to 41 (27)</p>	<ul style="list-style-type: none"> Change in quality of life 	<p>Included in Goldbeck 2014 SR</p> <p>Follow-up: 6 months</p>
Wolter 1997 (Australia) RCT and cross-over open study	<p>Intervention: home-based care spent 2 - 4 days in hospital before discharge</p> <p>people with CF were taught to prepare and administer their own intravenous antibiotics</p> <p>home visits were conducted</p> <p>Comparison: hospital-based care whole treatment was administered in the hospital</p>	<p>N=17 adults with CF with an infective exacerbation (31 admissions: home care: n=13 admissions hospital care: n=18 admissions)</p> <p>Age range (median): 19-41 (22)</p>	<ul style="list-style-type: none"> Change in FEV₁ (% predicted) at 21 days Weight gain (kg) at 10 days 	<p>Included in Balaguer 2015</p> <p>Follow-up: 21 days for FEV₁% predicted, 10 days after treatment for weight gain</p>
Additional primary studies				
Home care and community care: IV antibiotics for treatment of pulmonary exacerbations				
Bosworth (1997) USA Comparative cohort study	<p>Intervention: Home-based care</p> <ul style="list-style-type: none"> Prior to receiving home care, people with CF stayed in the hospital for up to 4 days IV antibiotics and chest physiotherapy administered at home either by self or family with the same 	<p>N= 40 people with CF who required IV antibiotic therapy for a pulmonary exacerbation (59 courses)</p> <ul style="list-style-type: none"> Home group: n=19 people with CF (27 courses). Age range: 7-31. Mean (SEM): 18.8 (1.2) Hospital group: 	<ul style="list-style-type: none"> Time to next exacerbation 	<p>Although not explicitly reported, likely to be a retrospective study.</p> <p>Mean follow-up: 18 days</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>frequency as in hospital</p> <ul style="list-style-type: none"> • Home visits from nurses at least once a week <p>Comparison: hospital-based care</p> <ul style="list-style-type: none"> • IV antibiotics in the hospital • Chest physiotherapy administered in the hospital 4 times a day 	<p>n= 21 people with CF (32 courses). Age range: 8-29. Mean (SEM): 17.05 (0.9)</p>		
<p>Donati 1987 (USA) Comparative prospective cohort study</p>	<p>Intervention: Home-based care</p> <ul style="list-style-type: none"> • Daily visits from nurses <p>Comparison: hospital-based care</p> <ul style="list-style-type: none"> • IV antibiotics administered at the hospital 	<p>N=64 people with CF (82 treatments) who required IV antibiotic therapy for a pulmonary exacerbation Age: ≥12 years</p> <ul style="list-style-type: none"> • home group: n= 26 people (41 treatments) • hospital group: n=38 people (41 treatments) 	<ul style="list-style-type: none"> • Change in FEV₁ (% predicted) • Change in weight (kg) 	<p>Follow-up: 18 days</p>
<p>Esmond 2006 (UK) Quasi-experimental design: prospective comparative cohort</p>	<p>Intervention: home-based care</p> <ul style="list-style-type: none"> • Home IV antibiotic treatment. • All patients who chose home therapy had previously self-administered IV antibiotics at home. • People performed their own chest physiotherapy <p>Comparison: hospital-based care</p> <ul style="list-style-type: none"> • IV antibiotic treatment administered at the hospital. • Chest physiotherapy performed by experienced 	<p>N=28 adults with CF experiencing an acute respiratory exacerbation. (30 treatments) Age: ≥18 years</p> <ul style="list-style-type: none"> • Home group: n=15 treatments • Hospital: n=15 treatments 	<ul style="list-style-type: none"> • Change in FEV₁ (% predicted) • Change in BMI <p>Change in quality of life measured with CFQoL questionnaire (Gee et al. 2000)</p>	<p>Follow-up: 15 days</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	respiratory physiotherapists.			
Home care/ community care versus hospital: IV antibiotics for patients with chronic infection				
Riethmueller 2002 (Germany) Prospective comparative cohort study	<p>Intervention: home-based care</p> <ul style="list-style-type: none"> • IV antibiotic treatment at home • People did their daily training and were supervised once per week by a physiotherapist specialized in CF. • People were offered a visit by a specialized nurse if intravenous line or other problems occurred. <p>Comparison: hospital-based care</p> <ul style="list-style-type: none"> • IV antibiotic treatment in hospital. • Two daily courses of supervised physiotherapy (1 h). • Diets were supervised by a dietitian specialized in CF care. 	<p>N=36 people with CF (58 courses)</p> <ul style="list-style-type: none"> • Home group: n=17 people (30 courses). Mean (SD) age: 16 (5) • Hospital group: n=19 people (28 courses). Mean (SD) age: 15 (4) <p>Inclusion criteria included P. aeruginosa in sputa over a time period of more than 6 months before entering, good compliance and regular home physiotherapy.</p>	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in weight (kg) • Change in weight for height (%) 	Follow-up: 14 days
CF centre care versus. shared care or local care				
Thomas 2006 (Australia) Cross-sectional study	<p>Intervention: CF centre care</p> <ul style="list-style-type: none"> • Children are reviewed at least 3 times a year and have full access to the MDT. <p>Comparison: shared care</p> <ul style="list-style-type: none"> • Children are managed by their local paediatrician or general practitioner and 	<p>N=162 people with CF (and/or their parents when relevant) Age range: 2-19</p> <ul style="list-style-type: none"> • Specialist centre: n=91 • Shared care: n=71 	<ul style="list-style-type: none"> • HRQOL measured with CFQ questionnaire (Quittner et al. 2005) 	<p>In the paper, different terminology is used compared to this review:</p> <ul style="list-style-type: none"> • CF centre care is called "Cystic Fibrosis Centre (CFC)" in the study • Shared care is called "Cystic Fibrosis Outreach

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>local hospital, and they also attend outreach clinics visited by CFOS.</p> <ul style="list-style-type: none"> Regional staff are invited to attend the clinics. Outreach clinics occur twice per year except for 1 site, which has 1 clinic and 2 tele health clinics per year. 			<p>Service (CFOS)" in the study</p>
<p>Thomas 2008 (Australia) Retrospective comparative cohort study</p>	<p>Intervention: CF centre care</p> <ul style="list-style-type: none"> All care is provided by the Cystic Fibrosis Centre (CFC) Admission to the CFC when required Outpatient review at CFC 3 or more times per year <p>Comparison 1: shared care, review 3+times a year (above UK standards)</p> <ul style="list-style-type: none"> Children living in regional centres and attending Cystic Fibrosis Outreach Service (CFOS) as well as attending CFC regularly. Admission to CFC or local hospital with local hospital care provided by local paediatrician. Outpatient review by CFC or CFOS 3 or more times per year. Children attending outreach clinics are also managed by their 	<p>N=150 people with CF born between 1982 and 2002. Age range: 0-20</p> <ul style="list-style-type: none"> Specialist centre: n=74 Shared care review 3+ times a year: n=21 Shared care review 2+ times a year: n=37 Local care: n=18 	<ul style="list-style-type: none"> Change in FEV₁ % predicted from first to last per year FEV₁ % slope per year 	<p>273 people were included in the study, however the analysis on FEV₁ % predicted focused on 150 people. Follow-up: 3 years Only longitudinal data on FEV₁ were extracted; cross-sectional data on FEV₁, height z scores and weight z scores were not extracted In the paper, different terminology is used compared to this review:</p> <ul style="list-style-type: none"> CF centre care is called "LOC1" in the study Shared care, review 3+times a year, is called "LOC2" in the study Shared care, review 2+

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>local paediatrician or general practitioner.</p> <p>Comparison 2: shared care, review 2+times a year (at UK standards)</p> <ul style="list-style-type: none"> • Care is predominantly provided by the local paediatrician with consultation with CFC • Admission to local hospital with care provided by local paediatrician. • Outpatient review by CFOS at least twice a year <p>Comparison 3: local care</p> <ul style="list-style-type: none"> • Involvement by CFC or CFOS once a year or no CFC/CFOS involvement. • Includes children seen by respiratory physicians but with no CFC or CFOS multidisciplinary health care involvement • Alternatively, care provided by local paediatrician or general practitioner or unknown 			<p>times a year, is called “LOC3” in the study</p> <ul style="list-style-type: none"> • Local care is called “LOC4” in the study
<p>Van Koolwijk 2002 (The Netherlands) Prospective comparative cohort study</p>	<p>Intervention: CF centre care</p> <ul style="list-style-type: none"> • Patients receive their treatment completely in the centre • Regular visits at minimum intervals of 3 	<p>N= 105 young people and children with CF. Age range: 5-17:</p> <ul style="list-style-type: none"> • CF centre care: n=41. Mean (SEM) age: 10.8 (0.5) years • Shared care: 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in BMI (kg/m²) 	<p>In the paper, similar terminology to this review is used:</p> <ul style="list-style-type: none"> • CF centre care is called “centralized care” in the

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>months</p> <p>Comparison 1: Shared care</p> <ul style="list-style-type: none"> Includes a half-yearly visit to the Centre (annual check-up and an MDT outpatient clinic visit) combined with regular visits to the local paediatrician The local paediatrician comes to the centre during the annual check-up and participates in the multidisciplinary consultation Regular visits at minimum intervals of 3 months <p>Comparison 2: Local care</p> <ul style="list-style-type: none"> Patients visit the centre only once a year at the annual check-up, but remain fully treated at their local hospitals. The local paediatrician comes to the centre during the annual check-up and participates in the multidisciplinary consultation Regular visits at minimum intervals of 3 months 	<p>n=41. Mean (SEM) age: 10.7 (0.5) years</p> <ul style="list-style-type: none"> Local care: n=23. Mean (SEM) age: 9.4 (0.5) years 		<p>study</p> <ul style="list-style-type: none"> Shared care is called the same ("shared care") in the study Local care is called "local care" in the study
CF centre versus usual care by non-specialists in CF				
Walters 1994 (UK) Cross-sectional study	<p>Intervention: CF centre care</p> <ul style="list-style-type: none"> Large special cystic fibrosis clinics 	<p>N= 746 people with CF aged ≥15 years</p> <ul style="list-style-type: none"> Specialist centre: n=494 General clinic: 	<ul style="list-style-type: none"> Patient satisfaction 	<p>Questionnaires were sent to all 1052 members of the Association of Cystic Fibrosis</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Comparison: Usual care <ul style="list-style-type: none"> • Non-CF general clinics at local hospitals 	n=252		Adults.
Telemedicine				
Finkelstein 1992 (USA) Retrospective comparative cohort study (follow-up of a RCT)	Intervention: Home monitoring with diary recording <ul style="list-style-type: none"> • Daily recording of physical measurements and symptoms • Diary sent to the data coordinating centre weekly for analysis. Comparison: usual care <ul style="list-style-type: none"> • No diary recording, no home monitoring 	N=50 people with CF <ul style="list-style-type: none"> • Intervention: n=25 • Control: n=25 Age range: 6 to 43 years	<ul style="list-style-type: none"> • Change in predicted FEV₁ (%) at 4 years 	Participants were randomly selected from a bigger RCT Follow-up: 4 years

1 Abbreviations: CF-Cystic Fibrosis; IV-intravenous; FEV-forced expiratory volume in 1 second; BMI-Body Mass
 2 Index; QoL-Quality of Life; HRQOL- Health-Related Quality of Life; CFC-Cystic Fibrosis Centre; CFOS-Cystic
 3 Fibrosis Outreach Service; LOC-Level of Care; RCT-Randomised Controlled Trial.

4 7.1.4 Clinical evidence profile

5 The summary clinical evidence profiles for this review question are presented in Table 32 to
 6 Table 40.

7 7.1.4.1 Home-based care

8 **Table 32: Summary clinical evidence profile: Comparison 1.1. Home versus hospital**
 9 **care for the administration of IV antibiotics in people with CF experiencing**
 10 **an acute pulmonary exacerbation**

Comparison 1.1. Home care compared to hospital care for the administration of IV antibiotics for people with CF experiencing an acute pulmonary exacerbation						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of treatments (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Hospital care for the administration of IV antibiotics	Home care for the administration of IV antibiotics				
Lung function: change in FEV ₁ % predicted Scale from: 0 to	The mean change in FEV ₁ % predicted in the hospital	The mean change in FEV ₁ % predicted in the home care groups was		31 ^a (Wolter 1997) ¹	⊕⊕⊕ ⊖ very low ^{2,3}	

Comparison 1.1. Home care compared to hospital care for the administration of IV antibiotics for people with CF experiencing an acute pulmonary exacerbation						
100. Follow-up: 21 days	care group was 7	3 lower (13.61 lower to 7.61 higher)				
Lung function: change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: mean 18 days	The mean change in FEV ₁ % predicted in the hospital care group was 12.3 ^c	The mean change in FEV ₁ % predicted in the home care groups was 5.60 lower (12.29 lower to 1.09 higher) ^a		63 ^b (Donati 1987)	⊕⊕⊕ ⊖ very low ⁴	
Lung function: change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 15 days	The mean change in FEV ₁ % predicted in the hospital care group was 5.1	The mean change in FEV ₁ % predicted in the intervention groups was 3.1 lower (6.93 lower to 0.73 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{4,5}	
Patients starting next course of antibiotics more than 12 weeks after completing the previous course (proxy outcome for time to next exacerbation) Follow-up: mean 18 days	875 per 1000	481 per 1000 (315 to 726)	RR 0.55 (0.36 to 0.83)	59 ^e (Bosworth 1997)	⊕⊕⊕ ⊖ very low ⁶	
Weight change (kg) Follow-up: mean 18 days	The mean weight change in the hospital care group was 1.6 ^c	The mean weight in the home care groups was 1.10 lower (4.29 lower to 2.09 higher) ^a		74 ^b (Donati 1987)	⊕⊕⊕ ⊖ very low ⁶	
Weight change (kg) Follow-up: 10 days post treatment	The mean weight change in the hospital care group was 0.7	The mean weight change (kg) in the home care groups was 0.5 lower (8.06 lower to 7.06 higher)		31 ^a (Wolter 1994)	⊕⊕⊕ ⊖ very low ⁷	
BMI change Follow-up: 15 days	The mean BMI change in the hospital care group was 0.7	The mean BMI change in the home care groups was 0.2 lower (0.63 lower to 0.23 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{5,6}	
Change in quality of life - CF-QOL-	The mean change in CF-QOL-physical	The mean change in CF-QOL-physical in		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very	

Comparison 1.1. Home care compared to hospital care for the administration of IV antibiotics for people with CF experiencing an acute pulmonary exacerbation						
Physical. Scale from: 0 to 100. Follow-up: 15 days	in the hospital care group was 12.2	the home care groups was 2.2 lower (13.21 lower to 8.81 higher)			low ^{3,5 f}	
Change in quality of life - CF-QOL-Social. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-social in the hospital care group was 6.4	The mean change in CF-QOL-social in the home care groups was 3.4 lower (18.87 lower to 12.07 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	
Change in quality of life - CF-QOL-Treatment. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-treatment in the hospital care group was 8.2	The mean change in CF-QOL-treatment in the home care groups was 2 lower (17.15 lower to 13.15 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	
Change in quality of life - CF-QOL-Symptoms. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-symptoms in the hospital care group was 23.3	The mean change in CF-QOL-symptoms in the home care groups was 17.1 lower (31.25 to 2.95 lower)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{4,5, f}	
Change in quality of life - CF-QOL-Emotional. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-emotional in the hospital care group was 8.3	The mean change in CF-QOL-emotional in the home care groups was 4.2 higher (8.67 lower to 17.07 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	
Change in quality of life - CF-QOL-Future. Scale from: 0 to 100. Follow-up: 15 days	The mean change in the hospital care group was 9.3	The mean change in CF-QOL-future in the home care groups was 5.5 lower (17.96 lower to 6.96 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	
Change in quality of life - CF-QOL-Relationships. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-relationships in the hospital care group was -0.5	The mean change in CF-QOL-relationships in the home care groups was 7.4 higher (5.6 lower to 20.4 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	
Change in quality of life -	The mean change in CF-	The mean change in CF-		30 ^d (Esmond	⊕⊕⊕ ⊖	

Comparison 1.1. Home care compared to hospital care for the administration of IV antibiotics for people with CF experiencing an acute pulmonary exacerbation

CF-QOL-Body image. Scale from: 0 to 100. Follow-up: 15 days	QOL-body image in the hospital care group was 1.8	QOL-body image in the home care groups was 0.9 higher (13.92 lower to 15.72 higher)		2006)	very low ^{3,5, f}	
Change in quality of life - CF-QOL-Career. Scale from: 0 to 100. Follow-up: 15 days	The mean change in CF-QOL-career in the hospital care group was 1.7	The mean change in CF-QOL-career in the home care groups was 8.3 higher (5.76 lower to 22.36 higher)		30 ^d (Esmond 2006)	⊕⊕⊕ ⊖ very low ^{3,5, f}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; CF-QOL: cystic fibrosis quality of life questionnaire; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference; RR: risk ratio

- 1 1 Cross-over trial
- 2 2 The quality of the evidence was downgraded by 1 as this is an open-label study
- 3 3 The quality of the evidence was downgraded by 2 because the 95% CI crosses 2 clinical MIDs.
- 4 4 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 clinical MID
- 5 5 The quality of the evidence was downgraded by 1 as there is a high-risk of bias in relation to the comparability of the groups
- 6 6 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 default MID
- 7 7 The quality of the evidence was downgraded by 2 because the 95% CI crosses 2 default MIDs
- 8 a Number of people in each group not reported
- 9 b Number of people included in the analysis in each group unclear
- 10 c The mean difference was calculated by the NGA technical team after calculating mean change and related SD in each group using the mean and SE at baseline and follow-up and assuming a correlation of 0.75
- 11 d There were 15 people in each group, but the total N of people is 28. Two people had both home care and hospital care.
- 12 e There were 19 people in the home group, 21 people in the hospital group (40 in total)
- 13 f Imprecision for quality of life was assessed using a clinical MID of 5 because the study by Esmond et al. used the CF-QOL questionnaire (Gee et al.2000)

Table 33: Summary clinical evidence profile: comparison 1.2. Home versus hospital care for the administration of IV antibiotics in people with CF and chronic pulmonary infection with *P aeruginosa*

Comparison 1.2 Home versus hospital care for the administration of IV antibiotics for people with CF and chronic pulmonary infection

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Treatments (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Hospital care for the administration of IV antibiotics	Home care for the administration of IV antibiotics				
Lung function: Change in FEV ₁ %	The mean change in the hospital care	The mean change in FEV ₁ in the home care		56 ^a (Riethmueler 2002)	⊕⊕⊕⊕ very low ^{1,2}	

Comparison 1.2 Home versus hospital care for the administration of IV antibiotics for people with CF and chronic pulmonary infection

predicted Scale from: 0 to 100. Follow-up: 14 days	group was 6	groups was 2 higher (9.81 lower to 13.81 higher)			
Nutritional status: change in weight (kg) Follow-up: 14 days	The mean change in the hospital care group was 1.1	The mean change in weight in the home care groups was 0 higher (4.38 lower to 4.38 higher)		57 ^a (Riethmuel ler 2002)	⊕⊕⊕⊕ very low ^{1,3}
Nutritional status: change in weight for height (%) Follow-up: 14 days	The mean change in the hospital care group was 4	The mean change in weight for height (%) in the home care groups was 1 lower (4.64 lower to 2.64 higher)		57 ^a (Riethmuel ler 2002)	⊕⊕⊕⊕ very low ^{1,4}

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference

- 1 The quality of the evidence was downgraded by 1 due to high risk of bias in relation to the comparability of the groups
- 2 The quality of the evidence was downgraded by 2 because the 95% CI crosses 2 clinical MIDs
- 3 The quality of the evidence was downgraded by 2 because the 95% CI crosses 2 default MIDs
- 4 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 default MID
- 5 A Number of people included in the analysis in each group unclear

7 **7.1.4.2 CF centre care**

8 **Table 34: Summary clinical evidence profile: Comparison 2.1. CF centre care versus**
9 **shared care (care by non-specialist in CF)**

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Shared care (UK equivalent)	CF centre care				
Change in FEV ₁ % predicted. Scale from: 0 to 100. Follow-up: 1 year	The mean change in FEV ₁ % predicted in the shared care group was -2.4	The mean change in FEV ₁ % predicted in the CF centre care groups was 0.5 lower (3.05 lower to 2.05 higher)		82 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ¹	
First to last FEV ₁ (% per	The mean first to last FEV ₁ (% per	The mean first to last FEV ₁ (% per year) in the CF		97 (Thomas	⊕⊕⊕⊕ very	

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF					
year) Scale from: 0 to 100. Follow-up: 3 years	year) in the shared care group was 1	centre care groups was 2.4 lower (5.72 lower to 0.92 higher)		2008)	low ^{2,3}
Slope FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean slope FEV ₁ (% per year) in the shared care group was 0.7	The mean slope FEV ₁ (% per year) in the CF centre care groups was 2.2 lower (5.37 lower to 0.97 higher)		97 (Thomas 2008)	⊕⊕⊕⊕ very low ^{2,3}
Change in BMI Follow-up: 1 year	The mean change in BMI in the shared care group was 0.54	The mean BMI in the CF centre care groups was 0.12 lower (0.44 lower to 0.2 higher)		82 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ¹
Quality of life: CFQ-Teen - Physical Scale from: 0 to 100.	The mean CF-QOL teen - physical in the control shared care was 90.4	The mean CF- QOL teen - physical in the CF centre care groups was 17.8 lower (30.28 to 5.32 lower)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,4, a}
Quality of life: CFQ-Teen - Role Scale from: 0 to 100.	The mean CF-QOL teen - role in the shared care group was 86.6	The mean CF- QOL teen - role in the CF centre care groups was 10.4 lower (26.45 lower to 5.65 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,4, a}
Quality of life: CFQ-Teen - Vitality Scale from: 0 to 100.	The mean CF-QOL teen - vitality in the shared care group was 74.2	The mean CF- QOL teen - vitality in the CF centre care groups was 18.2 lower (32.5 to 3.9 lower)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,4, a}
Quality of life: CFQ-Teen - Emotional Scale from: 0 to 100.	The mean CF-QOL teen - emotional in the shared care group was 82.7	The mean CF- QOL teen - emotional in the CF centre care groups was 5.5 lower (18.35 lower to 7.35 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,4, a}
Quality of life: CFQ-Teen - Social Scale from: 0 to 100.	The mean CF-QOL teen - social in the shared care group was 94	The mean CF- QOL teen - social in the CF centre care groups was 17.6 lower (26.71 to 8.49		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,4, a}

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF					
Quality of life: CFQ-Teen - Body Scale from: 0 to 100.	The mean CF-QOL teen - body in the shared care group was 76.7	lower) The mean CF-QOL teen - body in the CF centre care groups was 4.5 lower (21.56 lower to 12.56 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{5, 4, a}
Quality of life: CFQ-Teen - Eating Scale from: 0 to 100.	The mean CF-QOL teen - eating in the shared care group was 76.7	The mean CF-QOL teen - eating in the CF centre care groups was 4.5 lower (21.56 lower to 12.56 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{5, 4, a}
Quality of life: CFQ-Teen - TB Scale from: 0 to 100.	The mean CF-QOL teen - TB in the shared care group was 65.6	The mean CF-QOL teen - TB in the CF centre care groups was 9.6 lower (28.01 lower to 8.81 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{5, 4, a}
Quality of life: CFQ-Teen - Health Scale from: 0 to 100.	The mean CF-QOL teen - health in the shared care group was 72.2	The mean CF-QOL teen - health in the CF centre care groups was 14.8 lower (31.75 lower to 2.15 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,5, a}
Quality of life: CFQ-Teen - Weight Scale from: 0 to 100.	The mean CF-QOL teen - weight in the shared care group was 72.2	The mean CF-QOL teen - weight in the CF centre care groups was 12.5 lower (29.45 lower to 4.45 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3,5, a}
Quality of life: CFQ-Teen - Respiratory Scale from: 0 to 100.	The mean CF-QOL teen - respiratory in the shared care group was 72.8	The mean CF-QOL teen - respiratory in the CF centre care groups was 4.5 lower (15.25 lower to 6.25 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life: CFQ-Teen - Digestion Scale from: 0 to 100.	The mean CF-QOL teen - digestion in the shared care group was 92.2	The mean CF-QOL teen - digestion in the CF centre care groups was 7.9 lower (17.14 lower to 1.34 higher)		34 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life:	The mean CF-QOL child - physical in	The mean CF-QOL child - physical in the		83 (Thomas	⊕⊕⊕⊕ very low ^{5,}

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF						
CFQ-Child - Physical Scale from: 0 to 100.	the shared care group was 77.2	CF centre care groups was 1.2 lower (10.97 lower to 8.57 higher)		2006)	4, a	
Quality of life: CFQ-Child - Emotional Scale from: 0 to 100.	The mean mean CF-QOL child - emotional in the shared care group was 74.8	The mean CF-QOL child - emotional in the CF centre care groups was 1.3 higher (5.13 lower to 7.73 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{4, a}	
Quality of life: CFQ-Child - Social Scale from: 0 to 100.	The mean CF-QOL child - social in the shared care group was 71.9	The mean CF-QOL child - social in the CF centre care groups was 1.7 lower (9.46 lower to 6.06 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}	
Quality of life: CFQ-Child - Body Scale from: 0 to 100.	The mean CF-QOL child - body in the shared care group was 81.1	The mean CF-QOL child - body in the CF centre care groups was 2.8 lower (13.64 lower to 8.04 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}	
Quality of life: CFQ-Child - Eating Scale from: 0 to 100.	The mean score in the shared care group was 76.6	The mean CF-QOL child - eating in the CF centre care groups was 0.5 lower (11.94 lower to 10.94 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{5, 4, a}	
Quality of life: CFQ-Child - TB Scale from: 0 to 100.	The mean CF-QOL child - TB in the shared care group was 63.7	The mean CF-QOL child - TB in the CF centre care groups was 4.7 higher (5.88 lower to 15.28 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}	
Quality of life: CFQ-Child - Respiratory Scale from: 0 to 100.	The mean CF-QOL child - respiratory in the shared care group was 66.9	The mean CF-QOL child - respiratory in the CF centre care groups was 3.9 higher (5.69 lower to 13.49 higher)		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}	
Quality of life: CFQ-Child - Digestion Scale from: 0 to 100.	The mean CF-QOL child - digestion in the shared care group was	The mean CF-QOL child - digestion in the CF centre care groups was 4 higher		83 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}	

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF					
	72.1	(8.38 lower to 16.38 higher)			
Quality of life: CFQ-Parent - Physical Scale from: 0 to 100.	The mean CF-QOL parent - physical in the shared care group was 71.7	The mean CF-QOL parent - physical in the CF centre care groups was 2.5 higher (6.96 lower to 11.96 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life: CFQ-Parent - Vitality Scale from: 0 to 100.	The mean CF-QOL parent - vitality in the shared care group was 65	The mean CF-QOL parent - vitality in the CF centre care groups was 0.7 lower (7.78 lower to 6.38 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{4, a}
Quality of life: CFQ-Parent - Emotional Scale from: 0 to 100.	The mean CF-QOL parent - emotional in the shared care group was 75	The mean CF-QOL parent - emotional in the CF centre care groups was 1.1 higher (7.52 lower to 9.72 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life: CFQ-Parent - Body Scale from: 0 to 100.	The mean CF-QOL parent - body in the shared care group was 69.2	The mean CF-QOL parent - body in the CF centre care groups was 3 higher (9.12 lower to 15.12 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{5, 4, a}
Quality of life: CFQ-Parent - Eating Scale from: 0 to 100.	The mean CF-QOL parent - eating in the shared care group was 70.5	The mean CF-QOL parent - eating in the CF centre care groups was 7.5 lower (20.22 lower to 5.22 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life: CFQ-Parent - TB Scale from: 0 to 100.	The mean CF-QOL parent - TB in the control group was 51.4	The mean CF-QOL parent - TB in the CF centre care groups was 6.2 lower (14.63 lower to 2.23 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}
Quality of life: CFQ-Parent - Health Scale from: 0 to 100.	The mean CF-QOL parent - health in the shared care group was 68.3	The mean CF-QOL parent - health in the CF centre care groups was 1.1 higher (8.6 lower to 10.8 higher)		80 (Thomas 2006)	⊕⊕⊕⊕ very low ^{3, 4, a}

Comparison 2.1. CF centre care compared to shared care (UK equivalent) for people with CF

Quality of life: CFQ-Parent - Weight Scale from: 0 to 100.	The mean CF-QOL parent - weight in the shared care group was 57.1	The mean CF-QOL parent - weight in the CF centre care groups was 0.8 lower (16.4 lower to 14.8 higher)		80 (Thomas 2006)	⊕⊖⊖⊖ very low ⁵ . 4, a
Quality of life: CFQ-Parent - Respiratory Scale from: 0 to 100.	The mean mean CF-QOL parent - respiratory in the shared care group was 74.8	The mean CF-QOL parent - respiratory in the CF centre care groups was 0.5 lower (10.33 lower to 9.33 higher)		80 (Thomas 2006)	⊕⊖⊖⊖ very low ⁵ . 4, a
Quality of life: CFQ-Parent - Digestion Scale from: 0 to 100.	The mean CF-QOL parent - digestion in the shared care group was 77.1	The mean CF-QOL parent - digestion in the CF centre care groups was 0.6 lower (8.76 lower to 7.56 higher)		80 (Thomas 2006)	⊕⊖⊖⊖ very low ³ . 4, a
Quality of life: CFQ-Parent - School function Scale from: 0 to 100.	The mean score in the share care group was 65.1	The mean CF-QOL parent - school function in the CF centre care groups was 0.60 lower (11.63 lower to 10.43 higher)		80 (Thomas 2006)	⊕⊖⊖⊖ very low ⁵ . 4, a

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; CFQ: cystic fibrosis questionnaire; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference

- 1 1 The quality of the evidence was downgraded by 2 because of the differences between groups.
- 2 2 The quality of the evidence was downgraded by 2 due to high risk of bias in relation to the selection of the
- 3 population and high loss to follow-up
- 4 3 The quality of the evidence was downgraded by 1 as the 95% CI crosses 1 clinical MID
- 5 4 The quality of the study was downgraded by 2 due to high risk of bias in relation to comparability of the groups,
- 6 and significant differences at follow-up between groups
- 7 5 The quality of the evidence was downgraded by 2 as the 95% CI crosses 2 clinical MIDs
- 8 a Imprecision for quality of life was assessed using a clinical MID of 8.5 because the paper by Thomas et al. uses
- 9 the CFQ- Teen, CFQ-Child and CFQ-Parent (Quittner et al. 2005)
- 10 * CFQ-Teen: data not available for the following domain: school function
- 11 ** CFQ-Child: data no reported for the following domains: role, vitality, health and weight; data not available for
- 12 school function
- 13 *** CFQ-Parent: data no reported for the following domains: role and social

14 **Table 35: Summary clinical evidence profile: Comparison 2.2. CF centre care versus**
15 **local care (below CF Trust recommendations)**

Comparison 2.2. CF Centre compared to Local care (below CF Trust recs) for people with CF					
Outcomes	Illustrative comparative risks* (95% CI)	Relative effect	No of Participant	Quality of the	Comments

Comparison 2.2. CF Centre compared to Local care (below CF Trust recs) for people with CF						
	Assumed risk	Corresponding risk	(95% CI)	s (studies)	evidence (GRADE)	
	Local care (below CF Trust recs)	CF Centre				
Change in lung function: FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 1 year	The mean change in FEV ₁ % predicted in the local care group was -5.6	The mean change in FEV ₁ % predicted in the CF centre groups was 2.7 higher (0.55 lower to 5.95 higher)		64 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ^{1,2}	
Lung function: First to last FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean first to last FEV ₁ (% per year) in the local care group was 4.3	The mean first to last FEV ₁ (% per year) in the CF centre groups was 5.7 lower (10.99 to 0.41 lower)		78 (Thomas 2008)	⊕⊕⊕⊕ very low ^{2,3}	
Slope FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean slope FEV ₁ (% per year) in the local care group was 1.8	The mean slope FEV ₁ (% per year) in the CF centre groups was 3.3 lower (6.13 to 0.47 lower)		78 (Thomas 2008)	⊕⊕⊕⊕ very low ^{2,3}	
Change in BMI Follow-up: 1 year	The mean BMI change in the local care group was 0.51	The mean BMI change in the CF centre groups was 0.09 lower (0.42 lower to 0.24 higher)		64 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ¹	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference

- 1 The quality of the evidence was downgraded by 2 because of the differences between groups.
- 2 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 due to high risk of bias in relation to the selection of the population and high loss to follow-up

5 **Table 36: Summary clinical evidence profile: Comparison 2.3. CF centre care versus**
6 **general clinic (non-CF)**

Comparison 2.3. CF specialist clinic compared to general (not CF) clinic for people with CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	General	CF specialist				

Comparison 2.3. CF specialist clinic compared to general (not CF) clinic for people with CF						
	(not CF) clinic	clinic				
Patient satisfaction with care overall	The overall patient satisfaction rating in the general clinic group was 4.20	The overall patient satisfaction rating in the CF specialist clinic group was 3.76	0.44 higher (0.29 higher to 0.58 higher)	686 (Walters 1994)	⊕⊕⊕⊕ very low ¹	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; CF: cystic fibrosis; MD: mean difference						

1
2
1 The quality of the evidence was downgraded by 1 because the authors did not control the analysis for any of the confounding factors

3 7.1.4.3 Shared care

4
5
Table 37: Summary clinical evidence profile: Comparison 3.1. Local care (below CF Trust recs) compared to shared care (UK equivalent)

Comparison 3.1. Local care (below CF Trust recs) compared to shared care (UK equivalent) for people with CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Shared care (UK equivalent)	Local care (below CF Trust recs)				
Change in lung function: FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 1 year	The mean change in FEV ₁ % predicted in the shared care group was -2.4	The mean change in FEV ₁ % predicted in the local care groups was 3.2 lower (6.84 lower to 0.44 higher)		64 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ^{1,2}	
First to last FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean first to last FEV ₁ (% per year) in the shared care group was 1	The mean first to last FEV ₁ (% per year) in the local care groups was 3.3 higher (2.59 lower to 9.19 higher)		41 (Thomas 2008)	⊕⊕⊕⊕ very low ^{2,3}	
Slope FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean slope FEV ₁ (% per year) in the shared care group was 0.7	The mean slope FEV ₁ (% per year) in the local care groups was 1.1 higher (2.69 lower to 4.89 higher)		41 (Thomas 2008)	⊕⊕⊕⊕ very low ³	

Comparison 3.1. Local care (below CF Trust recs) compared to shared care (UK equivalent) for people with CF

Change in BMI Follow-up: 1 year	The mean BMI change in the shared care group was 0.54	The mean BMI change in the local care groups was 0.03 lower (0.43 lower to 0.37 higher)		64 (Van Koolwijk 2002)	⊕⊕⊕⊕ very low ¹	
---------------------------------------	---	---	--	------------------------	-------------------------------	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval;

- 1 *The quality of the evidence was downgraded by 2 because of the differences between groups.*
 2 *The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 clinical MID*
 3 *The quality of the evidence was downgraded by 2 due to high risk of bias in relation to the selection of the population and high loss to follow-up*

Table 38: Summary clinical evidence profile: Comparison 3.2. Shared care (above UK equivalent) compared to shared care (UK equivalent) for people with CF

Comparison 3.2. Shared care (above UK equivalent) compared to shared care (UK equivalent) for people with CF

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Shared care (UK equivalent)	Shared care (above UK equivalent)				
Lung function: First to last FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean in first to last (% per year) FEV ₁ % in the shared care group was 1	The mean change first to last FEV ₁ (% per year) in the shared care + groups was 0.5 lower (5.63 lower to 4.63 higher)		49 (Thomas 2008)	⊕⊕⊕⊕ very low ^{1,2,3}	
Lung function: Slope FEV ₁ (% per year) Scale from: 0 to 100. Follow-up: 3 years	The mean slope FEV ₁ (% per year) in the shared care group was 0.7	The mean slope FEV ₁ (% per year) in the shared care + groups was 2.1 lower (6.52 lower to 2.32 higher)		49 (Thomas 2008)	⊕⊕⊕⊕ very low ^{1,2,3}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; MD: mean difference

- 7 *The quality of the evidence was downgraded by 2 due to high risk of bias in relation to the selection of the population and high loss to follow-up*
 8
 9 *The quality of the evidence was downgraded by 1 because 1 of the comparators is not representative of current UK practice*
 10
 11 *The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 clinical MID*

1 **7.1.4.4 Telemedicine**

2 **Table 39: Summary clinical evidence profile: Comparison 4.1. Home monitoring**
3 **programme + diary recording versus usual care**

Comparison 4.1. Home monitoring program with diary and usual care compared to usual care for people with CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual care	Home monitoring program with diary and usual care				
Change in FEV ₁ (% predicted). Scale from: 0 to 100. Follow-up: 4 years	The mean FEV ₁ % predicted in the usual care group was 11.5	The mean FEV ₁ (% predicted) in the home monitoring groups was 8 lower (17.01 lower to 1.01 higher)		50 (Finkelstein 1992)	⊕⊕⊕⊕ very low ^{1,2}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; MD: mean difference

4 *1 The quality of the evidence was downgraded by 1 due to unclear comparability between groups*
5 *2 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 clinical MID*

6 **Table 40: Summary clinical evidence profile: Comparison 4.2. Telemedicine versus**
7 **usual care**

Comparison 4.2. Telemedicine compared to standard care for people with CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Standard care	Telemedicine				
Change in quality of life – CF-QOL. Scale from: 0 to 100. Follow-up: 6 months	Not reported	The mean in CQ-QOL body in the intervention groups was not reported. There was a significant improvement at 6 months, p=0.02		7 (Wilkinson 2008)	⊕⊕⊕⊕ very low ¹	The study does not report on differences between groups.

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; CFQOL: cystic fibrosis quality of life questionnaire

8 *1 The quality of the evidence was downgraded by 2 because of incomplete reporting and high-loss to follow-up*

1 7.1.5 Economic evidence

2 Three economic evaluations were identified that compared home-care IV antibiotic therapy to
3 hospital IV antibiotic therapy. Wolter 1998 conducted a cost–consequence analysis in
4 Australia based on a RCT that included 17 adult patients, whilst Thornton 2005 and Elliott
5 2005 performed a cost-effectiveness analysis and cost-benefit analysis, respectively, using
6 the same data collected retrospectively from 116 adults with cystic fibrosis at The
7 Manchester Adult CF Unit. The methods and results of these studies are described in more
8 detail in Appendix K.

9 Full details of the search and economic article selection flow chart can be found in Appendix
10 E and F, respectively. Data extraction tables and quality assessments of included studies
11 can be found in Appendix L and M, respectively.

12 This review question was not prioritised for de novo economic modelling. To aid
13 consideration of cost-effectiveness a costing tool that utilised a “what-if” approach was
14 developed to estimate the annual cost per patient per model of care for a given MDT
15 composition. The methods and results of this analysis are reported in Appendix K. The
16 results are reproduced in Table 41 for ease of reference, where the cheapest model of care
17 is the Specialist Centre (£9,191) followed by Outreach Care (£10,004) and Shared Care
18 (n=150, £14,347; n=250, £13,164).

19 **Table 41: Annual cost across the recognised models of care ^a**

Model of care	Clinic size ^b	Annual travel costs incurred by the Specialist Centre MDT	Annual MDT staff costs	Total annual cost/ clinic	Total annual cost/ person
Specialist Centre	250	None	£2,297,684 (Specialist Centre MDT)	£2,297,684	£9,191
Shared Care	250	£11,760 (25 days of travel totalling 3,000 miles @ 56p/ mile for 7 HCPs)	£1,384,445 (paediatric Shared Care MDT for 150 people) + £1,881,229 (Specialist Centre for 250 people ^c) + £13,567 (0.5 administrator)	£3,291,001	£13,164
Shared Care	150	£6,115 (13 days of travel totalling 1,560 miles @ 56p/ mile for 7 HCPs)	£747,951 (paediatric Shared Care MDT for 75 people) + £1,384,445 (Specialist Centre for 150 people ^c) + £13,567 (0.5 administrator)	£2,152,078	£14,347
Outreach Care	1	£627 (1 day of travel totalling 160 miles @ 56p/ miles for 7 HCPs) shared by 2 people	£9,191 (Specialist Centre MDT divided by a clinic size of 250) + £60 (administrator, 0.5 days) + £440 (capital costs, 0.5 days)	NC	£10,004

20 *HCP, health care professional; NC, not calculable; MDT, multidisciplinary team*

21 (a) *Excluding diagnosis and treatment costs*

22 (b) *Number of people with cystic fibrosis managed by the model of care*

23 (c) *Using paediatric WTE figures, see Appendix K*

1 **7.1.6 Evidence statements**

2 **7.1.6.1 Home-based care**

3 **37.1.6.1.1 Comparison 1.1. Home versus hospital care for the administration of IV antibiotics in**
4 **people with cystic fibrosis experiencing an acute pulmonary exacerbation**

5 **Lung function: FEV₁**

6 Very low quality evidence from 1 RCT demonstrated no clinically significant difference in lung
7 function (measured as change in FEV₁ % predicted) following 31 admissions of 17 adults
8 with cystic fibrosis who received IV antibiotic treatment at home or within hospital for the
9 treatment of an acute pulmonary exacerbation at 21 days follow-up.

10 Very low quality data from 1 prospective cohort study of 63 IV antibiotic treatments of people
11 with cystic fibrosis aged ≥ 12 years found no clinically significant difference in lung function
12 (measured as change in FEV₁% predicted) at 18 days following IV antibiotic treatment at
13 home and in hospital for the treatment of acute pulmonary exacerbation. The number of
14 people included in the analysis was not reported.

15 Very low quality evidence from 1 prospective cohort study of 30 IV antibiotic treatments in 28
16 adults with cystic fibrosis showed that there was no clinically significant difference in lung
17 function (measured as change in FEV₁ % predicted) at 15 days following home IV antibiotic
18 treatment (with self-performed chest physiotherapy) compared with hospital administered IV
19 antibiotic treatment with chest physiotherapy performed by experienced respiratory
20 physiotherapists.

21 **Mortality**

22 No evidence was found for this critical outcome.

23 **Patient and carer satisfaction**

24 No evidence was found for this critical outcome.

25 **Lung function: LCI**

26 No evidence was found for this important outcome.

27 **Time to next pulmonary exacerbation**

28 Very low quality evidence from 1 prospective cohort study of 59 IV antibiotic treatments in 40
29 children, young people and adults with cystic fibrosis aged ≥ 8 years receiving IV antibiotics
30 for acute pulmonary exacerbations identified that more patients treated in hospital did not
31 require a further course of antibiotics at 12 weeks compared to those receiving antibiotics at
32 home and this represents a clinically significant benefit.

33 **Nutritional status**

34 Very low quality evidence from 1 prospective cohort study of 74 antibiotic treatments in
35 people with cystic fibrosis aged ≥12 years found no clinically significant difference in weight
36 (measured as change in kg) following IV antibiotic treatment at home and in hospital for the
37 treatment of acute pulmonary exacerbation at 18 days follow-up. The number of people
38 included in the analysis was not reported.

39 Very low quality evidence from 1 RCT of 31 admissions with 17 adults with cystic fibrosis
40 found that at 10 days following treatment, there was no clinically significant difference in

1 weight change between those who received IV antibiotics at home compared to those
2 receiving IV antibiotics in hospital

3 Very low quality evidence from 1 prospective cohort study of 30 IV antibiotic courses for the
4 treatment of an acute respiratory exacerbation in 28 adults with cystic fibrosis showed no
5 clinically significant difference in BMI between the participants receiving therapy at home or
6 in the hospital at 15 days follow-up.

7 **Quality of life**

8 Very low quality evidence from 1 prospective cohort study of 30 IV antibiotic courses for the
9 treatment of acute respiratory exacerbations in 28 adults with cystic fibrosis showed no
10 clinically significant difference in relation to the quality of life (measured as change in the CF-
11 QOL domains – physical, social, treatment, emotional, future, relationships, body image and
12 career) between the participants receiving therapy at home or at hospital at 15 days follow-
13 up. The same evidence found a clinically significant lower score in quality of life (measured
14 as change in the domain CF-QOL – symptoms) in the group of participants receiving therapy
15 at home compared to those receiving therapy in the hospital at 15 days follow-up.

16 **Frequency of cross-infection**

17 No evidence was found for this important outcome.

18 **Staff experience**

19 No evidence was found for this important outcome.

20 **Adherence to treatment**

21 No evidence was found for this important outcome.

22 **227.1.6.1.2 Comparison 1.2: home versus hospital care for the administration of IV antibiotics in 23 people with cystic fibrosis and chronic pulmonary infection with *P aeruginosa***

24 **Lung function: FEV₁**

25 Very low quality evidence from 1 prospective cohort study of 56 courses of IV antibiotics for
26 the treatment of chronic pulmonary infection with *P aeruginosa* in children, young people and
27 adults with cystic fibrosis showed no clinically significant difference in lung function
28 (measured as change in FEV₁ % predicted) between receiving therapy at home or at hospital
29 at 14 days follow-up. The number of people included in the analysis was not reported.

30 **Mortality**

31 No evidence was found for this critical outcome.

32 **Patient and carer satisfaction**

33 No evidence was found for this critical outcome.

34 **Lung function: LCI**

35 No evidence was found for this important outcome.

36 **Time to next pulmonary exacerbation**

37 No evidence was found for this important outcome.

1 **Nutritional status**

2 Very low quality evidence from 1 prospective cohort study of 57 courses of IV antibiotics in
3 children, young people and adults with cystic fibrosis for the treatment of chronic pulmonary
4 infection with *P aeruginosa* showed no clinically significant difference in nutritional status
5 (measured as change in kg or as weight/ height %) between receiving therapy home or at
6 hospital at 14 days follow-up. The number of people included in the analysis was not
7 reported.

8 **Quality of life**

9 No evidence was found for this important outcome.

10 **Frequency of cross-infection**

11 No evidence was found for this important outcome.

12 **Staff experience**

13 No evidence was found for this important outcome.

14 **Adherence to treatment**

15 No evidence was found for this important outcome.

16 **7.1.6.2 Cystic fibrosis centre care**

17**7.1.6.2.1 Comparison 2.1. Cystic fibrosis centre care versus shared care**

18 **Lung function: FEV₁**

19 Very low quality evidence from 1 prospective cohort study with 82 children and young people
20 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in lung
21 function (measured as change in FEV% predicted) between the group of participants
22 attending a cystic fibrosis centre and the participants receiving shared care at 1 year follow-
23 up.

24 Very low quality evidence from 1 retrospective cohort study with 97 people with cystic fibrosis
25 aged 0 to 20 years showed no clinically significant differences in lung function (measured as
26 change from first to last FEV₁ % predicted per year and slope FEV₁% per year) between the
27 group of participants attending a cystic fibrosis centre and the participants receiving shared
28 care during the 3 years follow-up.

29 **Mortality**

30 No evidence was found for this critical outcome.

31 **Patient and carer satisfaction**

32 No evidence was found for this critical outcome.

33 **Lung function: LCI**

34 No evidence was found for this important outcome.

1 **Time to next pulmonary exacerbation**

2 No evidence was found for this important outcome.

3 **Nutritional status: BMI**

4 Very low quality evidence from 1 prospective cohort study with 82 children and young people
5 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in the
6 change in BMI between the group of participants attending a cystic fibrosis centre and the
7 participants receiving shared care at 1 year follow-up.

8 **Quality of life**

9 Very low quality evidence from 1 cross-sectional study with 34 young people with cystic
10 fibrosis showed a clinically significant lower score in the quality of life for the domains
11 physical, vitality and social(measured with the CFQ-R Teen questionnaire) in the group of
12 participants attending a cystic fibrosis centre compared to the group of participants receiving
13 shared care. However, no clinically significant differences were found in the rest of the
14 domains (role, emotional, body, eating, treatment burden, health, weight, respiratory and
15 digestion).

16 Very low quality evidence from 1 cross-sectional study with 83 children with cystic fibrosis
17 showed no clinically significant differences in the reported quality of life between the group of
18 participants attending a cystic fibrosis centre and the group of participants receiving shared
19 care (measured with the following CFQ-R child domains: physical, emotional, social, body,
20 eating, treatment burden, respiratory and digestion).

21 Very low quality evidence from 1 cross-sectional study with 80 parents of people with cystic
22 fibrosis showed no clinically significant differences in the parent-reported quality of life of
23 their children between the group of participants attending a cystic fibrosis centre and the
24 group of participants receiving shared care (measured with the following CFQ-R parents
25 domains: physical, vitality, emotional, body, eating, treatment burden, health, weight,
26 respiratory, digestion and school function).

27 **Frequency of cross-infection**

28 No evidence was found for this important outcome.

29 **Staff experience**

30 No evidence was found for this important outcome.

31 **Adherence to treatment**

32 No evidence was found for this important outcome.

337.1.6.2.2 **Comparison 2.2. Cystic fibrosis centre care versus local care (below CF Trust
34 recommendations)**

35 **Lung function: FEV₁**

36 Very low quality evidence from 1 prospective cohort study with 63 children and young people
37 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in lung
38 function (measured as change in FEV₁ % predicted) between those receiving cystic fibrosis
39 centre care and participants receiving local care (only annual check-up at the cystic fibrosis
40 centre) at 1 year follow-up.

1 Very low quality evidence from 1 retrospective cohort study with 78 children and young
2 people with cystic fibrosis aged 0 to 20 years showed a clinically significant lower lung
3 function (measured as change from first to last FEV₁% predicted per year) in the group of
4 participants receiving cystic fibrosis centre care compared to those receiving local care (care
5 by non-specialists in cystic fibrosis, with involvement by the cystic fibrosis centre once a year
6 for some people) during the 3 years follow-up. However, the same evidence found no
7 clinically significant difference in lung function (measured as slope FEV₁% predicted per
8 year) between those receiving cystic fibrosis centre care and participants receiving local care
9 during the 3 years follow-up.

10 **Mortality**

11 No evidence was found for this critical outcome.

12 **Patient and carer satisfaction**

13 No evidence was found for this critical outcome.

14 **Lung function: LCI**

15 No evidence was found for this important outcome.

16 **Time to next pulmonary exacerbation**

17 No evidence was found for this important outcome.

18 **Nutritional status: BMI**

19 Very low quality evidence from 1 prospective cohort study with 64 children and young people
20 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in the
21 change in BMI between those receiving cystic fibrosis centre care and the participants
22 receiving local care (only annual check-up at the cystic fibrosis centre) at 1 year follow-up.

23 **Quality of life**

24 No evidence was found for this important outcome.

25 **Frequency of cross-infection**

26 No evidence was found for this important outcome.

27 **Staff experience**

28 No evidence was found for this important outcome.

29 **Adherence to treatment**

30 No evidence was found for this important outcome.

317.1.6.2.3 **Comparison 2.3. Cystic fibrosis centre care versus general clinic (non-cystic fibrosis)**

32 **Lung function: FEV₁**

33 No evidence was found for this critical outcome.

1 **Mortality**

2 No evidence was found for this critical outcome.

3 **Patient and carer satisfaction**

4 Very low quality evidence from 1 cross-sectional study with 686 people with cystic fibrosis
5 aged ≥ 15 years showed a significant difference in overall patient satisfaction between the
6 participants receiving cystic fibrosis centre care and the participants receiving usual care
7 (care by non-specialists in cystic fibrosis). The clinical significance and the imprecision for
8 this outcome could not be calculated.

9 **Lung function: LCI**

10 No evidence was found for this important outcome.

11 **Time to next pulmonary exacerbation**

12 No evidence was found for this important outcome.

13 **Nutritional status: BMI**

14 No evidence was found for this important outcome.

15 **Quality of life**

16 No evidence was found for this important outcome.

17 **Frequency of cross-infection**

18 No evidence was found for this important outcome.

19 **Staff experience**

20 No evidence was found for this important outcome.

21 **Adherence to treatment**

22 No evidence was found for this important outcome.

23 **7.1.6.3 Shared care**

24 **247.1.6.3.1 Comparison 3.1. Local care (below CF Trust recommendations) versus shared care**
25 **(UK equivalent)**

26 **Lung function: FEV₁**

27 Very low quality evidence from 1 prospective cohort study with 64 children and young people
28 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in lung
29 function (measured as change in FEV₁ % predicted) between those receiving local care
30 (below the standards recommended by the CF Trust) and the participants receiving shared
31 care (as in current UK practice) at 1 year follow-up.

32 Very low quality evidence from 1 retrospective cohort study with 41 people with cystic fibrosis
33 aged 0 to 20 years showed no clinically significant difference in lung function (measured as
34 change from first to last FEV₁ % predicted per year and slope FEV₁% per year) between

1 those receiving local care (below the standards recommended by the CF Trust) and the
2 participants receiving shared care (as in current UK practice) during the 3 years follow-up.

3 **Mortality**

4 No evidence was found for this critical outcome.

5 **Patient and carer satisfaction**

6 No evidence was found for this critical outcome.

7 **Lung function: LCI**

8 No evidence was found for this important outcome.

9 **Time to next pulmonary exacerbation**

10 No evidence was found for this important outcome.

11 **Nutritional status: BMI**

12 Very low quality evidence from 1 prospective cohort study with 64 children and young people
13 with cystic fibrosis aged 5 to 17 years showed no clinically significant difference in the
14 change in BMI between the participants receiving local care (below the standards
15 recommended by the CF Trust) and the participants receiving shared care (as in current UK
16 practice) at 1 year follow-up.

17 **Quality of life**

18 No evidence was found for this important outcome.

19 **Frequency of cross-infection**

20 No evidence was found for this important outcome.

21 **Staff experience**

22 No evidence was found for this important outcome.

23 **Adherence to treatment**

24 No evidence was found for this important outcome.

257.1.6.3.2 **Comparison 3.2. Shared care (above UK equivalent) versus shared care (UK
26 equivalent)**

27 **Lung function: FEV₁**

28 Very low quality evidence from 1 retrospective cohort study with 49 people with cystic fibrosis
29 aged 0 to 20 years showed no clinically significant difference in lung function (measured as
30 change from first to last FEV₁% predicted per year and slope FEV₁% per year) between the
31 participants receiving intensive shared care (above UK equivalent) and usual shared care
32 (UK equivalent) during the 3 years follow-up.

33 **Mortality**

34 No evidence was found for this critical outcome.

- 1 **Patient and carer satisfaction**
- 2 No evidence was found for this critical outcome.
- 3 **Lung function: LCI**
- 4 No evidence was found for this important outcome.
- 5 **Time to next pulmonary exacerbation**
- 6 No evidence was found for this important outcome.
- 7 **Nutritional status**
- 8 No evidence was found for this important outcome.
- 9 **Quality of life**
- 10 No evidence was found for this important outcome.
- 11 **Frequency of cross-infection**
- 12 No evidence was found for this important outcome.
- 13 **Staff experience**
- 14 No evidence was found for this important outcome.
- 15 **Adherence to treatment**
- 16 No evidence was found for this important outcome.
- 17 **7.1.6.4 Telemedicine**
- 187.1.6.4.1 ***Comparison 4.1. Home monitoring programme + diary recording versus usual care***
- 19 **Lung function: FEV₁**
- 20 Very low quality evidence from 1 observational study with 50 people with cystic fibrosis > 6
- 21 years old showed no significant difference in lung function (measured as change in FEV₁ %
- 22 predicted) between the participants in the home monitoring programme and the participants
- 23 receiving usual care at 4 years follow-up. The uncertainty for this outcome could not be
- 24 calculated.
- 25 **Mortality**
- 26 No evidence was found for this critical outcome.
- 27 **Patient and carer satisfaction**
- 28 No evidence was found for this critical outcome.
- 29 **Lung function: LCI**
- 30 No evidence was found for this important outcome.

1 **Time to next pulmonary exacerbation**
2 No evidence was found for this important outcome.

3 **Nutritional status**
4 No evidence was found for this important outcome.

5 **Quality of life**
6 No evidence was found for this important outcome.

7 **Frequency of cross-infection**
8 No evidence was found for this important outcome.

9 **Staff experience**
10 No evidence was found for this important outcome.

11 **Adherence to treatment**
12 No evidence was found for this important outcome.

137.1.6.4.2 **Comparison 4.2. Telemedicine versus usual care**

14 **Lung function: FEV₁**
15 No evidence was found for this critical outcome.

16 **Mortality**
17 No evidence was found for this critical outcome.

18 **Patient and carer satisfaction**
19 No evidence was found for this critical outcome.

20 **Lung function: LCI**
21 No evidence was found for this important outcome.

22 **Time to next pulmonary exacerbation**
23 No evidence was found for this important outcome.

24 **Nutritional status**
25 No evidence was found for this important outcome.

26 **Quality of life**
27 Very low quality evidence from 1 RCT with 7 adults with cystic fibrosis on a transplantation
28 list showed that the participants in the telemedicine group experienced a significant
29 improvement in the quality of life, measured as the subjects' perception of body image (with
30 CF-QOL body domain), at 6 months follow-up. The clinical significance and imprecision for
31 this outcome could not be calculated.

1 **Frequency of cross-infection**

2 No evidence was found for this important outcome.

3 **Staff experience**

4 No evidence was found for this important outcome.

5 **Adherence to treatment**

6 No evidence was found for this important outcome.

7 **7.1.6.5 Economic evidence statements**

8 One cost-consequence analysis (Wolter 1997) on people with cystic fibrosis in Australia
9 found that managing exacerbations at home was less expensive compared to those
10 managed in hospital over 5 years. This analysis is partially applicable as the type of
11 economic evaluation applied is difficult to assess cost-effectiveness and a societal
12 perspective is inferred. This evidence is associated with serious limitations from the small
13 number of participants (n=17).

14 One cost-benefit analysis (Elliot 2005) on people with cystic fibrosis in the UK found that
15 hospital based care was more expensive than home based care over 1 year. This analysis is
16 partially applicable with minor limitations.

17 One cost-effectiveness analysis (Thornton 2005) on people with cystic fibrosis in the UK
18 found that the ICER (cost per decline in FEV₁% ≤0%) was £46,098 (the cost to obtain 1
19 more year of effective treatment with hospital care for 1 person) for hospital IV therapy
20 compared to home-care IV therapy. This analysis has minor limitations and is directly
21 applicable given that the type of economic evaluation is unlikely to change the conclusions
22 about cost-effectiveness and all other applicability criteria are met.

23 **7.1.7 Evidence to recommendations**

24 **7.1.7.1 Relative value placed on the outcomes considered**

25 The aim of this review was to assess the effectiveness of different models of care (for
26 example specialist centre, shared care, community care, telehealth and home care) for the
27 care of people with cystic fibrosis.

28 The Committee identified the following outcomes as critical: lung function (FEV₁% predicted),
29 mortality and patient satisfaction. LCI, time to next pulmonary exacerbation, nutritional status,
30 quality of life, carer satisfaction, frequency of cross-infections (*P aeruginosa*, *B cepacia*), staff
31 experience and adherence to treatment were considered to be important outcomes.

32 **7.1.7.2 Consideration of clinical benefits and harms**

33 A number of comparisons between different models of care were covered in this review.
34 Overall, no clinically significant differences in outcomes were found to favour a particular
35 model.

36 In relation to the administration of IV antibiotics for an acute pulmonary disease
37 exacerbation, all evidence was of very low quality; 3 studies found no significant difference in
38 lung function between home and hospital care; there was no evidence on the other 2 critical
39 outcomes of mortality and patient satisfaction. With regards to the important outcomes, the
40 evidence showed a clinically significant reduction in the risk of experiencing an exacerbation
41 at 12 weeks in the group treated in hospital. Very low quality evidence from 3 studies found
42 no clinically significant differences in nutritional status between home and hospital care. Very

1 low quality evidence from 1 study found no clinically significant difference in 8 domains of
2 quality of life but found a clinically significant lower score in the symptoms domain of quality
3 of life in the home care group compared to the hospital group. There was no evidence on
4 LCI, carer satisfaction, frequency of cross-infections, staff experience and adherence to
5 treatment. Likewise, in relation to the comparison between home and hospital administration
6 of IV antibiotics for chronic pulmonary infection, very low quality evidence from 1 study found
7 no clinically significant differences in lung function and nutritional status. There was no
8 evidence on the other outcomes. Following this, the Committee concluded that a
9 recommendation in favour of home IV therapy could only be supported if training and
10 equipment were available to people with cystic fibrosis who are willing and competent to self-
11 administer their treatment.

12 With regard to the comparison between cystic fibrosis management based on specialist
13 centre care and the use of shared care, all evidence was of very low quality; there was
14 evidence from 2 studies on the critical outcome of lung function, and evidence on nutritional
15 status and quality of life from 1 study. No clinically significant differences were found
16 between these 2 models of care except for a clinically significant difference in favour of
17 shared care for the physical, vitality and social domains of quality of life measured with the
18 CFQ-Teen questionnaire. However, the Committee noted that this study was conducted in
19 Australia, where people often need to travel long distances for review and management, and
20 this does not reflect the situation for most in the UK. There was therefore an issue of
21 indirectness with regard to this evidence, and it could not be readily applied to a typical UK
22 arrangement. There was no evidence on the other outcomes. With regards to the
23 comparison between cystic fibrosis specialist centre care and local care (below CF Trust
24 recommendations), very low quality evidence from 2 studies showed mixed results in relation
25 to lung function and very low quality evidence showed no clinically significant difference in
26 nutritional status between these 2 models of care. There was no evidence on the other
27 outcomes. With regards to the comparison between cystic fibrosis specialist centre care and
28 care in a general non-cystic fibrosis clinic, there was very low quality evidence from 1 study
29 on patient satisfaction which showed a significant difference in favour of centre care, but
30 there was no evidence on the other outcomes.

31 With regards to the comparison between local care below CF Trust recommendations and
32 shared care (UK equivalent), there was very low quality evidence from 2 studies which
33 showed no clinically significant difference in lung function between these two models of care,
34 and very low quality evidence from 1 study which showed no clinically significant difference
35 in nutritional status. With regards to the comparison between shared care and intensive
36 shared care, very low quality evidence from 1 study found no clinically significant difference
37 in lung function. There was no evidence on the other outcomes.

38 The Committee noted that there were 2 published studies (Doull 2012, Mahadeva 1998)
39 conducted in the UK that compared cystic fibrosis specialist care with other alternative
40 models of care. These studies had been identified in the evidence search, but were not
41 included in the review because they used a cross-sectional design, and therefore did not
42 meet the protocol criteria. However, the Committee considered that they were relevant, as
43 they reflected UK practice. In 1 of these studies (Doull 2012), the children who received
44 either full cystic fibrosis centre care or hybrid care (cystic fibrosis centre care + local clinic)
45 had better lung function than those who only received local hospital based care. The second
46 of these studies (Mahadeva 1998) found that adults attending a specialist cystic fibrosis
47 centre who had received treatment in a paediatric cystic fibrosis specialist centre (group A)
48 had better lung function and better nutritional status than those who received their paediatric
49 cystic fibrosis care at a non-specialist centre followed by an adult cystic fibrosis specialist
50 centre (group B).

51 In relation to telemedicine, very low quality evidence from 1 study on daily home monitoring
52 (daily recording of symptoms and physical measurements sent to the centre weekly), did not
53 show a clinically significant beneficial effect on lung function compared to usual care. No

1 evidence was available for the other outcomes. However, based on very low quality evidence
2 home telemedicine (weekly videoconference with the patient and members of the MDT to
3 provide psychological support) appeared to be beneficial in improving the quality of life of
4 people with cystic fibrosis that were terminally ill, compared to standard care alone. The
5 Committee noted that this benefit applied only to that specific population of people with cystic
6 fibrosis. However, based on their clinical experience and expertise, the Committee members
7 agreed that people with cystic fibrosis who are severely ill are likely to benefit from face-to-
8 face appointments for appropriate diagnosis and management. No evidence was available
9 on the other outcomes.

10 **7.1.7.3 Consideration of economic benefits and harms**

11 The Committee agreed that providing cystic fibrosis care outside of the specialist centre
12 incurs greater costs from travel and administration, and for shared care an additional MDT.
13 However, the Committee felt it was unrealistic and unethical to recommend the specialist
14 centre alone as people with cystic fibrosis may be unable to travel greater distances than
15 they would under a shared care, or outreach care model.

16 The Committee iterated that the burden from travel could negatively impact a patient's usual
17 activities and in some cases, increase their anxiety. Furthermore, long distances may reduce
18 adherence to hospital for treatment, potentially resulting in later downstream costs from
19 delayed diagnosis and management. The Committee added that this would include
20 conscientious people with cystic fibrosis as factors beyond their control may affect their
21 ability to travel when they are willing to do so. For these reasons, the Committee agreed that
22 a cost-effective recommendation would say how geography and the MDT indicate that care
23 could be delivered through a shared care network or outreach care clinic for paediatrics and
24 adults, respectively.

25 To reduce the burden of visits to a clinic, the Committee considered a role for telemedicine
26 as a supplementary model of care for routine monitoring in people with cystic fibrosis who
27 are clinically stable. The Committee also noted that this model could prevent cross-infection
28 at the clinic that can be costly to prevent and treat. However, people with cystic fibrosis who
29 are unwell must not utilise telemedicine as they would require additional face-to-face
30 appointments to obtain the correct diagnosis and management. Telemedicine would lead to
31 a change in clinical practice as it is currently performed in a small proportion of clinics.
32 According to the Committee, this would incur a small injection of resources to set up if people
33 with cystic fibrosis did not possess the necessary testing equipment such as spirometry.
34 However, the cost of equipment would be negligible (approximately £50) and would last
35 several years before it needed to be replaced (approximately 5 years).

36 The Committee advised that people with cystic fibrosis should be able to telephone for urgent
37 specialist advice as this could lead to more timely management. The Committee added that
38 such a recommendation would not lead to a change in current practice, but would increase
39 the awareness of this service to people with cystic fibrosis and their parents or carers.

40 The Committee agreed that home-care monitoring was an unsustainable model for routine
41 monitoring when they considered the opportunity cost of staff time travelling long distances.
42 Conversely, IV antibiotic therapy can be administered at home without health care
43 professional supervision, provided that training is given during routine clinic visits. The
44 Committee noted that the potential cost savings from this type of administration can only be
45 realised if people with cystic fibrosis are willing and competent to do so, as further costs will
46 be incurred to treat unsuccessful cases. Therefore, to ensure self-administered IV therapy is
47 cost-effective, the Committee made a recommendation to ensure that arrangements are in
48 place for people to have IV antibiotic therapy at home, by transferring the necessary
49 equipment and expert support provided in a clinic setting, to a home setting. Given that the
50 equipment and expert support from a healthcare professional is part of an attendance to the
51 clinic, no additional resources are anticipated from this move. The Committee also added

1 that if the IV training provided during routine clinic visits is successful, the frequency and
2 duration of expert support anticipated for this model, would be relatively low compared to the
3 level of contact provided during an attendance at the clinic.

4 The Committee discussed and agreed the review frequencies for people with cystic fibrosis
5 based on their knowledge and experience. After a diagnosis of cystic fibrosis it is important to
6 review frequently, given the immediate polypharmacy, to prevent the potentially lethal
7 consequences of giving an inappropriate treatment. Following those initial years, the
8 person's treatment schedule and condition is likely to stabilise; then, less frequent monitoring
9 would be considered cost-effective as the potential to miss a negative change in their
10 condition is unlikely to escalate between those reviews. Furthermore, the CF Trust Standards
11 of Care 2011 advises that people with cystic fibrosis are seen at least twice a year by the
12 Specialist Centre. Finally, given that the NHS service specifications for cystic fibrosis and
13 Committee agreed that review frequency should be based on individual circumstances, no
14 significant impact on resource is anticipated from the review frequencies provided by the
15 Committee.

16 **7.1.7.4 Quality of evidence**

17 The quality of the evidence presented in this review was very low as assessed by GRADE.

18 One of the main reasons that led to downgrading the quality of the evidence was the quality
19 of the studies. Most of the included studies used an observational design, and presented
20 problems in the comparability of the groups. Data was poorly reported across studies, and
21 some of them reported the results narratively only.

22 Another reason was the lack of generalisability of the interventions to the UK context. Many
23 of the studies were conducted in countries other than the UK, and the service delivery
24 models evaluated differed from current UK practice.

25 The Committee considered that it was important to note that no evidence was identified for
26 the critical outcomes for many of the comparisons.

27 **7.1.7.5 Other considerations**

28 Given the scarcity and poor quality of the evidence, the recommendations were mainly based
29 on the Committee members' clinical experience, and consensus on good clinical practice
30 (NHS service specifications for cystic fibrosis; CF Trust Standards of Care 2011, which are
31 mentioned in the NHS service specifications as the standards to follow; and European Cystic
32 Fibrosis Society Standards of Care 2014).

33 The Committee agreed that care for people with cystic fibrosis should be provided by a
34 specialist MDT based at a cystic fibrosis specialist centre. MDT care is current practice in
35 cystic fibrosis given that the condition is a multi-system chronic disease (please see the
36 review on MDTs in this guideline for further details). The expertise of the MDT is key for
37 effective management, therefore specialist cystic fibrosis centres should plan patient care,
38 including measures to avoid cross-infection; moreover, they should maintain local and
39 national registers of patients that include information about their clinical condition, treatment
40 and outcomes, as these are important for care, audit and research purposes; the specialist
41 centres should also audit practice and outcomes. These recommendations are consistent
42 with the NHS service specifications for cystic fibrosis, which state that care is to be directed
43 by the specialist centres and that individuals are to be seen by MDTs. The Committee noted
44 that local circumstances, such as the distance to the specialist centre for some patients
45 might support the need for arrangements in which members of the specialist MDT either
46 worked with the local hospital team or indeed came to the local hospital to provide specialist
47 care. Network based shared care might be appropriate for some. This could be provided with
48 local hospitals that have relatively small number of people with cystic fibrosis attending (for
49 example, fewer than 50, given that the CF Trust Standards of Care 2011 mention that the

1 number of patients in a specialist CF centre should not be less than 50). These would
2 receive their care at their local hospital, but with input from the specialist centre (members of
3 the specialist centre MDT coming to work with the local team in clinic) at least twice yearly,
4 which would remain responsible for ensuring the quality of the standard of care provided.
5 Such shared-care models are consistent with current practice for the management of
6 children and young people in the UK. The NHS service specifications for children with cystic
7 fibrosis mention that network care is provided in partnership with the specialist CF centre that
8 co-ordinates the network. The Committee noted that in adult practice shared care was not
9 the norm, and an outreach clinic model was an alternative and more accepted model. The
10 NHS service specifications for adults with cystic fibrosis state that centres serving more rural
11 areas should be able to demonstrate an ability to provide outreach care when appropriate.
12 With the outreach model the cystic fibrosis specialist team directly cares for adults with cystic
13 fibrosis but comes to their local hospital to provide a local cystic fibrosis clinic. In relation to
14 this, the Committee noted that this might pose an equality issue, and more importantly, that it
15 may increase the risk of cross-infection as local hospitals may not have the facilities required
16 to cohort people according to their microbiological profile.

17 In both cases, all people with cystic fibrosis should be managed by a cystic fibrosis core
18 MDT, with access to an extended MDT if necessary, as discussed in the MDT review.

19 With regard to monitoring in network based shared care, the Committee agreed that all
20 people with cystic fibrosis should have an annual assessment and at least one other review
21 per year with the specialist centre MDT. This is consistent with the CF Trust Standards of
22 Care 2011, which are referred to in the NHS service specification for cystic fibrosis. This
23 would be in addition to reviews by the local paediatric team (please see below for the
24 Committee's discussion on the frequency of regular routine reviews). They also agreed that
25 these reviews can be conducted either at the specialist centre, at the network clinic, by
26 telemedicine or at home.

27 Home visits were also considered an appropriate option for some people. Home care is
28 consistent with the NHS service specifications for cystic fibrosis, which state that where
29 appropriate the model of care must ensure access to care at home or close to home and
30 home treatment is encouraged.

31 Given the scarcity and very low quality of the evidence on telemedicine, the Committee used
32 their clinical experience and expertise to make a recommendation. The Committee agreed
33 that it can sometimes be a suitable alternative to outpatient visit for routine monitoring,
34 depending on individual circumstances. The NHS service specifications for cystic fibrosis do
35 not mention telemedicine, however, as explained in the economic benefits and harms
36 section, this can be a supplementary model of care for routine monitoring which has a
37 negligible cost and may be useful to prevent cross-infection. It was also discussed that, in
38 addition to this, people should be able to access specialist advice 24/7 for urgent situations.
39 This is consistent with current practice, with the NHS service specification for cystic fibrosis,
40 and adheres to good clinical practice recommendations.

41 In relation to the administration of intravenous antibiotics in people with cystic fibrosis
42 experiencing an acute exacerbation or with chronic pulmonary disease, this could sometimes
43 have advantages for the person with cystic fibrosis, but the choice between home or hospital
44 therapy would depend on individual circumstances. This is consistent with the NHS service
45 specifications for cystic fibrosis, which recommend that intravenous antibiotics may be
46 provided at home where appropriate. The Committee agreed that it was important to ensure
47 that appropriate training is provided for those involved in providing home IV therapy.

48 The Committee agreed that routine monitoring is key in ensuring effective management
49 because early detection of declining health can prompt early intervention, which in turn can
50 prevent or limit symptoms and complications of cystic fibrosis. Moreover, routine monitoring
51 allows healthcare professionals to regularly review and adjust management and self-
52 management interventions, which supports people's adherence to treatment. Additionally,

1 the Committee agreed that annual assessments are important to ensure that progress is
2 regularly assessed and that a 12-month management plan is agreed.

3 The Committee made a recommendation that detailed all the assessments that should be
4 included at annual review. This recommendation was based on evidence from a wide range
5 of reviews and consensus discussions about annual reviews and assessments that are
6 outlined more in detail in different sections of the guideline. The Committee agreed that they
7 wanted one overarching recommendation that summarised all reviews that were required
8 annually.

9 They also discussed the frequency with which people with cystic fibrosis should be reviewed.
10 Current practice is usually to review 6 times per year, but they agreed that the frequency
11 would vary considerably depending on individual circumstances. Elsewhere, for example, the
12 Committee had made specific recommendations for monitoring pulmonary disease, where
13 the advice varied depending on age and the clinical situation. Based on their clinical
14 experience and expertise, the Committee agreed to recommend more frequent routine
15 reviews after diagnosis and in early life, when carers are not yet familiar with the condition
16 and need more guidance to identify any issues and provide appropriate care. Moreover, as
17 mentioned above in the economic benefits and harms section, after a diagnosis of cystic
18 fibrosis frequent reviews are important to prevent the potentially lethal consequences of
19 inappropriate polypharmacy. Following those initial years, the person's condition is likely to
20 become more predictable. Therefore, the Committee gave some examples of review
21 frequency for different age groups, with frequency decreasing with age, from weekly reviews
22 in the first month of life to every 3 to 6 months for adults. The examples for each age group
23 were based on the Committee's clinical experience and expertise. The NHS service
24 specifications for cystic fibrosis state that routine outpatient appointments should be every 2
25 to 3 months when stable and more often if not for both adults and children. However the NHS
26 service specifications for cystic fibrosis also refer to the CF Trust Standards of Care 2011,
27 which specify that patients should be reviewed with a frequency appropriate to their
28 individual needs, and that newly diagnosed infants should be seen more frequently (initially
29 weekly). The Committee's examples indicated to review children up to 5 years old more
30 frequently (6 to 8 weeks) than the NHS service specifications for cystic fibrosis, and to review
31 adults less frequently than the NHS service specifications, however the NHS service
32 specifications and the Committee agreed that review frequency should be based on
33 individual circumstances.

34 The Committee discussed potential equality issues. They noted that people who live far from
35 a specialist centre may be disadvantaged. However they agreed no additional
36 recommendations were needed as the use of alternative models care had already
37 extensively been discussed (for example the access to shared care in paediatrics, outreach
38 clinics for adult care or telemedicine).

39 The Committee agreed that research recommendations were not prioritised for this topic
40 because clinical practice and the available models of care are already well established.
41 Models of care need to be responsive to and developed according to local needs and factors
42 (such as geography, as outlined above) which by definition are not suited to a general
43 research recommendation.

44 **7.1.7.6 Key conclusions**

45 The Committee concluded that care for all people with cystic fibrosis should be the
46 responsibility of a recognised specialist centre, but in paediatrics this could be provided
47 through shared care clinics as part of a network arrangement. All people should have an
48 annual assessment with their core MDT, and regular monitoring should be scheduled
49 depending on their individual needs. The core MDT should either include or have access to
50 relevant specialists with experience of cystic fibrosis.

1 **7.1.8 Recommendations**

2 **13. Care for people with cystic fibrosis should be provided by a specialist cystic**
3 **fibrosis multidisciplinary team based at a specialist cystic fibrosis centre. (See**
4 **multidisciplinary teams)**

5 **14. Specialist cystic fibrosis centres should:**

- 6 • plan patient care (including outpatient and inpatient care), taking into
7 account the risk of cross-infection (see Prevention of cross infection)
- 8 • maintain local and national registers of patients that include information
9 about their clinical condition, treatment and outcomes
- 10 • audit practice and outcomes.

11 **15. When a shared-care model is used for children and young people, it should**
12 **include:**

- 13 • formal arrangements between the local paediatric team at the shared
14 care centre and multidisciplinary team at the specialist cystic fibrosis
15 centre
- 16 • direct involvement of specialist cystic fibrosis multidisciplinary team
17 members
- 18 • an annual assessment and at least one other review per year by the
19 specialist cystic fibrosis multidisciplinary team in addition to reviews by
20 the local paediatric team (see annual and routine reviews).

21 **16. If available and when clinically appropriate, outreach care for adults with cystic**
22 **fibrosis may be provided by the specialist cystic fibrosis multidisciplinary team at**
23 **a local hospital.**

24 **17. The specialist cystic fibrosis multidisciplinary team should have a member**
25 **available for people with cystic fibrosis and their family members or carers (as**
26 **appropriate) to contact when they have urgent enquiries at any time (day and**
27 **night).**

28 **18. Consider telemedicine or home visits for routine monitoring where they are more**
29 **appropriate than outpatient visits and if the person with cystic fibrosis prefers it.**

30 **19. Make arrangements (including providing equipment and expert support) for**
31 **people to have intravenous antibiotic therapy at home, when this is appropriate.**

32 **Annual and routine reviews**

33 **20. Be aware that:**

- 34 • the aim of cystic fibrosis care is to prevent or limit symptoms and
35 complications of the condition
- 36 • routine monitoring and annual assessments are crucial in providing
37 effective care.

38 **21. Offer people with cystic fibrosis a comprehensive annual review that includes the**
39 **following:**

- 40 • a pulmonary assessment (see Pulmonary monitoring)

- 1 • an assessment of nutrition and intestinal absorption (see Nutritional
- 2 Interventions and Exocrine pancreatic insufficiency)
- 3 • as assessment for liver disease (see Monitoring for liver disease)
- 4 • testing for cystic fibrosis related diabetes, from 10 years of age (see
- 5 Monitoring for cystic fibrosis related diabetes)
- 6 • An assessment for other potential or existing cystic fibrosis
- 7 complications (see Complications of cystic fibrosis)
- 8 • assessments by a specialist physiotherapist, dietician, pharmacist and
- 9 clinical psychologist (see Service delivery)
- 10 • a review of the exercise programme (see Exercise).

11 **22. Provide regular routine reviews for people with cystic fibrosis, and do these more**
12 **frequently immediately after diagnosis and in early life. For example:**

- 13 • weekly in their first month of life
- 14 • every 4 weeks when they are between 1 and 12 months old
- 15 • every 6 or 8 weeks when they are between 1 and 5 years old
- 16 • every 8 or 12 weeks when they are over 5 years old
- 17 • every 3 or 6 months as adults.

18 **7.2 Multidisciplinary team**

19 **Review question: What is the clinical and cost-effectiveness of multidisciplinary teams**
20 **of various compositions?**

21 **7.2.1 Introduction**

22 Cystic fibrosis is a multi-system chronic disease that affects the respiratory tract and lungs,
23 digestive system, sweat glands and reproductive organs. The condition is typically identified
24 in infancy, and care is required throughout an individual's lifetime through to end of life. The
25 care aims to address the biological and psychosocial needs of the patient and their families
26 or carers. As cystic fibrosis is associated with poor quality of life and clinical outcomes, it is
27 important that care adequately addresses the needs of patients by allowing flexibility for
28 individual circumstances. This requires an experienced multidisciplinary team (MDT) of cystic
29 fibrosis-specialist healthcare professionals to ensure optimal care with continuity. This brings
30 up the question of what would represent an ideal MDT.

31 **7.2.2 Description of the clinical evidence**

32 The aim of this review was to compare different combinations of the individuals working
33 together as a core MDT with combinations of the individuals working together as an
34 extended MDT and examines key patient and clinical outcomes.

35 As per protocol, studies that assessed the effectiveness of multidisciplinary teams of various
36 compositions were eligible for inclusion in this review if they were randomised controlled
37 trials or comparative prospective and retrospective cohort studies from Western countries.
38 Studies based on registry and audit data from the UK were also eligible for inclusion.
39 Moreover, conference abstracts of randomised controlled trials (RCTs) were also considered
40 if the full papers were unavailable. However, no relevant studies were identified for inclusion
41 and no evidence was available to inform the review.

42 For full details see review protocol in Appendix D.

43 See also study selection flow chart in Appendix F, and excluded studies list in Appendix H.

1 7.2.3 Summary of included studies

2 Not applicable, as no studies were included in this review.

3 7.2.4 Clinical evidence profile

4 Not applicable, as no studies were included in this review.

5 7.2.5 Economic evidence

6 No economic evaluations of MDTs were identified in the literature search conducted for this
7 guideline. Full details of the search and economic article selection flow chart can be found in
8 Appendix E and F, respectively.

9 This review question was not prioritised for de novo economic modelling. To aid
10 considerations of cost-effectiveness a costing tool was developed that utilised a “what-if”
11 approach to estimate the cost of providing the core MDT specified in the protocol. This tool
12 enables the user to vary the composition of the MDT, the number of whole time equivalents
13 and the size of the clinic the MDT manages.

14 The cost of health care professionals included in the protocol are provided in Table 42, whilst
15 a full description of the methods and results of the analysis are presented in Appendix K.

16 **Table 42: Cost of providing the MDT at the Specialist Centre**

Health care professional	Cost per annum	Cost per hour	Source (bands informed by the Committee)
Core MDT			
Specialist CF Clinician	£190,408 ^a	£105 ^b	PSSRU 2016: Hospital-based doctors, medical consultant
Specialist Nurse	£83,628 ^c	£53 ^d	PSSRU 2016: Band 7, hospital-based nurses
Specialist Dietician	£85,739 ^f	£54 ^e	PSSRU 2016: Band 7, scientific and professional staff
Specialist Physiotherapist	£87,381 ^g	£55 ^e	PSSRU 2016: Band 7, scientific and professional staff
Specialist Pharmacist	£101,367 ^h	£64 ^e	PSSRU 2016: Band 8a, scientific and professional staff
Specialist Psychologist	£101,367 ^h	£64 ^e	PSSRU 2016: Band 8a, scientific and professional staff
Specialist Social worker	£61,730 ⁱ	£40 ^j	PSSRU 2016: Social worker (adult services)
Specialist Social worker	£58,947 ^k	£39 ^j	PSSRU 2016: Social worker (children’s services)
Extended MDT			
Paediatric Diabetic Medicine	£253 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance Follow-up, Consultant-led, Paediatric Diabetic Medicine 263
Diabetic Medicine	£159 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance Follow-up, Consultant-led, Diabetic Medicine 307
Paediatric ENT surgeon	£103 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance, Follow-up, Consultant led, Paediatric Ear Nose And

Health care professional	Cost per annum	Cost per hour	Source (bands informed by the Committee)
			Throat 215
ENT surgeon	£89 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance, Follow-up, Consultant led, Ear Nose And Throat 120
Obstetrician	£121 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance, Follow-up, Consultant-led, Obstetrics 501
General surgeon	£123 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance, Follow-up, Consultant led, General Surgery 100
Gastroenterologist/ hepatologist	£253 per attendance		NHS Reference Costs 2015/16: WF01A, Non-Admitted Face to Face Attendance, Follow-up, Consultant led, Hepatology 306

- 1 (a) Including wages/salary £87,449; salary oncosts £23,198; management, admin and estates staff overheads
2 £26,777; non-staff overheads £47,689 and capital overheads £5,295
3 (b) Working time 42.3 weeks (1,838 hours) per year 43.3 hours per week
4 (c) Including wages/salary £38,550; salary oncosts £9,605; management, admin and estates staff overheads
5 £11,653; non-staff overheads £20,755 and capital overheads £3,065
6 (d) Working time 42 weeks (1,572 hours) per year, 37.5 hours per week
7 (e) Working time 42.7 weeks (1,603 hours) per year, 37.5 hours per week
8 (f) Band 7 Dietitian. Including wages/salary £38,786; salary oncosts £9,670; management, admin and estates
9 staff overheads £11,726; non-staff overheads £20,885 and capital overheads £4,672
10 (g) Band 7 Physiotherapist. Including wages/salary £38,786; salary oncosts £9,670; management, admin and
11 estates staff overheads £11,726; non-staff overheads £20,885 and capital overheads £6,314
12 (h) Band 8a. Including wages/salary £46,095; salary oncosts £11,702; management, admin and estates staff
13 overheads £13,987; non-staff overheads £24,911 and capital overheads £4,672
14 (i) Including wages/salary £31,288; salary oncosts £9,463; direct overheads £11,818; indirect overheads £6,520
15 and capital overheads £2,641
16 (j) Working time 41 weeks (1,517) 37 hours per week
17 (k) Including wages/salary £29,854; salary oncosts £8,978; direct overheads £11,261; indirect overheads £6,213
18 and capital overheads £2,641

19 7.2.6 Evidence statements

20 Not applicable, as no clinical or economic studies were included in this review.

21 7.2.7 Evidence to recommendations

22 7.2.7.1 Relative value placed on the outcomes considered

23 The aim of this review was to compare different combinations of the individuals working
24 together as a core MDT with combinations of the individuals working together as an
25 extended MDT and examines key patient and clinical outcomes.

26 The Committee identified the following outcomes as critical: mortality, forced expiratory
27 volume in one second (FEV₁) and patient satisfaction. Lung clearance index (LCI), time to
28 next pulmonary exacerbation, nutritional status, quality of life, carer satisfaction, frequency of
29 cross-infections (*P aeruginosa*, *B cepacia*), staff experience and adherence to treatment
30 were rated as important outcomes.

31 7.2.7.2 Consideration of clinical benefits and harms

32 Cystic fibrosis is a multi-system chronic disease that affects the respiratory tract, digestive
33 system, sweat glands and reproductive organs. The condition is typically identified in infancy
34 and care is required throughout an individual's lifetime through to end of life. The care aims
35 to address the biological and psychosocial needs of the patient and their families or carers
36 and, in the UK, is provided by a specialist cystic fibrosis centre. As cystic fibrosis is

1 associated with poor quality of life and clinical outcomes, it is important that care adequately
2 addresses the needs of patients by allowing flexibility for individual circumstances.

3 Given the nature of this condition, the Committee considered it important to provide guidance
4 on the professionals that should be responsible for the provision of care. Due to the lack of
5 evidence, the recommendations were based on the Committee's knowledge and expertise
6 and were agreed by consensus.

7 In relation to the composition of the MDT, the Committee discussed which healthcare
8 professionals should be part of the core team. It was agreed the core team should include
9 the following specialists: cystic fibrosis clinicians, nurses, dietitians, physiotherapists,
10 pharmacists, psychologists and social workers. This is consistent with the NHS service
11 specifications for cystic fibrosis. Each of the specialist members should be a member of their
12 specialist cystic fibrosis group/organization and should have expert knowledge in cystic
13 fibrosis.

14 The Committee agreed that within the MDT there should be a specialist paediatrician or adult
15 physician that provides leadership for the MDT. Specialist paediatricians and adult
16 physicians would have the necessary breadth and depth of experience and knowledge
17 regarding cystic fibrosis to equip them for this role.

18 The Committee noted that specialist nurses communicate with people with cystic fibrosis and
19 their family members or carers about their overall care needs rather than focusing on one
20 area of specialisation, therefore are best placed to coordinate their care, coordinate
21 communication between the members of the team, and act as advocates for their best
22 possible care.

23 The Committee agreed that in their experience, people with cystic fibrosis benefit from
24 regular meetings with a specialist physiotherapist, who advises them on airway clearance,
25 nebuliser use, musculoskeletal disorders, exercise and physical activity, and urinary
26 incontinence. People with cystic fibrosis also benefit from regularly seeing a specialist
27 dietitian, who advises them on nutrition. Normally, a review from a physiotherapist and a
28 dietitian would be an integral part of a clinic visit. These reviews should be available at clinic
29 visits, however the frequency with which these reviews are necessary would be a matter for
30 individual consideration. The Committee considered that every person should have a
31 physiotherapist and a dietitian review as part of their annual review. Moreover, they agreed
32 that the input of a physiotherapist and a dietitian would be important during inpatient
33 admissions. Finally, they recommended physiotherapist assessments during pulmonary
34 exacerbations.

35 The Committee agreed that the specialist pharmacist should be available when necessary to
36 advise on matters related to medications, and people with cystic fibrosis should have a
37 pharmacist review as part of their annual review.

38 The Committee agreed that people with cystic fibrosis should have a specialist clinical
39 psychologist review as part of their annual review to identify psychological and behavioural
40 problems and offer advice; moreover, the specialist clinical psychologist should be available
41 to see the patient at outpatient visits and during inpatient admissions.

42 The Committee noted that the CF Trust Standards of Care 2011 (which are mentioned in the
43 NHS service specifications for cystic fibrosis as the standards to follow) offered a detailed
44 description of the role of each professional included in the core MDT.

45 In addition, the Committee discussed the health care professionals that should also be
46 involved in the MDT, depending on the person's circumstances. This extended team should
47 include: diabetologists, obstetricians, ENT surgeons, general surgeons and
48 gastroenterologists or hepatologists. The Committee emphasized the importance of the
49 health care professionals having cystic fibrosis expertise or interest.

1 Although not specifically covered in the protocol, the Committee agreed that microbiologists,
2 interventional radiologists, pulmonary physiologists, rheumatologists, thoracic surgeons and
3 palliative care specialists can play an important role in the management of people with cystic
4 fibrosis, and that the MDT should have access to them where appropriate. This is consistent
5 with current practice.

6 The Committee agreed that access to the aforementioned professionals would be necessary
7 to provide high quality care in the areas of microbiology, pulmonary physiology, diabetes,
8 gastroenterology, hepatology, rheumatology, psychiatry, interventional radiology, surgery
9 (gastrointestinal and thoracic), obstetrics and palliative care.

10 The Committee also discussed the need to have access to secretarial and data entry
11 support. This is already current practice, whether the staff is only dedicated to the MDT or
12 not, and it is supported by other consensus recommendations.

13 **7.2.7.3 Consideration of economic benefits and harms**

14 The Committee agreed that the core MDT composition should follow previous guidance
15 (such as recommendations of the Cystic Fibrosis Trust's and NHS service specifications for
16 cystic fibrosis) as this represents current clinical practice and resource use in England.
17 Moreover, the core MDT would have expertise in those specialities regularly sought by all
18 people with cystic fibrosis to provide a cost-effective full-time composition.

19 The Committee considered the WTE recommendations reported in the CF Trust Standards
20 of Care 2011 for a given clinic size. They concluded that the values they specified were
21 insufficient to provide a high quality service, given the need for the core MDT to be spread
22 over acute hospital work, outpatient clinics, network clinics and community work, such as
23 home visits. The Committee also highlighted that the number of WTEs did not allow for the
24 provision of annual leave or sick leave cover. However, the Committee were reluctant to
25 suggest a minimum WTE for each of the core members to prevent interpretation of a
26 maximum under austerity and because this would vary depending on the complexity of the
27 people with cystic fibrosis at each clinic.

28 The Committee highlighted the importance of administrative support to disseminate
29 information across the models of care and provide data entry into the Cystic Fibrosis
30 Registry, which is useful to monitor the complications and comorbidities of cystic fibrosis and
31 its management. The Committee considered this to be a cost-effective use of an
32 administrator's time when the collected data can be used to improve people's knowledge of
33 cystic fibrosis.

34 The Committee agreed the extended MDT would not be full-time members of the MDT as
35 their case load would include people without cystic fibrosis. One defined person within the
36 specialty would be called upon on a case-by-case basis as this would promote continuity,
37 cystic fibrosis expertise, and hence effectiveness. As a result, the Committee considered the
38 opportunity cost of the extended MDTs time to care for people without cystic fibrosis, and
39 made a recommendation for the core MDT to have access to those (extended) specialists
40 that can provide advice and accept referrals for people with cystic fibrosis.

41 In addition to the extended MDT outlined in the protocol, the Committee discussed how the
42 wider MDT service specification reported by NHS England should reflect current NHS
43 commissioning. However, it was noted that such service specifications by NHS England are
44 not evidence based and could not justify an increase in current resource use where they are
45 not followed, as there is no evidence that those specifications are a cost-effective use of
46 resources for wider implementation.

47 The Committee stated that people with cystic fibrosis should have access to a
48 physiotherapist, dietician, pharmacist and psychologist at outpatient clinic visits and annual
49 reviews, to ensure changes to their health and goals are identified and managed without

1 potentially harmful delays. Consequently, the Committee made recommendations to that
2 effect and added that the CF Trust Standards of Care 2011 (which are mentioned in the NHS
3 service specifications for cystic fibrosis as the standards to follow) offered a detailed
4 description of the role of each professional included in the core MDT.

5 Furthermore, during inpatient admissions, dietitians and physiotherapists need to ensure that
6 their patient's exercise programmes and nutritional requirements compliment their inpatient
7 treatment schedules (see recommendations in sections 10.1.8 and 10.8.8 for nutrition and
8 exercise, respectively). For these reasons, the Committee agreed their recommendations to
9 ensure access during inpatient admissions promoted a cost-effective use of resources as
10 management strategies usually need to be modified during inpatient care.

11 **7.2.7.4 Quality of evidence**

12 Not applicable as no evidence was found for this review.

13 **7.2.7.5 Other considerations**

14 Given the lack of evidence, the recommendations were based on the Committee members'
15 clinical experience, and consensus on good clinical practice (NHS service specifications for
16 cystic fibrosis; CF Trust Standards of Care 2011, which are mentioned in the NHS service
17 specifications as the standards to follow; and European Cystic Fibrosis Society Standards of
18 Care 2014).

19 No equality issues were identified by the Committee for this review question.

20 The Committee felt a research recommendation into the MDT was not needed as their
21 clinical experience and expertise was sufficient to show the need for all members outlined
22 above.

23 **7.2.7.6 Key conclusions**

24 The Committee concluded that the core MDT team should include the following specialist:
25 cystic fibrosis clinicians, nurses, dietitians, physiotherapists, pharmacists, psychologists and
26 social workers. In addition, the following professionals with expertise in cystic fibrosis may
27 also be included in the team if appropriate: diabetologists, obstetricians, ENT surgeons,
28 general surgeons and gastroenterologists or hepatologists. Although not strictly part of the
29 team, the MDT should have access to administrative and cystic fibrosis registry data entry
30 staff.

31 **7.2.8 Recommendations**

32 **23. The specialist cystic fibrosis multidisciplinary team should include the following**
33 **professionals who have specialist expertise in the condition:**

- 34 • specialist paediatrician or adult physician
- 35 • specialist nurse
- 36 • specialist physiotherapist
- 37 • specialist dietitian
- 38 • specialist pharmacist
- 39 • specialist clinical psychologist
- 40 • social worker.

41 **24. The specialist cystic fibrosis multidisciplinary team should be led by a specialist**
42 **paediatrician or adult physician.**

- 1 **25. Specialist nurses should coordinate care, coordinate communication between**
2 **other members of the team, and act as advocates for people with cystic fibrosis**
3 **and their family members or carers (as appropriate).**
- 4 **26. The specialist physiotherapist should assess and advise people with cystic**
5 **fibrosis at clinic, at inpatient admissions, during pulmonary exacerbations and at**
6 **their annual review on:**
- 7 • airway clearance
8 • nebuliser use
9 • musculoskeletal disorders
10 • exercise and physical activity
11 • urinary incontinence.
- 12 **27. The specialist dietitian should assess and advise people with cystic fibrosis about**
13 **nutrition at outpatient clinic visits, during inpatient admissions and at their annual**
14 **review.**
- 15 **28. The specialist pharmacist should advise people with cystic fibrosis on medicines**
16 **optimisation at outpatient clinic visits and at their annual review.**
- 17 **29. The specialist clinical psychologist should assess and advise people with cystic**
18 **fibrosis at their annual review, and when needed at outpatient and inpatient**
19 **clinics.**
- 20 **30. The specialist cystic fibrosis multidisciplinary team should either include or have**
21 **access to specialist expertise relevant to cystic fibrosis in the following areas:**
- 22 • microbiology
23 • pulmonary physiology
24 • diabetes
25 • gastroenterology
26 • hepatology
27 • rheumatology
28 • psychiatry
29 • interventional radiology
30 • surgery (gastrointestinal and thoracic)
31 • obstetrics
32 • palliative care.

33 **7.3 Transition to adult services**

34 **Review question: What parts of the transition from children's to adult services are**
35 **most important for young people with cystic fibrosis and their family members and**
36 **carers?**

37 **7.3.1 Introduction**

38 Transitional care from paediatrics to adult services is a complex process that involves a
39 multidisciplinary team approach. There is an obligation to ensure that the person's health
40 care needs are met at each step of the way. Young people should be transferred to the adult

1 service when they have completed growth and puberty. Where possible they should be
2 largely independent of parents and staff, for example with regards to communication and
3 decision making. Transition planning must begin well before the anticipated transfer time. A
4 transition programme is an essential part of quality care for young people with cystic fibrosis
5 and their parents or carers. In early adolescence it is favourable that a series of education
6 sessions take place. These should include disease understanding, the rationale behind
7 therapy and recognising and responding to symptoms and deterioration. Cystic fibrosis
8 teams, as well as people with cystic fibrosis, have found that the transfer of care from a
9 centre to another must be a gradual process rather than an abrupt shift. Transition is not
10 synonymous with transfer. It is an active process and not a single event. It must begin early,
11 be planned and regularly reviewed and be age and developmentally appropriate. There is a
12 lack of standardisation currently in transition practices, and people with cystic fibrosis and
13 their families report inconsistencies.

14 Young people themselves are nervous of change and worried about the new responsibility.
15 Parents are concerned about the level of care provided by a new team and fear the loss of
16 their child to adulthood with resultant loss of control over their care. Additionally, transition
17 often occurs during a time of emotional and social flux for the young person as in addition to
18 their change in treatment they are undergoing the same changes in schooling, work and
19 relationships which all people go through on maturity to adulthood.

20 Transition must be a collaboration between paediatric and adult centres, the person with
21 cystic fibrosis and their family. There must be input for psychological and practical
22 preparation and adjustment. We need to equip young adults with cystic fibrosis with the skills
23 to negotiate the adult system of health care. Paediatric centres cannot offer emotional
24 support nor offer the surroundings or service appropriate for adults. Paediatric centres focus
25 on the well-being of the whole family whereas adult centres focus on the patient. This can
26 lead to emotional strains as parents or care-givers begin to 'take a backseat' in the
27 management of their child's health. This fundamental difference in approach has major
28 implications for the emotional care of the young person and their families and adjustments or
29 interventions may need to be made. Preparation for the change can greatly affect the ease of
30 transition. The role of the paediatric team must be to not show bias, respect the rights of the
31 teenager, offer guidance and support, discuss the future, provide information about the adult
32 services and facilitate the move both practically and emotionally. The adult service needs to
33 be aware of these concerns and act to manage the transition not just from paediatrics to the
34 adult centre but also the transition of the young person from a dependent-child to
35 independent young adult, including support for primary care-givers.

36 The Committee recognises that effective transition is important for all young people with
37 health needs, as outlined in the NICE guideline on transition from children's to adult's care
38 (NG43). However, they were specifically focused on aspects that are particularly important or
39 unique for young people with cystic fibrosis, their families and carers.

40 **7.3.2 Description of clinical evidence**

41 The aim of this review was to identify elements of the transition process (for example,
42 transition planning involvement) from paediatric to adult services from perspectives of young
43 people with cystic fibrosis and their family and carers.

44 We looked for studies that collected data using qualitative methods (such as semi-structured
45 interviews, focus groups and surveys with open-ended questions) in which the authors
46 analysed the data qualitatively (including thematic analysis, framework thematic analysis or
47 content analysis). Survey studies restricted to reporting descriptive data that were analysed
48 quantitatively were excluded.

1 Given the nature of qualitative reviews, findings or themes were summarised from the
2 literature and were not restricted to those identified as likely themes by the guideline
3 Committee at protocol stage.

4 For full details see review protocol in Appendix D.

5 Eleven qualitative studies were included in this review (Al-Yateem 2012, Al-Yateem 2013,
6 Begley 2013, Brumfield 2004, Depuis 2011, Iles 2010, Moola 2011, Russell 1996, Tierney
7 2013, Tuchman 2008, Van Staa 2011).

8 Seven studies used semi-structured interviews (Dupuis 2011, Iles 2010, Moola 2011, Russell
9 1996, Tierney 2013, Tuchman 2008, van Staa 2011), 2 studies used in depth interviews (Al-
10 Yateem 2013, Brumfield 2004), 1 study used a postal survey (Begley 2013) and 1 study used
11 focus groups (Al-Yateem, 2013). The most common data analysis method employed across
12 studies was thematic analysis.

13 With regards to the population, 9 studies included young people and young adults (age
14 range: (Al-Yateem 2012, Brumfield 2004, Depuis 2011, Iles 2010, Moola 2011, Russell 1996,
15 Tierney 2013, Tuchman 2008, Van Staa 2011), 4 studies included parents (Dupuis 2011,
16 Moola 2011, Russell 1996, Van Staa 2011) and 4 studies included healthcare professionals
17 working with young people and young adults with cystic fibrosis (Al-Yateem 2013, Begley
18 2013, Iles 2010, Van Staa 2011).

19 The sample size of the studies ranged from 4 (Brumfield 2004) to 132 participants (Begley
20 2013). Three studies included a mixed population (Begley 2013, Moola 2011, Tuchman
21 2008), but only quotes from people with cystic fibrosis was retrieved.

22 Two studies were conducted in the UK (Iles 2010, Tierney 2013), 3 in Ireland (Al-Yateem
23 2012, Al-Yateem 2013, Begley 2013), 2 in Australia (Brumfield 2004, Russell 1996), 2 in
24 Canada (Dupuis 2011, Moola 2011), 1 in the USA (Tuchman 2008), and 1 in the Netherlands
25 (van Staa 2011).

26 See study selection flow chart in Appendix F, study evidence tables in Appendix G and list of
27 excluded studies in Appendix H. For presentation of findings, a theme map was generated
28 according to the themes emerged from studies (Figure 1). Due to the qualitative nature of
29 these studies, evidence is summarised in GRADE-CERQual tables within the evidence
30 report. Therefore no separate Appendix is provided for this.

31 7.3.3 Summary of included studies

32 A summary of the studies that were included in this review are presented in Table 43.

33 **Table 43: Summary of included studies**

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
Al-Yateem (2012) Ireland	In-depth interviews	N=15 young adults with CF Age of young adults: Not reported	To explore and understand the experience of young people before and after their transitional care, and the factors that both contribute to and hinder that experience.	<ul style="list-style-type: none"> • Overall quality: low • Age of young adult not reported. • Description of data collection method was vaguely described. • The analytical process was reported vaguely. • Description of emerging and overarching themes was reported, but saturation of data

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
				was not reported.
Al-Yateem (2013) Ireland	Focus groups	Health care professional working with young adults with CF Sample size and age group not reported	To develop relevant and feasible guidelines for transition care, based on perspectives of stakeholders.	<ul style="list-style-type: none"> Overall quality: low Number of participants and age group not reported. Description of data collection method was vaguely described. The analytical process poorly described.
Begley (2013) Ireland	Postal survey	N=132 healthcare professionals	To evaluate how the transition from child to adult healthcare is managed in young people with CF or diabetes in Ireland.	<ul style="list-style-type: none"> Overall quality: low The sample selection process was not clearly reported, although the authors tried to involve all participants caring after young people before the transition process, 56% response rate. Description of data collection method and data analysis was vaguely described. The authors do not explain whether data saturation was reached, but given the limited number of quotes presented in the study it is unlikely. This study includes healthcare professionals looking after young people with CF and diabetes. Only quotes relevant to CF have been extracted.
Brumfield (2004) Australia	In-depth interviews	N=4 young adults with CF Age: 19 to 22 years	To explore the experiences of Australian adolescents with CF as they made the transition from paediatric to adult care.	<ul style="list-style-type: none"> Overall quality: moderate Small sample size. Data collection was described in detail but no information on data saturation. Robustness of the finding not discussed considering the low sample.
Dupuis (2011) Canada	Semi-structured interviews	N=26 <ul style="list-style-type: none"> 7 young people with CF 7 mothers, 4 	To explore the experience of parents and adolescents living with cystic fibrosis prior to the transfer of	<ul style="list-style-type: none"> Overall quality: low No information on structure of interview or whether topic guide reported.

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
		fathers <ul style="list-style-type: none"> 8 members of the healthcare team Age of people with CF: 15 to 18 years	the adolescent's care from a paediatric to an adult healthcare facility.	<ul style="list-style-type: none"> Themes and categories used but no information on data saturation and full exploration of theme. Most of the findings unrelated to transitions process.
Iles (2010) UK	Semi-structured interviews	N=73 <ul style="list-style-type: none"> 50 young people and young adults with CF 23 healthcare professionals Age of people with CF: 13 to 24 years	To examine how young people and staff perceive the nature of parental care and support for those with CF who have left paediatric services.	<ul style="list-style-type: none"> Overall quality: moderate Sample selection was clearly reported. The relationship between the researcher and the respondents was reported. Information on structure of interview or topic guide reported. The analytical process was described with description of themes, categories and use of qualitative software.
Moola (2011) Canada	Semi-structured interviews	N=78 <ul style="list-style-type: none"> 50 young people; 28 parents affected by CF and coronary heart disease Age of young people: 11 to 17 years	To understand the transition experiences of young people with CF and coronary heart disease.	<ul style="list-style-type: none"> Overall quality: moderate The sample selection process was vaguely reported, and the relationship between the researcher and the respondents was not reported. All the steps of the thematic analysis were reported, but it was unclear whether data saturation was reached. Results were presented clearly, but very few themes referred to the transition from child to adult services. This study includes a mixed population. Only quotes relevant to CF have been extracted.
Russell (1996) Australia	Semi structured interviews	N=15 <ul style="list-style-type: none"> 7 young people and young adults with CF and 8 parents Age of young	To investigate the experience of transferring to adult healthcare from the perspective of adolescents with CF and their parents.	<ul style="list-style-type: none"> Overall quality: moderate Data collection method was well described and consistency in the interview process reported.

Study	Study design/ methods	Participants /respondent	Aim of the study	Comments
		people with CF: 11 to 20 years		<ul style="list-style-type: none"> Description of emerging and overarching themes was reported, but saturation of data was not reported.
Tierney (2013) UK	Semi-structured interviews	N=19 young people and young adults with CF Age: 17 to 19 years	To explore the experiences of young people with cystic fibrosis of transferring to adult services.	<ul style="list-style-type: none"> Overall quality: moderate Data collection and analysis method was well described and consistency in the interview process reported. Findings were presented clearly. Robustness of the overall findings discussed.
Tuchman (2008) USA	Semi-structured interviews	N=5 young people and young adults with CF Age: 15 to 21 years	To describe expectations and concerns of adolescents with chronic illness regarding transition from sub-speciality paediatric to adult-centred care during the transition process in order to guide effective programme design and implementation.	<ul style="list-style-type: none"> Overall quality: low Mixed sample with other chronic diseases. No inclusion criteria reported. Data collection and analysis was vaguely described and lacked detailed information.
van Staa (2011) Netherlands	Semi-structured interviews	N=8 <ul style="list-style-type: none"> 3 young people and adults with CF; 3 parents; 2 healthcare providers Age of people with CF: 15 to 22 years	<ul style="list-style-type: none"> To explore map experiences with the transfer to adult care of young adults with chronic conditions. To identify recommendations for transitional care of young adults, their parents and healthcare providers. 	<ul style="list-style-type: none"> Overall quality: moderate Mixed sample with other chronic diseases. Sample selection process was vaguely reported. Data collection and analysis process was well described. Findings were well presented but use of quotes was inadequate.

1 Abbreviations: CF: cystic fibrosis; UK: United Kingdom


2 7.3.4 Clinical evidence

3 7.3.4.1 Theme map

4 The theme map is presented in Figure 5.

5

Figure 5: Theme map: Needs of young adults in transition to adult health services



7.3.51 Clinical evidence profile

2 The clinical evidence (GRADE-CERQual) for the transition review question is presented in Table 44 to Table 48.

7.3.5.13 Overarching themes

4 Table 44: Summary of clinical evidence (GRADE-CERQual): Overarching themes

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: Transition as natural stage					
3 studies (Brumfield 2004, Moola 2011, van Staa 2001)	3 studies using interviews	<p>Two studies conducted in Australia, Canada and the Netherlands with young people and young adults with cystic fibrosis, parents and healthcare professionals noted that transition was seen as a natural process.</p> <p>Age appropriate treatment was considered and issue by people with cystic fibrosis: <i>"As I grew up, he (paediatric doctors) sort of treated me . . . as if I was older . . . he didn't start treating me like a little kid and stuff, so that was good." (19 year old with cystic fibrosis)</i></p> <p>However, another individual said that they were treated as children even when they were young adults ready for transition: <i>"I thought . . . if only the (paediatric) doctors knew that we were becoming adults, you know, we were in our mid-teens thinking, they still treat us as if we were ten years old." (20 year old with cystic fibrosis)</i></p> <p>Young people, young adults and their parents recognised the inevitability that they will be transferred to adult healthcare and the fact they will have to get more involved in their future care: <i>"My mum usually takes care of all the appointments and everything... it will probably change when I get older. When I'm an adult, I'll probably have to make all the appointments and everything" (young girl with cystic fibrosis, age 13).</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY

		<p><i>"I'll need to get used to it. I've known my doctor awfully long, for 18 years. But I'll just see what's going to happen. [. . .] Actually, I'm getting too old now for a children's hospital. Seems to be the right age [for transfer] because I'm an adult now, aren't I?" (19-year-old male, cystic fibrosis)</i></p> <p>Healthcare providers recognised transfer as 'a natural process' that is 'age-appropriate'.</p>			
Sub-theme 2: Parallel changes or transition to adulthood					
2 studies (Moola 2011, Tierney 2013)	2 studies using interviews	<p>In 1 study conducted in the UK with young people and young adults with cystic fibrosis, participants highlighted that relocating to a new healthcare system is a of several transitions young people with a long-term condition. Hence, transfer takes place against a backdrop of additional pressures and might not be a top priority:</p> <p><i>"It was very busy actually because you know you've got to come in here at some point but then you've also just started at college, a new course, you don't want to miss all the beginning of your course. It's a new start both sides so you don't want to miss either one." (young person with cystic fibrosis)</i></p> <p>In another study conducted in Canada with young people with chronic conditions and parents, the authors concluded that young people projected their lives into the future. Although they showed future aspirations, they contemplated if their health would deteriorate in the future, and the impact this would have on their lives.</p> <p>For example, they suggested it might be important to choose a school or work environment that is in close proximity to a hospital: <i>"I will probably not live in the residence in the future and I will probably live at home with all the medicine near and staff right near me... It is easier to get all of the medicine and stuff; my mum is always driving down to get the medicine. And then I can just bring it home and get it right away. Instead of me living somewhere else and then having to get it. It would be so much more complicated"</i> (young female with cystic fibrosis, age 15)</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY

In 1 study, young women expressed sadness and resignation regarding the potential impact that illness could have on fertility. They were worried that illness would preclude them from being able to have a normal pregnancy:

“I want to have a family, but I cannot have children because of cystic fibrosis” (young female with cystic fibrosis, age 16)

Young people with cystic fibrosis and their parents described they feel threatened by cystic fibrosis, and expressed sadness for the lost time that would result in an early death:

“I also know that I am not going to live as long as everybody else, so that is hard. I feel like it is out of my control, I feel helpless, how I used to be able to do it (physical activity), and now, I can't. It is kind of depressing. It makes me think that it is a progressive disease, and it makes me think that it is getting worse... it makes me worried” (young female with cystic fibrosis, age 15)

“A parent never wants to have a kid die before her and what is what she (wife) was upset about. That is why I was trying to tell her to spend as much time as you can, with her (child with cystic fibrosis). And just think; every waking moment that you have, spend it with her. Even if you have both of them, of one by themselves, spend that time with her. I had her quit her job and that is why I work 16 hours a day. So that she can spend more time with them) (parent of a young person with cystic fibrosis).

“It (cystic fibrosis) is part of who I am... I was thinking about life in general and how I knew that my disease was going to kill me off younger – I will probably not be able to see my grandchildren grow up, kind of thing... It really got to me that whole year and eventually, I just accepted the fact that everybody dies. It does not matter when you go, but you go. I just kind of got a positive attitude, that I might as well be happy and make the most of it” (young male with cystic fibrosis, age 17)

Sub-theme 3: independency in decision making

<p>5 studies (Al-Yateem 2013, Iles 2010, Russell 1996, Tuchman 2008, van Staa 2011)</p>	<p>4 studies using interviews and 1 study using focus groups</p>	<p>Five studies conducted in Australia, Ireland, the Netherlands, the UK and the USA with young people and young adults with cystic fibrosis, parents and healthcare professionals reported on the importance of being allowed to make choices, self-advocacy and build independence gradually.</p> <p>Promoting independence</p> <p>One participant suggested that young adults should be more involved in decision making process:</p> <p><i>"I think it is important to keep them involved ... and take part in all decisions and activities during clinic and so on". (healthcare professional working with young people)</i></p> <p>Increasing active involvement ("parent-as-partner")</p> <p>Young adults noted that more independence and self-reliance was expected of them. Parents wondered whether their children could take up the full responsibility for their treatment. Healthcare providers recognised cultural differences between the paediatric and adult specialities that complicated transfer. The adult care had a more 'business-like approach' which often contrasted with the paediatric 'holistic, system-oriented approach'</p> <p>Young adults want to create their own identity and perform task to gain independence. To gain independence, the young adults were in the process of developing autonomous and self-determined lives and separating from their parents.</p> <p>They also preferred active involvement in their clinic appointments and their desire for a confidential consultation. Many young people reported embracing opportunities to take the lead in adult clinic consultations, negotiating with their parents to facilitate this:</p> <p><i>"Since I went over to the adults' [clinic] it's been me more involved and she's just sat back and she'll take me if I want her to and she'll sit there and she won't say anything unless I ask her." (17 year old with cystic fibrosis)</i></p> <p>Some young adults did not even want their parents to be involved during hospital consultations:</p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Saturated</p>	<p>MODERATE QUALITY</p>
---	--	---	---	---	-------------------------

	<p><i>"My mum used to come with me to the clinic when I first transferred. I mean she'd come with me now if she could get time off work, if I'd let her, but sometimes I'd rather she wasn't there because there are obviously personal things you want to talk to the doctor about, like when I got my first boyfriend and stuff, I didn't want her to be there." (23 year old with cystic fibrosis)</i></p> <p><i>"I didn't tell my mum I was transferring. I didn't tell her, because my mum's a bit obsessive about the clinic and she feels she has to know everything, you know, even though I'm 17... I mean I know it's the duty of the parent, I know that's their job...But that's the good thing about the adult clinic, is the parents don't have to be there, just the child or the patient's wishes. That's the good thing about being in the adult [service]." (17 year old with cystic fibrosis)</i></p> <p>Health professionals stressed the need for understanding the family dynamics but also recognised that young adults are decision makers:</p> <p><i>"I think that when they move to the adult side we very much leave it up to the young person. Some of them leave their parents in the waiting room, others very firmly bring them in." (Specialist physician)</i></p> <p><i>"I make it very clear to the parents, when they come up, that I am quite happy to discuss anything with them that the patients want me to discuss with them, but that they are now adults and it's up to them if I speak to them. I mean I will ask my patients, 'Do you mind if I discuss your treatment options with your parents?' And nine times out of ten, they have no problem with it. But I make it very, very clear to the parents that I can't talk to them the way that I could do or [others] used to when they were children and it's really part of learning to let go for them." (Cystic fibrosis Nurse Specialist)</i></p> <p>In spite of this, the parents persisted in their attempts to be included until they were recognised as an integral part of the adolescent's care.</p> <p>One mother described this as, <i>"I don't know whether they realized that we just weren't going to go away, that they would</i></p>			
--	--	--	--	--

	<p><i>then have to get on with us and we were going to have to get on with them."</i></p> <p><i>"My mom doesn't want to let go. She has flat out told me. You put 18 years into your child's health and it becomes your health as well" (young adult with cystic fibrosis)</i></p> <p><i>"My mom has mixed emotions about it because she's not sitting back in the room with me anymore. She likes being able to put her two cents in. And I like being able to do it myself." (young adult with cystic fibrosis)</i></p> <p>However, in some situations parents were still regarded as protectors. The main way in which young people perceived their parents to protect them was in withholding information during childhood about the terminal nature of cystic fibrosis:</p> <p><i>"You know, I know a person [with cystic fibrosis], my mum knew him. And my mum never wanted to tell me [he had died] until I had turned 15." (16 year old with cystic fibrosis)</i></p> <p>Staff also acknowledged the difficulties communicating limited life expectancy with parents because of their protective nature, where it seemed that young people were perhaps more able than their parents to have these discussions with staff:</p> <p><i>"We had to have a discussion whether to go on to mechanical ventilation would be the right thing And she was able to have that discussion. And I asked her did she want me to let her mother know, if she wasn't going to tell her, that we had had that discussion? She asked me to tell her [mother]. And her mother was initially comfortable that we had had that discussion. Then over the space of about two or three hours, became very agitated and very upset that we'd had it... it was a huge stress for the mother, whereas her daughter, although finding it very difficult, was actually able to have [the conversation] and was – in the end, I think, glad she had had it." (Chest Physician)</i></p> <p>Although some young adults had also started to protect their parents from knowing the extent of their deteriorating health:</p> <p><i>"Mum always used to sit in on consultations until, until I could get rid of her about three years ago. That made it really hard to</i></p>			
--	---	--	--	--

		<i>talk about anything, because Mum obviously gets upset if you mention stuff like dying. So you have to be really careful. (23 year old with cystic fibrosis)</i>			
--	--	--	--	--	--

7.3.5.21 Support before transition to adult services

2 Table 45: Summary of clinical evidence (GRADE-CERQual): Theme 1. Provision of information/ sharing knowledge

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: adequate and adapted information					
3 studies (Al-Yateem 2012, Al-Yateem 2013, Tierney 2013)	2 studies using interviews, and 1 study using focus groups	<p>Three studies conducted in Ireland and the UK with young people and young adults with cystic fibrosis and healthcare professionals reported on the need for more information throughout the process of transition.</p> <p>Adequate information</p> <p>Participants noted a lack of adequate information about the transition process, especially about the different aspects of this process. One young adult with cystic fibrosis noticed a lack of detail: <i>"Ah, very little, almost nothing. I mean all they do is to tell you that you are transferring."</i></p> <p>Another was not aware of the hospital or cystic fibrosis service to which he would be referred. He said: <i>"I don't really know, it might be either X hospital or Y hospital. I don't know we have to go and see. I don't know how they will be like?"</i></p> <p>Lack of sharing knowledge led to unmet information needed for people with cystic fibrosis:</p> <p><i>"Well the thing that they don't talk with us enough about cystic fibrosis, how it will be ... you know, and what it will be like in the future."</i> (young adult with cystic fibrosis)</p> <p>Adapted information</p> <p>Participants suggested that information should be appropriate to the person's age and developmental stage and supplemented with extra printed or digital material.</p>	Limitation of evidence	Major limitations	VERY LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>"Anyway, providing information should be given extra attention, and possibly information sessions could solve this problem" (young adult with cystic fibrosis).</i></p> <p><i>Sources of information</i></p> <p>Online forums were a good source of information for young adults about to transfer to adult services.</p> <p>As a person with cystic fibrosis stated: <i>"...since it was mentioned in the [adult] clinic about the forums...I went on there and it's a big eye opener that there's loads of people going through it, been through it, and they can just offer you a lot more advice from a patient side of it..."</i></p>			
Sub-theme 2: timely and gradual information					
1 study (Al-Yateem 2012)	1 study using interviews	<p>In 1 study conducted in Ireland with young adults with cystic fibrosis, participants stated that information should be provided at a timely interval:</p> <p><i>"Actually, we haven't been told about the transfer yet, nobody talked to us about it..."</i></p> <p><i>"Yes, I know about transition...last month the cystic fibrosis nurse told me that I will be going to another hospital next year when I turn 18..."</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Major limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	VERY LOW QUALITY

1 Table 46: Summary of clinical evidence (GRADE-CERQual): Theme 2. Transition planning

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: gradual, timely and individualised transition					
5 studies (Al-Yateem 2012, Al-Yateem 2013, Brumfield)	4 studies using interviews, and 1 study using focus groups	Five studies conducted in Australia, Ireland, the UK and the USA with young people and young adults with cystic fibrosis and healthcare professionals reported the need of careful transition planning, especially with regards to the timing	<p>Limitation of evidence</p> <p>Coherence of findings</p>	<p>Major limitations</p> <p>Coherent</p>	LOW QUALITY

<p>2004, Tierney 2013, Tuchman 2008)</p>		<p>involved, The topic of transition with cystic fibrosis to adulthood should not be a new concept.</p>	<p>Applicability of evidence</p>	<p>Applicable</p>	
		<p>Timely transition Some participants noted that the timing of transition seemed arbitrary and unfair: <i>"Actually, we haven't been told about the transfer yet, nobody talked to us about it..." (young adult with cystic fibrosis)</i> <i>"Yes, I know about transition...last month the cystic fibrosis nurse told me that I will be going to another hospital next year when I turn 18..." (young adult with cystic fibrosis)</i> <i>"And right now it's like, maybe in a year, but right now I'm, I've just gone off to college. It's like, I don't want to make that transition now. And they're like, well you need to. It's like, and you have patients here that are 30 years old and you're telling me I have to go?" (young adult with cystic fibrosis)</i> <i>"I think maybe if it wasn't just an all of a sudden strange situation. Slam the door behind you, you know. You're at university, you can't come back." (young adult with cystic fibrosis)</i></p> <p>Structured transition service Participants stated that the service provided was amorphous and appeared to lack a comprehensive and coherent structure to transitional programming: <i>"There was no real discussion ... it was just the cystic fibrosis nurse who told us last year that it would be when I finish the leaving cert. It would not be straight away when I turned 18, so it would be after." (young adult with cystic fibrosis)</i> Good paediatric care not only optimizes the health of people with cystic fibrosis, but also prepares them for a change to adulthood: <i>"Sometimes it was like he (paediatric doctor) could read your mind . . . he always eased you, whatever your problem was, if you had a problem, you always went out feeling better." (22 year old with cystic fibrosis)</i> One 18-year-old young adult with cystic fibrosis anticipating transition explained: <i>"I think it would have been easier if they</i></p>	<p>Saturation</p>	<p>Saturated</p>	

		<p>would have really started pushing when I was younger. Like, even at 15 or 16, start really suggesting it. Really being like, you know you might want to look into this. You might want to start meeting some of the doctors over there."</p> <p>Positive and supportive attitude helped in transition from children to adult health services: <i>"He (paediatric doctor) really highly recommended Dr B. (adult doctor), which was good . . . the fact that he recommended him strongly would have helped." (21 year old with cystic fibrosis)</i></p> <p>Individualised assessment and care Planning and care tailored for the individual person for transition was considered important: <i>"The assessment will highlight to us what the adolescents actually need, and what might affect his or her transition in terms of information, family, or any other issues". (young adult with cystic fibrosis)</i> <i>"It will be good for everyone [planning] adolescent, parents, and even us... everybody will know what to do" (healthcare professional).</i></p>			
Sub-theme 2: care coordinator					
1 study (Tierney 2013)	1 study using interviews	<p>In 1 study conducted in the UK with young people and young adults, some interviewees believed their needs were neglected while they waited to move and described being uncertain about who was responsible for their care because they were straddled between 2 services: <i>".. they [paediatric staff] knew I was moving up so they'd gone a bit, put me last kind of thing, so I wasn't a priority because they knew I was going to be moving up soon that they'd lost interest but they were making sure that the younger ones were alright because they knew I was going soon" (young adult with cystic fibrosis).</i></p>	<p>Limitation of evidence</p> <p>Coherence of findings</p> <p>Applicability of evidence</p> <p>Saturation</p>	<p>Minor limitations</p> <p>Coherent</p> <p>Applicable</p> <p>Not saturated</p>	LOW QUALITY
Sub-theme 3: coordination with adult services					
1 study (Tierney 2013)	1 study using interviews	<p>In 1 study conducted in the UK with young people and young adults, some participants described their feelings about moving</p>	<p>Limitation of evidence</p>	<p>Minor limitations</p>	LOW QUALITY

		<p>to adult services as superseded by procedural tasks, such as gathering relevant documents to forward to the adult team.</p> <p><i>"... didn't seem to show interest in how you felt about moving over. It was more like we've sent your notes over to that side so we're just waiting for them to reply."</i> (young adult with cystic fibrosis).</p> <p>Young adults suggested that clinicians from paediatrics could be involved to provide continuous and optimal care:</p> <p><i>"...he [paediatric consultant] could tell if I was ill. I've got like a problem with my stomach and he could feel straight away if I was having, if it was worse or if it was manageable... So he's known me for a long time. That's why I was worried about moving here...he knew a lot of the problems that I've had..."</i> (young adult with cystic fibrosis).</p>	Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	

7.3.5.31 Support after transition to adult services

2 Table 47: Summary of clinical evidence (GRADE-CERQual): Theme 1. Healthcare professionals and services configuration

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: planning and continuity in co-ordinated care					
4 studies (Brumfield 2004, Tierney 2013, Tuchman 2008, van Staa 2011)	4 studies using interviews	<p>Four studies conducted in Australia, the Netherlands, the UK and the USA with young people and young adults with cystic fibrosis, parents and healthcare professionals reported that participants felt they have lost the resources available to them in the paediatric system. It is the abruptness of the transition that was most disruptive to participants, and lack of time and resources in the adult healthcare system.</p> <p>Familiarity Participants valued having a familiar face around after transition to adult services was reassuring for people with cystic fibrosis:</p>	Limitation of evidence	Minor limitations	MODERATE QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Saturated	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>"I knew her from before . . . even though it might not be a lot of contact with her, um, it was still a familiar face, and that was enough to give me reassurance in that regard." (20 year old with cystic fibrosis)</i></p> <p>Fracturing</p> <p>However, some participants reported that transition disrupts the bond which have developed between the person with cystic fibrosis and healthcare personnel looking after them. Some participants described being ignored and their need was not addressed during transition phase.</p> <p>As a young adult with cystic fibrosis stated: <i>"... didn't seem to show interest in how you felt about moving over. It was more like we've sent your notes over to that side so we're just waiting for them to reply."</i></p> <p>Some interviewees believed their needs were neglected while they waited to move and described being uncertain about who was responsible for their care because they were straddled between 2 services:</p> <p><i>".. they [paediatric staff] knew I was moving up so they'd gone a bit, put me last kind of thing, so I wasn't a priority because they knew I was going to be moving up soon that they'd lost interest but they were making sure that the younger ones were alright because they knew I was going soon" (young adult with cystic fibrosis).</i></p> <p>One group suggested that handover should be staggered over time to make transition easier for young adults.</p> <p><i>"I think staff from the ... hospital can come over here and do their first few clinics with adolescents...this might help... and we can give them better information about our patients"</i></p> <p>Appraisal of the experience of transition</p> <p>Some parents and young adults looked back at transfer as 'no</p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>big deal</i>' and even as <i>'peanuts'</i>, when the process had been smooth or <i>'seamless'</i>.</p> <p>But most young adults and especially parents said it had been more stressful and difficult than anticipated.</p> <p>Parents and young adults said that paediatric was more warm and friendly. On the other hand, they used metaphors like <i>'being lost'</i>, <i>'falling into a deep hole'</i>, <i>'feeling abandoned'</i> and even <i>'waking up in a horror movie'</i> for adult care. However, this was seen as temporary; transition was perceived as a rite of passage: <i>"you have to get used to it, that's all."</i></p> <p>A bad transition experience created a negative impact and discouraged people in seeking care.</p> <p><i>"I always say to Mum that it's a waste of time going . . . I don't . . . I don't really trust them, because of the way they treat you, because you are a number . . . if it was up to me I probably wouldn't go back to clinic, but Mum has always told me that you've got to keep your foot in the door in case we need them (the clinic) . . . So I sort of go back on the off chance that I may need them."</i> (22 year old with cystic fibrosis)</p> <p>Another person stated that the service provided was amorphous and appeared to lack a comprehensive and coherent structure to transitional programming:</p> <p><i>"There was no real discussion . . . it was just the cystic fibrosis nurse who told us last year that it would be when I finish the leaving cert. It would not be straight away when I turned 18, so it would be after."</i></p> <p>This lack of structure created anxiety and negative feeling:</p> <p><i>"I don't really like to come here...like, they don't do much for me...if they just listen I would have told them what I want to know...or how they can help...this is very annoying..."</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 2: adaptation and integration					
3 studies (Begley 2013, Tierney 2013, Van Staa 2011)	2 studies using interviews and 1 study using a survey	<p>In 3 studies conducted in Ireland, the Netherlands and the UK with young people and young adults with cystic fibrosis, parents and healthcare professionals, participants highlighted the challenges encountered during this process:</p> <p>Not being understood Integrating with the adult services was a of the challenges for young adults and was difficult to begin with. They believed that their issues would not be easily understood in adult health services. <i>"...I was with them for so long I got to know them really well and they knew I was a fussy person. But I'm sure it will all be the same with the adults." (person with cystic fibrosis)</i></p> <p>Different culture between paediatric and adult care Parents and young adults said that paediatric staff were more warm and friendly. On the other hand, they used metaphors like <i>'being lost'</i>, <i>'falling into a deep hole'</i>, <i>'feeling abandoned'</i> and even <i>'waking up in a horror movie'</i> for adult care. However, this was seen as temporary; transition was perceived as a rite of passage: <i>"you have to get used to it, that's all."</i></p> <p>Young adults noted that more independence and self-reliance was expected of them. Parents wondered whether their children could take up the full responsibility for their treatment. <i>"The adult care had a more 'business-like approach' which often contrasted with the paediatric 'holistic, system-oriented approach."</i></p> <p>Young adults appreciated the efficiency of adult services whereas others were concerned about the shorter appointment with less information and the less frequent appointments: <i>"We used to do it (sputum test) every month in x hospital, but</i></p>	Limitation of evidence	Minor limitations	MODERATE QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Unclear	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<p><i>they just do it every 3 months in here, so you kind of stay anxious for a long time ... you don't know when the infection is going to hit ... it will be late...</i></p> <p><i>"It's just different. I mean I like how you kind of rush through it. It's not like where you have to talk to a million people. They don't seem as qualified there, it seems like the people that I talk to at Children's really knew what they were doing. Maybe it's because they don't talk to me as much."</i></p> <p>Young people with cystic fibrosis and parents were also concerned about the lack of single rooms for people with cystic fibrosis, which lead to a fear of cross-infection: <i>"Young people and their parents have real concern re cross-infection in mixed wards. They do not want to share wards with elderly patients"</i> (healthcare professional)</p> <p>Transition at an early age Healthcare professionals found problematic for young people to be transferred at an "early age". <i>"Historically the age at which children attend adult services in this hospital is 14, which I consider too young and my training as an adult physician leaves me less able to deal with the 14-16 age group. I have difficulty in managing patients who in my mind are "children" and I find teenage/ adolescent emotional aspects of illness to be an unwelcome challenge"</i> (healthcare professional)</p> <p>Involvement in decision making The main positive differences that subjects noted between paediatric and adult care were feeling greater control over and more involvement in decision making. (See overarching theme) One participant with cystic fibrosis who was very resistant to transition was impressed with the pulmonologist: <i>"She gave me this big talk about some of the new things [I'm] going to</i></p>			

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<i>encounter as an adult with cystic fibrosis. And she just opened my eyes to a lot of things."</i>			
Sub-theme 3: holistic approach					
2 studies (Al-Yateem 2012, Depuis 2011)	2 studies using interviews	<p>In 2 studies conducted in Canada and Ireland with young adults with cystic fibrosis, parents and healthcare professionals, participants suggested that the focus of some health professionals on clinical activities, rather than focusing on the actual individual needs of the young adult, was source of frustration in people with cystic fibrosis.</p> <p><i>"People need to be looked at and listened to ...and then you make your diagnosis from there, but sometimes they are over reliant on paperwork that is obsolete and doesn't give any function. It doesn't solve problems..."</i>. (young adult with cystic fibrosis)</p> <p>Healthcare professionals in adult services appeared to concern themselves mainly with giving information around clinical parameters rather than dealing with emotional factors in young adults associated with transition.</p> <p>As stated by a healthcare professional: <i>"We talk to them about their medications, about physiotherapy, the respiratory therapist sees them at every visit to go over the techniques. But other times, we show them that their X-ray has deteriorated. We show them the film, we explain that it's important for them to take control. We shake them up a bit."</i></p>	Limitation of evidence	Major limitations	VERY LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	
Sub-theme 4: evaluation and follow up mechanisms					
1 study (Al-Yateem 2013)	1 study using focus groups	<p>In 1 study conducted in Ireland with healthcare professionals, it was suggested that monitoring the process can help to improve the care during transition.</p> <p>One participant said: <i>"Based on the transition plan you can later on evaluate whether the child has made any progress, or</i></p>	Limitation of evidence	Major limitations	VERY LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of	Applicable	

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
		<i>any further intervention might be needed".</i>	evidence		
			Saturation	Not saturated	

1 Table 48: Summary of clinical evidence (GRADE-CERQual): Theme 2. Psychosocial support

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Sub-theme 1: family support					
1 study (Tierney 2013)	1 study using interviews	In 1 study conducted in the UK with young people and young adults with cystic fibrosis, the presence of parents was described as reassuring, especially during the early transition phase: <i>"...if my mum hadn't come, I wouldn't have asked half as many questions. I don't think I'd have been as open with the doctors talking to me...I think I'd have been a little bit more introvert and worried...I know it sounds dead daft, me mum was there and I'm 19 but because me mum was there I was more confident in asking questions because I knew if I'd said something that had come out a funny way or the wrong way, mum would go well what she actually means is this" (young person with cystic fibrosis).</i>	Limitation of evidence	Minor limitations	LOW QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	
Sub-theme 2: support network					
1 study (Brumfield 2004)	1 study using interviews	In 1 study conducted in Australia with young adults with cystic fibrosis, a support network was considered important for transition. Participants noted they are anxious and the lack of support network does not help in fully utilising the services: <i>"I go to clinic, and I go in and I come out as quickly as I can." (22 year old with cystic fibrosis)</i>	Limitation of evidence	Minor limitations	VERY LOW QUALITY
			Coherence of findings	Unclear	
			Applicability of evidence	Unclear	
			Saturation	Not saturated	
Sub-theme 3: emotional support					
2 studies (AI-	2 studies using	Two studies conducted in Ireland and the UK with young adults	Limitation of	Major	VERY LOW

Study information		Description of theme or finding	Quality assessment		
Number of studies	Design		Criteria	Rating	Overall
Yateem 2012, Tierney 2013)	interviews	<p>with cystic fibrosis reported on the lack of understanding of mental state and their needs during the transition period.</p> <p>Transition from children to adult healthcare setting created anxiety in people with cystic fibrosis: <i>"I am worried about the cutbacks in health and all when we move there (the new hospital), you know that with cystic fibrosis there are some cuts that you don't know when it's going to knock off..." (young person with cystic fibrosis).</i></p> <p>Some participants who had transferred very recently felt ill-equipped emotionally to cope with this change. <i>"...the doctor did ask me "what do you want to do?"...in children's they wouldn't do that, they'd just say 'you need IVs, you should come in.' So it was quite a bit, it was a bit confusing because I didn't really know what to do myself. I was like, I don't know, you tell me what should I do" (young person with cystic fibrosis).</i></p>	evidence	limitations	QUALITY
			Coherence of findings	Coherent	
			Applicability of evidence	Applicable	
			Saturation	Not saturated	

1

2

3

1 **7.3.6 Economic evidence**

2 No economic evaluations related to the transition in cystic fibrosis were identified in the
3 literature search conducted for this guideline. Full details of the search and economic article
4 selection flow chart can be found in Appendix E and F, respectively.

5 This review question is not relevant for economic analysis because it does not involve a
6 decision between alternative courses of action.

7 **7.3.7 Evidence statements**

8 **7.3.7.1 Overarching themes**

9 **Transition as natural stage**

10 Low quality evidence from 3 qualitative studies conducted with young people and young
11 adults with cystic fibrosis, parents and healthcare professionals, noted that transition to adult
12 services was seen as a natural and inevitable process. Most people with cystic fibrosis
13 reported that as they grew up, healthcare professionals started to treat them as adults, and
14 this was perceived as positive.

15 **Parallel changes and planning**

16 Low quality evidence from 2 qualitative studies conducted with young people and young
17 adults with cystic fibrosis and parents highlighted that relocating to a new healthcare system
18 is a of several transitions young people with a long-term condition must face (for example
19 going to university). Hence, transfer takes place against a backdrop of additional pressures
20 and might not always be a top priority for the person. Young people and their parents are
21 concerned and sad about the future, and the impact the illness will have in their personal life.

22 **Independency in decision making**

23 Moderate quality evidence from 5 qualitative studies with young people and young adults
24 with cystic fibrosis, parents and healthcare professionals reported on the importance of being
25 allowed to make choices, self-advocacy and build independence gradually. Young people
26 and healthcare professionals recognised that young people should become more involved in
27 decision making as they become adults. Many young participants actively led this process,
28 and welcomed the opportunity to lead discussions with healthcare professionals. In some
29 instances, young people do not want parents to be involved in consultations or decision
30 making anymore. Parents, however, wanted to be recognised as an integral part of the
31 young person's care, and found it difficult to accept the new family dynamic.

32 **7.3.7.2 Support before transition to adult services**

33 **Provision of information or sharing of knowledge**

34 Very low quality evidence from 3 qualitative studies conducted with young people and young
35 adults with cystic fibrosis and healthcare professionals, reported on the information needs
36 during transition. Participants noted the need for information in relation to different aspects of
37 the process, such as the new place of care, and services structure. They also noted the
38 information has to be adapted to the age and developmental stage of the person, and it
39 should be provided at timely intervals.

1 **Transition planning**

2 Low quality evidence from 5 qualitative studies conducted with young people and young
3 adults with cystic fibrosis and healthcare professionals reported on the need of careful
4 transition planning, especially with regards to the timing involved. Participants felt transition
5 should happen at the right time, not just because of arbitrary criteria, such as the person
6 turning 18. They also highlighted the process should be clearly structured, with anticipated
7 planning and coordination between paediatric and adult care. They noted the importance of
8 having a care coordinator during the process, which helps the person navigating between
9 both services. Ultimately, planning and care tailored for the individual person for transition
10 was considered important.

11 **7.3.7.3 Support after transition to adult services**

12 **Healthcare professionals and services configuration**

13 Very low to moderate quality from 6 qualitative studies conducted with young people and
14 young adults with cystic fibrosis, parents and healthcare professionals discussed the
15 challenges faced during the transition process. Participants noted it would be useful to have
16 a familiar person in the new setting; instead, they had to face disruption in the care received
17 during the transition from paediatric to adult care. This made people with cystic fibrosis and
18 parents feel lost, abandoned and anxious. They attributed this to the different care culture.
19 These negative experiences had a negative impact in the long term for some people, but
20 most participants agreed they finally were able to adapt to the new situation. Some
21 participants, on the contrary, highlighted there were benefits, such as feeling greater control
22 over and more involvement in decision making. In addition, people with cystic fibrosis and
23 their parents noted healthcare professionals should not only focus on clinical information, but
24 also on the emotional aspects related to transition.

25 Healthcare professionals suggested that monitoring the process can help improve the
26 experience of transition to adult services.

27 **Psychosocial support**

28 Very low to low quality evidence from 4 qualitative studies conducted with young people and
29 young adults with cystic fibrosis, participants highlighted the importance of having
30 psychosocial support. Family and social support networks were seen as useful and were
31 described as reassuring. They also noted the importance of understanding the mental state
32 and the emotional needs of the people during this process.

33 **7.3.7.4 Economic evidence statements**

34 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

35 **7.3.8 Evidence to recommendations**

36 **7.3.8.1 Relative value placed on the outcomes considered**

37 The aim of this review was to identify elements of the transition process (for example,
38 transition planning involvement) from paediatric to adult services from perspectives of young
39 people with cystic fibrosis and their family and carers.

40 Although themes were mainly identified from the literature, the Committee identified some
41 expected themes that they thought would be important during protocol stage. They agreed
42 that the following themes would provide useful perspectives: transition plan, transition lead,
43 preparation period, involvement of young people and their family or carers in planning and

1 implementation, clarity about the process, coordination between paediatric and adult
2 services, timing of transition and availability of information.

3 **7.3.8.2 Consideration of clinical benefits and harms**

4 The Committee acknowledged the evidence, and they thought that the themes and their sub-
5 level themes which emerged or were derived from the evidence were useful and relevant,
6 and that the evidence matched their clinical observations. The Committee also
7 acknowledged the recommendations by the CF Trust and the NHS service specifications for
8 cystic fibrosis.

9 The Committee highlighted transition is a continual and stepwise process that does not stop
10 when the young person is transferred to adult services. With regards to the age, they agreed
11 the transition process should start at the age of 12 or 14 and they should have moved to
12 adult services before they reach 18 years. This is consistent with the recommendations
13 made by the NHS service specifications for cystic fibrosis and the CF Trust (CF Trust
14 Standards of Care 2011). They agreed paediatric services are not adequate for young
15 people over 16. However some flexibility should be allowed, for example if the young person
16 is very unwell or approaching end of life.

17 As raised in the literature, the Committed discussed about young people going through a
18 number of transitions, for example going to university, when additional support may be
19 needed. They also agreed it is important to respect the individual needs and to note that
20 young people mature at different rates.

21 When reviewing the evidence, the Committee discussed the role of parents. As identified in
22 the literature, the role of parents' changes during transition. It is very important to work with
23 both the young person and the parents for transition to go smooth. They agreed there is a
24 lack of recognition for parents in adult services. It was noted, for example, that in some cases
25 parents are not allowed in adult consultations and they felt parents may be especially needed
26 during the first visits. They agreed it is important to respect the family dynamics and to ask
27 parents how they want to be involved in their child's care. However, priority should always be
28 given to the young person when deciding who should be involved in decision making.

29 With regards to the healthcare professionals that should be involved, the Committee
30 underlined that transition should take account of the whole multidisciplinary team, and not
31 just the cystic fibrosis paediatricians. Paediatricians should play an important role in
32 providing information that it is adequate to the young person's needs. They also agreed it is
33 important to have a documented coordinated pathway to handover the information from
34 paediatric to adult services. As noted in the studies, some young people are concerned
35 about the differences between paediatric and adult services. For example, some young
36 people may be taken off prophylactic antibiotics after transition as it is not common practice
37 in adult treatment.

38 As highlighted in the evidence, the Committee noted the coordination between child and
39 adult services is challenging. Consistent with the recommendation in the NHS service
40 specifications for cystic fibrosis, they agreed a main specialist cystic fibrosis centre should
41 take responsibility for coordinating with all the services for transition, even for people
42 receiving shared care. This is to ensure that transition for specific conditions (for example
43 liver disease), happens concurrently.

44 Moreover, as noted in the evidence, the Committee thought it could be useful to have a
45 named person for point of contact to coordinate and facilitate transition. This is common
46 already practice, and it is consistent with the NHS Service Specifications for cystic fibrosis. In
47 current practice, nurses are mostly coordinating and taking the lead, but they agreed it could
48 be done by other members of multidisciplinary team.

1 Finally, the Committee agreed it is important to seek continuous feed-back to evaluate the
2 transition process. This was also highlighted in the evidence, as healthcare professionals
3 working with people with cystic fibrosis suggested that monitoring the transition process
4 could help to improve care during transition. In addition, assessing the progress on the
5 transition process allows to identify if progress is going according to plan, or if any further
6 interventions are needed.

7 **7.3.8.3 Consideration of economic benefits and harms**

8 This review question was not relevant for economic analysis because it does not involve a
9 decision between alternative courses of action. Even so, there may be resource implications
10 arising from the provision of any services that facilitate the transition from paediatric to adult
11 services.

12 The Committee stated that identifying an accountable healthcare professional to support a
13 person's transition is often considered as ideal practice by clinics, and consequently not in
14 place by all centres. Following this, the Committee did not think significant additional
15 expenditure was necessary to implement this across all centres. This was on the basis that
16 the named worker role comprises a set of tasks to be done by an existing worker, rather than
17 the creation of a new post. Furthermore, it is likely that the worker allocated these tasks will
18 be undertaking many of them already as there is a continual flow of transitioning young
19 people at every clinic. Instead, the Committee advised that the recommendation will
20 formalise the process and lead to more timely and accurate records to promote a co-
21 ordinated, transparent and cost-effective transition.

22 Following this, the Committee noted that existing guides on the process to prepare for
23 transition such as "Ready Steady Go" developed by Southampton Children's Hospital, can
24 be adapted for people with cystic fibrosis, providing a cost-effective solution to developing de
25 novo guides.

26 The Committee stated that paediatric and adult clinics see many transitions throughout the
27 year and may continue to use established methods to conduct the transition, without
28 reviewing the pathways they follow and considering the opportunity cost of resources a
29 transition requires. As a result, the Committee prioritised a recommendation to review
30 transition pathways, as pathways could be improved by de-duplicating work, or removing
31 unnecessary, or inefficient aspects and putting those resources into aspects that increase
32 satisfaction and care.

33 **7.3.8.4 Quality of evidence**

34 Very low to moderate quality evidence was presented in this review as assessed by GRADE-
35 CERqual. The main reasons leading to downgrading the quality of the evidence included:

- 36 • Many studies did not clearly report how they selected the participants, and a few studies
37 reported a low response rate from participants. The inclusion and exclusion criteria was
38 not reported in most studies nor were the characteristics of the population. In most
39 studies, the relationship between researchers, interviewers and participants was not
40 described.
- 41 • Data collection relied on interviews in most studies, but the process was not always
42 adequately described. Few studies included the list of questions used during the interview.
43 Data saturation was generally not discussed.
- 44 • Most studies conducted thematic analysis, although the process was not always
45 described in detail. For example there were no discussions about validity or robustness of
46 data. The role of the researchers in the analysis was not described. In general themes
47 were supported by quotes, but most studies did not indicate whether saturation was
48 achieved. When considering the evidence as a whole it did not appear very saturated, as
49 many themes were identified in a study and there were few quotes to support them.

1 **7.3.8.5 Other considerations**

2 The Committee agreed the NICE guideline on transition from children's to adult's care
3 (NG43) provides useful guidance. However, they noted that are some aspects that are
4 unique for young people with cystic fibrosis, their families and carers. Therefore the
5 recommendations related to this evidence review were based on the evidence and the
6 Committee's clinical experience. In addition, an overarching recommendation pointed to the
7 NICE guideline on transition from children's to adults' services for young people using health
8 or social care services (NICE NG43).

9 The Committee discussed potential equality issues. They noted that young people who live
10 far from a specialist centre may be disadvantaged. However they agreed no additional
11 recommendations were needed, as the use of alternative models care had already
12 extensively been discussed in the service delivery review. See section 7.1.

13 The Committee discussed whether a research recommendation was needed for this topic.
14 They agreed the only research recommendation would be to develop a cystic fibrosis specific
15 transition guideline or pathway for implementation across all centres but this is deemed an
16 unnecessary use of resources given the fact that funding is finite and there exists a NICE
17 transition guideline recently developed that would apply to cystic fibrosis with minor
18 adaptation.

19 **7.3.8.6 Key conclusions**

20 The Committee concluded that the transition from paediatric to adult services is an ongoing,
21 stepwise process that should start when the young person is 12-14 years and continue
22 beyond the age of 20. Unless contraindicated, transition to adult services should happen
23 between the age of 16 and 18 years old. Having a coordinated transition pathway is a useful
24 tool to achieve a successful transition.

25 **7.3.9 Recommendations**

26 **31. Begin the transition process to adult services with young people with cystic**
27 **fibrosis when they are 12 years old, and with their family members or carers (as**
28 **appropriate).**

29 **32. Ensure that young people with cystic fibrosis move to adults' services between**
30 **the ages of 16 and 18.**

31 **33. All cystic fibrosis services should have a coordinated and documented pathway**
32 **for transition from children's to adults' services that includes plans for managing**
33 **all cystic-fibrosis-related aspects of care.**

34 **34. Provide a named worker to lead the transition for each young person with cystic**
35 **fibrosis who is moving to adults' services, and if possible ensure that the named**
36 **worker is someone the young person already knows. For more guidance on**
37 **named workers see the section on named workers in the NICE guideline on**
38 **transition for young people using health or social care services.**

39 **35. The specialist cystic fibrosis centre should take responsibility for ensuring**
40 **transitions are successful.**

41 **36. Ask people with cystic fibrosis and their family members or carers (as**
42 **appropriate) for feedback on the quality of the transition service.**

- 1 **37. For more guidance on managing transition from children’s to adults’ services, see**
2 **the NICE guideline on transition for young people using health or social care**
3 **services.**
4

8 Complications of cystic fibrosis

Review question: What are the non-lower-respiratory complications of cystic fibrosis in infants, children, young people and adults?

8.1 Introduction

Cystic fibrosis is a multifaceted genetic disease with a wide range of complications which depend upon genotype, environment and disease progression. Traditionally progressive lung disease due to chronic infection and the impairment of pancreatic function have typified cystic fibrosis, resulting in premature death. However, the success of new-born screening programmes allows earlier intervention, slowing disease progression and altering the clinical manifestation of the condition. Also, with increasing longevity come unexpected complications due to the side-effects of treatment regimens, as well as those associated with older age.

Thus, a range of other co-morbidities are becoming increasingly apparent, and the traditional view of cystic fibrosis is increasingly disfavoured as other organ systems are involved and have a significant effect on patient health. This review question aims to determine the key health problems associated with a diagnosis of cystic fibrosis, and focuses on non-lower-respiratory complications since lower airway disease is addressed in detail in the rest of the guideline.

The results are presented separately for each complication.

See study selection flow chart in Appendix F, study evidence tables in Appendix G, and list of excluded studies in Appendix H.

8.2 Malnutrition

8.2.1 Description of clinical evidence: malnutrition

The aim of this review was to determine the prevalence of malnutrition among adults with cystic fibrosis, and the prevalence of vitamin deficiency among people of all ages. This section includes studies on adult or mixed populations, or studies on vitamin deficiency with people of all ages. Please see following section on impaired growth for studies focusing on children and young people.

We requested data to the UK CF registry on the percentage of adults with body mass index (BMI) under or over the cut-offs for normal nutritional status as outlined in the consensus document on nutritional management of cystic fibrosis published by the CF Trust in 2016 (CF Trust 2016). Data from the registry should be prioritized according to the protocol. We also looked for observational studies because they would provide data on indicators not included in the registry. We aimed to prioritize prospective cohort studies, but only included a prospective cross-sectional study and 4 retrospective studies. We prioritized studies from more relevant contexts (the UK in the first instance, then countries in Western Europe, Australia or North America). We also prioritized studies with more recent data for each indicator of malnutrition. Moreover, we prioritized studies that disaggregated data between infants, children, young people and adults; as well as studies that reported prevalence at different points in time and studies that were based on registries.

For full details see review protocol in Appendix D.

We included data obtained from the UK CF registry for the year 2015. This data referred to 5701 people with cystic fibrosis aged >16 years. In addition to these data, we included 3 studies on malnutrition defined by indicators such as BMI, height percentile or BMI

percentile. One study (Moen 2011) was a prospective multicentre study conducted in Denmark, Norway and Sweden from 2003 to 2006. This study included 347 adults with cystic fibrosis. One study (Stephenson 2013) used data from the Toronto CF registry for 3 time intervals: 1985-1990, 1991-1999, and 2000-2011. This study included 909 people with cystic fibrosis attending an adult clinic. 1 study (Vieni 2013) used data from 2 centres in Italy from 2007. This study included 393 people with cystic fibrosis aged >6 years.

We included 2 retrospective studies on vitamin deficiency. One study (Chavasse 2004) used data from a CF clinic in the UK for 1999-2001. This study included 290 people with cystic fibrosis aged 1-18. One study (Rana 2014) used data from 3 paediatric centres in Australia for 2007-2010. This study included 530 people with cystic fibrosis aged ≤18 years.

8.2.2 Summary of included studies and results: malnutrition

A summary of the studies that were included in this review are presented in Table 49.

Table 49: Summary of included studies

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Data request to UK CF registry	UK CF registry	2015	5701 people with cystic fibrosis aged >16 years	Age group: >16 BMI < 20 kg/m ² : 24.5% (1398/5701) BMI > 25 kg/m ² : 22.2% (1266/5701)	Cystic fibrosis care teams at every specialist centre and clinic across the UK entered data on weight, height and BMI for the registry. The following cut-offs were used: Requires additional nutritional support: <ul style="list-style-type: none"> • BMI <20 kg/m² • Overweight: • >25 kg/m² 	Moderate <ul style="list-style-type: none"> • Details on the people in the registry and on how data is collected for the registry are available in the UK Cystic Fibrosis Trust Registry 2015 Annual Data Report and on the CF Trust website. However it is unclear from these sources if weight and height were measured using objective, standard criteria or reliably.
Moen	Multicen	Septe	347	BMI<19.0 kg/m ² :	• Thresholds:	High

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
2011 Scandina- navia (Denm ark, Norwa y and Swede n) Prospe ctive cross- section al study	Register study - 7 centres	November 2003- May 2006	adults with cystic fibrosis Age: ≥ 18 years	18% (62/347) BMI<18.5 kg/m2: 13% (44/347)	<ul style="list-style-type: none"> • BMI < 19.0 kg/m2 • BMI < 18.5 kg/m2 • Measurements: • Weight was measured in the morning wearing undergarments. • Height was measured with no stockings or shoes and the means of 3 measurements were recorded. 	
Stephe nson 2013 Canad a Retros pective study	Toronto CF registry	1985- 1990; 1991- 1999; 2000- 2011	909 people with cystic fibrosis attending the Adult CF Clinic in Toronto Age not reported	<p>Nutritional status of people attending the clinic between 2000 and 2011 (n=651), using their last recorded measurement:</p> <ul style="list-style-type: none"> • Underweight: 17% • Adequate weight: 60% • Overweight: 18% • Obese: 3.8% <p>Prevalence calculated using 1000 random measurements per time interval:</p> <p>Underweight</p> <ul style="list-style-type: none"> • 1985-1990: 20.6% • 1991-1999: 11.6% • 2000-2011: 11.1% <p>Overweight:</p> <ul style="list-style-type: none"> • 1985-1990: 7.0% • 1991-1999: 15.8% • 2000-2011: 18.4% 	<ul style="list-style-type: none"> • Underweight status was defined as BMI<18.5 kg/m2 • Adequate weight was defined as 18.5-24.9 kg/m2 • Overweight was defined as BMI 25-29.9 kg/m2 • Obese was defined as BMI ≥30 kg/m2 <p>Definitions as per WHO BMI guidelines.</p>	<p>Low</p> <ul style="list-style-type: none"> • Age not reported for each cohort • No numerators provided
Vieni 2013	Regiona l Centre	Decem ber	393 people	Height percentile <5th: 24.4%	Height percentile <5th	<p>Low</p> <ul style="list-style-type: none"> • Data are

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Italy Retrospective study	in Palermo and Satellite Centre in Messina, Italy	2007	with cystic fibrosis Age: >6 years	BMI percentile <10th (paediatric) or BMI <18.5 kg/m ² (adults): 35.3%	BMI percentile <10th (paediatric) or BMI <18.5 kg/m ² (adults)	not disaggregated by age subgroup • No numerators provided
Vitamin deficiency						
Chavasse 2004 UK Retrospective study	Data from specialist paediatric CF clinic	August 1999-April 2001	290 people with cystic fibrosis Age: 1 to 18 years	25-hydroxyvitamin D (25-OHD) • < 15 nmol/l: 1% (4/290) • < 25 nmol/l: 6% (17/290)	25-OHD was measured by an in-house, competitive protein-binding assay following extraction and chromatography of 25-OHD on silicic acid, performed at Charing Cross Hospital.	Moderate • Data not disaggregated between infants, children, young people and adults
Rana 2014 Australia Retrospective audit	Data from 3 paediatric CF Centres	2007-2010	530 people with cystic fibrosis Age: ≤18 years • 301/470 took fat soluble-vitamin supplementation	Vitamin A deficiency: • 2007: 11.17% • 2010: 13.13% Vitamin D deficiency: • 2007: 22.11% • 2010: 20.22% Vitamin E deficiency: • 2007: 15.54% • 2010: 13.89% Deficiency of one or more fat-soluble vitamins on their first vitamin level test: 45% (240/530) Vitamin A levels: • Abnormal: 23.4% (123/526) * • Low: 15% (80/526) 25-OHD vitamin levels: • Abnormal: 19.8% (65/328) • Low: 19% (63/328) Vitamin E levels:	Vitamins A and E levels were performed using protein precipitation with high-performance liquid chromatography (HPLC) and ultraviolet detection (in-house method). Vitamins A and E reference ranges varied in each laboratory due to historical reasons. The reference range at CHW was vitamin A (0.8–2.5 mmol/L), vitamin E (12–36 mmol/L); at JHCH, vitamin A (1.05–2.50 mmol/L), vitamin E (8–30 mmol/L). The reference range for vitamins A and E at SCH varied based on	Very low • Numerators and denominators not provided for vitamin deficiency • Data not disaggregated between infants, children, young people and adults • Unclear why prevalence of low vitamin levels at first vitamin test is different from prevalence of vitamin deficiency in 2007 • The range

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
				<ul style="list-style-type: none"> Abnormal: 38.4% (201/523) Low: 20% (105/523) 	the child's age. 25-OHD was measured by radioimmunoassay (Diasorin, Stillwater, MN, USA). Deficiency of 25-OHD was defined as <50 nmol/L.	for normal levels of vitamins differs between the laboratories taking part in the study.

1 Abbreviations: BMI: Body Mass Index; CF: cystic fibrosis; HPLC: high-performance liquid chromatography; kg/m²:
2 kilograms per square metre; mmol/l: millimoles per litre; WHO: World Health Organization; 25-OHD: 25-
3 hydroxyvitamin D
4 * Percentage calculated by NGA technical team

5 8.2.3 Evidence statements: malnutrition

6 Moderate quality evidence from the UK CF Trust registry with 5701 people with cystic fibrosis
7 aged >16 found that the prevalence of those who required additional nutritional support
8 (defined as BMI < 20 kg/m²) was 24.5% and the prevalence of overweight (defined as BMI >
9 25 kg/m²) was 22.2% in 2015.

10 High quality evidence from 1 study with 347 adults with cystic fibrosis attending 7 centres in
11 Denmark, Norway and Sweden found that the prevalence of malnutrition defined as
12 BMI<19.0 kg/m² was 18% and the prevalence of malnutrition defined as BMI <18.5 kg/m²
13 was 13% between 2003 and 2006.

14 Low quality evidence from 1 study with 909 people with cystic fibrosis attending an adult
15 clinic in Toronto found that the prevalence of underweight (defined as BMI<18.5 kg/m²) was
16 17%, the prevalence of overweight (defined as BMI 25-29.9 kg/m²) was 18% and the
17 prevalence of obesity (defined as BMI≥30 kg/m²) was 3.8%, using the last recorded
18 measurement for each person between 2000 and 2011. Using 1000 random measurements
19 per time interval, the prevalence of underweight was 20.6% in 1985-1990, 11.6% in 1991-
20 1999, 11.1% in 2000-2011 and the prevalence of overweight was 7.0% in 1985-1990, 15.8%
21 in 1991-1999, and 18.4% in 2000-2011.

22 Low quality evidence from 1 study with 393 people with cystic fibrosis aged >6 years from 2
23 centres in Italy found that the prevalence of height percentile <5th was 24.4%, and the
24 prevalence of either BMI percentile <10th or BMI <18.5 kg/m² (depending on the age) was
25 35.3% in 2007.

26 Moderate quality evidence from 1 study with 290 people with cystic fibrosis aged 1-18
27 attending a clinic in the UK found that the prevalence of 25-hydroxyvitamin D (25-OHD)< 15
28 nmol/l was 1% and the prevalence of 25-OHD < 25 nmol/l was 6% between 1999 and 2001.

29 Very low quality from 1 study with 530 people with cystic fibrosis aged ≤18 years attending 3
30 paediatric centres in Australia found that the prevalence of vitamin A deficiency was 11.17%
31 in 2007 and 13.13% in 2010, the prevalence of vitamin D deficiency was 22.11% in 2007 and
32 20.22% in 2010, the prevalence of vitamin E deficiency was 15.54% in 2007 and 13.89% in
33 2010. Prevalence of deficiency of 1 or more fat-soluble vitamins was 45% on first vitamin
34 level test. Prevalence of abnormal vitamin A levels was 23.4% and of low vitamin A levels
35 was 15% on first vitamin level test. Prevalence of abnormal vitamin 25-OHD levels was
36 19.8% and of low vitamin 25-OHD levels was 19% on first vitamin level test. Prevalence of

1 abnormal vitamin E levels was 38.4% (201/523) and of low vitamin E levels was 20%
2 (105/523) on first vitamin level test.

3 8.2.3.1 Economic evidence statements

4 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

5 8.3 Impaired growth

6 8.3.1 Description of clinical evidence: impaired growth

7 The aim of this review was to determine the prevalence of impaired growth among infants,
8 children and young people with cystic fibrosis.

9 We requested data to the UK CF registry on the percentage of children and young people
10 with BMI percentile below the cut-off for additional nutritional support or above the cut-off for
11 overweight as outlined in the consensus document on nutritional management of cystic
12 fibrosis published by the CF Trust in 2016. Data from the registry should be prioritized
13 according to the protocol. We also looked for observational studies because they would
14 provide data on indicators not included in the registry. We aimed to prioritize prospective
15 cohort studies, but only included 1 prospective cross-sectional study and 4 retrospective
16 studies. We prioritized studies from more relevant contexts (the UK in the first instance, then
17 countries in Western Europe, Australia or North America). We also prioritized studies with
18 more recent data for each indicator of impaired growth. Moreover, we prioritized studies that
19 disaggregated data between infants, children, young people and adults; as well as studies
20 that reported prevalence at different points in time and studies that were based on registries.

21 For full details see review protocol in Appendix D.

22 We included data obtained from the UK CF registry for the year 2015. This data referred to
23 3383 people with cystic fibrosis aged 2 to 16 years. In addition to these data, we included 5
24 studies on impaired growth. Older studies were included because they provided data on
25 indicators or age sub-groups not covered by more recent studies. Four studies used data
26 from the CF Foundation National Registry in the United States. One study (Heltshe 2014)
27 used data for 2004-2009. This study included 1992 infants and children with cystic fibrosis
28 aged 0-24 months. One study (Zhang 2013) used data from 1994-2008 to calculate pubertal
29 peak height velocity. This study included 1862 people with cystic fibrosis aged 10 to 18 years
30 old. One study (Lai 2008) used data from 2002. This study included 14702 people with cystic
31 fibrosis aged less than 20 years old. One study (Zhang 2010) used data from the CF
32 Foundation Patient Registry in the United States from 1986 to 2005. This study included
33 3306 people with cystic fibrosis aged 2-18.5. This study provided the most recent data
34 relating to Himes adjusted percentiles. One study (Lucidi 2009) was a prospective
35 multicentre study with 10 Italian CF centres from 2005 to 2006. It included 892 people with
36 cystic fibrosis aged 1 month to 18 years.

37 8.3.2 Summary of included studies and results: impaired growth

38 A summary of the studies that were included in this review are presented in Table 50.

39 **Table 50: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Data	UK CF	2015	3383	• Age 2-11:	Cystic fibrosis	Moderate

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
request to UK CF registry	registry		people with cystic fibrosis aged 2 to 16 years	<ul style="list-style-type: none"> • BMIp < 25th: 17.3% (432/2498) • BMIp>91st: 10.1% (253/2498) • Age 12-16: • BMIp < 25th: 27.5% (243/885) • BMIp>91st: 5.9% (52/885) Age 2-16: <ul style="list-style-type: none"> • BMIp < 25th: 20.0% (675/3383) • BMIp>91st: 9.0% (305/3383) 	care teams at every specialist centre and clinic across the UK entered data on weight, height and BMI for the registry. The following cut-offs were used: Requires additional nutritional support: <ul style="list-style-type: none"> • <25th BMIp • Overweight: • > 91st BMIp 	<ul style="list-style-type: none"> • Details on the people in the registry and on how data is collected for the registry are available in the UK Cystic Fibrosis Trust Registry 2015 Annual Data Report and on the CF Trust website. However it is unclear from these sources if weight and height were measured using objective, standard criteria or reliably.
Heltsh 2014 United States Retrospective study	US CF Foundation National Registry	January 2004-December 2009	1992 infants and children with cystic fibrosis Age: 0 to 24 months	Age 12 months (N=374): <ul style="list-style-type: none"> • weight for age <10th: 11% • weight for age <5th: 8.3% • weight for age <2.5th: 3.7% • weight velocity (GUO-US) <50th: 48.1% • weight velocity (WHO)<50th: 30.5% • weight velocity <10th (WHO): 6.4% • weight velocity<5th 	Thresholds: <ul style="list-style-type: none"> • WHO standardized 2.5th, 5th, and 10th percentiles for weight and length for age • Guo et. al. US (Guo-US) 50th percentile for weight velocity and length velocity (as recommended by the CFF infant care guidelines) • WHO standardized 2.5th, 5th, 10th, 	Moderate. <ul style="list-style-type: none"> • Numerators are not provided

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
				<p>(WHO): 4.5%</p> <ul style="list-style-type: none"> • weight velocity < 2.5th (WHO): 3.5% • length for age < 10th: 26.8% • length for age < 5th: 17.7% • length for age < 2.5th: 10.4% • length velocity (GUO-US) < 50th: 45.7% • length velocity < 50th (WHO): 38.4% • length velocity < 10th (WHO): 20.7% • length velocity < 5th (WHO): 13.4% • length velocity < 2.5th (WHO): 13.4% <p>Age 24 months (N=317):</p> <ul style="list-style-type: none"> • weight for age < 10th: 6.9% • weight for age < 5th: 2.8% • weight for age < 2.5th: 1.9% • weight velocity (GUO-US) < 50th: 59.3% • weight velocity (WHO) < 50th: 51.1% • weight velocity < 10th (WHO): 18.3% • weight velocity < 5th (WHO): 12.6% • weight velocity < 2.5th (WHO): 8.2% • length for age < 10th: 24.9% • length for age 	<p>and 50th percentiles for weight and length velocity</p>	

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
				<p><5th: 17%</p> <ul style="list-style-type: none"> length for age <2.5th: 11.4% length velocity (GUO-US) <50th: 57.4% length velocity <50th (WHO): 58.7% length velocity <10th (WHO): 30.3% length velocity<5th (WHO): 24.3% length velocity<2.5th (WHO): 19.9% 		
Lai 2008 United States Retrospective cross-sectional study	US CF Foundation Patient Registry	2002	14702 people with cystic fibrosis Age: <20 years	2002 CFF definition of underweight (Height percentile<5th or BMIp<10th or % Ideal Body Weight<90): 33.0% Corrected classification of underweight ((Height percentile <5th or BMIp<10th): 26.8% BMIp<50th: 56.8%	2002 CFF definition: <ul style="list-style-type: none"> Age <2 y old: Height percentile <5th, % Ideal Body Weight <90%, or weight for height percentile <10th. Ages 2–20 y: Height percentile <5th, % Ideal Body Weight <90%, or BMIp <10th Corrected classification: elimination of % Ideal Body Weight <90% as an indicator of underweight Below BMI goal: <ul style="list-style-type: none"> Age<2 y old: Weight for height percentile <50th. Ages 2-20 y old: BMIp <50th. 	Very low <ul style="list-style-type: none"> Unclear why different definitions for infants were not taken into account in the results Unclear how many infants, children, young people and adults Results not disaggregated by age subgroup Unclear if weight and height were measured consistently
Lucidi 2009	Multicentre study – 10	January 2005-	892 people with cystic	0-2 years (Height for age percentile: n=104; Weight for	Nutritional failure was defined as Height for age	Moderate: <ul style="list-style-type: none"> No

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Italy Prospective cross-sectional study	Italian CF centres	December 2006	fibrosis Age range: 1 month to 18 years Mean age: 9.2	length percentile: n=101): Height for age percentile <5th: 15.4% Height for age percentile 5th-25th: 18.3% Weight for length percentile <10th: 12.9% Weight for length percentile 10th - 25th: 22.7% 2-18 years (n=788): Height for age percentile <5th: 11.8% Height for age percentile 5th-25th: 29.3% BMIp <15th: 20.9% BMIp 15th-25th: 9.6% BMIp <50th: 54.4% 10-14 years (n=179): Height for age percentile <5th: 11.7% BMIp <15th: 20.1% 14-18 years (n=183): Height for age percentile <5th: 21.9% BMIp <15th: 27.9% 0-18 years (n=892): Height for age percentile <5th: 12.2% Height for age percentile 5th-25th: 28%	percentile <5th (all ages), Weight for length percentile <10th (<2 years), BMIp <15th (2-18 years) Risk of malnutrition: Height for age percentile 5th-25th (all ages), Weight for length percentile 10th-25th (<2 years), BMIp 15th-25th (2-18 years) Height and weight were measured (when the patient was in a stable clinical condition) by specifically trained personnel. Reproducibility in anthropometric measurement was evaluated by comparing measures obtained with standard instruments in all centres with those obtained with reference instruments in a sample of patients.	numerators • Some age subgroups included less than 250 people
Zhang 2010 United States Retrospective study	CFF Patient Registry	1986-2005	3306 people with cystic fibrosis Age: 2 to 18.5 years	<ul style="list-style-type: none"> • Unadjusted height percentile <5th: 16% • Unadjusted height percentile <10th: 26% • Himes adjusted 	Procedure to calculate CFF target height and range (lower bound method): 1. Average 2 parental heights	Moderate • Self-reported parental heights

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
				<p>height percentile <5th: 18%</p> <ul style="list-style-type: none"> • Himes adjusted height percentile: <10th: 31% • CFF lower bound method: 24% 	<p>to obtain mid-parental height. Calculate the child's target adult height by adding 6.5 cm to mid-parental height for a boy, or subtracting 6.5 cm for a girl. Apply ± 10 cm for a boy or ± 9 cm for a girl to define the target height range.</p> <p>2. Plot target height and range at age 20 years on the 2000 CDC growth chart and estimate their respective percentiles.</p> <p>3. Extrapolate the percentiles of target height and range at age 20 to the child's current age.</p> <p>4. Plot the child's height on the 2000 CDC growth chart; if his/her height percentile is below the target height lower bound, he/she is considered to be below genetic potential.</p> <p>Procedure to calculate Himes adjusted height</p> <ol style="list-style-type: none"> 1. Calculate mid-parental height. 2. Based on the child's sex, age, height and mid-parent height, find the adjustment value from the reference tables. 3. Apply the adjustment value to the child's 	

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
					height to obtain adjusted height. 4. Plot adjusted height on the 2000 CDC growth chart to obtain adjusted height percentile.	
Zhang 2013 Unites States Retrospective study	US CF Foundation Registry	PHV was calculated in relation to time interval 1994-2008. Registry records from 1986-2008 were used to identify participants.	1862 people with cystic fibrosis born in 1984-87. Age: 10 to 18 years.	Pubertal peak height velocity (PHV) <ul style="list-style-type: none"> • Normal: 60.3% (1123/1862) • Delayed: 9.4% (175/1862) • Attenuated: 20.8% (387/1862) • Delayed and attenuated: 5.3% (98/1862) • Unknown: 4.2% (79/1862) 	Longitudinal standard for peak height velocity for North American children developed by Tanner and Davies were used to define normal PHV. PHV was classified either as normal, delayed (PHV age at 2 SD later than average), attenuated (magnitude<5th percentile), or both delayed and attenuated.	High

1 Abbreviations: BMIp: BMI percentile; BMI: Body Mass Index; CDC: Center for Disease Control and Prevention;
2 CF: Cystic Fibrosis; CFF: Cystic Fibrosis Foundation; PHV: Peak Height Velocity; WHO: World Health
3 Organization

4 8.3.3 Evidence statements: impaired growth

5 Moderate quality evidence from the UK CF Trust registry found that the prevalence of
6 children who required additional nutritional support (defined as BMIp<25th) was 17.3% and
7 the prevalence of overweight (defined as BMIp>91st) was 10.1% in 2015. The same
8 evidence found that the prevalence of young people aged 12-16 who required additional
9 nutritional support (defined as BMIp<25th) was 27.5% and the prevalence of overweight
10 (defined as BMIp>91st) was 5.9% in 2015. The same evidence found that the prevalence of
11 children and young people aged 2-16 requiring additional nutritional support (defined as
12 BMIp<25th) was 20.0% and the prevalence of overweight (defined as BMIp>91st) was 9.0%.

13 Moderate quality evidence from 1 study with infants and children with cystic fibrosis aged 0-
14 24 months in the US CF Foundation Registry found that the prevalence of impaired growth at
15 12 months was the following between 2004 and 2009, depending on the definition used:

- 16 • weight for age <10th: 11%
- 17 • weight for age <5th: 8.3%
- 18 • weight for age <2.5th: 3.7%
- 19 • weight velocity (GUO-US) <50th: 48.1%

- 1 • weight velocity (WHO)<50th: 30.5%
- 2 • weight velocity <10th (WHO): 6.4%
- 3 • weight velocity<5th (WHO): 4.5%
- 4 • weight velocity<2.5th (WHO): 3.5%
- 5 • length for age <10th: 26.8%
- 6 • length for age <5th: 17.7%
- 7 • length for age <2.5th: 10.4%
- 8 • length velocity (GUO-US) <50th: 45.7%
- 9 • length velocity <50th (WHO): 38.4%
- 10 • length velocity <10th (WHO): 20.7%
- 11 • length velocity<5th (WHO): 13.4%
- 12 • length velocity<2.5th (WHO): 13.4%
- 13 • The same study found that the prevalence of impaired growth at 24 months was the
- 14 following between 2004 and 2009, depending on the definition used:
- 15 • weight for age <10th: 6.9%
- 16 • weight for age <5th: 2.8%
- 17 • weight for age <2.5th: 1.9%
- 18 • weight velocity (GUO-US) <50th: 59.3%
- 19 • weight velocity (WHO)<50th: 51.1%
- 20 • weight velocity <10th (WHO): 18.3%
- 21 • weight velocity<5th (WHO): 12.6%
- 22 • weight velocity<2.5th (WHO): 8.2%
- 23 • length for age <10th: 24.9%
- 24 • length for age <5th: 17%
- 25 • length for age <2.5th: 11.4%
- 26 • length velocity (GUO-US) <50th: 57.4%
- 27 • length velocity <50th (WHO): 58.7%
- 28 • length velocity <10th (WHO): 30.3%
- 29 • length velocity<5th (WHO): 24.3%
- 30 • length velocity<2.5th (WHO): 19.9%

31 Very low quality evidence from 1 study with 14702 people aged less than 20 years old
32 registered in the US CF Foundation Patient Registry found that the prevalence of
33 underweight defined as either height percentile <5th or BMIp<10th or %IBW<90 was 33.0%,
34 the prevalence of underweight defined as height percentile <5th or BMIp < 10th was 26.8%,
35 and the prevalence of people below BMI goal defined as BMIp<50th was 56.8% in 2002.

36 Moderate quality evidence from 1 study with 892 people with cystic fibrosis aged 1 month to
37 18 years from 10 Italian centres found that between 2005 and 2006 the prevalence of
38 malnutrition defined as HAP<5th was 15.4% and the prevalence of malnutrition defined as
39 WLP<10th was 12.9% among infants and children aged 0-2 years. The prevalence of the risk
40 of malnutrition defined as HAP 5th-25th was 18.3% and the prevalence of the risk of
41 malnutrition defined as WLP 10th -25th was 22.7% in the same age group. Among people
42 aged 2-18 years, the prevalence of malnutrition defined as HAP<5th was 11.8% and the
43 prevalence of malnutrition defined as BMIp<15th was 20.9%. The prevalence of the risk of
44 malnutrition defined as HAP 5th-25th was 29.3% and the prevalence of the risk of malnutrition
45 defined as BMIp 15th-25th was 9.6% in the same age group. The prevalence of BMIp<50th
46 was 54.4% in the same age group. Among people aged 10-14, the prevalence of malnutrition

defined as HAP<5th was 11.7% and the prevalence of malnutrition defined as BMIp<15th was 20.1%. Among people aged 14-18 years, the prevalence of malnutrition defined as HAP<5th was 21.9% and the prevalence of malnutrition defined as BMIp<15th was 27.9%. Among people aged 0-18 years old, the prevalence of malnutrition defined as HAP<5th was 12.2% and the prevalence of the risk of malnutrition defined as HAP 5th-25th was 28%.

Moderate quality evidence from a study with 3306 people with cystic fibrosis aged 2-18.5 from the CF Foundation Patient Registry of the United States found that the prevalence of impaired growth was the following between 1986 and 2005, depending on the definition used: Unadjusted height percentile <5th: 16%; unadjusted height percentile <10th: 26%; Himes adjusted height percentile <5th: 18%; Himes adjusted height percentile: <10th: 31%; CFF lower bound method: 24%.

High quality evidence from a study with 1862 people with cystic fibrosis aged 10-18 registered in the CF Foundation Patient Registry of the United States found that the prevalence of attenuated but not delayed pubertal peak velocity (magnitude <5th percentile) was 20.8%, the prevalence of attenuated and delayed pubertal peak velocity (PHV age at 2SD later than average and magnitude <5th percentile) was 5.3% between 1994 and 2008.

8.4 Cystic fibrosis related renal disease

8.4.1 Description of clinical evidence: cystic fibrosis related renal disease

The aim of this review was to determine the prevalence of cystic fibrosis related renal disease among infants, children and young people with cystic fibrosis.

We extracted data from the UK CF registry on renal failure and kidney stones. Data from the UK CF registry should be prioritized according to the protocol, however the Committee believed that the registry was likely to underreport the prevalence of renal failure because routine measurements of renal function are not very reliable; moreover it does not provide prevalence data for chronic kidney disease and acute kidney injury; therefore we also looked for observational studies on these complications. We aimed to prioritize prospective cohort studies, but only included 2 retrospective studies. We aimed to prioritize studies from more relevant contexts (the UK in the first instance, then countries in Western Europe, Australia or North America) and more recent studies.

For full details see review protocol in Appendix D.

We included data from the UK CF Registry for the year 2015 on 9587 people with cystic fibrosis. In addition to these data, we included 2 studies. One study (Quon 2011) used data from the CF Foundation Registry in the United States for 2001-2008. This study included 11912 adults with cystic fibrosis. One study (Wilcock 2015) used data from a database of an adult CF centre in the UK for 1969-2009. This study included 1532 people with cystic fibrosis.

8.4.2 Summary of included studies and results: cystic fibrosis related renal disease

A summary of the studies that were included in this review are presented in Table 51.

Table 51: Summary of included studies

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust	UK CF registry	2015	9587 people with cystic	Kidney stones: • <16 years: 0.3%	Cystic fibrosis care teams at every specialist	Moderate • Criteria

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
UK, 2016 Registry annual data report			fibrosis <ul style="list-style-type: none"> • <16 years: 3845 • ≥16 years: 5742 	(12/3845) <ul style="list-style-type: none"> • ≥16 years: 1.5% (84/5742) • Overall: 1.0% (96/9587) • Renal failure: <ul style="list-style-type: none"> • <16 years: 0% (<5) • ≥16 years: 1.0% (55/5742) • Overall: 0.6% (57/9587) 	centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	used to diagnose the complication and consistency of diagnosis across centres were unclear
Quon 2011 United States Retrospective study	CF Foundation Registry	2001-2008	11912 adults with cystic fibrosis Age ≥18 years	Mean annual prevalence of chronic kidney disease (stage 3 or greater): <ul style="list-style-type: none"> • 18-25 years old: 0.6% • Older than 55 years old: 19.2% • All ages: 2.3% Mean annual prevalence of stage 4 or greater CKD: 0.7% Mean annual prevalence of stage 5 CKD: 0.6%	CKD was defined by eGFR measured less than 60 ml/min/1.73 m ² in 2 consecutive registry years. Based on National Kidney Foundation KDOQI guidelines, this corresponds to stage 3 CKD severity and is the earliest stage that can be diagnosed using serum creatinine alone. More advanced stages of CKD were defined as follows: stage 4, eGFR less than 30 ml/min/1.73 m ² ; and stage 5, eGFR less than 15 ml/min/1.73 m ² or need for hemodialysis.	Moderate <ul style="list-style-type: none"> • No numerators or denominators
Wilcock 2015 UK Retrospective study	Database of adult CF department at the Royal Brompton	1969-2009	1532 people with cystic fibrosis <ul style="list-style-type: none"> • Age: Not reported 	<ul style="list-style-type: none"> • Renal problem: 5.1% (78/1532) • Acute kidney injury: 1.1%* (17/1532) (9 cases were presumed to be drug-induced) • Chronic kidney 	<ul style="list-style-type: none"> • AKI was defined as an acute rise in creatinine above the person's established baseline; • CKD as an abnormal 	Moderate <ul style="list-style-type: none"> • Percentage not provided for some conditions (calculated by the NGA)

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
	Hospital			disease: 0.9% * (13/1532) (4 cases were presumed to be drug-induced) <ul style="list-style-type: none"> • Renal stone disease: 2.0% (30/1532) • Isolated proteinuria: 0.1% * (2/1532) • Isolated haematuria: 0.5% * (8/1532) • Nephrotic syndrome: 0.1%* (2/1532) • Miscellaneous disease: 0.5% * (8/1532) 	creatinine level for greater than 3 months; <ul style="list-style-type: none"> • Nephrotic syndrome as proteinuria (>3g/l), hypoalbuminaemia and peripheral oedema. Cases were excluded if: <ul style="list-style-type: none"> • isolated rise in plasma urea which resolved with no other evidence of renal impairment • impaired renal function in the last few weeks of life • nephrotoxic immunosuppressant therapy before any episode of renal disease 	technical team) <ul style="list-style-type: none"> • Study subjects were not described in detail

1 Abbreviations: AKI: Acute Kidney; CF: cystic fibrosis; CKD: Chronic Kidney Disease; Injury
2 * Percentage calculated by the NGA technical team

3 8.4.3 Evidence statements: cystic fibrosis related renal disease

4 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
5 Registry found that in 2015 the prevalence of kidney stones was 0.3% amongst children and
6 young people aged <16, 1.5% amongst young people and adults aged 16 years and over,
7 and 1.0% amongst people of all ages.

8 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
9 Registry found that in 2015 the prevalence of renal failure was 0% amongst children and
10 young people aged 10-15, 1.0% amongst young people and adults aged 16 years and over,
11 and 0.6% amongst people of all ages.

12 Moderate quality evidence from a study on 11912 people with cystic fibrosis aged 18 years
13 and over registered in the CF Foundation Registry in the United States found that between
14 2001 and 2008 the mean annual prevalence of chronic kidney disease (stage 3 or greater)
15 was 0.6% among people aged 18-25, 19.2% among people older than 55 years old, and
16 2.3% among people of all ages. The mean annual prevalence of stage 4 or greater CKD was
17 0.7% and the mean annual prevalence of stage 5 CKD was 0.6%.

18 Moderate quality evidence from a study on 1532 people with cystic fibrosis from the adult
19 department of a centre in the UK found that between 1969 and 2009 the prevalence of renal

1 problems was 5.1%; the prevalence of acute kidney injury was 1.1%; the prevalence of CKD
2 was 0.9%.

3 8.4.3.1 Economic evidence statements

4 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

5 8.5 Delayed puberty

6 8.5.1 Description of clinical evidence: delayed puberty

7 The aim of this review was to determine the prevalence of delayed puberty among people
8 with cystic fibrosis.

9 The UK CF registry did not provide prevalence data on this complication; therefore we looked
10 for observational studies. We only identified 1 study eligible for inclusion.

11 For full details see review protocol in Appendix D.

12 One study (Zhang 2013) used data from the CF Foundation Registry in the United States for
13 1994-2008. This study included 1862 people with cystic fibrosis aged 10-18 years old.

14 8.5.2 Summary of included studies and results: delayed puberty

15 A summary of the studies that were included in this review are presented in Table 52.

16 **Table 52: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age group - % (n/N)	Definitions of complication and measurement tools	Quality of the study
Zhang 2013 Unites States Retrospective study	US CF Foundat ion Registry	PHV was calcula ted in relatio n to time interva l 1994-2008. Registr y record s from 1986-2008 were used to identify partici pants.	1862 people with cystic fibrosis born in 1984-87. Age: 10 to 18 years.	Pubertal peak height velocity (PHV) <ul style="list-style-type: none"> • Normal: 60.3% (1123/1862) * • Delayed: 9.4% (175/1862) * • Attenuated: 20.8% (387/1862) * • D&A: 5.3% (98/1862) * • Unknown: 4.2% (79/1862) * 	Longitudinal standard for peak height velocity for North American children developed by Tanner and Davies were used to define normal PHV. PHV was classified either as normal, delayed (PHV age at 2 SD later than average), attenuated (magnitude<5 th percentile), or both delayed and attenuated (D&A).	High

17 Abbreviations: CF: cystic fibrosis; D&A – Delayed and Attenuated; PHV: Peak Height Velocity
18 *Numerator calculated by NGA technical team

1 **8.5.3 Evidence statements: delayed puberty**

2 High quality evidence from a study with 1862 people with cystic fibrosis aged 10-18 from the
3 CF Foundation Patient Registry of the United States found that the prevalence of delayed but
4 not attenuated pubertal peak velocity (PHV age at 2SD later than average) was 9.4% and the
5 prevalence of attenuated and delayed pubertal peak velocity (PHV age at 2SD later than
6 average and magnitude <5th percentile) was 5.3% between 1994 and 2008.

7 **8.5.3.1 Economic evidence statements**

8 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

9 **8.6 Abdominal pain**

10 **8.6.1 Description of clinical evidence: abdominal pain**

11 The aim of this review was to determine the prevalence of abdominal pain among infants,
12 children and young people with cystic fibrosis.

13 The UK CF registry does not provide prevalence data on this complication. Therefore, we
14 looked for relevant observational studies. However, we found no evidence for this
15 complication.

16 For full details see review protocol in Appendix D.

17 **8.6.2 Summary of included studies and results: abdominal pain**

18 No evidence was found on the prevalence of this complication among people with cystic
19 fibrosis.

20 **8.6.3 Evidence statements: abdominal pain**

21 No clinical or cost-effectiveness evidence was found on the prevalence of this complication
22 among people with cystic fibrosis.

23 **8.7 Cystic Fibrosis related diabetes**

24 **8.7.1 Description of clinical evidence: cystic fibrosis related diabetes**

25 The aim of this review was to determine the prevalence of Cystic Fibrosis Related Diabetes
26 (CFRD) amongst people with cystic fibrosis. We looked for data in the UK CF registry 2015.
27 This provided the prevalence of people on treatment for CFRD. This data was included
28 because data from the UK CF registry was prioritized according to the protocol. However
29 only considering people on treatment would underestimate the prevalence of CFRD.
30 Therefore we also looked for observational studies, excluding studies where CFRD was
31 defined based on treatment use only. We aimed to prioritize prospective cohort studies, but
32 only included retrospective studies. We prioritized studies from more relevant contexts (the
33 UK in the first instance, then countries in Western Europe, Australia or North America). We
34 also prioritized studies with more recent data and studies that disaggregated data between
35 infants, children, young people and adults; as well as studies that reported prevalence at
36 different points in time and studies that were based on registries.

37 We included data from the UK CF Registry for the year 2015 on 6970 people with cystic
38 fibrosis aged ≥ 10 years. In addition to data from the UK CF registry, we included 3 studies.
39 One study (Lewis 2015) used data from the Minnesota Cystic Fibrosis database (United
40 States) from 2008 to 2012. This study included 462 people with cystic fibrosis. One study

1 (Moran 2009) used the same database to calculate prevalence at different points in time
2 (1992-97, 1998-2000, 2003-2008) based on 872 people. Another study (Bell 2011) used the
3 Australian CF registry data from 2009. This study included 2986 people with cystic fibrosis.

4 8.7.2 Summary of included studies and results: cystic fibrosis related diabetes

5 A summary of the studies that were included in this review are presented in Table 53.

6 **Table 53: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	6970 people with cystic fibrosis aged ≥10 years	Treatment for CFRD: <ul style="list-style-type: none"> • 10-16 years: 10.0% (134/1624) • ≥16 years: 32.2% (1848/5346) • ≥10 years: 28.0% (1982/6970) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> • Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear
Bell 2011 Australia Retrospective study	Australian CF Registry	2009	2986 people with cystic fibrosis Median age: 17.6 years	Insulin-dependent diabetes (chronic): <ul style="list-style-type: none"> • 0-11 years: 0.5% (5/951) • 12-17 years: 13.6% (61/448) • ≥18 years: 20.7% (144/697) • All age groups: 10.0% (210/2096) Insulin-dependent diabetes (intermittent): <ul style="list-style-type: none"> • 0-11 years: 0% (0/951) • 12-17 years: 1.1% (5/448) • ≥18 years: 2.3% (16/697) • All age groups: 1.0% (21) 	Not reported	Moderate <ul style="list-style-type: none"> • Unclear criteria for measuring the condition • Unclear if different centres across Australia would measure the condition consistently
Lewis 2015 United States Retrospective	Data from people seen at the UNM CF	September 2008-December 2012	462 people Age: ≥ 20 years	CFRD: 48% (221/462)	Diabetes diagnosed by "standard criteria".	Moderate <ul style="list-style-type: none"> • Criteria to diagnose CFRD not reported • Unclear if

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
study	centre					there was consistency to diagnose CFRD across the centre
Moran 2009 United States Retrospective study	Minnesota Cystic Fibrosis Database	Prevalence at the end of 3 consecutive intervals: 1992-1997, 1998-2002, 2003-2008	872 people with cystic fibrosis followed at the University of Minnesota Cystic Fibrosis Centre Age: not reported	CFRD prevalence at the end of the interval: (%): <ul style="list-style-type: none"> • 1992-97: 20% +/-2% • 1998-2000: 30% +/-2% • 2003-2008: 33% +/-2% Prevalence of CFRD in September 2008: <ul style="list-style-type: none"> • Children < 11 years: 2% (2/93) (both without fasting hyperglycemia). • Young people aged 11-17, 19% (14/75) (4 with fasting hyperglycemia). • Adults aged ≥18: 43%* (155/359) 	CFRD was diagnosed by standard criteria including persistent random glucose levels >200mg/dl (11.1 mmol/l) and persistent fasting glucose levels >126 mg/dl (7.0 mmol/l) or by OGTT. People with a fasting glucose ≥126 mg/dl (7.0 mmol/l) were diagnosed with CFRD with fasting hyperglycemia. People with a fasting glucose 126 mg/dl (7.0 mmol/l) and a 2-h glucose ≥200 mg/dl (11.1 mmol/l) were diagnosed with CFRD without fasting hyperglycemia. Routine annual OGTT screening has been recommended at the University of Minnesota since the early 1990s for patients aged ≥6 years (1.75 g/kg glucose [maximum 75g]). OGTTs are performed when patients are	Moderate <ul style="list-style-type: none"> • No numerator or denominator provided for prevalence calculated at the end of the interval

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
					in their usual baseline state of health.	

Abbreviations: CF: cystic fibrosis; CFRD: Cystic Fibrosis Related Diabetes; g: grams; kg: kilograms; mmol/l: millimoles per litre; OGTT: Oral Glucose Tolerance Test; SD: Standard deviation.

* Percentage calculated by NGA technical team

1
2
3
4

5 8.7.3 Evidence statements: cystic fibrosis related diabetes

6 Moderate quality evidence from a report on from on 9587 people with cystic fibrosis from the
7 UK CF Registry found that in 2015 the prevalence of people on treatment for CFRD was
8 10.0% among children and young people aged 10-16, 32.2% among young people and
9 adults aged 16 years and over, and 28.0% among all people aged 10 years and over.

10 Moderate quality evidence from 1 study with 2986 people with cystic fibrosis from the
11 Australian CF Registry found that om 2009 the prevalence of chronic insulin-dependent
12 diabetes was 0.5% among infants and children aged 0-11, 13.6% among young people aged
13 12-17, 20.7% among adults aged 18 and over, and 10.0% among all age groups. The
14 prevalence of intermittent insulin-dependent diabetes was 0% among infants and children
15 aged 0-11, 1.1% among children aged 12-17, 2.3% among adults aged 18 years and over,
16 1.0% among all age groups.

17 Low quality evidence from 1 study with 462 people attending a CF centre in the United
18 States found that the prevalence of CFRD was 48% among adults aged 20 years and over
19 between 2008 and 2012.

20 Moderate quality evidence from 1 study with 872 people attending a CF centre in the United
21 States found that the prevalence of CFRD was 20%+2% at the end of the interval 1992-
22 1997, 30%+2% at the end of the interval 1998-2000, 33%+2% at the end of the interval
23 2003-2008. The prevalence of CFRD in 2008 was 2% among infants and children aged less
24 than 11 years, 19% amongst children and young people aged 11-17 and 43% among adults
25 aged 18 years and over.

26 8.7.3.1 Economic evidence statements

27 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

28 8.8 Upper airways disease

29 8.8.1 Description of clinical evidence: upper airways disease

30 The aim of this review was to determine the prevalence of upper airways disease among
31 people with cystic fibrosis. We included data from the UK CF Registry for the year 2015 on
32 9587 people with cystic fibrosis. Data provided by the registry was prioritized therefore we
33 did not include any studies from the published literature.

34 For full details see review protocol in Appendix D.

1 8.8.2 Summary of included studies and results: upper airways disease

2 A summary of the studies that were included in this review are presented in Table 54.

3 **Table 54: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> • <16 years: 3845 • ≥16 years: 5742 	Nasal polyps requiring surgery: <ul style="list-style-type: none"> • <16 years: 1.1% (44/3845) • ≥16 years: 3.1% (177/5742) • Overall: 2.3% (221/9587) Sinus disease <ul style="list-style-type: none"> • <16 years: 1.4% (53/3845) • ≥16 years: 15.4% (886/5742) • Overall: 9.8% (939/9587) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> • Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear

4 *Abbreviations: ABPA: Allergic bronchopulmonary aspergillosis; CF: cystic fibrosis*

5 8.8.3 Evidence statements: Upper airways disease

6 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
7 Registry found that in 2015 the prevalence of nasal polyps requiring surgery was 1.1%
8 among children and young people aged <16, 3.1% among young people and adults aged 16
9 years and over, and 2.3% among people of all ages.

10 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
11 Registry found that in 2015 the prevalence of sinus disease was 1.4% among children and
12 young people aged <16, 15.4% among young people and adults aged 16 years and over,
13 and 9.8% among people of all ages.

14 8.8.3.1 Economic evidence statements

15 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

16 8.9 Cystic fibrosis related musculoskeletal disorders

17 8.9.1 Description of clinical evidence: cystic fibrosis related musculoskeletal disorders

18 The aim of this review was to determine the prevalence of cystic fibrosis related
19 musculoskeletal disorders among people with cystic fibrosis.

20 We extracted data from the UK CF registry on the prevalence of arthritis and arthropathy.
21 Data from the UK CF registry should be prioritized according to the protocol, however the
22 Committee believed that the registry was likely to underreport the prevalence of these
23 complications as they are common in the general population and centres may not regard
24

1 them as a cystic fibrosis specific complication. Therefore we also looked for observational
2 studies. We aimed to prioritize prospective cohort studies but only found 1 retrospective
3 study eligible for inclusion.

4 For full details see review protocol in Appendix D.

5 We included data from the UK CF Registry for the year 2015 on 9587 people with cystic
6 fibrosis. In addition to these data, the included study (Watts 2009) used data from the CFF
7 Patient Registry in the United States from 2004. This study included 22714 people with cystic
8 fibrosis.

9 8.9.2 Summary of included studies and results: Cystic fibrosis related 10 musculoskeletal disorders

11 A summary of the studies that were included in this review are presented in Table 55.

12 **Table 55: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> <16 years: 3845 ≥16 years: 5742 	Arthritis: <ul style="list-style-type: none"> <16 years: 0.2% (7/3845) ≥16 years: 2.6% (151/5742) Overall: 1.6% (158/9587) Arthropathy: <ul style="list-style-type: none"> <16 years: 0.5% (18/3845) ≥16 years: 8.7% (499/5742) Overall: 5.4% (517/9587) • 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear
Watts 2009 United States Retrospective study	CFF Patient Registry	2004	22714 people with cystic fibrosis Age: not reported for the overall cohort	Bone and joint complications: 6.7% (1510/22714) *	Bone and joint complications include arthritis/arthropathy, bone fractures, osteopenia, osteoporosis	Very low <ul style="list-style-type: none"> No separate data for each complication Unclear how complications were diagnosed and whether they were diagnosed consistently Data not disaggregated

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
						ted by age subgroup

1 Abbreviations: CF: cystic fibrosis
2 *Prevalence calculated by the NGA technical team

3 8.9.3 Evidence statements: cystic fibrosis related musculoskeletal disorders

4 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
5 Registry found that in 2015 the prevalence of arthritis was 0.2% among children and young
6 people aged <16, 2.6% among young people and adults aged 16 years and over, and 1.6%
7 among people of all ages.

8 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
9 Registry found that in 2015 the prevalence of arthropathy was 0.5% among children and
10 young people aged <16, 8.7% among young people and adults aged 16 years and over, and
11 5.4% among people of all ages.

12 Very low quality from 1 study on 22714 people with cystic fibrosis from the CF Foundation
13 Patient Registry in the United States found that the prevalence of bone and joint
14 complications was 6.7% in 2004.

15 8.9.3.1 Economic evidence statements

16 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

17 8.10 Urinary stress incontinence

18 8.10.1 Description of clinical evidence: urinary stress incontinence

19 The aim of this review was to determine the prevalence of urinary stress incontinence among
20 people with cystic fibrosis.

21 The UK CF registry does not provide prevalence data on this complication. Therefore, we
22 looked for relevant observational studies. However, we found no evidence for this
23 complication.

24 For full details see review protocol in Appendix D.

25 8.10.2 Summary of included studies and results: urinary stress incontinence

26 No evidence was found on the prevalence of this complication among people with cystic
27 fibrosis.

28 8.10.3 Evidence statements: urinary stress incontinence

29 No evidence was found on the prevalence of this complication among people with cystic
30 fibrosis.

1 8.11 Reduced bone mineral density

2 8.11.1 Description of clinical evidence: reduced bone mineral density

3 The aim of this review was to determine the prevalence of reduced bone mineral density
4 among people with cystic fibrosis.

5 We extracted data from the UK CF registry on the prevalence of osteopenia, osteoporosis
6 and bone fractures. Data from the UK CF registry should be prioritized according to the
7 protocol, however the Committee believed that the registry was likely to underreport the
8 prevalence of reduced bone mineral density as these complications can only be diagnosed
9 by DEXA scan for which the take-up of these is not good, so many cases will be
10 undiscovered. Therefore we also looked for observational studies on this complication. We
11 aimed to prioritize prospective cohort studies, but only included 2 retrospective studies. We
12 aimed to prioritize studies from more relevant contexts (the UK in the first instance, then
13 countries in Western Europe, Australia or North America) and more recent studies but only 2
14 studies, both on data from 2009, were eligible for inclusion.

15 For full details see review protocol in Appendix D.

16 We included data from the UK CF Registry for the year 2015 on 9587 people with cystic
17 fibrosis. In addition to this data, we included 2 studies. One study (Bell 2011) used data from
18 the Australian CF registry for 2009. It included 2986 people with cystic fibrosis. One study
19 (Somerville 2013) used data from the CF Registry of Ireland. It included 859 people with
20 cystic fibrosis.

21 8.11.2 Summary of included studies and results: reduced bone mineral density

22 A summary of the studies that were included in this review are presented in Table 56.

23 **Table 56: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> <16 years: 3845 ≥16 years: 5742 	Osteopenia: <ul style="list-style-type: none"> <16 years: 0.9% (36/3845) ≥16 years: 22.0% (1261/5742) Overall: 13.5% (1297/9587) Osteoporosis: <ul style="list-style-type: none"> <16 years: 0% (<5/3845) ≥16 years: 8.8% (507/5742) Overall: 5.3% (511/9587) Bone fracture: <ul style="list-style-type: none"> <16 years: 0.4% (14/3845) ≥16 years: 0.6% (32/5742) Overall: 0.5% 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
				(46/9587)		
Bell 2011 Retrospective study	Australian CF Registry	2009	2986 people with cystic fibrosis Median age: 17.6 years	<p>Osteopenia:</p> <ul style="list-style-type: none"> • 0-11 years: 0.3% (3/951) • 12-17 years: 3.3% (15/448) • ≥18 years: 25.0% (174/697) • All age groups: 9.2% (192/2096) <p>Osteoporosis:</p> <ul style="list-style-type: none"> • 0-11 years: 0.2% (2/951) • 12-17 years: 1.3% (6/448) • ≥18 years: 9.5% (66/697) • All age groups: 3.7% (77/2096) <p>Fractures in 2009:</p> <ul style="list-style-type: none"> • 0-11 years: 0% (0/951) • 12-17 years: 0.4% (2/448) • ≥18 years: 1.3% (9/697) • All age groups: 0.5% (10/2096) 	Not reported	<p>Moderate</p> <ul style="list-style-type: none"> • Unclear criteria for measuring the condition • Unclear if different centres across Australia would measure the condition consistently
Somerville 2013 Ireland Retrospective study	CF Registry of Ireland	All people alive on 31/12/2009. Data recorded from 2001	859 people with cystic fibrosis*	<p>Osteopenia or osteoporosis:</p> <ul style="list-style-type: none"> • <18 years: 5.5% (25/454) ¥ • ≥18 years: 42.7% (173/405) 	Considered present if documented in the in the medical notes in the last year	<p>Low</p> <ul style="list-style-type: none"> • N used in the analysis was different from reported N but this was not mentioned by the authors • No denominator was provided for one age subgroup • No baseline characteristics were

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
						provided for people younger than 18

1 Abbreviations: CF: cystic fibrosis
 2 * N calculated by NGA technical team
 3 ‡Denominator calculated by NGA technical team

4 8.11.3 Evidence statements: reduced bone mineral density

- 5 • Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK
 6 CF Registry found that in 2015 the prevalence of osteopenia was 0.9% among children
 7 and young people aged <16, 22.0% among young people and adults aged 16 years and
 8 over, and 13.5% among people of all ages.
- 9 • Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK
 10 CF Registry found that in 2015 the prevalence of osteoporosis was 0% among children
 11 and young people aged <16, 8.8% among young people and adults aged 16 years and
 12 over, and 5.3% among people of all ages.
- 13 • Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK
 14 CF Registry found that in 2015 the prevalence of bone fractures was 0.4% among children
 15 and young people aged <16, 0.6% among young people and adults aged 16 years and
 16 over, and 0.5% among people of all ages.
- 17 • Moderate quality evidence from a study on 2986 people with cystic fibrosis from the
 18 Australian CF registry found that in 2009 the prevalence of osteopenia was 0.3% among
 19 infants and children aged 0-11, 3.3% among young people aged 12-17, 25.0% among
 20 adults, 9.2% among people of all ages. The prevalence of osteoporosis was 0.2% among
 21 infants and children aged 0-11, 1.3% among young people aged 12-17, 9.5% among
 22 adults, 3.7% among people of all ages. The prevalence of fractures was 0% among
 23 infants and children aged 0-11, 0.4% among young people aged 12-17, 1.3% among
 24 adults, 0.5% among people of all ages.
- 25 • Low quality evidence from a study on 859 people with cystic fibrosis from the CF Registry
 26 of Ireland found that in 2009 the prevalence of osteopenia or osteoporosis was 5.5%
 27 among people aged <18 years, and 42.7% among people aged ≥18 years.

28 8.11.3.1 Economic evidence statements

29 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

30 8.12 Cystic fibrosis related liver disease

31 8.12.1 Description of clinical evidence: cystic fibrosis related liver disease

32 The aim of this review was to determine the prevalence of cystic fibrosis related liver disease
 33 among people with cystic fibrosis. We included data from the UK CF Registry for the year
 34 2015 on 9587 people with cystic fibrosis. The data provided by the registry was prioritized
 35 therefore we did not include any studies from the published literature.

36 For full details see review protocol in Appendix D.

1 8.12.2 Summary of included studies and results: cystic fibrosis related liver disease

2 A summary of the studies that were included in this review are presented in Table 57.

3 **Table 57: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> • <16 years: 3845 • ≥16 years: 5742 	<p>Raised liver enzymes:</p> <ul style="list-style-type: none"> • <16 years: 6.9% (264/3845) • ≥16 years: 14.8% (852/5742) • Overall: 11.6% (1116/9587) <p>Liver disease:</p> <ul style="list-style-type: none"> • <16 years: 8.8% (340/3845) • ≥16 years: 18.0% (1031/5742) • Overall: 14.3% (1371/9587) <p>Cirrhosis with no portal hypertension:</p> <ul style="list-style-type: none"> • <16 years: 0.7% (26/3845) • ≥16 years: 1.6% (90/5742) • Overall: 1.2% (116/9587) <p>Cirrhosis with portal hypertension:</p> <ul style="list-style-type: none"> • <16 years: 0.7% (26/3845) • ≥16 years: 2.4% (138/5742) • Overall: 1.7% (164/9587) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> • Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear

4 *Abbreviations: CF: cystic fibrosis*

5 8.12.3 Evidence statements: cystic fibrosis related liver disease

6 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
7 Registry found that in 2015 the prevalence of raised liver enzymes was 6.9% among children
8 and young people aged <16, 14.8% among young people and adults aged 16 years and
9 over, and 11.6% among people of all ages.

10 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
11 Registry found that in 2015 the prevalence of liver disease was 8.8% among children and
12 young people aged <16, 18.0% among young people and adults aged 16 years and over,
13 and 14.3% among people of all ages.

1 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
2 Registry found that in 2015 the prevalence of cirrhosis with no portal hypertension was 0.7%
3 among children and young people aged <16, 1.6% among young people and adults aged 16
4 years and over, and 1.2% among people of all ages.

5 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
6 Registry found that in 2015 the prevalence of cirrhosis with portal hypertension was 0.7%
7 among children and young people aged <16, 2.4% among young people and adults aged 16
8 years and over, and 1.7% among people of all ages.

9 **8.12.3.1 Economic evidence statements**

10 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

11 **8.13 Infertility**

12 **8.13.1 Description of clinical evidence: infertility**

13 The aim of this review was to determine the prevalence of infertility among people with cystic
14 fibrosis.

15 The UK CF registry does not provide prevalence data on this complication. Therefore, we
16 looked for relevant observational studies. However, we found no evidence for this
17 complication.

18 For full details see review protocol in Appendix D.

19 **8.13.2 Summary of included studies and results: infertility**

20 No evidence was found on the prevalence of this complication among people with cystic
21 fibrosis.

22 **8.13.3 Evidence statements: infertility**

23 No evidence was found on the prevalence of this complication among people with cystic
24 fibrosis.

25 **8.14 Distal Intestinal Obstruction Syndrome (DIOS)**

26 **8.14.1 Description of clinical evidence: Distal Intestinal Obstruction Syndrome (DIOS)**

27 The aim of this review was to determine the prevalence of distal intestinal obstruction
28 syndrome among people with cystic fibrosis.

29 The UK CF registry provided data on intestinal obstruction, rather than on distal intestinal
30 obstruction syndrome; therefore we also looked for observational studies on this
31 complication. We aimed to prioritize prospective cohort studies, but only included 1
32 retrospective study. We prioritized studies from more relevant contexts (the UK in the first
33 instance, then countries in Western Europe, Australia or North America). We also requested
34 data to the registry on intestinal obstruction disaggregated by history of meconium ileus,
35 because the Committee agreed that prevalence may vary considerably between people with
36 or without a history of meconium ileus.

37 For full details see review protocol in Appendix D.

38 We included data from the UK CF Registry for the year 2015 on 9587 people with cystic
39 fibrosis. In addition to these data, we included 1 study. This study (Wiedemann 2001) used

1 data from the CF Quality Assurance Project Registry for 1997. This study included 3448
2 people with cystic fibrosis.

3 8.14.2 Summary of included studies and results: Distal Intestinal Obstruction 4 Syndrome (DIOS)

5 A summary of the studies that were included in this review are presented in Table 58.

6 **Table 58: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> • <16 years: 3845 • ≥16 years: 5742 	Intestinal obstruction: <ul style="list-style-type: none"> • <16 years: 3.0% (116/3845) • ≥16 years: 7.4% (423/5742) • Overall: 5.6% (539/9587) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> • Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear
Data request to UK CF registry	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> • Age <16: 3845 • Age ≥16: 5742 	Intestinal obstruction in age group <16: <ul style="list-style-type: none"> • Diagnosis of meconium ileus: 6.4% (41/643) • No diagnosis of meconium ileus: 2.3% (75/3202) Intestinal obstruction in age group ≥16: <ul style="list-style-type: none"> • Diagnosis of meconium ileus: 13.9% (113/815) • No diagnosis of meconium ileus: 6.3% (310/4927) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK entered data on meconium ileus and intestinal obstruction for the registry.	Moderate <ul style="list-style-type: none"> • Details from the people in the registry are available in the UK Cystic Fibrosis Trust Registry 2015 Annual Data Report and on the CF Trust website. However criteria used to diagnose the complication and consistency of diagnosis across

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
						centres were unclear from these sources
Wiedemann 2001 Germany Retrospective study	CF Quality Assurance Project registry	1997	3448 people with cystic fibrosis Age range: 0-58	DIOS <ul style="list-style-type: none"> • Children and young people (age not reported): 3.0% • Adults (age not reported): 3.5% • All ages: 3.2% 	Criteria used to diagnose DIOS not reported. Centres reported data for each person once a year from a routine visit near the person's birthday when the person was in a stable clinical condition.	Low <ul style="list-style-type: none"> • Unclear how DIOS was diagnosed and whether it was diagnosed consistently across centres. • Age cut-off to define age subgroups was unclear • Numerators and denominators not provided. • Data was not disaggregated between children and young people

1 Abbreviations: CF: cystic fibrosis; DIOS: Distal Intestinal Obstruction Syndrome

2 8.14.3 Evidence statements: Distal Intestinal Obstruction Syndrome (DIOS)

3 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
4 Registry found that in 2015 the prevalence of intestinal obstruction was 3.0% among children
5 and young people aged <16, 7.4% amongst young people and adults aged 16 years and
6 over, and 5.6% amongst people of all ages.

7 Moderate quality evidence from the UK CF Registry with 9587 people found that in 2015 the
8 prevalence of intestinal obstruction was 6.4% among children and young people aged <16
9 with a diagnosis of meconium ileus, and 2.3% among children and young people aged <16
10 without a diagnosis of meconium ileus. The same evidence found that the prevalence of
11 intestinal obstruction was 13.9% among people aged ≥16 with a diagnosis of meconium ileus
12 and 6.3% among people aged ≥16 without a diagnosis of meconium ileus.

1 Low quality evidence from a study on 3448 people with cystic fibrosis from the Quality
2 Assurance Project registry in Germany found that in 1997 prevalence of DIOS was 3.0%
3 among children and young people, 3.5% among adults, and 3.2% among all ages.

4 8.14.3.1 Economic evidence statements

5 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

6 8.15 Meconium ileus

7 8.15.1 Description of clinical evidence: meconium ileus

8 The aim of this review was to determine the prevalence of meconium ileus among people
9 with cystic fibrosis. We included data from the UK CF Registry for the year 2015 on 9587
10 people with cystic fibrosis. The data provided by the registry was prioritized therefore we did
11 not include any studies from the published literature.

12 For full details see review protocol in Appendix D.

13 8.15.2 Summary of included studies and results: meconium ileus

14 A summary of the studies that were included in this review are presented in Table 59.

15 **Table 59: Summary of included studies**

Study	Register / Data source	Dates	Participants	Prevalence: overall results and results by age subgroups - % (n/N)	Definition of complication and measurement tools	Quality of the study
Cystic Fibrosis Trust UK, 2016 Registry annual data report	UK CF registry	2015	9587 people with cystic fibrosis <ul style="list-style-type: none"> • <16 years: 3845 • ≥16 years: 5742 	Meconium ileus: <ul style="list-style-type: none"> • <16 years: 16.7% (643/3845) • ≥16 years: 14.2% (815/5742) • Overall: 15.2% (1458/9587) 	Cystic fibrosis care teams at every specialist centre and clinic across the UK filled in forms where they selected what complications (if any) each person with cystic fibrosis had. No definition of complications is provided in the form.	Moderate <ul style="list-style-type: none"> • Criteria used to diagnose the complication and consistency of diagnosis across centres were unclear

16 *Abbreviations: CF: cystic fibrosis*

17 8.15.3 Evidence statements: meconium ileus

18 Moderate quality evidence from a report on 9587 people with cystic fibrosis from the UK CF
19 Registry (year 2015) found that meconium ileus had occurred in 16.7% of children and young
20 people aged <16, 14.2% of young people and adults aged 16 years and over, and 15.2% of
21 people of all ages.

22 8.15.3.1 Economic evidence statements

23 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

1 **8.16 Economic evidence**

2 No economic evaluations related to the complications of cystic fibrosis were identified in the
3 literature search conducted for this guideline. Full details of the search and economic article
4 selection flow chart can be found in Appendix E and F, respectively.

5 This review question is not relevant for economic analysis because it does not involve a
6 decision between alternative courses of action.

7 **8.17 Evidence to recommendations**

8 **8.17.1 Relative value placed on the outcomes considered**

9 The Committee agreed that the prevalence of complications of cystic fibrosis was a critical
10 outcome.

11 **8.17.2 Consideration of clinical benefits and harms**

12 **8.17.2.1 Malnutrition and impaired growth**

13 The Committee noted that the evidence on prevalence of vitamin deficiency was of very low
14 quality in relation to vitamins A and E; there was no evidence in relation to vitamin K,
15 however based on their clinical expertise the Committee agreed that vitamin K deficiency
16 may occur; the quality of the evidence in relation to vitamin D ranged between very low and
17 moderate. The Committee agreed that pancreatic insufficiency (common in cystic fibrosis)
18 hinders the body's inability to digest fat hence fat-soluble vitamins are poorly absorbed. The
19 Committee noted that prevalence of vitamin deficiency would depend on whether people take
20 vitamin supplements; moreover the prevalence of vitamin D deficiency would change
21 depending on the time of the year. Therefore, the Committee decided not to specify a
22 numerical estimate of prevalence of vitamin deficiency in the recommendations. They
23 recommended to be aware that fat-soluble vitamin deficiencies (including vitamins A, D, E
24 and K) are common in people with cystic fibrosis.

25 The Committee noted that a multitude of anthropometric indicators of malnutrition and
26 impaired growth was used in the studies, and agreed that in the future consistency in
27 indicators is needed across studies in order to meaningfully compare results. The Committee
28 decided to prioritise the data from the UK registry on BMI (recorded as percentile until age 16
29 and in units of kg/m² for people older than 16) in order to formulate recommendations,
30 because this data was the most recent, and older studies were likely to show higher
31 prevalence due to treatment regimens that would not correspond to the current ones.
32 However, the Committee noted that the registry data indicated that children with cystic
33 fibrosis have better BMI compared to the non-cystic fibrosis population (17% of children
34 <25th percentile) and young people with cystic fibrosis have similar BMI compared to the
35 non-cystic fibrosis population (27.5% of young people <25th percentile) and this was in
36 contrast with their experience in clinical practice. Therefore the Committee decided not to
37 give a numerical estimate of the prevalence of reduced BMI and prioritised a
38 recommendation to be aware that underweight is common in people with cystic fibrosis.

39 **8.17.2.2 Meconium ileus**

40 The Committee used the data from the UK CF registry to formulate a recommendation on
41 meconium ileus. The prevalence in the overall population corresponds to 1 in 7 people.
42 Therefore they recommended to be aware that meconium ileus is common in people with
43 cystic fibrosis (affects 1 in 7 newborn babies).

1 8.17.2.3 Abdominal pain

2 There was no evidence on abdominal pain. Abdominal pain is common in school-age
3 children. The cause is often uncertain and most do not require investigation. Abdominal pain
4 may also occur in adults for example due to irritable bowel syndrome. Against this clinical
5 background, people with cystic fibrosis might also experience abdominal pain and in some
6 cases this might be due to cystic fibrosis. The Committee agreed that the gastrointestinal
7 complications of cystic fibrosis can cause abdominal pain. For example, pancreatic
8 insufficiency requires treatment with pancreatic enzyme replacement therapy and getting the
9 dosage right can be difficult in some patients and constipation or diarrhoea can be common
10 when titrating doses (which may cause abdominal pain); other causes of abdominal pain are:
11 distal intestinal obstruction syndrome; constant coughing; inflammatory bowel disease;
12 gastro-oesophageal reflux disease. Less common causes of abdominal pain in children with
13 cystic fibrosis include recurrent intussusception; volvulus; fibrosing colonopathy; and
14 appendiceal disease. The Committee agreed that abdominal pain can be very disabling,
15 however it is difficult to give an estimate of prevalence. For example, some people with cystic
16 fibrosis are continuously asked if they have abdominal pain, which may lead to over
17 reporting. Moreover, many people present with non-specific abdominal pain (generalised
18 abdominal pain with no specific focal point and no definable cause). The Committee did not
19 include abdominal pain as a specific complication of cystic fibrosis because in each case the
20 underlying likely explanation might differ and the cause would need consideration.

21 8.17.2.4 Distal ileal obstruction syndrome

22 The Committee prioritised the data from the UK CF registry because it was more recent than
23 the data from the included study from the published literature. The data from the registry was
24 on intestinal obstruction rather than on distal ileal obstruction syndrome, however the
25 Committee agreed that intestinal obstruction would mostly be due to distal ileal obstruction
26 syndrome in people with cystic fibrosis. The Committee noted that the data provided by the
27 UK CF registry showed an important difference in prevalence between those who have a
28 history of meconium ileus and those without a history of meconium ileus, as well as between
29 children and adults. However the Committee agreed that the recommendations should not go
30 into too much detail in terms of numerical estimates of prevalence because the risk of
31 complications would vary considerably from person to person. Therefore the Committee
32 decided to keep the recommendation general and recommended to be aware that distal ileal
33 obstruction syndrome is common in people with cystic fibrosis.

34 8.17.2.5 Cystic fibrosis related musculoskeletal disorders

35 The Committee noted that in their clinical experience people with cystic fibrosis often have
36 muscle pains and arthralgia. There was no evidence on these complications, therefore the
37 Committee decided to make a recommendation based on their expertise to be aware that
38 muscle pains and arthralgia are common in people with cystic fibrosis.

39 The Committee noted that evidence from the UK CF registry showed that arthritis was rare;
40 they noted that the prevalence varies considerably between the age groups used in the
41 registry (younger than 16 years or 16 years and over) , and reasoned that if additional age
42 cut-offs were used, prevalence would most likely vary between the smaller age sub-groups.
43 Therefore the Committee decided not to specify numerical estimates of prevalence and
44 decided to recommend to be aware that cystic fibrosis-related arthritis is less common than
45 other complications in people with cystic fibrosis.

46 8.17.2.6 Delayed puberty

47 There was some evidence on the prevalence of delayed puberty, however the Committee
48 believed that the evidence was too old to be relevant, because puberty occurs earlier on
49 nowadays. Delayed puberty is related to malnutrition and it occurs in people with severe

1 disease. Therefore, the Committee used their clinical experience to recommend to be aware
2 that delayed puberty is less common than other complications in people with cystic fibrosis
3 and is associated with severe cystic fibrosis.

4 **8.17.2.7 Infertility**

5 There was no evidence on the prevalence of infertility. The Committee used their clinical
6 experience and expertise to recommend to be aware that male infertility and reduced female
7 fertility are common in people with cystic fibrosis. Moreover, the Committee agreed that
8 almost all males with cystic fibrosis are infertile. They mentioned that the vas deferens is the
9 tube that connects the testis to the seminal vesicle. It is lined by mucus, which helps lubricate
10 the passage of sperm along the tube. In CF this mucus is too thick (the CF pathological
11 process) and the vas deferens becomes obstructed during its development in the foetus and
12 subsequently withers away. Thus, most males with CF are born with an absent vas deferens.
13 However, this process is not absolute, and in a very small proportion of males there is still a
14 functioning vas deferens, especially in those who have a lesser expression of the CF
15 condition (a very mild phenotype). Indeed, there are a group of CF males in which CF is only
16 suspected when they are found to have an absent vas deferens as part of investigations for
17 infertility when they are adults. The Committee noted that it was important to take into
18 account the small proportion of males with a functioning vas deferens, especially considering
19 the risk of unwanted pregnancies. Thus, the Committee agreed that the statement “almost all
20 males” was appropriate to use in the recommendations.

21 **8.17.2.8 Upper airways disease**

22 The Committee noted that the evidence from the UK CF registry showed that both the
23 prevalence of nasal polyps requiring surgery and the prevalence of sinus disease increase
24 with age. This age trend reflected their clinical experience. However the Committee also
25 noted that the prevalence of sinus disease and nasal polyps seemed low compared to their
26 clinical experience, and agreed that the data was likely to be based on symptomatic disease
27 only. The Committee also noted that underestimating the prevalence of nasal polyps in the
28 recommendations would lead to dismissal of this complication as minor and rare when in fact
29 early recognition and management can improve quality of life significantly. Therefore they
30 decided not to mention a numerical estimate of prevalence of sinus disease and nasal polyps
31 in the recommendations. The Committee recommended to be aware that upper airway
32 complications, including nasal polyps and sinusitis are common complications of cystic
33 fibrosis (prevalence increases with age).

34 **8.17.2.9 Cystic-fibrosis related diabetes**

35 The Committee noted that the included studies from the published literature were not from
36 the UK and prioritised the data from the UK registry on people on treatment for CFRD
37 because this data was based on the context where NICE guidelines are implemented. The
38 Committee noted that the registry did not have data on people younger than 10 on CFRD
39 treatment because this is uncommon in this age group. The Committee noted that the
40 prevalence was lower in the group of people aged 10 to 16 years (10%) compared to the
41 prevalence among people aged 16 or older (32%). The Committee agreed that the
42 prevalence of people on CFRD treatment was likely to underestimate the prevalence of
43 people with CFRD because many centres do not know how to accurately test for it.
44 Therefore, based on their clinical experience and expertise, the Committee recommended to
45 be aware that cystic fibrosis-related diabetes is common in people with cystic fibrosis
46 (uncommon in children under 10 years, but the prevalence increases with age and it affects
47 up to 1 in 2 adults).

B.17.2.10 Chronic liver disease

2 The evidence from the UK CF registry showed that 9% of people younger than 16 years and
3 18% of people aged 16 years and over have liver disease. The Committee noted that liver
4 disease is a chronic condition that develops slowly and for many it does not majorly impact
5 on overall health. Only a small number of people develop liver failure requiring transplant.
6 The Committee noted that it is rare to diagnose liver disease in adulthood; most people
7 present with abnormal liver function or abnormal ultrasound scan (AUSS) in childhood or
8 early teens. Therefore, the Committee recommended to be aware that chronic liver disease
9 is common in people with cystic fibrosis (the prevalence increases with age until early
10 adulthood).

1 B.17.2.11 Cystic fibrosis related renal disease

12 The Committee noted that the evidence from the UK CF registry was likely to underreport
13 prevalence of renal failure because routine measurements of renal function are not very
14 reliable. Moreover, the Committee decided not to include a recommendation on acute renal
15 failure because this is likely to be related to drug use (aminoglycosides; immunosuppressant
16 medications) and therefore it is debatable whether is a complication of cystic fibrosis (rather
17 than a complication of specific drugs). The Committee noted that there was some evidence
18 from the published literature on the prevalence of chronic kidney disease, however the
19 evidence from 1 study was only on chronic kidney disease of stage 3 or greater, and the
20 evidence from another study was based on records from 1969 to 2009 therefore unlikely to
21 be up to date; therefore the Committee decided not to include a recommendation about the
22 prevalence of chronic kidney disease.

23 The Committee noted that the UK CF registry data showed that the prevalence of kidney
24 stones was higher among people aged 16 years or older (1.5%) compared to people aged
25 less than 16 years (0.3%). The Committee noted that this age trend reflected their clinical
26 expertise. However the Committee agreed that the registry data on kidney stones (1%
27 prevalence for the overall population) was likely to underreport the prevalence of kidney
28 stones because they are often asymptomatic and are only discovered when they cause
29 problems. Based on the Committee's clinical expertise and experience prevalence of kidney
30 stones in adults was about 5%. Therefore, the Committee recommended to be aware that
31 although renal calculi are less common than other complications in people with cystic
32 fibrosis, the incidence increases with age and 1 in 20 adults are affected.

3 B.17.2.12 Urinary stress incontinence

34 There was no evidence on the prevalence of urinary stress incontinence. Therefore the
35 Committee decided not to include a numerical estimate of prevalence in the
36 recommendations and based on their clinical expertise recommended to be aware that
37 urinary stress incontinence is common in people with cystic fibrosis.

3 B.17.2.13 Reduced bone mineral density

39 There was evidence from the UK CF registry on the prevalence of osteopenia, osteoporosis
40 and bone fracture. The Committee agreed that given the low prevalence of bone fractures
41 there was no need to mention these in the recommendations. The Committee noted that the
42 registry was likely to underreport prevalence of osteopenia and osteoporosis as these
43 complications can only be diagnosed by DXA scan for which the take-up of these is not
44 good, so many cases will be undiscovered. There was some evidence from the published
45 literature on the prevalence of osteopenia or osteoporosis in Ireland and Australia; this
46 evidence was based on registries and did not specify how the conditions were diagnosed,
47 however diagnosing practices in these countries are likely to mirror UK practice, with the
48 same limitations leading to underreporting. Given the limitations in the evidence the
49 Committee decided not to include a numerical estimate of prevalence in the

1 recommendations and only recommended to be aware that reduced bone mineral density
2 and osteoporosis are common in people with cystic fibrosis.

3 **8.17.3 Consideration of economic benefits and harms**

4 This was an epidemiological review question and economic analysis to assess cost-
5 effectiveness is not applicable as it does not involve a comparison of competing alternatives.
6 Even so, the assessment, monitoring, referral and management of the comorbidities all have
7 cost implications.

8 The Committee recognised that if complications are not identified and management
9 appropriately, they can negatively impact on wellbeing, function and participation, and
10 increase their risk of further complications leading to additional treatment costs and
11 reductions in quality of life. Therefore, knowing the prevalence of important complications
12 may lead to increased vigilance and thus more timely management and has therefore,
13 indirectly, potentially important resource implications. Estimating the costs to manage those
14 complications would go beyond the scope of the guideline, but it was clear from the
15 Committee that such costs would offset the potential downstream costs from delayed or
16 inappropriate management.

17 **8.17.4 Quality of evidence**

18 This was a prevalence review, therefore the quality assessment with GRADE was not
19 performed. The quality of the studies was assessed with the checklist by Munn et al. 2014.
20 Often it was unclear from the evidence how a condition was diagnosed, and therefore
21 whether objective, standard criteria were used for the measurement of the condition or
22 whether the condition was measured reliably. Moreover, some of the evidence did not
23 disaggregate data by age groups.

24 **8.17.5 Other considerations**

25 The Committee discussed whether people with cystic fibrosis may feel overwhelmed if
26 presented with a long list of possible complications they are at risk of. However the
27 Committee agreed that people with cystic fibrosis have the right to receive all the available
28 information on their condition. Moreover, the Committee noted that the risk for each
29 individual largely depends on management, and this should be made clear in conversations
30 with people with cystic fibrosis.

31 No equality issues were identified by the Committee for this review question.

32 The Committee agreed a research recommendation was not needed. They noted that there
33 was sufficient awareness of both common and rarer complications of cystic fibrosis. It was
34 also felt that there are robust reporting mechanisms (CF Registry) should the prevalence of
35 certain complications change in the future. As people with cystic fibrosis survive longer they
36 are more likely to present with problems associated with organ damage – such as
37 retinopathy or kidney problems in those who have diabetes. However, these are
38 complications that are well established in the wider health-care setting and there are no
39 specific cystic fibrosis-related treatments.

40 **8.17.6 Key conclusions**

41 The guideline developers concluded that health care professionals should explain to each
42 person with cystic fibrosis and their family members or carers (as appropriate) what
43 complications they are at risk of, taking into account each individual's clinical risk factors.
44 Depending on the evidence available for each complication and the individual risk factors, it
45 may be appropriate to either use the terms common or less common, or provide a numerical
46 estimate.

1 8.18 Recommendations

2 **38. Be aware that people with cystic fibrosis are at risk of the following common**
3 **complications:**

- 4 • being underweight
- 5 • meconium ileus (affects 1 in 7 newborn babies)
- 6 • fat-soluble vitamin deficiencies (including vitamins A, D, E and K)
- 7 • distal ileal obstruction syndrome
- 8 • muscle pains and arthralgia
- 9 • male infertility (almost all males with cystic fibrosis are infertile)
- 10 • reduced female fertility
- 11 • upper airway complications, including nasal polyps and sinusitis
- 12 (prevalence increases with age)
- 13 • chronic liver disease (the prevalence increases with age until early
- 14 adulthood)
- 15 • urinary stress incontinence
- 16 • cystic-fibrosis-related diabetes (uncommon in children under 10 years,
- 17 but the prevalence increases with age and it affects up to 1 in 2 adults)
- 18 • reduced bone mineral density and osteoporosis.

19 **39. Be aware that people with cystic fibrosis are at risk of the following less common**
20 **complications:**

- 21 • cystic fibrosis-related arthritis
- 22 • delayed puberty (associated with severe cystic fibrosis)
- 23 • renal calculi (incidence increases with age and 1 in 20 adults are
- 24 affected).

9 Pulmonary monitoring, assessment and management

9.1 Pulmonary monitoring

Review questions:

1) What is the value of the following investigative strategies in monitoring the onset of pulmonary disease in people with CF without clinical signs or symptoms of lung disease?

- Non-invasive microbiological investigation- induced sputum samples, cough swab, throat swab, and nasopharyngeal aspiration
- Invasive microbiological investigation- broncho-alveolar lavage
- Lung physiological function tests- Cardiopulmonary exercise testing, Spirometry and Lung Clearance Index
- Imaging techniques- Chest x-ray and CT scan

2) What is the value of the following investigative strategies in monitoring evolving pulmonary disease in people with established lung disease?

- Non-invasive microbiological investigation- induced sputum samples, cough swab, throat swab, and nasopharyngeal aspiration
- Invasive microbiological investigation- broncho-alveolar lavage
- Lung physiological function tests- Cardiopulmonary exercise testing, Spirometry and Lung Clearance Index
- Imaging techniques- Chest x-ray and CT scan.

3) What is the added value of imaging and invasive microbiological testing in addition to non-invasive microbiological testing and lung function tests in monitoring the response to treatment following an acute exacerbation?

9.1.1 Introduction

Treatment for cystic fibrosis lung disease is based on the prevention of lung infection and subsequent colonisation by pathogenic organisms, long term maintenance therapies to ensure clinical stability and prevent progressive loss of lung function, and treatment of infective exacerbations. It is a condition which requires constant vigilance to monitor disease state with aggressive, early intervention to treat infection.

Infective exacerbations are associated with considerable morbidity and some episodes can lead to permanent reduction in lung function. Treatment response for pulmonary exacerbation is measured by a number of outcome measures, including analysis of non-invasive microbiological specimens, improvement in symptoms, oxygenation, inflammatory markers and pulmonary function. The treatment of an acute exacerbation is closely monitored by cystic fibrosis teams and therapy may be changed depending on the assessment of treatment response.

9.1.2 Description of clinical evidence

The aim of this review was to examine different monitoring strategies for pulmonary disease in people with cystic fibrosis and to determine their impact on improving subsequent

1 intervention and clinical outcomes, therefore the diagnostic accuracy of the different tests
2 was not prioritised for this review. Monitoring techniques were split into 4 categories:

- 3 • Monitoring technique 1: non-invasive microbiological investigation of respiratory tract
4 samples (including induced sputum samples, cough swabs, throat swabs and
5 nasopharyngeal aspiration);
- 6 • Monitoring technique 2: invasive microbiological investigation (i.e. bronchoalveolar lavage
7 - BAL);
- 8 • Monitoring technique 3: pulmonary function tests (including cardiopulmonary exercise
9 testing, spirometry and lung clearance index);
- 10 • Monitoring technique 4: imaging techniques (including chest X-ray and CT scanning).

11 The Committee considered the effects of monitoring considering three clinical scenarios
12 associated with lung disease and corresponding review questions and protocols were drafted
13 for people with cystic fibrosis:

- 14 • Protocol 1: without clinical signs or symptoms of lung disease
- 15 • Protocol 2: with established pulmonary disease
- 16 • Protocol 3: following an acute pulmonary exacerbation

17 The Committee recognised that those with no pulmonary disease would principally, but not
18 exclusively be young children and that this review would inform investigative strategies to
19 identify the onset of pulmonary disease, as opposed to identifying evolving pulmonary
20 disease in the second population. The Committee were interested in comparisons of
21 individual techniques within categories, individual techniques across categories and in
22 combinations of techniques within or across categories. Of particular interest was the
23 effectiveness of imaging techniques in addition to non-invasive microbiological techniques
24 and spirometry.

25 The Committee were specifically interested in the value of adding invasive microbiological
26 investigations and/or imaging techniques to non-invasive microbiological testing and lung
27 function tests to evaluate treatment response in those with an acute pulmonary exacerbation.

28 We aimed to include systematic reviews, test and treat RCTs and prospective and
29 retrospective observational studies.

30 For full details see review protocols in Appendix D.

31 One single literature search was run for the 3 protocols, and 2 studies were included. Neither
32 of these studies absolutely adhered to the clinical scenarios of the protocols.

33 **9.1.2.1 Review 1. Monitoring for pulmonary disease onset in people with cystic fibrosis** 34 **without clinical signs or symptoms of lung disease**

35 One study (Sanders 2015) was identified for this protocol. The authors used registry data to
36 follow up 60 children who were initially recruited to a RCT of pulmozyme. The authors
37 investigated whether chest CT and pulmonary function test scores (taken at the start and end
38 of the RCT) were associated with the rate of pulmonary exacerbations over the subsequent
39 10 year period.

40 **9.1.2.2 Review 2. Monitoring for evolving pulmonary disease in people with cystic fibrosis** 41 **with established lung disease**

42 No studies were identified for this protocol.

1 **9.1.2.3 Review 3. Monitoring for evolving pulmonary disease in people with cystic fibrosis**
2 **following an acute pulmonary exacerbation**

3 One study (Wainwright 2011) was identified for this protocol. This study was a multicentre
4 RCT (ACFBAL) which recruited 170 participants to determine whether monitoring using BAL
5 to direct therapy for pulmonary exacerbations in the first five years of life reduced *P*
6 *aeruginosa* infection and structural lung injury at age 5 years compared with standard
7 management based on clinical features and oropharyngeal culture results.

8 A summary of the studies included in the reviews is presented in Table 60 and Table 61. See
9 also study selection flow chart in Appendix F, study evidence tables in Appendix G, list of
10 excluded studies in Appendix H, and full GRADE profiles in Appendix J.

11 **9.1.3 Summary of included studies**

12 **9.1.3.1 Review 1. Monitoring for pulmonary disease onset in people with cystic fibrosis**
13 **without clinical signs or symptoms of lung disease**

14 A summary of the studies that were included in this review is presented in Table 60.

15 **Table 60: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
Sanders 2015 (USA) Cohort study	<p>Chest CT (Brody score)</p> <ul style="list-style-type: none"> Baseline data: For this study the authors used data obtained at the baseline and at end of the PEIT study (1997-1999) Follow-up data: Data from the time of chest CT in 1999 through 2009 were obtained from the CFFPR and linked to the original chest CT data <p>FEV₁% predicted</p> <ul style="list-style-type: none"> PFTs were obtained on the same day as the chest CT 	<p>N=60 children with CF who participated in the PEIT trial</p> <p>Age: 6 to 10 years</p>	<ul style="list-style-type: none"> Pulmonary exacerbations (proxy outcome for time to next exacerbation)* (pulmonary exacerbation defined as hospitalizations treated with IV AB and/ or if the "pulmonary exacerbation" box was checked in the CFFPR form FEV₁% predicted** <p>* multivariate Poisson model adjusted for sex, genotype, and FEV₁ and mucoid <i>P aeruginosa</i> status at the time of the chest CT</p> <p>**multivariate linear regression model adjusted for sex, genotype, and FEV₁ and mucoid <i>P aeruginosa</i> status at the time of the chest CT</p>	<p>No adjustments for the confounder of concurrent treatment with immunomodulatory and/or mucolytic agents</p>

16 *BAL: bronchoalveolar lavage; CFFPR: cystic fibrosis Foundation Patient Registry; MD: mean difference; SD:*
17 *standard deviation*

1 **9.1.3.2 Review 2. Monitoring for evolving pulmonary disease in people with cystic fibrosis**
2 **with established lung disease**

3 No studies were identified for this protocol.

4 **9.1.3.3 Review 3. Monitoring for evolving pulmonary disease in people with cystic fibrosis**
5 **following an acute pulmonary exacerbation**

6 A summary of the studies that were included in this review is presented in Table 61.

7 **Table 61: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
Wainwright 2011 (Australia & New Zealand) Multicentre RCT	Standard monitoring group Used clinical features and oropharyngeal cultures to direct therapy. BAL monitoring group standard monitoring + BAL performed before 6 months when well, when hospitalised for exacerbations, if <i>P aeruginosa</i> cultured from oropharyngeal specimen and following <i>P aeruginosa</i> eradication therapy. Culture results from the BAL fluid informed treatment decisions	N=168 infants with CF <6 months Mean age (SD): 3.6 (1.6) months • Standard therapy: n=84 • BAL therapy: n=86 • Children were followed until age 5 years	<ul style="list-style-type: none"> • FEV₁ z scores • Weight z scores • Height z scores • BMI z scores 	Indirectness: intervention in BAL monitoring group does not reflect that of current clinical practice

8 *BAL: bronchoalveolar lavage; CFFPR: cystic fibrosis Foundation Patient Registry; MD: mean difference; SD: standard deviation*

10 **9.1.4 Clinical evidence profiles**

11 **9.1.4.1 Review 1. Monitoring for pulmonary disease onset in people with cystic fibrosis**
12 **without clinical signs or symptoms of lung disease**

13 The clinical evidence profiles for this review question are presented in Table 62, Table 63
14 and Table 64.

15 **Table 62: Summary clinical evidence profile: Monitoring technique 3. Lung**
16 **physiological function tests (FEV₁% predicted at baseline) for prognosis of**
17 **pulmonary exacerbations and FEV₁ percent predicted at 10 years**

Prognostic factors	Included studies	Study design	Setting	n	Result (adjRR, MD)	Quality	Notes
Pulmonary exacerbations (defined as hospitalizations treated with IV AB), 10 year follow-up							
FEV ₁ % predicted, 5-point decrease	1 (Sanders 2015)	Cohort study	CF centres in Europe	60	adjRR: 1.19 (95% CI: 1.10 – 1.30)	Moderate ¹	Multiple Poisson model adjusted for sex, genotype, FEV ₁ and mucoid <i>P aeruginosa</i> status at time of chest CT. p value = < 0.001

Prognostic factors	Included studies	Study design	Setting	n	Result (adjRR, MD)	Quality	Notes
Difference in FEV₁ % predicted							
FEV ₁ % predicted, 5-point decrease	1 (Sanders 2015)	Cohort study	CF centres in Europe	60	MD: -4.47 (95% CI: -6.48 to -2.76)	Moderate ¹	Multiple linear model adjusted for sex, genotype, FEV ₁ and mucoid P aeruginosa status at time of chest CT. p value = < 0.001

Abbreviations: adjRR: adjusted rate ratio; CF: cystic fibrosis; CI: confidence interval; CT: computerised tomography; FEV₁: forced expiratory volume in 1 second; MD: mean difference
¹ The quality of the evidence was downgraded by 1 due to no adjustments for the confounder of concurrent treatment with immunomodulatory and/or mucolytic agents.

Table 63: Summary clinical evidence profile: Monitoring technique 4. Chest CT scan for prognosis of pulmonary exacerbations and FEV₁% predicted at 10 years

Prognostic factors	Included studies	Study design	Setting	n	Result (adjRR, MD)	Quality	Notes
Pulmonary exacerbations (defined as hospitalizations treated with IV AB), 10 year follow-up							
Brody chest CT score, 1-point increase	1 (Sanders 2015)	Cohort study	CF centres in Europe	60	adjRR: 1.39 (95% CI: 1.15 – 1.67)	Moderate ¹	Multiple Poisson model adjusted for sex, genotype, FEV ₁ and mucoid P aeruginosa status at time of chest CT. p value = < 0.001
Difference in FEV₁ % predicted, 10 year follow-up							
Brody chest CT score, 1-point increase	1 (Sanders 2015)	Cohort study	CF centres in Europe	60	MD: -4.76 (95% CI: -7.80 to -1.72)	Moderate ¹	Multiple linear model adjusted for sex, genotype, FEV ₁ and mucoid P aeruginosa status at time of chest CT. p value = 0.003

Abbreviations: adjRR: adjusted rate ratio; CF: cystic fibrosis; CI: confidence interval; CT: computerised tomography; FEV₁: forced expiratory volume in 1 second; MD: mean difference
¹ The quality of the evidence was downgraded by 1 due to no adjustments for the confounder of concurrent treatment with immunomodulatory and/or mucolytic agents

Table 64: Summary clinical evidence profile: Comparison 1. FEV₁% predicted versus chest CT scan for prognosis of pulmonary exacerbations and FEV₁% predicted at 10 years

Comparison 1. FEV₁% predicted versus chest CT scan for prognosis of pulmonary exacerbations and FEV₁% predicted at 10 years						
Outcomes	Illustrative comparative risks* (95% CI)		Difference between tests p-value	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	FEV ₁ % predicted, 5% decrease	Brody chest CT score, 1-point decrease				
Pulmonary exacerbations, defined as hospitalization treated with IV AB Follow-up: 10 years	adjRR = 1.19 (95% CI: 1.10 – 1.30) ²	adjRR = 1.39 (95% CI: 1.15 – 1.67) ²	RR = 0.86*, p-value = 0.037 By Chi-Square test ²	60 (Sanders 2015)	⊕⊕⊕ ⊖ moderate ¹	Multiple Poisson model adjusted for sex, genotype, FEV ₁ and mucoid P aeruginosa status at time

Comparison 1. FEV₁% predicted versus chest CT scan for prognosis of pulmonary exacerbations and FEV₁% predicted at 10 years

Change/ decline in FEV ₁ % predicted Follow-up: 10 years	Mean difference: -4.47 (95% CI: -6.48 to - 2.76)	Mean difference: - 4.76 (95% CI: -7.80 to - 1.72)	MD: 0.29*, p-value = 0.4 By F test ²	60 (Sanders 2015)	⊕⊕⊕ ⊖ moderat e ¹	of chest CT. Multiple linear model adjusted for sex, genotype, FEV ₁ and mucoi P aeruginosa status at time of chest CT.
--	--	---	--	-------------------------	---------------------------------------	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: AB: antibiotics; adjRR: adjusted rate ratio; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference

** Calculated by NGA technical team*

adjRR: adjusted rate ratio

1 The quality of the evidence was downgraded by 1 due to no adjustments for the confounder of concurrent treatment with immunomodulatory and/or mucolytic agents

2 This result was reported narratively only

1
2
3
4
5

9.1.4.2 Review 2. Monitoring for evolving pulmonary disease in people with cystic fibrosis with established lung disease

Not applicable, as no studies were included for this review.

9.1.4.3 Review 3. Monitoring for evolving pulmonary disease in people with cystic fibrosis following an acute pulmonary exacerbation

The clinical evidence profile for this review question is presented in Table 65.

Table 65: Summary clinical evidence profile: Comparison 1. BAL monitoring versus standard monitoring

Comparison 1. BAL monitoring versus standard monitoring						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Standard monitoring risk	BAL monitoring				
FEV ₁ z score Better indicated by higher values Follow-up: 5 years	The mean FEV ₁ in the control group was -0.41	The mean FEV ₁ in the intervention group was 0.15 lower (0.58 lower to 0.28 higher)		157 (Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	
Clearance of P aeruginosa following 1 or 2 courses of eradication therapy Better indicated by higher values	907 per 1000	970 per 1000 (871 to 1000)	RR 1.07 (0.96 to 1.2)	82 (Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	

Comparison 1. BAL monitoring versus standard monitoring						
Follow up: 5 years						
Weight z scores Better indicated by higher values Follow-up: 5 years	The mean weight in the control group was -0.21	The mean weight in the intervention group was 0.06 higher (0.21 lower to 0.32 higher)		157 (Wainwright 2011)	⊕⊕⊕⊖ low ^{1,2}	
Height z scores Better indicated by higher values Follow-up: 5 years	The mean height in the control group was -0.19	The mean height in the intervention group was 0.06 higher (0.23 to 0.35 lower)		157 (Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	
BMI z scores, BMI calculated as weight in kg divided by height in meters squared. Better indicated by higher values Follow-up: 5 years	The mean BMI in the control group was 0.01	The mean BMI in the intervention group was 0.02 higher (0.25 lower to 0.3 higher)		157 (Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: BAL: bronchoalveolar lavage; BMI: body mass index; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio</p>						

1 *The quality of the evidence was downgraded by 1 due to serious indirectness as intervention in BAL monitoring group does not reflect that of current clinical practice.*
2
3 *The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed 1 default MID.*

4 9.1.5 Economic evidence

5 One economic evaluation relevant to the protocol was identified in the literature search
6 conducted for this guideline. This study was a prospective cost-benefit analysis undertaken
7 on the RCT by Wainwright (2011) (Section 9.1.2.3). A second study has also been included
8 to aid consideration on the frequency of testing for people with established pulmonary
9 disease. Data extraction tables and quality assessments of included studies can be found in
10 Appendix L and M, respectively. Full details of the search and economic article selection flow
11 chart can be found in Appendix E and F, respectively.

12 This review question was not prioritised for de novo economic modelling. To aid
13 consideration of cost-effectiveness relevant resource and cost use data are presented in
14 Appendix K.

1 **9.1.6 Evidence statements**

2 **9.1.6.1 Review 1. Monitoring for pulmonary disease onset in people with cystic fibrosis**
3 **without clinical signs or symptoms of lung disease**

4 **49.1.6.1.1 Monitoring technique 1. Non-invasive microbiological investigation**

5 No evidence was found.

6 **69.1.6.1.2 Monitoring technique 2. Invasive microbiological investigation**

7 No evidence was found.

8 **89.1.6.1.3 Monitoring technique 3. Lung physiological function tests**

9 **Lung function**

10 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
11 that a 5-point decrease in FEV₁% predicted was associated with a reduction in FEV₁%
12 predicted at 10 years follow-up.

13 **Clearance of the organism from the cultures**

14 No evidence was found for this important outcome.

15 **Pulmonary exacerbations**

16 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
17 that a 5-point decrease in FEV₁% predicted was associated with a higher rate of pulmonary
18 exacerbations during the 10-year follow-up period.

19 **Nutritional parameters**

20 No evidence was found for this important outcome.

21 **Quality of life**

22 No evidence was found for this important outcome.

23 **239.1.6.1.4 Monitoring technique 4. Imaging tests**

24 **Lung function**

25 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
26 that a 1-point increase in Brody chest CT score was associated with a reduction in FEV₁%
27 predicted at 10-year follow-up.

28 **Clearance of the organism from the cultures**

29 No evidence was found for this important outcome.

30 **Pulmonary exacerbations**

31 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
32 that a 1-point increase in Brody chest CT score was associated with a higher rate of
33 pulmonary exacerbations during the 10-year follow-up period.

1 **Nutritional parameters**

2 No evidence was found for this important outcome.

3 **Quality of life**

4 No evidence was found for this important outcome.

5 **59.1.6.1.5 Comparison 1. Lung function tests versus imaging tests**

6 **Lung function**

7 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
8 that there were no differences in the strengths of the association between the Brody chest
9 CT score and FEV₁% predicted in 1999 with FEV₁% predicted in 2009. This result was
10 reported narratively only.

11 **Clearance of the organism from the cultures**

12 No evidence was found for this important outcome.

13 **Pulmonary exacerbations (proxy outcome for time to chronic infection)**

14 Moderate quality evidence from 1 cohort study with 60 children with cystic fibrosis showed
15 that a 1-point difference in the Brody chest CT score was more strongly associated with the
16 rate of pulmonary exacerbations between 1999 and 2009 than a 5% predicted difference in
17 FEV₁% predicted at the time of the chest CT. This result was reported narratively only.

18 **Nutritional parameters**

19 No evidence was found for this important outcome.

20 **Quality of life**

21 No evidence was found for this important outcome.

22 **9.1.6.2 Review 2. Monitoring for evolving pulmonary disease in people with cystic fibrosis**
23 **with established lung disease**

24 No evidence was found for this review.

25 **9.1.6.3 Review 3. Monitoring for evolving pulmonary disease in people with cystic fibrosis**
26 **following an acute pulmonary exacerbation**

27 **279.1.6.3.1 Monitoring strategy 1. Invasive microbiological investigations and/or imaging**
28 **techniques in addition to non-invasive microbiological investigations and/or lung**
29 **function test VERSUS non-invasive microbiological investigations**

30 **Comparison 1. Monitoring using bronchoalveolar lavage (BAL) versus standard**
31 **monitoring**

32 **Lung function**

33 Moderate quality evidence from 1 RCT with 157 infants with cystic fibrosis <6 months
34 showed no clinically significant difference between monitoring using BAL and standard
35 monitoring for FEV₁ z scores at 5 years follow-up.

36 **Clearance of the organism from the cultures**

1 Moderate quality evidence from 1 RCT with 157 infants with cystic fibrosis <6 months
2 showed no clinically significant difference between monitoring using BAL and standard
3 monitoring for clearance of *P aeruginosa* following 1 or 2 courses of eradication therapy at 5
4 years follow-up.

5 **Time to chronic infection**

6 No evidence was found for this important outcome

7 **Nutritional parameters**

8 Low to moderate quality evidence from 1 RCT with 157 infants with cystic fibrosis <6 months
9 showed no clinically significant difference in weight, height and BMI (measured as final z-
10 scores) between monitoring using BAL and standard monitoring at 5 years follow-up.

11 **Quality of life**

12 No evidence was found for this important outcome

139.1.6.3.2 **Monitoring strategy 2. Invasive microbiological investigations and/or imaging techniques in addition to non-invasive microbiological investigations and/or lung function test VERSUS lung function test**

16 No evidence was found for this strategy.

179.1.6.3.3 **Monitoring strategy 3. Invasive microbiological investigations and/or imaging techniques in addition to non-invasive microbiological investigations and/or lung function test VERSUS non-invasive microbiological investigations and lung function test**

21 No evidence was found for this strategy.

22 **9.1.6.4 Economic evidence statements**

23 One cost-benefit analysis (Moodie 2014) on people with cystic fibrosis in Australia and New
24 Zealand found that the additional cost of BAL therapy compared to standard therapy was not
25 offset by reductions in other healthcare expenditure over 5 years. This analysis has minor
26 limitations and is directly applicable given that the type of economic evaluation is unlikely to
27 change the conclusions about cost-effectiveness and all other applicability criteria are met.

28 One cost-consequence analysis (Etherington 2008) on people with cystic fibrosis in the UK
29 over 6 months, found that the number of routine susceptibility tests conducted on
30 *P.aeruginosa* isolates can be reduced to provide cost savings without adversely affecting
31 clinical outcomes. This analysis will be used as indirect evidence as the frequency of testing
32 was not a comparator included in the protocol. This evidence is characterised by potentially
33 serious limitations, including the before and after type study design and lack of detail
34 regarding the costs included and their sources.

35 **9.1.7 Evidence to recommendations**

36 **9.1.7.1 Relative value placed on the outcomes considered**

37 The aim of this review was to examine different monitoring strategies for pulmonary disease
38 in people with cystic fibrosis and to determine their impact on improving subsequent
39 intervention and clinical outcomes. The outcomes selected were different for each clinical
40 scenario:

- 41 • For monitoring for onset of pulmonary disease in people with cystic fibrosis without clinical
42 signs or symptoms of lung disease (review 1), the Committee chose lung function and

1 clearance of the organism from the cultures as critical outcomes for decision making;
2 whereas time to chronic infection, nutritional parameters and quality of life were rated as
3 important. Given that no evidence was found for time to chronic infection, pulmonary
4 exacerbations were considered a proxy outcome.

- 5 • For monitoring for evolving pulmonary disease in people with cystic fibrosis with
6 established lung disease (review 2), the Committee chose lung function and time to next
7 exacerbation as critical outcomes for decision making; whereas time to chronic infection,
8 mortality, nutritional parameters and quality of life were rated as important
- 9 • For monitoring for evolving pulmonary disease in people with cystic fibrosis following an
10 acute pulmonary exacerbation (review 3), the Committee chose lung function, time to next
11 exacerbation and clearance of the organism from as critical outcomes for decision
12 making; whereas time to chronic infection, inflammatory markers, nutritional parameters
13 and quality of life were rated as important.

14 **9.1.7.2 Consideration of clinical benefits and harms**

15 The Committee acknowledged the scarcity of the evidence, and therefore most of the
16 recommendations were based on their clinical expertise and experience, and good practice
17 recommendations.

18 The Committee discussed the recommendations for asymptomatic and symptomatic people,
19 to reflect the different scenarios associated with lung disease (people without clinical signs or
20 symptoms of lung disease and people with established lung disease or people presenting an
21 acute pulmonary exacerbation). A distinction was also made between children and adults,
22 were appropriate.

23 **Asymptomatic people**

24 The Committee noted that asymptomatic adults should have regular annual review including
25 clinical examination, oxygen saturation test, spirometry, chest X-ray, microbiological
26 investigations with sampling and culture of respiratory tract secretions for early asymptomatic
27 infection with cystic fibrosis pathogens, and blood testing to include white cell count and
28 markers or aspergillus, including aspergillus serology and serum IgE.

29 The Committee considered annual chest X-rays to be justified due to the low radiation dose
30 (particularly compared with CT scan), and the benefit of comparing serial films which may
31 pick up changes representing development or progression of lung disease before symptoms
32 develop. This aligned with the CF Trust consensus recommendations that recommends a
33 regular (annual) chest radiograph.

34 The Committee agreed that conducting microbiological tests was also very useful. This was
35 because detection of early infection is of key importance in cystic fibrosis, so that where
36 indicated, eradication therapy can be instituted promptly with the aim to prevent chronic
37 infection. Problematic pathogens may be found even in asymptomatic patients, and the
38 Committee consider knowledge of infection status essential for infection control purposes.

39 The Committee considered blood tests for aspergillus (specifically IgE and precipitins) to be
40 necessary to investigate for the presence of allergic broncho-pulmonary aspergillosis
41 (ABPA), a common complication of cystic fibrosis lung disease.

42 The usefulness of CT scan in this population was also discussed by the Committee. They
43 argued that the scan is a much more sensitive way of showing bronchiectasis than a plain
44 film, where it has to be fairly developed to see it at all. So in asymptomatic children where the
45 lungs are thought to be pretty healthy, a CT scan may find early bronchiectasis allowing
46 escalation of treatment, and therefore to prevent further deterioration. In addition, a CT scan
47 will also show other changes (e.g. mucus plugging, allowing targeting of physiotherapy).
48 They also noted that a CT scan gives a better idea of the structure of the lungs, and, as for

1 children, CT scans can be used to quantify deterioration in the future in adults. Given the
2 increased dose of radiation exposure associated to CT scans (compared with X-ray, for
3 example), the Committee agreed to recommend performing a baseline CT scan in
4 asymptomatic adults only when this has not previously been carried out.

5 Finally the Committee agreed that lung function testing should also be part of the annual
6 review, as they are useful to detect any deterioration in the lung function.

7 **Symptomatic people**

8 They noted that symptomatic adults should be reviewed at least every 3 months and should
9 have microbiological cultures and spirometry at each encounter. The rationale for
10 recommending these tests at each encounter in those with lung disease is that the objective
11 of the routine reviews (from the lung perspective) is to prevent deterioration in lung function
12 (which is measured by spirometry). Obtaining microbiological cultures helps to ensure that if
13 the cause of any such deterioration is an infection treatment can be tailored accordingly. This
14 aligned with the CF Trust consensus recommendations, that recommends that frequent (at
15 every clinic visit) microbiological surveillance of respiratory secretions should be undertaken
16 (e.g. cough swab, sputum culture, and induced sputum).

17 The Committee also noted that lung clearance index can be a useful tool to assess disease
18 progression, and the suggested it could be used in people in whom spirometry is normal.

19 Additionally, the Committee agreed that the annual review should include the same
20 investigations as for asymptomatic adults.

21 The Committee noted that more frequent assessment of symptomatic children may be
22 necessary to ensure resolution of symptoms. It was also noted that in children who are
23 responding poorly to treatment, and in the absence of identification of pathogens from cough
24 swabs and induced sputum, more invasive procedures including bronchoalveolar lavage
25 (BAL) or CT scanning can be considered.

26 Although the included study did not show clinically significant differences between the BAL-
27 directed therapy group and the standard group, BAL is still considered the gold standard.
28 The Committee discussed that it is likely that children allocated to the standard group did
29 actually receive BAL when they experience an exacerbation, which could explain the lack of
30 differences between both groups.

31 As evidence was found to suggest correlation between CT score and prognosis, CT may be
32 a useful to monitor disease progression. Where CT scores suggest worsening prognosis, a
33 more aggressive treatment approach may be required to limit or reverse deterioration and
34 improve the prognosis. Based on this, the Committee thought doing a chest CT scan for
35 children with cystic fibrosis could be useful, as it helps to monitor disease progression.

36 **Acute exacerbations**

37 The Committee noted that those with acute exacerbations need to have a separate, defined
38 protocol for monitoring during the exacerbation. As part of this monitoring process,
39 individuals with exacerbations should undergo clinical assessment, microbiological
40 investigations (sputum or cough swab for cystic fibrosis pathogens including selective media)
41 and spirometry.

42 The Committee noted that usually in practice, a chest X-ray is performed if FEV1 drops by
43 10% or more, although treatment of an exacerbation may be provided without reference to
44 an X-ray. The Committee therefore recommended that performance of X-ray for acute
45 exacerbation should be considered dependent on severity of the exacerbation, symptoms
46 (for example where there is suspicion of a pneumothorax), or where there is an element of

1 treatment failure. Where new radiological abnormalities are present on X-ray, this should be
2 repeated to confirm resolution following treatment.

3 The Committee noted that during and following an exacerbation, response to treatment
4 should be assessed using spirometry and microbiological investigation, time to next
5 exacerbation and patient reported outcomes.

6 As for asymptomatic adults, culture of respiratory secretions for early identification of
7 microbial pathogens is important to allow the most appropriate antibiotics to be selected in
8 line with good antibiotic stewardship. An acute exacerbation may be the initial presentation
9 of a newly acquired pathogen, so as for asymptomatic adults, it is important that any new
10 organisms are detected early to allow eradication to be attempted.

11 They also agreed lung function tests can be useful to assess response to treatment. They
12 noted height is necessary for accurate calculation of spirometric indices and should therefore
13 be confirmed when spirometry is undertaken where this has not recently been established.

14 They considered non-invasive oxygen saturation testing (pulse oximetry) to be part of the
15 clinical assessment, justifying the recommendation without specific evidence.

16 The Committee noted that inflammatory markers are helpful to indirectly determine lung
17 damage, and monitor response to treatment and as this is not specific to cystic fibrosis, the
18 Committee felt it was justified to include reference to this test in this section.

19 **9.1.7.3 Consideration of economic benefits and harm**

20 Spirometry was the cheapest lung function test under consideration and the Committee
21 noted that the accuracy of spirometry is demonstrated in both clinical practice and the study
22 by Sanders 2015. For these reasons, the Committee agreed spirometry was cost-effective
23 and should continue to be used to monitor for pulmonary disease at each clinic visit.

24 The Committee also considered a place for LCI investigations at the annual review given that
25 the additional respiratory information resulting from a LCI investigation compared to
26 spirometry justifies the additional cost of LCI. However, they noted that LCI is currently in its
27 infancy in the UK and although it is a promising investigative technique, its application is
28 currently limited to research rather than routine clinical practice.

29 The Committee noted that the intense monitoring schedules proposed during the first year of
30 identification (4-weekly) would put a strain on cystic fibrosis clinics. Moreover, face-to-face
31 contact at a clinic would be burdensome on the person with cystic fibrosis, and subject to
32 availability which may be too late during an exacerbation. For these reasons, the Committee
33 advised that visits could be performed outside of the clinic, as either home-visits or
34 telemedicine, where this is considered appropriate.

35 The Committee acknowledged that a CT scan costs considerably more than a chest X-ray
36 and involves greater exposure to radiation. However they believed a CT scan would show
37 subtle structural changes in the lungs that would not be evident from a chest X-ray, such as
38 early bronchiectasis. As a result, a CT scan could allow early escalation of treatment to
39 prevent further deterioration that could be more costly to treat. Furthermore, the Committee
40 noted that the accuracy of CT scans to predict pulmonary exacerbations was demonstrated
41 in the study by Sanders 2015. For these reasons, the Committee concluded they could justify
42 the use of CT scans as a cost-effective use of NHS resources and made a recommendation
43 to think about doing a chest CT scan in people who have not had one before.

44 The Committee considered the high cost of BAL and agreed that despite the lack of clinical
45 evidence in favour of BAL, an annual BAL in children would be more informative than several
46 non-invasive investigations throughout the year. The Committee also added that BAL is
47 considered as the gold standard test in clinical practice. However, combined with the high
48 cost of BAL and the potential adverse effects, the Committee agreed BAL could only be

1 considered cost-effective in symptomatic people with cystic fibrosis, when cheaper and less
2 invasive investigations such as sputum induction had been unsuccessful.

3 Monitoring for multi-resistant organisms was discussed by the Committee and they agreed it
4 would not be cost-effective use of resources to monitor for those organisms if there was not
5 an effective treatment for the organisms that are identified.

6 **9.1.7.4 Quality of evidence**

7 There was very little evidence available to inform these 3 reviews. Although many studies
8 examined monitoring techniques, only 2 studies were relevant to the protocol and presented
9 clinical outcomes, rather than, diagnostic outcomes or estimates of correlation between
10 monitoring techniques.

11 Only 1 prospective cohort study was available to inform the first protocol which focussed on
12 people with cystic fibrosis without clinical signs or symptoms of lung disease. This study
13 included children who had previously been participants in a randomised controlled trial. One
14 of the criteria for inclusion to the trial was having a FVC of 85% predicted or greater. The
15 participants actually had a mean FEV₁ of 99% predicted at the end of the trial which was
16 similar to the national (United States) average for children without cystic fibrosis and on this
17 basis the study was included to inform the first protocol.

18 The GRADE quality of this prognostic data was moderate, downgraded from high as the
19 adjustment of the rate ratios did not incorporate the potential confounders of concurrent
20 treatment with immunomodulatory and/or mucolytic agents during the 10 year follow up
21 period.

22 There was no available evidence that examined the effectiveness of different combinations of
23 monitoring techniques in adults or in children without clinical signs or symptoms of lung
24 disease.

25 Neither was there any available evidence that examined the effectiveness of different
26 combinations of monitoring techniques in adults or in children with established pulmonary
27 disease, nor any prognostic data for this clinical scenario.

28 The RCT which informed the third protocol did compare monitoring strategies, however, the
29 included population was aged from under 6 months to 5 years. The relevance of the
30 intervention to use BAL to direct therapy is questionable as this would not reflect current
31 clinical practice and the GRADE quality of outcomes was downgraded accordingly.
32 Moreover, the generalisability of the results to adults is questionable.

33 **9.1.7.5 Other considerations**

34 The Committee discussed potential equality issues. They noted that young people who live
35 far from a specialist centre may be disadvantaged. However they agreed no additional
36 recommendations were needed, as the use of alternative models care had already
37 extensively been discussed in the service delivery review. See section 7.1.

38 The Committee discussed the need to draft a research recommendation for this topic. They
39 agreed it would be important to evaluate if Lung Clearance Index (LCI) was a useful tool for
40 routine assessment and monitoring for changes in pulmonary status in people with cystic
41 fibrosis. This is because assessing the severity of lung disease is difficult in younger children,
42 as not all children under 5 years can do spirometry tests and they are not sufficiently
43 sensitive in people with good lung function, where CT scans can show pulmonary status
44 changes before spirometry changes. A simple, sensitive and reproducible measurement
45 such as lung clearance index allows assessment of respiratory status in people with cystic
46 fibrosis, and could improve clinical decision-making.

1 **9.1.7.6 Key conclusions**

2 The Guideline Committee concluded that monitoring for the onset and evolution of pulmonary
3 disease is key to be able to treat early infections.

4 They agreed that it is important to conduct regular routine reviews to both children and adults
5 with cystic fibrosis even if they are asymptomatic, and that these reviews should be more
6 frequent in early life. The Committee agreed reviews can be conducted more often if
7 necessary based on clinical judgement. During these routine reviews, it is important to carry
8 out a clinical assessment, conduct non-invasive microbiological investigations, and
9 pulmonary function tests. They also agreed it is useful to do a chest CT scan for all children
10 before the age of 12, even in the absence of lung disease. Likewise, they agreed it is
11 important to perform a baseline CT scan for cystic fibrosis people diagnosed in adulthood.

12 With regards to children who are symptomatic, the Committee agreed on recommending the
13 use of invasive microbiological investigations (such as BAL) when the cause of the disease
14 cannot be found using non-invasive microbiological tests, or if there is no response to
15 treatment.

16 Finally, the Committee also agreed that it is important to monitor the response to treatment
17 during and after a pulmonary exacerbation by assessing whether symptoms have resolved,
18 conducting microbiological investigations, and pulmonary function tests.

19 **9.1.8 Recommendations**

20 **40. For people with cystic fibrosis who have lung disease, base the frequency of**
21 **routine reviews on their clinical condition but review children and young people at**
22 **least every 8 weeks and adults at least every 3 months. If appropriate, think about**
23 **using the review schedules in the recommendation on routine reviews.**

24 **41. Include the following at each routine review, in relation to pulmonary assessment,**
25 **for people with cystic fibrosis:**

- 26
- 27 • a clinical assessment, including a review of the clinical history and a
28 physical examination, with measurement of weight and length (if less
29 than 2 years old) or height
 - 30 • measurement of oxygen saturation
 - 31 • taking respiratory secretion samples for microbiological investigations,
32 using sputum samples if possible, or a cough swab or nasal pharyngeal
33 aspirate (NPA) if not
 - 34 • lung function testing with spirometry (including forced expiratory volume
35 in 1 second [FEV₁], forced vital capacity [FVC], and forced expiratory
36 flow [FEF] 25–75%) in adults, and in children and young people who can
do this.

37 **42. Consider measuring lung clearance index at each routine review if spirometry is**
38 **normal.**

39 **43. Include the following at each annual review in relation to pulmonary assessment**
40 **for people with cystic fibrosis:**

- 41
- 42 • a clinical assessment, including a review of the clinical history and a
43 physical examination, with measurement of weight and length (if less
44 than 2 years old) or height
 - 45 • a physiotherapy assessment
 - measurement of oxygen saturation

- 1 • a chest X-ray
 - 2 • blood tests, including white cell count, aspergillus serology and serum
 - 3 IgE
 - 4 • taking respiratory secretion samples for microbiological investigations
 - 5 (including mycobacteria)
 - 6 • lung function testing (for example with spirometry, including FEV₁, FVC,
 - 7 and FEF 25–75%) in adults, and in children and young people who can
 - 8 do this, and
- 9 **44. Consider measuring lung clearance index at each annual review if spirometry is**
- 10 **normal.**
- 11 **45. For people with cystic fibrosis with lung disease who have symptoms that are**
- 12 **concerning them, their family members or carers (as appropriate) or a healthcare**
- 13 **professional, decide whether a remote telemedicine (see recommendation 17) or**
- 14 **face-to-face assessment is needed, and consider which of the following may be**
- 15 **useful:**
- 16 • review of the past history
 - 17 • physical examination, including measurement of weight and length (if
 - 18 less than 2 years old) or height
 - 19 • measurement of oxygen saturation
 - 20 • collection of respiratory secretion samples for microbiological
 - 21 investigations, using sputum samples if possible, or a cough swab or
 - 22 NPA if not
 - 23 • for adults, blood tests to measure white cell count and inflammatory
 - 24 markers such as C-reactive protein
 - 25 • lung function testing, for example with spirometry (including FEV₁, FVC,
 - 26 and FEF 25–75%) in adults, and in children and young people who can
 - 27 do this
 - 28 • lung clearance index for people with normal spirometry results.
- 29 **46. Think about doing a chest CT scan for children with cystic fibrosis who have not**
- 30 **had one before, to detect features that other tests would miss (for example early**
- 31 **bronchiectasis).**
- 32 **47. Think about doing a chest X-ray for people with cystic fibrosis who have had**
- 33 **treatment for an exacerbation of lung disease (taking account of severity), if:**
- 34 • the exacerbation does not respond to treatment or
 - 35 • a chest X-ray before treatment showed new radiological abnormalities.
- 36 **48. Monitor the treatment response during and after an exacerbation of lung disease**
- 37 **by assessing whether the symptoms and signs have resolved, and as appropriate:**
- 38 • take respiratory secretion samples for microbiological investigations,
 - 39 using sputum samples if possible, or a cough swab or NPA if not
 - 40 • test lung function, for example with spirometry (including FEV₁, FVC and
 - 41 FEF 25–75%) in adults, and in children and young people who can do
 - 42 this
 - 43 • measure oxygen saturation.

49. Think about using broncho-alveolar lavage to obtain airway samples for microbiological investigation in children and young people if:

- they have lung disease that has not responded adequately to treatment and
- the cause of the disease cannot be found with non-invasive upper airway respiratory secretion sampling.

9.1.9 Research recommendations

1. Is Lung Clearance Index (LCI) a useful and cost-effective tool for the routine assessment and monitoring of changes in pulmonary status in people with cystic fibrosis?

Table 66: Research recommendation rationale

Research question	Is Lung Clearance Index (LCI) a useful and cost-effective tool for the routine assessment and monitoring of changes in pulmonary status in people with cystic fibrosis?
Why this is needed	
Importance to 'patients' or the population	Lung clearance index (LCI) measured by multiple-breath washout (MBW) is a more sensitive test of impaired lung function compared to FEV ₁ , particularly in those with relatively normal pulmonary function. It may be a tool to help decisions on early intervention with treatment in children and adults with cystic fibrosis.
Relevance to NICE guidance	Validation of this tool for clinical care could change some recommendations for disease monitoring and treatment intervention. This is of particular relevance in pre-school children where FEV ₁ is not reliably performed. High: the research is essential to inform future updates of key recommendations in the guideline
Relevance to the NHS	If validated and adopted training in Physiology labs and new equipment would be required. In pre-school children it may identify those who require increased intervention, and those who potentially need less regular monitoring and less aggressive therapy. Targeted treatment may potentially have cost savings.
National priorities	No document identified
Current evidence base	FEV ₁ measurements are difficult to perform in preschool infants. FEV ₁ measurements often do not become abnormal until significant lung disease is established.
Equality	This test is of particular relevance to preschool children.
Feasibility	The proposed research can be carried out within a realistic timescale and at an acceptable cost. There are no ethical or technical issues.
Other comments	None

Table 67: Research recommendation statements

Criterion	Explanation
Population	Children with cystic fibrosis, up to 6 years old (pre-school children)
Intervention	• LCI, measure by multiple breath washout (MBW) testing at regular intervals
Comparators	Alternative lung function test: FEV ₁ (measured by spirometry)
Outcomes	<ul style="list-style-type: none"> • Change in lung function • Quality of life as measured by a validated tool, such as CF-QOL or CFQ-R) • Nutritional parameters (BMI, weight/ height) • Time to chronic infection • Resource use

Criterion	Explanation
	• Unit costs
Study design	RCT or cluster RCT
Timeframe	Within 2 years

1 9.2 Airway clearance techniques

2 **Review question: What is the effectiveness of airway clearance techniques in people**
3 **with cystic fibrosis?**

4 9.2.1 Introduction

5 The genetic defect in cystic fibrosis results in the dehydration of mucus, causing increased
6 viscosity and resultant difficulty in its clearance from the airways. Assisted airway clearance
7 has therefore featured in the treatment routines of people with cystic fibrosis for many
8 decades. A variety of techniques have been developed; some require equipment, some rely
9 on the assistance of others and some facilitate independence. The utility of each technique
10 depends upon the degree and extent of the pathophysiology within the lungs.

11 The role of the specialist cystic fibrosis physiotherapist is to evaluate this pathophysiology,
12 and in collaboration with the person with cystic fibrosis select the best airway clearance
13 technique that will overcome these challenges. The aim of treatment is not only to improve
14 the removal of bronchopulmonary secretions and reduce the risk of bacterial infection, but to
15 also reduce the burden of symptoms such as cough and breathlessness and ultimately slow
16 disease progression. Airway clearance techniques are often employed as part of a wider
17 airway treatment strategy which may also include mucolytic or anti-inflammatory drugs and
18 exercise. As a result, measuring the impact or success of airway clearance techniques alone
19 is not without difficulty.

20 Airway clearance approaches are individualised and physiologically reasoned and therefore
21 need to be frequently reviewed and modified by the physiotherapist according to the evolution
22 of lung disease. With the advent of newborn screening and the success of new medical
23 treatments, the person with cystic fibrosis who has little or no lung disease also needs careful
24 consideration. Airway clearance routines for the person with cystic fibrosis and their family or
25 carers can be considered a significant commitment and burden to achieving 'normal' life. It is
26 therefore essential that cystic fibrosis teams continue to question what is understood about
27 these frequently used techniques.

28 9.2.2 Description of clinical evidence

29 The aim of this review was to examine the effectiveness of airway clearance techniques in
30 people with cystic fibrosis.

31 The interventions reviewed were: manual physiotherapy techniques (including chest shaking
32 or vibrations, chest percussion); positive expiratory pressure (PEP) mask therapy; active
33 cycle of breathing techniques (ACBT); relaxation or breathing control forced expiration
34 techniques (FET) which includes huffing and breathing control; thoracic expansion exercises;
35 autogenic drainage (AD); oscillating devices (including acapella and flutter, cornet); high
36 frequency chest wall oscillation (e.g. the Vest) and non-invasive ventilation (NIV).

37 We aimed to compare each airway technique with no intervention, to ascertain effectiveness.
38 In addition, the Committee prioritised for inclusion the following comparisons between
39 techniques:

- 40 • Manual physiotherapy techniques versus oscillating devices (OD)
- 41 • Manual physiotherapy versus high frequency chest wall oscillation (HFCWO)

- 1 • Positive expiratory pressure mask (PEP) versus active cycle of breathing techniques
2 (ACBT)
- 3 • Positive expiratory pressure mask (PEP) versus oscillating devices (OD)
- 4 • Positive expiratory pressure mask (PEP) versus high frequency chest wall oscillation
5 (HFCWO)
- 6 • Active cycle breathing technique (ACBT) versus autogenic drainage (AD)
- 7 • Oscillating device (OD) versus high frequency chest wall oscillation (HFCWO).

8 We searched for systematic reviews of RCTs and RCTs aimed at assessing the
9 effectiveness of airway clearance techniques in people with cystic fibrosis. Observational
10 studies were not prioritised for inclusion in the review, as there was enough evidence from
11 published randomised trials.

12 For full details see review protocol in Appendix D.

13 Six Cochrane systematic reviews (McIlwaine 2015, Morrison 2014, Moran 2013, McKoy
14 2012, Main 2005, Warnock 2013) and 5 non-Cochrane systematic reviews (Boy 1994,
15 Bradley 2006, Flume 2009, Morgan 2015, Thomas 1995) were identified in our search for
16 potential inclusion. The quality of all reviews was assessed with AMSTAR.

17 All the Cochrane reviews obtained a total score equal or higher than 10 (out of 11) in the
18 AMSTAR quality checklist and were considered for inclusion. Each Cochrane review was
19 then checked for relevant potential comparisons, as the definitions of the airway clearance
20 techniques differ from the definitions proposed in our protocol.

21 Four Cochrane reviews were included, as they had relevant comparisons. Where possible,
22 data and quality assessment were extracted from the Cochrane reviews, although the
23 individual studies were retrieved full text for additional information and results.

24 The Cochrane reviews included were:

- 25 • McIlwaine (2015) evaluated the effectiveness and acceptability of PEP devices compared
26 to other physiotherapy techniques to improve airway clearance. 4 trials were included
27 from this review (McIlwaine 2013, McIlwaine 2001, Newbold 2005, Tannenbaum 2005), of
28 which 1 (Tannenbaum 2005) was an abstract and only information provided by the
29 Cochrane review was used as the paper could not be retrieved.
- 30 • Moran (2013) compared the effect of non-invasive ventilation versus no treatment in
31 people with cystic fibrosis. 2 trials were included from this review (Placidi 2006 and Young
32 2008).
- 33 • Morrison (2014) evaluated whether oscillatory devices, oral or chest wall were effective for
34 airway clearance and also compared them with other forms of airway clearance for the
35 management of secretions in people with cystic fibrosis. Seven trials were included from
36 this review (Darbee 2005, Grzincich 2008, Padman 1999, Oermann 2001, Homnick 1998,
37 van Winden, Warwick 2004), of which 1 was an abstract (Grzincich 2008)
- 38 • Warnok (2013) evaluated the effectiveness and acceptability of chest physiotherapy
39 compared to no treatment to improve airway clearance in people with cystic fibrosis. 1 trial
40 was included from this review (Braggion 1995).

41 Three Cochrane reviews were excluded:

- 42 • McKoy (2012) evaluated the clinical effectiveness of ACBT with other airway clearance
43 techniques, but it did not include comparisons relevant for this review. The individual
44 studies were also checked for potential inclusion, but none of them met the criteria for
45 inclusion.
- 46 • Main (2005) compared conventional chest physiotherapy to other airway clearance
47 techniques for cystic fibrosis, but it did not include comparisons relevant for this review.
48 The individual studies were also checked for potential inclusion, and only 1 was

1 considered relevant (Homnick 1998), but it had already been included in another review
2 (Morrison 2014).

3 • Robinson (2010) compared the clinical effectiveness of ACBT with other airway clearance
4 techniques. One trial (Miller 1995) was considered for inclusion, but the intervention
5 consisted in a combination of ACBT and postural drainage.

6 The quality of the non-Cochrane reviews obtained scores equal to or lower than 6 (out of 11)
7 in the AMSTAR checklist and were therefore excluded. The lists of included studies in these
8 reviews were checked in order to identify other studies that had not been already included,
9 but none of them met the inclusion criteria in our protocol.

10 No further trials were identified in our search.

11 The size of the studies ranged from 8 to 107 people with cystic fibrosis. Four studies
12 included adults (Grzincich, Newbold 2005, Young 2008, Warwick 2004), 2 studies included
13 young people and adults (Braggion 1995, Placidi 2006), 4 studies included children and
14 young people (McIlwaine 2001, Padman 1999, Tannenbaum 2005, van Winden 1998), 4
15 studies included children, young people and adults (Darbee 2005, Homnick 1998, McIlwaine
16 2013, Oermann 2001)

17 One study was conducted in Italy (Braggion 1995), 5 studies in the USA (Darbee 2005,
18 Homnick 1998, Oermann 2001, Padman 1999, Warwick 2004), 2 studies in Australia (Placidi
19 2006, Young 2008), 3 studies in Canada (McIlwaine 2001, McIlwaine 2013, Newbold 2005),
20 1 study in the Netherlands (van Winden 1998); the country was not reported in 2 abstracts
21 (Grzincich 2008, Tannenbaum 2005).

22 The included studies assess the effectiveness and acceptability of airway clearance
23 interventions with the following comparisons:

- 24 • manual physiotherapy techniques versus oscillating devices – 2 studies (Homnick 1998,
25 Padman 1999)
- 26 • manual physiotherapy versus high frequency chest wall oscillation (HFCWO) – 1 study
27 (Warwick 2004)
- 28 • positive expiratory pressure (PEP) mask versus no airway clearance - 2 studies (Braggion
29 1995, Placidi 2006)
- 30 • positive expiratory pressure (PEP) mask versus oscillating devices – 6 studies (McIlwaine
31 2001, McIlwaine 2013, Newbold 2005, Padman 1999, Tannenbaum 2005, van Winden
32 1998)
- 33 • positive expiratory pressure (PEP) mask versus high frequency chest wall oscillation
34 (HFCWO) – 4 studies (Braggion 1995, Darbee 2005, Grzincich 2008, McIlwaine 2013)
- 35 • oscillating device versus high frequency chest wall oscillation (HFCWO) - 1 study
36 (Oermann 2001)
- 37 • non-invasive ventilation versus no airway clearance technique – 1 study (Young 2008).

38 Where no evidence for sputum volume was found in the study, sputum weight (both dry and
39 wet) was taken as a proxy outcome for sputum volume.

40 Evidence from these are summarised in the clinical GRADE evidence profile below (Table
41 62Table 69 to Table 75). See also the study selection flow chart in Appendix F, forest plots in
42 Appendix I, study evidence tables in Appendix G and exclusion list in Appendix H.

43 **9.2.3 Summary of included studies**

44 A summary of the studies that were included in this review are presented in Table 68.

1

Table 68: Summary of included studies

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Mcllwaine 2015 Cochrane SR	Positive Expiratory Pressure (PEP) mask vs High Frequency Chest Wall Oscillation (HFCWO) (Mcllwaine 2013) Positive expiratory pressure (PEP) mask vs oscillating device (Mcllwaine 2001, Newbold 2005, Tannenbaum 2005)	People with CF of any age with any degree of disease severity. People with CF post-lung transplant were excluded.	<ul style="list-style-type: none"> • PEP vs HFCWO • Sputum weight (dry and wet) PEP vs oscillating device: <ul style="list-style-type: none"> • Patient preference (self-withdrawal due to lack of effectiveness) • Hospitalisations for respiratory exacerbation • Number of participants experiencing respiratory exacerbations • Lung function • Quality of life • Not reported: • Oxygen saturation 	
Moran 2013 Cochrane SR	Overnight non-invasive ventilation (NIV) vs no airway clearance technique (room air) (Young 2008) Positive Expiratory Pressure (PEP) mask vs no airway clearance (directed cough) (Placidi 2006)*	People with CF of any age with any type of acute and chronic respiratory failure.	NIV vs no airway clearance technique: <ul style="list-style-type: none"> • Lung function • Oxygen saturation (nocturnal) • Quality of life PEP vs no airway clearance technique: <ul style="list-style-type: none"> • Sputum weight • Lung function • Oxygen saturation • Not reported: • Patient preference • Pulmonary exacerbations • Hospitalisations 	*Comparison not in Cochrane SR; comparison made by the NGA technical team using data from controls in comparisons included in Cochrane SR.
Morrison 2014 Cochrane SR	Positive Expiratory Pressure (PEP) mask vs High Frequency Chest Wall Oscillation	Children, young people and adults with CF with any degree of disease severity.	PEP vs HFCWO: <ul style="list-style-type: none"> • Sputum volume • Lung function PEP vs oscillating device:	

Study	Intervention/Comparison	Population	Outcomes	Comments
	(HFCWO) (Braggion 1995, Derbee 2005, Grzincich 2008) Positive expiratory pressure mask vs oscillating device (flutter) (Padman 1999, van Winden 1998) Oscillating device vs High Frequency Chest Wall Oscillation (Oermann 2001) Manual physiotherapy techniques vs oscillating devices (flutter) (Homnick 1998, Padman 1999) Manual physiotherapy vs High Frequency Chest Wall Oscillation (Warwick 2004)		<ul style="list-style-type: none"> • Lung function Oscillating device vs HFCWO: <ul style="list-style-type: none"> • Lung function Manual techniques vs HFCWO: <ul style="list-style-type: none"> • Sputum weight. Manual techniques vs oscillating device: <ul style="list-style-type: none"> • Lung function • Not reported: • Patient preference • Pulmonary exacerbations • Hospitalisations • Oxygen saturation (nocturnal) • Quality of life 	
Warnock 2013 Cochrane SR	Positive expiratory pressure (PEP) mask vs no airway clearance technique (Braggion 1995)	People with CF of any age.	PEP vs control: <ul style="list-style-type: none"> • lung function test • Not reported: • Expectorated secretions • Sputum volume • Patient preference • Pulmonary exacerbations • Hospitalisations • Oxygen saturation (nocturnal) • Quality of life 	
Primary studies included in the Cochrane SR				
Braggion 1995 (Italy) RCT, crossover design	Intervention 1: Positive expiratory pressure (PEP) mask <ul style="list-style-type: none"> • 50-minute session (unclear if this 	N=16 young people and adults with CF Mean age (SD): 20.3 (4) years. Age range: 15 to 27	<ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted 	Included in Morrison 2014 SR and in Warnock 2013 SR 15 minutes of saline nebulised prior to treatment.

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>included 15 mins of nebulisation)</p> <p>Intervention 2: High Frequency Chest Wall Oscillation (HFCWO)</p> <ul style="list-style-type: none"> • 50-minute session (unclear if this included 15 mins of nebulisation) <p>Control: no airway clearance</p>			2 treatments per day for 2 days, then rest 1 day. Next intervention for 2 days, then rest 1 day. Then the final intervention.
Darbee 2005 (USA) RCT, crossover design	<p>Intervention 1: Positive expiratory pressure (PEP) mask</p> <ul style="list-style-type: none"> • 30-minute sessions <p>Intervention 2: High Frequency Chest Wall Oscillation (HFCWO)</p> <ul style="list-style-type: none"> • 30-minute sessions 	<p>N=15 children, young people and adults with CF</p> <p>Age ≥ 7 years.</p> <p>Mean (SD) age: 17.5 (4.2)</p> <p>Participants were admitted to hospital for acute exacerbation. All participants performed HFCWO 1 - 3 times daily as outpatients before admission, but none had performed PEP</p>	<ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted 	<p>Included in Morrison 2014 SR</p> <p>Both treatments were alternated within 48 hours of hospital admission and then reversed prior to discharge.</p>
Grzincich 2008 (Country not reported) RCT, unclear if crossover design	<p>Intervention 1: Positive expiratory pressure (PEP) mask</p> <ul style="list-style-type: none"> • 30-minute sessions <p>Intervention 2: High Frequency Chest Wall Oscillation (HFCWO)</p> <ul style="list-style-type: none"> • 30-minute sessions 	<p>N=23 adults with CF</p> <p>Mean age: 25 years.</p> <p>People hospitalized for an exacerbation</p>	<ul style="list-style-type: none"> • Sputum volume 	<p>Included in Morrison 2014 SR</p> <p>Interventions and control implemented during the first 3 days of hospitalisation for an exacerbation</p> <p>Abstract only</p>
Homnick 1998 (USA) RCT, crossover design	<p>Intervention 1. Manual physiotherapy</p> <ul style="list-style-type: none"> • 30-minute sessions, 4 times daily 	<p>N=22 children, young people and adults with CF (33 hospitalisations)</p> <p>Mean (range)</p>	<ul style="list-style-type: none"> • % change from baseline in FEV₁ • % change from baseline in FVC 	<p>Included in Morrison 2014 SR</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
	during hospitalisation Intervention 2. Oscillating device (flutter) <ul style="list-style-type: none"> 15-minute sessions, 4 times daily during hospitalisation 	age: 12 (7 to 44)		
Mcllwaine 2001 (Canada) RCT, parallel design	Intervention 1. Positive expiratory pressure (PEP) mask <ul style="list-style-type: none"> 20-minute sessions twice daily for 1 year Intervention 2. Oscillating device (flutter) <ul style="list-style-type: none"> Sessions of ≥ 15 minutes twice daily for 1 year 	N=40 children and young people with CF Age range: 7 to 17 <ul style="list-style-type: none"> Intervention 1: n=20 Intervention 2: n=20 Participants were excluded if they had been hospitalised within the past month for a pulmonary exacerbation, or if they were not stable on clinical evaluation, chest radiograph or pulmonary function	<ul style="list-style-type: none"> Patient preference: self-withdrawal due to lack of perceived effectiveness FEV₁ (% change from baseline) FVC (% change from baseline) 	Included in Mcllwaine 2015 SR
Mcllwaine 2013 (Canada) Multi-centre RCT, parallel design	Intervention 1: Positive expiratory pressure (PEP) mask <ul style="list-style-type: none"> 6 cycles; treatment for 1 year Intervention 2: High Frequency Chest Wall Oscillation (HFCWO) <ul style="list-style-type: none"> 6 sets of 5-minute cycles; treatment for 1 year 	N=107 children, young people and adults with CF from 12 CF centres. <ul style="list-style-type: none"> PEP: n=51 HFCWO: n=56 Age range: 6 to 47 years. FEV ₁ >40% predicted Participants were excluded if they had been hospitalised within the past month for a pulmonary exacerbation, or if they were not stable on clinical evaluation, chest radiograph or	<ul style="list-style-type: none"> FEV₁ (% change from baseline) FVC (% change from baseline) QoL (CFQ-R: physical domain, treatment burden, respiratory domain) 	Included in Mcllwaine 2015 SR On entering the study, participants performed a 2-month washout period before being allocated to an intervention

Study	Intervention/Comparison	Population	Outcomes	Comments
		pulmonary function.		
Newbold 2005 (Canada) RCT, parallel design	Intervention 1: Positive expiratory pressure (PEP) mask <ul style="list-style-type: none"> 20-minute session, twice daily for 13 months Intervention 2: Oscillating device (flutter) <ul style="list-style-type: none"> Duration approx. 20 minutes, twice daily for 13 months 	N=42 adults with CF <ul style="list-style-type: none"> PEP: n=21. Mean (SD) age: 28 (8.1) Flutter: n=21. Mean (SD) age: 31 (8.7) Participants were excluded if they had been hospitalised within the past month for a pulmonary exacerbation, had changed their medication within the past month, or did not have a daily cough or daily sputum	<ul style="list-style-type: none"> FEV₁ (% change from baseline) FVC (% change from baseline) Hospitalizations for respiratory exacerbations 	Included in McIlwaine 2015 SR
Oermann 2001 (USA) RCT, crossover design	Intervention 1: Oscillating device (flutter). <ul style="list-style-type: none"> Number of times per day not reported Intervention 2: High Frequency Chest Wall Oscillation. <ul style="list-style-type: none"> Number of times per day not reported. 	N=29 children, young people and adults with CF <ul style="list-style-type: none"> Mean (range) age: 23 (9 to 39) Baseline FVC range: 50 to 80 % predicted Clinically stable for 1 month prior to enrolment. 	<ul style="list-style-type: none"> FEV₁ % predicted FVC % predicted 	Included in Morrison 2014 SR <p>5 participants withdrew (4 exited due to illness and 1 due to non-compliance with clinic visits)</p> <p>4 weeks in each arm, 2-week lead-in/ wash out periods during which time they resumed their normal routine therapies which were not outlined</p>
Padman 1999 (USA) RCT, crossover design	Intervention 1: Manual chest physiotherapy Intervention 2: Positive expiratory pressure (PEP) mask <ul style="list-style-type: none"> 15-minute sessions, 3 times per day for 1 month 	N=15 children and young people with CF were randomized, 6 completed the study. <ul style="list-style-type: none"> Age range: 5 to 17 Participants were clinically stable and able to perform respiratory 	<ul style="list-style-type: none"> % change from baseline in FEV₁ 	Included in Morrison 2014 SR <p>5 participants excluded due to hospital admission for acute exacerbation, 4 withdrew (no reason given)</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
	Intervention 3: Oscillating device (flutter) • 15-minute sessions, 3 times per day for 1 month	function tests; no hospitalisations in the month prior to the study		
Placidi 2006 (Australia) RCT, crossover design	Intervention: Positive expiratory pressure (PEP) mask Control: no airway clearance technique (directed cough)	N= 17 young people and adults with CF Age: >15 years. Mean (SD) age: 28 (7) People had severe lung disease and were admitted for pulmonary exacerbation.	<ul style="list-style-type: none"> • Sputum dry weight • Sputum wet weight • Lung function - FEV₁ • Lung function - FVC • Oxygen saturation - Spo2 % 	Included in Moran 2013 SR Follow-up: mean 2 days
Tannenbaum 2005 (Country not reported) RCT, parallel design	Intervention 1: Positive expiratory pressure (PEP) mask Intervention 2: oscillating device (cornet)	N = 30 children and young people with CF Age range: 6 to 15 years	<ul style="list-style-type: none"> • % change from baseline FEV₁ 	Included in McIlwaine 2015 SR Information was provided from 3 abstracts, no further information obtained
Young 2008 (Australia) RCT, crossover design	Intervention: Overnight non-invasive ventilation for 6 weeks Control: No airway clearance technique (room air) for 6 weeks	N=8 adults with CF Mean (SD) age: 37 (8) years Participants with moderate lung disease.	<ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted • Oxygen saturation (nocturnal) (%) • Quality of life: CF QOL chest symptom score; CF QOL traditional dyspnoea index score 	Included in Moran 2013 SR 2-week washout period
van Winden 1998 (The Netherlands) RCT, crossover design	Intervention 1: Positive expiratory pressure (PEP) mask • Twice daily Intervention 2: Oscillating device (flutter) • Twice daily	N=22 children and young people with CF Mean age (range): 12 years (7 to 17). People were clinically stable for 2 weeks before the study	<ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted 	Included in Morrison 2014 SR 2 weeks in each arm, 1 week wash-in and wash-out period
Warwick 2004	Intervention 1:	N=12 adults with	<ul style="list-style-type: none"> • Sputum weight 	Included in

Study	Intervention/Comparison	Population	Outcomes	Comments
(USA) RCT, crossover design	Manual physiotherapy. <ul style="list-style-type: none"> 10 hand positions Duration: approximately 45 to 50 minutes daily for 4 weeks Intervention 2: High Frequency Chest Compression <ul style="list-style-type: none"> Duration: approximately 36 to 50 minutes daily for 4 weeks 	CF Age mean (range): 29.1 (19 to 50) Consistent sputum producers; all volunteers with no illness within 6 weeks of study	(dry) <ul style="list-style-type: none"> Sputum weight (wet) 	Morrison 2014 SR All treatments preceded by nebulisers.

1 CF: cystic fibrosis, HFCWO: high frequency chest wall oscillation; PEP: positive expiratory pressure; SR:
2 systematic review

3 9.2.4 Clinical evidence profile

4 The summary clinical evidence profiles are presented in Table 68 to Table 75.

5 **Table 69: Summary clinical evidence profile: Comparison 2. Manual physiotherapy**
6 **versus oscillating devices**

Comparison 2. Manual physiotherapy compared to oscillating device for CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Oscillating device (OD)	Manual physiotherapy				
Lung function Measured as FEV ₁ % change from baseline Scale from: 0 to 100 Follow-up: mean 8.8 days	The mean % change from baseline in FEV ₁ in the OD group was 43.7	The mean % change from baseline in FEV ₁ in the manual physiotherapy groups was 7.9 lower (31.04 lower to 15.24 higher)		22 (Homnick 1998)	⊕⊕⊕⊕ very low ^{1,2}	
Lung function Measured as FEV ₁ % change from baseline Scale	The mean % change from baseline in FEV ₁ in the OD group was 3.66	The mean % change from baseline in FEV ₁ in the manual physiotherapy groups was 2.59 higher (6.3 lower to		6 (Padman 1999)	⊕⊕⊕⊕ very low ^{3,4}	

Comparison 2. Manual physiotherapy compared to oscillating device for CF					
from: 0 to 100 Follow-up: mean 1 months		11.48 higher)			
Lung Function - FVC % change from baseline Scale from: 0 to 100 Follow-up: mean 2 weeks	The mean % change from baseline in FVC in the OD group was 27.2	The mean % change from baseline in FVC in the manual physiotherapy group was 2.9 higher (14.21 lower to 20.01 higher)		22 (Homnick 1998)	⊕⊕⊕⊕ very low ^{1,4}
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; FVC: forced vital capacity; MD: mean difference					

- 1 The quality of the evidence downgraded by 2 due to selection bias and attrition bias
- 2 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed 1 default MID
- 3 The quality of the evidence was downgraded by 2 due to attrition bias and reporting bias.
- 4 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default MIDs

6 **Table 70: Summary clinical evidence profile: Comparison 3. Manual physiotherapy**
7 **versus high frequency chest wall oscillation**

Comparison 3. Manual physiotherapy techniques compared to high frequency chest oscillation therapy for CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	High frequency chest oscillation therapy (HFCWO)	Manual physiotherapy				
Sputum weight (dry) (grams) Follow-up: 1-2 weeks	The mean sputum weight (dry) in the HFCWO groups was 0.57	The mean sputum weight (dry) in the manual physiotherapy groups was 0.13 lower (0.42 lower to 0.16 higher)		12 (Warwick 2004)	⊕⊕⊕⊕ low ^{1,2}	
Sputum weight (wet) (grams) Follow-up:	The mean sputum weight (wet) in the HFCWO groups was	The mean sputum weight (wet) in the manual physiotherapy		12 (Warwick 2004)	⊕⊕⊕⊕ low ^{1,2}	

Comparison 3. Manual physiotherapy techniques compared to high frequency chest oscillation therapy for CF

1-2 weeks	13.56	groups was 4.04 lower (10.77 lower to 2.69 higher)			
-----------	-------	---	--	--	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; HFCWO: high frequency chest wall oscillation; MD: mean difference

1 The quality of the evidence was downgraded by 1 due to lack of blinding.

2 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a default MID

Table 71: Summary clinical evidence profile: Comparison 4. Positive expiratory pressure mask (PEP) versus no airway clearance technique

Comparison 4. Positive expiratory pressure mask (PEP) compared to no airway clearance technique for CF

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No airway clearance technique	Positive expiratory pressure mask (PEP)				
Sputum dry weight (grams) Follow-up: mean 2 days	The mean sputum dry weight in the control groups was 0.97	The mean sputum dry weight in the PEP groups was 0.03 lower (0.48 lower to 0.42 higher)		17 (Placidi 2006)	⊕⊕⊕⊕ low ¹	
Sputum wet weight (grams) Follow-up: mean 2 days	The mean sputum wet weight in the control groups was 13.98	The mean sputum wet weight in the PEP groups was 1.8 higher (1.72 lower to 5.32 higher)		17 (Placidi 2006)	⊕⊕⊕⊕ moderate ²	
Lung function - FEV ₁ % predicted Scale from: 0 to 100 Follow-up: mean 2 days	The mean FEV ₁ % predicted in the control groups was 60.3	The mean FEV ₁ % predicted in the PEP groups was 2.1 higher (11.73 lower to 15.93 higher)		16 (Braggion 1995)	⊕⊕⊕⊕ very low ^{3,4}	
Lung function - FEV ₁ (litres) Follow-up: mean 2 days	The mean FEV ₁ (litres) in the control groups was 0.99	The mean FEV ₁ (litres) in the PEP groups was 0.01 higher (0.18 lower to 0.2 higher)		17 (Placidi 2006)	⊕⊕⊕⊕ low ¹	

Comparison 4. Positive expiratory pressure mask (PEP) compared to no airway clearance technique for CF

Lung Function FVC % predicted Scale from: 0 to 100 Follow-up: mean 2 days	The mean FVC % predicted in the control groups was 81.6 % predicted	The mean FVC % predicted in the PEP groups was 1.2 higher (12.88 lower to 15.28 higher)		16 (Braggion 1995)	⊕⊕⊕⊕ very low ^{1,3}	
Lung function - FVC (litres) Follow-up: mean 2 days	The mean FVC (litres) in the control groups was 1.96	The FVC in the PEP groups was 0.05 higher (0.35 lower to 0.45 higher)		17 (Placidi 2006)	⊕⊕⊕⊕ low ¹	
Oxygen saturation - Spo2 % Scale from: 0 to 100 Follow-up: mean 2 days	The mean oxygen saturation (spo2) in the control groups was 94.6	The mean oxygen saturation (spo2) in the PEP groups was 0.3 higher (0.58 lower to 1.18 higher)		17 (Placidi 2006)	⊕⊕⊕⊕ moderate ²	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; MD: mean difference; SpO₂: peripheral capillary oxygen saturation

- 1 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default MIDs
- 2 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a default MID
- 3 The quality of the evidence was downgraded by 2 due to lack of blinding, attrition bias and reporting bias
- 4 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 clinical MIDs

Table 72: Summary clinical evidence profile: Comparison 6. Positive expiratory pressure mask (PEP) versus oscillating devices

Comparison 6. Positive expiratory pressure mask (PEP) compared to oscillating device for CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	oscillating device (OD)	Positive expiratory pressure mask (PEP)				
Patient preference: self-withdrawal due to lack of perceived effectiveness Follow-up:	250 per 1000	22 per 1000 (2 to 385)	RR 0.09 (0.01 to 1.54)	40 (McIlwaine 2001)	⊕⊕⊕⊕ very low ^{1,2}	

Comparison 6. Positive expiratory pressure mask (PEP) compared to oscillating device for CF						
mean 1 years						
Hospitalizations for respiratory exacerbations number per participant Follow-up: mean 13 months	The mean number of hospitalizations for respiratory exacerbations in the OD group was 0.7 per participant	The mean hospitalizations for respiratory exacerbations in the PEP groups was 0.4 lower (0.92 lower to 0.12 higher)		42 (Newbold 2005)	⊕⊕⊕⊖ low ^{3,4}	
Lung function - FEV ₁ % change from baseline Scale from: 0 to 100 Follow-up: 2-4 weeks	The mean % change from baseline in FEV ₁ in the OD group was 3.66	The mean % change from baseline in FEV ₁ in the PEP groups was 4.08 higher (4.66 lower to 12.82 higher)		6 (Padman 1999)	⊕⊕⊕⊖ very low ^{4,5}	
Lung function - FEV ₁ % change from baseline Scale from: 0 to 100 Follow-up: mean 6-12 months	The mean % change from baseline in FEV ₁ in the OD groups was -10.95	The mean % change from baseline in FEV ₁ in the PEP groups was 9.71 higher (2.12 lower to 21.54 higher)		30 (McIlwaine 2001)	⊕⊕⊕⊖ low ^{1,4}	
Lung function - FEV ₁ % change from baseline Scale from: 0 to 100 Follow-up: 1-2 years	The mean % change from baseline in FEV ₁ in the OD groups was 2.78	The mean % change from baseline in FEV ₁ in the PEP groups was 2.82 lower (6.36 lower to 0.72 higher)		160 (McIlwaine 2013, Newbold 2005, Tannenbaum 2005)	⊕⊕⊕⊖ low ^{4,6}	
Lung function - FVC % change from baseline Scale from: 0 to 100 Follow-up: mean 1 years	The mean % change from baseline in FVC in the OD groups was -0.07	The mean % change from baseline in FVC in the PEP groups was 0.44 lower (6.66 lower to 5.78 higher)		160 (McIlwaine 2001, McIlwaine 2013, Newbold 2005)	⊕⊕⊕⊖ low ^{6,7}	
Lung function - FVC (% predicted) Scale from: 0 to 100 Follow-up: 2-4 weeks	The mean FVC % predicted in the OD groups was 99 % predicted	The mean FVC % predicted in the PEP groups was 2 lower (4.09 lower to 0.09 higher)		22 (van Winden 1998)	⊕⊕⊕⊖ moderate ⁴	
QOL – CFQ-R: physical domain Scale from: 0 to	The mean CFQ-R - physical domain in the	The mean CFQ-R - physical domain in the		107 (McIlwaine 2013)	⊕⊕⊕⊕ high ⁸	

Comparison 6. Positive expiratory pressure mask (PEP) compared to oscillating device for CF						
100 Follow-up: mean 1 years	OD groups was -3.04	PEP groups was 2.2 higher (1.32 lower to 5.72 higher)				
QOL – CFQ-R: treatment burden Scale from: 0 to 100 Follow-up: mean 1 years	The mean QOL-CFQ-R: treatment burden in the OD groups was -3.6	The mean QOL – CFQ-R: treatment burden in the PEP groups was 1.05 higher (6.35 lower to 8.45 higher)		107 (McIlwaine 2013)	⊕⊕⊕⊕ high ⁸	
QOL – CFQ-R: respiratory domain Scale from: 0 to 100 Follow-up: mean 1 years	The mean CFQ-R - respiratory domain in the OD groups was 0.19	The mean CFQ-R - respiratory domain in the PEP groups was 2.79 higher (3.68 lower to 9.26 higher)		107 (McIlwaine 2013)	⊕⊕⊕⊖ moderate 8,9	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; CFQ-R: cystic fibrosis questionnaire revised; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; MD: mean difference; OD: oscillating device; PEP: positive expiratory pressure; RR: risk ratio</p>						

- 1 The quality of the evidence was downgraded by 1 due to reporting bias.
- 2 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default MIDs
- 3 The quality of the evidence was downgraded by 1 due to differences in baseline characteristics (pulmonary function values) between both groups.
- 4 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a default MID
- 5 The quality of the evidence was downgraded by 2 due to attrition bias and reporting bias.
- 6 Taking into account weighting in a meta-analysis and the likely contribution from each component, the quality of the evidence was downgraded by 1 due differences in baseline participant characteristics.
- 7 The quality of the evidence was downgraded by 1 due to serious heterogeneity (I-squared inconsistency statistic of 69%) and no plausible explanation was found with sensitivity analysis.
- 8 Clinical MID=8.5 was used to assess imprecision because the CFQ-R questionnaire (Quittner et al. 2009) was used
- 9 The quality of the evidence was downgraded by 1 as 95% CI crossed 1 clinical MID

Table 73: Summary clinical evidence profile: Comparison 7. Positive expiratory pressure mask (PEP) versus High Frequency Chest Wall Oscillation (HFCWO)

Comparison 7. Positive expiratory pressure mask (PEP) compared to high frequency chest wall oscillation (HFCWO) for CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	High frequency oscillation	Positive expiratory pressure				

Comparison 7. Positive expiratory pressure mask (PEP) compared to high frequency chest wall oscillation (HFCWO) for CF						
	therapy (HFCWO)	mask (PEP)				
Sputum volume (ml) Follow-up: mean 1 weeks	The mean sputum volume in the HFCWO groups was 6.7	The mean sputum volume in the PEP groups was 1.8 higher (3 lower to 6.6 higher)		23 (Grzincich 2008)	⊕⊕⊕⊕ low ^{1,2}	
Respiratory exacerbations: number of patients Follow-up: mean 1 years	833 per 1000	608 per 1000 (458 to 792)	RR 0.73 (0.55 to 0.95)	91 (McIlwaine 2013)	⊕⊕⊕⊕ moderate ²	
Pulmonary exacerbations (patients requiring antibiotics) Follow-up: mean 1 years	870 per 1000	615 per 1000 (348 to 824)	RR 0.71 (0.55 to 0.93)	88 (McIlwaine 2013)	⊕⊕⊕⊕ moderate ²	
Lung function - FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1 weeks	The mean FEV ₁ % predicted in the HFCWO groups was 64.8	The mean FEV ₁ % predicted in the PEP groups was 0.67 higher (8.04 lower to 9.38 higher)		39 (Braggion 1995; Grzincich 2008)	⊕⊕⊕⊕ very low ^{3,4}	
Lung Function - FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1-2 weeks	The mean FEV ₁ % predicted in the HFCWO groups was 69	The mean FEV ₁ % predicted in the PEP groups was 3 lower (20.54 lower to 14.54 higher)		15 (Darbee 2005)	⊕⊕⊕⊕ very low ^{4,5}	
Lung function - FEV ₁ change from baseline in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1 years	The mean change in FEV ₁ % predicted in the HFCWO groups was 9.4	The mean change in FEV ₁ % predicted in the PEP groups was 3.59 lower (9.29 lower to 2.11 higher)		88 (McIlwaine 2013)	⊕⊕⊕⊕ moderate ⁶	
FVC % predicted Scale from: 0 to 100 Follow-up: 1-2 weeks	The mean FVC % predicted in the HFCWO groups was 83	The mean FVC in the PEP groups was 3 lower (16.6 lower to 10.6 higher)		15 (Darbee 2005)	⊕⊕⊕⊕ very low ^{5,7}	

Comparison 7. Positive expiratory pressure mask (PEP) compared to high frequency chest wall oscillation (HFCWO) for CF

FVC % predicted Scale from: 0 to 100 Follow-up: 1 weeks	The mean FVC % predicted in the HFCWO groups was 86.4	The mean FVC % predicted in the PEP groups was 0.66 higher (7.4 lower to 8.71 higher)		39 (Braggion 1995, Grzincich 2008)	⊕⊕⊕⊖ moderate ³	
Lung function - FVC change from baseline in % predicted Scale from: 0 to 100 Follow-up: 1 years	The mean change in FVC % predicted in the HFCWO groups was 11.39	The mean change in FVC % predicted in the PEP groups was 5 lower (10.3 lower to 0.3 higher)		88 (McIlwaine 2013)	⊕⊕⊕⊖ moderate ²	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; HFCWO: high frequency chest wall oscillation; MD: mean difference; PEP: positive expiratory pressure; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1 as risk of bias could not be fully assessed from abstract paper which did not discuss method in detail.
- 2 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a default MID.
- 3 Taking into account weighting in a meta-analysis and the likely contribution from each component, the quality of the evidence was downgraded by 1 as risk of bias could not be fully assessed from abstract paper which did not discuss method in detail.
- 4 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 clinical MIDs.
- 5 The quality of the evidence was downgraded by 1 due to selection bias.
- 6 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a clinical MID
- 7 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default MIDs

Table 74: Summary clinical evidence profile: Comparison 12. Oscillating device versus High Frequency Chest Wall Oscillation (HFCWO)

Comparison 12. Oscillating device compared to high frequency chest wall oscillation for CF						
Outcome s	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	High frequency chest wall oscillation (HFCWO)	Oscillating device (OD)				
Lung function - FEV ₁ % predicted Follow-up: 2-4	The mean FEV ₁ % predicted in the HFCWO groups was 56.5	The mean FEV ₁ % predicted in the OD groups was 1.6 lower (3.44 lower to 0.24 higher)		24 (Oermann 2001)	⊕⊕⊕⊖ moderate ₁	

Comparison 12. Oscillating device compared to high frequency chest wall oscillation for CF					
weeks					
Lung function - FVC % predicted Follow-up: 2-4 weeks	The mean FVC % predicted in the HFCWO groups was 74	The mean FVC% predicted in the OD groups was 1.4 lower (3.07 lower to 0.27 higher)		24 (Oermann 2001)	⊕⊕⊕⊕ low ^{1,2}
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; FVC: forced vital capacity; HFCWO: high frequency chest wall oscillation; MD: mean difference; OD: oscillating device					

- 1 The quality of the evidence was downgraded by 1 due to reporting bias.
2 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed a default MID.

3 **Table 75: Summary clinical evidence profile: Comparison 14. Non-invasive ventilation**
4 **(NIV) versus no airway clearance technique**

Comparison 14. Non-invasive ventilation (NIV) compared to no airway clearance technique for CF						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No airway clearance technique	Non-invasive ventilation (NIV)				
Lung function - FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 weeks	The mean FEV ₁ % predicted in the control groups was 32	The mean FEV ₁ % predicted in the NIV groups was 1 higher (8.62 lower to 10.62 higher)		8 (Young 2008)	⊕⊕⊕⊕ low ¹	
Lung function - FVC % predicted Scale from: 0 to 100 Follow-up: 6 weeks	The mean FVC % predicted in the control groups was 54	The FVC% predicted in the NIV groups was 4 higher (10.3 lower to 18.3 higher)		8 (Young 2008)	⊕⊕⊕⊕ low ²	
Oxygen saturation (nocturnal) % Scale from: 0 to 100 Follow-up: 6 weeks	The mean oxygen saturation (%) (nocturnal) in the control groups was 89	The mean oxygen saturation (nocturnal) (%) in the NIV groups was 3 higher (1.12 lower to 7.12 higher)		8 (Young 2008)	⊕⊕⊕⊕ moderate ³	

Comparison 14. Non-invasive ventilation (NIV) compared to no airway clearance technique for CF

Quality of life - CF QOL chest symptom score Scale from: 0 to 100 Follow-up: 6 weeks	The mean CF-QOL chest symptom score in the control groups was 64	The mean CF-QOL chest symptom score in the NIV groups was 7 higher (11.73 lower to 25.73 higher)		8 (Young 2008)	⊕⊕⊕⊖ low ^{1,4}	
Quality of life - CF QOL traditional dyspnoea index score Scale from: 0 to 100 Follow-up: 6 weeks	The mean CF-QOL traditional dyspnoea index score in the control groups was -1.9	The mean CF-QOL traditional dyspnoea index score in the NIV groups was 2.9 higher (0.71 to 5.09 higher)		8 (Young 2008)	⊕⊕⊕⊖ moderate ^{4,5}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; MD: mean difference; NIV: non-invasive ventilation

- 1 *1 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 clinical*
- 2 *MIDs*
- 3 *2 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default*
- 4 *MIDs*
- 5 *3 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed 1 default MID*
- 6 *4 Clinical MID=5 was used to assess imprecision for quality of life because the CF QOL questionnaire (Gee et al.*
- 7 *2000) was used*
- 8 *5 The quality of the evidence was downgraded by 1 due to serious imprecision as 95% CI crossed 1 clinical MID*

9 9.2.5 Economic evidence

10 One conference abstract identified in the literature search conducted for this guideline was
11 considered relevant to this review question. This paper undertook a cost-consequence
12 analysis to compare positive expiratory pressure (PEP) to high frequency chest wall
13 oscillation (HFCWO) in 107 cystic fibrosis patients in Canada (McIlwaine 2014). They
14 concluded that PEP was less expensive and more effective (dominant) at reducing the
15 number of exacerbations than HFCWO. The methods and results from this analysis are
16 provided in Appendix K.

17 This review question was not prioritised for de novo economic modelling. To aid
18 consideration of cost-effectiveness, relevant resource and cost use data are presented in
19 Appendix K.

20 Full details of the search and economic article selection flow chart can be found in Appendix
21 E and F, respectively. Data extraction tables and quality assessments and of included
22 studies can be found in Appendix L and M, respectively.

23 9.2.6 Evidence statements

24 9.2.6.1 Comparison 1. Manual physiotherapy versus no airway clearance techniques

25 No evidence was found for this comparison.

1 **9.2.6.2 Comparison 2. Manual physiotherapy techniques versus oscillating devices**

2 **Sputum volume**

3 No evidence was found for this critical outcome.

4 **Patient preference**

5 No evidence was found for this critical outcome.

6 **Pulmonary exacerbations**

7 No evidence was found for this critical outcome.

8 **Pulmonary function tests: FEV₁ and FVC**

9 Very low quality evidence from 1 crossover RCT with 22 children, young people and adults
10 with cystic fibrosis showed no clinically significant difference in FEV₁ percent change from
11 baseline between manual physiotherapy techniques and oscillating device after 8.8 days
12 follow-up.

13 Very low quality evidence from 1 crossover RCT with 6 children and young people with cystic
14 fibrosis showed no clinically significant difference in FEV₁ percent change from baseline
15 between manual physiotherapy techniques and oscillating device after 1 month follow-up.

16 Very low quality evidence from 1 crossover RCT with 22 children, young people and adults
17 with cystic fibrosis showed no clinically significant difference in FVC percent change from
18 baseline between manual physiotherapy techniques and oscillating device after 2 weeks
19 follow-up.

20 **Oxygen saturation**

21 No evidence was found for this important outcome.

22 **Quality of life**

23 No evidence was found for this important outcome.

24 **Hospitalisations**

25 No evidence was found for this important outcome.

26 **9.2.6.3 Comparison 3. Manual physiotherapy versus High Frequency Chest Wall Oscillation**

27 **Sputum volume**

28 Low quality evidence from 1 crossover RCT with 12 adults with cystic fibrosis showed no
29 clinically significant difference in sputum dry weight between manual physiotherapy and
30 HFCWO after 1 to 2 weeks follow-up.

31 Low quality evidence from 1 crossover RCT with 12 adults with cystic fibrosis showed no
32 clinically significant difference in sputum wet weight between manual physiotherapy and
33 HFCWO after 1 to 2 weeks follow-up.

34 **Patient preference**

35 No evidence was found for this critical outcome.

- 1 **Pulmonary exacerbations**
- 2 No evidence was found for this critical outcome.
- 3 **Pulmonary function tests**
- 4 No evidence was found for this important outcome.
- 5 **Oxygen saturation**
- 6 No evidence was found for this important outcome.
- 7 **Quality of life**
- 8 No evidence was found for this important outcome.
- 9 **Hospitalisations**
- 10 No evidence was found for this important outcome.
- 11 **9.2.6.4 Comparison 4. Positive expiratory pressure mask (PEP) versus no airway clearance**
- 12 **technique**
- 13 **Sputum volume**
- 14 Low quality evidence from 1 crossover RCT with 17 young people and adults with cystic
- 15 fibrosis showed no clinically significant difference in sputum dry weight between PEP and
- 16 control after 2 days follow-up.
- 17 Moderate quality evidence from 1 crossover RCT with 17 young people and adults with cystic
- 18 fibrosis showed no clinically significant difference in sputum wet weight between PEP and
- 19 control after 2 days follow-up.
- 20 **Patient preference**
- 21 No evidence was found for this critical outcome.
- 22 **Pulmonary exacerbations**
- 23 No evidence was found for this critical outcome.
- 24 **Pulmonary function tests: FEV₁ and FVC**
- 25 Very low quality evidence from 1 crossover RCT with 16 young people and adults with cystic
- 26 fibrosis showed no clinically significant difference in percent predicted FEV₁ between PEP
- 27 and control after 2 days follow-up.
- 28 Low quality evidence from 1 crossover RCT with 17 young people and adults showed no
- 29 clinically significant difference in FEV₁ (L) between PEP and control after 2 days follow-up.
- 30 Very low quality evidence from 1 crossover RCT with 16 young people and adults with cystic
- 31 fibrosis showed no clinically significant difference in percent predicted FVC between PEP
- 32 and control after 2 days follow-up.
- 33 Low quality evidence from 1 crossover RCT with 17 young people and adults showed no
- 34 clinically significant difference in FVC between PEP and control after 2 days follow-up.

1 **Oxygen saturation**

2 Moderate quality evidence from 1 crossover RCT with 17 young people and adults showed
3 no clinically significant difference in oxygen saturation between PEP and control after 2 days
4 follow-up.

5 **Quality of life**

6 No evidence was found for this important outcome.

7 **Hospitalisations**

8 No evidence was found for this important outcome.

9 **9.2.6.5 Comparison 5. Positive expiratory pressure mask (PEP) versus active cycle of**
10 **breathing techniques (ACBT)**

11 No evidence was found for this comparison.

12 **9.2.6.6 Comparison 6. Positive expiratory pressure mask (PEP) versus oscillating devices**

13 **Sputum volume**

14 No evidence was found for this critical outcome.

15 **Patient preference**

16 Very low quality evidence from 1 parallel RCT with 40 children and young people with cystic
17 fibrosis showed no clinically significant difference in self-withdrawal due to lack of perceived
18 effectiveness between PEP and oscillating device after 1 month follow-up.

19 **Pulmonary exacerbations**

20 Low quality evidence from 1 parallel RCT with 42 adults with cystic fibrosis showed no
21 clinically significant difference in number of hospitalisation for respiratory exacerbations per
22 participant between PEP and oscillating device after 13 months follow-up.

23 **Pulmonary function tests: FEV₁ and FVC**

24 Very low quality evidence from 1 crossover RCT with 6 children and young people with cystic
25 fibrosis showed no clinically significant difference in FEV₁ percent change from baseline
26 between PEP and oscillating device after 2 to 4 weeks follow-up.

27 Low quality evidence from 1 parallel RCT with 30 children and young people with cystic
28 fibrosis showed no clinically significant difference in FEV₁ percent change from baseline
29 between PEP and oscillating device after 6 to 12 months follow-up.

30 Low quality evidence from 3 parallel RCTs with 160 children, young people and adults with
31 cystic fibrosis showed no clinically significant difference in FEV₁ percent change from
32 baseline between PEP and oscillating device after 1 to 2 years follow-up.

33 Low quality evidence from 3 RCTs with 160 children, young people and adults with cystic
34 fibrosis showed no clinically significant difference in FVC percent change from baseline
35 between PEP and oscillating device after 1 year follow-up. Moderate inconsistency was
36 observed between the trials, but all of them showed a no clinically significant difference
37 between both treatment groups.

1 Moderate quality evidence from 1 crossover RCT with 22 children and young people with
2 cystic fibrosis showed no clinically significant difference in percent predicted FVC between
3 PEP and oscillating device after 2 to 4 weeks follow-up.

4 **Oxygen saturation**

5 No evidence was found for this important outcome.

6 **Quality of life**

7 High quality evidence from 1 parallel RCT with 107 children, young people and adults with
8 cystic fibrosis showed no clinically significant difference in the physical domain of the CFQ-R
9 questionnaire between PEP and oscillating device after 1 year follow-up.

10 High quality evidence from 1 parallel RCT with 107 children, young people and adults with
11 cystic fibrosis showed no clinically significant difference in the treatment burden domain of
12 the CFQ-R questionnaire between PEP and oscillating device after 1 year follow-up.

13 Moderate quality evidence from 1 parallel RCT with 107 children, young people and adults
14 with cystic fibrosis showed no clinically significant difference in the respiratory domain of the
15 CFQ-R questionnaire between PEP and oscillating device after 1 year follow-up.

16 **Hospitalisations**

17 No evidence was found for this important outcome.

18 **9.2.6.7 Comparison 7. Positive expiratory pressure mask (PEP) versus High Frequency Chest** 19 **Wall Oscillation (HFCWO)**

20 **Sputum volume**

21 Low quality evidence from 1 RCT with 23 adults with cystic fibrosis showed no clinically
22 significant difference in sputum volume between PEP and HFCWO after 1 week follow-up.

23 **Patient preference**

24 No evidence was found for this critical outcome.

25 **Pulmonary exacerbations**

26 Moderate quality evidence from 1 parallel RCT with 91 children, young people and adults
27 with cystic fibrosis showed a clinically significant beneficial effect of PEP compared to
28 HFCWO in number of patients with respiratory exacerbation after 1 year follow-up.

29 Moderate quality evidence from 1 parallel RCT with 88 children, young people and adults
30 with cystic fibrosis showed a clinically significant beneficial effect of PEP compared to
31 HFCWO in number of patients requiring antibiotics for respiratory exacerbation after 1 year
32 follow-up.

33 **Pulmonary function tests: FEV₁ and FVC**

34 Very low quality evidence from a 2 RCTs with 39 young people and adults with cystic fibrosis
35 showed no clinically significant difference in percent predicted FEV₁ between PEP and
36 HFCWO after 1 week follow-up.

37 Very low quality evidence from 1 crossover RCT with 15 children, young people and adults
38 with cystic fibrosis showed no clinically significant difference in percent predicted FEV₁
39 between PEP and HFCWO after 1 to 2 weeks follow-up.

1 Moderate quality evidence from 1 parallel RCT with 88 children, young people and adults
2 with cystic fibrosis showed no clinically significant difference in change from baseline percent
3 predicted FEV₁ between PEP and HFCWO after 1 year follow-up.

4 Very low quality evidence from 1 crossover RCT with 15 children, young people and adults
5 with cystic fibrosis showed no clinically significant difference in percent predicted FVC
6 between PEP and HFCWO after 1 to 2 weeks follow-up.

7 Moderate quality evidence from a 2 RCTs with 39 young people and adults with cystic
8 fibrosis showed no clinically significant difference in percent predicted FVC between PEP
9 and HFCWO after 1 week follow-up.

10 Moderate quality evidence from 1 parallel RCT with 88 children, young people and adults
11 with cystic fibrosis showed no clinically significant difference in change from baseline percent
12 predicted FVC between PEP and HFCWO after 1 year follow-up.

13 **Oxygen saturation**

14 No evidence was found for this important outcome.

15 **Quality of life**

16 No evidence was found for this important outcome.

17 **Hospitalisations**

18 No evidence was found for this important outcome.

19 **9.2.6.8 Comparison 8. Active cycle of breathing technique (ACBT) versus no airway clearance 20 technique**

21 No evidence was retrieved for this comparison.

22 **9.2.6.9 Comparison 9. Active cycle breathing technique (ACBT) versus autogenic drainage 23 (AD)**

24 No evidence was retrieved for this comparison.

25 **9.2.6.10 Comparison 10. Autogenic drainage (AD) versus no airway clearance technique**

26 No evidence was retrieved for this comparison.

27 **9.2.6.11 Comparison 11. Oscillating device versus no airway clearance technique**

28 No evidence was retrieved for this comparison.

29 **9.2.6.12 Comparison 12. Oscillating device versus High Frequency Chest Wall Oscillation 30 (HFCWO)**

31 **Sputum volume**

32 No evidence was found for this critical outcome.

33 **Patient preference**

34 No evidence was found for this critical outcome.

1 **Pulmonary exacerbations**

2 No evidence was found for this critical outcome.

3 **Pulmonary function tests: FEV₁ and FVC**

4 Moderate quality evidence from 1 crossover RCT with 24 children, young people and adults
5 with cystic fibrosis showed no clinically significant difference in percent predicted FEV₁
6 between oscillating device and HFCWO after 2 to 4 weeks follow-up.

7 Low quality evidence from 1 crossover RCT with 24 participants showed no clinically
8 significant difference in percent predicted FVC between oscillating device and HFCWO after
9 2 to 4 weeks follow-up.

10 **Oxygen saturation**

11 No evidence was found for this important outcome.

12 **Quality of life**

13 No evidence was found for this important outcome.

14 **Hospitalisations**

15 No evidence was found for this important outcome.

16 **9.2.6.13 Comparison 13. High Frequency Chest Wall Oscillation (HFCWO) versus no clearance**
17 **technique**

18 No evidence was retrieved for this comparison.

19 **9.2.6.14 Comparison 14. Non-invasive ventilation (NIV) versus no airway clearance technique**

20 **Sputum volume**

21 No evidence was found for this critical outcome.

22 **Patient preference**

23 No evidence was found for this critical outcome.

24 **Pulmonary exacerbations**

25 No evidence was found for this critical outcome.

26 **Pulmonary function tests: FEV₁ and FVC**

27 Low quality evidence from 1 crossover RCT with 8 adults with cystic fibrosis showed no
28 clinically significant difference in percent predicted FEV₁ between NIV and control after 6
29 weeks follow-up.

30 Low quality evidence from 1 crossover RCT with 8 adults with cystic fibrosis showed no
31 clinically significant difference in percent predicted FVC between NIV and control after 6
32 weeks follow-up.

1 **Oxygen saturation**

2 Moderate quality evidence from 1 crossover RCT with 8 adults with cystic fibrosis showed no
3 clinically significant difference in nocturnal oxygen saturation between NIV and control after 6
4 weeks follow-up.

5 **Quality of life**

6 Low quality evidence from 1 crossover RCT with 8 adults with cystic fibrosis showed no
7 clinically significant difference in quality of life chest symptom score using the CF-QOL
8 questionnaire between NIV and control after 6 weeks follow-up.

9 Moderate quality evidence from 1 crossover RCT with 8 adults with cystic fibrosis showed no
10 clinically significant difference in quality of life traditional dyspnoea index score using the CF-
11 QOL questionnaire between NIV and control after 6 weeks follow-up.

12 **Hospitalisations**

13 No evidence was found for this important outcome.

14 **9.2.6.15 Economic evidence statements**

15 One cost-benefit analysis on people with cystic fibrosis in Canada found that PEP was less
16 expensive and more effective at reducing exacerbations than HFCWO over 1 year. This
17 analysis is partially applicable with serious limitations, given the limited details reported in the
18 conference abstract.

19 **9.2.7 Evidence to recommendations**

20 **9.2.7.1 Relative value placed on the outcomes considered**

21 The aim of this review was to examine the effectiveness of airway clearance techniques in
22 people with cystic fibrosis.

23 The Committee chose sputum volume, pulmonary exacerbations and patient preference as
24 critical outcomes for decision making; whereas pulmonary function (FEV₁, FVC), oxygen
25 saturation, hospitalizations and quality of life were rated as important outcomes.

26 **9.2.7.2 Consideration of clinical benefits and harms**

27 The Committee acknowledged the evidence, but they showed some concerns regarding the
28 quality of the studies, and its usefulness to make recommendations. They noted that the
29 studies included in the clinical review had a low number of participants and were conducted
30 over a short time frame. Moreover, none of the studies included children under the age of 6
31 and several studies excluded patients who were unstable, or recovering from an
32 exacerbation. The Committee also stressed that RCTs would not reflect clinical practice, as
33 techniques are normally individualised because they do not treat the same physiological
34 causes. Given that, the Committee noted that it would be difficult for the benefits from an
35 airway clearance technique to be demonstrated in randomised trials.

36 Based on the review, the Committee acknowledged that there was limited evidence either in
37 favour or against the use of routine airway clearance techniques in people with cystic
38 fibrosis. Apart from a clinically significant beneficial effect which favoured the use of PEP
39 mask over high frequency chest wall oscillation in the number of pulmonary exacerbations
40 (moderate quality evidence), no other clinically significant findings were found. The
41 Committee noted that there was low to moderate quality evidence that showed that using a
42 PEP mask was no better than no airway clearance technique in sputum volume, lung

1 function or oxygen saturation. With regards to the comparisons between different techniques,
2 very low quality evidence showed no clinically significant differences between manual
3 physiotherapy techniques and oscillating devices in lung function. Likewise, very low to high
4 quality evidence showed no clinically significant differences between PEP and oscillatory
5 devices in patient preference, pulmonary exacerbations, lung function, or quality of life.

6 It was also noted by the Committee that the effectiveness of airway clearance in other
7 conditions is not generalisable to those with cystic fibrosis, as cystic fibrosis is a condition
8 with specific clinical manifestations.

9 However, the Committee discussed that despite the lack of evidence showing effectiveness,
10 there was a strong rationale (physiological) that airway clearance techniques are useful in
11 children and adults who produce sputum. This is based on the knowledge that in cystic
12 fibrosis the normal mucociliary transport system is impaired and ineffective, so airway
13 clearance techniques are utilised to make up for the defects in this system and promote the
14 mobilisation of sputum from the airways to allow expectoration. Due to the dehydrated mucus
15 and airway damage, airway clearance techniques may reduce the risk of infection by
16 assisting the removal of bacteria in the sputum. The Committee suggested that the number
17 of trials comparing combinations of airway clearance techniques infers an underlying
18 knowledge and experience that individual airway clearance techniques are useful.

19 Moreover, the Committee argued that the benefits of airway clearance may not be
20 demonstrated by the amount of sputum produced. In other words the benefits of airway
21 clearance techniques are not always measurable because the person may just feel better
22 within themselves. For these reasons the Committee agreed not to make a “do not do
23 recommendation” despite the lack of favourable evidence.

24 Based on this, the Committee agreed it is important to discuss the use of airway clearance
25 techniques with people with cystic fibrosis who do not have lung disease.

26 The Committee also discussed which patients would benefit the most from airway clearance
27 techniques depending on their disease trajectory. The Committee noted that paediatric
28 practice has considerably changed in the last few years and some centres are more
29 comfortable with not instigating routine airway clearance with children that are asymptomatic
30 but rather teaching an airway clearance technique for use when needed and using more
31 structured exercise to promote airflow in the lungs and encourage close parental assessment
32 of the lungs. It was recognised that some clinicians believe that learning airway clearance
33 techniques at a young age helps to establish a daily routine to carry forward during
34 adulthood.

35 On the other hand, the Committee agreed that when a patient is symptomatic (i.e. has a
36 productive cough), or has received a treatment that produces sputum (such as mucolytic
37 treatment) airway clearance techniques have a strong rationale and are often helpful in
38 relieving symptoms of cough and breathlessness. The Committee agreed that there are a
39 number of factors that should be taken into account when choosing an airway clearance
40 technique.

41 First, they noted it was important to assess the patient’s symptoms (including stage of lung
42 disease and current health), and the ability of the person to clear mucus from their lungs. In
43 addition, they highlighted the difficulties of understanding the impact of other treatments and
44 lifestyle choices in patients with cystic fibrosis on airway clearance outcomes. They also
45 noted it is very important to take into account the individual preferences of the person and of
46 their parents or carers, as these may influence adherence.

47 The Committee agreed it is important to assess the effectiveness of the airway clearance
48 technique, and choose a different one if needed.

49 The Committee discussed at length the high frequency chest wall oscillation (trade name:
50 The Vest), as this technique was popular among patients in the USA (although not so in the

1 UK), but also has a high associated cost. They also noted that an increasing number of
2 patients and carers are buying The Vest privately, because it is only funded by the NHS in
3 exceptional circumstances when all other techniques have been exhausted. However the
4 evidence retrieved for this review did not support the use of high frequency chest wall
5 oscillation. No clinically significant differences was found between manual physiotherapy
6 techniques and high frequency chest wall oscillation in sputum volume (low quality
7 evidence). No clinically significant differences were found either between using oscillating
8 devices and high frequency chest wall oscillation in lung function (low to moderate quality).
9 Likewise, no clinically significant differences were found between PEP and high frequency
10 chest wall oscillation in sputum volume (low quality evidence), or lung function (very low to
11 moderate). In fact, moderate quality evidence from 2 trials showed that PEP was better at
12 reducing pulmonary exacerbations. Based on this, the Committee agreed that given the
13 current evidence The Vest should not be recommended as part of this guideline.

14 In relation to non-invasive ventilation (NIV), the Committee noted that this technique is not
15 widely used in clinical practice. Low to moderate quality evidence from 1 trial showed no
16 clinically significant beneficial effect of NIV over control in lung function, oxygen saturation
17 and quality of life. However, based on their experience and expertise, the Committee
18 considered that non-invasive ventilation could be used as an adjunct to support other airway
19 clearance techniques in severely symptomatic patients. This is because it is known that non-
20 invasive ventilation helps the lungs to eliminate carbon dioxide, and therefore it reduces the
21 symptoms associated with reduced oxygen such as headaches, drowsiness or poor quality
22 of sleep.

23 9.2.7.3 Consideration of economic benefits and harms

24 Techniques including ACBT, oscillating devices, and PEP can be performed at home after an
25 initial visit with a physiotherapist to issue the device and teach the techniques; hence, the
26 cost of performing combinations would be similar to single techniques. Moreover, there is no
27 increase in cost if those techniques are performed more frequently as no additional
28 resources are required.

29 The Committee noted that those relatively inexpensive techniques should not be performed if
30 they are ineffective, particularly when the opportunity cost of the patient's time is considered.
31 Performing long, regular periods of airway clearance could reduce adherence to other
32 treatments and that could potentially reduce the effects of those treatments. However, the
33 Committee agreed that lack of evidence does not mean lack of effect, and hence, lack of
34 cost-effectiveness and made a recommendation to take account preferences and any factors
35 that may influence adherence, if an airway clearance technique is considered appropriate.

36 The Vest and NIV were the most expensive airway clearance techniques included in this
37 review. The Committee agreed there was clinical and cost-effectiveness evidence to suggest
38 The Vest was dominated (more expensive and less effective) by PEP. Subsequently, the
39 Committee made a recommendation against its use, to prevent a cost ineffective use of NHS
40 resources. The Committee added that their recommendation does not prevent private
41 funding which may be sought in cases, such as neuro-disability, where The Vest may be
42 considered as a last resort by parents and carers.

43 The Committee noted that the additional benefits of NIV outweigh its additional cost
44 compared to the other techniques, in people who have severe lung disease. As a result, the
45 Committee made a recommendation to think about the use of NIV in people with severe lung
46 disease, as a subgroup where NIV would be cost-effective. Following this, the Committee
47 added that not all cystic fibrosis centres will have access to NIV machines for airway
48 clearance, but would have access to NIV for respiratory failure. Consequently, the
49 recommendation would prompt readers to consider its alternative use to support airway
50 clearance techniques in those that are using it for sleep.

1 **9.2.7.4 Quality of evidence**

2 The quality of the evidence presented in this report ranged from very low to high as assessed
3 by GRADE. The main reasons that led to downgrading the quality of the evidence were:

4 For the domain risk of bias, the studies were assigned the same risk of bias as in the
5 Cochrane reviews, and were not individually reviewed. The main biases that lead to
6 downgrading the quality of the evidence were selection, attrition, and reporting bias.

7 Another reason that lead to downgrading the quality of the evidence was the imprecision, as
8 confidence intervals crossed 1 or 2 MIDs. The Committee noted that many trials were
9 underpowered to detect a clinically important difference.

10 No serious issues were found regarding inconsistency (heterogeneity) and the directness of
11 the population (generability of the results).

12 **9.2.7.5 Other considerations**

13 No equality issues were identified by the Committee for this review question.

14 The Committee discussed the need to draft a research recommendation for this topic. Since
15 the advent of newborn screening for cystic fibrosis there has been international debate about
16 the level of physiotherapy intervention required from diagnosis to preserve lung health. Some
17 clinical teams opt to teach and recommend daily airway clearance techniques, others, for
18 example, use parental respiratory assessment tools with structured exercise. Routine airway
19 clearance from diagnosis places considerable responsibility and time burden on the parents
20 and carers at a time when such techniques are challenging to perform and negotiate with the
21 infant and child. It is important that we fully understand if routine practice is providing benefit
22 to maintain lung health or in fact creating unnecessary burden. Future research must seek to
23 understand the impact not only on long term clinical outcomes, but on the lives of parents,
24 families and carers of infants and children with cystic fibrosis.

25 **9.2.7.6 Key conclusions**

26 The Committee concluded that there was limited evidence either in favour or against the use
27 of airway clearance techniques in patients with cystic fibrosis however, there is a strong
28 physiological rationale for airway clearance techniques and they continue to be used
29 routinely in the symptomatic patient. The use of manual physiotherapy techniques, PEP
30 mask, ACBT and AD were prioritised by the Committee; whereas they agreed The Vest
31 should not be recommended due to evidence that is inferior to other airway clearance
32 techniques and is costly. The decision to choose one technique over another would be
33 mainly based on individual factors (for example age) and the physiological problem or
34 circumstances at the time, rather than one technique being superior to another. Individual
35 preferences should also be taken into account, as this may impact adherence. Non-invasive
36 ventilation could be used as an adjunct and in combination with airway clearance techniques
37 in people who are limited by symptoms such as breathlessness and fatigue due to moderate
38 or severe lung disease.

39 **9.2.8 Recommendations**

40 **50. Discuss the use of airway clearance techniques with people with cystic fibrosis**
41 **who do not have lung disease and with their parents or carers (as appropriate).**

42 **51. Offer training in an airway clearance technique to people with cystic fibrosis who**
43 **have lung disease and their parents or carers (as appropriate).**

44 **52. When choosing an airway clearance technique for people with cystic fibrosis:**

- 1 • assess their ability to clear mucus from their lungs, and offer an
 - 2 individualised plan to optimise this
 - 3 • take account of the preferences of the person and (if appropriate) their
 - 4 parents and carers
 - 5 • take account of any factors that may influence adherence.
- 6 **53. Regularly assess the effectiveness of airway clearance techniques, and modify the**
- 7 **technique or use a different one if needed.**
- 8 **54. Do not offer high-frequency chest wall oscillation as an airway clearance**
- 9 **technique for people with cystic fibrosis.**
- 10 **55. Think about using non-invasive ventilation to help with airway clearance**
- 11 **techniques in people who have moderate or severe lung disease.**

12 9.2.9 Research recommendations

- 13 **2. What is the clinical and cost effectiveness of daily airway clearance techniques in**
- 14 **maintaining lung function in infants and children with cystic fibrosis?**

15 **Table 76: Research Recommendation justification**

Research question	What is the clinical and cost effectiveness of daily airway clearance techniques in maintaining lung function in infants and children with cystic fibrosis?
Why this is needed	
Importance to 'patients' or the population	Performing daily airway clearance techniques places considerable responsibility and time burden on parents and carers at a time when such techniques are challenging to perform and negotiate with the infant and child with cystic fibrosis. It is essential that parents, carers and people with CF are reassured that this level of input is based on proven clinical benefit. Parent, carers and people with CF frequently report in the literature that daily airway clearance routines are difficult to adhere to and incorporate into daily life. If routine airway clearance is not deemed necessary until a certain time point, this could have a significant impact on the quality of life of parents, carers and people with CF.
Relevance to NICE guidance	High: With this evidence, definitive recommendations could be made regarding 'routine' vs 'when required' use of airway clearance techniques – at present there is varied clinical practice throughout the UK
Relevance to the NHS	CF services need to be able to direct and plan the physiotherapy input to people with CF and their families/carers to ensure that the resources are placed where there is proven clinical benefit rather than basing service delivery on historical behaviours. Further evidence in this area will eliminate the need to provide an intervention 'just in case' it preserves lung health but in fact physiotherapists will be able to use the evidence to target the right patients with the right intervention based on evidence of the likelihood of improved health outcomes. There will be a financial advantage if less airway clearance devices are required in the future due to a reduction in routine airway clearance approaches in the UK. There may be a small financial disadvantage if more airway clearance devices are required if routine airway clearance is proven from diagnosis to benefit health outcomes. There is unlikely to be any change in the physiotherapy input required as interventions will continue to include exercise guidance, advice on assessment/monitoring of lung health and teaching of airway clearance

Research question	What is the clinical and cost effectiveness of daily airway clearance techniques in maintaining lung function in infants and children with cystic fibrosis?
	techniques for when they are required.
National priorities	This research question is similar to one of the top 10 research priorities published by the James Lind Alliance in January 2017 – ‘Can exercise replace chest physiotherapy for people with CF’ The comparative group to ‘routine airway clearance’ is likely to include structured exercise so could help answer the above question for infants and children.
Current evidence base	The current evidence base includes mostly small, short term studies with limited methodological rigour and are unable to draw conclusions regarding the long-term impact of airway clearance techniques on lung health in people with CF. Most of the studies include people with respiratory symptoms and established lung disease and therefore it is also not possible to apply this knowledge to the asymptomatic person without evidence of lung disease. The current evidence base compares the different types of airway clearance techniques but has yet to answer the question whether airway clearance as an intervention provides any long term clinical benefit to lung health either in the symptomatic or indeed the asymptomatic person with CF.
Equality	CF centres across the UK have different approaches to routine airway clearance – further evidence will help direct care and potentially improve equity of access of evidence based physiotherapy interventions.
Feasibility	The research could be carried out within a realistic timescale and with realistic funding
Other comments	None

1

Table 77: Research Recommendation Statements

Criterion	Explanation
Population	Infants and children with a diagnosis of cystic fibrosis – to be confirmed by specified criteria
Intervention	<ul style="list-style-type: none"> o Daily airway clearance techniques
Comparators	Airway clearance techniques only when specified criteria met. For example, based on: <ul style="list-style-type: none"> - Disease severity - Asymptomatic/ symptomatic
Outcomes	<ul style="list-style-type: none"> • Expectorated secretions (mucus, sputum, phlegm) • Sputum volume • Hospitalisations, change in frequency • Pulmonary exacerbations, change in frequency • Lung function (FEV₁, FVC) <ul style="list-style-type: none"> o forced vital capacity (FVC) • Oxygen saturation measured by pulse or transcutaneous oximetry • Quality of Life (using a validated tool, such CFQ-R or CF-QOL) • Patient preference • Resource use • Unit costs
Study design	Multicentre RCT
Timeframe	2-6 years

1 9.3 Mucoactive agents

2 **Review question: What is the effectiveness of mucoactive or mucolytic agents,**
3 **including dornase alfa, nebulised sodium chloride (isotonic and hypertonic) and**
4 **mannitol?**

5 9.3.1 Introduction

6 In lungs with cystic fibrosis, the underlying defect leads to an increase in water absorption
7 from the epithelial surface, resulting in a reduced airway surface liquid layer, and a more
8 viscous mucus layer on the surface of the airways. This mucus accumulates due to reduced
9 clearance, which supports the retention of micro-organisms. In turn, this leads to infection,
10 and the destructive inflammatory processes which lead to bronchiectasis.

11 The primary aims of pulmonary disease management in people with cystic fibrosis is to
12 stabilise or prevent decline in pulmonary function, and to prevent the occurrence of acute
13 pulmonary exacerbations. Therefore, people with cystic fibrosis who have evidence of
14 pulmonary disease commonly employ airway clearance techniques to reduce the burden of
15 viscid mucus, and break the destructive cycle of mucus stasis, infection and inflammation.

16 Mucoactive agents are often employed as adjuncts to airway clearance techniques. These
17 agents change the properties of mucus, through a number of mechanisms, rendering it more
18 easy to expectorate: dornase alfa is a recombinant human enzyme which acts by cleaving
19 extracellular DNA (a by-product of neutrophil degeneration) in the mucus; whilst osmotic
20 agents such as mannitol and hypertonic sodium chloride draw water onto the airway surface
21 to rehydrate the airway surface liquid layer; solutions of hypertonic sodium chloride also
22 disrupt ionic bonds within the mucus gel.

23 With a number of mucoactive agents available, this review question aims to determine the
24 effectiveness of mucoactive or mucolytic agents – including dornase alfa, inhaled sodium
25 chloride solutions (both isotonic, and hypertonic) and inhaled mannitol – and therefore
26 determine their place in the management of cystic fibrosis pulmonary disease.

27 9.3.2 Description of clinical evidence

28 The aim of this review was to establish the clinical and cost effectiveness of mucoactive or
29 mucolytic agents in improving airway clearance in children, young people and adults with
30 cystic fibrosis.

31 The nebulised and inhaled mucoactive and mucolytic agents reviewed were: acetylcysteine,
32 dornase alfa, nebulised sodium chloride (hypertonic and isotonic) and mannitol (only in
33 children and young people up to the age of 18 years as Technology Appraisal (TA) in adults
34 will be included).

35 NICE TA266 has been published to provide guidance on the use of mannitol dry powder for
36 inhalation for the treatment of cystic fibrosis in adults. The following comparisons were
37 considered:

- 38 • Mannitol versus placebo
 - 39 ○ 2 trials (DPM-CF-301, DPM-CF-302) were included in the TA to assess the
 - 40 effectiveness of mannitol
 - 41 ○ another 4 trials were excluded from the TA because of their short duration (DPM-CF-
 - 42 201, DPM-CF-202), population (children only) (DPM-CF-203), and low dose of mannitol
 - 43 and short duration of treatment (Robinson 1999); these trials have been retrieved for
 - 44 potential inclusion in this review.
- 45 • Mannitol versus other treatments

- 1 ○ no trials were included
- 2 – 1 trial (Robinson 1999) was identified in the TA for the comparison mannitol versus
- 3 hypertonic sodium chloride, but was excluded due to low dosage, short treatment
- 4 duration and small population
- 5 – 2 trials (CF-301, Jaques 2008) compared mannitol to control (control = low dose
- 6 mannitol) and 8 trials (Button 1996, Chadwick 1997, Elkins 2006, Eng 1996, Riedler
- 7 1996, Robinson 1996, Robinson 1997, Robinson 1999) compared hypertonic
- 8 sodium chloride to control; however, the Technology Assessment Group (TAG,
- 9 Riemsma 2011) agreed that the results were not comparable due to the difference in
- 10 duration (26 weeks versus < 2 weeks).

11 A report from the National Horizon Scanning Centre (NHSC 2008) was also identified. This

12 report included 4 trials that have been retrieved for assessment.

13 Systematic reviews of randomised controlled trials (RCTs) and RCTs, including cross-over

14 trials were considered for this review. Systematic reviews were assessed for inclusion

15 against the protocol, and if relevant, their quality was assessed using AMSTAR. High-quality

16 systematic reviews were included in our review, and where possible, data was taken directly

17 from the review. Individual studies were also retrieved for completeness and accuracy. Low-

18 quality systematic reviews were excluded from the review, but the list of included studies was

19 checked to identify relevant trials.

20 Three Cochrane reviews were identified and included in this review:

- 21 • Yang 2016 evaluated the effectiveness of dornase alfa compared to placebo and other
- 22 mucoactive or mucolytic agents. Twelve trials were included from this review (Amin 2011;
- 23 the trial reported in both Ballmann 1998 and Ballmann 2002; Fuchs 1994, Laube 1996,
- 24 McCoy 1996, Minasian 2010; Quan 2011; Ramsey 1993a; Ranasinha 1993, Shah 1995;
- 25 the trial reported in Suri 2001, Suri 2002a and Suri 2002b; Wilmott 1996).
- 26 • Nolan 2015 evaluated the effectiveness and safety of inhaled mannitol in people with
- 27 cystic fibrosis. Four trials have been included from this review (Aitken 2012, Bilton 2011,
- 28 Jaques 2008, Minasian 2010). Two of the trials had already been included in the TA report
- 29 (Aitken 2012, Bilton 2011) and were used to retrieve additional outcomes.
- 30 • Wark 2010 evaluated the effectiveness of nebulised hypertonic sodium chloride compared
- 31 to placebo and other mucoactive or mucolytic agents. Three trials have been included
- 32 from this review (the trial reported in both Ballmann 1998 and Ballmann 2002; Elkins
- 33 2006; and the trial reported in Suri 2001, Suri 2002a and Suri 2002b).

34 Four (non-Cochrane) systematic reviews were also identified (Christopher 1999, Cramer

35 1996, Duijvestijn 1999, Taylor 200). All of them were excluded due to their low

36 methodological quality, and the list of included studies checked for their potential inclusion.

37 In addition, 9 trials were identified for inclusion (Amin 2010, Conrad 2015, Dentice 2016,

38 Gupta 2012, Mainz 2016, Ratjen 1985, Rosenfeld 2012, Shah 1996, Skov 2015).

39 The size of the trials ranged from 14 to 968 participants. Twelve trials included children,

40 young people and adults (Aitken 2012, Bilton 2011, Conrad 2015, Elkins 2006, Fuchs 1994,

41 Jaques 2008, Mainz 2016, McCoy 1996; Ramsey 1993a; Ratjen 1985, Shah 1995a, Wilmott

42 1996), 1 trial included infants and children (Rosenfeld 2012), 1 trial included children only

43 (Quan 2001), 2 studies included children and young people (Gupta 2012; and the trial

44 reported in Suri 2001, Suri 2002a and Suri 2002b), 3 trials included children and young

45 people and adults aged 18 only (Amin 2010, Amin 2011, Minasian 2010), 2 trials included

46 adults only (Laube 1996, Skov 2015), 3 trials included young people and adults (Dentice

47 2016, Ranasinha 1993, Shah 1996). The age range was not reported for 1 trial (reported in

48 both Ballmann 1998 and Ballmann 2002 – mean age 13.3).

49 Five trials were conducted in the United States (Conrad 2015, Fuchs 1994, Laube 1996,

50 McCoy 1996, Wilmott 1996), 3 in the UK (Minasian 2010; the trial reported in Suri 2001, Suri

1 2002a and Suri 2002b; and Shah 1996), 2 in Canada (Amin 2010, Amin 2011), 2 in Australia
2 (Dentice 2016, Elkins 2006), 1 in India (Gupta 2012), 1 in Denmark (Skov 2015) and 3 in
3 Germany (Mainz 2016, Ratjen 1985; and the trial reported in both Ballmann 1998 and
4 Ballmann 2002). The following 6 studies were conducted in multiple countries: 1 in the United
5 States, Canada, Argentina and Europe (Aitken 2012), 1 in Australia, New Zealand, UK and
6 Ireland (Bilton 2011), 1 in Australia and New Zealand (Jaques 2008), 1 in the United States,
7 Canada and the UK (Shah 1995a), 1 in the United States and Canada (Rosenfeld 2012), 1 in
8 Australia, Belgium, Canada, Denmark, Germany, Ireland, Israel, Netherlands, Norway,
9 Spain, Switzerland and the United States (Quan 2001). For 2 trials the country was not
10 reported (Ramsey 1993a and Ranasinha 1993).

11 The included studies assessed the effectiveness of mucoactive or mucolytic agents based on
12 the following comparisons:

- 13 • mannitol versus placebo – 3 trials (Aitken 2012, Bilton 2011, Jaques 2008)
- 14 • mannitol versus dornase alfa – 1 trial (Minasian 2010)
- 15 • mannitol plus dornase alfa versus dornase alfa – 1 trial (Minasian 2010)
- 16 • dornase alfa versus placebo – 10 trials (Amin 2011, Fuchs 1994, Laube 1996, McCoy
17 1996, Ramsey 1993a, Ranasinha 1993, Quan 2001, Shah 1995, Shah 1996, Wilmott
18 1996)
- 19 • dornase alfa versus nebulised sodium chloride – 2 trials (trial reported in both Ballman
20 1998 and Ballmann 2002; and trial reported in Suri 2001, Suri 2002a and Suri 2002b)
- 21 • nebulised sodium chloride (>3% concentration) versus placebo (0.9% to 0.12%) or low
22 concentration ($\leq 3\%$)
 - 23 ○ nebulised sodium chloride (7% concentration) versus placebo (0.9% sodium chloride
24 concentration) – 3 trials (Amin 2010, Elkins 2006, Rosenfeld 2012)
 - 25 ○ nebulised sodium chloride (7% concentration) versus placebo (0.12% sodium chloride
26 concentration) – 1 trial (Dentice 2016)
 - 27 ○ nebulised sodium chloride (7% concentration) versus low concentration (3%) – 1 trial
28 (Gupta 2016)
 - 29 ○ nebulised sodium chloride (6% concentration) versus placebo (0.9% sodium chloride
30 concentration) – 1 trial (Mainz 2016)
- 31 • acetylcysteine versus placebo – 3 trials (Conrad 2015, Ratjen 1985, Skov 2015)

32 A summary of the studies included in this review are presented in Table 78. See study
33 selection flow chart in Appendix F, study evidence tables in Appendix G, list of excluded
34 studies in Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

35 9.3.3 Summary of included studies

36 A summary of the studies that were included in this review are presented in Table 78.

37 **Table 78: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
NICE Technology Appraisal				
NICE Technical Appraisal (TA Mannitol 2012)	Comparison 1: Mannitol versus placebo • CF-301 • CF-302 Comparison 2:	Adults with CF.	Comparison 1: Mannitol versus placebo • Change in FEV ₁ • Reduction in pulmonary exacerbations • Reduction of AB and	CF-301: Bilton 2011 CF-302: Aitken 2012 This TA was reviewed in Riemsma et al., Mannitol dry

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Mannitol versus other treatments No studies were included.		hospitalization days <ul style="list-style-type: none"> Quality of life Safety Comparison 2: Mannitol versus other treatments n/a	powder for inhalation for the treatment of cystic fibrosis. Kleijnen Systematic Reviews Ltd., 2011
Cochrane systematic reviews				
Yang 2016 Cochrane SR	Comparison 1: Dornase Alfa versus placebo <ul style="list-style-type: none"> Amin 2011 Ballmann 1998 and Ballmann 2002 Fuchs 1994 Laube 1996 McCoy 1996 Minasian 2010 Quan 2001 Ramsey 1993a Ranasinha 1993 Shah 1995a Wilmott 1996 (non-NMA) Comparison 2: Dornase alfa versus hypertonic sodium chloride <ul style="list-style-type: none"> Suri 2001, Suri 2002 and Suri 2002a 	Children and adults with CF (diagnosed clinically and by sweat or genetic testing) and with all stages of lung disease were included.	Comparison 1: Dornase Alfa versus placebo <ul style="list-style-type: none"> FEV₁ change People experiencing exacerbations Use of antibiotics Adverse events Change in quality of life Comparison 2: Dornase alfa versus hypertonic sodium chloride <ul style="list-style-type: none"> FEV₁ change Quality of life Days of inpatient treatment 	AMSTAR: 10/11
Nolan 2015 Cochrane SR	Comparison 1: Mannitol versus placebo <ul style="list-style-type: none"> Aitken 2012 Bilten 2011 Comparison 2: Mannitol versus Dornase Alfa <ul style="list-style-type: none"> Minasian 2010 Comparison 3: Mannitol + Dornase Alfa versus Dornase Alfa <ul style="list-style-type: none"> Jaques 2008 	Children and adults with CF (diagnosed clinically and by sweat or genetic testing) and including all degrees of disease severity.	Comparison 1: Mannitol versus placebo <ul style="list-style-type: none"> FEV₁ ml, % predicted Pulmonary exacerbations Time to first pulmonary exacerbation Number of patients needing AB/hospitalisation HRQOL – 11 domains Adverse events Comparison 2:	AMSTAR: 11/11

Study	Intervention/ Comparison	Population	Outcomes	Comments
			Mannitol versus Dornase Alfa <ul style="list-style-type: none"> • FEV₁% change Comparison 3: Mannitol + Dornase Alfa versus Dornase Alfa <ul style="list-style-type: none"> • FEV₁% change 	
Wark 2010 Cochrane SR	Comparison 1: Nebulised hypertonic sodium chloride versus isotonic sodium chloride <ul style="list-style-type: none"> • Elkins 2006 Comparison 2: hypertonic sodium chloride versus dornase alfa (dornase alfa) <ul style="list-style-type: none"> • Ballmann 1998 and Ballmann 2002 • Suri 2001, Suri 2002a and Suri 2002b 	Children and adults with CF (diagnosed clinically and by sweat or genetic testing) and including all degrees of disease severity.	Comparison 1: Nebulised hypertonic sodium chloride versus isotonic sodium chloride <ul style="list-style-type: none"> • FEV₁ • Inflammatory markers • Quality of life • Hospital admissions • Adverse events Comparison 2: hypertonic sodium chloride versus dornase alfa (dornase alfa) <ul style="list-style-type: none"> • FEV₁ • Quality of life • Hospital admissions 	AMSTAR: 9/11 The update for this review was not yet published when we carried out our review. Cochrane kindly provided us with a copy of the update. We concluded that the relevant studies included in the update were also included in Wark 2010.
Individual studies included in the TA or the Cochrane SR				
CF-301: Bilton 2011 (Australia, New Zealand, UK, Ireland) Multi-centre RCT, parallel design	Intervention: Inhaled dry powder mannitol, 400 mg BD for 26 weeks Comparison: Subtherapeutic mannitol (mannitol 50 mg)	N=295 people with CF were randomized Mean (SD) age: 23 (11.3) years. Age ≥6 years. Participants were clinically stable at start of study.	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Time to first protocol defined pulmonary exacerbation • Number of patients needing additional IV antibiotics • Change in quality of life (HRQOL – CFQOL) • Adverse events: <ul style="list-style-type: none"> ○ bronchospasm (mild, moderate and severe) ○ haemoptysis (mild, moderate and severe) 	In NICE TA on Mannitol & in Cochrane SR Nolan 2015 Participants receiving hypertonic saline were excluded. High dropout rate, however sensitivity analysis showed a consistent treatment effect with no change to conclusions.
CF-302: Aitken 2012 (USA, Canada,	Intervention: Inhaled dry mannitol, 400 mg BD for 26 weeks	N= 305 people with CF from 53 sites were randomized	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Time to first protocol defined pulmonary 	In NICE TA on Mannitol & in Cochrane SR Nolan 2015

Study	Intervention/ Comparison	Population	Outcomes	Comments
Argentina, Europe) Multi-centre RCT, parallel design	Comparison: Subtherapeutic mannitol (mannitol 50 mg)	Mean age: 20 years. Age range: 6 to 53. Participants were clinically stable at start of study.	exacerbation <ul style="list-style-type: none"> • Number of patients needing additional IV antibiotics • Change in quality of life (HRQOL – CFQOL) • Adverse events: <ul style="list-style-type: none"> ◦ haemoptysis (mild, moderate and severe) 	Participants receiving hypertonic saline were excluded. High dropout rate, however sensitivity analysis showed a consistent treatment effect with no change to conclusions.
Amin 2011 (Canada) RCT, crossover design	Intervention: Dornase alfa, 2.5 mg OD for 4 weeks intervention/comparison, 4-week washout period Comparison: Placebo	N=19 people with CF were randomized. Age range: 6 to 18years Baseline FEV ₁ ≥ 80% predicted	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in quality of life (measured with CFQ-R questionnaires) 	In Cochrane SR Yang 2016 Data only reported on 17 people who completed the trial.
Ballmann 1998 and Ballmann 2002 (Germany) RCT, crossover design	Intervention 1: Nebulization of 2.5 mg dornase alfa OD 3 weeks intervention, 3-week washout period Intervention 2: Nebulization of 10 ml 5.85% sodium chloride OD	N=14 people with CF Mean age: 13.3. Age range not reported. With mild to moderate pulmonary involvement.	<ul style="list-style-type: none"> • % change in FEV₁ 	In Wark 2010 and Yang 2016 Withdrawals were not discussed within the paper. Both interventions preceded by 2 puffs salbutamol via a spacer
Elkins 2006 (Australia) RCT, parallel design	Intervention: 7% sodium chloride BD 48 weeks Comparison: 0.9% sodium chloride BD	N=164 people with CF from 16 hospitals were randomized. Age ≥6 years Participants were in clinically stable condition.	<ul style="list-style-type: none"> • % change in FEV₁ • Change in quality of life (measured with CFQ-R questionnaires) 	In Wark 2010 2 people were excluded from the analysis: 1 person in each group voluntarily withdrew before first dose. 17 people were lost to follow-up. 15 people stopped inhalation but continued visits.
Fuchs 1994 (USA) RCT, parallel design	Intervention: Nebulized dornase alfa 2.5 mg OD or BD	N= 968 people with CF were randomized. Age: Over 5.	<ul style="list-style-type: none"> • Relative mean % change in FEV₁ • Number of people experiencing 	In Cochrane SR Yang 2016 25 people withdrew from

Study	Intervention/ Comparison	Population	Outcomes	Comments
	24 weeks Comparison: Placebo	FVC: > 40% predicted Clinically stable	exacerbations • Adverse events: haemoptysis, voice alteration	the study, 8 in the placebo group and once- daily group and 9 in the twice- daily group. All participants were included in the analysis.
Jaques 2008 (Australia and New Zealand) Multi-centre RCT, crossover design	Intervention: Inhaled dry powder mannitol 420mg BD for 2 weeks, 2- week washout period Comparison: Non-respirable Mannitol	39 people with CF from 7 sites were randomized. Mean (range)age: 19.1 (range 8 to 48) years Participants were clinically stable at start of study	• Change in FEV ₁ % predicted • Adverse events: haemoptysis (mild and severe)	In Cochrane SR Nolan 2015 No hypertonic saline within 2 weeks of start of study. 4 withdrawals.
Laube 1996 (USA) RCT, parallel design	Intervention: 2.5 mg nebulized dornase alfa BD for 6 days Comparison: placebo	N= 20 adults with CF Age: Over 18 With stable stage CF. FVC: 35%-75% predicted No withdrawals.	• Relative mean % change in FEV ₁ (overall and for subgroup with moderate disease)	In Cochrane SR Yang 2016 There were no withdrawals.
McCoy 1996 (USA) RCT, parallel design	Intervention: Nebulized dornase alfa 2.5 mg OD for 12 weeks Comparison: Placebo	N= 320 people with CF Age: 7 to 57 FVC: < 40% predicted.	• Relative mean % change in FEV ₁ • Number of days of IV antibiotic use • Adverse events: voice alteration	In Cochrane SR Yang 2016 40 participants withdrew from the trial.
Minasian 2010 (UK) RCT, crossover design	Intervention1: Mannitol 400mg BD Intervention 2: Dornase alfa alone: 2.5 mg BD Intervention 3: Mannitol (as above) plus dornase alfa (dose unclear) For 12 weeks	N=28 people with CF randomised Mean (SD) age: 13.3 (2.24) years. Age for eligibility was between 8 and 18 years. Participants were clinically stable at start of study. Currently receiving dornase alfa or having an FEV ₁ >40% and <70% predicted (and therefore eligible to receive	• FEV ₁ % change from baseline	In Cochrane SR Nolan 2015 and Cochrane SR Yang 2016 Participants using hypertonic saline were excluded. 45 were recruited but only 28 were randomised. 8 participants withdrew.

Study	Intervention/ Comparison	Population	Outcomes	Comments
		dornase alfa).		
Quan 2001 (Australia, Belgium, Canada, Denmark, Germany, Ireland, Israel, Netherlands, Norway, Spain, Switzerland and the United States) Multi-centre RCT, parallel design	Intervention: 2.5 mg dornase alfa OD for 96 weeks Comparison: Placebo	N=474 children with CF randomized from 49 centres Age: 6 to 10 (mean age 8.4) FVC>85% predicted	<ul style="list-style-type: none"> • Absolute mean % change in FEV₁ • Number of people experiencing exacerbations • Adverse events: voice alteration 	In Cochrane SR Yang 2016 <ul style="list-style-type: none"> • 410 completed the study. 60 participants withdrew from the study. The ITT population was 470.
Ramsey 1993a (Country not reported) RCT, parallel design	Intervention: Nebulized dornase alfa 0.6 mg, 2.5 mg or 10 mg BD for 10 days Comparison: placebo	N= 181 people with CF Age: 8 to 65 Stable stage CF FVC>= 40% predicted	<ul style="list-style-type: none"> • Relative mean % change in FEV₁ (overall results and subgroup analysis for people with moderate disease severity) • Adverse events: voice alteration 	In Cochrane SR Yang 2016 No withdrawals.
Ranasinha 1993 (Country not reported) RCT, parallel design	Intervention: Nebulized dornase alfa 2.5 mg BD for 10 days Comparison: placebo	N= 71 people with CF Age range: 16 to 55 All participants had stable disease FVC > 40% predicted	<ul style="list-style-type: none"> • Relative mean % change in FEV₁ (overall results and subgroup analysis for people with moderate disease severity) • Adverse events: haemoptysis, voice alteration 	In Cochrane SR Yang 2016
Shah 1995a (USA, Canada, UK) Multi-centre RCT, parallel design	Intervention: 2.5 mg nebulised dornase alfa BD for 14 days Comparison: Placebo	N: 70 people with CF randomized from 3 sites. Age: ≥5 years Severe (FVC < 40% predicted) lung disease	<ul style="list-style-type: none"> • Relative mean % change in FEV₁ (Overall results and subgroup analysis for people with severe disease) • Adverse events: haemoptysis, voice alteration 	In Cochrane SR Yang 2016 5 dropouts (2 died, 2 withdrew consent, 1 had a heart lung transplant)
Suri 2001, Suri 2002a and Suri 2002b (UK) RCT, crossover design	Intervention 1: 2.5 mg dornase alfa OD; Intervention 2: alternate day 2.5 mg dornase alfa; Intervention 3:	N= 48 children and young people with CF were randomised. Age range: 7.3 to 17.	<ul style="list-style-type: none"> • Mean % change in FEV₁ • Number of days of inpatient treatment 	In Wark 2010 and Yang 2016 45 completed first treatment period, 44 completed the second treatment period and 40

Study	Intervention/ Comparison	Population	Outcomes	Comments
	5 mL 7%hypertonic saline BD 12-week treatment periods with a 2-week washout period between each period			completed the third treatment period.
Wilmott 1996 (USA) RCT, parallel design	Intervention: Dornase alfa 2.5 mg nebulised BD over 15 days Comparison: Placebo	N= 80 people with CF from 11 CF centres Age: Over 5 Admitted to hospital for at least 1 night for treatment of a chest exacerbation (protocol defined) with FVC > 35% predicted Participants in both groups had a moderate clinical degree of dyspnoea.	<ul style="list-style-type: none"> • Mean % change in FEV₁ 	In Cochrane SR Yang 2016 No withdrawals mentioned in the paper
Additional primary studies				
Amin 2010 (Canada) RCT, crossover design	Intervention: 7% sodium chloride BD for 4 weeks, 4-week washout period Comparison: 0.9% sodium chloride	N=20 people with CF randomized. Mean (SD) age: 10.5 (3.1). Age range for eligibility: 6 to 18 years. Baseline FEV ₁ % predicted ≥80%	<ul style="list-style-type: none"> • Quality of life (Measured with CFQ-R questionnaire) 	1 person was excluded from the analysis.
Conrad 2015 (USA) RCT, parallel design	Intervention: Acetylcysteine 3 times daily for 24 weeks Comparison: Placebo	N0=70 people with CF were randomized. Age range: 9 to 59 years. Stable mild-moderate lung disease; FEV ₁ ≥40% and ≤85% predicted	<ul style="list-style-type: none"> • Change in FEV₁ • Change in sputum IL-8 (log10) • Incidence of pulmonary exacerbations • Quality of life (Measured with CFQ-R) 	6 in the acetylcysteine group and 2 in the placebo group were withdrawn or lost to follow-up. Of these, 1 in the placebo group was excluded from the analysis.
Dentice 2016 (Australia) RCT, parallel	Intervention: 7% sodium chloride 4ml	N=132 people with CF admitted for management of	<ul style="list-style-type: none"> • Failed to regain pre-exacerbation FEV₁% predicted • Change in quality of 	All participants were included in the analysis related to

Study	Intervention/ Comparison	Population	Outcomes	Comments
design	3 times daily throughout hospital admission Control: 0.12% sodium chloride Both groups received usual care.	a pulmonary exacerbation. Mean age (SD), range: <ul style="list-style-type: none">• Intervention: 28 (9), 17 to 62 years• Control: 27 (9), 18 to 63 years	life <ul style="list-style-type: none">• Time to first pulmonary exacerbation	relevant outcomes. On average, hospital admission lasted 12 days in intervention group and 13 in control group
Gupta 2012 (India) RCT, parallel design	Intervention: 7% sodium chloride BD for 28 days (high dose) Comparison: 3% sodium chloride	N=31 children and young people with CF were randomized. Age range: 6 to 16 years Able to perform reproducible pulmonary function test was among inclusion criteria. No inclusion criteria relating to severity of lung disease.	<ul style="list-style-type: none">• % change in FEV₁	30 people completed the study and were included in the analysis..
Mainz 2016 (Germany) RCT, crossover design	Intervention: 6% sodium chloride OD for 28 days, 28-day washout period Comparison: 0.9% sodium chloride	N=69 people with CF with chronic rhinosinusitis Age ≥6 years. Mean age (SD): 22.8 (12).	<ul style="list-style-type: none">• FEV₁ % predicted	5 people were excluded from the analysis.
Ratjen 1985 (Germany) RCT, parallel design	Intervention: Acetylcysteine 3 times daily for 12 weeks Comparison: Placebo	N=21 people with CF were included in the analysis. Mean (range) age: 13.9 (6 to 21) People with mild to moderate lung disease.	<ul style="list-style-type: none">• Change in FEV₁	Total N in the study was 36 people who were randomized to 3 interventions (1 not relevant to this review). There were 4 withdrawals out of 36 people; 1 in placebo group, 3 unclear.
Rosenfeld 2012 (USA and Canada)	Intervention: 7% sodium chloride BD for 48 weeks	N=321 infants and children with CF were randomized.	<ul style="list-style-type: none">• Time to first pulmonary exacerbation• Number of days of	29 withdrew (15 in intervention group and 14 in comparison group) but

Study	Intervention/ Comparison	Population	Outcomes	Comments
Multicentre RCT, parallel design	Comparison: 0.9% sodium chloride	Age: < 6 years	treatment for a pulmonary exacerbation • Change in quality of life (Measured with CFQ-R)	everyone was included in the analysis.
Shah 1996 (UK) RCT, parallel design	Intervention 1: 2.5 mg dornase alfa bd BD for 10 days Placebo: 150 mmol sodium chloride, 1.5 mmol calcium chloride	N: 71 people with CF randomized. Age: >15 years Stable condition prior for 14 days prior to enrolment	• Relative mean % change in FEV ₁	30 people were excluded from the analysis due to inadequate sputum samples.
Skov 2015 (Denmark) Open trial, parallel design	Intervention: Acetylcysteine 2400 mg/day for 4 weeks Comparison: Placebo	N=21 people with CF n=11 in the NAC group and n=10 in the placebo group Median age: 39 years (range 25 to 61)	• Change in FEV ₁ % predicted	There were 2 withdrawals, 1 in the intervention and 1 in the placebo group.

1 Abbreviations: BD: twice daily; CF: cystic fibrosis; FEV: forced expiratory volume; N: number; OD: once daily;
2 RCT: randomised controlled trial; SR: systematic review

3 9.3.4 Clinical evidence profile

4 The clinical evidence profiles for this review question are presented in Table 79 to Table 85.

5 9.3.4.1 Mannitol

6 **Table 79: Summary clinical evidence profile: Comparison 1.1. Mannitol versus placebo**

Comparison 1.1. Mannitol versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Mannitol				
FEV ₁ % predicted (repeated measures, change from baseline) Range of scores: 0 to 100 Follow-up: 2 weeks	Not reported	The mean FEV ₁ % predicted (repeated measures, change from baseline) in the mannitol groups was 3.95 higher (0.96 to 6.94 higher)		36 (Jaques 2008) ¹	⊕⊕⊕⊕ low ^{2,3}	
FEV ₁ %	Not reported	The mean		600	⊕⊕⊕⊕	

Comparison 1.1. Mannitol versus placebo						
predicted (repeated measures, change from baseline) Range of scores: 0 to 100 Follow-up: 2 months		FEV ₁ % predicted (repeated measures, change from baseline) in the mannitol groups was 2.98 higher (1.04 to 4.92 higher)		(Aitken 2012, Bilton 2011)	moderate ²	
FEV ₁ % predicted (repeated measures, change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	Not reported	The mean FEV ₁ % predicted (repeated measures, change from baseline) in the mannitol groups was 3.26 higher (1.16 to 5.35 higher)		600 (Aitken 2012, Bilton 2011)	⊕⊕⊖⊖ low ^{2,3}	
FEV ₁ % predicted (repeated measures, change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	Not reported	The mean FEV ₁ % predicted (repeated measures, change from baseline) in the mannitol groups was 3.89 higher (1.69 to 6.08 higher)		600 (Aitken 2012, Bilton 2011)	⊕⊕⊖⊖ low ^{2,3}	
Time to first protocol defined pulmonary exacerbation Follow-up: 6 months	Not reported	Not reported	HR 0.7 (0.48 to 1.02)	600 (Aitken 2012, Bilton 2011)	⊕⊕⊖⊖ low ^{2,4}	
Number of patients needing additional IV antibiotics Follow-up: 6 months	Study population		RR 0.81 (0.7 to 0.95)	600 (Aitken 2012, Bilton 2011)	⊕⊖⊖⊖ very low ^{2,5,6}	
	561 per 1000	454 per 1000 (392 to 533)				
	Moderate					
	560 per 1000	454 per 1000 (392 to 532)				
Quality of life - CFQOL respiratory domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL respiratory in the placebo group was 3.8 in 1 study, 0.1 in the other	The mean change in CFQOL respiratory domain in the mannitol groups was 1.54 lower (4.69 lower to 1.61 higher)		507 (Aitken 2012, Bilton 2011)	⊕⊖⊖⊖ very low ^{2,3,7}	

Comparison 1.1. Mannitol versus placebo					
	study				
Quality of life – CFQOL respiratory domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL respiratory in the placebo group was 5.6 in 1 study, -2.5 in the other study	The mean change in CFQOL respiratory in the mannitol groups was 0.99 lower (4.5 lower to 2.52 higher)		465 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ very low ^{2,8,9}
Quality of life - CFQOL vitality domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL vitality in the placebo group was -5.4 in 1 study, -3.5 in the other study	The mean change in CFQOL vitality in the mannitol groups was 3.42 higher (0.21 lower to 7.04 higher)		361 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ low ^{2,3}
Quality of life - CFQOL vitality domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL change in CFQOL vitality in the placebo group was -4.2 in 1 study, -5.1 in the other study	The mean change in CFQOL vitality in the mannitol groups was 4.84 higher (0.86 to 8.82 higher)		325 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ low ^{2,3}
Quality of life - CFQOL physical domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL physical in the placebo group was 2.3 in 1 study, -1.5 in the other study	The mean change in CFQOL physical in the mannitol groups was 1.8 lower (4.72 lower to 1.11 higher)		505 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ moderate ²
Quality of life - CFQOL physical domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL physical in the placebo group was 1.1 in 1 study, -4.7 in the other study	The mean change in CFQOL physical in the mannitol groups was 0.52 higher (2.75 lower to 3.79 higher)		465 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9,10}
Quality of life - CFQOL emotion domain (change	The mean change in CFQOL	The mean change in CFQOL		506 (Aitken 2012,	⊕⊕⊕⊕ moderate ²

Comparison 1.1. Mannitol versus placebo						
from baseline) Range of scores: 0 to 100 Follow-up: 4 months	emotion in the placebo group was 2.9 in 1 study, -0.1 in the other study	emotion in the mannitol groups was 2.11 lower (4.56 lower to 0.34 higher)		Bilton 2011)		
Quality of life - CFQOL emotion domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL emotion in the placebo group was 2.1 in 1 study, 0.5 in the other study	The mean change in CFQOL emotion in the mannitol groups was 1.27 lower (3.74 lower to 1.2 higher)		465 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate 2	
Quality of life - CFQOL eating domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL eating in the placebo group was -3.3 in 1 study, 0.6 in the other study	The mean change in CFQOL eating in the mannitol groups was 0.81 higher (1.96 lower to 3.58 higher)		505 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate 2	
Quality of life - CFQOL eating domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL eating in the placebo group was -1.4 in 1 study, 1.9 in the other study	The mean change in CFQOL eating in the mannitol groups was 0.68 higher (2.29 lower to 3.65 higher)		466 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate 2	
Quality of life - CFQOL health domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL health in the placebo group was -1 in 1 study, 2.3 in the other study	The mean change in CFQOL health in the mannitol groups was 0.43 lower (4.18 lower to 3.32 higher)		360 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate 2	
Quality of life - CFQOL health domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL health in the placebo group was -0.9 in 1 study, 1.1 in the other study	The mean change in CFQOL health in the mannitol groups was 0.21 lower (4.14 lower to 3.72 higher)		325 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate 2	

Comparison 1.1. Mannitol versus placebo						
Quality of life - CFQOL social domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL social in the placebo group was 0.2 in 1 study, -0.8 in the other study	The mean change in CFQOL social (change from baseline) in the mannitol groups was 1.2 lower (3.7 lower to 1.3 higher)		504 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate ²	
Quality of life - CFQOL social domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL social in the placebo group was 0.9 in 1 study, -0.7 in the other study	The mean change in CFQOL social in the mannitol groups was 1.47 lower (4.25 lower to 1.32 higher)		465 (Aitken 2012, Bilton 2011)	⊕⊖⊖⊖ very low ^{2,3,11}	
Quality of life - CFQOL body domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL body in the placebo group was 1.5 in 1 study, 1.6 in the other study	The mean change in CFQOL body in the mannitol groups was 3.1 lower (6.49 lower to 0.29 higher)		500 (Aitken 2012, Bilton 2011)	⊕⊕⊖⊖ low ^{2,3}	
Quality of life - CFQOL body domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL body in the placebo group was 2.9 in 1 study, 1.8 in the other study	The mean change in CFQOL body in the mannitol groups was 1.19 lower (4.51 lower to 2.13 higher)		461 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate ²	
Quality of life - CFQOL role domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL role in the placebo group was -0.8 in 1 study, -2.4 in the other study	The mean change in CFQOL role in the mannitol groups was 1.22 higher (2.21 lower to 4.66 higher)		358 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊖ moderate ²	
Quality of life - CFQOL role domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL role in the placebo group was 1.1 in 1 study, -1.6	The mean change in CFQOL role in the mannitol groups was 1.43 lower (4.87 lower to 2 higher)		324 (Aitken 2012, Bilton 2011)	⊕⊖⊖⊖ very low ^{2,3,12}	

Comparison 1.1. Mannitol versus placebo						
	in the other study					
Quality of life - CFQOL digestion domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL digestion in the placebo group was 2.1 in 1 study, 0.2 in the other study	The mean change in CFQOL digestion in the mannitol groups was 1.49 lower (4.77 lower to 1.78 higher)		505 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ moderate ²	
Quality of life - CFQOL digestion domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL digestion in the placebo group was 2.8 in 1 study, 0 in the other study	The mean change in CFQOL digestion in the mannitol groups was 1.07 lower (5.04 lower to 2.9 higher)		465 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ low ^{2,3}	
Quality of life - CFQOL weight domain (change from baseline) Range of scores: 0 to 100 Follow-up: 4 months	The mean change in CFQOL weight in the placebo group was 4.6 in 1 study, 7.3 in the other study	The mean change in CFQOL weight in the mannitol groups was 4.23 lower (10.28 lower to 1.83 higher)		360 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ low ^{2,3}	
Quality of life - CFQOL weight domain (change from baseline) Range of scores: 0 to 100 Follow-up: 6 months	The mean change in CFQOL weight in the placebo group was 7.8 in 1 study, 6.5 in the other study	The mean change in CFQOL weight in the mannitol groups was 3.27 lower (9.84 lower to 3.31 higher)		325 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ low ^{2,3}	
Adverse events: haemoptysis (mild) Follow-up: 2 weeks	0%	0%	Not estimable	36 (Jaques 2008) ¹	⊕⊕⊕⊕ moderate ²	
Adverse events: haemoptysis (severe) Follow-up: 2 weeks	53 per 1000	53 per 1000 (8 to 355)	RR 1 (0.15 to 6.74)	(Jaques 2008) ¹	⊕⊕⊕⊕ very low ^{2,9}	
Adverse events: Bronchospasm (mild) Follow-up: 6 months	0 events in each group	0 events in each group	Not estimable	295 (Bilton 2011)	⊕⊕⊕⊕ moderate ²	

Comparison 1.1. Mannitol versus placebo					
Adverse events: Haemoptysis (mild) Follow-up: 6 months	Study population		RR 1.73 (0.26 to 11.62)	600 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9}
	8 per 1000	14 per 1000 (2 to 97)			
	Moderate				
	9 per 1000	16 per 1000 (2 to 105)			
Adverse events: Bronchospasm (moderate) Follow-up: 6 months	0 per 1000	0 per 1000 (0 to 0)	RR 2.01 (0.03 to 133.11)	295 (Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9}
Adverse events: Haemoptysis (moderate) Follow-up: 6 months	Study population		RR 4.66 (0.5 to 43.49)	600 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9}
	4 per 1000	19 per 1000 (2 to 182)			
	Moderate				
	4 per 1000	19 per 1000 (2 to 174)			
Adverse events: Bronchospasm (severe) Follow-up: 6 months	0 per 1000	0 per 1000 (0 to 0)	RR 2.01 (0.03 to 133.11)	295 (Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9}
Adverse events: Haemoptysis (severe) Follow-up: 6 months	Study population		RR 1.55 (0.13 to 18.99)	600 (Aitken 2012, Bilton 2011)	⊕⊕⊕⊕ very low ^{2,9}
	4 per 1000	6 per 1000 (1 to 79)			
	Moderate				
	4 per 1000	6 per 1000 (1 to 76)			
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: CFQOL: cystic fibrosis quality of life questionnaire; CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; HR: hazard ratio; MD: mean difference; RR: risk ratio					

- 1 1 Cross-over design
- 2 2 The quality of the evidence was downgraded by 1 as the participants in the trial underwent a tolerance test at screening. Those who failed were not entered in the study, and this limits the generalisability of the results to the general CF population.
- 3
- 4
- 5 3 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 6 4 The quality of the evidence was downgraded by 1, as the 95% CI crossed the null effect
- 7 5 The quality of the evidence was downgraded by 1 due to moderate heterogeneity (I²=59%)
- 8 6 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 9 7 The quality of the evidence was downgraded by 1 due to moderate heterogeneity (I²=37%).
- 10 8 The quality of the evidence was downgraded by 2 due to high heterogeneity (I²=89%)
- 11 9 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs
- 12 10 The quality of the evidence was downgraded by 1 due to high heterogeneity (I²=77%). It was not downgraded further as both studies showed no differences between groups.
- 13 11 The quality of the evidence was downgraded by 2 due to high heterogeneity (I²=70%). Studies show conflicting results.
- 14
- 15
- 16 12 The quality of the evidence was downgraded by 1 due to moderate heterogeneity (I²=41%)

Table 80: Summary clinical evidence profile: Comparison 1.2.1. Mannitol versus dornase alfa

Comparison 1.2.1. Mannitol versus dornase alfa					
Outcomes	Illustrative comparative risks* (95% CI)	Relative effect	No of Participant	Quality of the	Comments

Comparison 1.2.1. Mannitol versus dornase alfa						
	Assumed risk	Corresponding risk	(95% CI)	Studies (studies)	evidence (GRADE)	
	Dornase alfa	Mannitol				
FEV ₁ (% change from baseline) - Range of scores: 0 to 100 Follow-up: 3 months	Not reported	The mean FEV ₁ (% change from baseline) in the intervention mannitol was 2.8 higher (4.8 lower to 10.4 higher)		20 (Minasian 2010) ¹	⊕⊕⊕⊕ very low ^{2,3,4}	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference</p>						

- 1 *1 Cross-over design*
2 *2 The quality of the evidence was downgraded by 1 because this is an open trial, and there is high risk of*
3 *incomplete reporting*
4 *3 The quality of the evidence was downgraded by 1 as the participants in the trial underwent a tolerance test at*
5 *screening. Those who fail were not entered in the study, and this limits the generalisability of the results to the*
6 *general CF population*
7 *4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs*

8 **Table 81: Summary clinical evidence profile: Comparison 1.2.2. Mannitol plus dornase**
9 **alfa versus dornase alfa**

Comparison 1.2.2. Mannitol plus dornase alfa versus dornase alfa						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Dornase alfa alone	Mannitol + Dornase alfa				
FEV ₁ (% change from baseline) - Range of scores: 0 to 100 Follow-up: 3 months	Not reported	The mean FEV ₁ (% change from baseline) in the mannitol + dornase alfa groups was 4.3 lower (14.1 lower to 5.5 higher)		20 (Minasian 2010) ¹	⊕⊕⊕⊕ very low ^{2,3,4}	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference</p>						

- 10 *1 Cross-over design*
11 *2 The quality of the evidence was downgraded by 1 because this is an open trial, and there is high risk of*
12 *incomplete reporting*
13 *3 The quality of the evidence was downgraded by 1 as the participants in the trial underwent a tolerance test at*
14 *screening. Those who fail were not entered in the study, and this limits the generalisability of the results to the*
15 *general CF population*
16 *4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs*

1 9.3.4.2 Dornase alfa

2 Table 82: Summary clinical evidence profile: Comparison 2.1. Dornase alfa versus
3 placebo

Comparison 2.1. Dornase alfa versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Dornase alfa				
Lung function: relative mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 10 days	The relative mean % change in FEV ₁ in the placebo group was 0.15	The relative mean % change in FEV ₁ the dornase alfa group was 13.17 higher (0.70 to 25.64 higher)		41 (Shah 1996)	⊕⊕⊕⊕ very low ^{1,7}	
Lung function: relative mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 1 months	The relative mean % change in FEV ₁ in the placebo group was -1.9 in the first study, -1.6 in the second study, -1.5 in the third study, 4.2 in the fourth study	The relative mean % change in FEV ₁ in the dornase alfa groups was 9.52 higher (0.59 to 18.46 higher)		248 (Laube 1996, Ramsey 1993a, Ranasinha 1993, Shah 1995)	⊕⊕⊕⊕ very low ^{3,4,7}	
Lung function: relative mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 3 months	The relative mean % change in FEV ₁ in the placebo group was 0.76 in 1 study, 2.1 in the other study	The relative mean % change in FEV ₁ in the dornase alfa groups was 6.7 higher (3.72 to 9.67 higher)		319 (Amin 2011, McCoy 1996) ⁵	⊕⊕⊕⊕ very low ^{5,6,7}	
Lung function: relative mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 6 months	The relative mean % change in FEV ₁ in the placebo group was 0	The relative % mean change in FEV ₁ in the dornase alfa groups was 5.8 higher (4.41 to 7.19 higher)		647 (Fuchs 1994)	⊕⊕⊕⊕ low ^{7,8}	
subgroup analysis based on disease severity: moderate	The relative mean % change in FEV ₁ in the placebo	The mean % change in FEV ₁ in the dornase alfa groups was 14.32 higher		183 (Laube 1996, Ramsey 1993a, Ranasinha	⊕⊕⊕⊕ low ⁹	

Comparison 2.1. Dornase alfa versus placebo						
relative mean % change in FEV ₁ – Range of scores: 0 to 100 Follow-up: 1 months	group was -1.8 in the first study, -1.6 in the second study, -1.5 in the third study	(10.81 to 17.83 higher)		1993)		
subgroup analysis based on disease severity: severe FEV ₁ relative mean % change in FEV ₁ - Range of scores: 0 to 100 Follow-up: 1 months	The relative mean % change in FEV ₁ in the placebo group was 4.2	The mean % change in FEV ₁ in the dornase alfa groups was 2.8 lower (8.76 lower to 3.16 higher)		65 (Shah 1995)	⊕⊕⊕⊕ very low ⁷	
subgroup analysis based on disease severity: acute pulmonary exacerbation mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 1 months	The mean % change in FEV ₁ in the placebo group was 19	The mean % change in FEV ₁ in the dornase alfa groups was 1 higher (13.93 lower to 15.93 higher)		80 (Wilmott 1996)	⊕⊕⊕⊕ very low ^{2,11}	
Lung function: absolute mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 2 years	The absolute mean % change in FEV ₁ in the placebo group was -3.2	The mean % change in FEV ₁ in the dornase alfa groups was 3.24 higher (1.03 to 5.45 higher)		410 (Quan 2001)	⊕⊕⊕⊕ moderate ⁷	
Number of people experiencing exacerbations Follow-up: 6 months	274 per 1000	222 per 1000 (167 to 290)	RR 0.81 (0.61 to 1.06)	647 (Fuchs 1994)	⊕⊕⊕⊕ low ^{8,12}	
Number of people experiencing exacerbations Follow-up: 2 years	239 per 1000	170 per 1000 (117 to 244)	RR 0.71 (0.49 to 1.02)	470 (Quan 2001)	⊕⊕⊕⊕ moderate ^{1 2}	
Number of days of IV antibiotics use Follow-up: 3 months	The mean number of days of IV antibiotics use in the placebo group was	The mean number of days of IV antibiotics use in the dornase alfa groups was		320 (McCoy 1996)	⊕⊕⊕⊕ very low ^{13,14}	

Comparison 2.1. Dornase alfa versus placebo						
	28.31	2.96 lower (7.29 lower to 1.37 higher)				
Adverse events: haemoptysis Follow-up: 1 months	Study population		RR 1.23 (0.20 to 7.64)	141 (Ranasinh a 1993, Shah 1995)	⊕⊕⊕⊕ very low ^{14,15}	
	43 per 1000	53 per 1000 (9 to 327)				
	Moderate					
	43 per 1000	53 per 1000 (9 to 328)				
Adverse events: haemoptysis Follow-up: 6 months	Study population		RR 0.82 (0.44 to 1.52)	647 (Fuchs 1996)	⊕⊕⊕⊕ very low ^{8,14}	
	65 per 1000	53 per 1000 (28 to 98)				
Adverse events: voice alteration Follow-up: 1 months	Study population		RR 2.79 (0.03 to 278.07)	233 (Ramsey 1993a, Ranasinha 1993, Shah 1995)	⊕⊕⊕⊕ very low ^{14,16,17}	
	25 per 1000	71 per 1000 (1 to 1000)				
	Moderate					
	0 per 1000	0 per 1000 (0 to 0)				
Adverse events: voice alteration Follow-up: 3 months	62 per 1000	177 per 1000 (89 to 352)	RR 2.87 (1.44 to 5.71)	320 (McCoy 1996)	⊕⊕⊕⊕ moderate ¹³	
Adverse events: voice alteration Follow-up: 6 months	22 per 1000	37 per 1000 (15 to 93)	RR 1.73 (0.69 to 4.34)	647 (Fuchs 1994)	⊕⊕⊕⊕ very low ^{8,14}	
Adverse events: voice alteration Follow-up: 2 years	115 per 1000	110 per 1000 (66 to 183)	RR 0.95 (0.57 to 1.59)	470 (Quan 2001)	⊕⊕⊕⊕ low ¹⁴	
Quality of life: change in QFQ- R (QFQ-R parents) Range of scores: 0 to 100 Follow-up: 3 months	The change in QFQ-R parents in the placebo group was 0.89	The mean change in QFQ-R parents in the dornase alfa groups was 5.45 lower (15.23 lower to 4.33 higher)		17 (Amin 2011) ⁵	⊕⊕⊕⊕ moderate ⁷	
Quality of life: change in QFQ- R - QFQ-R 14+ Range of scores: 0 to 100 Follow-up: 3 months	The change in QFQ-R 14+ in the placebo group was -5.28	The mean change in QFQ-R 14+ in the dornase alfa groups was 5.21 lower (15.5 lower to 5.08 higher)		17 (Amin 2011) ⁵	⊕⊕⊕⊕ moderate ⁷	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in

Comparison 2.1. Dornase alfa versus placebo

the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFQ-R: cystic fibrosis questionnaire revised; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference; RR: risk ratio

- 1 The quality of the evidence was downgraded by due to unclear sequence generation, allocation concealment, blinding and reporting
- 2
- 3 2 The quality of the evidence was downgraded by 2 as the CI crossed 2 clinical MID
- 4 3 The quality of the evidence was downgraded by 2 due to unclear sequence generation, blinding, allocation concealment and reporting in 3 of the trials, and unclear blinding and reporting in the fourth trial
- 5 4 The quality of the evidence was downgraded by 1 due to high heterogeneity (I²=88%). See sensitivity analysis.
- 6 5 Amin 2011: cross-over trial
- 7
- 8 6 The quality of the evidence was downgraded by 1 due to unclear sequence generation, blinding, allocation concealment and reporting in the 1 of the trial
- 9
- 10 7 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 11 8 The quality of the evidence was downgraded by 1 due to unclear blinding, allocation, concealment and reporting
- 12 9 The quality of the evidence was downgraded by 2 due to unclear sequence generation, blinding, allocation concealment and reporting in 2 of the trials, and unclear blinding and reporting in the third trial
- 13 10 The quality of the evidence was downgraded by 2 due to unclear sequence generation, blinding, allocation concealment and reporting
- 14 11 The quality of the evidence was downgraded by 2 due to unclear sequence generation, blinding, allocation concealment and reporting
- 15
- 16 12 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 17 13 The quality of the evidence was downgraded by 2 due to unclear randomization, blinding, allocation concealment and reporting
- 18
- 19 14 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MID
- 20 15 The quality of the evidence was downgraded by 2 due to unclear sequence generation, blinding, allocation concealment and reporting in both trials
- 21
- 22 16 The quality of the evidence was downgraded by 2 due to unclear blinding, allocation concealment and reporting in 2 of the trials, and unclear blinding and reporting in the third trial
- 23
- 24 17 The quality of the evidence was downgraded by 1 due to high heterogeneity (I²=85%)
- 25
- 26

Table 83: Summary clinical evidence profile: Comparison 2.2. Dornase alfa versus nebulised sodium chloride

Comparison 2.2. Dornase alfa versus nebulised sodium chloride						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Nebulised sodium chloride	Dornase alfa				
Lung function: mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 3 weeks	The mean % change in FEV ₁ in the nebulised sodium chloride group was 7.7	The mean % change in FEV ₁ in the dornase alfa groups was 1.6 higher (7.96 lower to 11.16 higher)		48 (Ballmann 1998) ¹	⊕⊕⊕⊕ very low ^{2,3}	
Lung function: mean % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 3 weeks	Not reported	The mean % change in FEV ₁ in the dornase alfa groups was 8 higher (2 to 14 higher)		14 (Suri 2001) ¹	⊕⊕⊕⊕ low ^{2,4}	

Comparison 2.2. Dornase alfa versus nebulised sodium chloride					
months					
Number of days inpatient treatment Follow-up: 3 months	Not reported	The mean number of days inpatient treatment in the dornase alfa groups was 0.4 lower (2.32 lower to 1.52 higher)		14 (Suri 2001) ¹	⊕⊕⊕⊖ moderate ²
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; MD: mean difference					

- 1 1 Cross-over study
 2 2 The quality of the evidence was downgraded by 1 due to unclear blinding, allocation, concealment and reporting
 3 3 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
 4 4 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID

5 9.3.4.3 Nebulised sodium chloride

6 **Table 84: Summary clinical evidence profile: Comparison 3.1. Nebulised sodium**
 7 **chloride (> 3% concentration) versus placebo (0.9% to 0.12%) or low-**
 8 **concentration (≤ 3%)**

Comparison 3.1. Nebulised sodium chloride (> 3% concentration) versus placebo (0.9% to 0.12%) or low-concentration (≤ 3%)						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Low concentration (≤ 3% sodium chloride)	High concentration (>3% sodium chloride)				
Failed to regain pre-exacerbation FEV ₁ % predicted Range of scores: 0 to 100 Follow-up: at hospital discharge	431 per 1000	254 per 1000 (155 to 418)	RR 0.59 (0.36 to 0.97)	132 (Dentice 2016)	⊕⊕⊕⊖ moderate ¹	
Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 2 weeks	The mean % change in FEV ₁ in the ≤3% sodium chloride group was 13.81	The mean % change in FEV ₁ in the >3% sodium chloride groups was 14.35 lower (27.8 to 0.9 lower)		30 (Gupta 2012)	⊕⊕⊕⊖ moderate ¹	

Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 4 weeks	The mean % change in FEV ₁ in the ≤3% sodium chloride group was 12.53 in 1 study, -0.3 in the other study	The mean % change in FEV ₁ in the >3% sodium chloride groups was 4.92 lower (17.69 lower to 7.86 higher)		93 (Gupta 2012, Mainz 2016) ²	⊕⊕⊕ very low ^{3,4,5}	
Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 12 weeks	The mean % change in FEV ₁ in the ≤3% sodium chloride group was 3.96	The mean % change in FEV ₁ in the >3% sodium chloride groups was 4.1 higher (0.08 lower to 8.28 higher)		149 (Elkins 2006)	⊕⊕⊕⊕ moderate ¹	
Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 24 weeks	The mean: % change in FEV ₁ in the ≤3% sodium chloride group was 4.46	The mean % change in FEV ₁ in the >3% sodium chloride groups was 5.37 higher (1.03 to 9.71 higher)		140 (Elkins 2006)	⊕⊕⊕⊕ moderate ¹	
Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 36 weeks	The mean % change in FEV ₁ in the ≤3% sodium chloride group was 5	The mean % change in FEV ₁ in the >3% sodium chloride groups was 3.63 higher (1.56 lower to 8.82 higher)		134 (Elkins 2006)	⊕⊕⊕⊕ moderate ¹	
Lung function: % change in FEV ₁ Range of scores: 0 to 100 Follow-up: 48 weeks	The mean % change in FEV ₁ in the ≤3% sodium chloride group was 4.75	The mean % change in FEV ₁ in the >3% sodium chloride groups was 2.31 higher (2.72 lower to 7.34 higher)		134 (Elkins 2006)	⊕⊕⊕⊕ moderate ¹	
Time to first pulmonary exacerbation Follow-up: mean 1 years	Not reported	Not reported	HR 0.92 (0.74 to 1.14)	453 (Dentice 2016, Rosenfeld 2012)	⊕⊕⊕⊕ moderate ⁶	
Number of days of treatment for a pulmonary exacerbation Follow-up: 48 weeks	The mean number of days of treatment for a pulmonary exacerbation in the ≤3% sodium chloride group was 52	The mean number of days of treatment for a pulmonary exacerbation in the >3% sodium		321 (Rosenfeld 2012)	⊕⊕⊕⊕ high	

		chloride groups was 1.11 higher (0.89 to 1.33 higher)			
Change in quality of life following treatment - CFQOL - physical domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change CFQOL physical domain in the ≤3% sodium chloride groups was 9	The mean change in CFQOL - physical domain in the >3% sodium chloride groups was 2.00 higher (3.12 lower to 7.12 higher)		132 (Dentice 2016)	⊕⊕⊕⊕ moderate ¹
Change in quality of life following treatment - CFQOL - burden domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change CFQOL burden domain in the ≤3% sodium chloride groups was 9	The mean change in CFQOL - burden domain in the >3% sodium chloride groups was 0.00 higher (4.78 lower to 4.78 higher)		132 (Dentice 2016)	⊕⊕⊕⊕ high
Change in quality of life following treatment - CFQOL - health domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change in CFQOL health domain in the ≤3% sodium chloride groups was 14	The mean change in CFQOL - health domain in the >3% sodium chloride groups was 2.00 lower (8.15 lower to 4.15 higher)		132 (Dentice 2016)	⊕⊕⊕⊕ moderate ¹
Change in quality of life following treatment - CFQOL - respiratory domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change in CFQOL - respiratory domain in the ≤3% sodium chloride groups was 12	The mean change in CFQOL - respiratory domain in the >3% sodium chloride groups was 1.00 higher (4.99 lower to 6.99 higher)		132 (Dentice 2016)	⊕⊕⊕⊕ moderate ¹
Change in quality of life following treatment - CFQOL - physical domain Range of	The mean change in CFQOL-physical domain in the ≤3% sodium chloride groups was	The mean change in CFQOL - physical domain in the >3% sodium chloride groups was		132 (Dentice 2016)	⊕⊕⊕⊕ moderate ¹

scores: 0 to 100 Follow-up: 7 days	14	2.00 higher (4.15 lower to 8.15 higher)				
Change in quality of life following treatment - CFQOL - burden domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change CFQOL - burden domain in the ≤3% sodium chloride groups was -1	The mean change in CFQOL - burden domain in the >3% sodium chloride groups was 2.00 higher (4.04 lower to 8.04 higher)		132 (Dentice 2016)	⊕⊕⊕⊖ moderate ¹	
Change in quality of life following treatment - CFQOL - health domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change in CFQOL - health domain in the ≤3% sodium chloride groups was 18	The mean change in CFQOL - health domain in the >3% sodium chloride groups was 2.00 higher (4.99 lower to 8.99 higher)		132 (Dentice 2016)	⊕⊕⊕⊖ moderate ¹	
Change in quality of life following treatment - CFQOL - respiratory domain Range of scores: 0 to 100 Follow-up: 7 days	The mean change in CFQOL - respiratory domain in the ≤3% sodium chloride groups was 21	The mean change in CFQOL - respiratory domain in the >3% sodium chloride groups was 2.00 lower (8.67 lower to 4.67 higher)		132 (Dentice 2016)	⊕⊕⊕⊖ moderate ¹	
Quality of life: CFQ-R respiratory: CFQ parent Range of scores: 0 to 100 Follow-up: 4 weeks	Not reported	The mean CFQ-R respiratory, CFQ parent in the >3% sodium chloride group was 5.9 higher (3.1 lower to 14.9 higher)		20 (Amin 2010) ⁷	⊕⊕⊕⊖ moderate ¹	
Quality of life: CFQ-R respiratory 14+ Range of scores: 0 to 100 Follow-up: 4 weeks	Not reported	The mean CFQ-R respiratory, CFQ 14+ in the >3% sodium chloride groups was 5.2 higher (7 lower to 17.4 higher)		20 (Amin 2010) ⁷	⊕⊕⊕⊖ low ⁵	

Change in quality of life: CFQ - CFQ-R parents Range of scores: 0 to 100 Follow-up: 48 weeks	The mean change in CFQ-R parents in the ≤3% sodium chloride group was 0.9	The mean change in CFQ-R parents in the >3% sodium chloride groups was 1.13 lower (7.49 lower to 5.23 higher)		67 (Elkins 2006)	⊕⊕⊕⊕ low ⁵	
Change in quality of life: CFQ-R 14+ Range of scores: 0 to 100 Follow-up: 48 weeks	The mean change in CFQ-R 14+ in the ≤3% sodium chloride group was 1.09	The mean change in CFQ-R 14+ in the >3% sodium chloride groups was 7.77 higher (1.86 to 13.68 higher)		92 (Elkins 2006)	⊕⊕⊕⊕ moderate ¹	
Change in quality of life: CFQ-R respiratory domain Range of scores: 0 to 100 Follow-up: 48 weeks	The mean change in CFQ-R respiratory domain in the ≤3% sodium chloride group was -3.2	The mean change in CFQ-R respiratory domain in the >3% sodium chloride groups was 3.3 higher (0 to 6.6 higher)		321 (Rosenfeld 2012)	⊕⊕⊕⊕ moderate ¹	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFQ-R: cystic fibrosis questionnaire revised; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; HR: hazard ratio, MD: mean difference; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 2 Mainz 2016: Cross-over study
- 3 The quality of the study was downgraded by 1 due to unclear risk of bias in relation to random sequence generation, allocation concealment and selective reporting in 1 study
- 4 The quality of the evidence was downgraded by 2 due to serious inconsistency (I²=77%)
- 5 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
- 6 The quality of the evidence was downgraded by 1 as the 95% CI crossed the null effect
- 7 Amin 2010: cross-over study

9 9.3.4.4 Acetylcysteine

10 **Table 85: Summary clinical evidence profile: Comparison 4. Acetylcysteine versus**
11 **placebo**

Comparison 4. Acetylcysteine versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Acetylcysteine				
Lung function: change in FEV ₁	The mean change in FEV ₁ in	The mean change in FEV ₁		21 (Skov)	⊕⊕⊕⊕ very low ^{1,2}	

Comparison 4. Acetylcysteine versus placebo						
% predicted Range of scores: 0 to 100 Follow-up: 4 weeks	the placebo group was -1.41	% predicted in the acetylcysteine groups was 3.51 higher (0.65 lower to 7.67 higher)		2015)		
Lung function: change in FEV ₁ % predicted Range of scores: 0 to 100 Follow-up: 12 weeks	The mean change in FEV ₁ in the placebo group was -8.6	The mean change in FEV ₁ % predicted in the acetylcysteine groups was 5 higher (10.84 lower to 20.84 higher)		21 (Ratjen 1985)	⊕⊕⊕⊕ low ³	
Lung function: change in FEV ₁ % predicted Range of scores: 0 to 100 Follow-up: 24 weeks	Not reported	The mean change in FEV ₁ % predicted in the acetylcysteine groups was 4.4 higher (0.83 to 7.97 higher)		70 (Conrad 2015)	⊕⊕⊕⊕ moderate ²	
Inflammatory markers: change in sputum IL-8 (log ₁₀) Follow-up: 24 weeks	Not reported	The mean change in sputum IL-8 (log ₁₀) in the acetylcysteine groups was MD 0.19 higher (0.03 lower to 0.42 higher)		70 (Conrad 2015)	⊕⊕⊕⊕ high ⁴	
Incidence of pulmonary exacerbations Follow-up: 24 weeks	500 per 1000	415 per 1000 (250 to 695)	RR 0.83 (0.5 to 1.39)	70 (Conrad 2015)	⊕⊕⊕⊕ low ³	
Quality of life: QFQ-R respiratory Follow-up: 24 weeks	Not reported	The mean QFQ-R respiratory in the acetylcysteine groups was 0.34 lower (6.3 lower to 5.62 higher)		70 (Conrad 2015)	⊕⊕⊕⊕ low ³	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFQ-R: cystic fibrosis questionnaire revised; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IL-8: interleukin 8; MD: mean difference; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1 as this is an open trial, and there was unclear randomization and allocation concealment.
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
- 4 Imprecision not calculable, as SD for the control group was not available in the study

1 **9.3.5 Economic evidence**

2 Six economic evaluations of mucoactive or mucolytic agents to manage cystic fibrosis were
3 identified in the literature search conducted for this guideline. Five of those 6 studies included
4 dornase alfa as an intervention compared with either no dornase alfa or hypertonic sodium
5 chloride, the remaining economic evaluation assessed mannitol (with and without dornase
6 alfa) against best supportive care (control). No economic evaluations were identified that
7 included acetylcysteine. A description of the methods and results of those economics
8 evaluations can be found in Appendix K.

9 Full details of the search and economic article selection flow chart can be found in Appendix
10 E and F, respectively. Data extraction tables and quality assessments of included studies
11 can be found in Appendix L and M, respectively.

12 Based on the available evidence, the Committee agreed additional economic analysis would
13 be superfluous. Instead, a cost description of the interventions was undertaken in Appendix
14 K.

15 **9.3.6 Evidence statements**

16 **9.3.6.1 Mannitol**

17 **9.3.6.1.1 Comparison 1.1 Mannitol versus placebo**

18 **Lung function: FEV₁**

19 Low quality evidence from 1 cross-over trial with 18 children and young people with cystic
20 fibrosis up to 18 years showed no clinically significant difference in the lung function
21 (measured as change in FEV₁ % predicted) between the group of participants receiving
22 mannitol (420 mg twice daily) and those in the control group (non-respirable mannitol <2%)
23 at 2 weeks follow-up.

24 Moderate quality evidence from 2 RCTs with 600 people with cystic fibrosis ≥6 years
25 showed no clinically significant difference in the lung function (measured as change in FEV₁
26 % predicted) between the group of participants receiving mannitol (400 mg twice daily) and
27 those in the control group (mannitol 50 mg twice daily) at 2 months follow-up.

28 Low quality evidence from 2 RCTs with 600 people with cystic fibrosis ≥6 years showed no
29 clinically significant difference in the lung function (measured as change in FEV₁ %
30 predicted) between the group of participants receiving mannitol (400 mg twice daily) and
31 those in the control group (mannitol 50 mg twice daily) at 4 and 6 months follow-up.

32 **Time to next exacerbation**

33 Low quality evidence from 2 RCTs with 600 people with cystic fibrosis ≥6 years showed no
34 clinically significant difference in the time to first pulmonary exacerbation (protocol defined
35 exacerbation) between the participants receiving mannitol (400 mg twice daily) and those in
36 the control group (mannitol 50 mg twice daily) at 6 months follow-up.

37 **Need for additional intravenous antibiotics for pulmonary exacerbation**

38 Very low quality evidence from 2 RCTs with 600 people with cystic fibrosis ≥6 years showed
39 no clinically significant difference in the number of people requiring intravenous antibiotics
40 between the group of participants receiving mannitol (400 mg twice daily) and those in the
41 control group (mannitol 50 mg twice daily) at 6 months follow-up. Moderate heterogeneity
42 was observed between both trials. One trial (n=305) showed no clinically significant

1 differences, whereas the other trial (n=295) showed that there was a clinically significant
2 lower number of people who needed additional intravenous antibiotics in the mannitol group.

3 **Inflammatory markers**

4 No evidence was found for this outcome.

5 **Quality of life**

6 Very low to moderate quality evidence from 2 RCTs with 600 people with cystic fibrosis ≥ 6
7 years showed no clinically significant difference in the quality of life (measured with CF-QOL
8 respiratory, vitality, physical, emotion, eating, health, social, body, role, digestion and weight
9 domains) between the participants receiving mannitol (400 mg twice daily) and those in the
10 control group (mannitol 50 mg twice daily) at 4 and 6 months follow-up.

11 Moderate to high heterogeneity was observed for the respiratory domain at 4 and 6 months,
12 and for the physical, social and role domains at 6 months. In spite of the heterogeneity, both
13 trials showed that the differences between groups were not clinically significant in either trial
14 for the respiratory domain at 4 months, and for the physical and social domains at 6 months.

15 However, for the respiratory domain, 1 trial (n=278) showed a clinically significant
16 improvement in the control group compared to the mannitol group, whereas the other trial
17 (n=229) showed no clinically significant differences at 6 months follow-up.

18 **Adverse events**

19 Moderate quality evidence from 1 cross-over trial with 18 children and young people with
20 cystic fibrosis up to 18 years reported that none of the participants in the intervention group
21 (Mannitol 420 mg twice daily) or the control group (non-respirable mannitol <2%)
22 experienced mild haemoptysis at 2 weeks follow-up.

23 Very low quality evidence from the same trial with 18 children and young people with cystic
24 fibrosis up to 18 years showed no clinically significant difference in the occurrence of severe
25 haemoptysis between the participants receiving Mannitol (420 mg twice daily) and those in
26 the control group (non-respirable mannitol <2%) at 2 weeks follow-up.

27 Moderate quality evidence from 1 trial with 295 people with cystic fibrosis ≥ 6 years reported
28 that none of the participants in the intervention group (Mannitol 400 mg twice daily) or in the
29 control group (50 mg twice daily) experienced mild bronchospasm at 6 months follow-up.

30 Moderate quality evidence from the same 2 trials with 600 people with cystic fibrosis ≥ 6
31 years reported that 1 participant in the mannitol group experienced moderate bronchospasm,
32 and 1 participant experienced severe bronchospasm in the mannitol group. No events of
33 moderate or severe bronchospasm were observed in the control group at 6 months follow-
34 up. These differences were not clinically significant.

35 Very low quality evidence from 2 trials with 600 people with cystic fibrosis ≥ 6 years showed
36 no clinically significant differences in the occurrence of mild, moderate or severe
37 haemoptysis between the participants receiving Mannitol (420 mg twice daily) and those in
38 the control group (non-respirable mannitol <2%) at 2 weeks follow-up.

399.3.6.1.2 **Comparison 1.2.1. Mannitol versus dornase alfa**

40 **Lung function: FEV₁**

41 Very low quality evidence from 1 cross-over trial with 20 children with cystic fibrosis (mean
42 age 13.2 years) showed no clinically significant difference in the lung function (measured as

1 FEV₁ % change from baseline) between the group of participants receiving mannitol (400 mg
2 twice daily) and the participants receiving dornase alfa (2.5 mg twice daily).

3 **Time to next exacerbation**

4 No evidence was found for this outcome.

5 **Need for additional intravenous antibiotics for pulmonary exacerbation**

6 No evidence was found for this outcome.

7 **Inflammatory markers**

8 No evidence was found for this outcome.

9 **Quality of life**

10 No evidence was found for this outcome.

11 **Adverse events**

12 No evidence was found for this outcome.

139.3.6.1.3 ***Comparison 1.2.1. Mannitol plus dornase alfa versus dornase alfa***

14 **Lung function: FEV₁**

15 Very low quality evidence from 1 cross-over trial with 20 children with cystic fibrosis (mean
16 age 13.2 years) showed no clinically significant difference in the lung function (measured as
17 FEV₁ % change from baseline) between the group of participants receiving a combination of
18 mannitol (400 mg mannitol twice daily) and dornase alfa (2.5 mg twice daily) and the
19 participants receiving dornase alfa alone (2.5 mg twice daily).

20 **Time to next exacerbation**

21 No evidence was found for this outcome.

22 **Need for additional intravenous antibiotics for pulmonary exacerbation**

23 No evidence was found for this outcome.

24 **Inflammatory markers**

25 No evidence was found for this outcome.

26 **Quality of life**

27 No evidence was found for this outcome.

28 **Adverse events**

29 No evidence was found for this outcome.

309.3.6.1.4 ***Comparison 1.3: Mannitol versus nebulised sodium chloride***

31 No evidence was found for this comparison.

19.3.6.1.5 **Comparison 1.4. Mannitol versus acetylcysteine**

2 No evidence was found for this comparison.

3 **9.3.6.2 Dornase alfa**

49.3.6.2.1 **Comparison 2.1: Dornase alfa versus placebo**

5 **Lung function: FEV₁**

6 Very low quality evidence from 1 RCT with 41 people with cystic fibrosis aged >15 years
7 showed a clinically significant improvement in lung function (measured as relative mean %
8 change in FEV₁) in the group of participants receiving dornase alfa (2.5 mg twice daily)
9 compared to those who were receiving placebo at 10 days follow-up.

10 Very low quality evidence from 4 RCTs with 248 people with cystic fibrosis >5 years showed
11 a clinically significant improvement in lung function (measured as relative mean % change in
12 FEV₁) in the group of participants receiving dornase alfa (2.5 mg once daily or twice daily)
13 compared to those who were receiving placebo at 1 month follow-up. However, a high level
14 of heterogeneity was found. A subgroup analysis showed that that this improvement in lung
15 function was significant in people with moderate lung disease (3 RCTs, n=183, low quality),
16 whereas no differences were found in the group of participants with severe lung disease (1
17 RCT, n=65, very low quality).

18 Very low quality evidence from 1 RCT with 80 people with cystic fibrosis and acute
19 pulmonary exacerbation >5 years showed no clinically significant difference in lung function
20 (measured as % mean change in FEV₁) between the group of participants receiving dornase
21 alfa (2.5 mg) and those who were receiving placebo at 1 month follow-up.

22 Very low quality evidence from 2 RCTs with 319 people with cystic fibrosis >6 years showed
23 a clinically significant improvement in lung function (measured as relative mean % change in
24 FEV₁) in the group of participants receiving dornase alfa (2.5 mg twice daily) compared to
25 those who were receiving placebo at 3 months follow-up.

26 Low quality evidence from 1 RCT with 647 people with cystic fibrosis >6 years showed a
27 clinically significant improvement in lung function (measured as relative mean % change in
28 FEV₁) in the group of participants receiving dornase alfa (2.5 mg once daily or twice daily)
29 compared to those who were receiving placebo at 6 months follow-up.

30 Moderate quality evidence from 1 RCT with 410 children with cystic fibrosis aged 6 to 10
31 years showed no clinically significant difference in lung function (measured as absolute mean
32 % change in FEV₁) between the group of participants receiving dornase alfa (2.5 mg) and
33 those who were receiving placebo at 2 years follow-up.

34 **People experiencing exacerbations (proxy outcome for time to next exacerbation)**

35 Low quality evidence from 1 RCT with 647 people with cystic fibrosis ≥5 years showed no
36 clinically significant difference in the number of people experiencing pulmonary
37 exacerbations between the group of participants receiving dornase alfa (2.5 mg once daily or
38 twice daily) and those who were receiving placebo at 6 months follow-up.

39 Moderate quality evidence from 1 RCT with 470 children with cystic fibrosis aged 6 to 10
40 years showed no clinically significant difference in the number of children experiencing
41 pulmonary exacerbations between the group of participants receiving dornase alfa (2.5 mg)
42 and those who were receiving placebo at 6 months follow-up.

1 **Number of days of intravenous antibiotics use (proxy outcome for need for additional**
2 **intravenous antibiotics for pulmonary exacerbation)**

3 Very low quality evidence from 1 RCT with 320 people with cystic fibrosis ≥ 7 years showed
4 no clinically significant difference in the number of days of intravenous antibiotic use between
5 the group of participants receiving dornase alfa (2.5 mg once daily or twice daily) and those
6 who were receiving placebo at 3 months follow-up.

7 **Inflammatory markers**

8 No evidence was found for this outcome.

9 **Quality of life**

10 Moderate quality evidence from 1 cross-over RCT with 17 children and young people with
11 cystic fibrosis aged 6 to 18 years showed no clinically significant difference in the quality of
12 life (QFQ-R parents and QFQ-R 14+) between the group of participants receiving dornase
13 alfa (2.5 mg) and those who were receiving placebo at 3 months follow-up.

14 **Adverse events**

15 Very low quality evidence from 2 RCTs with 141 people with cystic fibrosis > 5 years showed
16 no clinically significant difference in the occurrence of haemoptysis between the group of
17 participants receiving dornase alfa (2.5 mg) and those who were receiving placebo at 1
18 month follow-up.

19 Likewise, very low quality evidence from another RCT with 647 people with cystic fibrosis ≥ 5
20 years showed no clinically significant difference in the occurrence of haemoptysis between
21 the group of participants receiving dornase alfa (2.5 mg) and those who were receiving
22 placebo at 6 months follow-up.

23 Very low quality evidence from 3 RCTs with 233 people with cystic fibrosis > 5 years showed
24 a clinically significant higher occurrence of voice alteration in the group of participants
25 receiving dornase alfa (2.5 mg) compared to those who were receiving placebo at 1 month
26 follow-up. Significant heterogeneity was noted between the 2 trials that could be pooled in
27 the meta-analysis. One trial (n=92) noted a harmful effect of dornase alfa, whereas the other
28 did not show any differences. The third trials reported no events in either group.

29 Likewise, moderate quality evidence from another RCT with 320 children with cystic fibrosis
30 aged 6 to 10 years showed a clinically significant higher occurrence of voice alteration in the
31 group of participants receiving dornase alfa (2.5 mg) compared to those who were receiving
32 placebo at 3 months follow-up.

33 However, very low quality evidence from 1 RCT with 647 people with cystic fibrosis ≥ 5 years
34 showed no clinically significant difference in the occurrence of voice alteration between the
35 group of participants receiving dornase alfa (2.5 mg once daily or twice daily) and those who
36 were receiving placebo at 6 months follow-up.

37 Similarly, low quality evidence from another RCT with 470 children with cystic fibrosis aged 6
38 to 10 years showed no clinically significant difference in the occurrence of voice alteration
39 between the group of participants receiving dornase alfa (2.5 mg once daily or twice daily)
40 and those who were receiving placebo at 2 years follow-up.

19.3.6.2.2 **Comparison 2.2. Dornase alfa versus nebulised sodium chloride - hypertonic or isotonic (NaCL HS or IS)**

3 **Lung function: FEV₁%**

4 Very low quality evidence from 1 cross-over trial with 48 children with cystic fibrosis (mean
5 age 13.3) showed no clinically significant difference in the lung function (measured as mean
6 % change in FEV₁) between the participants receiving dornase alfa (2.5 mg) and those
7 receiving 5.85% sodium chloride at 3 weeks follow-up.

8 Low quality evidence from 1 cross-over trial with 14 people with cystic fibrosis >7 years
9 showed a clinically significant improvement in the lung function (measured as mean %
10 change in FEV₁) in the participants receiving dornase alfa (2.5 mg) compared to those
11 receiving 7% sodium chloride at 3 months follow-up.

12 **Time to next exacerbation**

13 No evidence was found for this outcome.

14 **Number of days of inpatient treatment (proxy outcome for need for additional
15 intravenous antibiotics for pulmonary exacerbation)**

16 Moderate quality evidence from 1 cross-over trial with 14 people with cystic fibrosis >7 years
17 showed no clinically significant difference in the number of days of inpatient treatment
18 between the participants receiving dornase alfa (2.5 mg) and those receiving 7% sodium
19 chloride at 3 months follow-up.

20 **Inflammatory markers**

21 No evidence was found for this outcome.

22 **Quality of life**

23 No evidence was found for this outcome.

24 **Adverse events**

25 No evidence was found for this outcome.

269.3.6.2.3 **Comparison 2.3. Dornase alfa versus acetylcysteine**

27 No evidence was found for this comparison.

28 **9.3.6.3 Nebulised sodium chloride: hypertonic or isotonic**

299.3.6.3.1 **Comparison 3.1. Nebulised sodium chloride (> 3% concentration) versus placebo
30 (0.9%) or low-concentration (≤ 3%)**

31 **Lung function: FEV₁ %**

32 Moderate quality evidence from 1 RCT with 132 adults with cystic fibrosis showed a clinically
33 significant higher likelihood of regaining pre-exacerbation FEV₁% predicted in the group of
34 participants receiving 7% sodium chloride compared to those who were receiving 3% sodium
35 chloride at hospital discharge.

36 Moderate quality evidence from 1 RCT with 30 children and young people with cystic fibrosis
37 aged 6 to 16 years showed a clinically significant decrease in the lung function (measured as

1 % change in FEV₁) between the group of participants receiving 7% sodium chloride
2 compared to those who were receiving 3% sodium chloride at 2 weeks follow-up.

3 Very low quality evidence from 2 RCTs with 93 people with cystic fibrosis aged ≥6 years
4 showed no clinically significant difference in lung function (measured as % change in FEV₁)
5 between the group of participants receiving 6 to 7% sodium chloride compared to those who
6 were receiving 3% sodium chloride at 4 weeks follow-up. Significant heterogeneity was
7 observed between both trials. The larger trial (n=123) showed no clinically significant
8 difference between both groups, whereas the smallest trial (n=30) showed a clinical
9 significant difference in favour of low-dose concentration.

10 Moderate quality evidence from 1 RCT with 148 people with cystic fibrosis ≥6 years showed
11 no clinically significant difference in the lung function (measured as % change in FEV₁)
12 between the group of participants receiving 7% sodium chloride and those who were
13 receiving 0.9% sodium chloride at 12, 36 and 48 weeks follow-up.

14 Moderate quality evidence from 1 RCT with 140 people with cystic fibrosis ≥5 years showed
15 a clinically significant improvement in the lung function (measured as % change in FEV₁) in
16 the group of participants receiving 7% sodium chloride compared to those who were
17 receiving 0.9% sodium chloride at 24 weeks follow-up.

18 **Time to next exacerbation**

19 Moderate quality evidence from 2 RCTs with 453 infants, children, young people and adults
20 with cystic fibrosis showed no clinically significant difference in the time to first pulmonary
21 exacerbation between the group of participants receiving 7% sodium chloride and those who
22 were receiving <3% sodium chloride at 48 weeks follow-up.

23 **Need for additional intravenous antibiotics for pulmonary exacerbation**

24 High quality evidence from 1 RCT with 321 children with cystic fibrosis ≤6 years showed a
25 clinically significant increase in the number of days of treatment 7% sodium chloride -dose
26 NaCl (7% HS) compared to those who were receiving 0.9% sodium chloride at 48 weeks
27 follow-up.

28 **Inflammatory markers**

29 No evidence was found for this outcome.

30 **Quality of life**

31 Moderate to high quality evidence from 1 RCT with 132 adults with cystic fibrosis showed no
32 clinically significant difference in the quality of life (measured with CF-QOL physical, burden,
33 health and respiratory domains) between the group of participants receiving 7% sodium
34 chloride and those who were receiving 3% sodium chloride at hospital discharge and at 7
35 weeks follow-up.

36 Low to moderate quality evidence from 1 cross-over trial with 20 children and young people
37 with cystic fibrosis aged 6 to 18 years showed no clinically significant difference in the quality
38 of life (measured with CFQ-R 14+ or CFQ-R parent respiratory domain) between the group of
39 participants receiving 7% sodium chloride and those who were receiving 0.9% sodium
40 chloride at 4 weeks follow-up.

41 High quality evidence from 1 RCT with 67 people with cystic fibrosis ≥6 years showed no
42 clinically significant difference in the quality of life (measured as change in CFQ-R parents)
43 between the group of participants receiving 7% sodium chloride and those who were
44 receiving 0.9% sodium chloride at 48 weeks follow-up.

1 Moderate quality evidence from 1 RCT with 92 people with cystic fibrosis ≥ 6 years showed a
2 clinically significant beneficial effect in the quality of life (measured as change in CFQ-R 14+)
3 in the group of participants receiving 7% sodium chloride compared to those who were
4 receiving 0.9% sodium chloride at 48 weeks follow-up.

5 Moderate quality evidence from 1 RCT with 321 children with cystic fibrosis ≤ 6 years showed
6 no clinically difference in the quality of life (measured as change in CFQ-R respiratory)
7 between the group of participants receiving 7% sodium chloride compared and those who
8 were receiving 0.9% sodium chloride at 48 weeks follow-up.

9 **Adverse events**

10 No evidence was found for this outcome.

119.3.6.3.2 **Comparison 3.2. Nebulised sodium chloride versus acetylcysteine**

12 No evidence was found for this comparison.

13 **9.3.6.4 Acetylcysteine**

149.3.6.4.1 **Comparison 4. Acetylcysteine versus placebo**

15 **Lung function: FEV₁**

16 Very low quality evidence from 1 RCT with 21 adults with cystic fibrosis showed no clinically
17 significant difference in the lung function (measured as change in FEV₁ % predicted)
18 between the participants receiving acetylcysteine (2400 mg per day) and those receiving
19 placebo at 4 weeks follow-up

20 Low quality evidence from 1 RCT with 22 children and young people with cystic fibrosis aged
21 6 to 21 years showed no clinically significant difference in the lung function (measured as
22 change in FEV₁ % predicted) between the participants receiving acetylcysteine (200 mg x 3
23 times per day) and those receiving placebo at 12 weeks follow-up.

24 Similarly, moderate quality evidence from 1 RCT with 70 people with cystic fibrosis >9 years
25 showed no clinically significant difference in the lung function (measured as change in FEV₁
26 % predicted) between the participants receiving acetylcysteine (900mg/ 3 times per day) and
27 those receiving placebo at 24 weeks follow-up.

28 **Time to next exacerbation**

29 No evidence was found for this outcome.

30 **Incidence of exacerbations (proxy outcome for need for additional intravenous 31 antibiotics for pulmonary exacerbation)**

32 Low quality evidence from 1 RCT with 70 people with cystic fibrosis >9 years showed no
33 clinically significant difference in the incidence of pulmonary exacerbations between the
34 participants receiving acetylcysteine (900mg/ 3 times per day) and those receiving placebo at
35 24 weeks follow-up.

36 **Inflammatory markers**

37 High quality evidence from 1 RCT with 70 people with cystic fibrosis >9 years showed no
38 significant difference in the inflammatory markers (measured as sputum IL-8 log₁₀) between
39 the participants receiving acetylcysteine (900mg/ 3 times per day) and those receiving
40 placebo at 24 weeks follow-up. The uncertainty for this outcome could not be calculated.

1 **Quality of life**

2 Low quality evidence from 1 RCT with 70 people with cystic fibrosis >9 years showed no
3 clinically significant difference in the quality of life (measured with QFQ-R respiratory domain)
4 between the participants receiving acetylcysteine (900mg/ 3 times per day) and those
5 receiving placebo at 24 weeks follow-up.

6 **Adverse events**

7 No evidence was found for this outcome.

8 **9.3.6.5 Economic evidence statements**

9 One cost-benefit analysis (Menzin 1996) undertaken in the UK, on people with cystic fibrosis,
10 found that daily dornase alfa may reduce the cost of respiratory tract infection related care
11 compared to placebo, over 24 weeks. This analysis is partially applicable as clinical
12 effectiveness data was taken from an old US trial that may reflect outdated practices and
13 practices that may not be generalisable to the UK. The evidence was also associated with
14 very serious limitations including the omission of dornase alfa in their costs.

15 One cost-effectiveness analysis (Christopher 1999) undertaken in the UK, on people with
16 cystic fibrosis, using a lifetime horizon, found that the cost per life year gained for daily
17 dornase alfa compared to placebo was £52,550 for all participants and £16,110 for the
18 subgroup of participants with FEV1≤70%. This analysis is partially applicable with very
19 serious limitations, namely as clinical effectiveness data was taken from an old US trial that
20 may reflect outdated practices and practices that may not be generalisable to the UK.

21 One cost-benefit analysis (McIntyre 1996) undertaken in the UK, on people with cystic
22 fibrosis, using a lifetime horizon, found that the cost per life year gained for daily dornase alfa
23 compared to placebo could range from £10,311 to £45,234. This analysis is partially
24 applicable with very serious limitations, namely as clinical effectiveness data was taken from
25 an old US trial that may reflect outdated practices and practices that may not be
26 generalisable to the UK.

27 One cost-benefit analysis (Suri 2002) on people with cystic fibrosis in the UK, over 12 weeks,
28 found that daily dornase alfa was more effective than hypertonic saline, but significantly
29 increased health care costs. Administering dornase alfa on alternate days, rather than daily,
30 was as effective, with a potential for cost savings. This analysis is partially applicable with
31 minor limitations.

32 One cost-effectiveness analysis (Grieve 2003) on people with cystic fibrosis in the UK found
33 that the cost per 1% gain in FEV1% over 12 weeks, for daily dornase alpha compared to
34 hypertonic saline was £110; for daily dornase alpha compared to alternate day dornase alpha
35 £214 and for alternate day dornase alpha compared to hypertonic saline £89. This analysis
36 has minor limitations and is directly applicable given that the type of economic evaluation is
37 unlikely to change the conclusions about cost-effectiveness and all other applicability criteria
38 are met.

39 One cost-utility analysis (NICE TA266) on people with cystic fibrosis in the UK using a
40 lifetime horizon, found that the ICER for mannitol compared to best supportive care was
41 £41,074 and for mannitol plus dornase alfa compared to best supportive care plus dornase
42 alfa £47,095. Amendments to the original analysis found that mannitol plus dornase alfa
43 compared dornase alfa plus best supportive care for people with cystic fibrosis using dornase
44 alfa (i.e. mannitol as add-on therapy) produced an ICER of £80,098. For mannitol compared
45 to best supportive care for people with cystic fibrosis who are ineligible, intolerant or
46 inadequately responsive to rhDNase (i.e. mannitol as second-line therapy) the ICER was
47 £29,883. This analysis is directly applicable. The initial analysis is associated with serious
48 limitations as both comparisons use clinical effectiveness data taken from the whole adult

1 population, irrespective of dornase alfa use which underestimates the effectiveness of
2 dornase alfa use.

3 **9.3.7 Evidence to recommendations**

4 **9.3.7.1 Relative value placed on the outcomes considered**

5 The aim of this review was to establish the clinical and cost effectiveness of mucoactive or
6 mucolytic agents in improving airway clearance in children, young people and adults with
7 cystic fibrosis.

8 The Committee identified lung function (FEV₁% predicted), time to pulmonary exacerbation
9 and need for intravenous antibiotics for pulmonary exacerbation as critical outcomes for this
10 evidence review. Where no evidence was found for time to pulmonary exacerbation, the
11 number of people experiencing a pulmonary exacerbation was taken as a proxy outcome.
12 Inflammatory markers, quality of life and adverse events were rated as important outcomes.

13 **9.3.7.2 Consideration of clinical benefits and harms**

14 The Committee discussed whether a mucoactive or mucolytic agent should be prescribed to
15 everyone who has cystic fibrosis. However, taking into account the potential adverse effects,
16 as well as the inconvenience and the cost of treatment, it was agreed not to recommend it to
17 everyone; instead, the Committee agreed that it should be offered to people with cystic
18 fibrosis who have respiratory symptoms or other evidence of lung disease based on
19 radiological imaging or lung function testing.

20 The Committee reviewed the evidence comparing dornase alfa to placebo, which shows
21 significant differences in FEV₁ in favour of dornase alfa at 1, 3, 6 and 24 months follow-up,
22 but also a lack of significant differences in FEV₁ in people with severe lung disease at 1
23 month follow-up. The Committee discussed the recommendations from NICE TA266 that
24 provides guidance on the use of mannitol dry powder for inhalation for the treatment of cystic
25 fibrosis in adults. They agreed that mannitol could be recommended as an option in adults
26 who cannot use dornase alfa because of ineligibility, intolerance or inadequate response to
27 dornase alfa, and in those whose lung function is rapidly declining (forced expiratory volume
28 in 1 second [FEV₁] decline greater than 2% annually) for whom other osmotic agents are not
29 considered appropriate. They also agreed that people currently receiving mannitol whose
30 cystic fibrosis does not meet the cited criteria should be able to continue treatment until they
31 and their clinician consider it appropriate to stop. Therefore the Committee adopted these
32 recommendations from NICE TA266.

33 The Committee discussed the evidence comparing nebulised sodium chloride with control
34 (0.9%) or low-concentration (< 3%). After reviewing the conflicting evidence comparing 7%
35 sodium chloride to 0.9% sodium chloride, the Committee relied on their expertise and
36 experience to recommend hypertonic sodium chloride instead of isotonic sodium chloride.
37 The Committee also reviewed the evidence comparing 7% sodium chloride to 3% sodium
38 chloride. A moderate quality RCT found a clinically significant improvement in FEV₁ in the
39 group of participants receiving 7% sodium chloride compared to those who were receiving
40 3% sodium chloride at 2 and 4 weeks follow-up. It was discussed whether a specific
41 concentration of hypertonic sodium chloride should be specified in the recommendations.
42 The Committee concluded that it was appropriate not to mention a specific concentration
43 because the highest concentration tolerable for the individual patient should be used (to
44 maximum 7%).

45 The Committee reviewed the evidence comparing acetylcysteine to placebo. Very low to
46 moderate quality evidence showed no clinically significant differences in FEV₁ between
47 acetylcysteine and placebo at 4, 12 and 24 weeks follow-up. Likewise, low quality evidence
48 showed no differences in need for additional intravenous antibiotics for pulmonary

1 exacerbation at 24 weeks follow-up. No clinically significant differences were found in
2 inflammatory markers, or quality of life either. The Committee also noted that acetylcysteine
3 was not commonly used in clinical practice because of the very unpleasant smell and taste.
4 Moreover, acetylcysteine needs to be taken up to four times a day, so overall it is much less
5 tolerable and more burdensome than other mucoactive agents. Based on this, the
6 Committee agreed not to make a recommendation in favour of acetylcysteine.

7 The Committee reviewed the evidence comparing nebulised dornase alfa to hypertonic
8 sodium chloride, which showed significant differences in FEV₁ in favour of dornase alfa at 3
9 months follow-up but not at 3 weeks. The evidence was low or very low quality. Due to the
10 limited evidence, the Committee also relied on their expertise and experience to guide their
11 decision as to whether dornase alfa or hypertonic sodium chloride should be the first-line
12 treatment. On balance, they agreed that dornase alfa was more effective and tolerable, and
13 insufficient evidence was presented to change currently accepted practice. Therefore, the
14 Committee recommended dornase alfa as first choice treatment and hypertonic sodium
15 chloride as second choice treatment.

16 The Committee recommended to use hypertonic sodium chloride if there is an inadequate
17 response to dornase alfa based on clinical assessment or lung function testing, or if dornase
18 alfa cannot be used because of intolerance. The Committee noted that treatment should be
19 tailored to the individual patient, taking into account their previous experience of mucoactive
20 agents and any previously demonstrated efficacy.

21 No evidence was found for children under 5 years in the evidence review. The Committee
22 noted that dornase alfa is not licensed for this age group, however it is current practice to
23 prescribe dornase alfa to children under 5.

24 **9.3.7.3 Consideration of economic benefits and harms**

25 The economic evidence found that dornase alfa was more expensive and more effective than
26 placebo. Although those subjective measures of cost-effectiveness cannot be compared to
27 NICE's threshold, the Committee concluded that the evidence did not infer dornase alfa was
28 cost-ineffective to warrant a change from current clinical practice. Furthermore, the clinical
29 evidence showed clinically significant improvements in participants receiving dornase alfa,
30 compared to those receiving placebo, providing evidence that the benefits of dornase alfa
31 could justify the costs. The Committee also added that there is some evidence that early use
32 of dornase alfa is associated with better survival rates. Therefore, despite the high
33 acquisition cost of dornase alfa, the Committee believed, based on the evidence, and their
34 knowledge and expertise, that a recommendation in favour of dornase alfa as the first line
35 treatment, would be a cost-effective use of resources

36 In light of the economic evidence from Suri 2002 and Grieve 2003, the Committee agreed
37 that substantial cost savings could be made by reducing dornase alfa from once daily
38 (current practice in England) to alternate day use. However, the Committee noted that the
39 trial by Suri 2002 aimed to identify a change in FEV₁% and was not powered to measure a
40 change in exacerbations which have a greater treatment cost and impact on health-related
41 quality of life. The Committee noted that alternate day use is encouraged in Wales based on
42 the findings in those studies, but agreed that additional research was needed to justify a
43 deviation to the licensed dose.

44 For these reasons, the Committee agreed that a research recommendation should be made
45 to analyse the clinical and cost-effectiveness of once daily dornase alfa compare to alternate
46 day dornase alfa. To acknowledge this uncertainty, the Committee did not state whether
47 dornase alfa should be offered daily or on alternate days in their recommendation, given that
48 potential cost savings from alternate day use should not be discouraged, where they are
49 considered effective.

1 It was noted that a high proportion of the participants in the trial by Suri 2002 were already
2 receiving dornase alfa at enrolment. When questioned if hypertonic sodium chloride should
3 be used as a first line treatment, the Committee stated that dornase alfa and hypertonic
4 sodium chloride have different mechanisms of action and there was clinical and cost-
5 effective evidence to suggest that the former should be targeted first. The Committee also
6 highlighted that the unpleasant taste and experience of hypertonic saline can lead to poor
7 adherence, subsequently reducing the potential benefits of treatment.

8 However, the Committee agreed that when dornase alfa begins to stabilise respiratory
9 symptoms, there was a role for hypertonic sodium chloride as an add-on, or second-line
10 therapy to dornase alfa, to improve their symptoms, before more costly treatment (mannitol)
11 is considered. The Committee stated higher concentrations of hypertonic sodium chloride are
12 more effective than lower concentrations; although higher concentrations are less well
13 tolerated. Given the unit cost of hypertonic sodium chloride (£0.45 per 4ml, NHS Electronic
14 Drug Tariff November 2016) is equivalent regardless of concentration (7%, 6% or 3%) the
15 Committee wanted to recommend the highest tolerable concentration, before mannitol is
16 considered.

17 When presented with the HTA on mannitol (NICE TA266), the Committee acknowledged that
18 the Appraisal Committee did not find mannitol to be cost-effective as add-on therapy to
19 dornase alfa. However, the Committee accepted the population identified by the Appraisal
20 Committee (people with cystic fibrosis for whom hypertonic saline is not considered
21 appropriate, who cannot use rhDNase because of ineligibility, intolerance or inadequate
22 response to rhDNase, and whose lung function is rapidly declining) as a cost-effective use of
23 resources. Following this, the Committee agreed a recommendation in favour of dornase alfa
24 as first line therapy, and hypertonic as the subsequent add-on, or second line therapy, would
25 reflect the sequence of treatments inferred from the HTA on mannitol.

26 The Committee advised that unlike dornase alfa, acetylcysteine is less well tolerated and is
27 more burdensome to take which may reduce its effects. Moreover, despite the low
28 acquisition cost of acetylcysteine, the clinical evidence review found no significant benefits
29 compared to placebo to make a recommendation in favour of acetylcysteine. Instead, the
30 Committee adopted the recommendation from NICE TA266 that appropriate osmotic agents
31 should be considered before mannitol is considered as an option.

32 Following the review of the clinical and economic evidence, the Committee concluded that
33 additional economic analysis in this area would have limited value to influence their
34 recommendations given that current practice, inferred largely by NICE TA266 was followed.
35 Following this, the Committee iterated the importance of a research recommendation to
36 identify the most clinical and cost effectiveness dose of dornase alfa in people with cystic
37 fibrosis.

38 **9.3.7.4 Quality of evidence**

39 The quality of the evidence presented in this report ranged from very low to high as assessed
40 by GRADE. The main reasons that led to downgrading the quality of the evidence were:

41 For the domain risk of bias, the studies were assigned the same risk of bias as in the
42 Cochrane reviews, and were not individually reviewed. The main biases that lead to
43 downgrading the quality of the evidence were selection, attrition, and reporting bias.

44 Another reason that lead to downgrading the quality of the evidence was the imprecision, as
45 confidence intervals crossed 1 or 2 clinical or default MIDs.

46 High heterogeneity was also a reason to downgrade the quality of the evidence. The
47 Committee noted some studies were underpowered to detect differences between groups.

1 With regards to indirectness, the Committee noted that the participants in the trials
2 comparing mannitol versus placebo, mannitol versus dornase alfa, and mannitol + dornase
3 alfa versus dornase alfa alone had undergone a tolerance test at screening. Those who
4 failed were not entered in the studies, and this limits the generalisability of the results to the
5 general cystic fibrosis population. No serious issues were found regarding the directness of
6 the population or intervention for the other comparisons.

7 **9.3.7.5 Other considerations**

8 No equality issues were identified by the Committee for this review question.

9 The Committee agreed to draft a research recommendation for this topic. They noted the trial
10 by Suri 2002 (comparing daily dornase alfa, alternate day dornase alfa and hypertonic saline)
11 was underpowered to detect a difference in the number of exacerbations. They discussed
12 that reducing the dose, for example from once daily to alternate day dornase alfa, would
13 reduce the burden of treatment, and potentially increase adherence. In addition, substantial
14 cost savings are also anticipated.

15 **9.3.7.6 Key conclusions**

16 The Committee concluded that a mucoactive agent should be given to everyone who has
17 respiratory symptoms or other evidence of lung disease. Dornase alfa should be
18 recommended as first-line treatment, and hypertonic sodium chloride as second-line
19 treatment. The guideline should recommend hypertonic sodium chloride instead of isotonic
20 sodium chloride, but should not mention the concentration of hypertonic sodium chloride.
21 They agreed that Mannitol dry powder should be recommended to adults that fulfil the criteria
22 outlined by the HTA (NICE TA266).

23 **9.3.8 Recommendations**

24 **56. Offer a mucoactive agent to people with cystic fibrosis who have respiratory**
25 **symptoms, or other evidence of lung disease.**

26 **57. Offer rhDNase (dornase alfa; recombinant human deoxyribonuclease) as the first**
27 **choice of mucoactive agent.**

28 **58. If clinical evaluation or lung function testing indicates an inadequate response to**
29 **rhDNase, consider both rhDNase and hypertonic sodium chloride or hypertonic**
30 **sodium chloride alone.**

31 **59. Mannitol dry powder for inhalation is recommended as an option for treating**
32 **cystic fibrosis in adults:**

- 33 • who cannot use rhDNase because of ineligibility, intolerance or
34 inadequate response to rhDNase and
- 35 • whose lung function is rapidly declining (forced expiratory volume in 1
36 second [FEV1] decline greater than 2% annually) and
- 37 • for whom other osmotic agents are not considered appropriate.

38 **[This recommendation is from Mannitol dry powder for inhalation for treating**
39 **cystic fibrosis (NICE technology appraisal 266).]**

40 **60. People currently receiving mannitol whose cystic fibrosis does not meet the**
41 **criteria in recommendation 59 should be able to continue treatment until they and**
42 **their clinician consider it appropriate to stop.**

1 [This recommendation is from Mannitol dry powder for inhalation for treating
2 cystic fibrosis (NICE technology appraisal 266).]

3 **9.3.9 Research recommendations**

4 **3. What is the most clinical and cost effectiveness dose of dornase alfa in people**
5 **with cystic fibrosis?**

6 **Table 86: Research Recommendation justification**

Research question	What is the most clinical and cost effectiveness dose of dornase alfa in people with cystic fibrosis?
Why this is needed	
Importance to 'patients' or the population	Taking daily dornase alfa increases the treatment burden on people with cystic fibrosis who already have complex treatment schedules, including multiple nebulised treatments. It is essential that parents, carers and people with cystic fibrosis are reassured that this dosing frequency is necessary to provide clinical benefit. Parents, carers and people with cystic fibrosis frequently report that reducing the treatment frequency burden increases overall adherence and this will improve the treatment effect. There is some evidence that alternate day dornase alfa is as effective as daily administration, if this is confirmed then overall treatment adherence may improve and moreover cost savings would be made. There is also some evidence that early use of dornase alfa is associated with better survival rates; hence, if alternate day dornase alfa was an option this would potentially lower the threshold for its use.
Relevance to NICE guidance	With this evidence, a definitive recommendation could be made regarding dosing frequency – at present there is varied clinical practice between England and Wales.
Relevance to the NHS	Further evidence in this area will eliminate the need to provide an intense treatment schedule 'just in case' it improves respiratory symptoms. There will be a financial advantage if administration frequency can be reduced in the future.
National priorities	Research priorities from the James Lind Alliance highlighted reducing treatment burden as a priority which would fit with this, in addition to its economic advantages.
Current evidence base	The current evidence base includes mostly small, underpowered, short-term trials with limited methodological rigour and are unable to draw conclusions regarding the long-term impact of dornase alfa in people with cystic fibrosis. Most of the studies assess the licensed dose in people who are not naïve to treatment. There is insufficient evidence as yet to answer the question whether reduced dosing frequency of dornase alfa as a first line treatment is cost-effective, in people with cystic fibrosis who have respiratory symptoms.
Equality	None
Feasibility	The proposed research can be carried out within a realistic timescale and at an acceptable cost. There are no ethical or technical issues.
Other comments	If improvements on alternate day dornase alfa are similar to that given daily, it may be cost-effective to titrate the dose further.

7 **Table 87: Research Recommendation Statements**

Criterion	Explanation
Population	People with a diagnosis of cystic fibrosis with respiratory symptoms
Intervention	Once daily dornase alfa (2.5 mg)

Criterion	Explanation
Comparators	Reduced dosing frequency of dornase alfa, including alternate day dornase alfa (2.5 mg)
Outcomes	<ul style="list-style-type: none"> • Hospitalisations, change in frequency • Pulmonary exacerbations, change in frequency and severity • Lung function (FEV₁, FVC) • Quality of Life (using a validated tool, such CFQ-R or CF-QOL) • Patient preference • Adverse events • Resource use • Unit costs
Study design	Multicentre RCT
Timeframe	Five years of randomisation and 5 year of follow up, providing recruitment numbers are sufficient to achieve population numbers of sufficient size to answer the research question.

1 9.4 Pulmonary Infection

2 Pulmonary infection is the cause of much of the morbidity and mortality associated with
3 cystic fibrosis.

4 Antimicrobial treatment strategies aim to prevent acquisition of infection, eradicate early
5 infection, and suppress chronic respiratory infections where chronic infection ensues from
6 organisms with known or suspected pathogenicity. There is a low threshold to treat
7 respiratory infection. To minimise the risk of antimicrobial resistance, treatment is guided and
8 informed by known or expected microbiological results, based on local surveillance data
9 where necessary.

10 Chronic infection with *P aeruginosa* and *B cepacia* complex leads to a worsening clinical
11 picture, a reduction in respiratory function, more hospital admissions and increased
12 treatment costs. Prompt and aggressive treatment of these organisms is therefore imperative
13 for first and recurrent isolates, following a period free from infection.

14 The scope of these review questions is to review the evidence in people with cystic fibrosis
15 for the prevention of *S aureus* infection, and the treatment of respiratory exacerbations and
16 chronic infection caused by *S aureus*, *P aeruginosa*, *B cepacia* complex and *Aspergillus*
17 species.

18 This chapter will also review evidence for the use of antifungal treatments. The use of
19 antifungal agents is increasing in the management of cystic fibrosis with recognition that
20 fungi such as *Aspergillus* spp may lead to infection as well as a damaging immune response
21 associated with allergic bronchopulmonary aspergillosis.

22 Other organisms such as *Achromobacter xylosoxidans* and *Stenotrophomonas maltophilia*
23 are encountered in people with cystic fibrosis; these have a relatively lower prevalence and
24 are beyond the scope of this current guideline.

1 9.4.1 Prophylaxis

2 **Review question: What is the effectiveness of long-term antimicrobial prophylaxis to**
3 **prevent pulmonary bacterial colonisation with *S aureus* in people with CF?**

4 9.4.1.1 Introduction

5 Pulmonary infection is the cause of much of the morbidity and mortality associated with
6 cystic fibrosis.

7 Antimicrobial treatment strategies aim to prevent acquisition of infection, eradicate early
8 infection and suppress chronic respiratory infections where chronic infection ensues from
9 organisms with known or suspected pathogenicity. There is a low threshold to treat
10 respiratory infection. To minimise the risk of antimicrobial resistance, treatment is guided and
11 informed by known or expected microbiological results, based on local surveillance data
12 where necessary.

13 Chronic infection with *P aeruginosa* and *B cepacia* complex leads to a worsening clinical
14 picture, a reduction in respiratory function, more hospital admissions and increased
15 treatment costs. Prompt and aggressive treatment of these organisms is therefore imperative
16 for first and recurrent isolates, following a period free from infection.

17 The scope of these review questions is to review the evidence in people with cystic fibrosis
18 for the prevention of *S aureus* infection, and the treatment of respiratory exacerbations and
19 chronic infection caused by *S aureus*, *P aeruginosa*, *B cepacia* complex and *Aspergillus*
20 species.

21 This chapter will also review evidence for the use of antifungal treatments. The use of
22 antifungal agents is increasing in the management of cystic fibrosis with recognition that
23 fungi such as *Aspergillus* may lead to infection as well as a damaging immune response
24 associated with allergic bronchopulmonary *aspergillosis*.

25 Other organisms such as *Achromobacter xylosoxidans* and *Stenotrophomonas maltophilia*
26 are encountered in people with cystic fibrosis; these have a relatively lower prevalence and
27 are beyond the scope of this current guideline.

28 9.4.1.2 Description of clinical evidence

29 The aim of this review was to compare the clinical and cost effectiveness of various
30 antimicrobials given as long-term prophylaxis (for more than 3 months) against bacterial
31 infection in people with cystic fibrosis.

32 We searched for systematic reviews of RCTs and RCTs aimed at assessing the
33 effectiveness of long-term antimicrobial prophylaxis to prevent pulmonary bacterial infection
34 with *S aureus* in people with cystic fibrosis. Cross-over trials were not considered for
35 inclusion, as this study design does not allow evaluation of the effects of prophylaxis on long-
36 term outcomes measures.

37 For full details see review protocol in Appendix D.

38 Two systematic reviews were identified for potential inclusion in this review (McCaffery 1999,
39 Smyth 2014).

40 One systematic review (McCaffery 1999) was finally excluded as the quality was assessed
41 as low according to AMSTAR checklist (score of 5 out of 11). The individual studies included
42 in this review were checked for potential inclusion.

43 One Cochrane systematic review (Smyth 2014) was included as the quality was assessed as
44 high according to AMSTAR checklist (score of 10 out of 11). This systematic review included

1 4 RCTs (Chatfield 1991, Schlesinger 1984, Stutman 2002, Weaver 1994, Beardsmore 1995).
2 One study (Schlesinger 1984) was excluded from our review as it included treatments that
3 were not specified in the evidence review protocol and it did not report on any outcomes of
4 interest.

5 The data and risk of bias assessment from the systematic review were used where possible.
6 The individual studies were also retrieved full copy for completeness. Data for other
7 outcomes of interests included in the protocol, but not included in the Cochrane SR, was
8 directly extracted from the individual studies.

9 No further studies were identified in our search.

10 With regards to the population, the studies included in the Cochrane review included infants
11 or small children (under 2 years) with confirmed cystic fibrosis.

12 The trials compared the effectiveness of long-term prophylactic antibiotic treatment with
13 placebo and treatment “as required”. The treatments evaluated included continuous oral
14 Flucloxacillin and continuous oral Cephalexin.

15 In relation to the outcomes, evidence was found for number of children who *S aureus* was
16 identified at least once; lung function, measured as FEV₁, minor adverse events and
17 identification of *P aeruginosa*. Where no evidence was retrieved for time to next pulmonary
18 exacerbation, number of children experiencing a pulmonary exacerbations and number of
19 children requiring hospitalisation due to infection was taken as a proxy outcome.

20 No results were identified for time to identification of *S aureus*, major adverse events and
21 emergence of the resistant organisms.

22 Evidence from these are summarised in the clinical GRADE evidence profile below (Table 89
23 and Table 90). See also the study selection flow chart in Appendix F, forest plots in Appendix
24 I, study evidence tables in Appendix G and exclusion list in Appendix H.

25 9.4.1.3 Summary of included studies

26 A summary of the studies that were included in this review are presented in Table 88.

27 **Table 88: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Smyth 2014 Cochrane SR	Intervention Prophylactic anti- staphylococcal antibiotic + antibiotic 'as required'	People with a confirmed diagnosis of cystic fibrosis, of any age.	<ul style="list-style-type: none"> • Number of positive pathogen cultures (<i>S aureus</i>) identified during study period - measured as number of children from whom <i>S aureus</i> was isolated at least once by year of age (n/N) • Lung function measured as FEV₁ • Evidence of inflammation in CT scan - Proxy outcome, X-ray scores (Crispin-Norman score at 1.3 years) • Pulmonary exacerbation - Proxy outcome, 	SR AMSTAR score: 10/11
	Comparison Placebo + antibiotic 'as required'			

Study	Intervention/ Comparison	Population	Outcomes	Comments
			<ul style="list-style-type: none"> number of children requiring admission Adherence to treatment Adverse events, minor events Number of children in whom <i>P aeruginosa</i> was identified 	
Primary studies included in the Cochrane SR				
Chatfield 1991 (UK) RCT	<p>Intervention</p> <ul style="list-style-type: none"> Continuous oral Flucloxacillin <p>Comparison</p> <p>Placebo + antibiotic 'as required'</p>	<p>Infants with a confirmed diagnosis of cystic fibrosis.</p> <p>Age: not reported</p> <ul style="list-style-type: none"> 	<ul style="list-style-type: none"> Number of positive pathogen cultures (<i>S aureus</i>) identified during study period - measured as number of children from whom <i>S aureus</i> was isolated at least once by year of age (n/N) Evidence of inflammation in CT scan (only for < 5 yrs) - Proxy outcome, X-ray scores (Crispin-Norman score at 1.3 years) Pulmonary exacerbation - Proxy outcome, number of children requiring admission (annualised rates) Number of children in whom <i>P aeruginosa</i> was identified 	Included in Smyth 2014
Stutman 2002 (USA) RCT	<p>Intervention</p> <ul style="list-style-type: none"> Continuous cephalixin <p>Comparison</p> <p>Placebo + antibiotic 'as required'</p>	<p>Infants and children with a confirmed diagnosis of cystic fibrosis.</p> <p>Age: 4 to 24 months</p> <ul style="list-style-type: none"> 	<ul style="list-style-type: none"> Number of positive pathogen cultures (<i>S aureus</i>) identified during study period - measured as number of children from whom <i>S aureus</i> was isolated at least once by year of age (n/N) Lung function - measured as FEV₁ at 6 years Pulmonary exacerbation - Proxy outcome, number of children requiring admission (annualised rates) Minor adverse events, including generalised rash, nappy rash, increased stool frequency 	Included in Smyth 2014

Study	Intervention/ Comparison	Population	Outcomes	Comments
			<ul style="list-style-type: none"> Number of children in whom <i>P aeruginosa</i> was identified 	
Weaver 1994 (UK) RCT	Intervention <ul style="list-style-type: none"> continuous oral Flucloxacillin Comparison Placebo + antibiotic 'as required'	Infants with a confirmed diagnosis of cystic fibrosis, of any age. Age: not reported <ul style="list-style-type: none"> 	<ul style="list-style-type: none"> Number of positive pathogen cultures (<i>S aureus</i>) identified during study period - measured as number of children from whom <i>S aureus</i> was isolated at least once by year of age (n/N) Pulmonary exacerbation - Proxy outcome, number of children requiring admission (annualised rates) Number of children in whom <i>P aeruginosa</i> was identified 	Included in Smyth 2014 Additional data obtained from Beardsmore 1995

1 AB: antibiotics; SR: systematic review

2 9.4.1.4 Clinical evidence profile

3 The summary clinical evidence profiles for this review question are presented in Table 89
4 and Table 90.

5 **Table 89: Summary clinical evidence profile: Comparison 1. Continuous oral**
6 **Flucloxacillin versus antibiotics 'as required'**

Comparison 1. Continuous oral Flucloxacillin versus antibiotics 'as required'						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Continuous oral Flucloxacillin, antibiotic prophylaxis				
Number of children from whom <i>S aureus</i> isolated at least once Follow-up: mean 1 year	373 per 1000	201 per 1000 (101 to 395)	RR 0.54 (0.27 to 1.06)	96 (Chatfield 1991)	⊕⊕⊕⊖ very low ^{1,2}	
Number of children from whom <i>S aureus</i> isolated at least once Follow-up: mean 2 years	Study population		RR 0.44 (0.25 to 0.77)	149 (Chatfield 1991, Weaver 1994)	⊕⊕⊕⊖ low ³	
	425 per 1000	187 per 1000 (106 to 327)				
	Moderate 483 per 1000	213 per 1000 (121 to 372)				

Comparison 1. Continuous oral Flucloxacillin versus antibiotics 'as required'						
Number of children from whom <i>S aureus</i> isolated at least once Follow-up: mean 3 years	431 per 1000	224 per 1000 (125 to 392)	RR 0.52 (0.29 to 0.91)	119 (Chatfield 1991)	⊕⊕⊕⊕ very low ^{1,2}	
Number of children from whom <i>P aeruginosa</i> isolated at least once Follow-up: mean 1 year	59 per 1000	136 per 1000 (36 to 514)	RR 2.32 (0.62 to 8.73)	95 (Chatfield 1991)	⊕⊕⊕⊕ very low ^{1,4}	
Number of children from whom <i>P aeruginosa</i> isolated at least once Follow-up: mean 2 years	Study population		RR 0.74 (0.34 to 1.61)	149 (Chatfield 1991, Weaver 1994)	⊕⊕⊕⊕ very low ^{3,4}	
	175 per 1000	129 per 1000 (60 to 282)				
	Moderate					
	217 per 1000	161 per 1000 (74 to 349)				
Number of children from whom <i>P aeruginosa</i> isolated at least once Follow-up: mean 3 years	212 per 1000	168 per 1000 (78 to 354)	RR 0.79 (0.37 to 1.67)	120 (Chatfield 1991)	⊕⊕⊕⊕ very low ^{1,4}	
Number of children requiring admission due to pulmonary exacerbations (annualised rates) Follow-up: mean 3 years	333 per 1000	327 per 1000 (197 to 540)	RR 0.98 (0.59 to 1.62)	124 (Chatfield 1991, Weaver 1994)	⊕⊕⊕⊕ very low ^{3,4}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; RR: Risk ratio						

- 1 The quality of the evidence was downgraded by 2 as this study was rated as high risk of bias (Cochrane review Smyth 2014)
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crosses 1 default MID
- 3 The quality of the evidence was downgraded by 2 as these studies was rated as high and moderate risk of bias (Cochrane review Smyth 2014)
- 4 The quality of the evidence was downgraded by 2 as the 95% CI crosses 2 default MIDs

7 **Table 90: Summary clinical evidence profile: Comparison 2. Continuous oral**
8 **Cephalexin versus antibiotics 'as required'**

Comparison 2. Continuous oral Cephalexin versus antibiotics 'as required'						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Antibiotics 'as required'	Continuous oral Cephalexin, antibiotic prophylaxis				
Number of children from	468 per 1000	145 per 1000 (79 to 266)	RR 0.31	152 (Stutman)	⊕⊕⊕⊕ moderate ¹	

Comparison 2. Continuous oral Cephalexin versus antibiotics 'as required'						
whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 1 years			(0.17 to 0.57)	2002)		
Number of children from whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 2 years	658 per 1000	217 per 1000 (145 to 336)	RR 0.33 (0.22 to 0.51)	166 (Stutman 2002)	⊕⊕⊕⊖ moderate ²	
Number of children from whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 3 years	688 per 1000	289 per 1000 (199 to 406)	RR 0.42 (0.29 to 0.59)	141 (Stutman 2002)	⊕⊕⊕⊖ moderate ³	
Number of children from whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 4 years	839 per 1000	352 per 1000 (252 to 495)	RR 0.42 (0.3 to 0.59)	127 (Stutman 2002)	⊕⊕⊕⊖ moderate ⁴	
Number of children from whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 5 years	850 per 1000	348 per 1000 (238 to 501)	RR 0.41 (0.28 to 0.59)	98 (Stutman 2002)	⊕⊕⊕⊖ low ⁵	
Number of children from whom <i>S aureus</i> isolated at least once Respiratory cultures Follow-up: mean 6 years	778 per 1000	280 per 1000 (140 to 552)	RR 0.36 (0.18 to 0.71)	43 (Stutman 2002)	⊕⊕⊕⊖ low ⁶	
Lung function FEV ₁ Follow-up: mean 6 years	The mean lung function in the control group was 115.8	The mean lung function in the intervention groups was 2.3 lower (13.59 lower to		119 (Stutman 2002)	⊕⊕⊕⊖ very low ^{7,8}	

Comparison 2. Continuous oral Cephalexin versus antibiotics 'as required'						
Any pulmonary exacerbations % Follow-up: mean 6 years	The mean number of pulmonary exacerbation in the control group was 66.8	8.99 higher) The mean number of pulmonary exacerbations in the intervention groups was 4.9 lower (22.24 lower to 12.44 higher)		119 (Stutman 2002)	⊕⊕⊕⊕ very low ^{7,9}	
Number of children requiring admission due to pulmonary exacerbations (annualised rates) not reported Follow-up: mean 6 years	78 per 1000	74 per 1000 (20 to 260)	RR 0.94 (0.26 to 3.32)	119 (Stutman 2002)	⊕⊕⊕⊕ very low ^{7,9}	
Adherence to treatment Parents self-report Follow-up: mean 6 years	The mean adherence to treatment in the control groups was 85 %	The mean adherence to treatment in the intervention groups was 5 higher (0 to 0 higher)		119 (Stutman 2002)	⊕⊕⊕⊕ moderate ^{7,10}	
Minor adverse events - generalised rash Parents self-report Follow-up: mean 6 years	The mean number of generalised rash events in the control group was 0.2	The mean – number of generalised rash events in the intervention groups was 0.4 higher (0.07 lower to 0.87 higher)		119 (Stutman 2002)	⊕⊕⊕⊕ moderate ⁷	
Minor adverse events - nappy rash Parents self-report Follow-up: mean 6 years	The mean number of nappy rash events in the control group was 3.1	The mean – number of nappy rash events in the intervention groups was 0.9 higher (1.06 lower to 2.86 higher)		119 (Stutman 2002)	⊕⊕⊕⊕ moderate ⁷	
Minor adverse events - increased stool frequency Parents self-report Follow-up: mean 6 years	The mean number of increased stool frequency events in the control group was 4.1	The mean minor increased stool frequency events in the intervention groups was 0.2 higher (2.18 lower to 2.58 higher)		119 (Stutman 2002)	⊕⊕⊕⊕ moderate ⁷	

Comparison 2. Continuous oral Cephalexin versus antibiotics 'as required'						
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 1 years	312 per 1000	358 per 1000 (231 to 564)	RR 1.15 (0.74 to 1.81)	152 (Stutman 2002)	⊕⊕⊕⊕ very low ^{1,9}	
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 2 years	506 per 1000	435 per 1000 (314 to 603)	RR 0.86 (0.62 to 1.19)	166 (Stutman 2002)	⊕⊕⊕⊕ low ^{2,11}	
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 3 years	594 per 1000	582 per 1000 (445 to 772)	RR 0.98 (0.75 to 1.3)	141 (Stutman 2002)	⊕⊕⊕⊕ very low ^{3,9}	
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 4 years	Study population		RR 1.1 (0.83 to 1.45)	127 (Stutman 2002)	⊕⊕⊕⊕ low ^{4,11}	
	589 per 1000	648 per 1000 (489 to 854)				
	Moderate					
	589 per 1000	648 per 1000 (489 to 854)				
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 5 years	550 per 1000	709 per 1000 (512 to 979)	RR 1.29 (0.93 to 1.78)	98 (Stutman 2002)	⊕⊕⊕⊕ very low ^{5,11}	
Number of children from whom <i>P. aeruginosa</i> identified at least once Follow-up: mean 6 years	667 per 1000	880 per 1000 (613 to 1000)	RR 1.32 (0.92 to 1.89)	43 (Stutman 2002)	⊕⊕⊕⊕ very low ^{6,11}	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio</p>						

- 1 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the evidence was downgraded by 1 for this outcome, as the losses to follow up are over 20% (n=152; N=209).
- 2 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the evidence was downgraded by 1 for this outcome, as the losses to follow up are over 20% (n=166; N=209).
- 3
- 4

1 3 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the
2 evidence was downgraded by 1 for this outcome, as the losses to follow up are over 20% (n=141; N=209).
3 4 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the
4 evidence was downgraded by 1 for this outcome, as the losses to follow up are over 20% (n=127; N=209).
5 5 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the
6 evidence was downgraded by 2 for this outcome, as the losses to follow up are over 50% (n=98; N=209).
7 6 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the
8 evidence was downgraded by 2 for this outcome, as the losses to follow up are over 50% (n=43; N=209).
9 7 This study was assessed by the Cochrane review Smyth 2014 as low risk of bias. However, the quality of the
10 evidence was downgraded by 1 for this outcome, as the losses to follow up are over 20% (n=119; N=209).
11 8 The quality of the evidence was downgraded by 2, as the 95% CI crossed 2 clinical MIDs
12 9 The quality of the evidence was downgraded by 2, as the 95% CI crossed 2 default MIDs
13 10 Imprecision is not calculable with the data reported
14 11 The quality of the evidence was downgraded by 1, as the 95% CI crossed 1 default MID for dichotomous
15 outcomes

16 9.4.1.5 Economic evidence

17 No economic evaluations of prophylaxis treatment were identified in the literature search
18 conducted for this guideline. Full details of the search and economic article selection flow
19 chart can be found in Appendix E and F, respectively.

20 This review question was not prioritised for de novo economic modelling. To aid
21 consideration of cost-effectiveness relevant resource and cost use data are presented in
22 Appendix K.

23 9.4.1.6 Evidence statements

24 249.4.1.6.1 Comparison 1. Continuous oral Flucloxacillin versus antibiotics 'as required'

25 Time to identification of the pathogen (*S aureus*) in sputum culture

26 No evidence was found for this critical outcome.

27 Number of positive pathogen cultures (*S aureus*) identified

28 Very low quality evidence from 1 RCT with 96 infants with cystic fibrosis showed no clinically
29 significant difference in the number of children in whom *S aureus* was identified between the
30 group who were receiving continuous oral Flucloxacillin prophylaxis and the group who were
31 receiving antibiotics 'as required' during the first year of follow-up.

32 Low quality evidence from 2 RCTs with 149 infants with cystic fibrosis showed that there was
33 a clinically significant lower number of children in whom *S aureus* was identified in the group
34 who were receiving continuous oral Flucloxacillin prophylaxis compared to the group who
35 were receiving antibiotics 'as required' during the first 2 years of follow-up.

36 Very low quality evidence from 1 RCT with 119 infants with cystic fibrosis showed that there
37 was a clinically significant lower number of children in whom *S aureus* was identified
38 between the group who were receiving continuous oral Flucloxacillin prophylaxis and the
39 group who were receiving antibiotics 'as required' during the first 3 years of follow-up.

40 Lung function

41 No evidence was found for this important outcome.

42 Pulmonary exacerbation

43 Very low quality evidence from 2 RCTs with 124 infants with cystic fibrosis showed no
44 clinically significant difference in the number of annual hospital admissions due to pulmonary
45 exacerbations between the children who were receiving continuous oral Flucloxacillin

1 prophylaxis and the children who were receiving antibiotics 'as required' during the 3 years
2 follow-up.

3 **Quality of life**

4 No evidence was found for this important outcome.

5 **Minor adverse events**

6 No evidence was found for this important outcome.

7 **Major adverse events**

8 No evidence was found for this important outcome.

9 **Identification of *P aeruginosa***

10 Very low quality evidence from 1 RCT with 95 infants with cystic fibrosis showed no clinically
11 significant difference in the number of children in whom *P aeruginosa* was identified between
12 the group who were receiving continuous oral Flucloxacillin prophylaxis and the group who
13 were receiving antibiotics 'as required' during the first year of follow-up.

14 Very low quality evidence from 2 RCTs with 149 infants with cystic fibrosis showed no
15 clinically significant difference in the number of children in whom *P aeruginosa* was identified
16 between the group who were receiving continuous oral Flucloxacillin prophylaxis and the
17 group who were receiving antibiotics 'as required' during the first 2 years of follow-up.

18 Very low quality evidence from 1 RCT with 120 infants with cystic fibrosis showed no
19 clinically significant difference in the number of children in whom *P aeruginosa* was identified
20 between the group who were receiving continuous oral Flucloxacillin prophylaxis and the
21 group who were receiving antibiotics 'as required' during the first 3 years of follow-up.

22 **Adherence to treatment**

23 No evidence was found for this important outcome.

24 **Emergence of resistant organisms**

25 No evidence was found for this important outcome.

269.4.1.6.2 **Comparison 2. Continuous oral Cephalexin versus antibiotics 'as required'**

27 **Time to identification of the pathogen (*S aureus*) in sputum culture**

28 No evidence was found for this critical outcome.

29 **Number of positive pathogen cultures (*S aureus*) identified**

30 Moderate quality evidence from 1 RCT with 152 children with cystic fibrosis < 2 years
31 (n=152) showed that there was a clinically significant lower number of children in whom *S*
32 *aureus* was identified in the group who were receiving continuous oral Cephalexin
33 prophylaxis compared to the group who were receiving antibiotics 'as required' during the
34 first year of follow-up.

35 Moderate quality evidence from 1 RCT (n=166) showed that there was a clinically significant
36 lower number of children in whom *S aureus* was identified in the group who were receiving
37 continuous oral Cephalexin prophylaxis compared to the group who were receiving
38 antibiotics 'as required' during the first 2 years of follow-up.

1 Moderate quality evidence from 1 RCT (n=141) showed that there was a clinically significant
2 lower number of children in whom *S aureus* was identified in the group who were receiving
3 continuous oral Cephalexin prophylaxis compared to the group who were receiving
4 antibiotics 'as required' during the first 3 years of follow-up.

5 Moderate quality evidence from 1 RCT (n=127) showed that there was a clinically significant
6 lower number of children in whom *S aureus* was identified in the group who were receiving
7 continuous oral Cephalexin prophylaxis compared to the group who were receiving
8 antibiotics 'as required' during the first 4 years of follow-up.

9 Low quality evidence from 1 RCT (n=98) showed that there was a clinically significant lower
10 number of children in whom *S aureus* was identified in the group who were receiving
11 continuous oral Cephalexin prophylaxis compared to the group who were receiving
12 antibiotics 'as required' during the first 5 years of follow-up.

13 Low quality evidence from 1 RCT (n=43) showed that there was a clinically significant lower
14 number of children in whom *S aureus* was identified in the group who were receiving
15 continuous oral Cephalexin prophylaxis compared to the group who were receiving
16 antibiotics 'as required' during the 6 years of follow-up.

17 **Lung function: FEV₁**

18 Very low quality evidence from 1 RCT (n=119) showed no clinically significant difference in
19 lung function, measured as FEV₁, between the children who were receiving continuous oral
20 Cephalexin prophylaxis and the children who were receiving antibiotics 'as required' at 6
21 years follow-up.

22 **Pulmonary exacerbation**

23 Very low quality evidence from 1 RCT (n=119) showed no clinically significant difference in
24 the percentage of pulmonary exacerbations between the children who were receiving
25 continuous oral Cephalexin prophylaxis and the children who were receiving antibiotics 'as
26 required' during the six years follow-up.

27 Very low quality evidence from 1 RCT (n=119) showed no clinically significant difference in
28 the number of annual hospital admissions due to pulmonary exacerbations between the
29 children who were receiving continuous oral Cephalexin prophylaxis and the children who
30 were receiving antibiotics 'as required' during the 6 years follow-up.

31 **Quality of life**

32 No evidence was found for this important outcome.

33 **Minor adverse events**

34 Moderate quality evidence from 1 RCT (n=119) showed no clinically significant difference in
35 the report of generalised rash, nappy rash and stool frequency between the children who
36 were receiving continuous oral Cephalexin prophylaxis and the children who were receiving
37 antibiotics 'as required' during the 6 year study duration.

38 **Major adverse events**

39 No evidence was found for this important outcome.

40 **Identification of *P aeruginosa***

41 Very low quality evidence from 1 RCT (n=152) showed no clinically significant difference in
42 the number of children in whom *P aeruginosa* was identified between the group who were

1 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
2 antibiotics 'as required' during the first year of follow-up.

3 Low quality evidence from 1 RCT (n=166) showed no clinically significant difference in the
4 number of children in whom *P aeruginosa* was identified between the group who were
5 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
6 antibiotics 'as required' during the first 2 years of follow-up.

7 Very low quality evidence from 1 RCT (n=141) showed no clinically significant difference in
8 the number of children in whom *P aeruginosa* was identified between the group who were
9 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
10 antibiotics 'as required' during the first 3 years of follow-up.

11 Low quality evidence from 1 RCT (n=127) showed no clinically significant difference in the
12 number of children in whom *P aeruginosa* was identified between the group who were
13 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
14 antibiotics 'as required' during the first 4 years of follow-up.

15 Very low quality evidence from 1 RCT (n=98) showed no clinically significant difference in the
16 number of children in whom *P aeruginosa* was identified between the group who were
17 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
18 antibiotics 'as required' during the first 5 years of follow-up.

19 Very low quality evidence from 1 RCT (n=43) showed no clinically significant difference in the
20 number of children in whom *P aeruginosa* was identified between the group who were
21 receiving continuous oral Cephalexin prophylaxis and the group who were receiving
22 antibiotics 'as required' during the 6 years of follow-up.

23 **Adherence to treatment**

24 Moderate quality evidence from 1 RCT (n=119) showed a higher level of adherence to
25 treatment in the group of children who were receiving continuous oral Cephalexin prophylaxis
26 (85% vs. 80%). The uncertainty around this could not be calculated.

27 **Emergence of resistant organisms**

28 No evidence was found for this important outcome.

299.4.1.6.3 **Economic evidence statements**

30 **9.4.1.7 No evidence on cost-effectiveness in people with cystic fibrosis was available for this**
31 **review.Evidence to recommendations**

329.4.1.7.1 **Relative value placed on the outcomes considered**

33 The aim of this review was to compare the clinical and cost effectiveness of various
34 antimicrobials given as long-term prophylaxis (for more than 3 months) against bacterial
35 infection in people with cystic fibrosis.

36 The Committee identified time to identification of the pathogen (*S aureus*) in respiratory
37 samples, the time to the next pulmonary exacerbation and the development of *P aeruginosa*
38 infection as critical outcomes for this evidence review. If there was no evidence for time to
39 next pulmonary exacerbation, the number of people with cystic fibrosis experiencing a
40 pulmonary exacerbations and number of people with cystic fibrosis being admitted to hospital
41 with to pulmonary exacerbations were taken as alternative proxy outcomes.

42 In addition to the critical outcomes, lung function (measured by FEV₁ or LCI), evidence of
43 inflammation on CT scanning (in children under 5 years of age), quality of life, adherence to

1 treatment (or patient preference), adverse events and the emergence of resistant organisms
2 were considered important outcomes.

39.4.1.7.2 **Consideration of clinical benefits and harms**

4 The Committee noted that it is accepted that *S aureus* can cause serious lung disease in
5 cystic fibrosis. The isolation of *S aureus* from respiratory samples means that the respiratory
6 tract is colonised or infected with this pathogen and this can lead to pulmonary inflammation
7 and progressive lung disease.

8 The Committee noted that the evidence showed anti-staphylococcal prophylaxis with an
9 antimicrobial agent (either flucoxacin or cephalixin) led to a decreased number of children
10 in whom *S aureus* was isolated. The quality of the evidence for this outcome ranged from
11 very low to moderate. Despite the fact that the evidence did not show this was associated
12 with clinical benefit, there being no improvement in lung function or reduction in
13 exacerbations in children given prophylaxis compared with those who were not, the reduction
14 in *S aureus* was nevertheless a critically important outcome. Overall, the evidence did not
15 reveal the occurrence of adverse events with prophylaxis.

16 The Committee was concerned about the theoretical possibility that long-term antimicrobial
17 prophylaxis for *S aureus* might be associated with an increased risk infection with *P*
18 *aeruginosa*. Although the evidence did not demonstrate this, the Committee noted that the
19 quality of the evidence for this outcome ranged from very low to low. The Committee
20 observed that given the widespread expert consensus that this risk is a concern, they agreed
21 that it could be mitigated by recommending that flucloxacillin be used rather than cephalixin.
22 The fact that cephalosporins are broad spectrum is postulated to be the reason why an
23 increase in pseudomonas isolation may be seen but this is not known with certainty.
24 Recommending flucloxacillin rather than a cephalosporin was in keeping with current usual
25 practice. Even with flucloxacillin, the Committee recognised that the benefits of antimicrobial
26 prophylaxis in terms of preventing colonisation with *S aureus* might possibly be outweighed
27 by the risk of *P aeruginosa* and so a strong recommendation in favour such prophylaxis
28 could not be justified.

299.4.1.7.3 **Consideration of economic benefits and harms**

30 The Committee believed, based on their own expertise, that colonisation with *S. aureus* was
31 likely to be associated with an increased risk of pulmonary disease and with a worse
32 prognosis compared to no colonisation. The clinical benefits of prophylaxis, although not
33 demonstrated by the available evidence, might well be important, and if so prophylaxis is
34 likely to be cost-effective.

35 In contrast to current recommendations in the United States, the Committee did not think
36 there were grounds to advise the non-use of prophylaxis, given the low cost of prophylaxis
37 treatment and the potentially serious consequences of *S.aureus* pulmonary infection that
38 would outweigh the cost of prophylaxis.

39 The Committee believed cephalixin and flucloxacillin were similar in terms of efficacy but
40 noted that cephalixin is a broader spectrum antibiotic that could increase the risk of
41 *P.aeruginosa* infection. For this reason, the Committee agreed that using the more expensive
42 flucloxacillin, would be a cost-effective choice, because it is a narrower spectrum antibiotic
43 reducing the expected cost of a *P.aeruginosa* infection.

44 The Committee noted that the cost of flucloxacillin varied substantially according to the
45 preparation used, with oral solutions costing more than capsules (NHS Electronic Drug Tariff
46 November 2016: 250mg/5ml oral solution sugar free, £1.32/5ml versus 250mg capsules,
47 £0.05). Prophylaxis in infants would require the use of oral solutions and the Committee
48 believed that sugar free solutions were preferable although more expensive. However, once

1 children were old enough to take capsule preparations they believed these should be used if
2 they are cheaper and equally effective.

3 The Committee recognised that long-term prophylaxis could be burdensome for the person
4 with cystic fibrosis and their parents or carers; particularly when products are unpalatable,
5 and as a medications for cystic fibrosis increase with age. Moreover, long-term use of
6 flucloxacillin would be costly at approximately £67 per month (costed on a BNF
7 recommended dose of 125 mg twice daily: 125mg/5ml oral solution sugar free; basic price,
8 £21.97; quantity 100ml). For this reason, the Committee made a recommendation to offer
9 prophylaxis treatment up to the age of 6 years, as there was no evidence it provided benefit
10 beyond this age, to justify the cost and burden of treatment beyond this time. The Committee
11 also added that this could reduce resource use in this area, if children who are receiving
12 prophylactic treatment over this age, discontinue treatment following the recommendation.

139.4.1.7.4 **Quality of evidence**

14 Two of the studies included in the Cochrane review (Smyth 2014) compared oral flucloxacillin
15 versus antibiotics as required. The quality of the evidence was moderate to very low quality
16 as assessed by GRADE. The main reasons that lead to downgrading the quality of the
17 evidence were the moderate risk of bias found in the studies and the levels of imprecision.

18 One study included in the Cochrane review (Smyth 2014) compared oral cephalixin versus
19 antibiotics as required. The quality of the evidence was moderate to very low quality as
20 assessed by GRADE. The main reason for downgrading the quality of the evidence was the
21 large number of losses to follow-up. It is known that the participants who leave the study do
22 normally differ to those who remain, therefore the results have to be interpreted with caution.
23 The outcome about adherence to treatment was reported narratively only.

249.4.1.7.5 **Other considerations**

25 In the absence of strong evidence in favour of the use of antimicrobial prophylaxis to prevent
26 pulmonary bacterial colonisation with *S Aureus*, the Committee agreed that other aspects
27 were to be considered when deciding whether to start treatment or not. The preferences of
28 the parents were discussed. It was acknowledged that some parents want their children to
29 take anti-staphylococcal treatment; whereas others are more reluctant as they cannot see a
30 clear benefit of having treatment. The Committee agreed that it was important to discuss
31 prophylaxis with the parents and a recommendation was made to this effect.

32 Patient tolerance of prophylaxis was discussed by the Committee. As mentioned above,
33 clinical practice in the UK is to favour the use of flucloxacillin, but other alternatives (for
34 example Cephalixin) can be considered if there are significant side effects.

35 An important concern raised by the Committee was the duration of treatment. Currently there
36 is considerable variation in clinical practice. It was noted that although there is a common
37 view that prophylaxis should continue until the age of three (as suggested by the CF Trust),
38 in many cases it is continued beyond that age. The study included in the review followed
39 children up to the age of 6. There was no conclusive evidence that there was continuing
40 benefit of prophylaxis treatment up to this age, but it was the consensus view of the
41 Committee that this was a reasonable duration of treatment. Given the lack of evidence
42 comparing different lengths of treatment, and the concerns raised by patients and parents
43 alike, the Committee concluded that this should be a priority for research. However, they
44 agreed not to draft a research recommendation as there is an ongoing trial that will address
45 this issue (cystic fibrosis CF STAR study).

46 No equality issues were identified by the Committee for this review question.

19.4.1.7.6 **Key conclusions**

2 The Committee concluded that given the quality of the evidence, it was not possible to
3 strongly recommend anti-staphylococcal prophylaxis for children with cystic fibrosis. Instead,
4 the Committee agreed that it should be considered in discussion with parents or carers,
5 taking account of the potential benefits and disadvantages, flucloxacillin should be the first
6 choice given that cephalexin may be associated with a higher rate of *P aeruginosa* growth or
7 isolation.

8 **9.4.2 Acute**

9 **Review question: What is the effectiveness of antimicrobial treatment for acute**
10 **pulmonary infection or those with an exacerbation in children and adults with cystis**
11 **fibrosis?**

12 **9.4.2.1 Description of clinical evidence**

13 The aim of this review was to compare the clinical and cost effectiveness of different
14 antimicrobial regimens in achieving clinical resolution of acute pulmonary infection or
15 exacerbation in children and adults with cystic fibrosis.

16 We looked for studies that included children and adults with cystic fibrosis who presented
17 with clinical manifestations suggesting development of an acute pulmonary infection or those
18 with an exacerbation and who are already known to have a positive sputum or airway culture
19 for one of the following pathogens at entry to the trial:

- 20 • *S aureus*
- 21 • *P aeruginosa*
- 22 • *B cepacia complex*
- 23 • *Haemophilus influenza*
- 24 • Nontuberculous *mycobacteria* (*Mycobacterium avium* complex and *Mycobacterium*
25 *abscessus*).
- 26 • We also looked for studies that included children and adults with cystic fibrosis who
27 present with clinical manifestations suggesting development of an acute pulmonary
28 infection or those with an exacerbation without an identified pathogen at trial entry.

29 Pulmonary exacerbation was defined as:

- 30 1. Fuchs definition (original form (4/16 symptoms leading to IV antibiotic treatment) or
31 modified form (4/16 symptoms leading to any change in antibiotic therapy).
- 32 2. or
- 33 3. European Cystic Fibrosis Society Consensus definition: “need for additional antibiotic
34 treatment as indicated by a recent change in at least 2 of 6 defined symptoms”.

35 However, the definition of pulmonary exacerbation used in the study was also accepted.

36 Additionally, acute infection was defined as a person with cystic fibrosis who is found, on
37 routine microbiological investigation, to have a significant respiratory pathogen (newly
38 identified infection).

39 We searched for systematic reviews of RCTs and RCTs. Systematic reviews were assessed
40 for inclusion against the protocol, and if relevant, their quality was assessed using AMSTAR.
41 High-quality systematic reviews were included in our review, and where possible, data and
42 quality assessment was taken directly from the review. Individual studies were also retrieved
43 for completeness and accuracy, and were also checked for additional outcomes of interest.
44 Low-quality SR were excluded from our review, but the list of included studies was checked
45 to identify relevant trials.

1 For full details see review protocol in Appendix D.

2 The results are presented separately for each pathogen.

39.4.2.1.1 *P aeruginosa*

4 The interventions that were included in the protocol for the treatment of infection with or
5 exacerbation due to *P aeruginosa* were: Ciprofloxacin (oral), Aztreonam (inhaled or IV),
6 Ceftazidime (IV), Meropenem (IV), Piperacillin-Tazobactam (IV), Fosfomycin (IV), Ticarcillin-
7 Clavulanate (IV), Aztreonam (inhaled or IV), Chloramphenicol (oral). For first infection only
8 we also considered sequencing antibiotics: Ciprofloxacin (oral) then either Colistin or
9 Tobramycin (inhaled).

10 Four Cochrane systematic reviews were identified in the search (Elphick 2016, Hurley 2015,
11 Hewer 2014, Remington 2016).

12 Two reviews were included:

- 13 • Hurley (2015) evaluated if intravenous antibiotics for the treatment of pulmonary
14 exacerbations in people with CYSTIC FIBROSIS improve short- and long-term clinical
15 outcomes. 13 randomised control trials (RCTs) from this Cochrane review were included
16 (Blumer 2005; Conway 1997; De Boeck 1989, Elborn 1992; Gold 1985; Macystis
17 fibrosisarlane 1985; Master 2001; McCarty 1988; Richard 1997; Salh 1992; Schaad 1987;
18 Schaad 1989; Wesley 1988). The outcome 'eradication of pathogen' was not included in
19 this Cochrane review. Each paper was checked for this outcome and 2 RCTs (Richard
20 1997 and Schaad 1989) were included.
- 21 • Langton (2014) evaluated whether antibiotic treatment of early *P aeruginosa* infection in
22 children and adults with CYSTIC FIBROSIS alters clinical and microbiological outcomes.
23 Two RCTs from this Cochrane review were included (Proesmans 2013, Taccetti 2012).
24 These RCTs aimed to eradicate acute infection (first positive isolate of *P aeruginosa*).

25 Two reviews were excluded.

- 26 • Elphick (2016): no additional RCTs were included from this Cochrane review.
- 27 • Remington (2016): no additional RCTs were included from this Cochrane review.

28 No additional relevant trials were identified in our search.

- 29 • The 13 included studies from the Hurley (2015) Cochrane systematic review evaluated the
30 effectiveness of intravenous antibiotics for the treatment of pulmonary exacerbations
31 based on the following comparisons:
 - 32 • Comparison 1. Single IV agents compared
 - 33 ○ Ceftazidime versus Aztreonam (Elborn 1992, Salh 1992)
 - 34 • Comparison 2. Single IV antibiotic (with placebo) versus combination IV antibiotic
 - 35 ○ Tobramycin+placebo versus tobramycin+ceftazidime (Master 2001)
 - 36 ○ Tobramycin+placebo versus piperacillin + tobramycin (Macystis fibrosisarlane 1985)
 - 37 • Comparison 3. Single IV antibiotic versus combination IV antibiotic
 - 38 ○ Piperacillin versus piperacillin+tobramycin (McCarty 1988)
 - 39 ○ Ceftazidime versus tobramycin + ticarcillin (Gold 1985, Wesley 1988)
 - 40 ○ Ceftazidime versus tobramycin + piperacillin (De Boeck 1989)
 - 41 ○ Colistin versus colistin + another anti-pseudomonal antibiotic (Conway 1997)
 - 42 • Comparison 4. Combination IV antibiotics versus combination IV antibiotics
 - 43 ○ Aztreonam+IV amikacin versus IV ceftazidime+IV amikacin (Schaad 1989)
 - 44 ○ IV meropenem + IV tobramycin versus IV ceftazidime + IV tobramycin (Blumer 2005)
 - 45 • Comparison 5. Combination of two IV antibiotics + inhaled antibiotic versus 2 IV antibiotics
46 without inhaled antibiotic

- 1 ○ IV ceftazidime + IV amikacin versus IV ceftazidime + IV amikacin + inhaled amikacin
- 2 (Schaad 1987)
- 3 • Comparison 6. Combination of IV ceftazidime + IV tobramycin versus oral ciprofloxacin
- 4 (Richard 1997)
- 5 • The 2 included studies from the Langton (2014) Cochrane systematic review evaluated the
- 6 effectiveness of antibiotic treatment for acute infection with the first positive isolate of *P*
- 7 *aeruginosa* based on the following comparisons:
- 8 • Comparison 7. Oral ciprofloxacin + inhaled colistin versus inhaled tobramycin (Proesmans
- 9 2013)
- 10 • Comparison 8. Inhaled colistin + oral ciprofloxacin versus inhaled tobramycin + oral
- 11 ciprofloxacin (Taccetti 2012)
- 12 • The size of studies ranged from 19 to 223 participants with cystic fibrosis. Five studies
- 13 included children, young people and adults (Blumer 2005, Macystis fibrosisarlane 1985,
- 14 Schaad 1987, Schaad 1989, Taccetti 2012), 1 study included adults only (Conway 1997),
- 15 3 studies included young people and adults (Elborn 1992, Gold 1985, Salh 1992), 4
- 16 studies included children and young people (McCarty 1988, Proesmans 2013, Richard
- 17 1997, Wesley 1988), 2 studies did not report the age range (De Boeck 1989, Master
- 18 2001).
- 19 • Three studies were conducted in the UK (Conway 1997, Elborn 1992, Salh 1992), 2 in the
- 20 USA (Blumer 2005, McCarty 1988), 2 in Belgium (De Boeck 1989, Proesmans 2013), 1 in
- 21 Canada (Gold 1985), 2 in Australia (Macystis fibrosisarlane 1985, Master 2001), 3 in
- 22 Switzerland (Richard 1997, Schaad 1987, Schaad 1989), 1 in New Zealand (Wesley
- 23 1988), 1 in Italy (Taccetti 2012).

249.4.2.1.2 **S aureus**

25 The interventions that were included in the protocol for the treatment of infection with or

26 exacerbation due to *S aureus* were: Flucloxacillin (oral or IV), Cotrimoxazole (oral or IV),

27 Doxycycline (oral) (not for under 12's) and Cefradrine (oral).

28 One Cochrane systematic review was identified for inclusion (Lo 2015). This review

29 examined interventions for eradicating methicillin-resistant *S aureus*, however no published

30 RCTs were identified.

31 An additional Cochrane systematic review (Southern 2012) on the use of macrolide

32 antibiotics was assessed. The RCTs included in this review have comparisons of placebo

33 and different dosages. As these comparisons are not included in our protocol, no RCTs from

34 this Cochrane review were included.

35 No relevant trials were identified in our search.

369.4.2.1.3 **B cepacia complex**

37 The interventions that were included in the protocol for the treatment of infection with or

38 exacerbation due *B Cepacia* Complex were: Cotrimoxazole (oral or IV), Meropenem (IV or

39 inhaled), Ceftazidime (IV or inhaled), Temocillicin (oral or IV), Imipenem (oral or IV),

40 Trimethoprim (oral or IV) and Tobramycin (oral or IV).

41 Two Cochrane systematic reviews were identified for inclusion. One Cochrane review

42 examined eradication of *B Cepacia* Complex (Regan 2016) and another Cochrane review

43 examined treatment of exacerbations in people with *B Cepacia* Complex (Horsley 2016).

44 Neither of these Cochrane reviews included any RCTs.

45 No relevant trials were identified in our search.

19.4.2.1.4 *H influenza*

2 The interventions that were included in the protocol for the treatment of infection with or
3 exacerbation due to *H influenza* were: Co-amoxiclav (oral or IV), Cefuroxime (IV), Cefaclor,
4 Cefixime, Doxycycline (>12 years), and Macrolide (clarithromycin/azithromycin).

5 No relevant systematic reviews or trials were identified in our search.

69.4.2.1.5 *Nontuberculous mycobacteria*

7 We considered *Mycobacterium Avium* Complex and *Mycobacterium Abscessus*.

8 The interventions that were included in the protocol for the treatment of infection with or
9 exacerbation due to *M. Avium* Complex were: Clarithromycin (oral), Azithromycin (oral),
10 Rifampicin (oral), Ethambutol (oral) and Amikacin (inhaled and potentially IV).

11 The interventions that were included in the protocol for the treatment of infection with or
12 exacerbation due to *M. Abscessus* were: Cefoxitin (IV), Clarithromycin (IV), Amikacin (IV and
13 inhaled), Meropenem (IV and inhaled), Tigecycline, Co-trimoxazole (oral), Moxifloxacin (oral),
14 Ciprofloxacin (oral), Doxycycline/minocycline (tetracyclines) (oral), Linezolid (oral) and
15 Clofazimine (oral).

16 A Cochrane SR was identified in the search (Waters 2016). This review aimed to compare
17 antibiotic treatment *versus* non-antibiotic treatment, or different combinations of antibiotics,
18 for non-tuberculous *mycobacteria* lung infection in people with CYSTIS FIBROSIS. No trials
19 were identified for inclusion in this review.

20 No relevant trials were identified in our search.

219.4.2.1.6 *No identified pathogen*

22 The interventions that were included in the protocol for the treatment of infection with or
23 exacerbation without an identified pathogen at trial entry were any of the above.

24 No relevant systematic reviews or trials were identified in our search.

25 Evidence from these are summarised in the clinical GRADE evidence profile below (Table 65
26 -Table 103). See also the study selection flow chart in Appendix F, forest plots in Appendix I,
27 study evidence tables in Appendix G and exclusion list in Appendix H.

28 9.4.2.2 Summary of included studies

29 A summary of the studies that were included in this review are presented in Table 91 to
30 Table 95.

319.4.2.2.1 *P aeruginosa*

32 **Table 91: Summary of included studies for antimicrobials for pulmonary exacerbations**
33 **with *P aeruginosa***

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Hurley 2015 Cochrane SR	Comparison 1. Single IV agents compared • (Elborn 1992, Salh 1992) Comparison 2. Single IV (with placebo) vs	Participants diagnosed with cystic fibrosis using the Cystic Fibrosis Foundation diagnostic consensus statement, of all ages and all levels of severity. All studies that explicitly aimed to trial	<ul style="list-style-type: none"> Lung function (FEV₁) Time to next exacerbatio n Quality of 	AMSTAR score: 11/11

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>combination of IV antibiotics</p> <ul style="list-style-type: none"> (Macystis fibrosisarlane 1985, Master 2001) <p>Comparison 3. Single IV vs combination of IV antibiotics</p> <ul style="list-style-type: none"> (De Boeck 1989, Conway 1997, Gold 1985, McCarty 1988, Wesley 1988) <p>Comparison 4. Combination of antibiotics vs combination of antibiotics</p> <ul style="list-style-type: none"> (Blumer 2005, Richard 1997, Schaad 1989) 	<p>an IV antibiotic for the treatment of pulmonary exacerbation were considered.</p>	<p>life (CFQ-R)</p> <ul style="list-style-type: none"> Mortality (cystic fibrosis-related and all causes) Adverse events 	
Primary studies included in the Cochrane SR				
Blumer 2005 (USA) RCT	<p>Intervention 1: IV meropenem 40 mg/kg up to a maximum dose of 2 g and IV tobramycin(given for a mean of 13.5 days)</p> <p>Intervention 2: IV ceftazidime 50 mg/kg up to a maximum dose of 2 g and I V tobramycin(given for a mean of 14.1 days)</p> <ul style="list-style-type: none"> Tobramycin dose adjusted to give a peak serum concentration of ≥ 8 $\mu\text{g/mL}$ and trough concentration of < 2 $\mu\text{g/mL}$ 	<p>N=121 participants with a recent (usually < 1 month) culture of <i>P aeruginosa</i> or <i>B cepacia</i> complex recruited at a protocol-de fined exacerbation.</p> <ul style="list-style-type: none"> Age: ≥ 5 years of age 	<ul style="list-style-type: none"> Lung function (FEV₁) 	Included in Cochrane SR Hurley 2015
Conway 1997 (UK) RCT	<p>Intervention 1: IV colistin (2 MU 3x daily).</p> <p>Intervention 2: IV colistin (2 MU 3x daily) and a second anti-pseudomonal antibiotic</p>	<p>N=71 adults with cystic fibrosis and chronic <i>P aeruginosa</i> experiencing a protocol-defined exacerbation.</p> <ul style="list-style-type: none"> Mean age (SD): 21 (4.2) years. 	<ul style="list-style-type: none"> FEV₁ Mortality Adverse effects 	Included in Cochrane SR Hurley 2015
De Boeck 1989 (Belgium)	<p>Intervention 1: IV ceftazidime 50 mg/kg 3x daily.</p>	<p>N=21 participants with cystic fibrosis and a protocol-defined pulmonary exacerbation, chronically infected with</p>	<ul style="list-style-type: none"> FEV₁ Time to readmission Mortality 	Included in Cochrane SR Hurley 2015

Study	Intervention/Comparison	Population	Outcomes	Comments
RCT	Intervention 2: IV piperacillin 75 mg/kg 4x daily and IV tobramycin 10 mg/kg/day in 3 doses	<i>P aeruginosa</i> that was sensitive to piperacillin, tobramycin and ceftazidime • Mean age 14.8 years		
Elborn 1992 (UK) RCT	Intervention 1: IV ceftazidime 2 g 3x daily. Intervention 2: IV aztreonam 2 g 3x daily.	N=24 participants with cystic fibrosis and chronic <i>P aeruginosa</i> infection experiencing exacerbations. Mean (range) age: 20 (14 to 48) years	• FEV ₁	Included in Cochrane SR Hurley 2015
Gold 1985 (Canada) RCT	Intervention 1: IV ceftazidime 200 mg/kg/day in 4 doses. Intervention 2: IV ticarcillin 300 mg/kg/day in 4 doses and IV tobramycin 10 mg/kg/day in 3 doses	N=30 participants with cystic fibrosis and <i>P aeruginosa</i> infection present at the previous clinic visit, experiencing an acute respiratory exacerbation. • Age >12 years. Mean age (SD): 18.9 (1.1) in group 1; 17.8 (0.8) in group 2	• FEV ₁ • Adverse effects	Included in Cochrane SR Hurley 2015
Macystis fibrosis arlane 1985 (Australia) RCT	Intervention 1: IV piperacillin 50 mg/kg 4-hourly. Intervention 2: IV placebo 5% dextrose 4-hourly Intervention 3: IV piperacillin 100 mg/kg 8-hourly. Intervention 4: IV placebo 5% dextrose 8-hourly. All participants received IV tobramycin 2.5 mg/kg 3x daily, oral flucloxacillin 25 mg/kg/day in 4 doses and oral probenecid (suggested to increase antibiotic concentrations) 250 - 500 mg 3x daily Duration: 14 days.	N=19 participants aged over 8 years with cystic fibrosis with <i>P aeruginosa</i> in sputum admitted to hospital for worsening respiratory status. • Mean age: 13.7 to 15.6 years	• FEV ₁ • Adverse effects	Included in Cochrane SR Hurley 2015 Pseudomonas was not eradicated from the sputum in any of the patients
Master 2001 (Australia) RCT	Intervention 1: IV ceftazidime 50 mg/kg/dose 3x daily and IV tobramycin 3 mg/kg/dose 3x daily	N=51 participants with cystic fibrosis experiencing a protocol-defined exacerbation with <i>P aeruginosa</i>	• FEV ₁ • Adverse effects	Included in Cochrane SR Hurley 2015

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>Intervention 2: IV tobramycin 9 mg/kg/day 1x daily.</p> <p>Duration: at least 10 days.</p>	<p>isolated from sputum. Participants with an FVC lower than 40% predicted were excluded.</p> <ul style="list-style-type: none"> • Mean age (SD): 16 (7) years in group 1; 14 (5) years in group 2 		
McCarty 1988 (USA) RCT	<p>Intervention 1: IV piperacillin 600 mg/kg/day (regimen not detailed)</p> <p>Intervention 2: IV piperacillin 600 mg/kg/day and tobramycin 8 - 10 mg/kg/day (regimen not detailed)</p> <p>Duration: at least 10 days.</p>	<p>N=17 children with cystic fibrosis admitted for treatment of pulmonary exacerbations with <i>P aeruginosa</i>.</p> <ul style="list-style-type: none"> • Age range: 2 to 12 years. 	<ul style="list-style-type: none"> • FEV₁ • Eradication of pseudomonas • Adverse effects 	Included in Cochrane SR Hurley 2015
Richard 1997 (Switzerland) RCT	<p>Intervention 1: oral ciprofloxacin 15 mg/kg 2x daily.</p> <p>Intervention 2: IV ceftazidime 50 mg/kg 3x daily and IV tobramycin 3 mg/kg 3x daily</p> <p>Duration: 14 days.</p>	<p>N=108 children with cystic fibrosis and <i>P aeruginosa</i> infection and experiencing a protocol-defined pulmonary exacerbation with <i>P aeruginosa</i>.</p> <ul style="list-style-type: none"> • Mean age: 10.2 years in group 1; 11 years in group 2. Age range: 5 to <17. 	<ul style="list-style-type: none"> • Eradication of pseudomonas • Adverse effects 	Included in Cochrane SR Hurley 2015
Salh 1992 (UK) RCT	<p>Intervention 1; IV aztreonam 8 g/day in 4 doses.</p> <p>Intervention 2: IV ceftazidime 8 g/day in 4 doses.</p> <p>Duration: 2 weeks</p>	<p>N=22 participants with cystic fibrosis and <i>P aeruginosa</i> sensitive to the study drugs who were admitted to hospital due to an infective exacerbation</p> <ul style="list-style-type: none"> • Age range: 16 to 32 years. 	<ul style="list-style-type: none"> • FEV₁ 	Included in Cochrane SR Hurley 2015
Schaad 1987 (Switzerland) RCT	<p>Intervention 1: IV ceftazidime 250 mg/kg/day in 4 doses and IV amikacin 33 mg/kg/day in 3 doses</p> <p>Intervention 2: IV ceftazidime 250 mg/kg/day in 4 doses and IV amikacin 33 mg/kg/day in 3 doses and nebulised amikacin 100 mg 2x</p>	<p>N=62 participants with cystic fibrosis admitted with an acute pulmonary exacerbation who had <i>P aeruginosa</i> isolated on admission. Those who had been admitted to hospital in the recent 6 months were excluded</p> <ul style="list-style-type: none"> • (n=87 courses of therapy by random 	<ul style="list-style-type: none"> • Adverse effects 	Included in Cochrane SR Hurley 2015

Study	Intervention/Comparison	Population	Outcomes	Comments
	daily Duration: 15 days	assignment) • Age range: 3 to 24 years.		
Schaad 1989 (Switzerland) RCT	Intervention 1: IV aztreonam 300 mg/kg/day in 4 doses and IV amikacin 36 mg/kg/day in 3 doses Intervention 2: IV ceftazidime 300 mg/kg/day in 4 doses and IV amikacin 36 mg/kg/day in 3 doses for 2 weeks followed by oral ciprofloxacin 30 mg/kg/day for 4 weeks Duration: 2 weeks IV treatment, with oral treatment extended for a further 4 weeks in 1 group	N=42 participants with cystic fibrosis admitted with a protocol-defined pulmonary exacerbation and <i>P aeruginosa</i> isolated at admission. Those who had been admitted to hospital in previous 4 months were excluded • (n=56 treatment courses by random assignment) • Mean age (SD): 15.4 (6) years (range 2.3 to 25.4 years).	• Eradication of <i>Pseudomonas</i> • FEV ₁ • Adverse effects	Included in Cochrane SR Hurley 2015
Wesley 1988 (New Zealand) RCT	Intervention 1: IV ceftazidime 150 mg/kg/day (regimen not detailed) Intervention 2: IV tobramycin 7.5 mg/kg/day and IV ticarcillin 300 mg/kg/day (regimen not detailed) Duration: 14 days	N=13 children with cystic fibrosis and severe chest disease with pseudomonas chest exacerbation. • Age range: 9 to 15 years.	• Number of admission requiring iv antibiotics or death • Adverse effects	Included in Cochrane SR Hurley 2015 Conference abstract

1 FEV₁: Forced Expiratory volume. Information provided in the table is adapted from Hurley 2015.

2 **Table 92: Summary of included studies for antimicrobials for the treatment of acute**
3 **infection with *P aeruginosa***

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Langton 2014 Cochrane SR	Comparisons of combinations of oral, inhaled or intravenous antibiotics (Proesman 2013, Tacetti 2012)	Children and adults with confirmed cystic fibrosis with a first ever positive microbiological isolate of <i>P aeruginosa</i> from a respiratory tract specimen.	• FEV ₁ • Trial discontinuation due to lack of compliance (proxy outcome for treatment failure) • Adverse effects	AMSTAR score: 10/11
Primary studies included in the Cochrane systematic reviews				
Proesmans	Intervention	N=58 children with	• Adverse effects	Included in

Study	Intervention/Comparison	Population	Outcomes	Comments
2013 (Belgium) RCT	Inhaled TSI (300 mg 2x daily for 28 days), 3 months Control Combination therapy with inhaled colistin (2 MU 2x daily)+ oral ciprofloxacin (10 mg/kg 3x daily), 3 months	cystic fibrosis, all with new isolation of <i>P aeruginosa</i> (sputum or cough swabs) <ul style="list-style-type: none"> • Treatment: n=29 • Control: n=29 • Age: median age 9 years, interquartile range (4.7 - 13.1 years) 		Langton 2014 Cochrane SR This study did not report the outcome 'eradication of pathogen' but reported 'number of positive respiratory culture for pseudomonas. There was no difference between groups in positive pseudomonas culture 6 months and 24 months follow-up
Taccetti 2012 (Italy) RCT	Group A 28 days therapy 2 x daily inhalation of 2 MU colistin with 2 x daily doses of ciprofloxacin 15 mg/kg/dose. Group B 28 days therapy TSI (300 mg 2 x daily) with 2 x daily doses of ciprofloxacin 15 mg/kg/dose	N=223 participants with cystic fibrosis with first ever or new <i>P aeruginosa</i> infection. New infection defined as <i>P aeruginosa</i> isolation following bacterial clearance documented by 3 negative cultures within the previous 6 months <ul style="list-style-type: none"> • Group A: n=105 (52 male and 53 female) • Group B: n=118 (64 male and 54 female) • Age: over 1 year. 	<ul style="list-style-type: none"> • FEV₁ • Trial discontinuation due to lack of compliance (proxy outcome for treatment failure) 	Included in Langton 2014 Cochrane SR This study did not report the outcome 'eradication of pathogen' but reported 'number of positive respiratory culture for pseudomonas'. No difference was found in positive respiratory cultures of pseudomonas at end

Study	Intervention/Comparison	Population	Outcomes	Comments
				of treatment between both regimes. An increased incidence of the emerging pathogen <i>S.maltophilia</i> was reported following treatment,

19.4.2.2.2 *S aureus*

2 **Table 93: Summary of included studies for the antimicrobial treatment of infection with**
3 **or exacerbation due to *S aureus***

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Lo 2015 Cochrane SR	Intervention Any combination of topical, inhaled, oral or intravenous antibiotics with the primary aim of eradicating MRSA Comparison • Placebo • Standard treatment • No treatment	Children and adults with confirmed diagnosis of cystic fibrosis with a confirmed positive microbiological isolate of MRSA on clinically relevant cystic fibrosis respiratory cultures.	No studies were identified for inclusion in this review	AMSTAR score:10/11

4 *CYSTIS FIBROSIS: cystis fibrosis; SR: systematic review; MRSA: methicillin-resistant S aureus*

59.4.2.2.3 *B cepacia complex*

6 **Table 94: Summary of included studies for the antimicrobial treatment of infection with**
7 **or exacerbation due *B Cepacia Complex (BCC)***

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Horsley 2016 Cochrane SR	Intervention Any antibiotic treatment regimen for treating an exacerbation of CF lung disease Comparison • Placebo • Different antibiotic	Patients with cystic fibrosis of any age with evidence of pulmonary infection with organisms of the BCC (defined as at least 2 positive sputum or bronchoalveolar lavage specimens within the last 6 months)	No studies were identified for inclusion in this review	AMSTAR score:11/11

Study	Intervention/Comparison	Population	Outcomes	Comments
	regimen			
Regan 2016 Cochrane SR	<p>Intervention Any antibiotic or antibiotic adjuvant therapy used alone or in combination to eradicate BCC infection</p> <p>Comparison</p> <ul style="list-style-type: none"> • Placebo • No treatment • Alternative antimicrobial agent (excluding the participant's usual therapeutic regimen) 	<p>Any person of any age with a confirmed clinical diagnosis of cystic fibrosis who acquires a new infection or a re-infection with BCC. People with all disease severity were included.</p>	No studies were identified for inclusion in this review	AMSTAR score: 10/11

1 BCC: *B Cepacia Complex*; CYSTIS FIBROSIS: *cystis fibrosis*; SR: *systematic review*

29.4.2.2.4 Non-tuberculous mycobacteria

3 **Table 95: Summary of included studies for the antimicrobial treatment of infection with**
4 **or exacerbation due to non-tuberculous mycobacteria (NTM)**

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Waters 2016 Cochrane SR	<p>Intervention Antibiotics to treat NTM pulmonary infection</p> <p>Comparison No antibiotic treatment Different NTM antibiotic regimen</p>	<p>Children and adults with a confirmed clinical diagnosis of cystic fibrosis, who have NMT pulmonary infection. People with all disease severity were included.</p>	No studies were identified for inclusion in this review	AMSTAR score: 10/11 Last search September 2016

59.4.2.2.5 Non-identified pathogen

6 No studies were identified for inclusion.

7 9.4.2.3 Clinical evidence profile

8 The clinical evidence profiles for the review question addressing antimicrobials for pulmonary
9 exacerbation are presented in Table 65 - Table 101. The clinical evidence profiles for the
10 review question addressing antimicrobials for acute exacerbations are presented in Table
11 102 and Table 103.

129.4.2.3.1 *P aeruginosa*

13 Antimicrobials for pulmonary exacerbations due to *P aeruginosa*

14 **Table 96: Summary clinical evidence profile: Comparison 1. Single IV agents compared**
15 **for pulmonary exacerbations with *P aeruginosa***

Comparison 1. Single IV agents compared for pulmonary exacerbations with *P aeruginosa*

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Single IV agents	Single IV agents				
[ceftazidime versus aztreonam] FEV ₁ (absolute change) litres Follow-up: 2 weeks	The mean absolute change in FEV ₁ litres in the IV aztreonam group ranged between 0.27 and 0.54	The mean absolute change in FEV ₁ litres in the IV ceftazidime group was 0.06 lower (0.44 lower to 0.32 higher)		46 (Elborn 1992, Salh 1992)	⊕⊕⊕⊕ low ^{1,2}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference

- 1 The quality of the evidence was downgraded by 1 as 4 participants received both drugs in Salh 1992 study,
 2 The quality of the evidence was downgraded by 1 due to serious heterogeneity (chi-squared p<0.1, I-squared inconsistency statistic of 50%-74.99%)

4 **Table 97: Comparison 2. Single IV antibiotic (with placebo) versus combination IV**
 5 **antibiotic for pulmonary exacerbations with *P aeruginosa***

Comparison 2. Single IV antibiotic (with placebo) versus combination IV antibiotic for pulmonary exacerbations with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Combination IV antibiotics	Single IV antibiotic (with placebo)				
[tobramycin + placebo versus tobramycin + ceftazidime] FEV ₁ % predicted (absolute change) - Follow-up: 10 days	The mean FEV ₁ % predicted (absolute change) in the tobramycin + ceftazidime group was 12.8	The mean FEV ₁ % predicted (absolute change) in the tobramycin + placebo groups was 2.2 lower (6.63 lower to 2.23 higher)		98 (Master 2001)	⊕⊕⊕⊕ low ^{1,2}	
[tobramycin + placebo versus piperacillin + tobramycin] FEV ₁ % predicted (relative change) - Follow-up: 2	The mean FEV ₁ % predicted (relative change) in the piperacillin + tobramycin group was	The mean FEV ₁ % predicted (relative change) in the tobramycin + placebo groups was 4.2 lower (26.5 lower to 18.1 higher)		9 (Macystis fibrosisarlane 1985)	⊕⊕⊕⊕ very low ^{3,4}	

weeks	12.2					
[tobramycin + placebo <i>versus</i> piperacillin + tobramycin] FEV ₁ % predicted (relative change) - Follow-up: 2 weeks	The mean FEV ₁ % predicted (relative change) in the piperacillin + tobramycin group was 1.8	The mean FEV ₁ % predicted (relative change) - in the tobramycin + placebo groups was 7.95 higher (8.78 lower to 24.68 higher)		9 (Macystis fibrosisarlane 1985)	⊕⊕⊕⊕ very low ^{3,4}	
[tobramycin + placebo <i>versus</i> piperacillin all regimens] Adverse effects - sensitivity reaction - number of participants Follow-up: 2 weeks	300 per 1000	51 per 1000 (3 to 888)	RR 0.17 (0.01 to 2.96)	18 (Macystis fibrosisarlane 1985)	⊕⊕⊕⊕ low ^{3,5}	
[tobramycin + placebo <i>versus</i> tobramycin + ceftazidime] Adverse effects - Number of hospital admissions due to Tinnitus Follow-up: 2 weeks	39 per 1000	43 per 1000 (6 to 290)	RR 1.09 (0.16 to 7.4)	98 (Master 2001)	⊕⊕⊕⊕ very low ^{1,6}	
[tobramycin + placebo <i>versus</i> tobramycin + ceftazidime] Adverse effects - serum - Creatinine Follow-up: 2 weeks	The mean serum creatinine in the tobramycin + ceftazidime groups was 0	The mean serum creatinine in the Tobramycin + placebo groups was 4 lower (9.38 lower to 1.38 higher)		44 (Master 2001)	⊕⊕⊕⊕ very low ^{1,6}	
[tobramycin + placebo <i>versus</i> tobramycin + ceftazidime] Adverse effects - serum - NAG Follow-up: 2 weeks	The mean serum NAG in the tobramycin + ceftazidime groups was	The mean serum NAG in the Tobramycin + placebo groups was 2.1 lower (3.46 lower to 0.74 lower)		44 (Master 2001)	⊕⊕⊕⊕ moderate ¹	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference; NAG: N-acetyl glucosamide; RR: risk ratio</p>						

1 The quality of the evidence was downgraded by 1 as each participant contributed to multiple treatment episodes.
2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID

- 1 3 The quality of the evidence was downgraded by 1 due to attrition bias (2 participants withdrew and did not
2 contribute to analysis) and 1 participant received 2 treatment courses.
3 4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
4 5 The quality of the evidence was downgraded by 1 due to very serious imprecision as 95%CI crossed 1 default
5 MIDs
6 6 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs

7 **Table 98: Comparison 3. Single IV antibiotic versus combination IV antibiotic for**
8 **pulmonary exacerbations with *P aeruginosa***

Comparison 3. Single IV antibiotic versus combination IV antibiotic for pulmonary exacerbations with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Combination IV antibiotic	Single IV antibiotic				
[piperacillin versus piperacillin + tobramycin] Eradication: number people in whom pseudomonas isolates were eradicated at end of course - Follow-up: 10 days	632 per 1000	265 per 1000 (114 to 600)	RR 0.42 (0.18 to 0.95)	38 (McCarty 1988)	⊕⊕⊕⊕ low ^{1,2}	
[ceftazidime versus IV tobramycin + ticarcillin] FEV ₁ (relative change) - % Follow-up: 10 to 14 days	The mean FEV ₁ (relative change) in the tobramycin + ticarcillin groups was 33.3	The mean FEV ₁ (relative change) in the ceftazidime groups was 19.6 lower (38.26 to 0.94 lower)		30 (Gold 1985)	⊕⊕⊕⊕ low ^{3,4}	
[colistin versus colistin + "other"] FEV ₁ (absolute change) ml Follow-up: 12 days	The mean FEV ₁ (absolute change) in the colistin + "other" groups was 300	The mean FEV ₁ (absolute change) in the colistin groups was 160 lower (309.72 to 10.28 lower)		71 (Conway 1997)	⊕⊕⊕⊕ low ⁵	
[ceftazidime versus tobramycin + piperacillin] FEV ₁ % predicted (absolute change) Follow-up: 12 days	The mean FEV ₁ % predicted (absolute change) in the ceftazidime groups was 1 higher (8.85 lower to 10.85 higher)	The mean FEV ₁ % predicted (absolute change) in the ceftazidime groups was 1 higher (8.85 lower to 10.85 higher)		21 (De Boeck 1989)	⊕⊕⊕⊕ very low ^{3,6}	
[ceftazidime	The mean time	The mean time		19	⊕⊕⊕⊕	

Comparison 3. Single IV antibiotic versus combination IV antibiotic for pulmonary exacerbations with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
<i>versus</i> tobramycin + piperacillin Time to readmission - months Follow-up: 24 to 26 months	to readmission in the tobramycin + piperacillin groups was 9 months	to readmission in the piperacillin groups was 1 month lower (5.52 lower to 3.52 higher)		(De Boeck 1989)	very low ^{3,7}	
[ceftazidime <i>versus</i> tobramycin + ticarcillin Number of admissions, requiring IV antibiotics or death - Follow-up: 3 months	500 per 1000	585 per 1000 (265 to 1000)	RR 1.17 (0.53 to 2.55)	22 (Wesley 1988)	⊕⊕⊕⊕ very low ^{7,8}	
[ceftazidime <i>versus</i> tobramycin + piperacillin Mortality - Follow-up: 4 months	91 per 1000	100 per 1000 (7 to 1000)	RR 1.1 (0.08 to 15.36)	21 (De Boeck 1989)	⊕⊕⊕⊕ low ^{9,10}	
[IV colistin <i>versus</i> IV colistin + "other"] Mortality – Follow-up: 12 weeks	29 per 1000	9 per 1000 (0 to 220)	RR 0.32 (0.01 to 7.7)	71 (Conway 1997)	⊕⊕⊕⊕ low ^{5,10}	
[ceftazidime <i>versus</i> IV tobramycin + ticarcillin Adverse effects: liver transaminase enzyme elevation Follow-up: 10-14 days	87 per 1000	133 per 1000 (29 to 618)	RR 1.53 (0.33 to 7.11)	52 (Gold 1987 and Wesley 1988)a,b	⊕⊕⊕⊕ very low ^{7,11}	
[colistin <i>versus</i> combination anti-pseudomonal antibiotics] Adverse effects: neurological	1000 per 1000	940 per 1000 (860 to 1000)	RR 0.94 (0.86 to 1.04)	17 (McCarty 1988)	⊕⊕⊕⊕ low ⁵	

Comparison 3. Single IV antibiotic versus combination IV antibiotic for pulmonary exacerbations with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
adverse effects Follow-up: 12 days						
[piperacillin <i>versus</i> piperacillin + tobramycin] Adverse effects: rash Follow-up: 10 days	111 per 1000	41 per 1000 (2 to 888)	(McCarthy 1988)	(McCarthy 1988)	⊕⊕⊕⊕ very low ^{1,7}	
[piperacillin <i>versus</i> piperacillin + tobramycin] Adverse effects -fever Follow-up: 10 days	111 per 1000	124 per 1000 (9 to 1000)	RR 1.12 (0.08 to 15.19)	17 (McCarthy 1988)	⊕⊕⊕⊕ very low ^{1,7}	
[ceftazidime <i>versus</i> tobramycin + ticarcillin] Adverse effects – proteinuria Follow-up: 10 - 14 days	59 per 1000	59 per 1000 (4 to 866)	RR 1 (0.07 to 14.72)	34 (Gold 1985)a	⊕⊕⊕⊕ very low ^{3,7}	
[colistin <i>versus</i> combination anti-pseudomonal antibiotics] Adverse effects - renal toxicity - Change in blood urea (mmol/l) Follow-up: 12 days	The mean renal toxicity - change in blood urea (mmol/l) in the combination anti-pseudomonal AB groups was 0.83	The mean renal toxicity - change in blood urea (mmol/l) in the colistin groups was 0.26 lower (0.93 lower to 0.41 higher)		71 (Conway 1997)	⊕⊕⊕⊕ very low ^{5,12}	
[colistin <i>versus</i> combination anti-pseudomonal antibiotics] Adverse effects: renal toxicity - Change in serum creatinine (mol/l) Follow-up: 12	The mean change in serum creatinine (mol/l) in the combination anti-pseudomonal AB groups was -5.85	The mean change in serum creatinine (mol/l) in the colistin groups was 8.85 higher (0.66 lower to 18.36 higher)		71 (Conway 1997)	⊕⊕⊕⊕ very low ^{5,7}	

Comparison 3. Single IV antibiotic versus combination IV antibiotic for pulmonary exacerbations with *P aeruginosa*

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference; mmol/l: millimoles per litre; RR: risk ratio

a Gold 1985: total of 34 treatment observations in N=30

b Wesley 1988: total of 23 observations in N=13

1 The quality of the evidence was downgraded by 2 due to no blinding and 3 participants were included twice in analysis

2 Minimal important difference for this outcome (MID) = any difference is clinically significant

3 The quality of the evidence was downgraded by 1 due to no blinding.

4 The quality of the evidence was downgraded by 1 as 95% CI crossed 1 clinical MID

5 The quality of the evidence was downgraded by 2 due to single blinding and 18 participants were enrolled twice.

6 The quality of the evidence was downgraded by 2 due as 95%CI crossed 2 clinical MIDs.

7 The quality of the evidence was downgraded by 2 as 95% CI crossed 2 default MIDs

8 The quality of the evidence was downgraded by 1 as 13 participants received 23 courses of treatment.

9 The quality of the evidence was downgraded by 1 due to multiple enrolment of participants (40 participants contribute to 46 treatment episodes).

10 The quality of the evidence was downgraded by 1, as the 95% CI crossed the null effect (mortality could either decrease or increase)

11 The quality of the evidence was downgraded by 1 due lack of blinding in 1 trial, and because some participants were enrolled twice

12 The quality of the evidence was downgraded by 1 as 95% CI crossed 1 default MID

Table 99: Comparison 4. Combination IV antibiotics versus combination IV antibiotics for pulmonary exacerbations with *P aeruginosa*

Comparison 4. Combination IV antibiotics versus combination IV antibiotics for pulmonary exacerbations with *P aeruginosa*

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Combination IV	Combination IV				
[aztreonam + amikacin versus ceftazidime + amikacin] Eradication of pathogen Follow-up: 2 weeks	571 per 1000	606 per 1000 (394 to 943)	RR 1.06 (0.69 to 1.65)	56 a (Schaad 1989)	⊕⊕⊕ ⊖ very low ^{1,2}	
[aztreonam + amikacin versus ceftazidime + amikacin] FEV ₁ % predicted (absolute change) - (combination B) Follow-up: 2 weeks	The mean FEV ₁ % predicted (absolute change) in the versus IV ceftazidime + IV amikacin	The mean FEV ₁ % predicted (absolute change) in the Aztreonam + IV amikacin groups was		49a (Schaad 1989)	⊕⊕⊕ ⊖ low ^{1,3}	

Comparison 4. Combination IV antibiotics versus combination IV antibiotics for pulmonary exacerbations with P aeruginosa						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	groups was 9	4 higher (0.25 lower to 8.25 higher)				
[meropenem + tobramycin <i>versus</i> ceftazidime + tobramycin] FEV ₁ % predicted (absolute change) - Follow-up: 2 to 4 weeks ^b	The mean FEV ₁ % predicted (absolute change) in the <i>versus</i> IV ceftazidime + IV tobramycin groups was 11.1	The mean FEV ₁ % predicted (absolute change) in the IV meropenem + IV tobramycin groups was 2.7 higher (0.76 lower to 6.16 higher)		97 (Blumer 2005)	⊕⊕⊕ ⊖ low ^{3,4}	
[meropenem + tobramycin <i>versus</i> ceftazidime + tobramycin] FEV ₁ % predicted (relative % change) - Follow-up: 2 to 4 weeks ^b	The mean FEV ₁ % predicted (relative % change) in the <i>versus</i> IV ceftazidime + IV tobramycin groups was 29.4	The mean FEV ₁ % predicted (relative % change) in the IV meropenem + IV tobramycin groups was 9.4 higher (8.44 lower to 27.24 higher)		97 (Blumer 2005)	⊕⊕⊕ ⊖ very low ^{4,5}	
[aztreonam + amikacin <i>versus</i> ceftazidime + amikacin] Adverse effects - Rash Follow-up: 2 weeks	71 per 1000	14 per 1000 (1 to 285)	RR 0.2 (0.01 to 3.99)	56a (Schaad 1989)	⊕⊕⊕ ⊖ very low ^{1,6}	
[aztreonam + amikacin <i>versus</i> ceftazidime + amikacin] Adverse effects - Liver transaminases - AST + ALT Follow-up: 2 weeks	71 per 1000	143 per 1000 (29 to 718)	RR 2 (0.4 to 10.05)	56a (Schaad 1989)	⊕⊕⊕ ⊖ very low ^{1,6}	
[aztreonam + amikacin <i>versus</i> ceftazidime + amikacin] Adverse effects - Thrombocytopenia Follow-up: 2 weeks	0 per 1000	0 per 1000 (0 to 0)	RR 7 (0.38 to 129.55)	56a (Schaad 1989)	⊕⊕⊕ ⊖ very low ^{1,6}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in

Comparison 4. Combination IV antibiotics versus combination IV antibiotics for pulmonary exacerbations with *P aeruginosa*

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				

footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: AST: aminotransferase, ALT: alanine aminotransferase; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IV: intravenous; MD: mean difference; RR: risk ratio

a total of 56 treatment courses were randomised, N=42 participants

b 2 to 4 weeks after discontinuation of 2 week course.

1 The quality of the evidence was downgraded by 1 due to attrition bias (clinical outcomes available for only around 50% of participants).

2 The quality of the evidence was downgraded by 2, as the 95% CI crossed the null effect and the CI was very wide

3 The quality of the evidence was downgraded by 1 as 95%CI crossed 1 clinical MID.

4 The quality of the evidence was downgraded by 1 due to attrition bias (some data missing).

5 The quality of the evidence was downgraded by 2 as 95%CI crossed 2 clinical MIDs.

6 The quality of the evidence was downgraded by 2 as 95%CI crossed 2 default MIDs.

Table 100: Comparison 5. Two IV antibiotics + inhaled antibiotic versus 2 IV without inhaled antibiotic for pulmonary exacerbations with *P aeruginosa*

Comparison 5. Two IV antibiotics + inhaled antibiotic versus 2 IV without inhaled antibiotic for pulmonary exacerbations with *P aeruginosa*

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	2 IV without inhaled antibiotic	2 IV antibiotic + inhaled antibiotic				
[IV ceftazidime + IV amikacin + inhaled amikacin <i>versus</i> IV ceftazidime + IV amikacin] Eradication of <i>P aeruginosa</i> Follow-up: 15 days	409 per 1000	749 per 1000 (503 to 1000)	RR 1.83 (1.23 to 2.73)	84 (Schaad 1987)	⊕⊕⊕⊖ moderate ¹	
[IV ceftazidime + IV amikacin + inhaled amikacin <i>versus</i> IV ceftazidime + IV amikacin] Adverse effects: raised liver transaminases Follow-up: 4 to 6 weeks	250 per 1000	168 per 1000 (58 to 480)	RR 0.67 (0.23 to 1.92)	54 (Schaad 1987)	⊕⊖⊖⊖ very low ^{1,2}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio;

1 The quality of the evidence was downgraded by 1 as 18 participants were recruited twice and 6 participants enrolled 3 times.
2 The quality of the evidence was downgraded by 2 due to serious imprecision as 95% CI crosses 2 default MID.

Table 101: Comparison 6. IV ceftazidime + IV tobramycin versus oral ciprofloxacin for pulmonary exacerbations with *P aeruginosa*

Comparison 6. IV ceftazidime + IV tobramycin versus oral ciprofloxacin for pulmonary exacerbations with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Oral ciprofloxacin	IV ceftazidime + IV tobramycin				
Eradication of <i>P aeruginosa</i> Follow-up: 2 weeks	245 per 1000	624 per 1000 (365 to 1000)	RR 2.55 (1.49 to 4.39)	89 (Richard 1997)	⊕⊕⊕⊖ moderate ¹	
Adverse effects - Treatment-related events Follow-up: 2 weeks	164 per 1000	188 per 1000 (83 to 427)	RR 1.15 (0.51 to 2.61)	108 (Richard 1997)	⊕⊖⊖⊖ very low ^{1,2}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; IV: intravenous; RR: risk ratio						

1 The quality of the evidence was downgraded by 1 due to no blinding.
2 The quality of the evidence was downgraded by 2 as 95% CI crossed 2 default MID.

Antimicrobials for acute infection with *P aeruginosa*

Table 102: Comparison 7. Oral ciprofloxacin + inhaled colistin versus inhaled tobramycin for acute infection with *P aeruginosa*

Comparison 7. Oral ciprofloxacin + inhaled colistin versus inhaled tobramycin for acute infection with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Inhaled tobramycin	Oral ciprofloxacin + inhaled colistin				
Adverse events - Severe cough Follow-up: 3 months	34 per 1000	11 per 1000 (0 to 271)	RR 0.33 (0.01 to 7.86)	58 (Proesmans 2013)	⊕⊖⊖⊖ very low ^{1,2}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; RR: risk ratio						

1 The quality of the evidence was downgraded by 1 due to no blinding. Blinding was not possible due to route of administration (oral versus inhaled).
2 The quality of the evidence was downgraded by 2 due to very serious imprecision as 95% CI crossed 2 default MIDs.

Table 103: Comparison 8. Inhaled colistin + oral ciprofloxacin versus inhaled tobramycin + oral ciprofloxacin for acute infection with *P aeruginosa*

Comparison 8. Inhaled colistin + oral ciprofloxacin versus inhaled tobramycin + oral ciprofloxacin for acute infection with <i>P aeruginosa</i>						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Inhaled tobramycin + oral ciprofloxacin	Inhaled colistin + oral ciprofloxacin				
Relative change in % predicted FEV ₁ from baseline Follow-up: 54 days	The mean relative change in % predicted FEV ₁ from baseline in the control groups was 2.15	The mean relative change in % predicted FEV ₁ from baseline in the intervention groups was 2.4 lower (5.885 lower to 1.0855 higher)		128 (Taccetti 2012)	⊕⊕⊕ ⊖ very low ^{1,2}	
Treatment failure - trial discontinuation due to Lack of compliance Follow-up: 28 days	110 per 1000	105 per 1000 (50 to 224)	RR 0.95 (0.45 to 2.03)	223 (Taccetti 2012)	⊕⊕⊕ ⊖ very low ^{1,3,4}	
Adverse events - Vomiting Follow-up: 28 days	17 per 1000	9 per 1000 (1 to 104)	RR 0.56 (0.05 to 6.11)	223 (Taccetti 2012)	⊕⊕⊕ ⊖ very low ^{1,5}	
Adverse events - Photosensitivity Follow-up: 28 days	0 per 1000	0 per 1000 (0 to 0)	RR 3.37 (0.14 to 81.79)	223 (Taccetti 2012)	⊕⊕⊕ ⊖ very low ^{1,5}	
Adverse events - Wheeze Follow-up: 28 days	8 per 1000	3 per 1000 (0 to 77)	RR 0.37 (0.02 to 9.09)	223 (Taccetti 2012)	⊕⊕⊕ ⊖ very low ^{1,5}	
Adverse events leading to trial discontinuation - Pulmonary exacerbation during early	42 per 1000	38 per 1000 (11 to 138)	RR 0.9 (0.25 to 3.26)	223 (1 study)	⊕⊕⊕ ⊖ very low ^{1,4}	

eradication treatment leading to treatment failure Follow-up: 28 days						
--	--	--	--	--	--	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; IV: intravenous; RR: risk ratio

- 1 1 The quality of the evidence was downgraded by 1 due to serious imprecision as there was no blinding (open-label).
2
3 2 The quality of the evidence was downgraded by 2 due to serious imprecision as 95% CI crosses two clinical
4 MIDs.
5 3 The quality of the evidence was downgraded due to indirect outcome for discontinuation due to adverse events.
6 It is unclear if discontinuation is due to adverse events or other factors.
7 4 The quality of the evidence was downgraded by 2, as the 95% CI crossed the null effect and the CI was very
8 wide
9 5 The quality of the evidence was downgraded by 2 due to serious imprecision as 95% CI crosses 2 default MIDs.

109.4.2.3.2 **S aureus**

11 No studies were identified for inclusion.

129.4.2.3.3 **B cepacia complex**

13 No studies were identified for inclusion.

149.4.2.3.4 **Non-tuberculous mycobacteria**

15 No studies were identified for inclusion.

169.4.2.3.5 **Non-identified pathogen**

17 No studies were identified for inclusion.

18 **9.4.2.4 Economic evidence**

19 No economic evaluations of interventions relevant to acute antimicrobial treatment were
20 identified in the literature search conducted for this guideline. Full details of the search and
21 economic article selection flow chart can be found in Appendix E and F, respectively.

22 This review question was not prioritised for de novo economic modelling. Instead additional
23 economic analysis has been undertaken on chronic antimicrobial treatment as this was
24 considered to have a larger impact on resources and current clinical practice.

25 To aid their recommendations, the Committee requested a cost description on antimicrobials
26 to manage acute pulmonary infection with *P aeruginosa*. Unlike the other pathogens under
27 consideration, *P Aeruginosa* was considered to be one of the most prevalent pathogens that
28 require urgent treatment.

29 Antimicrobials to manage pulmonary infection with *P aeruginosa* include ceftazidime,
30 meropenem and imipenem amongst others. But for these antimicrobials, several brands are
31 available resulting in a variety of acquisition costs. As outlined in NICE's Guide to the
32 methods of technology appraisal 2013, the reduced price should be used in the reference-
33 case analysis to best reflect the price relevant to the NHS. For this reason the lowest cost
34 brand is presented in Table 104. Basic prices are taken from the NHS Electronic Drug Tariff
35 November 2016, unless otherwise stated.

1 **Table 104: Cost of antimicrobials to resolve acute pulmonary infection with *P***
2 ***aeruginosa***

Antimicrobial (quantity, basic price)	Unit cost
Ciprofloxacin (oral)	
100mg tablets (6, £1.86)	£0.31
250mg tablets (10, £0.74)	£0.07
250mg/5ml oral suspension (100ml, £19.80)	£0.99/5ml
500mg tablets (10, £0.87)	£0.09
750mg tablets (10, £8.00)	£0.80
Chloramphenicol (oral)	
250mg capsules (60, £377.00)	£6.28
Aztreonam (inhaled or oral) a	
Cayston 75mg powder and solvent for nebuliser solution vials with Altera Nebuliser Handset (84, £2,181.53)	£25.97
Azactam 1g powder for solution for injection vials (1, £9.40)	£9.40
Azactam 2g powder for solution for injection vials (1, £18.82)	£18.82
Meropenem IV a	
1g powder for solution for injection vials (10, £153.50)	£15.35
500mg powder for solution for injection vials (10, £76.90)	£7.69
Ceftazidime IV a	
500mg powder for solution for injection vials (1, £4.25)	£4.25
1g powder for solution for injection vials (10, £13.90)	£1.39
2g powder for solution for injection vials (10, £27.70)	£2.77
3g powder for solution for injection vials (1, £25.76)	£25.76
Piperacillin-Tazobactam IV a	
Piperacillin 2g / Tazobactam 250mg powder for solution for injection vials (1, £7.91)	£7.91
Piperacillin 4g / Tazobactam 500mg powder for solution for injection vials (1, £12.90)	£12.90
Tazocin 4.5g powder for solution for injection vials (1, £15.17)	£15.17
Fosfomicin IV a	
Fomicyt 2g powder for solution for infusion vials (10, £150.00)	£15.00
Fomicyt 4g powder for solution for infusion vials (10, £300.00)	£30.00
Ticarcillin-Clavulanate IV a	
Timentin 3.2g powder for solution for infusion vials (4, £21.32)	£5.33

3 (l) Taken from the BNF November 2016

4 9.4.2.5 Evidence statements

59.4.2.5.1 *P aeruginosa*

6 Antimicrobial treatment for pulmonary exacerbations due to *P aeruginosa*

7 Comparison 1. Single IV agents compared for pulmonary exacerbations with *P* 8 *aeruginosa*

9 Lung function: FEV₁

10 Low quality evidence from 2 RCTs with 46 young people and adults with cystic fibrosis
11 experiencing a pulmonary exacerbation with *P aeruginosa* showed no clinically significant
12 difference in absolute change of FEV₁ litres between the participants receiving 2 week
13 courses of both ceftazidime (2g 3/day or 8g/day in 4 doses) and those receiving aztreonam

1 (2g 3/day or 8g/day in 4 doses) at 2 weeks follow-up. Moderate inconsistency was observed
2 between both trials, but both trials showed no differences between groups. In addition, the
3 difference in the absolute change in FEV₁ litres was minimal in both groups.

4 **Eradication**

5 No evidence was found for this critical outcome.

6 **Resolution of infection/exacerbation or measure of treatment failure (e.g. need for 7 additional antibiotics)**

8 No evidence was found for this important outcome.

9 **Duration of the acute episode**

10 No evidence was found for this important outcome.

11 **Quality of life**

12 No evidence was found for this important outcome.

13 **Mortality**

14 No evidence was found for this important outcome.

15 **Adverse events**

16 No evidence was found for this important outcome.

17 **Comparison 2. Single IV antibiotic (with placebo) versus combination IV antibiotic for 18 pulmonary exacerbations with *P aeruginosa***

19 **Lung function: FEV₁**

20 Low quality evidence from 1 RCT with 98 young people with cystic fibrosis experiencing a
21 pulmonary exacerbation with *P aeruginosa* showed no clinically significant difference in FEV₁
22 % predicted (absolute change) between the participants who received a 10 day course of IV
23 tobramycin (9mg/kg 3x daily) with placebo and those who received a 10 day course of IV
24 ceftazidime (50mg/kg/dose 3x daily and IV tobramycin 3mg/kg 3x daily) at 10 days follow-up.

25 Very low quality evidence from 1 RCT with 18 young people with cystic fibrosis with *P*
26 *aeruginosa* in sputum admitted to hospital for worsening respiratory status showed no
27 clinically significant difference in FEV₁ % predicted (relative change) between the participants
28 who received a 2 week course of both tobramycin with placebo (5% dextrose 4-hourly) and
29 piperacillin at both 50mg/kg 4-hourly and 100mg/kg 8-hourly at 2 weeks follow-up. All
30 participants received tobramycin 2.5mg/kg 3x daily, oral flucloxacillin 25 mg/kg/day in 4
31 doses and oral probenecid (suggested to increase antibiotic concentrations) at 250 - 500mg
32 3x daily.

33 **Adverse events**

34 Low quality evidence from 1 RCT with 18 young people with cystic fibrosis with *P aeruginosa*
35 in sputum admitted to hospital for worsening respiratory status showed no clinically
36 significant difference in sensitivity reactions between the participants who received a 2-week
37 course of tobramycin with placebo (5% dextrose 4-hourly) and those who received a 2-week
38 course of all regimens of piperacillin (both 50 mg/kg 4-hourly and 100 mg/kg 8-hourly) at 2
39 weeks follow-up. All participants received tobramycin 2.5 mg/kg 3x daily, oral flucloxacillin
40 25 mg/kg/day in 4 doses and oral probenecid (suggested to increase antibiotic
41 concentrations) at 250 - 500 mg 3x daily.

1 Very low quality evidence from 1 RCT with 18 young people with cystic fibrosis with *P*
2 *aeruginosa* in sputum admitted to hospital for worsening respiratory status showed no
3 clinically significant difference in number of hospital admissions due to tinnitus between the
4 participants who received a 10 day course of both tobramycin (9 mg/kg 3x daily) with placebo
5 and those who received ceftazidime (50 mg/kg/dose 3x daily and IV tobramycin 3 mg/kg 3x
6 daily) at 2 weeks follow-up.

7 Very low quality evidence from 1 RCT with 44 young people with cystic fibrosis experiencing
8 an exacerbation with *P aeruginosa* showed no clinically significant difference in serum or
9 creatinine levels between the participants who received a 10 day course of both tobramycin
10 (9 mg/kg 3x daily) with placebo and those who received ceftazidime (50 mg/kg/dose 3x daily
11 and IV tobramycin 3 mg/kg 3x daily) at 2 weeks follow-up.

12 However, moderate quality evidence from the same trial showed a clinically significant lower
13 levels of NAG in the participants who received a 10 day course of tobramycin (9 mg/kg 3x
14 daily) with placebo compared with who received a 10 day course combination of IV
15 ceftazidime (50 mg/kg/dose 3x daily) and IV tobramycin (3 mg/kg 3x daily) at 2 weeks follow-
16 up.

17 **Comparison 3. Single IV antibiotic versus combination IV antibiotic for pulmonary** 18 **exacerbations with *P aeruginosa***

19 **Lung function: FEV₁**

20 Low quality evidence from 1 RCT with 30 young people with cystic fibrosis experiencing an
21 acute exacerbation due to *P aeruginosa* showed a clinically significant beneficial effect of a
22 10-14 day course of combination ticarcillin (300 mg/kg/day in 4 doses) and tobramycin (10
23 mg/kg/day in 3 doses) in FEV₁ % relative change compared with a 10-14 day course
24 ceftazidime (200 mg/kg/day in 4 doses) at 2 weeks follow-up.

25 Low quality evidence from 1 RCT with 71 adults with cystic fibrosis experiencing a pulmonary
26 exacerbation with *P aeruginosa* showed a clinically significant beneficial effect of a 12 day
27 course of a combination of colistin (2 MU 3x daily) and a second anti-pseudomonal antibiotic
28 in FEV₁ (ml) absolute change compared with colistin (2 MU 3x daily) alone at 12 days follow-
29 up.

30 Very low quality evidence from 1 RCT with 21 young people with cystic fibrosis experiencing
31 a pulmonary exacerbation with *P aeruginosa* showed no clinically significant difference in
32 FEV₁ % predicted (absolute change) between a 2 week course of ceftazidime 50 mg/kg 3x
33 daily and a combination of piperacillin (75 mg/kg 4x daily) and tobramycin (10 mg/kg/day in 3
34 doses) at 2 weeks follow-up.

35 **Eradication**

36 Low quality evidence from 1 RCT with 38 children with cystic fibrosis showed a clinically
37 significant beneficial effect of combination of piperacillin (600 mg/kg/day) and (tobramycin 8 -
38 10 mg/kg/day) in eradicating *P aeruginosa* compared with piperacillin alone (600 mg/kg/day).

39 **Resolution of infection/exacerbation or measure of treatment failure (e.g. need for** 40 **additional antibiotics)**

41 Very low quality evidence from 1 RCT with 19 children with cystic fibrosis admitted to hospital
42 for treatment of a pulmonary exacerbation with *P aeruginosa* showed no clinically significant
43 difference in time to readmission (months) between a 2 week course of ceftazidime (50
44 mg/kg 3x daily) and a 2 week course combination piperacillin (75 mg/kg 4x daily) and
45 tobramycin (10 mg/kg/day in 3 doses) at 3 months follow-up.

1 Very low quality evidence from 1 RCT with 22 children and young people with cystic fibrosis
2 and severe chest infection treated for an exacerbation with *P aeruginosa* showed no clinically
3 significant difference in number of admissions to hospital requiring IV antibiotics or mortality
4 between a 2 week course of ceftazidime (150 mg/kg/day) and a 2 week course of
5 combination tobramycin (7.5 mg/kg/day) and ticarcillin (300 mg/kg/day) at 3 months follow-
6 up.

7 **Duration of the acute episode**

8 No evidence was found for this important outcome.

9 **Quality of life**

10 No evidence was found for this important outcome.

11 **Mortality**

12 Low quality evidence from 1 RCT with 21 young people with cystic fibrosis experiencing a
13 pulmonary exacerbation with *P aeruginosa* showed no clinically significant difference in
14 mortality between a 2 week course of ceftazidime 50 mg/kg 3x daily and 2 week course
15 combination piperacillin (75 mg/kg 4x daily) and tobramycin (10 mg/kg/day in 3 doses) at 4
16 months follow-up.

17 Low quality evidence from 1 RCT with 71 adults with cystic fibrosis experiencing a pulmonary
18 exacerbation with *P aeruginosa* showed no clinically significant difference in mortality
19 between a 12 day course of a combination of colistin (2 MU 3x daily) and a second anti-
20 pseudomonal antibiotic in FEV₁ (ml) absolute change compared with 12 day course colistin
21 (2 MU 3x daily) alone at 12 weeks follow-up.

22 **Adverse events**

23 Very low quality evidence from 2 RCTs with 52 children and young people with cystic fibrosis
24 experiencing an exacerbation due to *P aeruginosa* showed no clinically significant difference
25 in liver transaminase enzyme elevation between a 10-14 day course of ceftazidime and
26 combination ticarcillin and tobramycin.

27 Low quality evidence from 1 RCT with 71 adults with cystic fibrosis experiencing a pulmonary
28 exacerbation with *P aeruginosa* showed no clinically significant difference in neurological
29 adverse effects between a 12 day course colistin (2 MU 3x daily) alone and a 12 day course
30 combination of colistin (2 MU 3x daily) and a second anti-pseudomonal antibiotic at 12 days
31 follow-up.

32 Very low quality evidence from 1 RCT with 17 children with cystic fibrosis admitted for
33 treatment of pulmonary exacerbations showed no clinically significant difference in rash and
34 fever between a 10 day course of piperacillin alone (600 mg/kg/day) and a 10 day course
35 combination piperacillin (600 mg/kg/day) and tobramycin (8 - 10 mg/kg/day) at 10 days
36 follow-up.

37 Very low quality from 1 RCT with 30 young people with cystic fibrosis and *P aeruginosa*
38 infection (34 treatment observations) showed no clinically significant difference in proteinuria
39 between a 10-14 day course of ceftazidime (200 mg/kg/day in 4 doses) and combination
40 ticarcillin (300 mg/kg/day in 4 doses) and 10-14 day course tobramycin (10 mg/kg/day in 3
41 doses) in FEV₁ % relative change compared with ceftazidime.

42 Low and very low quality evidence from 1 RCT with 71 adults with cystic fibrosis
43 experiencing a pulmonary exacerbation with *P aeruginosa* showed no clinically significant
44 difference in change in blood urea (mmol/L) and change in serum creatine (mol/L) between a
45 12 day course of a combination of colistin (2 MU 3x daily) and a 12 day course second anti-
46 pseudomonal antibiotic and colistin (2 MU 3x daily) alone at 12 days follow-up.

1 **Comparison 4. Combination IV antibiotics versus combination IV antibiotics for**
2 **pulmonary exacerbations with *P aeruginosa***

3 **Lung function: FEV₁**

4 Low quality evidence from 1 RCT including observations for 49 courses of IV combination
5 therapy (≈42 people with cystic fibrosis aged 3 to 24 years admitted to hospital due to a
6 pulmonary exacerbation with *P aeruginosa*) showed no clinically significant difference in
7 FEV₁ % predicted (absolute change) between a 2 week course of combination of aztreonam
8 (300 mg/kg/day in 4 doses) and amikacin (36 mg/kg/day in 3 doses) and a 2 week course
9 combination of ceftazidime (300 mg/kg/day in 4 doses) and amikacin (36 mg/kg/day in 3
10 doses) at 2 weeks follow-up.

11 Very low to low quality evidence from 1 RCT with 97 people with cystic fibrosis ≥5 years
12 treated for pulmonary exacerbation showed no clinically significant difference in both FEV₁ %
13 predicted absolute change and relative change between a 2 week course of combination of
14 meropenem (40 mg/kg up to a maximum dose of 2 g) and tobramycin and a 2 week course
15 combination of ceftazidime (50 mg/kg) and tobramycin. Tobramycin dose adjusted to give a
16 peak serum concentration of ≥ 8 µg/mL and trough concentration of < 2 µg/mL.

17 **Eradication**

18 Very low quality evidence from 1 RCT including observations for 49 courses of IV
19 combination therapy (≈42 people with cystic fibrosis aged 3 to 24 years admitted to hospital
20 due to a pulmonary exacerbation with *P aeruginosa*) showed no clinically significant
21 difference in eradication of *P aeruginosa* between a 2 week course of combination of
22 aztreonam (300 mg/kg/day in 4 doses) and amikacin (36 mg/kg/day in 3 doses) and a 2
23 week course combination of ceftazidime (300 mg/kg/day in 4 doses) and amikacin (36
24 mg/kg/day in 3 doses) at 2 weeks follow-up.

25 **Resolution of infection/exacerbation or measure of treatment failure (e.g. need for**
26 **additional antibiotics)**

27 No evidence was found for this important outcome.

28 **Duration of the acute episode**

29 No evidence was found for this important outcome.

30 **Quality of life**

31 No evidence was found for this important outcome.

32 **Mortality**

33 No evidence was found for this important outcome.

34 **Adverse events**

35 Very low quality evidence from 1 RCT including observations for 56 courses of IV
36 combination therapy (≈42 people with cystic fibrosis aged 3 to 24 years admitted to hospital
37 due to a pulmonary exacerbation with *P aeruginosa*) showed no clinically significant
38 difference in rash, liver transaminase levels and thrombocytopenia between a 2 week course
39 of combination of aztreonam (300 mg/kg/day in 4 doses) and amikacin (36 mg/kg/day in 3
40 doses) compared with a 2 week course of combination of ceftazidime (300 mg/kg/day in 4
41 doses) and amikacin (36 mg/kg/day in 3 doses) at 2 weeks follow-up.

1 **Comparison 5. Combination of 2 IV antibiotics + inhaled antibiotic versus combination**
2 **of 2 IV antibiotics without inhaled antibiotic for pulmonary exacerbations with P**
3 **aeruginosa**

4 **Lung function**

5 No evidence was found for this critical outcome.

6 **Eradication of pathogen**

7 Moderate quality evidence from 1 RCT including observations for 84 courses of treatment (≈
8 62 people with cystic fibrosis aged 3 to 24 years admitted to hospital due to a pulmonary
9 exacerbation with *P aeruginosa*) showed a clinically significant beneficial effect of a 15 day
10 course of IV ceftazidime (250 mg/kg/day in 4 doses) and IV amikacin (33 mg/kg/day in 3
11 doses) and nebulised amikacin (100 mg 2x daily) compared with a 15 day course IV
12 ceftazidime and IV amikacin without nebulised amikacin at 15 days follow-up.

13 **Duration of the acute episode**

14 No evidence was found for this important outcome.

15 **Quality of life**

16 No evidence was found for this important outcome.

17 **Mortality**

18 No evidence was found for this important outcome.

19 **Adverse events**

20 Very low quality evidence from 1 RCT with including observations for 54 courses of treatment
21 (≈62 people with cystic fibrosis aged 3 to 24 years admitted to hospital due to a pulmonary
22 exacerbation with *P aeruginosa*) found no clinically significant difference in raised liver
23 transaminases between a 15 day course of IV ceftazidime (250 mg/kg/day in 4 doses) and IV
24 amikacin (33 mg/kg/day in 3 doses) and nebulised amikacin (100 mg 2x daily) compared with
25 IV ceftazidime and IV amikacin without nebulised amikacin.

26 **Comparison 6. Combination IV antibiotics versus oral antibiotics for pulmonary**
27 **exacerbations with P aeruginosa**

28 **Lung function**

29 No evidence was found for this critical outcome.

30 **Eradication of pathogen**

31 Moderate quality evidence from 1 RCT with 89 children with cystic fibrosis and *P aeruginosa*
32 infection showed a clinically significant beneficial effect of a 2 week course of combination of
33 IV ceftazidime (50 mg/kg 3x daily) and a 2 week IV tobramycin (3 mg/kg 3x daily) in
34 eradicating *P aeruginosa* compared with oral ciprofloxacin (15 mg/kg 2x daily) at 2 weeks
35 follow-up.

36 **Duration of the acute episode**

37 No evidence was found for this important outcome.

38 **Quality of life**

39 No evidence was found for this important outcome.

40 **Mortality**

1 No evidence was found for this important outcome.

2 **Adverse events**

3 Very low quality evidence from 1 RCT with 108 children with cystic fibrosis and *P aeruginosa*
4 infection showed no difference in treatment-related adverse events between a 2 week course
5 of combination IV ceftazidime (50 mg/kg 3x daily) and IV tobramycin (3 mg/kg 3x daily) and
6 oral ciprofloxacin (15 mg/kg 2x daily).

7 **Antimicrobial treatment for acute infection with *P aeruginosa***

8 **Comparison 7. Oral ciprofloxacin and inhaled colistin versus inhaled tobramycin for**
9 **acute infection with *P aeruginosa***

10 **Lung function**

11 No evidence was found for this critical outcome.

12 **Eradication of pathogen**

13 No evidence was found for this critical outcome.

14 **Time to next pulmonary exacerbation**

15 No evidence was found for this important outcome.

16 **Resolution of infection/exacerbation or measure of treatment failure**

17 No evidence was found for this important outcome.

18 **Quality of life**

19 No evidence was found for this important outcome.

20 **Adverse events**

21 Very low quality evidence from 1 RCT with 58 children with cystic fibrosis with new isolation
22 of *P aeruginosa* showed no clinically significant difference in adverse events (severe cough)
23 between a 3 month course of combination inhaled colistin (2 MU 2x daily) and oral
24 ciprofloxacin (10 mg/kg 3x daily) compared with a 3 month course of inhaled tobramycin (300
25 mg 2x daily for 28 days) at 3 months follow-up.

26 **Comparison 8. Inhaled colistin and oral ciprofloxacin versus inhaled tobramycin and**
27 **oral ciprofloxacin for acute infection with *P aeruginosa***

28 **Lung function: FEV₁**

29 Very low quality evidence from 1 RCT with 128 people with cystic fibrosis >1 year with first
30 ever or new *P aeruginosa* infection showed no clinically significant difference in FEV₁ %
31 predicted (relative change) between a 28 day course of combination colistin (2x daily
32 inhalation of 2 MU) and ciprofloxacin (2x daily of 15 mg/kg/dose) compared with a 28 day
33 course combination of tobramycin inhaled solution (300 mg 2x daily) and ciprofloxacin (2x
34 daily doses of 15 mg/kg/dose) at 54 days follow-up.

35 **Eradication of pathogen**

36 No evidence was found for this critical outcome.

37 **Time to next pulmonary exacerbation**

38 No evidence was found reporting this important outcome.

1 **Resolution of infection/exacerbation or measure of treatment failure**

2 Very low quality evidence from 1 RCT with 128 people with cystic fibrosis >1 year with first
3 ever or new *P aeruginosa* infection showed no clinically significant difference in
4 discontinuation due to lack of compliance between a 28 day course of combination colistin
5 (2x daily inhalation of 2 MU) and ciprofloxacin (2x daily of 15 mg/kg/dose) compared with a
6 28 day course of combination of tobramycin inhaled solution (300 mg 2x daily) and
7 ciprofloxacin (2x daily doses of 15 mg/kg/dose) at 28 days follow-up.

8 Very low quality evidence from 1 RCT with 128 people with cystic fibrosis >1 year with first
9 ever or new *P aeruginosa* infection showed no clinically significant difference in pulmonary
10 exacerbation during early eradication treatment leading to treatment failure between a 28 day
11 course of combination colistin (2x daily inhalation of 2 MU) and ciprofloxacin (2x daily of 15
12 mg/kg/dose) compared with a 28 day course combination of tobramycin inhaled solution (300
13 mg 2x daily) and ciprofloxacin (2x daily doses of 15 mg/kg/dose) at 28 days follow-up.

14 **Quality of life (QOL)**

15 No evidence was found for this important outcome.

16 **Adverse events**

17 Very low quality evidence from 1 RCT with 128 people with cystic fibrosis >1 year with first
18 ever or new *P aeruginosa* infection showed no clinically significant difference in adverse
19 events (vomiting, photosensitivity, wheeze) between a 28 day course of combination colistin
20 (2x daily inhalation of 2 MU) and ciprofloxacin (2x daily of 15 mg/kg/dose) compared with a
21 28 day course combination of tobramycin inhaled solution (300 mg 2x daily) and ciprofloxacin
22 (2x daily doses of 15 mg/kg/dose) at 28 days follow-up.

239.4.2.5.2 ***S aureus***

24 No evidence was found.

259.4.2.5.3 ***B cepacia complex***

26 No evidence was found.

279.4.2.5.4 ***H influenza***

28 No evidence was found.

299.4.2.5.5 ***Nontuberculous mycobacteria***

30 No evidence was found.

319.4.2.5.6 ***Non-identified pathogen***

32 No evidence was found.

339.4.2.5.7 ***Economic evidence statements***

34 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

35 **9.4.2.6 Linking evidence to recommendations**

369.4.2.6.1 ***Relative value placed on the outcomes considered***

37 The aim of this review was to compare the clinical and cost effectiveness of different
38 antimicrobial regimens in achieving clinical resolution of acute pulmonary infection or
39 exacerbation in children and adults with cystic fibrosis.

1 The Committee chose eradication of specific pathogens where present and improvement in
2 lung function measured using either FEV₁ or the lung Clearance Index (LCI) as critical
3 outcomes for decision making for both acute infection and pulmonary exacerbation.

4 In addition they chose the following as important outcomes, resolution of exacerbation or
5 measures of treatment failure (e.g. need for additional antibiotics), duration of the episode,
6 quality of life, adverse events and mortality for pulmonary exacerbations. For acute
7 pulmonary infection, the important outcomes were time to next acute infection, resolution of
8 infection or measure of treatment failure (e.g. need for additional antibiotics), quality of life
9 and adverse events.

109.4.2.6.2 **Consideration of clinical benefits and harms**

11 The Committee discussed the recommendations for each pathogen separately.

12 ***P aeruginosa***

13 The Committee acknowledged the evidence presented to them, and discussed it in the light
14 of their clinical expertise and experience.

15 The Committee discussed 2 different scenarios: acute infection and pulmonary exacerbation.

16 If a person with cystic fibrosis develops a new infection with *P aeruginosa* (meaning a
17 positive respiratory secretion sample culture, previous cultures in the recent past having
18 been negative) the Committee agreed antibiotic treatment is needed. They emphasized early
19 treatment of *P aeruginosa* infection is very important, as it is recognised chronic infection
20 with this pathogen has a negative impact in the quality of life.

21 The Committee noted the management will differ depending on the severity of the symptoms.

22 If the person is clinically well, the Committee suggested it should be treated in order to try to
23 eradicate it using a combination of systemic antibiotics (oral or intravenous), together with an
24 inhaled antibiotic. They discussed the use, for example of oral ciprofloxacin combined with
25 inhaled colistin or nebulised tobramycin. The recommendation to treat this group was based
26 on the Committee's recognition that *P aeruginosa* is an important pathogen in cystic fibrosis
27 and in their expert opinion intensive treatment with systemic and inhaled antibiotics should
28 improve the chances of eradication. The Committee made this recommendation based on
29 their clinical experience, as the available evidence was very scarce and of very low quality,
30 and therefore not very useful to make recommendations.

31 If the person is clinically unwell, for example with new respiratory symptoms and signs or a
32 worsening of existing respiratory symptoms and signs, the approach might be different. The
33 Committee recommended that in that situation the initial therapy should consist of a course
34 of intravenous antibiotics with an inhaled antibiotic. They discussed, for example, the use of
35 two anti-pseudomonal antibiotics - such as ceftazidime and tobramycin - given intravenously
36 together with the inhaled antibiotic. This is because moderate quality evidence showed that
37 participants who received an inhaled antibiotic in addition to a combination of 2 intravenous
38 antibiotics were less likely to be admitted to hospital due to a pulmonary exacerbation.

39 In both groups, based on the consensus of the committee, they advised giving extended
40 treatment in order to try to increase the likelihood of eradication.

41 In the event that eradication was unsuccessful, the Committee agreed that prolonged
42 treatment with an inhaled antibiotic should be given to try and suppress it and they
43 recommendations using colistimethate as the first line choice for this inhalation therapy.
44 Please see antimicrobial treatment for the management of chronic *P aeruginosa*.

45 For the management of pulmonary exacerbations due to *P aeruginosa* in people who are
46 chronically infected with pseudomonas, the Committee agreed to recommend oral or

1 intravenous antibiotics, depending on the severity of the illness. The intravenous treatment
2 should consist in a combination of 2 agents, as supported by the evidence included in this
3 review. The Committee noted that low quality evidence showed a clinically significant
4 beneficial effect in lung function and in eradication of the organism of a combination of 2
5 intravenous antibiotics compared to a single antibiotic. In addition, the evidence showed no
6 clinically significant difference in the occurrence of adverse events.

7 Finally, the Committee recommended that if people with chronic *P aeruginosa* infection suffer
8 from repeated pulmonary exacerbations, consideration should be given to altering the
9 antibiotic regimen used at intervals, taking account of the individual's pseudomonas antibiotic
10 sensitivity testing, in order to reduce the possibility of non-response due to emergence of
11 resistance. This recommendation was based on the consensus of the Committee.

12 ***S aureus***

13 No evidence was found for the treatment of *S aureus*, therefore recommendations were
14 based on Committee's clinical expertise.

15 The Committee discussed 2 possible scenarios, depending on whether the child is on
16 prophylaxis treatment.

17 The Committee recognised that a potential reason for emergence of *S aureus* infection was
18 non-adherence to the prophylaxis regimen, and so they advised this should be reviewed with
19 parents and carers. They also advised that following attempted eradication with antibiotics
20 the prophylaxis should be reinstated even if the eradication was unsuccessful, as they
21 believed that suppression of the infection was likely to be clinically beneficial.

22 If a child is not on prophylaxis against *S aureus*, and is then found to have developed an
23 infection with this pathogen, the Committee recommended oral antibiotic treatment if they are
24 well, and they discussed as potentially useful antibiotic choices flucloxacillin, co-amoxiclav or
25 doxycycline for persons over 12 years. If, however, they are unwell (for example, with
26 symptoms such as cough) or have evidence of pulmonary disease (for example, reduced
27 lung function based on testing) then either oral or intravenous antibiotic treatment, depending
28 on disease severity is recommended. The antibiotic used should be broad spectrum to take
29 account of the possibility that *S aureus* might not be the cause of the illness, but the
30 treatment should include anti-*S aureus* cover.

31 ***B cepacia* complex**

32 No evidence was found for the treatment of *B cepacia* complex, therefore recommendations
33 were based on Committee's clinical expertise.

34 The Committee agreed that if a person develops a new infection with *B cepacia* complex, an
35 attempt should be made to eradicate the infection with antibiotic therapy whether or not the
36 person was unwell with the infection. They considered that specialist advice should be
37 sought on this treatment. They suggested a combination of 2 or 3 appropriate intravenous
38 antibiotics would usually be given. Example of intravenous antibiotics that might be advised
39 included but are not limited to ceftazidime, meropenem, amikacin, and temocillin and
40 specialist advice on the exact regimen was required. The Committee noted that it is
41 important to treat new *B cepacia* complex infections effectively as chronic infection can
42 cause a deterioration in lung function, and in some people an overwhelming and even fatal
43 infection called 'cepacia syndrome' may occur. Persistent isolation of *B cepacia* complex
44 may also adversely impact on a persons' eligibility for transplantation.

45 The Committee also discussed the case of people with chronic *B cepacia* complex infection
46 despite attempts at eradication. They did not recommend treatment for those with chronic *B*
47 *cepacia* complex who are clinically well, as they noted that *B cepacia* complex is very
48 resistant to most treatments once established, and therefore unlikely to work. However, if

1 they became unwell with a pulmonary exacerbation, the Committee recommended again that
2 specialist advice be sought regarding the use of oral or intravenous antibiotics. They
3 discussed that this would usually be with a course of intravenous antibiotics, although oral
4 antibiotics might also be used for a less severe exacerbation.

5 ***H influenza***

6 No evidence was found for the treatment of *H influenza*, therefore recommendations were
7 based on Committee's clinical expertise.

8 The Committee agreed it is important to treating *H influenza* in order to prevent chronic
9 infection with this pathogen. This is because although there might not be detectable evidence
10 of disease due to it, the belief is that it will cause lung damage so it should be eradicated.
11 They discussed 2 possible scenarios if a person develops an infection.

12 If the person is clinically well (asymptomatic), the Committee recommended giving an oral
13 antibiotic agent; whereas if the person is clinically unwell (for example with cough or reduced
14 lung function), they recommended the use of an oral or intravenous antibiotic treatment
15 depending on the severity of the illness.

16 These recommendations are consistent with clinical practice and with the CF Trust
17 recommendations (CF Consensus document: antibiotic treatment for Cystic Fibrosis, 2009).

18 **Nontuberculous *mycobacteria***

19 No evidence was found for the treatment of nontuberculous *mycobacteria*, therefore
20 recommendations were based on Committee's clinical expertise. The Committee noted that
21 treatment is complex and guidelines are evolving, and there is still uncertainty about the best
22 approach to treatment.

23 The Committee emphasised the importance of confirming the presence on nontuberculous
24 *mycobacteria* by repeating respiratory secretion cultures. This is because nontuberculous
25 *mycobacteria* are often just sporadic commensal organisms – that is, they just come and go
26 without causing disease. Therefore, the first step when this pathogen is found, is to ensure
27 that it persists before actually considering whether it is causing disease. This diagnostic
28 issue was also raised by the CF Trust Consensus document on antibiotic treatment for cystic
29 fibrosis (CF Consensus document: antibiotic treatment for Cystic Fibrosis, 2009).

30 The Committee noted in a person with respiratory disease who is found to be NTM positive it
31 was not always easy to determine whether or not the infection was contributing to the
32 disease. Therefore they recommended that a chest CT scan should be performed because
33 this may show changes that would support the likely importance of nontuberculous
34 *mycobacteria* to the disease.

35 The Committee recommended that a discussion should take place with the person affected
36 and if appropriate with parents or carers about the uncertain benefits of attempting
37 nontuberculous *mycobacteria* eradication, and regarding the fact that when considering a
38 decision to treat it was important to be aware of the potential toxicities associated with the
39 drugs used. Potential toxic effects included vomiting, nephrotoxicity, and ototoxicity. This is
40 particularly pertinent for persons who are positive for nontuberculous *mycobacteria* but
41 clinically well, where the benefits are less certain.

42 The Committee recommended that consideration be given to treatment for those who are
43 positive for nontuberculous *mycobacterium* respiratory infection and who have a chest CT
44 scan showing changes consistent with it and who are unwell with respiratory disease, despite
45 optimisation of other treatment. The recommendation made by the Committee to treat based
46 on clinical grounds is consistent with the CF Trust recommendations (CF Consensus
47 document: antibiotic treatment for Cystic Fibrosis, 2009).

1 The Committee recognised that evidence regarding the optimal antibiotic regimen and
2 duration of treatment was lacking. The Committee discussed the fact that the approaches to
3 treating *M avium* complex and *M abscessus* differ. Currently, treatment for *M avium* typically
4 includes a combination of 3 oral anti-mycobacterial agents including a macrolide and
5 rifampicin and ethambutol. Current eradication treatment for *M abscessus* is more intensive
6 and may include repeated courses of triple antibiotic therapy administered intravenously,
7 together with a combination of oral and inhaled antibiotics. Antibiotics used include, but are
8 not limited, to ceftazidime, tigecycline, amikacin, carbapenems and macrolides. They
9 recommended that specialist advice be sought when considering treatment.

10 **Non-identified pathogen**

11 No evidence was found for the treatment of unidentified infections, therefore
12 recommendations were based on Committee's clinical expertise.

13 The Committee agreed that if a person presents with clinical manifestations suggesting
14 development of an acute pulmonary infection or an exacerbation without an identified
15 pathogen, the Committee agreed it would be appropriate to treat empirically based on likely
16 causative organisms which may be influenced by symptoms and previous microbiology
17 results. The choice of oral or IV treatment will depend on the severity of the symptoms.

18 Treatment should be modified once the pathogen causing the infection or the exacerbation is
19 identified.

209.4.2.6.3 **Consideration of economic benefits and harms**

21 The clinical evidence review demonstrated that the benefits of combination antibiotics to treat
22 exacerbations due to *P aeruginosa* could justify their additional cost relative to single
23 antibiotics. For example, clinically significant beneficial effects in lung function were found for
24 combination IV antibiotics compared to single IV antibiotics, and clinically significant benefits
25 in eradication were found for the more intensive regimen; specifically, combination IV
26 antibiotics compared to single antibiotic therapy, 2 IV antibiotics plus an inhaled antibiotic
27 compared to IV antibiotics without an inhaled antibiotic, and combination IV antibiotics
28 compared to oral antibiotics.

29 No evidence was found for the other pathogens listed in the protocol, thus the Committee
30 made recommendations to reflect current clinical practice and resource use as they
31 considered this to be a cost-effective use of resources.

32 The specific antibiotics administered in the trials were of less importance to the Committee
33 as antibiotics received by a person with cystic fibrosis to treat an acute infection, or
34 exacerbation, need to be varied in accordance with cultures, sensitivities and local resistance
35 patterns of isolated pathogens. Consequently, the Committee stated it would be
36 inappropriate to recommend specific antibiotics or a number of antibiotics as this would limit
37 the variation in antibiotics used by healthcare professionals and subsequently, the
38 effectiveness and cost-effectiveness of the antibiotic over time. Instead, the Committee were
39 interested in the combination and preparation of antibiotics that were administered to infer
40 which regimens were cost-effective.

41 When the same number of different combinations of antibiotics, of the same preparation (for
42 example two IV antibiotics vs. a different combination of two IV antibiotics), were compared,
43 no significant difference was demonstrated for any of the proposed outcomes. However, the
44 Committee iterated that no significance difference does not infer that there was no clinical
45 change pre- versus post treatment, as the trials included a comparator that was an
46 antimicrobial agent that would be expected to issue a treatment effect.

47 In light of the findings from the clinical evidence review and their own clinical experience, the
48 Committee wanted to recommend 2 antibiotics in different classes and consider changing

1 regimens over time when treating exacerbations associated with *P aeruginosa*. The
2 Committee also added, when discussing cost implications, that ceftazidime is used more
3 often and is less expensive than aztreonam and meropenem.

4 The Committee agreed IV antibiotics are generally more expensive than oral preparations for
5 many reasons regardless of the active agent, including their purchase price, and in some
6 cases sterile production and healthcare professional administration. Additionally, they are
7 more invasive and have a greater potential for associated adverse effects. Consequently, the
8 Committee recommended that where appropriate, if the person is clinically well
9 (asymptomatic), treatment with an oral agent should be considered; whereas if the person is
10 clinically unwell (for example presents with cough or reduced lung function), the use of an
11 appropriate oral or IV antibiotic, could also be considered depending on the severity of
12 symptoms.

13 The Committee were reluctant to recommend “no treatment” in people with cystic fibrosis for
14 whom a pathogen has been isolated as the lack of evidence should not infer lack of effect.
15 Given that the expected downstream costs from an untreated infection would outweigh the
16 cost of treatment, the Committee agreed a broad-spectrum antibiotic should be offered.

179.4.2.6.4 **Quality of evidence**

18 ***P aeruginosa***

19 The quality of the evidence was rated as very low to moderate as assessed by GRADE for
20 the antimicrobial treatment due to pulmonary exacerbations with *P aeruginosa*, and very low
21 for the antimicrobial treatment of acute infections with *P aeruginosa*. No high quality
22 evidence was found. The main reasons that led to downgrading the quality of the evidence
23 were:

- 24 • For the domain risk of bias, the studies were assigned the same risk of bias as in the
25 Cochrane reviews, and were not individually reviewed. The main biases that lead to
26 downgrading the quality of the evidence were attrition bias and lack of blinding.
- 27 • Another reason that lead to downgrading the quality of the evidence was imprecision, as
28 confidence intervals crossed 1 or 2 MIDs.

29 No serious issues were found regarding the directness of the population or the interventions.

30 ***S aureus***

31 Not applicable, as no evidence was found for this pathogen.

32 ***B cepacia* complex**

33 Not applicable, as no evidence was found for this pathogen.

34 ***Non-tuberculous mycobacteria***

35 Not applicable, as no evidence was found for this pathogen.

36 **Non-identified pathogen**

37 Not applicable, as no evidence was found for this pathogen.

389.4.2.6.5 **Other considerations**

39 No equality issues were identified by the Committee for this review question.

40 The Committee agreed a research recommendation should not be prioritised for this topic.
41 They noted there is sufficient evidence available on the management of acute exacerbations

1 with *P aeruginosa*. They also noted studies on the management of acute infection with other
2 pathogens are difficult to conduct. Recommendations are consistent with clinical practice,
3 and new research is unlikely to lead to significant changes.

49.4.2.6.6 **Key conclusions**

5 The Committee concluded that:

- 6 • People with cystic fibrosis who present with a new infection with *P aeruginosa* should be
7 treated with a combination of oral or IV antibiotics together with inhaled antibiotics,
8 regardless of whether the person is symptomatic or not. A follow-up antibiotic treatment
9 could be considered with the aim to eradicate the pathogen.
- 10 • People with cystic fibrosis who present with an acute exacerbation due to *P aeruginosa*
11 should be treated with a combination of oral or IV antibiotics, depending on the severity of
12 the illness. Agents should be changed over time.
- 13 • People with cystic fibrosis who present with a new infection with *S aureus* and are
14 clinically well should be treated with an oral staphylococcal-agent, or have their dose
15 increased if they are on prophylactic treatment.
- 16 • People with cystic fibrosis who present with a new infection with *S aureus* and are
17 clinically unwell may require treatment with an additional oral or intravenous anti-
18 staphylococcal antibiotic.
- 19 • People with cystic fibrosis who present with a new infection with *S aureus* when taking
20 anti-staphylococcal prophylaxis should have their adherence checked before increasing
21 the dose of the antibiotic or changing to a different agent.
- 22 • People with cystic fibrosis who present with a new infection with *B cepacia* complex
23 should be given a combination of IV antibiotics, regardless of whether they are
24 symptomatic or not. A follow-up treatment with oral or inhaled antibiotics could be
25 considered with the aim to eradicate the pathogen.
- 26 • People with cystic fibrosis who present with an acute exacerbation due to *B cepacia*
27 complex should be treated with appropriate oral or IV antibiotics, depending on the
28 severity of the illness.
- 29 • People with cystic fibrosis who present with a new infection with *H influenzae* should be
30 treated with appropriate oral or IV antibiotics, depending on the severity of the symptoms.
- 31 • People with cystic fibrosis who present with a new infection where non-tuberculous
32 *mycobacteria* is suspected should have the diagnosis confirmed prior to commencing
33 treatment. Combination anti-mycobacterial treatment should be considered. Specialist
34 microbiologist advice should be sought.
- 35 • People with cystic fibrosis who present with an acute exacerbation due to non-tuberculous
36 *mycobacteria* should be treated with appropriate oral or IV antibiotics, depending on the
37 severity of the illness.
- 38 • People with cystic fibrosis who present with an exacerbation without a known pathogen
39 should be treated empirically with an oral or IV antibiotic. Treatment should be changed
40 accordingly, once the pathogen has been identified.

1 9.4.3 Chronic

2 **Review question: What is the effectiveness of antimicrobial regimens in suppressing**
3 **chronic pulmonary infection in children and adults with cystic fibrosis with any of the**
4 **following pathogens: *P aeruginosa*, *B cepacia* Complex, *S aureus* and *Aspergillus***
5 ***Fumigatus*?**

6 9.4.3.1 Description of clinical evidence

7 The aim of this review was to determine the clinical and cost-effectiveness of different
8 antimicrobial treatment regimens to suppress chronic pulmonary infection in children and
9 adults with cystic fibrosis and one of the following pathogens:

- 10 • *P aeruginosa*
- 11 • *S aureus*
- 12 • *B cepacia* complex
- 13 • *A fumigatus*

14 We searched for systematic reviews of RCTs and RCTs, including cross-over trials.
15 Systematic reviews were assessed for inclusion against the protocol, and if relevant, their
16 quality was assessed using AMSTAR. High-quality systematic reviews were included in our
17 review, and where possible, data and quality assessment was taken directly from the review.
18 Individual studies were also retrieved for completeness and accuracy, and were also
19 checked for additional outcomes of interest. Low-quality SR were excluded from our review,
20 but their lists of included studies were checked to identify relevant trials.

21 For full details see review protocol in Appendix D.

22 The results are presented separately for each pathogen.

239.4.3.1.1 *P Aeruginosa*

24 The interventions that were included in the protocol for the treatment of chronic infection with
25 *P aeruginosa* were: Aztreonam lysine (nebulised), Azythromycin (oral, immunomodulatory-
26 dose only), Ciprofloxacin (oral), Colistimethate sodium (dry powder, inhaled), Fosfomycin
27 (inhaled) and Tobramycin (dry powder, inhaled).

28 One NICE TA report 276 (Tappenden 2013) has been published to provide guidance on the
29 treatment of chronic *P Aeruginosa*. This systematic review evaluated the effectiveness and
30 cost-effectiveness of Colistimethase sodium dry powder inhalation and Tobramycin dry
31 powder inhalation for the treatment of chronic *P Aeruginosa* lung infection in people with
32 cystic fibrosis over the age of 6 years. Three trials were included in the review
33 (COLO/DPI/02/05, COLO/DPI/02/06, Konstan 2011a/ EAGER trial).

34 Four Cochrane systematic reviews were identified in the search.

- 35 • Two reviews were included:
 - 36 ○ Remington (2016) evaluated the effectiveness of oral anti-pseudomonal antibiotics
 - 37 for cystic fibrosis. One trial was included from this review (Sheldon 1993).
 - 38 ○ Ryan (2011) evaluated the effectiveness of inhaled antibiotics for long-term therapy in
 - 39 cystic fibrosis. Eight trials were included from this review (Chuchalin 2007, Hodson
 - 40 2002, Jensen 1987, Lenoir 2007, McCoy 2008, Murphy 2004, Ramsay 1993, Ramsey
 - 41 1999).
- 42 • Two reviews were excluded:
 - 43 ○ Elphick (2016) evaluated the effectiveness of single versus combination IV antibiotic
 - 44 therapy for treating people with cystic fibrosis. One trial included people with cystic
 - 45 fibrosis and chronic infection with *P Aeruginosa*, but it was not included in the review as

1 it evaluated the effectiveness of Ceftazidime, a treatment that was not prioritised by the
2 Committee in the evidence review protocol.

3 ○ Southern (2012) evaluated the effectiveness and safety of macrolide antibiotics. All the
4 included studies used low dose Azithromycin (lower than the therapeutic range for
5 antimicrobial action), and they were not relevant for this review.

6 Seven further systematic reviews were identified. Six of them (Cai 2011, Carr 2004, Florescu
7 2009, Littlewood 2012, Maiz 2013 and Mukhopadhyay 1996) were assessed as low quality
8 according to AMSTAR, and were therefore excluded from our review. The included papers
9 were checked for inclusion. Utteley (2013) was excluded as it reported the same data as the
10 TA report.

11 In addition, 8 primary studies have also been identified (Assael 2013, Flume 2016, Galeva
12 2013, Konstan 2011/ EVOLVE trial, Retsch-Bogart 2009, Schuster 2013, Trapnell 2012,
13 Wainwright 2011).

14 The size of the studies ranged between 16 and 520 people. Thirteen studies included
15 children, young people and adults (Assael 2013, Chuchalin 2007, COLO/DPI/02/05,
16 COLO/DPI/02/06, Flume 2016, Galeva 2013, Jensen 1987, Konstan 2011/EVOLVE trial,
17 Konstan 2011a/EAGER trial, Lenoir 2007, McCoy 2008, Retsch-Bogart 2009, Schuster
18 2013), 2 studies included children and young people (Murphy 2004, Wainwright 2011), 1
19 included young people and adults (Hodson 2002), 3 studies included adults only (Ramsey
20 1999, Sheldon 1993, Trapnell 2012). 1 study (Ramsey 1993) did not report the age range,
21 the mean age was 17.7 years.

22 Two studies were conducted in the UK (COLO/DPI/02/05, Hodson 2002), 6 in the USA
23 (Flume 2016, Murphy 2004, Retsch-Bogart 2009, Ramsey 1993, Ramsey 1999, Trapnell
24 2012), 1 in Canada (Sheldon 1993), 1 in Denmark (Jensen 1987), 10 studies in multiple
25 countries: 1 in the EU, Russia and Ukraine (COLO/DPI/02/06), 1 in Europe and the United
26 States (Assael 2013), 1 in Hungary, Poland and Russia (Chuchalin 2007), 1 in Bulgaria,
27 Estonia, Latvia, Lithuania, Romania, Russia, Egypt, and India (Galeva 2013), 1 in Bulgaria,
28 Lithuania, Serbia, Argentina, Brazil, Chile, Mexico and the United States (Konstan
29 2011/EVOLVE trial), 1 in 15 unspecified countries (Konstan 2011a/EAGER trial), 1 in France,
30 Italy, Moldova, Ukraine (Lenoir 2007), 1 in Australia, Canada, New Zealand and the United
31 States (McCoy 2008), 1 in Europe (Schuster 2013), 1 in Australia and the United States
32 (Wainwright 2011).

33 The included studies evaluated their effectiveness based on the following comparisons:

- 34 • Aztreonam lysine vs placebo – 3 studies (McCoy 2009, Retsch-Bogart 2009, Wainwright
35 2011)
- 36 • Ciprofloxacin vs placebo – 1 study (Sheldon 1993)
- 37 • Colistin vs placebo – 1 study (Jensen 1987)
- 38 • Colistin inhalation powder vs colistin inhalation solution – 1 study (COLO/DPI/02/05)
- 39 • Colistin vs tobramycin – 3 studies (Hodson 2002, COLO/DPI/02/06, Schuster 2013)
- 40 • Tobramycin vs placebo – 6 studies (Chuchalin 2007, Galeva 2013, Konstan
41 2011/EVOLVE trial, Lenoir 2007, Ramsey 1993, Ramsey 1999)
- 42 • Tobramycin inhalation powder vs tobramycin inhalation solution – 1 study (Konstan
43 2011a/EAGER trial)
- 44 • Tobramycin vs Aztreonam lysine – 1 study (Assael 2013)
- 45 • Fosfomycin + tobramycin vs placebo – 1 study (Trapnell 2012)
- 46 • Tobramycin + azithromycin vs tobramycin alone (Flume 2016)
- 47 • Continuous alternating therapy vs intermittent treatment: aztreonam lysine + tobramycin or
48 placebo + tobramycin (Flume 2016)

1 A report from the National Horizon Scanning Centre (NHSC 2010) was also identified. This
2 report included 5 trials that have been retrieved for assessment.

3 The presentation of evidence synthesis will be divided in 2 parts based on the type of
4 analysis which was used to produce these syntheses:

5 A network meta-analysis was conducted for the treatment of chronic *P Aeruginosa*. It
6 included the critical outcomes listed in the protocol: lung function (FEV₁) and number of
7 people with ≥ 1 exacerbations. The results for these will be provided at a later stage.

8 Pairwise comparisons have been performed for the rest of the outcomes included in the
9 review protocol and are presented in this review.

109.4.3.1.2 **S Aureus**

11 The interventions that were included in the protocol for the treatment of chronic infection with
12 *S aureus* were: Cefradine (oral), Cotrimoxazole (oral), Doxycycline (oral) and Flucloxacillin
13 (oral).

14 One Cochrane review (Lo 2015) was identified for potential inclusion. This review aimed to
15 evaluate the effectiveness of antimicrobial treatment regimens to eradicate methicillin-
16 resistant *S aureus* (MRSA) in people with cystic fibrosis and all disease severities. No trials
17 were identified for inclusion. The list of excluded studies was also checked. None of the 48
18 excluded studies were relevant.

19 No other trials relevant trials were identified in our search.

209.4.3.1.3 **B Cepacia Complex**

21 The interventions that were included in the protocol for the treatment of chronic infection with
22 *B cepacia* complex were: Ceftazidime (inhaled), Cotrimoxazole (oral), Imipenem,
23 Meropenem (inhaled), and Trimethoprim (oral).

24 One Cochrane review (Ryan 2011) was identified for potential inclusion. This review included
25 trials that evaluated the effectiveness of inhaled antibiotics for long-term therapy in cystic
26 fibrosis and included one cross-over trial that looked at people with cystic fibrosis infected
27 with *B cepacia*. This study was not considered for inclusion in our review as it assessed the
28 effectiveness of inhaled Taurolidine, an intervention that is not routinely used in clinical
29 practice.

30 No other trials relevant trials were identified in our search.

319.4.3.1.4 **A Fumigatus**

32 The interventions that were included in the protocol for the treatment of chronic infection with
33 *A fumigatus* were: Amphotericin (inhaled), Itraconazole (oral), Posaconazole (oral) and
34 Voriconazole (oral).

35 One Cochrane review (Elphick 2014) was identified for potential inclusion. This review
36 evaluated the effectiveness of antifungal interventions for the treatment of allergic
37 bronchopulmonary aspergillosis (ABPA) in people with cystic fibrosis. No trials were
38 identified for inclusion. The list of excluded studies was also checked. One study (Aaron
39 2012) had already been identified in our search and it is included in the review, and the other
40 three studies were not relevant (due to study design or intervention evaluated). This RCT
41 included 35 people with cystic fibrosis over the age of 6 years, and chronically colonised
42 with *A fumigatus*. It was conducted in Canada. It compared the effectiveness of oral
43 Itraconazole versus placebo for a 24-week treatment period. In relation to the outcomes, it
44 included lung function, pulmonary exacerbations, quality of life and adverse events.

1 9.4.3.2 Summary of included studies

2 A summary of the studies that were included in this review are presented in Table 105 to
3 Table 107.

4 **Table 105: Summary of included studies for antimicrobials for chronic pulmonary**
5 **infection with *P aeruginosa***

Study	Intervention/ Comparison	Population	Outcomes	Comments
Technology Appraisal				
NICE TA 276 (Tappenden 2013)	<p>Comparison 1: Tobramycin DPI vs Tobramycin nebulised (EAGER trial)</p> <p>Comparison 2: Colisthimethate sodium DPI vs tobramycin nebulised (COLO/DPI/02/06)</p> <p>Comparison 3: Colisthimethate sodium DPI vs colisthimethate sodium nebulised (Davies 2004, COLO/DPI/02/05)</p>	People with cystic fibrosis \geq 6 years and chronic <i>P aeruginosa</i> pulmonary colonisation.	<ul style="list-style-type: none"> • Lung function, FEV₁% • Frequency and severity of respiratory exacerbations • <i>proxy for Time to next pulmonary exacerbation</i> • Rate and extent of microbiological response (Eradication of the specified organism from sputum/airway cultures) • Quality of life • Adverse events <p>Not reported:</p> <ul style="list-style-type: none"> • Nutritional status • Emergence of resistant organisms/ antibiotic resistance 	All studies are RCTs, open label EAGER trial: Konstan 2011a Other outcomes included in the TA: Respiratory symptoms
Cochrane systematic reviews				
Remington 2016 Cochrane SR	Comparison 1: Ciprofloxacin (500 mg) vs placebo (Sheldon 1993)	People with cystic fibrosis diagnosed by clinical features and all levels of severity of lung disease.	<ul style="list-style-type: none"> • Comparison 1: Ciprofloxacin vs placebo • FEV₁ • Eradication of the organism • Adverse events 	AMSTAR: 11/11
Ryan 2011 Cochrane SR	<p>Comparison 1: Aztreonam lysine vs placebo (McCoy 2008)</p> <p>Comparison 2: Colistin vs placebo (Hodson 2002, Jensen 1987)</p> <p>Comparison 3: Tobramycin vs placebo (Chuchalin 2007, Hodson 2002,</p>	People with cystic fibrosis diagnosed by clinical features and all levels of severity of lung disease.	<p>Comparison 1:</p> <ul style="list-style-type: none"> • Aztreonam lysine vs placebo • FEV₁ %: Included in NMA • Exacerbations: Included in NMA • QoL • Adverse events <p>Not reported:</p> <ul style="list-style-type: none"> • Eradication of the organism • Nutritional status • Emergence of 	AMSTAR: 11/11 *reported in individual studies

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Lenoir 2007, Murphy 2004, Ramsay 1993, Ramsay 1999)		<p>resistant organisms/ antibiotic resistance</p> <p>Comparison 2: Colistin vs placebo</p> <ul style="list-style-type: none"> • FEV₁ %: Included in NMA • Exacerbations: Included in NMA <p>Not reported:</p> <ul style="list-style-type: none"> • Eradication of the organisms* • Nutritional status* • QoL • Adverse events • Emergence of resistant organisms/ antibiotic resistance <p>Comparison 3: Tobramycin vs placebo</p> <ul style="list-style-type: none"> • FEV₁ %: Included in NMA • Exacerbations: Included in NMA • Adverse events <p>Not reported:</p> <ul style="list-style-type: none"> • Eradication of the organisms* • Nutritional status* • QoL • Emergence of resistant organisms/ antibiotic resistance* 	
Primary studies included in the TA or in the Cochrane SR				
Chuchalin 2007 (Hungary, Poland, Russia) RCT	<p>Intervention Tobramycin (nebulised) 300 mg. 24 weeks: 4 weeks “on treatment”, followed by 4 weeks “off treatment”</p> <p>Comparison Placebo</p>	<p>N=247 people with cystic fibrosis and <i>P aeruginosa</i> ≥6 years Age range: 6 to 45</p>	<ul style="list-style-type: none"> • Lung function (FEV₁) • Exacerbations • Nutritional status • Eradication of the organism • Adverse events • Mortality • Emergence of resistant organisms 	<p>Included in Cochrane SR Ryan 2011 Included in NMA and review</p>
COLO/DP I/02/05 (UK) Open label RCT, with	<p>Intervention Colistin sodium DPI 125 mg, twice daily</p> <p>Comparison Colistin sodium</p>	<p>N=16 people with cystic fibrosis with chronic <i>P aeruginosa</i> infection ≥8</p>	<ul style="list-style-type: none"> • Lung function (FEV₁) • Adverse events 	<p>Included in NICE TA 276 Included in the review</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
cross-over	solution 2 MU, twice daily Duration: 8 weeks	years Mean (SD) age: 20.3 (12.87) years		
COLO/DP I/02/06 (EU, Russia, Ukraine) Open label RCT	Intervention Colistin sodium DPI 125 mg; twice daily Comparison Tobramycin inhalation solution 300 mg/ 5 ml; twice daily Duration: 24 weeks	N=380 people with cystic fibrosis and <i>P aeruginosa</i> ≥6 years Mean (SD) age: 21.3 (9.72) vs 20.9 (9.30) years	<ul style="list-style-type: none"> • Lung function (FEV₁) • Time to next exacerbation • Nutritional status • Quality of life • Adverse events 	Included in NICE TA 276 Included in the review
Hodson 2002 (UK) Open label RCT	Intervention Tobramycin (nebulised) 300 mg daily in 5ml twice daily Comparison Colistin (nebulised) 1MU in 3ml in saline twice daily Duration: 28 days	N=126 randomised (n=115 treated) people with cystic fibrosis Age range: 17 to 50 years	<ul style="list-style-type: none"> • Pulmonary function (FEV₁) • Eradication of the organism • Adverse effects • Emergence of resistant organisms 	Included in Cochrane SR Ryan 2011 Included in NMA and review
Jensen 1987 (Denmark) RCT	Intervention Colistin (nebulised) 1 million units, twice daily for 3 months Comparison Placebo (normal saline)	N=40 people with cystic fibrosis and chronic <i>P aeruginosa</i> infection ≥6 years Age range: 7 to 35 years	<ul style="list-style-type: none"> • Lung function (FEV₁) • Eradication of the organism • Emergence of resistant organisms 	Included in Cochrane SR Ryan 2011 Included in NMA and review
Konstan 2011a (EAGER trial) (15 countries, not specified)	Intervention Tobramycin inhalation powder 112 mg, 4-capsules, twice daily Comparison Tobramycin (nebulised) 300mg/ 5 ml, twice daily	N=121 people with cystic fibrosis ≥ 6 years and positive cultures of <i>P aeruginosa</i> within 6 months of screening Mean age (SD): 26 (11.4) vs 25 (10.2)	<ul style="list-style-type: none"> • Lung function (FEV₁) • Exacerbations (hospitalization) • Eradication of the organism • Adverse events 	Included in NICE TA 276 Included in NMA and the review
Lenoir 2007 (France, Italy, Moldova, Ukraine) RCT	Intervention Tobramycin (nebulised) 300 mg twice daily for 4 weeks followed by a 4-week run-out phase	N=59 with cystic fibrosis and <i>P aeruginosa</i> infection ≥6 years Age range: 6 to 30 years	<ul style="list-style-type: none"> • Lung function (FEV₁) • Eradication of the organism • Adverse effects 	Included in Cochrane SR Ryan 2011 Included in NMA and review

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Comparison Placebo			
McCoy 2008 (Australia, Canada, New Zealand and USA) RCT	Intervention Aztreonam lysine 75 mg, for 4 weeks, 2 or 3- times daily Comparison Placebo (5 mg lactose in 1 ml 0.17% sodium chloride)	N=246 people with cystic fibrosis and documented <i>P aeruginosa</i> infection Age range: 7 to 65 years	<ul style="list-style-type: none"> • FEV₁ • Time to next exacerbation • Adverse events • Note: FEV₁ was not included in the review as it was reported narratively only, and could not be meta-analysed 	Included in Cochrane SR Ryan 2011 Included in NMA and review
Murphy 2004 (USA) Open label RCT	Intervention Tobramycin (nebulised), 300 mg twice daily. Alternating 4-weekly cycles for 56 weeks Comparison No treatment	N=184 children and young people with cystic fibrosis with ≥2 cultures of <i>P aeruginosa</i> Age range: 6 to 15 years	<ul style="list-style-type: none"> • Exacerbations (hospitalization) 	Included in NMA only
Ramsey 1993 (USA) Cross- over RCT	Intervention Tobramycin (nebulised) 600 mg, 3-times daily for 28 days, then cross-over for 2 28- days periods Comparison Placebo (0.5 normal saline)	N=71 people with cystic fibrosis and <i>P aeruginosa</i> sputum culture susceptible to tobramycin Mean age (SD): 17.7 years (1.25)	<ul style="list-style-type: none"> • Lung function (FEV₁) • Exacerbations 	Included in Cochrane SR Ryan 2011 Included in NMA and review
Ramsey 1999 (USA) RCT	Intervention Tobramycin (nebulised) 300mg twice daily for three 28-day on-off cycles Comparison Placebo (0.225 normal saline and 1.25 mg quinine)	N=520 people with cystic fibrosis infected with <i>P aeruginosa</i> Age: 18 years or older	<ul style="list-style-type: none"> • Lung function (FEV₁) • Exacerbations (hospitalization) • Adverse events • Note: FEV₁ was not included in the review as it was reported narratively only, and could not be meta-analysed 	Included in Cochrane SR Ryan 2011 Included in NMA and review
Sheldon 1993 (Canada) RCT	Intervention Ciprofloxacin 500 mg, for 10 days every 3 months for 4 courses Comparison Placebo	N=40 adults with cystic fibrosis and chronically infected with <i>P aeruginosa</i> (31 completed the trial) Age ≥ 18 years	<ul style="list-style-type: none"> • Weight • Adverse events • Mortality • Emergence of resistant organisms 	Included in Cochrane SR Remington 2013 Included in review and review
Additional primary studies				

Study	Intervention/ Comparison	Population	Outcomes	Comments
Assael 2013 (Europe and USA) Open label RCT	Intervention Aztreonam lysine 28-day course x 3 Comparison Tobramycin (inhaled) 28 days course, 3000 mg, 2- times/day	N=273 people with cystic fibrosis ≥6 years and PA-positive sputum culture within the previous 3 months Mean age (SD): 25.5 years (9.0)	<ul style="list-style-type: none"> • Pulmonary function (FEV₁) • Exacerbations (requiring i.v. and/or additional antibiotics for respiratory events) • Eradication of the organism • Adverse effects 	Open label Included in NMA and review
Flume 2016 (USA) RCT	Intervention Aztreonam lysine plus tobramycin (nebulised) Comparison Tobramycin (nebulised) Enrolled subjects received TIS 300 mg twice daily (BID) during a 28-day run-in phase This was followed by randomization to a 24- week comparative phase. Subjects received 3 cycles of 28-days of double-blind AZLI or placebo (1:1 randomization) alternating with 28- days of open-label TIS.	N=88 people with cystic fibrosis ≥ 6 years and documented <i>P aeruginosa</i> infection Mean age (SD): 28.4 years (11.4)	<ul style="list-style-type: none"> • Exacerbations (requiring i.v. and/or additional antibiotics for respiratory events, time to first event and rate of hospitalisations) • Adverse effects • Adjusted mean cystic fibrosisQ-R scores averaged from weeks 4, 12, and 20 • Adjusted mean FEV₁% predicted average from weeks 4, 12 and 20 	Included in NMA and review
Galeva 2013 (EDIT trial) (Bulgaria, Estonia, Latvia, Lithuania, Romania, Russia, Egypt, and India) RCT	Intervention Tobramycin inhalation powder 112mg twice daily, as capsules administered via the T-326 dry powder inhaler Comparison Placebo	N=62 people with cystic fibrosis ≥6 years, and a positive sputum or throat culture for P.A within 6 months of screening and positive sputum culture for P.A at the screening visit Mean age (SD), years 12.9 (4.3) vs. 12.9 (4.7)	<ul style="list-style-type: none"> • Pulmonary function (FEV₁) • Exacerbations (hospitalization) • Adverse effects 	Included in NMA and review
Konstan 2011 (EVOLVE trial) (Bulgaria, Lithuania,	Intervention Tobramycin inhalation powder (112 mg), 28 day cycle followed by 2 28-day cycles open- label tobramycin	N=95 children, young people and adults with cystic fibrosis with a positive sputum or throat	<ul style="list-style-type: none"> • Pulmonary function (FEV₁) • Suppression of the organism • Adverse events 	Included in NMA and review

Study	Intervention/ Comparison	Population	Outcomes	Comments
Serbia, Argentina, Brazil, Chile, Mexico, USA) RCT	inhalation powder Comparison Placebo, 28-day cycle followed by 2 28-day cycles open-label tobramycin inhalation powder Total duration 24 weeks	culture for <i>P aeruginosa</i> within 6 months of screening and a positive sputum culture for <i>P aeruginosa</i> at the screening visit Age: 6 to 21 years		
Retsch- Bogart 2009 (USA) RCT	Intervention Aztreonam lysine 75mg aztreonam, 52.5mg of lysine monohydrate Comparison Placebo (5mg lactulose)	N=164 people with cystic fibrosis and <i>P aeruginosa</i> documented infection ≥ 6 years Mean age (range): 31.7 (11-74); 27.4 (7- 54)	<ul style="list-style-type: none"> • Exacerbations (hospitalization) • Eradication of the organism • Quality of life • Adverse effects 	Included in NMA and review
Schuster 2013 (Europe - countries not specified) Open label RCT	Intervention Colistin DPI 1.6625 MU twice daily, 24 weeks Comparison Tobramycin inhalation solution 300 mg/5 ml twice- daily, three 28-day cycles	N=380 people with cystic fibrosis chronically colonised with <i>P aeruginosa</i> infection, ≥ 6 years Mean (SD) age: 21.1 (9.49) years	<ul style="list-style-type: none"> • FEV₁% • Adverse events 	Open label Included in NMA and review
Trapnell 2012 (USA) RCT	Intervention Fosfomycin/ Tobramycin (160/ 40 mg or 80/ 20 mg) Comparison Placebo	N=119 people with cystic fibrosis ≥ 18 years and confirmed <i>P aeruginosa</i> infection <ul style="list-style-type: none"> • 80/ 20 mg: n=38 • 160/ 40 mg: n=41 • Placebo: n=40 Mean age: 32 years (10.1)	<ul style="list-style-type: none"> • FEV₁ • Exacerbations (hospitalization) 	Included in NMA and review
Wainwrig ht 2011 (Australia, USA) RCT	Intervention Aztreonam lysine 28 days + 14 days follow-up 75 mg/ day, 3-times	N=157 people with cystic fibrosis ≥ 6 years Age range: 6 to 17 years	<ul style="list-style-type: none"> • Exacerbations (hospitalization) • Eradication of the organism • Adverse effects 	Included in NMA and review

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Comparison Placebo			

cystic fibrosis: cystic fibrosis; cystic fibrosisQ-R: Cystic Fibrosis Questionnaire-Revised; DPI: dry powder for inhalation; FEV₁: forced expiratory volume; ITT: intention to treat analysis; RCT: randomized controlled trial

Table 106: Summary of included studies for antimicrobials for chronic pulmonary infection with *S aureus*

Study	Intervention/ Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Ahmed 2016 Cochrane SR	Intervention Any combination of topical, inhaled, oral or IV antimicrobials used with the objective of suppressive therapy for chronic Comparison Placebo or no treatment	• Children and adults with confirmed diagnosis of cystic fibrosis and confirmed microbiological evidence of <i>S aureus</i> (MSSA strains only).	No studies were identified for inclusion in this review	AMSTAR score:10/11
Lo 2015 Cochrane SR	Intervention Any combination of topical, inhaled, oral or IV antimicrobials to eradicate MRSA Comparison Placebo, standard treatment or not treatment	Children and adults with cystic fibrosis with a confirmed positive microbiological isolate of methicillin-resistant <i>S aureus</i> (MRSA).	No studies were identified for inclusion in this review	This review includes people with different disease severity.

Table 107: Summary of included studies for antimicrobials for chronic pulmonary infection with *A fumigatus*

Study	Intervention/ Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Elphick 2014 Cochrane SR	Intervention Antifungal treatments, including major treatments such as oral azoles and nebulised amphotericin Comparison No treatment, placebo or different dosages	Children and adults with cystic fibrosis and allergic bronchopulmonary aspergillosis.	No studies were identified for inclusion in this review	
Additional primary studies				
Aaron	Intervention	N=35	• Lung function,	(+) low risk of

Study	Intervention/ Comparison	Population	Outcomes	Comments
2012 (Canada) RCT	Itraconazole capsules daily dose of 5 mg/kg once daily; if the dose exceeded 200 mg/day it was given twice daily Comparison Placebo	People with cystic fibrosis ≥ 6 years of age and chronically colonised with <i>A. Fumigatus</i> (defined as at least 2 positive sputum cultures within the last 12 months)	measured as FEV ₁ % predicted at 24 and 48 weeks <ul style="list-style-type: none"> • Time to next pulmonary exacerbation • <i>proxy outcomes</i>: <ul style="list-style-type: none"> ○ number of patients that experienced pulmonary exacerbations requiring oral or IV AB ○ number of patients that experienced pulmonary exacerbations requiring hospitalization • Quality of life, measured with the tool cystic fibrosisQ-R at 24 weeks • Adverse events, reported during the 24-week study duration Not reported: <ul style="list-style-type: none"> • Eradication of the specified organism from sputum/airway cultures • Nutritional status • Emergence of resistant organisms/ antibiotic resistance 	bias for sequence generation, blinding and selective reporting (+) first prospective RCT (+) ITT analysis (?) unclear risk of bias for allocation concealment (-) pilot study small sample size, authors failed to recruit more patients to extend the study (-) Failure to achieve therapeutic levels of Itraconazole in many patients

1 9.4.3.3 Clinical evidence profile

2 The clinical evidence profiles for this review question are presented separately for each
3 pathogen.

49.4.3.3.1 *P aeruginosa*

5 Results from the NMA and pairwise comparisons are presented separately in this section.

69.4.3.3.2 **Clinical evidence profile for NMA outcomes (FEV₁ % predicted and number of participants experiencing at least one pulmonary exacerbation)**

8 As treatment effects were found to vary over time, NMAs were conducted separately for
9 short (4-10 weeks) and long (>10 weeks) of treatment.

- 10
- FEV₁ % Predicted


11 Nine studies of 7 treatments tested in 1346 participants were included in the review. Due to
12 very high unexplained heterogeneity between studies, it was felt that the studies should not

1 be meta-analysed. Therefore the studies have been evaluated individually, and are reported
2 without pooled effects (section 9.4.3.4).

- 3 • Number of patients experiencing at least one pulmonary exacerbation – Short-term
4 treatment (4-10 weeks)

5 Three studies of 354 participants were included in the network of 4 interventions (placebo,
6 aztreonam lysine (nebulised), tobramycin (nebulised), tobramycin plus fosfomycin
7 (nebulised)) (Figure 6). The evidence for this analysis was of low quality. For all three studies
8 the risk of bias was unclear.

9 **Figure 6: Network for number of patients experiencing at least one exacerbation with**
10 **short-term (4-10 weeks) treatment**



11 *Source/Note: Size of nodes are proportional to the number of patients in the network treated with a particular*
12 *intervention. Thickness of connecting lines are proportional to the number of studies comparing two*
13 *interventions.*
14

15 Table 108 presents the results of the conventional pair-wise meta-analyses (head to head
16 comparisons) (upper-right section of table), together with the results computed by the NMA
17 for every possible treatment comparison (lower-left section of table). Both results are
18 presented as odds ratios (95% CrI). These results were derived from a fixed effects model
19 (see Appendix N – Model Fit).

20 There was considerable uncertainty throughout the network. Aztreonam lysine was found to
21 be significantly more effective than placebo and tobramycin (nebulised) at reducing the odds
22 of experiencing at least one exacerbation. No other significant effects were found.
23 Inconsistency could not be assessed as there were no closed loops of treatments.

24 In this analysis, aztreonam lysine was found to have the highest probability (88.30%) of
25 being the best treatment to reduce the odds of experiencing at least one exacerbation,
26 followed by tobramycin plus fosfomycin (nebulised) (10.00%) (Table 109).

Table 108: Odds ratios (95% CrI) from conventional (white area) and network meta-analysis (grey area) for the number of people experiencing at least one exacerbation with short-term (4-10 weeks) treatment

	Placebo	Aztreonam lysine	Tobramycin (nebulised)	Tobramycin + Fosfomycin (nebulised)
Placebo		0.3 (0.08, 0.92)	3 (0.55, 24.66)	0.9 (0.25, 3.81)
Aztreonam lysine	0.3 (0.08, 0.92)			
Tobramycin (nebulised)	3 (0.55, 24.66)	10.43 (1.31, 122.2)		
Tobramycin + Fosfomycin (nebulised)	0.9 (0.25, 3.81)	3.11 (0.55, 21.38)	0.3 (0.03, 2.78)	

(m) Results in the top right diagonal of the table are the mean differences and 95% CrI from the conventional meta-analyses of direct evidence between the column-defined treatments compared to the row-defined treatment. Mean differences greater than 0 favour the column-defined treatment.

(n) Results in the bottom left are the mean differences and 95% CrI from the NMA model of direct and indirect evidence between the row-defined treatments compared to the column-defined treatments. Mean differences greater than 0 favour the row-defined treatment.

(o) Numbers in bold denote results for which the 95% CrI does not include the null effect of 0

Table 109: Median treatment ranking (with their 95% CrI) of all interventions in the network and the probability of being the best treatment for reducing the number of people experiencing at least one exacerbation with short-term (4-10 weeks) treatment

	Median (95% CrI) treatment rank	Probability of being the best treatment (%)
Placebo	3 (2-4)	0.70%
Aztreonam lysine	1 (1-2)	88.30%
Tobramycin (nebulised)	4 (2-4)	1.00%
Tobramycin + Fosfomycin (nebulised)	2 (1-4)	10.00%

Table 110: Quality assessment of the evidence for the NMA – number of patients with at least one exacerbation in the short-term

NMA	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Quality
Short-term (4-10 weeks) number of patients with at least one exacerbation (3 studies)	Serious ¹	No serious inconsistency	No serious indirectness	Serious ²	None	Low


(p) 1 - For all three studies the risk of bias was unclear

(q) 2 – No intervention has rank credible intervals $\leq 33\%$ of total distribution of comparators

- Number of patients experiencing at least one pulmonary exacerbation – Long-term treatment (>10 weeks)

1 Six studies of 1,749 participants were included in the network of 6 interventions (placebo,
2 aztreonam lysine (nebulised), tobramycin (nebulised), no treatment, tobramycin (powder), 28
3 days aztreonam lysine (nebulised) alternating with 28 days tobramycin (nebulised) (Figure
4 7)The evidence for this analysis was of moderate quality. Two studies were at high risk of
5 bias and for the other 4 studies the risk of bias was unclear.

6 **Figure 7: Network for number of patients experiencing at least one exacerbation with**
7 **long-term (>10 weeks) treatment**



8 *Source/Note: Size of nodes are proportional to the number of patients in the network treated with a particular*
9 *intervention. Thickness of connecting lines are proportional to the number of studies comparing two*
10 *interventions.*
11

12 Table 111 presents the results of the conventional pair-wise meta-analyses (head to head
13 comparisons) (upper-right section of table), together with the results computed by the NMA
14 for every possible treatment comparison (lower-left section of table). Both results are
15 presented as odds ratios (95% CrI). These results were derived from a fixed effects model
16 (see Appendix G – Model Fit).

17 Inconsistency could not be assessed as there were no closed loops of treatments.

18 In this analysis, aztreonam lysine was found to have the highest probability (85.01%) of
19 being the best treatment to reduce the odds of experiencing at least one exacerbation,
20 followed by the combination treatment (14.83%) (Table 112).

21 **Table 111: Odds ratios (95% CrI) from conventional (white area) and network meta-**
22 **analysis (grey area) for the number of people experiencing at least one**
23 **exacerbation with long-term (>10 weeks) treatment**

	Placebo	Aztreonam lysine	Tobramycin (nebulised)	No treatment	Tobramycin (powder)	Combination
Placebo			0.88 (0.65, 1.20)			
Aztreonam lysine	0.40 (0.22, 0.71)		2.20 (1.36, 3.61)			
Tobramycin (nebulised)	0.88 (0.65, 1.20)	2.20 (1.36, 3.61)		2.84 (1.28, 6.71)	1.14 (0.75, 1.75)	0.77 (0.33, 1.78)
No	2.51	6.27	2.84			

	Placebo	Aztreonam lysine	Tobramycin (nebulised)	No treatment	Tobramycin (powder)	Combination a
treatment	(1.07, 6.22)	(2.46, 16.81)	(1.28, 6.71)			
Tobramycin (powder)	1.01 (0.60, 1.70)	2.52 (1.33, 4.84)	1.14 (0.75, 1.75)	0.40 (0.16, 1.00)		
Combination a	0.67 (0.27, 1.65)	1.68 (0.63, 4.48)	0.77 (0.33, 1.78)	0.27 (0.08, 0.86)	0.67 (0.26, 1.72)	

(r) Results in the top right diagonal of the table are the mean differences and 95% CrI from the conventional meta-analyses of direct evidence between the column-defined treatments compared to the row-defined treatment. Mean differences greater than 0 favour the column-defined treatment.

(s) Results in the bottom left are the mean differences and 95% CrI from the NMA model of direct and indirect evidence between the row-defined treatments compared to the column-defined treatments. Mean differences greater than 0 favour the row-defined treatment.

(t) Numbers in bold denote results for which the 95% CrI does not include the null effect of 0

(u) (a) 28 days aztreonam lysine (nebulised) alternating with 28 days tobramycin (nebulised)

Table 112: Median treatment ranking (with their 95% CrI) of all interventions in the network and the probability of being the best treatment for reducing the number of people experiencing at least one exacerbation with long-term (>10 weeks) treatment

	Median (95% CrI) treatment rank	Probability of being the best treatment (%)
Placebo	4 (2, 5)	0.03%
Aztreonam lysine	1 (1, 2)	85.01%
Tobramycin (nebulised)	3 (2, 5)	0.01%
No treatment	6 (5, 6)	0.00%
Tobramycin (powder)	4 (2, 5)	0.12%
Combination a	2 (1, 5)	14.83%

(v) 28 days aztreonam lysine (nebulised) alternating with 28 days tobramycin (nebulised)

Table 113: Quality assessment of the evidence for the NMA – number of patients with at least one exacerbation in the long-term

NMA	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Quality
Long-term (>10 weeks) number of patients with at least one exacerbation (6 studies)	Serious1	No serious inconsistency	No serious indirectness	No serious imprecision	None	Moderate

(w) 1 – For two studies the risk of bias was high and for four studies the risk of bias was unclear

9.4.3.4 Clinical evidence profile for non-NMA outcomes

The summary clinical evidence profiles for this review question are presented in Table 114 to Table 123.

Table 114: Summary clinical evidence profile: Comparison 1. Aztreonam lysine versus placebo

Comparison 1.: Aztreonam versus placebo

Comparison 1.: Aztreonam versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Aztreonam lysine				
Lung function: relative change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 28 days	The mean relative change in FEV ₁ % predicted at 28 days follow up in the placebo group was -2.5	The mean relative change in FEV ₁ % predicted at 28 days follow up in the aztreonam lysine groups was 2.79 higher (0.48 to 5.1 higher)		157 (Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	
Number of patients with 1 or more exacerbations	NMA outcome					
Suppression of the organism: adjusted mean change sputum density log ₁₀ cystic fibrosisU/g Follow-up: 28 days	Not reported	The mean adjusted mean change sputum density in the aztreonam lysine groups was 1.40 lower (1.94 lower to 0.85 higher)		321 (Retsch-Bogart 2009, Wainwright 2011)	⊕⊕⊕⊕ high	
Nutritional status % weight change (kg) Follow-up: 28 days	The mean % weight change (kg) in the placebo groups ranged between 1 and 1.1	The mean % weight change (kg) in the aztreonam lysine groups was 1 higher (0.33 to 1.67 higher)		164 (Retsch-Bogart 2009)	⊕⊕⊕⊕ high	
Quality of life: change in cystic fibrosisQ-R body image Follow-up: 28 days	The mean change in cystic fibrosisQ-R body image in the placebo groups ranged between 1 and 1.1	The mean change in cystic fibrosisQ-R body image in the aztreonam lysine groups was 2.44 higher (0.35 lower to 5.23 higher)		320 (Retsch-Bogart 2009, Wainwright 2011)	⊕⊕⊕⊖ moderate ¹	
Quality of life: change in cystic fibrosisQ-R	The mean change in cystic	The mean change in cystic		321 (Retsch-Bogart	⊕⊕⊕⊕ high	

Comparison 1.: Aztreonam versus placebo						
digestion Follow-up: 28 days	fibrosisQ-R digestion in the placebo groups ranged between 1.9 and 5.5	fibrosisQ-R digestion in the aztreonam lysine groups was 0.45 lower (3.53 lower to 2.63 higher)		2009, Wainwrig t 2011)		
Quality of life: change in cystic fibrosisQ-R eating Scale from: 0 to 100 Follow-up: 28 days	The mean change in cystic fibrosisQ-R eating in the placebo groups ranged between -4.7 and -1.2	The mean change in cystic fibrosisQ-R eating in the aztreonam lysine groups was 4.99 higher (1.47 lower to 11.46 higher)		321 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R emotional functioning Scale from: 0 to 100 Follow-up: 28 days	The mean change in cystic fibrosisQ-R emotional functioning in the placebo groups ranged between -1.3 and 3.7	The mean change in cystic fibrosisQ-R emotional functioning in the aztreonam lysine groups was 2.36 higher (3.13 lower to 7.84 higher)		320 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R health perceptions Follow-up: 28 days	The mean change in cystic fibrosisQ-R health perceptions in the placebo groups ranged between -4.8 and -1.8	The mean change in cystic fibrosisQ-R health perceptions in the aztreonam lysine groups was 6.82 higher (0.75 to 12.89 higher)		272 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R physical functioning Follow-up: 28 days	The mean change in cystic fibrosisQ-R health perceptions in the placebo groups ranged between -6.9 and -0.69	The mean change in cystic fibrosisQ-R health perceptions in the aztreonam lysine groups was 5.60 higher (0.96 lower to 12.15 higher)		320 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R respiratory	The mean change in cystic fibrosisQ-R	The mean change in cystic fibrosisQ-R		321 (Retsch- Bogart 2009,	⊕⊕⊕⊕ very low ^{1,2}	

Comparison 1.: Aztreonam versus placebo						
symptoms Follow-up: 28 days	respiratory symptoms in the placebo groups ranged between -2.6 and 2.9	respiratory symptoms in the aztreonam lysine groups was 4.81 higher (4.6 lower to 14.21 higher)		Wainwrig t 2011)		
Quality of life: change in cystic fibrosisQ-R role/school Follow-up: 28 days	The mean change in cystic fibrosisQ-R role/school in the placebo groups ranged between -4.2 and 0.29	The mean change in cystic fibrosisQ-R role/school in the aztreonam lysine groups was 2.97 higher (3.2 lower to 9.13 higher)		272 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R social functioning Follow-up: 28 days	The mean change in cystic fibrosisQ-R social functioning in the placebo groups ranged between -3.6 and -2.6	The mean change in cystic fibrosisQ-R social functioning in the aztreonam lysine groups was 3.54 higher (0.78 to 6.31 higher)		319 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ moderate ¹	
Quality of life: change in cystic fibrosisQ-R treatment burden Follow-up: 28 days	The mean change in cystic fibrosisQ-R treatment burden in the placebo groups ranged between -3.1 and 5.1	The mean change in cystic fibrosisQ-R treatment burden in the aztreonam lysine groups was 0.36 lower (7.42 lower to 6.69 higher)		321 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ very low ^{2,3}	
Quality of life: change in cystic fibrosisQ-R vitality Follow-up: 28 days	The mean change in cystic fibrosisQ-R vitality in the placebo groups ranged between -4.4 and -2.2	The mean change in cystic fibrosisQ-R vitality in the aztreonam groups was 5.46 higher (0.16 to 10.76 higher)		272 (Retsch- Bogart 2009, Wainwrig t 2011)	⊕⊕⊕⊕ low ^{1,2}	
Quality of life: change in cystic fibrosisQ-R weight Follow-up: 28	The change in cystic fibrosisQ-R weight in the placebo groups	The change in cystic fibrosisQ-R weight in the aztreonam lysine groups		272 (Retsch- Bogart 2009, Wainwrig	⊕⊕⊕⊕ moderate ¹	

Comparison 1.: Aztreonam versus placebo						
days	ranged between 1.4 and 2.6	was 2.58 higher (2.83 lower to 7.98 higher)		t 2011)		
Minor adverse events: chest discomfort Follow-up: 28 days	48 per 1000	62 per 1000 (18 to 224)	RR 1.31 (0.37 to 4.71)	164 (Retsch-Bogart 2009)	⊕⊕⊕⊕ low ⁴	
Minor adverse events: cough Follow-up: 28 days	Study population		RR 1.10 (0.87 to 1.38)	532 (Retsch-Bogart 2009, McCoy 2009, Wainwright 2011)	⊕⊕⊕⊕ low ⁴	
	340 per 1000	374 per 1000 (296 to 470)				
	Moderate					
	342 per 1000	376 per 1000 (298 to 472)				
Minor adverse events: headache Follow-up: 28 days	Study population		RR 0.94 (0.34 to 2.61)	321 (Retsch-Bogart 2009, Wainwright 2011)	⊕⊕⊕⊕ very low ^{4,6}	
	121 per 1000	114 per 1000 (41 to 316)				
	Moderate					
	121 per 1000	114 per 1000 (41 to 316)				
Major adverse events: dyspnoea Follow-up: 28 days	95 per 1000	63 per 1000 (21 to 183)	RR 0.66 (0.22 to 1.92)	164 (Retsch-Bogart 2009)	⊕⊕⊕⊕ low ⁴	
Major adverse events: haemoptysis Follow-up: 28 days	Study population		RR 0.86 (0.44 to 1.7)	375 (Retsch-Bogart 2009, McCoy 2009)	⊕⊕⊕⊕ low ⁴	
	94 per 1000	81 per 1000 (41 to 159)				
	Moderate					
	94 per 1000	81 per 1000 (41 to 160)				
Mortality Follow-up: 28 days	No events	No events	-	211 (McCoy 2009)	⊕⊕⊕⊕ high	
Emergence of resistant organisms: persistent isolation of <i>S aureus</i> Follow-up: 42 days	62 per 1000	27 per 1000 (6 to 135)	RR 0.44 (0.09 to 2.19)	155 (Retsch-Bogart 2009)	⊕⊕⊕⊕ moderate ⁵	
Emergence of resistant organisms : persistent isolation of <i>B cepacia</i> Follow-up: 42 days	No events	No events	-	155 (Retsch-Bogart 2009)	⊕⊕⊕⊕ high	
Emergence of	0 per 1000	0 per 1000	RR 5.47	155	⊕⊕⊕⊕	

Comparison 1.: Aztreonam versus placebo						
resistant organisms: persistent isolation of <i>S. maltophilia</i> Follow-up: 42 days		(0 to 0)	(0.27 to 112.04)	(Retsch-Bogart 2009)	low ⁴	
Emergence of resistant organisms: persistent isolation of <i>A. xilosidans</i> Follow-up: 42 days	25 per 1000	14 per 1000 (1 to 146)	RR 0.55 (0.05 to 5.91)	155 (Retsch-Bogart 2009)	⊕⊕⊕⊕ low ⁴	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: cystic fibrosisQ-R: cystic fibrosis questionnaire revised; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 2 The quality of the evidence was downgraded by 1 or by 2 due to the moderate of high heterogeneity in the different cystic fibrosisQ-R domains (eating I²=79%; emotional functioning I²=80%; health perceptions I²=62%; respiratory symptoms I²=85%; role/ school I²=73%; treatment burden I²=79%; vitality I²=40%)
- 3 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
- 4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs
- 5 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 6 The quality of the evidence was downgraded by 2 due to high heterogeneity (I²=62%)

Table 115: Summary clinical evidence profile: Comparison 2. Ciprofloxacin versus placebo

Comparison 2.: Ciprofloxacin versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Ciprofloxacin				
Lung function	Not reported					
Number of people with one or more exacerbations	NMA outcome					
Nutritional status: weight kg Follow-up: 6 to 12 months	The mean weight in the placebo group was 51.3	The mean change in weight in the ciprofloxacin groups was 4.4 higher (3.7 lower to 12.5 higher)		31 (Sheldon 1993)	⊕⊕⊕⊕ very low ^{1,2}	
Minor adverse events - gastrointestinal Follow-up: 12 months	0 per 1000	0 per 1000 (0 to 0)	RR 5 (0.26 to 98)	40 (Sheldon 1993)	⊕⊕⊕⊕ very low ^{3,4}	
Mortality	50 per	50 per 1000	RR 1	40	⊕⊕⊕⊕	

Comparison 2.: Ciprofloxacin versus placebo						
Follow-up: 12 months	1000	(4 to 745)	(0.07 to 14.9)	(Sheldon 1993)	low ⁵	
Emergence of resistant organisms - isolation of resistant strains of <i>P aeruginosa</i> Follow-up: 12 months	312 per 1000	666 per 1000 (297 to 1000)	RR 2.13 (0.95 to 4.8)	(Sheldon 1993)	⊕⊕⊕⊕ very low ^{1,2}	
Emergence of resistant organisms - isolation of resistant strains of <i>S aureus</i> Follow-up: 12 months	375 per 1000	266 per 1000 (94 to 761)	RR 0.71 (0.25 to 2.03)	31 (Sheldon 1993)	⊕⊕⊕⊕ very low ^{1,4}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio

- 1 The quality of the evidence was downgraded by 2 due to unclear blinding and reporting and high loss to follow-up
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 3 The quality of the evidence was downgraded by 1 due to unclear blinding and reporting
- 4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs
- 5 The quality of the evidence was downgraded by 2 as the 95% CI crossed the line of null effect, and the CI is very wide (trial underpowered to detect a difference)

8 **Table 116: Summary clinical evidence profile: Comparison 3.1. Colistin versus**
9 **placebo**

Comparison 3.1.: Colistin versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Colistin				
Lung function: change in FEV ₁ % predicted Follow-up: 3 months	The mean change in FEV ₁ % predicted in the placebo groups was -17	The mean change in FEV ₁ % predicted in the colistin groups was 6.00 higher (1.07 lower to 13.07 higher)		40 (Jensen 1987)	⊕⊕⊕⊕ low ^{1,2}	
Number of patients with 1 or more exacerbations	NMA outcome					
Suppression of the organism: eradication of <i>P aeruginosa</i> from the sputum Follow-up: 3 months	PA was not eradicated from the sputum of any patient during the 3-month trial		-	40 (Jensen 1987)	⊕⊕⊕⊕ moderate ^{1,3}	

Comparison 3.1.: Colistin versus placebo					
Emergence of resistant organisms: superinfection with other colistin-resistant organisms Follow-up: 3 months	No super infection with other colistin-resistant microorganisms, including Ps. Cepacia, Serratia Macencens, Proteus mirabilis, Gram-positive organisms or funghi during the 3-month trial	-	40 (Jensen 1987)	⊕⊕⊕⊕ moderate ^{1,3}	
Emergence of resistant organisms: resistance to colistin Follow-up: 3 months	Resistance to colistin did not develop in any strain during the 3-month trial	-	40 (Jensen 1987)	⊕⊕⊕⊕ moderate ^{1,3}	
Emergence of resistant organisms: resistance to other commonly used anti-pseudomonas treatment Follow-up: 3 months	No change in resistance pattern to other commonly used anti-pseudomonas during the 3-month trial	-	40 (Jensen 1987)	⊕⊕⊕⊕ moderate ^{1,3}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
CI: Confidence interval;					

- 1 The quality of the evidence was downgraded by 1 due to unclear randomization, allocation and blinding methods. Poor reporting.
2 The quality of the evidence was downgraded by 1 due to serious imprecision, as the 95% CI crossed 1 clinical MID
3 Not calculable, as data reported narratively only.

Table 117: Summary clinical evidence profile: Comparison 3.2. Colistin inhalation powder versus colistin inhalation solution

Comparison 3.2. Colistin inhalation powder versus colistin inhalation solution						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Colistin inhalation solution (COLI neb)	Colistin inhalation powder (COLI DPI)				
Lung function: % mean change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 4 weeks	Not reported	The % mean change in FEV ₁ % predicted in the COLI DPI groups was 3.01 lower (18.71 lower to 12.69 higher)		31 (COLO/DPI /02/05)	⊕⊕⊕⊕ very low ^{1,2}	
Number of patients with 1 or more exacerbations	NMA outcome					
Minor adverse events: vomiting	0 per 1000	0 per 1000 (0 to 0)	RR 4.71 (0.24 to	31 (COLO/DPI	⊕⊕⊕⊕ very	

Comparison 3.2. Colistin inhalation powder versus colistin inhalation solution						
Follow-up: 8 weeks			90.69)	/02/05)	low ^{1,3}	
Minor adverse events: productive cough Follow-up: 8 weeks	67 per 1000	125 per 1000 (13 to 1000)	RR 1.88 (0.19 to 18.6)	31 (COLO/DPI /02/05)	⊕⊕⊕⊕ very low ^{1,3}	
Minor adverse events: chest discomfort Follow-up: 8 weeks	133 per 1000	251 per 1000 (53 to 1000)	RR 1.88 (0.4 to 8.78)	31 (COLO/DPI /02/05)	⊕⊕⊕⊕ very low ^{1,3}	
Serious adverse events - AE: dyspnoea Follow-up: 8 weeks	267 per 1000	187 per 1000 (51 to 701)	RR 0.7 (0.19 to 2.63)	31 (COLO/DPI /02/05)	⊕⊕⊕⊕ very low ^{1,3}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; RR: Risk ratio;						

- 1 The quality of the evidence was downgraded by 1 as this is an open trial, and the randomization is unclear
 2 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs
 3 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs

4 **Table 118: Summary clinical evidence profile: Comparison 3.3. Colistin versus**
 5 **Tobramycin**

Comparison 3.3. Colistin versus Tobramycin						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Tobramycin	Colistin				
[COLI nebulised versus TOBI nebulised] Lung function: mean % change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1 to 3 months	The mean % change in FEV ₁ % predicted in the tobramycin groups was 6.7	The mean % change in FEV ₁ % predicted in the intervention groups was 6.33 lower (12.7 lower to 0.04 higher)		109 (Hodson 2002)	⊕⊕⊕⊕ very low ^{1,2}	
[COLI DPI versus TOBI nebulised] Lung function: mean % change in FEV ₁ % predicted Scale from: 0 to 100	Not reported	The mean % change in FEV ₁ % predicted in the intervention groups was 1.67 lower (5.43 lower to 2.09 higher)		374 (COLO/DPI/02/06)	⊕⊕⊕⊕ low ^{2,3}	

Comparison 3.3. Colistin versus Tobramycin						
Follow-up: 4 weeks						
[COLI DPI versus TOBI nebulised] Lung function: mean % change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 12 weeks	Not reported	The mean % change in FEV ₁ % predicted in the intervention groups was 2.63 lower (6.67 lower to 1.41 higher)		374 (COLO/D PI/02/06)	⊕⊕⊕⊖ low ^{2,3}	
[COLI versus TOBI] Lung function: mean % change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 24 weeks	Not reported	The mean % change in FEV ₁ % predicted in the intervention groups was 0.99 higher (0.95 to 1.03 higher)		658 (COLO/D PI/02/06, Schuster 2013)	⊕⊕⊕⊖ low ⁴	
Number of patients with 1 or more exacerbations	NMA outcome					
[COLI DPI versus TOBI nebulised] Time to next pulmonary exacerbation: time to first additional anti <i>P aeruginosa</i> treatment (days)	The mean time to first additional anti <i>P aeruginosa</i> treatment in the tobramycin groups was 51.79 days	The mean time to first additional anti <i>P aeruginosa</i> treatment in the colistin groups was 3.49 higher (5.14 lower to 12.12 higher)		374 (COLO/D PI/02/06)	⊕⊕⊕⊖ very low ^{3,5}	
[COLI DPI versus TOBI nebulised] Suppression of the organism: change in sputum PA density Log ₁₀ cystic fibrosisU/ml Follow-up: 4 weeks	The mean change in sputum pa density log ₁₀ cystic fibrosisU/ML in the tobramycin groups was -0.79	The mean change in sputum pa density log ₁₀ cystic fibrosisU/ML in the colistin groups was 0.32 higher (0.32 lower to 0.96 higher)		79 (COLO/D PI/02/06)	⊕⊕⊕⊖ low ¹	
[COLI DPI versus TOBI nebulised] Nutritional status: BMI change kg	The mean BMI change in the colistin groups was 0.17	The mean BMI change in the colistin groups was 0.09 lower (0.26 lower to 0.88 higher)		374 (COLO/D PI/02/06)	⊕⊕⊕⊖ low ^{3,6}	

Comparison 3.3. Colistin versus Tobramycin						
Follow-up: 24 weeks						
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R physical Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life cystic fibrosisQ-R physical in the tobramycin groups was -1.56	The mean change in quality of life: cystic fibrosisQ-R physical in the colistin groups was 1.82 higher	P=0.353	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R vitality Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R vitality in the tobramycin groups was -1.40	The mean change in quality of life: cystic fibrosisQ-R vitality in the colistin groups was 2.27 higher	P=0.293	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R emotion Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R emotion in the tobramycin groups was 0.47	The mean change in quality of life: cystic fibrosisQ-R emotion in the colistin groups was 1.75 higher	P=0.244	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R eating Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R eating in the tobramycin groups was 0.66	The mean change in quality of life: cystic fibrosisQ-R eating in the colistin groups was 0.19 lower	P=0.925	372 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R treatment burden in the control	The mean change in quality of life: cystic fibrosisQ-R treatment burden in the control	The mean change in quality of life: cystic fibrosisQ-R treatment burden in the colistin groups was	P=0.091	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	

Comparison 3.3. Colistin versus Tobramycin						
burden Scale from: 0 to 100. Follow-up: 24 weeks	tobramycin was 2.75	2.87 higher				
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R health perception Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R health perception in the tobramycin groups was -2.71	The mean change in quality of life: cystic fibrosisQ-R health perception in the intervention groups was 2.96 higher	P=0.159	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3,7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R social Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R social in the tobramycin groups was 2.18	The mean change in quality of life: cystic fibrosisQ-R social in the colistin groups was 0.92 higher	P=0.153	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3,7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R body image Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in change in quality of life: cystic fibrosisQ-R body image in the tobramycin groups was 5.98	The mean change in quality of life: cystic fibrosisQ-R body image in the colistin groups was 1.85 higher	P=0.385	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3,7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R role Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R role in the tobramycin groups was 1.87	The mean change in quality of life: cystic fibrosisQ-R role in the colistin groups was 1.22 lower	P=0.607	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3,7	
[COLI DPI versus TOBI nebulised] Quality of life: change in	The mean change in quality of life: cystic fibrosisQ-R	The mean change in quality of life: cystic fibrosisQ-R weight in the	P=0.461	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3,7	

Comparison 3.3. Colistin versus Tobramycin						
cystic fibrosisQ-R weight Scale from: 0 to 100. Follow-up: 24 weeks	weight in the tobramycin groups was -1.93	colistin groups was 2.81 higher				
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R respiratory Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R respiratory in the tobramycin groups was 3.51	The mean change in quality of life: cystic fibrosisQ-R respiratory in the colistin groups was 0.53 lower	P=0.756	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI DPI versus TOBI nebulised] Quality of life: change in cystic fibrosisQ-R digestion Scale from: 0 to 100. Follow-up: 24 weeks	The mean change in quality of life: cystic fibrosisQ-R digestion in the tobramycin groups was 2.89	The mean change in quality of life: cystic fibrosisQ-R digestion in the colistin groups was 3.22 higher	P=0.077	374 (COLO/D PI/02/06)	⊕⊕⊕⊖ moderate 3.7	
[COLI nebulised versus TOBI nebulised] Minor adverse events: sputum Follow-up: 4 weeks	113 per 1000	129 per 1000 (48 to 349)	RR 1.14 (0.42 to 3.08)	115 (Hodson 2002)	⊕⊖⊖⊖ very low ^{1,8}	
[COLI nebulised versus TOBI nebulised] Minor adverse events: pharyngitis Follow-up: 4 weeks	132 per 1000	49 per 1000 (13 to 178)	RR 0.37 (0.1 to 1.35)	115 (Hodson 2002)	⊕⊖⊖⊖ very low ^{1,8}	
[COLI nebulised versus TOBI nebulised] Minor adverse events: cough Follow-up: 4 weeks	94 per 1000	177 per 1000 (66 to 478)	RR 1.88 (0.7 to 5.07)	115 (Hodson 2002)	⊕⊖⊖⊖ very low ^{1,8}	
[COLI DPI	228 per	203 per 1000	RR 0.89	380	⊕⊖⊖⊖	

Comparison 3.3. Colistin versus Tobramycin						
versus TOBI nebulised] Minor adverse events: productive cough Follow-up: 24 weeks	1000	(139 to 299)	(0.61 to 1.31)	(COLO/D PI/02/06)	very low ^{3,8}	
[COLI DPI versus TOBI nebulised] Minor adverse events: chest discomfort Follow-up: 24 weeks	176 per 1000	139 per 1000 (86 to 222)	RR 0.79 (0.49 to 1.26)	380 (COLO/D PI/02/06)	⊕⊕⊕⊕ very low ^{3,8}	
[COLI DPI versus TOBI nebulised] Minor adverse events: vomiting Follow-up: 24 weeks	41 per 1000	32 per 1000 (11 to 91)	RR 0.77 (0.27 to 2.19)	380 (COLO/D PI/02/06)	⊕⊕⊕⊕ very low ^{3,8}	
[COLI nebulised versus TOBI nebulised] Serious adverse events: patients with >1 serious AE Follow-up: 4 weeks	151 per 1000	113 per 1000 (44 to 291)	RR 0.75 (0.29 to 1.93)	115 (Hodson 2002)	⊕⊕⊕⊕ very low ^{1,8}	
[COLI DPI versus TOBI nebulised] Serious adverse events: patients withdrawn Follow-up: 24 weeks	26 per 1000	118 per 1000 (46 to 304)	RR 4.54 (1.76 to 11.74)	380 (COLO/D PI/02/06)	⊕⊕⊕⊕ moderate ³	
[COLI DPI versus TOBI nebulised] Serious adverse events: haemoptysis Follow-up: 24 weeks	67 per 1000	107 per 1000 (55 to 209)	RR 1.59 (0.81 to 3.1)	380 (COLO/D PI/02/06)	⊕⊕⊕⊕ very low ^{1,6}	
[COLI nebulised versus TOBI nebulised]	94 per 1000	113 per 1000 (38 to 335)	RR 1.2 (0.4 to 3.55)	115 (Hodson 2002)	⊕⊕⊕⊕ very low ^{1,8}	

Comparison 3.3. Colistin versus Tobramycin					
Serious adverse events: dyspnoea - Follow-up: 4 weeks					
Serious adverse events: dyspnoea Follow-up: 24 weeks	269 per 1000	261 per 1000 (189 to 366)	RR 0.97 (0.7 to 1.36)	380 (COLO/D PI/02/06)	⊕⊕⊕⊕ very low ^{3,8}
[COLI nebulised versus TOBI nebulised] Emergence of resistant organisms: emergence of highly tobramycin-resistant <i>P aeruginosa</i> Follow-up: 24 weeks	0 events	0 events	-	115 (Hodson 2002)	⊕⊕⊕⊕ low ¹

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: cystic fibrosis Q-R: cystic fibrosis questionnaire revised; CI: confidence interval; COLI: colistin; DPI: dry powder for inhalation; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio; TOBI: tobramycin

- 1 The quality of the evidence was downgraded by 2 because this is an open trial, and risk of bias for randomisation and allocation concealment was unclear
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 1 because this is an open trial, and risk of bias for randomisation was unclear
- 4 The quality of the evidence was downgraded by 2 because both studies were open trials, and risk of bias for randomisation and allocation concealment was unclear
- 5 The quality of the evidence was downgraded by 2, as the 95% CI is very large and crosses the line of no effect
- 6 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 7 Not calculable, p-value > 0.05
- 8 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs

Table 119: Summary clinical evidence profile: Comparison 4.1. Tobramycin versus placebo

Comparison 4.1. Tobramycin versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Tobramycin				
Lung function: mean % change in FEV ₁ % predicted Scale from: 0 to	Not reported	The mean lung function: mean % change in FEV ₁ % predicted in the		516 (Galeva 2013, Konstan 2011/)	⊕⊕⊕⊕ low ^{1,2}	

Comparison 4.1. Tobramycin versus placebo						
100 Follow-up: 1 to 3 months		tobramycin groups was 9.36 higher (5.01 to 13.7 higher)		EVOLVE trial, Lenoir 2007, Ramsey 1993)		
Number of patients with 1 or more exacerbations	NMA outcome					
Suppression of the organism: eradication of the organism (negative culture) Follow-up: 4 weeks	Study population		RR 2.46 (1.20 to 5.04)	357 (Chuchalin 2007, Galeva 2013, Lenoir 2007)	⊕⊕⊕⊕ high	
	121 per 1000	299 per 1000 (146 to 612)				
	Moderate					
	143 per 1000	352 per 1000 (172 to 721)				
Suppression of the organism: eradication of the organism (negative culture) Follow-up: 6 weeks	100 per 1000	103 per 1000 (23 to 471)	RR 1.03 (0.23 to 4.71)	59 (Lenoir 2007)	⊕⊕⊕⊖ moderate ³	
Suppression of the organism: eradication of the organism (negative culture) Follow-up: 8 weeks	120 per 1000	145 per 1000 (72 to 289)	RR 1.2 (0.6 to 2.4)	242 (Chuchalin 2007)	⊕⊕⊕⊖ moderate ³	
Suppression of the organism: eradication of the organism (negative culture) Follow-up: 20 weeks	165 per 1000	334 per 1000 (194 to 574)	RR 2.03 (1.18 to 3.49)	235 (Chuchalin 2007)	⊕⊕⊕⊕ high	
Suppression of the organism: eradication of the organism (negative culture) Follow-up: 24 weeks	202 per 1000	239 per 1000 (144 to 397)	RR 1.18 (0.71 to 1.96)	243 (Chuchalin 2007)	⊕⊕⊕⊖ moderate ³	
Suppression of the organism: change in <i>P aeruginosa</i> sputum density log ₁₀ cystic fibrosisU/g Follow-up: 4	The mean change in <i>P aeruginosa</i> sputum density log ₁₀ cystic fibrosisU/g in the placebo	The mean change in <i>P aeruginosa</i> sputum density log ₁₀ cystic fibrosisU/g in the tobramycin groups was		55 (Galeva 2013)	⊕⊕⊕⊖ moderate ⁴	

Comparison 4.1. Tobramycin versus placebo						
weeks	groups was 0	1.2 lower (2.03 to 0.37 lower)				
Suppression of the organism: change in non-mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g Follow-up: 4 weeks	The mean change in non-mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g in the placebo groups was -0.15	The mean change in non-mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g in the tobramycin groups was 1.76 lower (2.52 to 1 lower)		95 (Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ low ⁵	
Suppression of the organism: change in mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g Follow-up: 4 weeks	The mean suppression of the organism: change in mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g in the placebo groups was -0.43	The mean suppression of the organism: change in mucoid <i>P aeruginosa</i> sputum density log10 cystic fibrosisU/g in the tobramycin groups was 2.18 lower (2.97 to 1.39 lower)		95 (Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ low ⁵	
Nutritional status: body weight change kg Follow-up: 12 weeks	The mean nutritional status: body weight change in the control groups was 0.16	The mean body weight change in the tobramycin groups was 0.23 higher (0.23 lower to 0.69 higher)		59 (Lenoir 2007)	⊕⊕⊕⊕ high	
Nutritional status: body weight change kg Follow-up: 24 weeks	The mean body weight change in the tobramycin groups was 1.05	The mean body weight change in the tobramycin groups was 0.75 higher (0.22 to 1.28 higher)		245 (Chuchalin 2007)	⊕⊕⊕⊕ moderate ⁴	
Minor adverse events: minor adverse (any) Follow-up: 4 weeks	Study population		RR 0.66 (0.49 to 0.89)	150 (Galeva 2013, Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ very low ^{4,6}	
	640 per 1000	422 per 1000 (314 to 570)				
	Moderate					
	423 per 1000	279 per 1000 (207 to 376)				
Minor adverse events: minor adverse events (any) Follow-up: 24 weeks	153 per 1000	156 per 1000 (84 to 288)	RR 1.02 (0.55 to 1.88)	246 (Chuchalin 2007)	⊕⊕⊕⊕ low ⁷	

Comparison 4.1. Tobramycin versus placebo						
Minor adverse events: auditory impairment Follow-up: 4 weeks	77 per 1000	103 per 1000 (18 to 572)	RR 1.34 (0.24 to 7.43)	55 (Galeva 2013)	⊕⊕⊕⊕ low ⁷	
Minor adverse events: auditory impairment Follow-up: 24 weeks	0 events	0 events	-	300 (Ramsey 1999)	⊕⊕⊕⊕ high	
Minor adverse events: auditory impairment Follow-up: 42 weeks	0 events	0 events	-	71 (Ramsey 1993)	⊕⊕⊕⊕ high	
Minor adverse events: cough Follow-up: 4 weeks	173 per 1000	289 per 1000 (14 to 1000)	RR 1.67 (0.08 to 36.11)	150 (Galeva 2013, Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ very low ^{6,7,8}	
Minor adverse events: tinnitus Follow-up: 24 weeks	0 events	-	RR 17.26 (1 to 297.54)	520 (Ramsey 1999)	⊕⊕⊕⊕ moderate ⁴	
Headaches Follow-up: 4 weeks	20 per 1000	7 per 1000 (1 to 67)	RR 0.36 (0.04 to 3.29)	95 (Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ very low ^{5,7}	
Major adverse events: any Follow-up: 4 weeks	Study population		RR 0.52 (0.16 to 1.64)	150 (Galeva 2013, Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ very low ^{6,7}	
	107 per 1000	55 per 1000 (17 to 175)				
	Moderate					
	39 per 1000	20 per 1000 (6 to 64)				
Major adverse events: any Follow-up: 24 weeks	259 per 1000	106 per 1000 (60 to 189)	RR 0.41 (0.23 to 0.73)	246 (Chuchalin 2007)	⊕⊕⊕⊕ high	
Major adverse events: haemoptysis Follow-up: 4 weeks	20 per 1000	22 per 1000 (1 to 338)	RR 1.07 (0.07 to 16.54)	95 (Konstan 2011/ EVOLVE trial)	⊕⊕⊕⊕ very low ^{5,7}	
Major adverse events: haemoptysis Follow-up: 24 weeks	309 per 1000	269 per 1000 (204 to 349)	RR 0.87 (0.66 to 1.13)	520 (Ramsey 1999)	⊕⊕⊕⊕ moderate ⁴	
Major adverse events: pneumothorax Follow-up: 24 weeks	15 per 1000	4 per 1000 (0 to 35)	RR 0.25 (0.03 to 2.26)	520 (Ramsey 1999)	⊕⊕⊕⊕ low ⁷	

Comparison 4.1. Tobramycin versus placebo						
Mortality Follow-up: 4 weeks	20 per 1000	7 per 1000 (0 to 173)	RR 0.35 (0.01 to 8.49)	95 (1 study)	⊕⊕⊕⊕ low ⁹	
Mortality Follow-up: 3 to 12 months	17 per 1000	3 per 1000 (1 to 19)	RR 0.17 (0.03 to 1.09)	767 (Chuchalin 2007, Ramsey 1999)	⊕⊕⊕⊕ moderate ³	
Emergence of resistant organisms: frequency of Tobramycin-resistant <i>P. aeruginosa</i> Follow-up: 24 weeks	105 per 1000	204 per 1000 (90 to 463)	RR 1.95 (0.86 to 4.42)	672 (Chuchalin 2007, Ramsey 1999)	⊕⊕⊕⊕ very low ^{4,10}	
Emergence of resistant organisms: frequency of new isolates of drug resistant <i>B. cepacia</i> Follow-up: 24 weeks	0 events	0 events	-	258 (Ramsey 1999)	⊕⊕⊕⊕ high	
Emergence of resistant organisms: frequency of new isolates of drug resistant <i>S. maltophilia</i> Follow-up: 24 weeks	4 per 1000	12 per 1000 (1 to 111)	RR 3.05 (0.32 to 29.1)	520 (Ramsey 1999)	⊕⊕⊕⊕ low ⁷	
Emergence of resistant organisms: frequency of new isolates of drug resistant <i>A. xylosoxidans</i> Follow-up: 24 weeks	4 per 1000	4 per 1000 (0 to 62)	RR 1.02 (0.06 to 16.15)	520 (Ramsey 1999)	⊕⊕⊕⊕ low ⁷	
Emergence of resistant organisms: frequency of new isolates of drug resistant <i>aspergillus</i> Follow-up: 24 weeks	104 per 1000	21 per 1000 (7 to 59)	RR 0.2 (0.07 to 0.57)	389 (Ramsey 1999)	⊕⊕⊕⊕ high	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: cystic fibrosisU/G: colony forming units per gram; CI: confidence interval; FEV ₁ :						

Comparison 4.1. Tobramycin versus placebo

- forced expiratory volume in 1 second; kg: kilogrammes; MD: mean difference; RR: risk ratio
- 1 *1 The quality of the evidence was downgraded by 1, as 1 of the trials had unclear risk of bias for the domains*
2 *randomisation, allocation concealment, and blinding and another trial had unclear risk of bias for the domains*
3 *randomisation, allocation concealment and high risk of bias for blinding*
4 *2 The quality of the evidence was downgraded by 1 due to moderate inconsistency (I²=51%). Sub-group analysis*
5 *was not conducted, as all of the trials showed a beneficial effect of tobramycin*
- 6 *3 The quality of the evidence was downgraded by 1 as the 95% CI crossed the null effect*
7 *4 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID*
8 *5 The quality of the evidence was downgraded by 2 due to unclear risk of bias for the domains randomisation,*
9 *allocation concealment and high risk of bias for blinding*
- 10 *6 The quality of the evidence was downgraded by 2, as the largest trial had unclear risk of*
11 *bias for the domains randomisation, allocation concealment and high risk of bias for blinding*
- 12 *7 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs*
- 13 *8 The quality of the evidence was downgraded by 2 due to very serious inconsistency*
14 *(I²=77%).*
- 15 *9 The quality of the evidence was downgraded by 2 as the 95% CI is very wide and it*
16 *crossed the null effect. The study is underpowered to detect differences*
- 17 *10 The quality of the evidence was downgraded by 2 due to very serious inconsistency*
18 *(I²=79%)*

Table 120: Summary clinical evidence profile: Comparison 4.2. Tobramycin inhalation powder versus tobramycin inhalation solution

Comparison 4.2. Tobramycin inhalation powder versus tobramycin inhalation solution						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Tobramycin inhalation solution (TOBI neb)	Tobramycin inhalation powder (TOBI DPI)				
Lung function: % mean change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 4 weeks	Not reported	The mean % mean change in FEV ₁ % predicted in the TOBI DPI groups was 0.8 lower (3.9 lower to 2.3 higher)		517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊖ low ^{1,2}	
Lung function: % mean change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 20 weeks	Not reported	The mean % mean change in FEV ₁ % predicted in the TOBI DPI groups was 1.10 higher (2.33 lower to 4.53 higher)		517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊖ low ^{1,2}	
Lung function: % mean change in FEV ₁	Not reported	The mean % mean change in FEV ₁ %		517 (Konstan 2011a/EA	⊕⊕⊕⊖ low ^{1,2}	

Comparison 4.2. Tobramycin inhalation powder versus tobramycin inhalation solution						
% predicted Scale from: 0 to 100 Follow-up: 24 weeks		predicted in the TOBI DPI groups was 2.20 higher (1.11 lower to 5.51 higher)		GER trial)		
Number of patients with 1 or more exacerbations	NMA outcome					
Suppression of the organism: mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU Follow-up: 4 weeks	The mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU in the TOBI neb groups was -1.32	The mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU in the TOBI DPI groups was 0.44 lower (0.79 to 0.09 lower)		517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ moderate ¹	
Suppression of the organism: mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU Follow-up: 20 weeks	The mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU in the TOBI neb groups was -0.77	The mean change in <i>P aeruginosa</i> sputum density log10 cystic fibrosisU in the TOBI DPI groups was 0.84 lower (1.17 to 0.51 lower)		517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ low ^{1,3}	
Adverse events - any mild or moderate AE Follow-up: 24 weeks	684 per 1000	732 per 1000 (657 to 821)	RR 1.07 (0.96 to 1.2)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ moderate ¹	
Adverse events - any serious AE Follow-up: 24 weeks	292 per 1000	271 per 1000 (207 to 362)	RR 0.93 (0.71 to 1.24)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ low ^{1,3}	
mild AE: productive cough Follow-up: 24 weeks	196 per 1000	182 per 1000 (126 to 261)	RR 0.93 (0.64 to 1.33)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ very low ^{1,4}	
mild AE: headache	120 per 1000	114 per 1000 (71 to 184)	RR 0.95 (0.59 to 1.54)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ very low ^{1,4}	
mild AE: vomiting Follow-up: 24 weeks	57 per 1000	61 per 1000 (30 to 125)	RR 1.07 (0.53 to 2.17)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ very low ^{1,4}	
serious AE: dyspnoea	124 per 1000	156 per 1000 (100 to 243)	RR 1.25 (0.8 to	517 (Konstan	⊕⊕⊕⊕ very	

Comparison 4.2. Tobramycin inhalation powder versus tobramycin inhalation solution						
Follow-up: 24 weeks			1.95)	2011a/EA GER trial)	low ^{1,4}	
serious AE: haemoptysis Follow-up: 24 weeks	124 per 1000	129 per 1000 (82 to 207)	RR 1.04 (0.66 to 1.66)	517 (Konstan 2011a/EA GER trial)	⊕⊕⊕⊕ very low ^{1,4}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: cystic fibrosisU: colony forming units; CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio						

- 1 The quality of the evidence was downgraded by 1 as this is an open trial, and randomization is unclear
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MID

5 **Table 121: Summary clinical evidence profile: Comparison 4.3. Tobramycin versus**
6 **Aztreonam lysine**

Comparison 4.3. Tobramycin versus Aztreonam						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Aztreonam lysine	Tobramycin				
[TOBI nebulised versus AZLI inhaled] Lung function: % change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 3 months	The mean % change in FEV ₁ % predicted in the aztreonam lysine groups was 2.05	The mean % change in FEV ₁ % predicted in the tobramycin groups was 2.71 lower (2.88 to 2.54 lower)		268 (Assael 2013)	⊕⊕⊕⊕ moderate ¹	
Number of patients with 1 or more exacerbations	NMA outcome					
[TOBI nebulised versus AZLI inhaled] Suppression of the organism: adj mean change sputum density log ₁₀ PA cystic fibrosisU/g- Follow-up: 20 weeks	The adjusted mean change sputum density log ₁₀ PA cystic fibrosisU/g in the aztreonam lysine groups was -0.55	The adjusted mean change sputum density log ₁₀ PA cystic fibrosisU/g in the tobramycin groups was 0.23 higher (0.3 lower to 0.76 higher)		194 (Assael 2013)	⊕⊕⊕⊕ low ^{1,2}	
[TOBI nebulised versus AZLI	The adjusted	The adjusted mean weight		268 (Assael	⊕⊕⊕⊕ low ^{1,2}	

Comparison 4.3. Tobramycin versus Aztreonam						
inhaled] Nutritional status: % adj mean weight change – Follow-up: 20 weeks	mean weight change in the aztreonam lysine groups was 0.58	change in the tobramycin groups was 0.52 lower (1.68 lower to 0.64 higher)		2013)		
[TOBI nebulised versus AZLI inhaled] Quality of life: cystic fibrosisQR respiratory, adj mean change – Follow-up: 20 weeks	The adjusted mean change in cystic fibrosisQR respiratory in the aztreonam lysine groups was 6.3	The adjusted mean change in cystic fibrosisQR respiratory in the tobramycin groups was 4.1 lower (8.59 lower to 0.39 higher)		268 (Assael 2013)	⊕⊕⊕⊕ low ^{1,3}	
[TOBI nebulised versus AZLI inhaled] Minor adverse events: chest discomfort – Follow-up: 3 months	103 per 1000	99 per 1000 (48 to 202)	RR 0.96 (0.47 to 1.96)	268 (Assael 2013)	⊕⊕⊕⊕ very low ^{1,4}	
[TOBI nebulised versus AZLI inhaled] Minor adverse events: cough – Follow-up: 3 months	706 per 1000	791 per 1000 (685 to 904)	RR 1.12 (0.97 to 1.28)	268 (Assael 2013)	⊕⊕⊕⊕ low ^{1,2}	
[TOBI nebulised versus AZLI inhaled] Minor adverse events: headache - Follow-up: 3 months	213 per 1000	205 per 1000 (128 to 326)	RR 0.96 (0.6 to 1.53)	268 (Assael 2013)	⊕⊕⊕⊕ very low ^{1,4}	
[TOBI nebulised versus AZLI inhaled] Minor adverse events: vomiting - Follow-up: 3 months	103 per 1000	106 per 1000 (53 to 214)	RR 1.03 (0.51 to 2.08)	268 (Assael 2013)	⊕⊕⊕⊕ very low ^{1,4}	
[TOBI nebulised versus AZLI inhaled] Major adverse events: dyspnoea -	228 per 1000	160 per 1000 (96 to 262)	RR 0.7 (0.42 to 1.15)	268 (Assael 2013)	⊕⊕⊕⊕ low ^{1,2}	

Comparison 4.3. Tobramycin versus Aztreonam						
Follow-up: 3 months						
[TOBI nebulised versus AZLI inhaled]	228 per 1000	160 per 1000 (96 to 262)	RR 0.7 (0.42 to 1.15)	268 (Assael 2013)	⊕⊕⊕⊖	low ^{1,2}
Major adverse events: haemoptysis						
Follow-up: 3 months						
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: AZLI: aztreonam lysine; cystic fibrosisQ-R: cystic fibrosis questionnaire revised; cystic fibrosisU/g: colony forming units per gram; CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio; TOBI: tobramycin						

- 1 The quality of the evidence was downgraded by 1 because this is an open trial
- 2 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID
- 3 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 4 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs

5 **Table 122: Summary clinical evidence profile: Comparison 5. Combination of**
6 **fosfomycin + tobramycin versus placebo**

Comparison 5. Combination of fosfomycin + tobramycin versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Combination of fosfomycin + tobramycin (FTI)				
Lung function: relative change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 4 weeks [FTI 80/20 mg]	Not reported	The relative change in FEV ₁ % predicted in the fosfomycin + tobramycin groups was 7.5 higher (3.6 to 11.4 higher)		70 (Trapnell 2012)	⊕⊕⊕⊖ moderate ¹	
Lung function: relative change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 4 weeks [FTI160/40 mg]	Not reported	The mean relative change in FEV ₁ % predicted in the fosfomycin + tobramycin groups was 6.2 higher (2.42 to 9.98 higher)		73 (Trapnell 2012)	⊕⊕⊕⊖ low ^{1,2}	
Suppression of the organism: change in sputum <i>P aeruginosa</i> density, log 10	Not reported	The mean change in sputum <i>P aeruginosa</i> density, log 10 cystic fibrosisU/g in the fosfomycin + tobramycin groups		70 (Trapnell 2012)	⊕⊕⊕⊖ low ^{1,3}	

Comparison 5. Combination of fosfomycin + tobramycin versus placebo					
cystic fibrosisU/g Follow-up: 4 weeks [FTI 80/20 mg]		was 1.04 lower (1.82 to 0.26 lower)			
Suppression of the organism: change in sputum <i>P aeruginosa</i> density, log 10 cystic fibrosisU/g mg Follow-up: 4 weeks [FTI160/40 mg]	Not reported	The mean change in sputum <i>P aeruginosa</i> density, log 10 cystic fibrosisU/g in the fosfomacyn + tobramycin groups was 0.28 lower (1.06 lower to 0.5 higher)	73 (Trapnell 2012)	⊕⊕⊕⊖ low ^{1,3}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: cystic fibrosisU: colony forming units; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; FTI: Fosfomycin/ tobramycin inhaled; MD: mean difference; mg: milligrams; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1 due to unclear risk of bias for allocation concealment and data reporting
- 2 The quality of the evidence was downgraded by as the 95 % CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by as the 95% CI crossed 1 default MID

5 **Table 123: Summary clinical evidence profile: Comparison 6. Continuous**
6 **alternating therapy versus intermittent treatment: aztreonam lysine +**
7 **tobramycin or placebo + tobramycin**

Comparison 6. Continuous alternating therapy versus intermittent treatment: aztreonam + tobramycin or placebo + tobramycin						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Intermittent treatment: aztreonam lysine + tobramycin or placebo + tobramycin	Continuous alternating therapy				
Lung function: % change in FEV ₁ predicted Scale from: 0 to 100 Follow-up: 20 weeks ¹	The mean % change in FEV ₁ % predicted in the intermittent treatment groups was 0.04	The mean % change in FEV ₁ % predicted in the continuous alternating therapy groups was 1.33 higher (1.05 to 1.61 higher)		88 (Flume 2016)	⊕⊕⊕⊖ moderate ²	

Comparison 6. Continuous alternating therapy versus intermittent treatment: aztreonam + tobramycin or placebo + tobramycin						
Time to next pulmonary exacerbation	-	-	HR 0.89 (0.49 to 1.6)	88 (Flume 2016)	⊕⊕⊕⊕ low ^{2,3}	
Quality of life: change in cystic fibrosisQ-R Scale from: 0 to 100 Follow-up: 20 weeks ¹	The mean change in cystic fibrosisQ-R in the intermittent treatment groups was -2.06	The mean change in cystic fibrosisQ-R in the continuous alternating therapy groups was 3.06 higher (2.35 to 3.77 higher)		88 (Flume 2016)	⊕⊕⊕⊕ low ^{2,4}	
Minor adverse events: cough Follow-up: 3 months	435 per 1000	761 per 1000 (526 to 1000)	RR 1.75 (1.21 to 2.54)	88 (Flume 2016)	⊕⊕⊕⊕ low ^{2,5}	
Serious adverse events: dyspnoea Follow-up: 3 months	522 per 1000	308 per 1000 (183 to 527)	RR 0.59 (0.35 to 1.01)	88 (Flume 2016)	⊕⊕⊕⊕ low ^{2,5}	
Serious adverse events (not treatment related) Follow-up: 3 months	522 per 1000	501 per 1000 (334 to 751)	RR 0.96 (0.64 to 1.44)	88 (Flume 2016)	⊕⊕⊕⊕ very low ^{2,6}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: cystic fibrosisQ-R: cystic fibrosis questionnaire reviewed; CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; MD: mean difference; mg: milligrams; RR: risk ratio						

- 1 Values at 4, 12 and 20 weeks were averaged
- 2 The quality of the evidence was downgraded by 1 due to unclear allocation concealment, blinding, and data collection/ reporting
- 3 The quality of the evidence was downgraded by 1 as the 95% CI crossed the null effect line
- 4 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 clinical MID
- 5 The quality of the evidence was downgraded by 1 as the 95%CI crossed 1 default MID
- 6 The quality of the evidence was downgraded by 2 as the 95%CI crossed 2 default MIDs

89.4.3.4.1 **S Aureus**

9 No relevant studies were identified.

109.4.3.4.2 **B cepacia complex**

11 No relevant studies were identified.

19.4.3.4.3 *A fumigatus*

2 The clinical evidence profile for the antimicrobial treatment of chronic infection with *A*
3 *fumigatus* is presented in Table 65.

4 **Table 124: Summary clinical evidence profile: Comparison 7. Itraconazole versus**
5 **placebo**

Comparison 7. Itraconazole versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Itraconazole				
Lung function: percentage change in FEV ₁ predicted from baseline Scale from: 0 to 100 Follow-up: mean 24 weeks	The mean lung function in the placebo group was: 0.32	The mean lung function change in the itraconazole group was 4.94 lower (15.33 lower to 5.45 higher)	-	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,3}	
Lung function: percentage change in FEV ₁ predicted from baseline Scale from: 0 to 100 Follow-up: mean 48 weeks	Not reported	The mean lung function change in the itraconazole group was 3.71 lower (-13.26 lower to 20.68 higher)	Not estimable	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,3}	
Time to next pulmonary exacerbation Follow-up: mean 24 weeks	The median time to next exacerbation in the placebo group was: 134 days	The median time to next exacerbation in the itraconazole group was: 77 days	adjHR 1.34 (0.57 to 3.14)	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,4}	
proxy: number of patients with an exacerbation requiring AB Follow-up: mean 24 weeks	389 per 1000	665 per 1000 (342 to 1000)	RR 1.71 (0.88 to 3.33)	36 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,5}	
proxy: number of patients with an exacerbation requiring AB Follow-up: mean 48 weeks	611 per 1000	831 per 1000 (544 to 1000)	RR 1.36 (0.89 to 2.08)	36 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
proxy: number of patients with an exacerbation	176 per 1000	166 per 1000 (39 to 715)	RR 0.94 (0.22 to 4.05)	35 (Aaron 2012)	⊕⊕⊕⊕ very	

Comparison 7. Itraconazole versus placebo						
admitted to hospital Follow-up: mean 24 weeks						low ^{1,2,6}
proxy: number of patients with an exacerbation admitted to hospital Follow-up: mean 48 weeks	176 per 1000	222 per 1000 (58 to 851)	RR 1.26 (0.33 to 4.82)	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,4}	
Quality of life - all domains cystic fibrosisQ-R Scale from: 0 to 100 Follow-up: mean 24 weeks	Not reported	Not reported	Not estimabl e, but no significa nt differenc es found	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,7}	
Quality of life – change in respiratory domain cystic fibrosisQ-R Scale from: 0 to 100 Follow-up: mean 24 weeks	The mean change in QFQ- R score for the respirator y domain in the placebo group was: 4.77	The mean change in QFQ- R score for the respiratory domain in the itraconazole group was: 3.76	Not estimabl e, but no significa nt differenc es (p=0.87)	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,7}	
Minor adverse events - increased dyspnoea Follow-up: mean 24 weeks	125 per 1000	111 per 1000 (18 to 700)	RR 0.89 (0.14 to 5.6)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
Minor adverse events - rash Follow-up: mean 24 weeks	62 per 1000	111 per 1000 (11 to 1000)	RR 1.78 (0.18 to 17.8)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
Minor adverse events - hyperglycaemia Follow-up: mean 24 weeks	0 per 1000	0 per 1000 (0 to 0)	RR 2.68 (0.12 to 61.58)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
Minor adverse events - flu-like illness Follow-up: mean 24 weeks	0 per 1000	0 per 1000 (0 to 0)	RR 6.26 (0.35 to 112.7)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
Minor adverse events - diarrhoea Follow-up: mean 24 weeks	62 per 1000	19 per 1000 (1 to 428)	RR 0.3 (0.01 to 6.84)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	
Minor adverse events (lower scores are better) - conjunctivitis Follow-up: mean	62 per 1000	19 per 1000 (1 to 428)	RR 0.3 (0.01 to 6.84)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}	

Comparison 7. Itraconazole versus placebo					
24 weeks					
Major adverse events - haemoptysis Follow-up: mean 24 weeks	62 per 1000	111 per 1000 (11 to 1000)	RR 1.78 (0.18 to 17.8)	34 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}
Major adverse events - spontaneous pneumothorax Follow-up: mean 24 weeks	0 per 1000	0 per 1000 (0 to 0)	RR 2.84 (0.12 to 65.34)	35 (Aaron 2012)	⊕⊕⊕⊕ very low ^{1,2,6}
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: cystic fibrosisQ-R: cystic fibrosis questionnaire reviewed; CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MD: mean difference; RR: risk ratio</p>					

- 1 *1 The quality of the evidence was downgraded by 1 due to unclear allocation, data reporting and sample size*
- 2 *2 The quality of the evidence was downgraded by 1 due to indirectness, as the therapeutic dosages were not*
- 3 *achieved in 2/3 of the participants*
- 4 *3 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 clinical MIDs.*
- 5 *4 The quality of the evidence was downgraded by 2 as the 95% CI crossed the null effect and it is very wide. The*
- 6 *study is underpowered to detect differences between groups.*
- 7 *5 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID.*
- 8 *6 The quality of the evidence was downgraded by 2 as the 95% CI crossed 2 default MIDs*
- 9 *7 Not calculable, as no data was provided in the study.*

10 9.4.3.5 Economic evidence

11 Four economic evaluations of antimicrobial agents to suppress chronic infection with *P*

12 *aeruginosa* were identified in the literature search conducted for this guideline. The methods

13 and results of those studies are described in Appendix K. Full details of the search and

14 economic article selection flow chart can be found in Appendix E and F, respectively. Data

15 extraction tables and quality assessments of included studies can be found in Appendix L

16 and M, respectively.

17 This area was prioritised for de novo economic modelling; consequently, a cost-utility model

18 was developed. Due to study heterogeneity, it was considered inappropriate to undertake

19 one reliable, fully incremental analysis; hence, 4 comparisons within the model were

20 developed. The model uses a lifetime horizon based on the assumption that antimicrobials to

21 suppress *P aeruginosa* are given on a long-term basis.

22 The model takes the form of a state transition model to estimate transitions between 3 lung

23 function (FEV₁% predicted) strata. Transition probabilities between the 3 FEV₁% strata and

24 the probability of experiencing an exacerbation each cycle were taken from the clinical

25 evidence review. A post lung transplant health state was also included in the model to reflect

26 the clinical pathway.

27 A series of scenario analyses were undertaken in order to test how sensitive the results were

28 to uncertainty in individual parameters. The methods used to construct the model and the

29 results of all analyses are reported in Appendix K.

30 Manufacturers of antimicrobials included in the model have agreed Patient Access Schemes

31 (PAS) with the Department of Health to reduce the cost of their drug, to subsequently

32 increase cost-effectiveness. To account for these discounts on the drug acquisition cost,

33 there is an option in the model to apply them. For completeness, the Committee applied

1 those discounts in the model, to reassess their cost-effectiveness. Table 125 below provides
2 the base case result using list prices and PAS prices over a lifetime horizon.

3 **Table 125: Results from the economic model**

Treatment	Total costs	Total QALYs	Inc. costs	Inc. QALYs	ICER
Comparison 1					
List price					
Placebo	£92,040	11.25	-	-	-
Nebulised colistimethate sodium	£105,872	11.52	£13,833	0.33	£52,168
Nebulised tobramycin	£244,919	11.57	£139,047	0.05	£2,824,240
Tobramycin dry powder	£274,658	11.43	£29,739	-0.14	Dominated
PAS price					
Placebo	No change	No change	-	-	-
Nebulised colistimethate sodium	No change	No change	No change	No change	No change
Nebulised tobramycin	Reduced	No change	Reduced	No change	Reduced (>£30,000)
Tobramycin dry powder	Reduced	No change	Reduced	No change	No change
Comparison 2					
List price					
Nebulised colistimethate sodium	£107,149	11.35	-	-	-
Nebulised tobramycin	£244,890	11.58	£137,741	0.23	£600,472
PAS price					
Nebulised colistimethate sodium	No change	No change	-	-	-
Nebulised tobramycin	Reduced	No change	Reduced	No change	Reduced (>£30,000)
Comparison 3					
List price					
Colistimethate sodium dry powder	£276,593	11.37	-	-	-
Nebulised tobramycin	£245,561	11.41	-£31,032	0.04	Dominant
PAS price					
Colistimethate sodium dry powder	Reduced	No change	-	-	-
Nebulised tobramycin	Reduced	No change	Increased	No change	Increased (>£20,000)
Comparison 4					
List price					
Nebulised tobramycin	£245,830	11.35	-	-	-
Nebulised aztreonam lysine	£265,151	11.91	£19,321	0.56	£34,348
Combination ^a	£340,265	11.53	£75,114	-0.38	Dominated
PAS price					
Nebulised tobramycin	Reduced	No change	-	-	-
Nebulised aztreonam lysine	Reduced	No change	Reduced	No change	Reduced (<£30,000)

Treatment	Total costs	Total QALYs	Inc. costs	Inc. QALYs	ICER
Combination ^a	Reduced	No change	Reduced	No change	No change

1 (a) 28 days aztreonam lysine (nebulised) alternating with 28 days tobramycin (nebulised)

2 9.4.3.6 Evidence statements

39.4.3.6.1 Antimicrobial regimens for the treatment of chronic *P Aeruginosa*

4 Aztreonam lysine

5 Comparison 1: Aztreonam lysine versus placebo

6 Lung function: FEV₁

7 Moderate quality evidence from 1 RCT with 157 people with cystic fibrosis and chronic *P*
8 *aeruginosa* infection ≥ 6 years showed a clinically significant improvement in lung function
9 (measured as relative change in FEV₁% predicted) in the group of participants receiving
10 Aztreonam lysine (75 mg/ day) compared to those in the placebo group at 28 days follow-up.

11 Number of people with exacerbations

12 NMA outcome

13 Time to next exacerbation

14 No evidence was found for this critical outcome.

15 Suppression of the organism

16 High quality evidence from two RCTs with 321 people with cystic fibrosis and chronic *P*
17 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the suppression
18 of *P Aeruginosa* (measured as change in sputum density log₁₀ cystic fibrosisU/G) between
19 the participants who were receiving Aztreonam lysine (75 mg/day) and those who were
20 receiving placebo at 4 weeks follow-up.

21 Nutritional status: weight

22 High quality evidence from one RCT with 164 people with cystic fibrosis and chronic *P*
23 *aeruginosa* infection ≥ 6 years showed difference clinically significant improvement in the
24 nutritional status (measured as % weight change in kg) between the participants who were
25 receiving Aztreonam lysine (75 mg/day) compared to those who were receiving placebo at 4
26 weeks follow-up.

27 Quality of life

28 Very low to high quality evidence from two RCTs with 321 people with cystic fibrosis and
29 chronic *P aeruginosa* infection ≥ 6 years showed a clinically significant beneficial effect of
30 Aztreonam lysine (75 mg/day) in the following domains of quality of life: eating and vitality
31 (measured with the CFQ-R questionnaire) compared to placebo, at 4 weeks follow-up.
32 However, very low to moderate quality evidence from the same trials showed no clinically
33 difference in the following domains of quality of life: body image, digestion, eating, emotional
34 functioning, physical functioning, respiratory symptoms, role/ school, social functioning,
35 treatment burden and weight ((measured with the CFQ-R questionnaire) between both
36 groups.

1 Moderate to high unexplained heterogeneity was found for the following domains: eating,
2 emotional functioning, health perceptions, physical functioning, respiratory symptoms, role or
3 school, treatment burden and vitality.

4 **Mild adverse events**

5 Low quality evidence from 1 RCT with 164 people with cystic fibrosis and chronic *P*
6 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
7 chest discomfort between the participants who were receiving Aztreonam lysine (75 mg/day)
8 and those who were receiving placebo at 4 weeks follow-up.

9 Low quality evidence from 3 RCTs with 532 people with cystic fibrosis and chronic *P*
10 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
11 cough between the participants who were receiving Aztreonam lysine (75 mg/day) and those
12 who were receiving placebo at 4 weeks follow-up.

13 Very low quality evidence from 2 RCTs with 321 people with cystic fibrosis and chronic *P*
14 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
15 headache between the participants who were receiving Aztreonam lysine (75 mg/day) and
16 those who were receiving placebo at 4 weeks follow-up. Significant unexplained
17 heterogeneity was found between both trials, although both showed no clinically significant
18 differences between both treatment groups.

19 **Serious adverse events**

20 Low quality evidence from 1 RCT with 164 people with cystic fibrosis and chronic *P*
21 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
22 dyspnoea between the participants who were receiving Aztreonam lysine (75 mg/day) and
23 those who were receiving placebo at 4 weeks follow-up.

24 Low quality evidence from 2 RCTs with 375 people with cystic fibrosis and chronic *P*
25 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
26 haemoptysis between the participants who were receiving Aztreonam lysine (75 mg/day) and
27 those who were receiving placebo at 4 weeks follow-up.

28 **Mortality**

29 High quality evidence from 1 RCT with 211 people with cystic fibrosis and chronic *P*
30 *aeruginosa* infection $>$ 7 years showed that there were no deaths in either group (Aztreonam
31 lysine 75 mg/day or placebo) at 4 weeks follow-up.

32 **Emergence of resistant organisms**

33 Low to moderate quality evidence from 1 RCT with 155 people with cystic fibrosis and
34 chronic *P aeruginosa* infection \geq 6 years showed no clinically significant difference in the
35 number of people in whom *S aureus*, *S maltophilia* or *A xilosidans* was persistently isolated
36 between the participants who were receiving Aztreonam lysine (75 mg/day) and those who
37 were receiving placebo at 42 days follow-up.

38 High quality evidence from 1 RCT with 155 people with cystic fibrosis and chronic *P*
39 *aeruginosa* infection \geq 6 years showed that *B Cepacia* was not isolated in the participants of
40 either group (Aztreonam lysine 75 mg/day or placebo) at 42 days follow-up.

419.4.3.6.2 **Azithromycin (high-dose only)**

42 No evidence was found for this treatment.

19.4.3.6.3 **Ciprofloxacin (oral)**

2 **Comparison 2: Ciprofloxacin versus placebo**

3 **Lung function**

4 No evidence was found for this critical outcome.

5 **Number of people with exacerbations**

6 NMA outcome.

7 **Time to next exacerbation**

8 No evidence was found for this critical outcome.

9 **Suppression of the organism**

10 No evidence was found for this important outcome.

11 **Nutritional status: weight**

12 Very low quality evidence from 1 RCT with 40 adults with cystic fibrosis and chronic *P*
13 *aeruginosa* infection showed no clinically significant difference in weight (kg) between the
14 participants who were receiving Ciprofloxacin (500 mg. for 10 days/ every 3 months) and
15 those who were receiving placebo at 12 months follow-up.

16 **Quality of life**

17 No evidence was found for this important outcome.

18 **Mild adverse events**

19 Very low quality evidence from 1 RCT with 40 adults with cystic fibrosis and chronic *P*
20 *aeruginosa* infection showed no clinically significant difference in the occurrence of
21 gastrointestinal adverse events between the participants who were receiving Ciprofloxacin
22 (500 mg. for 10 days/ every 3 months) and those who were receiving placebo at 12 months
23 follow-up.

24 **Serious adverse events**

25 No evidence was found for this important outcome.

26 **Mortality**

27 Low quality evidence from 1 RCT with 40 adults with cystic fibrosis and chronic *P aeruginosa*
28 infection showed no clinically significant difference in the mortality rate between the
29 participants who were receiving Ciprofloxacin (500 mg. for 10 days/ every 3 months) and
30 those who were receiving placebo at 12 months follow-up.

31 **Emergence of resistant organisms**

32 Very low quality evidence from 1 RCT with 40 adults with cystic fibrosis and chronic *P*
33 *aeruginosa* infection showed no clinically significant difference in the number of people in
34 whom resistant strains of *P Aeruginosa* were isolated between the participants who were
35 receiving Ciprofloxacin (500 mg. for 10 days/ every 3 months) and those who were receiving
36 placebo at 12 months follow-up.

1 Very low quality evidence from 1 RCT with 40 adults with cystic fibrosis and chronic *P*
2 *aeruginosa* infection showed no clinically significant difference in the number of people in
3 whom resistant strains of *S aureus* were isolated between the participants who were
4 receiving Ciprofloxacin (500 mg. for 10 days/ every 3 months) and those who were receiving
5 placebo at 12 months follow-up.

69.4.3.6.4 **Colistimethate sodium (dry powder, inhaled)**

7 **Comparison 3.1: Colistin versus placebo**

8 **Lung function: FEV₁**

9 Low quality evidence from 1 RTC with 29 people with cystic fibrosis and chronic *P*
10 *aeruginosa* infection >7 years showed no clinically significant difference in lung function
11 (measure as change in FEV₁ % predicted) between the participants receiving colistin (1
12 million units, twice daily for 3 months) and those receiving placebo at 3 months follow-up.

13 **Number of people with exacerbations**

14 NMA outcome

15 **Time to next exacerbation**

16 No evidence was found for this critical outcome.

17 **Suppression of the organism**

18 Moderate quality evidence from 1 RCT with 40 people with cystic fibrosis and chronic *P*
19 *aeruginosa* infection >7 years showed that *P Aeruginosa* was not eradicated of the sputum of
20 any patient (Colistin solution 1 million units, twice daily for 3 months or placebo) during the 3
21 month trial.

22 **Nutritional status**

23 No evidence was found for this important outcome.

24 **Quality of life**

25 No evidence was found for this important outcome.

26 **Mild adverse events**

27 No evidence was found for this important outcome.

28 **Serious adverse events**

29 No evidence was found for this important outcome.

30 **Mortality**

31 No evidence was found for this important outcome.

32 **Emergence of resistant organisms**

33 Moderate quality evidence from 1 RCT with 40 people with cystic fibrosis and chronic *P*
34 *aeruginosa* infection >7 years reported that none of the patients in either group (Colistin
35 solution 1 million units, twice daily for 3 months or placebo) were infected with other colistin-

1 resistant microorganisms (*Ps. Cepacia*, *Serratia marcescens*, *Preteus mirabilis*, Gram-positive
2 organisms or fungi) during the 3 month trial.

3 Moderate quality evidence from 1 RCT with 40 people with cystic fibrosis and chronic *P*
4 *aeruginosa* infection >7 years reported that resistance to colistin was not developed in any
5 patient (Colistin solution 1 million units, twice daily for 3 months or placebo) during the 3
6 month trial.

7 Moderate quality evidence from 1 RCT with 40 people with cystic fibrosis and chronic *P*
8 *aeruginosa* infection >7 years reported no change in resistant pattern to other commonly
9 used anti-pseudomonas treatments in any patient (Colistin solution 1 million units, twice daily
10 for 3 months or placebo) during the 3 month trial.

11 **Comparison 3.2: Colistin DPI versus colistin nebulised**

12 **Lung function: FEV₁**

13 Very low quality evidence from 1 RCT with 31 people with cystic fibrosis and chronic *P*
14 *aeruginosa* infection ≥8 years showed no clinically significant difference in lung function
15 (measured as % mean change in FEV₁ % predicted) between the participants who were
16 receiving Colistin DPI (125 mg/ twice daily) and those receiving Colistin inhalation solution (2
17 MU/ twice daily) at 4 weeks follow-up.

18 **Number of people with exacerbations**

19 NMA outcome

20 **Time to next exacerbation**

21 No evidence was found for this critical outcome.

22 **Suppression of the organism**

23 No evidence was found for this important outcome.

24 **Nutritional status**

25 No evidence was found for this important outcome.

26 **Quality of life**

27 No evidence was found for this important outcome.

28 **Mild adverse events**

29 Very low quality evidence from 1 RCT with 31 people with cystic fibrosis and chronic *P*
30 *aeruginosa* infection ≥8 years showed no clinically significant difference in the occurrence of
31 chest discomfort, cough and vomiting between the participants who were receiving Colistin
32 DPI (125 mg/ twice daily) and those receiving Colistin inhalation solution (2 MU/ twice daily)
33 at 8 weeks follow-up.

34 **Serious adverse events**

35 Very low quality evidence from 1 RCT with 31 people with cystic fibrosis and chronic *P*
36 *aeruginosa* infection ≥8 years showed no clinically significant difference in the occurrence of
37 dyspnoea between the participants who were receiving Colistin DPI (125 mg/ twice daily) and
38 those receiving Colistin inhalation solution (2 MU/ twice daily) at 8 weeks follow-up.

1 **Mortality**

2 No evidence was found for this important outcome.

3 **Emergence of resistant organisms**

4 No evidence was found for this important outcome.

5 **Comparison 3.3: Colistin versus Tobramycin**

6 **Lung function: FEV₁**

7 Very low quality evidence from 1 RCT with 109 people with cystic fibrosis and chronic *P*
8 *aeruginosa* infection > 7 years showed no clinically significant difference in lung function
9 (measured as mean % change in FEV₁ % predicted) between the participants receiving
10 Colistin (COLI neb 1MU/3 ml. twice daily) and those who were receiving Tobramycin (TOBI
11 neb 300 mg/5ml twice daily) at 1 to 3 months follow-up.

12 Low quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
13 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in lung function
14 (measured as mean % change in FEV₁ % predicted) between the participants receiving
15 Colistin (COLI DPI 120 mg. twice daily) and those who were receiving Tobramycin (TOBI neb
16 300 mg/5ml twice daily) at 4 and 12 weeks follow-up.

17 Low quality evidence from 2 RCTs with 658 people with cystic fibrosis and chronic *P*
18 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in lung function
19 (measured as mean % change in FEV₁ % predicted) between the participants receiving
20 Colistin and those who were receiving Tobramycin (COLI neb 1MU/3 ml. twice daily versus
21 TOBI neb 300 mg/5ml twice daily and COLI DPI 120 mg. twice daily versus TOBI neb 300
22 mg/5ml twice daily) at 24 weeks follow-up.

23 **Number of people with exacerbations**

24 NMA outcome

25 **Time to next exacerbation**

26 Very low quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
27 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the time to next
28 pulmonary exacerbation (measured as mean time to first additional anti-pseudomonal
29 treatment) between the participants receiving Colistin (COLI DPI 120 mg. twice daily) and
30 those who were receiving Tobramycin (TOBI neb 300 mg/5ml twice daily) during the 24
31 weeks duration of the trial.

32 **Suppression of the organism**

33 Low quality evidence from 1 RCT with 79 people with cystic fibrosis and chronic *P*
34 *aeruginosa* infection > 7 years showed no clinically significant difference in the suppression
35 of *P aeruginosa* (measured as change in sputum PA density log₁₀ cystic fibrosisU/ml)
36 between the participants receiving Colistin (COLI neb 1MU/3 ml. twice daily) and those who
37 were receiving Tobramycin (TOBI neb 300 mg/5ml twice daily) at 4 weeks follow-up.

38 **Nutritional status**

39 Low quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
40 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the BMI change

1 between the participants receiving Colistin (COLI DPI 120 mg. twice daily) and those who
2 were receiving Tobramycin (TOBI neb 300 mg/5ml twice daily) at 24 weeks follow-up.

3 **Quality of life**

4 Moderate quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
5 *aeruginosa* infection \geq 6 years showed no difference in change in quality of life (measured
6 with the individual domains of the CFQ-R questionnaire) between the participants receiving
7 Colistin (COLI DPI 120 mg. twice daily) and those who were receiving Tobramycin (TOBI neb
8 300 mg/5ml twice daily) at 24 weeks follow-up. The uncertainty for this outcome could not be
9 calculated.

10 **Mild adverse events**

11 Very low quality evidence from 1 RCT with 115 people with cystic fibrosis and chronic *P*
12 *aeruginosa* infection >7 years showed no clinically significant difference in the occurrence of
13 sputum changes, pharyngitis or cough between the participants receiving Colistin (COLI neb
14 1MU/3 ml. twice daily) and those who were receiving Tobramycin (TOBI neb 300 mg/5ml
15 twice daily) at 4 weeks follow-up.

16 Very low quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
17 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
18 productive cough, chest discomfort or vomiting between the participants receiving Colistin
19 (COLI DPI 120 mg. twice daily) and those who were receiving Tobramycin (TOBI neb 300
20 mg/5ml twice daily) at 24 weeks follow-up.

21 **Serious adverse events**

22 Very low quality evidence from 1 RCT with 115 people with cystic fibrosis and chronic *P*
23 *aeruginosa* infection >7 years showed no clinically significant difference in the number of
24 participants who experienced more than 1 serious adverse event between the participants
25 receiving Colistin (COLI neb 1MU/3 ml. twice daily) and those who were receiving
26 Tobramycin (TOBI neb 300 mg/5ml twice daily) at 4 weeks follow-up.

27 Very low quality evidence from 1 RCT with 115 people with cystic fibrosis and chronic *P*
28 *aeruginosa* infection >7 years showed no clinically significant difference in the occurrence of
29 dyspnoea between the participants receiving Colistin (COLI neb 1MU/3 ml. twice daily) and
30 those who were receiving Tobramycin (TOBI neb 300 mg/5ml twice daily) at 4 weeks follow-
31 up.

32 Moderate evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P aeruginosa*
33 infection \geq 6 years showed a clinically significant higher number of people withdrawn from
34 the study due to a serious adverse effect in the group of participants receiving Colistin (COLI
35 DPI 120 mg. twice daily) compared to those who were receiving Tobramycin (TOBI neb 300
36 mg/5ml twice daily) at 24 weeks follow-up.

37 Very quality evidence from 1 RCT with 374 people with cystic fibrosis and chronic *P*
38 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
39 dyspnoea or haemoptysis between the participants receiving Colistin (COLI DPI 120 mg.
40 twice daily) and those who were receiving Tobramycin (TOBI neb 300 mg/5ml twice daily) at
41 24 weeks follow-up.

42 **Mortality**

43 No evidence was found for this outcome.

1 **Emergence of resistant organisms**

2 Low quality evidence from 1 RCT with 115 people with cystic fibrosis and chronic *P*
3 *aeruginosa* infection >7 years reported that none of the patients in either treatment group
4 (COLI neb 1MU/3 ml. twice daily and TOBI neb 300 mg/5ml twice daily) developed highly
5 tobramycin-resistant *P aeruginosa* at 24 weeks follow-up.

6 **9.4.3.7 Fosfomycin (inhaled)**

7 No evidence was found for this treatment.

8 **9.4.3.8 Tobramycin (dry powder, inhaled)**

9 **Comparison 4.1: Tobramycin versus placebo**

10 **Lung function: FEV₁**

11 Low quality evidence from 4 RCTs with 516 children, young people and adults with cystic
12 fibrosis and chronic *P aeruginosa* infection showed a clinically significant improvement in
13 lung function (measured as mean % change in FEV₁ % predicted) in the group of participants
14 who were receiving tobramycin (TOBI DPI 112 mg daily, TOBI nebulised 300 mg or 600 mg
15 daily) compared to those who were receiving placebo at 1 to 3 months follow-up. Moderate
16 heterogeneity was found between the trials. Three trials showed a clinically significant
17 improvement in the tobramycin group, whereas 1 trial showed no differences.

18 **Number of people with exacerbations**

19 NMA outcome

20 **Time to next exacerbation**

21 No evidence was found for this outcome.

22 **Suppression of the organism**

23 High quality evidence from 3 RCTs with 357 people with cystic fibrosis and chronic *P*
24 *aeruginosa* infection ≥ 6 years showed a clinically significant higher number of people in
25 whom *P aeruginosa* was eradicated (measured as negative culture) in the group of
26 participants receiving Tobramycin (TOBI neb 300 mg or TOBI DPI 112 mg daily) compared
27 to those who were receiving placebo at 4 weeks follow-up. Low heterogeneity was observed
28 between the 3 trials, but all of them were consistent in showing a beneficial effect of
29 tobramycin compared to placebo.

30 High quality evidence from 1 RCT with 242 people with cystic fibrosis and chronic *P*
31 *aeruginosa* infection ≥ 6 years showed a clinically significant higher number of people in
32 whom *P aeruginosa* was eradicated (measured as negative culture) in the group of
33 participants receiving Tobramycin (TOBI neb 300 mg daily) compared to those who were
34 receiving placebo at 20 weeks follow-up. However, moderate quality evidence from the same
35 trial showed no clinically significant difference in the eradication of *P aeruginosa* 8, and 24
36 weeks follow-up.

37 Moderate quality evidence from 1 RCT with 59 people with cystic fibrosis and chronic *P*
38 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the eradication of
39 *P aeruginosa* (measured as negative culture) between the participants receiving Tobramycin
40 (TOBI neb 300 mg daily) and those who were receiving placebo at 6 weeks follow-up.

1 Moderate quality evidence from 1 RCT with 55 people with cystic fibrosis and chronic *P*
2 *aeruginosa* infection \geq 6 years showed a clinically significant decrease in *P*
3 *aeruginosa* sputum density (log₁₀ cystic fibrosisU/ml) in the group of participants receiving Tobramycin
4 (TOBI DPI 112 mg daily) compared to those who were receiving placebo at 4 weeks follow-
5 up.

6 Likewise, low quality evidence from another RCT with 95 people with cystic fibrosis and
7 chronic *P aeruginosa* infection > 6 years showed a clinically significant decrease in *P*
8 *aeruginosa* mucoid and non-mucoid sputum density (log₁₀ cystic fibrosisU/ml) in the group
9 of participants receiving tobramycin (TOBI DPI 112 mg) compared to those who were
10 receiving placebo at 4 weeks follow-up.

11 **Nutritional status: weight**

12 High quality evidence from 1 RCT with 59 people with cystic fibrosis and chronic *P*
13 *aeruginosa* infection \geq 6 years showed no clinically significant difference in weight (kg)
14 between the participants receiving Tobramycin (TOBI neb 300 mg daily) and those who were
15 receiving placebo at 12 weeks follow-up.

16 Likewise, moderate quality evidence from another RCT with 245 people with cystic fibrosis
17 and chronic *P aeruginosa* infection \geq 6 years showed no clinically significant difference in
18 weight (kg) between the participants receiving Tobramycin (TOBI neb 300 mg daily) and
19 those who were receiving placebo at 24 weeks follow-up.

20 **Quality of life**

21 No evidence was found for this outcome.

22 **Mild adverse events**

23 Very low quality evidence from 2 RCTs with 150 people with cystic fibrosis and chronic *P*
24 *aeruginosa* infection \geq 6 years showed a clinically significant lower occurrence of mild
25 adverse events in the group of participants receiving Tobramycin (TOBI DPI 112 mg daily)
26 compared to those who were receiving placebo at 4 weeks follow-up.

27 However, low quality evidence from 1 RCT with 245 people with cystic fibrosis and chronic *P*
28 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
29 minor adverse events between the participants receiving Tobramycin (TOBI neb 300 mg
30 daily) and those who were receiving placebo at 24 weeks follow-up.

31 Low quality evidence from 1 RCT with 55 people with cystic fibrosis and chronic *P*
32 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
33 auditory impairment between the participants receiving tobramycin (TOBI DPI 112 mg) and
34 those who were receiving placebo at 4 weeks follow-up. In addition,

35 Moreover, high quality evidence from 1 RCT with 300 people with cystic fibrosis and chronic
36 *P aeruginosa* infection \geq 6 years showed no cases of auditory impairment in either group
37 (TOBI neb 300 mg daily or placebo) at 24 weeks follow-up. No cases were identified either at
38 42 weeks follow-up (n=71).

39 Moderate quality evidence from 1 RCT with 520 people with cystic fibrosis and chronic *P*
40 *aeruginosa* infection \geq 6 years showed a clinically significant higher occurrence of tinnitus in
41 the group of participants receiving Tobramycin (TOBI neb 300 mg daily) compared to those
42 who were receiving placebo at 24 weeks follow-up.

43 Very low quality evidence from 2 RCTs with 150 people with cystic fibrosis and chronic *P*
44 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
45 cough between the participants receiving tobramycin (TOBI DPI 112 mg) and those who

1 were receiving placebo at 4 weeks follow-up. High unexplained heterogeneity was found
2 between both trials, although both of them showed no clinically significant differences
3 between both groups.

4 Moderate quality evidence from 1 RCT with 300 people with cystic fibrosis and chronic *P*
5 *aeruginosa* infection ≥ 6 years no clinically significant difference in the occurrence of tinnitus
6 between the participants receiving tobramycin (TOBI neb 300 mg daily) and those who were
7 receiving placebo 24 weeks follow-up.

8 Very low quality evidence from 1 RCT with 95 people with cystic fibrosis and chronic *P*
9 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
10 headaches between the participants receiving tobramycin (TOBI DPI 112 mg) and those who
11 were receiving placebo 4 weeks follow-up.

12 Very low quality evidence from 2 RCTs with 150 people with cystic fibrosis and chronic *P*
13 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
14 serious adverse events between the participants receiving tobramycin (TOBI DPI 112 mg)
15 and those who were receiving placebo at 4 weeks follow-up.

16 However, high quality evidence from 1 RCT with 246 people with cystic fibrosis and chronic
17 *P aeruginosa* infection ≥ 6 years showed a clinically significant lower occurrence of serious
18 adverse events in the group of participants receiving Tobramycin (TOBI neb 300 mg daily)
19 compared to those who were receiving placebo at 24 weeks follow-up.

20 Very low quality evidence from 1 RCT with 95 people with cystic fibrosis and chronic *P*
21 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
22 haemoptysis between the participants receiving tobramycin (TOBI DPI 112 mg) and those
23 who were receiving placebo 4 weeks follow-up.

24 Likewise, moderate quality evidence from one RCT with 520 people with cystic fibrosis and
25 chronic *P aeruginosa* infection ≥ 6 years showed no clinically significant difference in the
26 occurrence haemoptysis between the participants receiving Tobramycin (TOBI neb 300 mg
27 daily) and those who were receiving placebo at 24 weeks follow-up.

28 Low quality evidence from 1 RCT with 520 people with cystic fibrosis and chronic *P*
29 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence
30 pneumothorax between the participants receiving Tobramycin (TOBI neb 300 mg daily) and
31 those who were receiving placebo at 24 weeks follow-up.

32 **Mortality**

33 Low quality evidence from 1 RCT with 95 people with cystic fibrosis and chronic *P*
34 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in mortality between
35 the participants receiving tobramycin (TOBI DPI 112 mg) and those who were receiving
36 placebo 4 weeks follow-up.

37 Likewise, moderate quality evidence from two RTCs with 839 people with cystic fibrosis and
38 chronic *P aeruginosa* infection ≥ 6 years showed no clinically significant difference in the
39 mortality rate between the participants receiving Tobramycin (TOBI neb 300 mg daily) and
40 those who were receiving placebo at 24 weeks follow-up.

41 **Emergence of resistant organisms**

42 Very low quality evidence from 2 RCTs with 672 people with cystic fibrosis and chronic *P*
43 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the number of
44 participants in whom tobramycin-resistant *P aeruginosa* was isolated between the group of
45 participants receiving Tobramycin (TOBI neb 300 mg daily) and those who were receiving
46 placebo at 24 weeks follow-up. Significant unexplained heterogeneity was found between

1 both trials, with 1 of them showing a clinically significant harmful effect of tobramycin, and the
2 other trial showing no differences between both groups.

3 High quality evidence from 1 RCT with 520 people with cystic fibrosis and chronic *P*
4 *aeruginosa* infection \geq 6 years showed that drug-resistant *B Cepacia* was not isolated in the
5 participants of either group (TOBI neb 300 mg daily or placebo) at 24 weeks follow-up

6 Low quality evidence from 1 RCT with 520 people with cystic fibrosis and chronic *P*
7 *aeruginosa* infection \geq 6 years showed that drug-resistant *B Cepacia* was not isolated in the
8 participants of either group (TOBI neb 300 mg daily or placebo) at 24 weeks follow-up

9 Low quality evidence from 1 RCT with 520 people with cystic fibrosis and chronic *P*
10 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the frequency of
11 new isolates of drug-resistant *A. Xiloxidans* between the participants who were receiving
12 Tobramycin (TOBI neb 300 mg daily) and those who were receiving placebo at 24 weeks
13 follow-up.

14 High quality evidence from 1 RCT with 389 people with cystic fibrosis and chronic *P*
15 *aeruginosa* infection \geq 6 years showed a clinically significant lower number of patients in
16 whom new isolates of drug-resistant *Aspergillus* was isolated in the group of participants
17 receiving Tobramycin (TOBI neb 300 mg daily) compared to those who were receiving
18 placebo at 24 weeks follow-up.

19 **Comparison 4.2: Tobramycin DPI versus tobramycin nebulised**

20 **Lung function: FEV₁**

21 Low quality evidence from 1 RCT with 517 people with cystic fibrosis and chronic *P*
22 *aeruginosa* infection \geq 6 years showed no clinically significant difference in lung function
23 (measured as % mean change in FEV₁% predicted) between the participants receiving
24 tobramycin DPI (112 mg/ twice daily) and those receiving tobramycin inhalation solution (300
25 mg/ 5ml twice daily) at 4, 20 and 24 weeks follow-up.

26 **Number of people with exacerbations**

27 NMA outcome

28 **Time to next exacerbation**

29 No evidence was found for this critical outcome.

30 **Suppression of the organism**

31 Low to moderate quality evidence from 1 RCT with 517 people with cystic fibrosis and
32 chronic *P aeruginosa* infection \geq 6 years showed no clinically significant difference in change
33 in sputum density (measured as log₁₀ CFU) in the group of participants receiving tobramycin
34 DPI (112 mg/ twice daily) compared with those receiving tobramycin inhalation solution (300
35 mg/ 5ml twice daily) at 4 and at 20 weeks follow-up.

36 **Nutritional status**

37 No evidence was found for this important outcome.

38 **Quality of life**

39 No evidence was found for this important outcome.

1 **Mild adverse events**

2 Moderate quality evidence from 1 RCT with 517 people with cystic fibrosis and chronic *P*
3 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
4 mild adverse events (any) between the participants receiving tobramycin DPI (112 mg/ twice
5 daily) and those receiving tobramycin inhalation solution (300 mg/ 5ml twice daily) at 24
6 weeks follow-up.

7 Very low quality evidence from 1 RCT with 517 people with cystic fibrosis and chronic *P*
8 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
9 productive cough, headache or vomiting between the participants receiving tobramycin DPI
10 (112 mg/ twice daily) and those receiving tobramycin inhalation solution (300 mg/ 5ml twice
11 daily) at 24 weeks follow-up.

12 **Serious adverse events**

13 Low quality evidence from 1 RCT with 517 people with cystic fibrosis and chronic *P*
14 *aeruginosa* infection ≥ 6 years no clinically significant difference in the occurrence of serious
15 adverse events (any) in the group of participants receiving tobramycin DPI (112 mg/ twice
16 daily) compared with those receiving tobramycin inhalation solution (300 mg/ 5ml twice daily)
17 at 24 weeks follow-up.

18 Very low quality evidence from 1 RCT with 517 people with cystic fibrosis and chronic *P*
19 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
20 dyspnoea and haemoptysis between the participants receiving tobramycin DPI (112 mg/
21 twice daily) and those receiving tobramycin inhalation solution (300 mg/ 5ml twice daily) at 24
22 weeks follow-up.

23 **Mortality**

24 No evidence was found for this important outcome.

25 **Emergence of resistant organisms**

26 No evidence was found for this important outcome.

27 **Comparison 4.3: Tobramycin versus Aztreonam lysine**

28 **Lung function: FEV₁**

29 Moderate quality evidence from 1 RCT with 268 people with cystic fibrosis and chronic *P*
30 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in lung function
31 (measured as % change in FEV₁ % predicted) between the group of patients receiving
32 tobramycin inhalation solution (300 mg/ twice daily) and the patients receiving aztreonam
33 lysine (75g/ 3-times daily) at 3 months follow-up.

34 **Number of people with exacerbations**

35 NMA outcome

36 **Time to next exacerbation**

37 No evidence was found for this critical outcome.

1 **Suppression of the organism**

2 Low quality evidence from 1 RCT with 194 people with cystic fibrosis and chronic *P*
3 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the change of
4 sputum density (measured as log₁₀ cystic fibrosisU) between the group of patients receiving
5 Tobramycin inhalation solution (300 mg/ twice daily) and the patients receiving Aztreonam
6 lysine (75g/ 3-times daily) at 20 weeks follow-up.

7 **Nutritional status: weight**

8 Low quality evidence from 1 RCT with 268 people with cystic fibrosis and chronic *P*
9 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the percentage of
10 weight change (kg) between the group of patients receiving Tobramycin inhalation solution
11 (300 mg/ twice daily) and the patients receiving Aztreonam lysine (75g/ 3-times daily) at 24
12 weeks follow-up.

13 **Quality of life**

14 Low quality evidence from 1 RCT with 261 people with cystic fibrosis and chronic *P*
15 *aeruginosa* infection ≥ 6 years showed a clinically significant decrease in quality of life
16 (measured as change from baseline in the CFQ-R questionnaire respiratory domain) in the
17 group of participants receiving tobramycin inhalation solution (300 mg/ twice daily) compared
18 to those receiving Aztreonam lysine (75g/ 3-times daily) at 20 weeks follow-up.

19 **Mild adverse events**

20 Very low to low quality evidence with 268 people with cystic fibrosis and chronic *P*
21 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in the occurrence of
22 chest discomfort, cough, headache and vomiting between the group of patients receiving
23 Tobramycin inhalation solution (300 mg/ twice daily) and the patients receiving Aztreonam
24 lysine (75g/ 3-times daily) at 3 months follow-up.

25 **Serious adverse events**

26 Low quality evidence with 268 people with cystic fibrosis and chronic *P aeruginosa* infection
27 ≥ 6 years showed no clinically significant difference in the occurrence of dyspnoea and
28 haemoptysis between the group of patients receiving Tobramycin inhalation solution (300
29 mg/ twice daily) and the patients receiving Aztreonam lysine (75g/ 3-times daily) at 3 months
30 follow-up.

31 **Mortality**

32 No evidence was found for this outcome.

33 **Emergence of resistant organisms**

34 No evidence was found for this outcome.

35 **Comparison 5. Combination of fosfomycin + tobramycin versus placebo**

36 **Lung function: FEV₁**

37 Moderate quality evidence from 1 RCT with 70 adults with cystic fibrosis and chronic *P*
38 *aeruginosa* infection showed a clinically significant improvement in lung function (measured
39 as relative change in FEV₁ % predicted) in the group of participants receiving combination of
40 fosfomacy and tobramacyin (80/ 20 mg) compared to those receiving placebo at 4 weeks
41 follow-up.

1 Likewise, low quality evidence from the same trial (N=73) showed a clinically significant
2 improvement in lung function (measured as relative change in FEV₁ % predicted) in the
3 group of participants receiving combination of fosfomacy and tobramacyin (160/ 40 mg)
4 compared to those receiving placebo at 4 weeks follow-up.

5 **Number of people with exacerbations**

6 NMA outcome

7 **Time to next exacerbation**

8 No evidence was found for this critical outcome.

9 **Suppression of the organism**

10 Low quality evidence from 1 RCT with 70 adults with cystic fibrosis and chronic *P aeruginosa*
11 infection showed a clinically significant decrease in sputum *P aeruginosa* density (log₁₀
12 CFU/g) in the group of participants receiving combination of fosfomacy and tobramacyin (80/
13 20 mg) compared to those receiving placebo at 4 weeks follow-up.

14 However, low quality evidence from the same trial (N=73) showed no clinically significant
15 difference between a combination of fosfomacy and tobramacyin (160/ 40 mg) and placebo
16 at 4 weeks follow-up.

17 **Nutritional status**

18 No evidence was found for this important outcome.

19 **Quality of life**

20 No evidence was found for this important outcome.

21 **Mild adverse events**

22 No evidence was found for this important outcome.

23 **Serious adverse events**

24 No evidence was found for this important outcome.

25 **Mortality**

26 No evidence was found for this important outcome.

27 **Emergence of resistant organisms**

28 No evidence was found for this important outcome.

29 **Comparison 6. Continuous alternating therapy versus intermittent treatment:**
30 **aztreonam lysine + tobramycin or placebo + tobramycin**

31 **Lung function: FEV₁**

32 Moderate quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
33 *aeruginosa* infection ≥ 6 years showed no clinically significant difference in lung function
34 (measure as % change in FEV₁ % predicted) between the participants receiving continuous
35 alternating therapy (tobramycin inhalation solution 300 mg daily for 28 days, followed by

1 aztreonam lysine) and those on an intermittent regimen (tobramycin inhalation solution 300
2 mg daily for 28 days, followed by placebo) at 20 weeks follow-up. Values at 4, 12 and 20
3 weeks were averaged.

4 **Number of people with exacerbations**

5 NMA outcome

6 **Time to next exacerbation**

7 Low quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
8 *aeruginosa* infection \geq 6 years showed no clinically significant difference in time to next
9 exacerbation between the participants receiving continuous alternating therapy (tobramycin
10 inhalation solution 300 mg daily for 28 days, followed by aztreonam lysine) and those on an
11 intermittent regimen (tobramycin inhalation solution 300 mg daily for 28 days, followed by
12 placebo).

13 **Suppression of the organism**

14 No evidence was found for this important outcome.

15 **Nutritional status**

16 No evidence was found for this important outcome.

17 **Quality of life**

18 Low quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
19 *aeruginosa* infection \geq 6 years showed no clinically significant difference in quality of life
20 (measured with the CFQ-R questionnaire) between the participants receiving continuous
21 alternating therapy (tobramycin inhalation solution 300 mg daily for 28 days, followed by
22 aztreonam lysine) and those on an intermittent regimen (tobramycin inhalation solution 300
23 mg daily for 28 days, followed by placebo) at 20 weeks follow-up. Values at 4, 12 and 20
24 weeks were averaged.

25 **Mild adverse events**

26 Low quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
27 *aeruginosa* infection \geq 6 years showed a clinically significant increase in the occurrence of
28 cough in the group of participants receiving continuous alternating therapy (tobramycin
29 inhalation solution 300 mg daily for 28 days, followed by aztreonam lysine) compared to
30 those on an intermittent regimen (tobramycin inhalation solution 300 mg daily for 28 days,
31 followed by placebo) at 3 months follow-up.

32 **Serious adverse events**

33 Low quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
34 *aeruginosa* infection \geq 6 years showed no clinically significant difference in the occurrence of
35 dyspnoea between the participants receiving continuous alternating therapy (tobramycin
36 inhalation solution 300 mg daily for 28 days, followed by aztreonam) and those on an
37 intermittent regimen (tobramycin inhalation solution 300 mg daily for 28 days, followed by
38 placebo) at 3 months follow-up.

39 Very low quality evidence from 1 RCT with 88 people with cystic fibrosis and chronic *P*
40 *aeruginosa* infection \geq 6 years showed a clinically significant increase in the occurrence of
41 serious adverse events (no treatment related) in the group of participants receiving
42 continuous alternating therapy (tobramycin inhalation solution 300 mg daily for 28 days,

1 followed by aztreonam lysine) compared to those on an intermittent regimen (tobramycin
2 inhalation solution 300 mg daily for 28 days, followed by placebo) at 3 months follow-up.

3 **Mortality**

4 No evidence was found for this important outcome.

5 **Emergence of resistant organisms**

6 No evidence was found for this important outcome.

7 **79.4.3.8.1 Antimicrobial regimens for the treatment of chronic *S aureus***

8 **Cefradine (oral)**

9 No evidence was found for this treatment.

10 **Cotrimoxazole (oral)**

11 No evidence was found for this treatment.

12 **Doxycycline (oral)**

13 No evidence was found for this treatment.

14 **Flucloxacillin (oral)**

15 No evidence was found for this treatment.

16 **169.4.3.8.2 Antimicrobial regimens for the treatment of chronic *B cepacia* complex**

17 **Ceftazidime (inhaled)**

18 No evidence was found for this treatment.

19 **Cotrimoxazole (oral)**

20 No evidence was found for this treatment.

21 **Imipenem (oral)**

22 No evidence was found for this treatment.

23 **Meropenem (inhaled)**

24 No evidence was found for this treatment.

25 **Trimethoprim (oral)**

26 No evidence was found for this treatment.

27 **279.4.3.8.3 Antimicrobial regimens for the treatment of chronic *A fumigatus***

28 **Amphotericin (inhaled)**

29 No evidence was found for this treatment.

1 **Itraconazole (oral)**

2 **Comparison 7: Itraconazole versus placebo**

3 **Lung function: FEV₁**

4 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis and chronically
5 colonised with *A fumigatus* ≥ 6 years showed no clinically significant difference in lung
6 function (measured as percentage change in FEV₁ predicted from baseline) between the
7 participants who were receiving oral Itraconazole (5mg/kg once or twice daily) and those who
8 were receiving placebo after 24 or 48 weeks follow-up.

9 **Exacerbations**

10 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis and chronically
11 colonised with *A fumigatus* ≥ 6 years showed no clinically significant difference in the time to
12 next exacerbation between the participants who were receiving oral Itraconazole (5mg/kg
13 once or twice daily) and those who were receiving placebo.

14 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis and chronically
15 colonised with *A fumigatus* ≥ 6 years showed no clinically significant difference in the number
16 of patients requiring oral or IV antibiotics due to a pulmonary exacerbation between the
17 participants who were receiving oral Itraconazole (5mg/kg once or twice daily) and those who
18 were receiving placebo after 24 or 48 weeks follow-up.

19 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis and chronically
20 colonised with *A fumigatus* ≥ 6 years showed no clinically significant difference in the number
21 of patients admitted to hospital due to a pulmonary exacerbation between the participants
22 who were receiving oral Itraconazole (5mg/kg once or twice daily) and those who were
23 receiving placebo after 24 or 48 weeks follow-up.

24 **Suppression of the organism**

25 No evidence was found for this important outcome.

26 **Nutritional status**

27 No evidence was found for this important outcome.

28 **Quality of life**

29 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis and chronically
30 colonised with *A fumigatus* ≥ 6 years showed no differences in quality of life (measured as
31 change in CFQ-R total score and respiratory domain) between the participants who were
32 receiving oral Itraconazole (5mg/kg once or twice daily) and those who were receiving
33 placebo after the 24-week treatment duration. The uncertainty around this outcome could not
34 be calculated.

35 **Mild adverse events**

36 Very low evidence from one RCT with 35 people with cystic fibrosis and chronically colonised
37 with *A fumigatus* ≥ 6 years showed no clinically significant difference in the occurrence of
38 minor adverse events (including: increased dyspnoea, rash, hyperglycaemia, flu-like illness,
39 diarrhoea and conjunctivitis) between the participants who were receiving oral Itraconazole
40 (5mg/kg once or twice daily) and those who were receiving placebo during the 24-week
41 treatment duration.

1 **Serious adverse events**

2 Very low quality evidence from one RCT with 35 people with cystic fibrosis and chronically
3 colonised with *A fumigatus* \geq 6 years showed no clinically significant difference in the
4 occurrence of major adverse events (including: haemoptysis and spontaneous
5 pneumothorax) between the participants who were receiving oral Itraconazole (5mg/kg once
6 or twice daily) and those who were receiving placebo during the 24-week treatment duration.

7 **Mortality**

8 No evidence was found for this important outcome.

9 **Emergence of resistant organisms**

10 No evidence was found for this important outcome.

11 **Posaconazole (oral)**

12 No evidence was found for this treatment.

13 **Voriconazole (oral)**

14 No evidence was found for this treatment.

159.4.3.8.4 **Economic evidence statements**

16 One cost-benefit analysis (Iles 2003) on people with cystic fibrosis in the UK, compared 12
17 months before nebulised tobramycin use with 12 months during nebulised tobramycin use.
18 They found that the introduction of nebulised tobramycin reduced the cost of hospital
19 attendances and parenteral antibiotics, but did not offset the cost of nebulised tobramycin.
20 This analysis is directly applicable given that the type of economic evaluation is unlikely to
21 change the conclusions about cost-effectiveness and all other applicability criteria are met.
22 The evidence is associated with serious limitations from the before and after type study used
23 to inform the analysis.

24 One cost-utility analysis (Tappenden 2013) on people with cystic fibrosis in the UK, found
25 that if colistimethate sodium dry powder is priced lower than that of nebulised tobramycin the
26 ICER lies in the south-west quadrant of the cost-effectiveness plane reflecting a QALY loss
27 and cost savings for colistimethate sodium dry powder compared with nebulised tobramycin.
28 However, if colistimethate sodium dry powder is priced higher than that of nebulised
29 tobramycin the incremental cost is positive, and colistimethate sodium dry powder is
30 dominated by nebulised tobramycin. The lifetime horizon and 'within-trial' analysis resulted in
31 smaller incremental differences in both costs and QALYs, but led to the same conclusions.
32 This analysis is directly applicable with minor limitations.

33 One cost-utility analysis (Tappenden 2014) on people with cystic fibrosis in the UK, over a
34 lifetime horizon, found that using list prices, nebulised tobramycin dominated colistimethate
35 sound dry powder and tobramycin dry powder has an ICER of £123,563 compared to
36 nebulised tobramycin. When the revised patient access scheme discount was applied to
37 colistimethate sodium dry powder, nebulised tobramycin was more expensive and less
38 effective with an ICER of £288,563. When the revised patient access scheme discount was
39 applied to tobramycin dry powder, it dominated nebulised tobramycin. This analysis is
40 directly applicable with minor limitations.

41 One cost-utility analysis (Schechter 2015) on people with cystic fibrosis in the US, found that
42 aztreonam dominated nebulised tobramycin over a 3 year time horizon. This analysis is
43 partially applicable, due to the US third party payer perspective taken. The evidence is

1 associated with serious limitations, including the potential conflict of interest and lack of detail
2 in their methods.

3 The economic model developed for this review, found that nebulised colistimethate sodium,
4 nebulised tobramycin and tobramycin dry powder were not cost-effective compared to
5 placebo in the base case (list prices). Tobramycin dry powder was also dominated (more
6 expensive and less effective) by the treatments in this comparison.

7 The economic model developed for this review, found that nebulised tobramycin is not cost-
8 effective compared to nebulised colistimethate sodium, in the base case (list prices), with an
9 ICER of £602,472.

10 The economic model developed for this review, found that nebulised tobramycin dominated
11 (more effective and less expensive) colistimethate sodium dry powder, in the base case (list
12 prices).

13 The economic model developed for this review, found that nebulised aztreonam is not cost-
14 effective compared to nebulised tobramycin, in the base case (list prices), with an ICER of
15 £34,348. The combination treatment (28 days aztreonam lysine alternating with 28 days
16 nebulised tobramycin) was also dominated (more expensive and less effective) by
17 aztreonam lysine in this comparison.

18 **9.4.3.9 Evidence to recommendations**

19 **9.4.3.9.1 Relative value placed on the outcomes considered**

20 The aim of this review was to determine the clinical and cost-effectiveness of different
21 antimicrobial treatment regimens to suppress chronic pulmonary infection in people with
22 cystic fibrosis.

23 The Committee identified lung function and time to next pulmonary exacerbation as critical
24 outcomes for this evidence review. Where no evidence was found for time to next pulmonary
25 exacerbation, number of people experiencing a pulmonary exacerbation and number of
26 hospital due to a pulmonary exacerbation were taken as proxy outcomes instead.
27 Suppression of the organism, nutritional status, quality of life, adverse events and
28 emergence of resistant organisms were rated as important outcomes.

29 **9.4.3.9.2 Consideration of clinical benefits and harms**

30 The Committee discussed the recommendations for each pathogen separately.

31 **Chronic *P. Aeruginosa***

32 The Committee discussed the results from the network meta-analysis. They noted that it was
33 not possible to conduct network meta-analysis for the critical outcome FEV1 due to high
34 unexplained heterogeneity.

35 The results from the NMA suggested that Aztreonam lysine was more effective than placebo
36 or tobramycin at reducing the odds of experiencing a pulmonary exacerbation. This result
37 has to be taken with caution, as there was considerable uncertainty in the network.

38 The Committee also discussed the results from the review (conventional pair-wise meta-
39 analysis). They first reviewed the evidence comparing treatments against placebo.

40 With regards to colistin, the evidence showed that inhaled colistin was no better than placebo
41 with regards to lung function. The Committee noted this evidence was of low quality, and
42 came from a small single trial. Moderate quality evidence showed that inhaled colistin was
43 not associated with the emergence of resistant organisms when compared to placebo. They
44 also looked at the evidence comparing different routes of administration. Very low quality

1 evidence from 1 trial showed no differences between colistin DPI and inhaled colistin in lung
2 function and in the occurrence of adverse events.

3 The Committee also discussed the evidence comparing tobramycin and placebo, the
4 Committee noted tobramycin was associated with a clinically significant improvement in lung
5 function at 1 to 3 months. However moderate heterogeneity was found between trials, and
6 therefore they agreed this result should be interpreted with caution. With regards to
7 suppressing the organisms, the evidence showed contradicting results. High quality evidence
8 showed that tobramycin was better than placebo at suppressing the organism (measured as
9 having a negative culture) at 4 and 20 weeks follow-up. In addition, low to moderate quality
10 evidence showed a clinically significant reduction in sputum density at 4 weeks follow-up.
11 However, moderate quality evidence showed no clinically significant difference in the
12 eradication of *P aeruginosa* 6, 8, and 24 weeks follow-up. The evidence regarding side
13 effects was not conclusive either, but tobramycin was associated with an increased risk in
14 the occurrence of mild adverse events, including tinnitus. They also looked at the evidence
15 comparing different routes of administration. They noted there were no clinically significant
16 differences between tobramycin DPI and inhaled tobramycin in lung function, change in
17 sputum density, or adverse events. The quality of the evidence range from very low to
18 moderate.

19 With regards to ciprofloxacin, the Committee noted there was no evidence for the critical
20 outcome lung function. No differences were found in weight.

21 They noted there was moderate quality evidence showing that aztreonam lysine was better
22 than placebo for lung function, weight and quality of life. In addition, no clinically significant
23 differences were found in the occurrence of adverse events, or in the emergence of resistant
24 organisms.

25 The Committee also reviewed the evidence comparing colistin and tobramycin. Very low to
26 low quality evidence showed no clinically significant differences in lung function at 4, 12 and
27 24 weeks. Likewise, very low quality evidence showed no differences in time to next
28 additional anti-pseudomonal treatment. With regards to the important outcomes, no clinically
29 significant differences were found for suppression of the organism (measured as change in
30 sputum density), weight, quality of life, adverse events and emergence of resistant
31 organisms. The quality of the evidence ranged from very low to moderate.

32 The Committee reviewed the evidence comparing tobramycin and aztreonam. Moderate
33 quality evidence showed no clinically significant differences in lung function at 3 months
34 follow-up. Low quality evidence showed no differences in change in sputum density or
35 quality of life at 20 weeks follow-up, and in weight at 24 weeks. Finally, no differences were
36 found in the occurrence of adverse events at 3 months.

37 The Committee noted the combination of fosfomycin and high low-dose tobramycin was
38 found to be significantly better than placebo in lung function at 4 weeks follow-up. This
39 evidence came from a single trial, and was rated as of low to moderate quality. Low quality
40 evidence from the same trial showed a clinically significant decrease in sputum density in the
41 group receiving low-dose tobramycin, but this different was not clinically significant in the
42 high-dose group.

43 Finally the Committee discussed the comparison between continuous alternating therapy
44 with tobramycin followed by aztreonam and intermittent treatment with tobramycin followed
45 by placebo. The evidence showed no differences between both treatment regimens in lung
46 function (moderate quality), time to next exacerbation (low quality) and quality of life (low
47 quality) at 20 weeks. However, they noted the risk of adverse events was higher in the
48 participants in the continuous alternating therapy group.

49 The Committee noted no evidence was found for high-dose azithromycin and fosfomycin.

1 The findings of the clinical evidence were discussed in the light of the economic evidence. In
2 addition, the Committee discussed the current recommendations from the NICE TA report
3 276 (Tappenden 2013) that looked at colistin and tobramycin for the treatment of chronic P
4 aeruginosa in people with cystic fibrosis over the age of 6 years.

5 As recommended in the NICE TA report 276 the Committee agreed that colistin should be
6 used as first line treatment and can be given as dry powder for inhalation to those people
7 who cannot tolerate it in its nebulised form. This recommendation is consistent with clinical
8 practice, and it also consistent with the CF Trust consensus recommendations.

9 The Committee discussed whether aztreonam lysine or tobramycin should be given as
10 second line treatment in case clinical deterioration occurs despite regular colistin. Clinical
11 deterioration was considered either an increase in the number of exacerbations or a decline
12 in pulmonary function (determined by spirometry). In line with NICE TA report 276, the
13 Committee agreed to recommend either aztreonam lysine, nebulised tobramycin, or
14 tobramycin dry powder (see also section on economic benefits and harms). This is because
15 although they noted the NMA suggested aztreonam was better than placebo or tobramycin,
16 there was lots of uncertainty regarding the results. In addition, the available direct evidence
17 comparing aztreonam lysine and tobramycin did not favour either treatment.

18 The Committee discussed their understanding that the effect of an antimicrobial may
19 diminish over time with repeated exposure, which may account for improvements seen in
20 clinical trials with new agents in treatment-naïve populations. Therefore it would be
21 appropriate to change between agents in line with an individual patient's response.

22 The Committee noted that adherence to treatment can be a relevant issue, and it should be
23 considered when prescribing treatment.

24 The Committee noted that in practice, combinations of inhaled antimicrobials may be
25 prescribed on alternate cycles, for example, colistin alternating with tobramycin or aztreonam
26 lysine, or tobramycin alternating with aztreonam lysine. However in the absence of evidence,
27 they did not write a recommendation.

28 The Committee noted that other inhaled antimicrobials were in development, but did not form
29 part of this review as the data was not available when the review was undertaken (for
30 example, levofloxacin inhalation solution). The omission of levofloxacin from the
31 recommendations does not reflect a decision not to recommend it, but rather the fact that it
32 was not included in this review.

33 **Chronic *S aureus***

34 No evidence was found for the treatment of chronic *S aureus*, therefore recommendations
35 were based on Committee's clinical expertise. They noted the Cochrane review is empty.

36 The Committee made separate recommendations for MSSA and MRSA. This is because
37 MRSA treatment is more complex than MSSA treatment.

38 The Committee's consensus was that long term oral antibiotic treatment to suppress MSSA
39 could be justified whether the person is unwell or not, because this is a recognised and
40 important pulmonary pathogen in children and adults. Suppressing it might be expected to
41 reduce the risk of progressive lung disease and of acute exacerbations caused by this
42 infection. The choice of treatment would depend on disease severity.

43 With regards to MRSA, the Committee agreed that there is no need to routinely give
44 antibiotic treatment to suppress infection in people with chronic MRSA who are stable. They
45 noted it is important to explain the risks and benefits of treatment to the person with cystic
46 fibrosis and their families. However, they agreed that if *S aureus* is repeatedly isolated from a
47 patient's respiratory samples and the lung function is deteriorating a course of antibiotics
48 could be considered. This is because MRSA can be fatal in people who are unwell./

1 The Committee acknowledged the infection control guideline (CG 139), and agreed that
2 many of the principles are also applicable to people with cystic fibrosis.

3 **Chronic *B cepacia* complex**

4 No evidence was found for the treatment of *B cepacia* complex, therefore recommendations
5 were based on Committee's clinical expertise. The Committee noted that only a small
6 number of people are infected with *B cepacia* complex, and most published studies are
7 anecdotal reports.

8 The Committee agreed that there is a strong emphasis in cross-infection prevention to
9 prevent the spread of *B cepacia* complex between people with cystic fibrosis. However, they
10 noted variability in the way people are treated.

11 The Committee agreed there is no need to give treatment to people with chronic *B cepacia*
12 complex who are stable. They noted it is important to discuss this with the person with cystic
13 fibrosis and their family, and to provide reassurance.

14 The Committee agreed treatment could be considered for people with chronic *B cepacia*
15 complex who are experiencing an exacerbation, or whose lung function is deteriorating.

16 The Committee noted the used of inhaled antibiotics can be considered, however they
17 agreed it is important to seek specialist microbiological advice on which antibiotic to use, as
18 healthcare professionals may have limited experience dealing with BCC.

19 Finally the Committee agreed treatment should be stopped if no benefit is observed.

20 **Chronic *A fumigatus***

21 The Committee acknowledged the evidence was scarce and of poor quality. They noted
22 therapeutic azole levels were not achieved in many participants. Therefore recommendations
23 were mainly based on the Committee's clinical knowledge and experience.

24 The Committee agreed that in people who are stable, there is no need to treat with antifungal
25 agents to suppress infection. This is because it is known that chronic aspergillus colonisation
26 can exist without associated deterioration in lung function, and treatment of this organism
27 can be difficult and associated with adverse events.

28 The Committee discussed that in people who are chronically infected with *A fumigatus* and
29 deteriorating without an obvious explanation, a trial with an antifungal agent can be
30 considered. They agreed to recommend a trial with an antifungal agent, because this
31 pathogen could be the cause of the deterioration in the lung function. They noted that their
32 first line choice was itraconazole but that therapeutic levels of this agent are particularly
33 difficult to achieve and it may be necessary to change to another antifungal such as
34 voriconazole or posaconazole. Regardless, the choice of the antifungal agent should take *in*
35 *vitro* sensitivities into account to ensure the optimal drug is used. They also acknowledged
36 that advice may be sought from a specialist microbiologist to inform choice. Finally, they
37 highlighted that clinical response should be appropriately assessed, and that treatment could
38 be stopped or modified if no benefit was observed.

39 The Committee acknowledge that it was important to distinguish between those who have
40 respiratory symptoms due to infection with aspergillus and those who have clinical
41 manifestations due to allergic sensitisation to aspergillus. For people with cystic fibrosis who
42 have elevated aspergillus serology and declining pulmonary function and whose pulmonary
43 treatment is optimised, the Committee agreed that clinicians should think about a diagnosis
44 of allergic bronchopulmonary aspergillosis (ABPA), especially if a chest x-ray or CT scan
45 shows changes consistent with ABPA.

19.4.3.9.3 **Consideration of economic benefits and harms**

2 **Antimicrobial regimens for the treatment of chronic *P aeruginosa***

3 The Committee agreed that the use of a lifetime horizon in the economic model was
4 appropriate, but acknowledged the limitation of extrapolating short-term trial results over a
5 lifetime horizon. The Committee also noted that the model did not reflect the current
6 treatment pathway where some people switch their treatment or receive a combination of
7 treatments. However, the Committee agreed that there was no clinical effectiveness data
8 available on treatment switching or combinations to inform the model beyond the trial by
9 Flume 2016. As a result, the Committee also considered the “with-in” trial analysis that did
10 not extrapolate data to a lifetime horizon to address those limitations. The Committee also
11 noted that trial participants were not treatment naïve which may underestimate the benefits
12 and cost-effectiveness of those treatments in a naïve population.

13 The Committee considered the impact of treatment adherence on cost-effectiveness, as
14 cost-effectiveness may be overestimated beyond a trial setting, if the same benefits are not
15 achieved. The Committee advised that colistimethate sodium has a lower adherence than
16 tobramycin because it requires more time and effort to administer. On the other hand, the
17 Committee also believed colistimethate tasted better than tobramycin which may increase
18 adherence to colistimethate compared to tobramycin, especially in children. As a result, there
19 are issues in both directions that could cancel out with little influence on the model. The
20 Committee also added that people with cystic fibrosis may be more likely to adhere to a dry
21 powder inhalation treatment than a nebulised treatment in view of the speed and
22 convenience of drug delivery. However, newer nebulisers with quicker delivery time, such as
23 the PARI eFlow jet nebuliser, are increasing in use. For these reasons, it remains unclear
24 whether dry powder inhalers would reduce treatment burden compared to newer, quicker
25 nebulisers.

26 It is accepted current practice to offer people infected with chronic *P aeruginosa*,
27 antimicrobial treatment to prevent deterioration in their lung function. Despite this, the
28 Committee discussed how the benefits of antimicrobial treatment may not outweigh their
29 costs. As a result, the Committee questioned if current practice should be changed based on
30 the economic model that found a small decrease in effectiveness for a large cost saving.
31 Following this, the Committee acknowledged that current NICE HTA recommendations state
32 nebulised colistimethate sodium should be offered as a first line option. However, the
33 sources of evidence considered in NICE TA276 did not compare nebulised colistimethate
34 sodium to “no treatment” which again questions if nebulised colistimethate sodium should be
35 recommended.

36 If “no treatment” cannot be accepted as an option, the Committee agreed that nebulised
37 colistimethate sodium would be the most cost-effective antimicrobial and tobramycin dry
38 powder would be the least as it was dominated (less effective and more expensive) by the
39 other options in the economic model.

40 The Committee agreed that it was reasonable to assume the exacerbation rate for nebulised
41 colistimethate sodium was equal to nebulised tobramycin in the absence of data. The
42 Committee also agreed that given the current clinical pathway, they would have liked to have
43 seen effectiveness evidence comparing nebulised colistimethate sodium, colistimethate dry
44 powder and “no treatment”, especially as this evidence was not available during the
45 submission of evidence in NICE TA276. To reduce this uncertainty, the Committee
46 considered a research recommendation to assess the clinical and cost-effectiveness of those
47 options. However, the Committee acknowledged a trial with a placebo arm would be unlikely
48 to be approved.

49 The Committee highlighted that current practice for inhaled therapies in cystic fibrosis follows
50 the Clinical Commissioning Policy by NHS England who propose aztreonam lysine as a third
51 line treatment following tobramycin. However, this recommendation is based primarily on

1 cost, and not cost-effectiveness that assesses if the additional benefit from aztreonam lysine
2 outweigh its additional cost. Moreover, the cost-effectiveness of aztreonam lysine compared
3 to nebulised tobramycin, was published during the development of this guideline by
4 Schechter 2015. Consequently, the comparison between nebulised tobramycin and
5 aztreonam lysine was of greatest interest to the Committee.

6 They agreed that the analysis by Schechter 2015 was favoured towards aztreonam lysine for
7 several reasons. For example, the analysis included a much higher cost to manage an
8 exacerbation (potentially as the analysis took a US third party perspective) and used a high
9 drug acquisition cost for nebulised tobramycin than aztreonam lysine, influencing aztreonam
10 lysine's "dominant" result. Furthermore, participants included in the trial by Assael 2013 were
11 not naïve to tobramycin, potentially with less to scope to benefit from tobramycin compared
12 to aztreonam lysine.

13 However, the clinical evidence review also favoured aztreonam lysine over the other
14 treatments under consideration. Based on the results from the economic model that was
15 developed to reflect UK clinical practice, the Committee agreed that aztreonam lysine could
16 displace nebulised tobramycin as many of the analysis explored provided an ICER below
17 NICE's upper threshold for cost-effectiveness. However, given current NICE HTA
18 recommendations, the Committee agreed they could not recommend aztreonam lysine as
19 the sole second line option following colistimethate sodium. As a result, the Committee
20 prioritised a recommendation to consider aztreonam lysine or tobramycin when the person
21 with a chronic infection is clinically deteriorating, despite regular inhaled colistimethate
22 sodium.

23 The Committee continued to discuss the combination treatment (28 days aztreonam lysine
24 alternating with 28 days nebulised tobramycin) that was dominated by aztreonam lysine in
25 the model. Given that a combination treatment incurs a continuous drug cost and
26 demonstrated clinical effectiveness above nebulised tobramycin and below aztreonam lysine,
27 the result was considered to accurately represent the available evidence. However, the
28 Committee vocalised their concerns that the RCTs identified in the clinical evidence review
29 were flawed as they were often underpowered and included participants who were not
30 treatment naïve. The Committee agreed a research recommendation was needed to
31 demonstrate the clinical and cost effectiveness of combination treatments, but acknowledged
32 that such a study would be impossible to conduct as the number of eligible participants would
33 lead to another underpowered study. Overall, the Committee acknowledged the limitations of
34 the available evidence and agreed not to make a recommendation regarding combination
35 treatments, as they believed clinical practice was clinically effective and cost-effective.

36 **Antimicrobial regimens for the treatment of *A fumigatus*, *S aureus* and *B cepacia*** 37 **complex**

38 The Committee advised that people with cystic fibrosis who have a stable chronic infection
39 with *A fumigatus*, *S aureus* or *B cepacia* complex, and leave hospital untreated, feel anxious.
40 To reduce their anxiety, the Committee agreed that clinicians should inform their patients that
41 there is no evidence on the effectiveness of suppressive antimicrobial treatment, with the aim
42 to reduce the number of people who insist to receive treatment that is potentially cost-
43 ineffective.

44 The Committee advised that people who are chronically infected with *S aureus* are given
45 flucloxacillin in UK clinical practice, noting that there is some variation when the treatment is
46 initiated and stopped with regards to the severity of their symptoms. The Committee
47 acknowledged the higher price of oral solutions compared to capsules (NHS Electronic Drug
48 Tariff September 2016: flucloxacillin 500mg capsules; £2.22/28 capsules, £0.08/500mg vs.
49 flucloxacillin 250mg/5ml oral solution sugar free; £27.24/100ml, £2.72/500mg/10ml) and
50 agreed that people who are chronically infected would be given the cheaper capsule
51 preparation as they are unlikely to have swallowing difficulties at the age of chronic infection.

1 Based on a dose of 4g/day, the cost of flucloxacillin in capsule form would be less than
2 £1/day. Despite such low acquisition costs, the Committee agreed that they should not be
3 prescribed if they do not benefit the person with the chronic infection, especially as cystic
4 fibrosis is a multi-system disorder associated with complex daily regimens. As a result, the
5 Committee concluded that antibiotics to suppress chronic MRSA should only be considered
6 in people who are deteriorating, but not in people who have a stable chronic infection as the
7 cost and burden of treatment is likely to outweigh the benefits of treatment. Conversely,
8 treatment should be considered to suppress chronic MSSA in people with stable pulmonary
9 status given that the expected downstream costs associated with an unmanaged infection
10 would outweigh the cost and burden of suppressive therapy.

11 The Committee advised that the long-term use of drugs used to suppress *B cepacia* complex
12 can have adverse effects associated with additional treatment costs and quality of life
13 decrements. Consequently, the Committee did not want to recommend the use of those
14 drugs in people who are chronically infected and stable, adding that reducing their treatment
15 burden, may subsequently promote adherence to their existing regimens and outweigh the
16 benefits of suppressive treatment. However, the Committee agreed that people who are
17 deteriorating should consider a trial of chronic suppressive treatment with an inhaled
18 antibiotic. The Committee also stated that clinicians should observe their patient's response
19 to treatment to ensure the cost of treatment is justified by suppression of the infection and to
20 discontinue their treatment when they suspect the expected cost to exceed the expected
21 benefit.

22 The Committee advised that the first line treatment for people who are chronically infected
23 with *A fumigatus* is itraconazole. The Committee discussed the clinical evidence that found
24 no significant difference between itraconazole and usual care and concluded that the study
25 was low quality with an undefined population. The Committee also noted that the
26 antimicrobials used to suppress *A fumigatus* are relatively expensive compared to those used
27 to suppress *B cepacia* complex and *S aureus*. As a result, the Committee concluded that
28 treatment in people who are deteriorating should take a stepwise approach, starting with the
29 cheapest treatment first (NHS Electronic Drug Tariff September 2016: itraconazole 100mg
30 capsules; £3.42/15 capsules, £0.23/100mg). Based on a dose of 200mg twice daily, the cost
31 of itraconazole in capsule form would be less than £1/day. Similarly to a stable chronic
32 infection with *B cepacia* complex or *S aureus*, the Committee agreed that people who have a
33 stable chronic infection with *A fumigatus* should not be given routine antimicrobials treatment
34 in an attempt to suppress the chronic infection.

35 Overall, cost data has little use without associated benefits; hence, while the cost of long-
36 term suppressive antimicrobial treatment could be significant, without knowing the benefits of
37 treatment we cannot know if they will be cost-effective. Therefore a research
38 recommendation to assess the clinical effectiveness of suppressive antimicrobial treatment in
39 people chronically infected with *S aureus*, *B cepacia* complex or *A fumigatus* will assess if
40 the benefits can justify the costs, to reduce current uncertainty in this area.

419.4.3.9.4 **Quality of evidence**

42 ***P aeruginosa***

43 The quality of the evidence ranged from very low to high as assessed by GRADE. For the
44 domain risk of bias, the studies were assigned the same risk of bias as in the Cochrane
45 reviews, and were not individually reviewed.

46 The main reasons that lead to downgrading the quality of the evidence was the risk of bias.
47 Many of the studies were open trials. There were also issues in relation to data reporting,
48 randomisation and allocation concealment.

49 Another reason that lead to downgrading the quality of the evidence was imprecision, as
50 confidence intervals crossed 1 or 2 clinical or default MIDs.

1 No issues were identified in relation to the directness of the population.

2 ***S aureus***

3 Not applicable, as no evidence was found for this pathogen.

4 ***B Cepacia Complex***

5 Not applicable, as no evidence was found for this pathogen.

6 ***A fumigatus***

7 One study was found for the treatment of chronic *A fumigatus*. The evidence was considered
8 of low to very low quality as assessed by GRADE, and the main reasons that lead to
9 downgrading the quality of the evidence were the moderate risk of bias found in the study,
10 and the levels of imprecision. The evidence was downgraded further, because the
11 Committee noted therapeutic dosages were not achieved in 2 out of 3 of the participants.

12 **129.4.3.9.5 Other considerations**

13 No equality issues were identified by the Committee for this review question.

14 The Committee agreed a research recommendation should not be prioritised for this topic.
15 There is sufficient evidence to inform the management of chronic *P aeruginosa*. Studies on
16 the management of *S aureus*, *B cepacia* complex, and *A fumigatus* are difficult to conduct.

17 **179.4.3.9.6 Key conclusions**

18 The guideline Committee concluded that the recommendations from the NICE TA report 276
19 (Tappenden 2013) are current, and they agreed colistin should be the first line treatment for
20 the management of chronic *P aeruginosa*. Aztreonam lysine or tobramycin can be given if
21 the person continues to deteriorate, although they noted aztreonam lysine has shown to be
22 more cost-effective. The Committee agreed there is no need to routinely give treatment to
23 people who are chronically infected with *S aureus*, *B cepacia* complex or *A fumigatus*.
24 Treatment should be considered if clinical deterioration is observed, and the response to
25 treatment should be assessed. The Committee also agreed it is important discuss with the
26 person with cystic fibrosis and the families about the benefits and harms of giving treatment.

27 **9.4.4 Recommendations**

28 ***S Aureus***

29 **61. Offer flucloxacillin as antibiotic prophylaxis against respiratory staphylococcus**
30 **aureus infection for children with cystic fibrosis from the point of diagnosis and**
31 **consider continuing up to 6 years of age. Before starting flucloxacillin, discuss the**
32 **uncertainties and possible adverse effects with their parents or carers (as**
33 **appropriate).**

34 **62. For children who are taking antibiotic prophylaxis and have a respiratory sample**
35 **culture that is positive for staphylococcus aureus:**

- 36
- 37 • review prophylaxis adherence and help the child's parents or carers (as
38 appropriate) with any difficulties they are having
 - 39 • start eradication therapy with antibiotics
 - 40 • restart prophylaxis after treatment, even if eradication has not been
successful.

- 1 **63. For people who are not taking prophylaxis and have a new staphylococcus aureus**
2 **infection (that is, previous respiratory sample cultures did not show**
3 **staphylococcus aureus infection):**
- 4 • if they are clinically well, consider an oral anti-staphylococcus aureus
5 agent
 - 6 • if they are clinically unwell and have pulmonary disease, consider oral or
7 intravenous (depending on infection severity) broad-spectrum antibiotics
8 that include an oral anti-staphylococcus aureus agent.
- 9 **64. Consider a long-term antibiotic to suppress chronic methicillin-sensitive**
10 **staphylococcus aureus (MSSA) respiratory infection in people whose pulmonary**
11 **disease is stable.**
- 12 **65. For people with chronic MSSA respiratory infection who become clinically unwell**
13 **with pulmonary disease, consider oral or intravenous broad-spectrum antibiotics**
14 **(depending on infection severity) that include an anti-staphylococcus aureus**
15 **agent.**
- 16 **66. Do not routinely use antibiotics to suppress methicillin-resistant staphylococcus**
17 **aureus (MRSA) in people whose pulmonary disease is stable.**
- 18 **67. If a person with cystic fibrosis and chronic MRSA respiratory infection becomes**
19 **unwell with a pulmonary exacerbation or shows decline in pulmonary function,**
20 **seek specialist microbiological advice.**
- 21 **68. For guidance on preventing the spread of infection, refer to the NICE guideline on**
22 **healthcare-associated infections**
- 23 ***P aeruginosa***
- 24 **69. If a person with cystic fibrosis develops a new pseudomonas aeruginosa infection**
25 **(that is, recent respiratory secretion sample cultures showed no infection):**
- 26 • if they are clinically well:
 - 27 ○ commence eradication therapy with a course of oral or intravenous
28 antibiotics, together with an inhaled antibiotic
 - 29 ○ follow this with an extended course of oral and inhaled antibiotics
- 30 • if they are clinically unwell:
 - 31 ○ commence eradication therapy with a course of intravenous
32 antibiotics together with an inhaled antibiotic
- 33 ○ follow this with an extended course of oral and inhaled antibiotics.
- 34 **70. If eradication treatment is not successful, despite treatment as recommended in**
35 **69, offer sustained treatment with an inhaled antibiotic. Consider nebulised**
36 **colistimethate sodium as the preferred first-line treatment. (See recommendations**
37 **on using colistimethate dry powder for inhalation)**
- 38 **71. Depending on infection severity, use either an oral antibiotic or a combination of 2**
39 **intravenous antibiotics of different classes for people:**
- 40 • who have chronic pseudomonas aeruginosa infection (when treatment
41 has not eradicated the infection) and
 - 42 • who become clinically unwell with a pulmonary disease exacerbation.

1 **72. If a person with chronic pseudomonas aeruginosa infection repeatedly becomes**
2 **clinically unwell with pulmonary disease exacerbations, consider changing the**
3 **antibiotic regimens used to treat exacerbations.**

4 **73. Colistimethate sodium dry powder for inhalation (DPI) is recommended as an**
5 **option for treating chronic pulmonary infection caused by pseudomonas**
6 **aeruginosa in people with cystic fibrosis only if:**

- 7 • they would clinically benefit from continued colistimethate sodium but do
8 not tolerate it in its nebulised form and thus tobramycin therapy would
9 otherwise be considered and
- 10 • the manufacturer provides colistimethate sodium DPI with the discount
11 agreed as part of the patient access scheme to primary, secondary and
12 tertiary care in the NHS.

13 **[This recommendation is from Colistimethate sodium and tobramycin dry**
14 **powders for inhalation for treating pseudomonas lung infection in cystic fibrosis**
15 **(NICE technology appraisal 276)]**

16 **74. For people who are clinically deteriorating despite regular inhaled colistimethate**
17 **sodium, consider nebulised aztreonam, nebulised tobramycin, or tobramycin DPI**
18 **(see recommendation on using tobramycin DPI).**

19 **75. Tobramycin DPI is recommended as an option for treating chronic pulmonary**
20 **infection caused by pseudomonas aeruginosa in people with cystic fibrosis only**
21 **if:**

- 22 • nebulised tobramycin is considered an appropriate treatment, that is,
23 when colistimethate sodium is contraindicated, is not tolerated or has not
24 produced an adequate clinical response and
- 25 • the manufacturer provides tobramycin DPI with the discount agreed as
26 part of the patient access scheme to primary, secondary and tertiary
27 care in the NHS.

28 **[This recommendation is from Colistimethate sodium and tobramycin dry**
29 **powders for inhalation for treating pseudomonas lung infection in cystic fibrosis**
30 **(NICE technology appraisal 276)]**

31 **76. People currently using tobramycin DPI or colistimethate sodium DPI that is not**
32 **recommended according to the recommendations on colistimethate sodium or**
33 **tobramycin DPI should be able to continue treatment until they and their clinician**
34 **consider it appropriate to stop. For children and young people this decision**
35 **should be made jointly by the clinician, the child or young person and their**
36 **parents or carers.**

37 **[This recommendation is from Colistimethate sodium and tobramycin dry**
38 **powders for inhalation for treating pseudomonas lung infection in cystic fibrosis**
39 **(NICE technology appraisal 276)]**

40 ***Burkholderia cepacia* complex**

41 **77. For people with cystic fibrosis who develop a new *Burkholderia cepacia* complex**
42 **infection (that is, recent respiratory sample cultures showed no *Burkholderia***
43 ***cepacia* infection)):**

- 1 • whether they are clinically well or not, give antibiotic eradication therapy
- 2 using a combination of intravenous antibiotics
- 3 • seek specialist microbiological advice on the choice of antibiotics to use.

4 **78. Do not use antibiotics to suppress chronic Burkholderia cepacia complex**
5 **infection in people with cystic fibrosis who have stable pulmonary status.**

6 **79. For people with cystic fibrosis who have chronic Burkholderia cepacia complex**
7 **infection (when treatment has not eradicated the infection) and who become**
8 **clinically unwell with a pulmonary disease exacerbation:**

- 9 • give a combination of oral or intravenous antibiotics, the choice of
- 10 antibiotics being based on specialist microbiological advice.:

11 **80. For people with cystic fibrosis who have chronic Burkholderia cepacia complex**
12 **infection and declining pulmonary status:**

- 13 • consider sustained treatment with an inhaled antibiotic to suppress the
- 14 infection
- 15 • seek specialist microbiological advice on which antibiotic to use
- 16 • stop this treatment if there is no observed benefit.

17 ***H influenza***

18 **81. For people with cystic fibrosis who develop a haemophilus influenza infection**
19 **(diagnosed by a positive respiratory sample culture) but do not have clinical**
20 **evidence of pulmonary infection, treat with an appropriate oral antibiotic.**

21 **82. For people with cystic fibrosis who develop a haemophilus influenza infection**
22 **(diagnosed by a positive respiratory sample culture) and are unwell with clinical**
23 **evidence of pulmonary infection, treat with an appropriate antibiotic, given orally**
24 **or intravenously depending on the severity of the illness**

25 ***Non-tuberculous mycobacteria***

26 **83. For people with cystic fibrosis who are clinically well but whose airway secretions**
27 **are persistently positive for non-tuberculous mycobacteria, discuss with them and**
28 **their family members or carers (as appropriate):**

- 29 • the clinical uncertainties about non-tuberculous mycobacterial infection
- 30 and
- 31 • the possible benefits and risks (for example, drug toxicity) of treating it.

32 **84. If a person with cystic fibrosis has a respiratory sample test positive for new non-**
33 **tuberculous mycobacteria infection, repeat the test for confirmation.**

34 **85. If repeat testing confirms persistent non-tuberculous mycobacteria, do a chest CT**
35 **scan to look for changes consistent with non-tuberculous mycobacteria disease.**

36 **86. Consider non-tuberculous mycobacterial eradication therapy in people with cystic**
37 **fibrosis:**

- 38 • whose airway secretions persistently test positive for non-tuberculous
- 39 mycobacteria and

- 1 • who are clinically unwell with respiratory disease, or who have a chest
- 2 CT scan showing changes consistent with non-tuberculous
- 3 mycobacteria disease and
- 4 • whose pulmonary disease has not responded to other recommended
- 5 treatments.

6 **87. Consider the following eradication regimen for people with cystic fibrosis who are**

7 **having treatment for non-tuberculous mycobacteria.**

- 8 • Initial treatment: a combination of appropriate oral and intravenous
- 9 antibiotics based. Seek specialist microbiological advice on which
- 10 antibiotics to use.
- 11 • Continued treatment: prolonged oral and inhaled antibiotics. Seek
- 12 specialist microbiological advice on which antibiotics to use.

13 ***A fumigatus* complex**

14 **88. Do not routinely use antifungal agents to suppress chronic aspergillus fumigatus**

15 **complex respiratory infection (diagnosed by persistently positive respiratory**

16 **secretion sample cultures) in people with cystic fibrosis and stable pulmonary**

17 **status.**

18 **89. For people with cystic fibrosis with chronic aspergillus fumigatus complex**

19 **respiratory infection and declining pulmonary status:**

- 20 • consider sustained treatment with an antifungal agent to suppress the
- 21 infection
- 22 • seek specialist microbiological advice on which antifungal agent to use
- 23 • stop treatment or change to a different agent if there is no benefit.

24 **90. For people with cystic fibrosis with elevated aspergillus serology and declining**

25 **pulmonary function despite optimised pulmonary treatment, think about treating**

26 **for a possible diagnosis of allergic bronchopulmonary aspergillosis, especially if**

27 **there are consistent chest X-ray or CT scan changes.**

28 **Unidentified Infections**

29 **91. For people with cystic fibrosis who have a pulmonary disease exacerbation and**

30 **no clear cause in their respiratory secretion samples:**

- 31 • use an oral or intravenous (depending on the exacerbation severity)
- 32 broad-spectrum antibiotic
- 33 • continue collecting respiratory secretion samples, and change
- 34 treatments if a pathogen is identified and a more appropriate treatment
- 35 is available.

1 **9.5 Immunomodulatory agents**

2 **Review question: What is the effectiveness of immunomodulatory agents in the**
3 **management of lung disease?**

4 **9.5.1 Introduction**

5 Progressive pulmonary disease is the primary cause of morbidity and mortality in adults with
6 cystic fibrosis (CF). The decrease in lung function associated with chronic infection by a
7 variety of organisms has been related to the severity of pulmonary inflammation.

8 In cystic fibrosis the cystic fibrosis transmembrane conductance regulator (CFTR) is
9 defective, resulting in alterations in the way airway epithelial cells direct the inflammatory
10 response in the airways. Defects in CFTR are associated with increased production of pro-
11 inflammatory mediators, including IL-8, which is a potent neutrophil attractor and so large
12 number of these inflammation-causing cells are directed to the airways resulting in high
13 levels of inflammation and damage. These neutrophils are the primary effector cells
14 responsible for the pathological manifestations of cystic fibrosis lung disease. Additionally,
15 deficiencies in molecules regulating the immune response - such as IL-10, likely contribute to
16 the generation of the excessive and persistent inflammatory response.

17 Therapies which reduce pulmonary inflammation may prove to be clinically efficacious and
18 so reduce the damage caused by persistent infection and improve patient outcomes.

19 **9.5.2 Description of clinical evidence**

20 The aim of this review was to determine the clinical and cost effectiveness of
21 immunomodulatory agents in children and young people and adults with cystic fibrosis.

22 We aimed to look at different immunomodulatory treatments, including inhaled
23 corticosteroids, oral and IV corticosteroids, macrolide antibiotics, NSAIDs and monoclonal
24 antibodies, compared to placebo or other immunomodulatory treatment.

25 Use of Azithromycin in an antimicrobial dose (greater than 250 mg 3 times a week or 500 mg
26 3 times a week for body weight over 40kg) was excluded from this evidence review and
27 considered in the evidence review on antimicrobials for acute pulmonary infections.

28 We searched for systematic reviews of randomised controlled trials (RCTs) and RCTs
29 assessing the effectiveness of immunomodulatory agents in people with cystic fibrosis.

30 For full details see review protocol in Appendix D.

31 Five Cochrane systematic reviews were identified in the search (Balfour-Lynn 2016, Cheng
32 2015, Jat 2013, Lands 2016, Southern 2012).

33 Four reviews were included in this evidence review. Where possible, data and risk of bias
34 assessment was extracted directly from the Cochrane systematic reviews. Individual studies
35 were retrieved for completeness and accuracy, and were also checked for additional
36 outcomes of interest.

- 37 • Balfour-Lynn 2016 evaluated the effectiveness of inhaled corticosteroids compared to
38 placebo or standard treatment. 3 RCTs on Fluticasone use were included from this review
39 (Balfour-Lynn 1997, Balfour-Lynn 2006, De Boeck 2007).
- 40 • Cheng 2015 evaluated the effectiveness of oral corticosteroids compared to placebo or
41 existing conventional therapy. 3 RCTs were included from this review: 2 trials examined
42 use of Prednisone (Eigen 1995 and Auerbach 1985) and 1 trial examined use of
43 Prednisolone (Greally 1994).

- Lands 2016 evaluated the effectiveness of oral NSAIDs compared to placebo or existing therapy. 3 RCTs were included from this review: 2 trials on use of ibuprofen (Konstan 1995 and Lands 2007) and 1 trial on piroxicam (Sordelli 1994).
- Southern 2012 evaluated the effectiveness of macrolide antibiotics compared to placebo, other antibiotic class, or other macrolide. 5 RCTs on Azithromycin were included from this review (Clement 2006, Equi 2002, Saiman 2003, Saiman 2010, Wolter 2002).

An additional Cochrane review (Jat 2015) evaluated the effectiveness of anti-IgE therapy for allergic bronchopulmonary aspergillosis in people with cystic fibrosis. The review reported on 1 clinical trial (Novartis 2008) of Omalizumab which terminated early. This clinical trial (Novartis 2008) was not included as the adverse events reported were not listed in the review protocol.

In addition, 1 observational study (Lai 2000) on the use of Prednisone and 1 RCT on the use of clarithromycin (Robinson 2012) were identified for inclusion in this evidence review.

No evidence was identified which reported on IV methylprednisolone use.

The presentation of evidence synthesis was divided in two parts based on the type of analysis which was used to produce these syntheses:

Two outcomes, the forced expiratory volume in 1 second (FEV₁) % predicted and the rate of pulmonary exacerbations, were considered for network meta-analysis (NMA). These were each split into short (1-10 months) and long-term (>10 months) treatment. Ten studies were included in the NMAs for FEV₁ % predicted and eight studies were included in the NMAs for rate of pulmonary exacerbations (see Section 1.1.2.1) Pairwise comparisons were performed for the rest of the outcomes included in the review protocol. Ten RCTs were included from the Cochrane systematic reviews for the outcomes of nutritional status, time to next pulmonary exacerbation, adverse effects and quality of life.

- If a study did not report standardised height or weight scores (such as BMI, z scores or standard deviation scores), the absolute value (in centimetres or kilograms) was included. Standard deviation scores reported were compared to a population of children without cystic fibrosis.
- The size of the studies ranged between 23 and 285 people with cystic fibrosis. Two studies included children (Auerbach 1985, De Boeck 2007), 8 studies included children and young people (Balfour-Lynn 1997, Clement 2006, Eigen 1995, Equi 2002, Grealley 1994, Lai 2000, Lands 2007, Saiman 2010), 1 study included young people and adults (Wolter 2002), 5 studies included children, young people and adults (Balfour-Lynn 2006, Konstan 1995, Robinson 2012, Saiman 2003, Sordelli 1994).
- Three studies were conducted in the UK (Balfour-Lynn 1997, Balfour-Lynn 2006, Equi 2002), 6 in the USA (Auerbach 1985, Eigen 1995, Konstan 1995, Lai 2000, Saiman 2003, Saiman 2010), 1 in Belgium (De Boeck 2007), 1 in France (Clement 2006), 1 in Ireland (Grealley 1994), 1 in Canada (Lands 2007), 1 in Argentina (Sordelli 1994), 1 in Australia (Wolter 2002), 1 in the USA and Australia (Robinson 2012).

A summary of the studies included in this review is presented in Table 159. See study selection flow chart in Appendix F, study evidence tables in Appendix G, list of excluded studies in Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

9.5.3 Summary of included studies

A summary of the included studies is presented in Table 126.

Table 126: Summary of included studies (NMA and non-NMA outcomes)

Study	Intervention/ Comparison	Population	Outcomes	Comments
-------	--------------------------	------------	----------	----------

Cochrane systematic reviews				
Balfour-Lynn 2016 Cochrane SR	Intervention: Any inhaled corticosteroid, using any inhalation device, for a period of ≥ 2 weeks Comparison: Placebo or standard treatment	People of any age, with confirmed diagnosis of cystic fibrosis, regardless of clinical severity.	<ul style="list-style-type: none"> • FEV₁ % predicted • time to next exacerbation • change in height (cm) • Change in height 	AMSTAR score: 11/11 • Studies included from this review: Balfour-Lynn 1997, Balfour-Lynn 2006, De Boeck 2007
Cheng 2015 Cochrane SR	Intervention: Oral corticosteroids Comparison: Placebo or existing conventional therapy	People with defined cystic fibrosis, of any age, at all stages of lung disease.	<ul style="list-style-type: none"> • Mortality, at 6 months • Adverse effects, • Absolute change in weight (kg) 	AMSTAR score: 10/11 Studies included from this review: Auerbach 1985, Eigen 1995, Greally 1994.
Lands 2016 Cochrane SR	Intervention: Oral NSAIDs for a minimum period of 2 months Comparison: Placebo or existing therapy	People with defined cystic fibrosis, of any age, and any stage of lung disease.	<ul style="list-style-type: none"> • Adverse effects • Annual rate of change in % ideal body weight overall and by age 	AMSTAR score: 10/11 Studies included from this review: Konstan 1995, Lands 2007, Sordelli 1994.
Southern 2012 Cochrane SR	Intervention: Short or long term use of a macrolide antibiotic Comparison: Placebo, other antibiotic class, other macrolide	People with confirmed diagnosis of cystic fibrosis	<ul style="list-style-type: none"> • Time to next exacerbation • Change in weight • Exacerbation* • Adverse effects • Quality of life using CFQ-R 	AMSTAR score: 11/11 Studies included from this review: Clement 2006, Equi 2002, Saiman 2003, Saiman 2010, Wolter 2002.
Primary studies included in the Cochrane SR				
Auerbach 1985 (USA) RCT	Intervention: Prednisone 2 mg/kg to a maximum of 60 mg Comparison: Placebo	N=45 children with cystic fibrosis Age range: 1 to 12 years Participants with mild to moderate pulmonary disease	<ul style="list-style-type: none"> • Mortality, at 6 months 	Included in Cochrane SR Cheng 201 Included in pairwise comparisons only.
Balfour-Lynn 1997 (UK) Cross-over RCT	Intervention: Fluticasone (dry powder) 200 micrograms twice per day Comparison: Placebo	N=23 Mean age 10.3 years (range 7 to 17 years)	<ul style="list-style-type: none"> • Short-term FEV₁ % predicted (6 weeks) 	Included in Cochrane SR Balfour-Lynn 2016 • Included in NMA only

Balfour-Lynn 2006 (UK) Cross-over RCT	<p>Intervention: Fluticasone propionate via volumatic spacer given at equivalent dose taken (55% had 400 g/d fluticasone and 45% took lower dose); before trial entry</p> <p>Comparison: Placebo via volumatic spacer. If previously given budesonide beclometasone, switched to fluticasone to 2:1 ratio.</p>	<p>N=171 people with cystic fibrosis Age (mean, range): 14 (6 to 53) years in fluticasone group; 15.8 years in placebo</p>	<ul style="list-style-type: none"> • Time to next exacerbation • Change in height (cm) 	<p>Included in Cochrane SR Balfour-Lynn 2016 Included in pairwise comparisons only.</p>
De Boeck 2007 (Belgium) RCT	<p>Intervention: Fluticasone 500 mcg dry powder inhaler twice daily</p> <p>Comparison: Lactose placebo dispensed in identical canister</p>	<p>N=29 children with cystic fibrosis Mean age (SD): 8.2 (0.6) years intervention group; 9.0 (0.5) in the control group</p>	<ul style="list-style-type: none"> • Change in height 	<p>Included in Cochrane SR Balfour-Lynn 2016 Included in NMA and pairwise comparisons.</p>
Clement 2006 (France) RCT	<p>Intervention: Azithromycin 250 mg tablets 3 times a week (>40 kg, 500 mg)</p> <p>Comparison: Placebo</p>	<p>N=82 children and young people with cystic fibrosis Age (mean, SD): 11 (3.3) years</p>	<ul style="list-style-type: none"> • Time to next exacerbation • Change in BMI, z score at 12 months follow-up 	<p>Included in Cochrane SR Southern 2012 Included in NMA and pairwise comparisons.</p>
Eigen 1995 (USA) RCT	<p>Intervention: Prednisone 2 mg/kg or 1 mg/kg on alternate days to a maximum of 60 mg.</p> <p>Comparison: Placebo</p>	<p>N=285 children and young people with cystic fibrosis Age range: 6 to 14 years</p>	<ul style="list-style-type: none"> • Adverse effects, up to 3 and 4 years follow-up 	<p>Included in Cochrane SR Cheng 2015 Included in NMA and pairwise comparisons.</p>
Equi 2002 (UK) RCT	<p>Intervention: Azithromycin 250 mg tablets (>40 kg, 500 mg) per day</p> <p>Comparison: Placebo</p>	<p>N=41 Aged: 8 to 18 years</p>	<ul style="list-style-type: none"> • Short-term FEV₁ % predicted (26 weeks) • Short-term exacerbations per patient (26 weeks) 	<p>Included in Cochrane SR Southern 2012 • Included in NMA only</p>
Greally 1994 (Ireland) RCT	<p>Intervention: Soluble prednisolone 2 mg/kg daily for 14 days then 1</p>	<p>N=24 children and young people with cystic fibrosis</p>	<ul style="list-style-type: none"> • Absolute change in weight (kg) 	<p>Included in Cochrane SR Cheng 2015 Included in</p>

	mg/kg/day on alternate days for 10 weeks (maximum dose 40 mg)	Age range: 5.5 to 19.5 years		pairwise comparisons only.
	Comparison: Placebo			
Konstan 1995 (USA) RCT	Intervention: Ibuprofen 20 – 30 mg/kg to a maximum of 1600 mg, determined by pharmacokinetic study	N=85 people with cystic fibrosis Age range: 5 to 39	<ul style="list-style-type: none"> • Adverse effects up to 4 years follow-up • Annual rate of change in % ideal body weight overall and by age 	Included in Cochrane SR Lands 2016 Included in NMA and pairwise comparisons.
	Comparison: Placebo			
Lands 2007 (Canada) RCT	After pharmacokinetic study: Intervention: Ibuprofen 200 mg tablets at a dose of 20 to 30 mg/kg to a maximum of 1600 mg	N=142 children and young people with cystic fibrosis Age: 6 to 18 years	<ul style="list-style-type: none"> • Adverse effects, up to 2 years follow-up 	Included in Cochrane SR Lands 2016 Included in NMA and pairwise comparisons.
	Comparison: Placebo			
Saiman 2003 (USA) RCT	Intervention: Azithromycin 250 mg tablets 3 days a week (>40 kg, 500 mg)	N=185 people with cystic fibrosis Age: > 6 years	<ul style="list-style-type: none"> • Time to next exacerbation • Adverse effects at 6 months follow-up • Change in weight at 6 months follow-up, and quality of life using CFQ-R 	Included in Cochrane SR Southern 2012 Included in NMA and pairwise comparisons.
	Comparison: Placebo			
Saiman 2010 (USA) RCT	Intervention: Azithromycin 250 mg tablets 3 times a week (>36 kg, 500 mg)	N=263 children and young people with cystic fibrosis Age range: 6 to 18 years	<ul style="list-style-type: none"> • Time to next exacerbation* • Change in weight at 6 months follow-up 	Included in Cochrane SR Southern 2012 Included in NMA and pairwise comparisons.
	Comparison: Placebo			
Sordelli 1994 (Argentina) RCT	Intervention: Piroxicam (16-25kg: 10mg/d; 26-45kg: 15mg/d; >46kg: 20mg/d)	N = 41 children, young people and adults with cystic fibrosis Age range: 5 to 37 years	Long-term exacerbations per patient (12-19 months)	Included in Cochrane SR Lands 2016 Included in NMA only
	Comparison: Placebo			
Wolter 2002 (Australia)	Intervention: Azithromycin 250mg	N = 60 young people and adults	Short-term FEV ₁ % predicted (12	Included in Cochrane SR

RCT	daily Comparison: Placebo	with cystic fibrosis Mean age (SD): 27.9 (6.5) years	weeks) Short-term exacerbations per patient (12 weeks)	Southern 2012 Included in NMA only
Additional primary studies				
Lai 2000 Retrospective cohort study (10 years follow up of a Double-blind multicentre RCT - Eigen 1995)	Group 1: placebo Group 2: 1 mg Prednisone/kg Group 3: 2 mg Prednisone/kg	N = 224 Age range: 6 to 14 years	• Height and weight at 18 years of age	Included in pairwise comparisons only.
Robinson 2012 (USA and Australia) Cross-over RCT	Clarithromycin 500mg per day or placebo	N=63 children, young people and adults with cystic fibrosis Mean age (SD): 16 (10.5) years	• Short-term FEV ₁ % predicted (22 weeks) • Short-term exacerbations per patient (22 weeks)	• Included in NMA only

1 CFQ-R: cystic fibrosis respiratory questionnaire; NSAIDS: non-steroidal anti-inflammatory drugs

2 9.5.4 Clinical evidence profile


3 9.5.4.1 Clinical evidence profile for NMA outcomes (FEV₁ % predicted and rate of pulmonary 4 exacerbations)

5 As treatment effects were found to vary over time, NMAs were conducted separately for
6 short (1-10 months) and long (>10 months) of treatment.

7 9.5.4.1.1 FEV₁ % Predicted – Short-term (1-10 month) treatment

8 Six studies of 735 participants were included in the network of three classes of interventions
9 – placebo, macrolide antibiotics (azithromycin, clarithromycin), inhaled corticosteroids
10 (fluticasone) (Figure 8). The evidence was of moderate quality. Four studies were at low risk
11 of bias and for the two other studies the risk of bias was unclear.

Figure 8: Network for FEV₁ % predicted short-term treatment



Note: Size of nodes are proportional to the number of patients in the network treated with a particular intervention. Thickness of connecting lines are proportional to the number of studies comparing two interventions.

1 Table 127 presents the results of the conventional pair-wise meta-analyses (head to head
2 comparisons) (upper-right section of table), together with the results computed by the NMA
3 for every possible treatment comparison (lower-left section of table). Both results are
4 presented as mean differences (95% CrI). These results were derived from a fixed effects
5 mode I(Appendix N).

6 Macrolide antibiotics were found to have a clinically significant improvement versus placebo
7 for limiting the decline in FEV₁ % predicted for people with cystic fibrosis treated for 1-10
8 months (Figure 9). Inhaled corticosteroids were not found to have a clinically significant effect
9 on FEV₁ % predicted compared with placebo. Macrolide antibiotics were also not found to
10 have a clinically significant effect over inhaled corticosteroids for short-term treatment.
11 Incoherence could not be assessed as there were no closed loops of treatments.


12 In this analysis, macrolide antibiotics were found to have the highest probability (91.21%) of
13 being the best treatment to improve/limit the decline in FEV₁ % predicted among
14 interventions with a duration of 1-10 months (Table 128).

15 **Table 127: Mean differences (95% CrI) from conventional (white area) and network
16 meta-analysis (grey area) for FEV₁ % predicted with short-term treatment**

	Placebo	Macrolide antibiotics	Inhaled corticosteroids
Placebo		2.16 (0.52, 3.79)	-2.00 (-7.65, 3.67)
Macrolide antibiotics	2.16 (0.52, 3.79)		
Inhaled corticosteroids	-2.00 (-7.65, 3.67)	-4.16 (-10.04, 1.75)	

17 *Results in the top right diagonal of the table are the mean differences and 95% CrI from the conventional meta-*
18 *analyses of direct evidence between the column-defined treatments compared to the row-defined treatment.*
19 *Mean differences greater than 0 favour the column-defined treatment.*
20 *Results in the bottom left are the mean differences and 95% CrI from the NMA model of direct and indirect*
21 *evidence between the row-defined treatments compared to the column-defined treatments. Mean differences*
22 *greater than 0 favour the row-defined treatment.*
23 *Numbers in bold denote results for which the 95% CrI does not include the null effect of 0*

1 **Figure 9: Forest plot showing mean differences (with their 95% CrI) of NMA estimates**
 2 **for each intervention versus placebo for FEV₁ % predicted with short-term**
 3 **treatment.**



4
5 *Note: Vertical dashed line shows the line of no effect*

6 **Table 128: Median treatment ranking (with their 95% CrI) of all interventions in the**
 7 **network and the probability of being the best treatment for improving/limiting the**
 8 **decline of FEV₁ % predicted in the short-term**

	Median (95% CrI) treatment rank	Probability of being the best treatment (%)
Placebo	2 (2-3)	0.37%
Macrolide antibiotics	1 (1-2)	91.21%
Inhaled corticosteroid	3 (1-3)	8.41%

9 **Table 129: Quality assessment of the evidence for the NMA for FEV₁ % predicted in**
 10 **the short-term**


NMA	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Quality
Short term (1-10 months) FEV % predicted (6 studies)	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious ¹	None	Moderate

11 ¹ – No intervention has rank credible intervals ≤33% of total distribution of comparators

12 **129.5.4.1.2 FEV₁ % Predicted - Long-term (>10 month) treatment**

13 Five studies of 511 participants were included in the network of four classes of interventions
 14 – placebo, NSAIDs (ibuprofen), macrolide antibiotics (azithromycin), inhaled corticosteroids
 15 (fluticasone), and oral corticosteroids (prednisolone) (Figure 10). The evidence was of low
 16 quality. Two studies were at low risk of bias and for the other three studies the risk of bias
 17 was unclear.

Figure 10: Network for FEV₁ % predicted long-term treatment



Note: Size of nodes are proportional to the number of patients in the network treated with a particular intervention. Thickness of connecting lines are proportional to the number of studies comparing two interventions.

1 Table 130 presents the results of the conventional pair-wise meta-analyses (head to head
2 comparisons) (upper-right section of table), together with the results computed by the NMA
3 for every possible treatment comparison (lower-left section of table). Both results are
4 presented as mean differences (95% CrI). These results were derived from a fixed effects
5 model (Appendix N).

6 NSAIDs and oral corticosteroids were found to have a clinically significant improvement
7 versus placebo for limiting the decline in FEV₁ % predicted for people with cystic fibrosis for
8 >10 months of treatment (Figure 11). Long-term macrolide antibiotic treatment was not
9 found to have a clinically significant effect on FEV₁ % predicted compared with placebo.
10 There may be a clinically significant improvement of long-term oral corticosteroid treatment
11 versus long-term macrolide antibiotic treatment, though there was insufficient evidence to
12 confirm this. Incoherence could not be assessed as there were no closed loops of
13 treatments.

14 In this analysis, long-term NSAID treatment was found to have the highest probability
15 (65.2%) of being the best treatment to improve/limit the decline in FEV₁ %, followed by long-
16 term oral corticosteroid treatment (12.9%) (Table 131).

17 **Table 130: Mean differences (95% CrI) from conventional (white area) and network
18 meta-analysis (grey area) for FEV₁ % predicted with long-term treatment**

	Placebo	NSAIDs	Macrolide antibiotics	Inhaled corticosteroids	Oral corticosteroids
Placebo		2.26 (1.40, 3.11)	-2.8 (-10.02, 4.44)	7.17 (-3.64, 17.95)	4.07 (0.7, 7.49)
NSAIDs	2.26 (1.41, 3.11)				


	Placebo	NSAIDs	Macrolide antibiotics	Inhaled corticosteroids	Oral corticosteroids
Macrolide antibiotics	-2.81 (-10.05, 4.45)	-5.07 (-12.34, 2.24)			
Inhaled corticosteroids	7.17 (-3.64, 17.95)	4.9 (-5.94, 15.73)	9.96 (-2.98, 22.95)		
Oral corticosteroids	4.07 (0.7, 7.49)	1.81 (-1.67, 5.33)	6.88 (-1.11, 14.92)	-3.09 (-14.39, 8.24)	

Results in the top right diagonal of the table are the mean differences and 95% CrI from the conventional meta-analyses of direct evidence between the column-defined treatments compared to the row-defined treatment. Mean differences greater than 0 favour the column-defined treatment.

Results in the bottom left are the mean differences and 95% CrI from the NMA model of direct and indirect evidence between the row-defined treatments compared to the column-defined treatments. Mean differences greater than 0 favour the row-defined treatment.

Numbers in bold denote results for which the 95% CrI does not include the null effect of 0

Figure 11: Forest plot showing mean differences (with their 95% CrI) of NMA estimates for each intervention versus placebo for FEV₁ % predicted with long-term treatment.



Note: Vertical dashed line shows the line of no effect

Table 131: Median treatment ranking (with their 95% CrI) of all interventions in the network and the probability of being the best treatment for improving/limiting the decline of FEV₁ % predicted in the long-term

	Median (95% CrI) treatment rank	Probability of being the best treatment (%)
Placebo	4 (3-5)	8.12%
NSAIDs	3 (1-4)	65.20%
Macrolide antibiotics	5 (2-5)	5.99%
Inhaled corticosteroids	1 (1-5)	7.82%
Oral corticosteroids	2 (1-3)	12.88%

1 **Table 132: Quality assessment of the evidence for the NMA for FEV₁ % predicted in the**
2 **long-term**


NMA	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Quality
Long term (>10 months) FEV % predicted (3 studies)	Serious ¹	No serious inconsistency	No serious indirectness	Serious ²	None	Low

3 1 – Two included studies were at low risk of bias and for the other three studies the risk of bias was unclear.
4 2 – No intervention has rank credible intervals ≤33% of total distribution of comparators

59.5.4.1.3 **Rate of pulmonary exacerbations - Short-term (1-10 month) treatment**

6 Three studies of 226 participants were included in the network of two classes of interventions
7 – placebo and macrolide antibiotics (azithromycin, clarithromycin) (Figure 12). The evidence
8 for this analysis was of moderate quality. One study was at low risk of bias and for the other
9 two studies the risk of bias was unclear.

Figure 12: Network for rate of pulmonary exacerbations short-term treatment



Note: Size of nodes are proportional to the number of patients in the network treated with a particular intervention. Thickness of connecting lines are proportional to the number of studies comparing two interventions.

10 As there was only evidence on two classes of short-term treatments to reduce the rate of
11 pulmonary exacerbations no NMA was performed for this outcome.

12 Very low quality evidence showed no clinically significant difference in the rate of
13 exacerbations after short-term treatment between macrolide antibiotics and placebo (see
14 Appendix N). This result was derived from a random effects pairwise analysis.

15 **Table 133: Quality assessment of the evidence for the NMA for rate of**
16 **exacerbations in the short-term**

Fluticasone compared to placebo for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect	No of Participants	Quality of the evidence	Comments
	Assumed	Corresponding				


Fluticasone compared to placebo for cystic fibrosis						
	rate	rate	(95% CI)	(studies)	(GRADE)	
	Placebo	Macrolide antibiotics				
Rate of exacerbations after short-term (10 month) treatment	The mean rate of exacerbations per patient in the control groups was 0.56	The mean rate of exacerbations in the intervention groups was 0.42 (0.21 to 0.85)	Rate ratio 0.75 (0.37 to 1.51)	226 (Equi 2002, Robnson 2012, Wolter 2002)	⊕⊖⊕⊖ very low ^{1,2}	
*The basis for the assumed rate (e.g. the median control group rate across studies) is provided in footnotes. The corresponding rate (and its 95% confidence interval) is based on the assumed rate in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval;						

1 Evidence was downgraded by 2 due to very serious inconsistency between studies
2 Evidence was downgraded by 2 due to very serious imprecision as 95%CI crossed two default MIDs

39.5.4.1.4 Rate of pulmonary exacerbations - Long-term (>10 month) treatment

4 Three studies of 321 participants were included in the network of four classes of
5 interventions – placebo, macrolide antibiotics (azithromycin), inhaled corticosteroids
6 (fluticasone), and oral corticosteroids (prednisolone) (Figure 13). The evidence was of low
7 quality. One study was at low risk of bias and for the other two the risk of bias was unclear.

8 **Figure 13: Network for rate of pulmonary exacerbations long-term treatment**



9
10 Note: Size of nodes are proportional to the number of patients in the network treated with a particular
11 intervention. Thickness of connecting lines are proportional to the number of studies comparing two interventions.

12 Table 134 presents the results of the conventional pair-wise meta-analyses (head to head
13 comparisons) (upper-right section of table), together with the results computed by the NMA
14 for every possible treatment comparison (lower-left section of table). Both results are

1 presented as mean differences (95% CrI). These results were derived from a fixed effects
2 model (Appendix N).

3 There was considerable uncertainty throughout the network. No clinically significant
4 differences were found between any of the treatments in the network. Incoherence could not
5 be assessed as there were no closed loops of treatments.

6 In this analysis, long-term macrolide antibiotic treatment was found to have the highest
7 probability (56.8%) of being the best treatment to reduce the rate of exacerbations, followed
8 by long-term oral corticosteroid treatment (25.5%) (Table 135).

9 One study (Sordelli 1994) that provided information on NSAID efficacy was a candidate for
10 inclusion into the network. However, as the NSAID used in the study (piroxicam) was
11 considered to have potentially severe side effects and as the study was at high risk of bias
12 (trial was unblinded and neither randomisation nor allocation methods were sufficiently
13 described) it was not included in the final network.

14 As this was the only study providing information on NSAIDs, the results of the network were
15 highly sensitive to it. Inclusion of this study did not affect estimates for other classes in the
16 network, but provided an estimate for NSAID efficacy. This quality of evidence for this new
17 network worsened from low to very low quality, with NSAIDs having the highest probability
18 (61.2%) of being the best treatment, followed by macrolide antibiotics (24.7%). Further
19 results of this sensitivity analysis are reported in Appendix N.

20 **Table 134: Rate ratios (95% CrI) from conventional (white area) and network meta-**
21 **analysis (grey area) for the rate of exacerbations with long-term treatment**


	Placebo	Macrolide antibiotics	Inhaled corticosteroids	Oral corticosteroids
Placebo		0.44 (0.03, 3.29)	1.34 (0.22, 9.37)	0.91 (0.08, 10.1)
Macrolide antibiotics	0.44 (0.03, 3.29)			
Inhaled corticosteroids	1.34 (0.22, 9.37)	3.14 (0.2, 72.84)		
Oral corticosteroids	0.91 (0.08, 10.1)	2.13 (0.09, 66.33)	0.67 (0.03, 13.54)	

22 *Results in the top right diagonal of the table are the rate ratios and 95% CrI from the conventional meta-analyses*
23 *of direct evidence between the column-defined treatments compared to the row-defined treatment. Rate*
24 *ratios greater than 1 favour the column-defined treatment.*

25 *Results in the bottom left are the rate ratios and 95% CrI from the NMA model of direct and indirect evidence*
26 *between the row-defined treatments compared to the column-defined treatments. Rate ratios greater than 1*
27 *favour the row-defined treatment.*

28 *Numbers in bold denote results for which the 95% CrI does not include the null effect of 1*

1 **Figure 14: Forest plot showing rate ratio (with their 95% CrI) of NMA estimates for**
 2 **each intervention versus placebo for the rate of exacerbations with long-term**
 3 **treatment.**



4
5 *Note: Vertical dashed line shows the line of no effect*

6 **Table 135: Median treatment ranking (with their 95% CrI) of all interventions in the**
 7 **network and the probability of being the best treatment for reducing the rate of**
 8 **exacerbations in the long-term**

	Median (95% CrI) treatment rank	Probability of being the best treatment (%)
Placebo	3 (1-4)	6.37%
Macrolide antibiotics	1 (1-4)	56.82%
Inhaled corticosteroids	3 (1-4)	11.30%
Oral corticosteroids	2 (1-4)	25.51%

9
10 **Table 136: Quality assessment of the evidence for the NMA for rate of**
 11 **exacerbations in the long-term**

NMA	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Quality
Long term (>10 months) rate of exacerbations (4 studies)	Serious ¹	No serious inconsistency	No serious indirectness	Serious ²	None	Low

12 (x) 1 – One included study was at low risk of bias, one study was at high risk of bias, and for the other two the
 13 risk of bias was unclear
 14 (y) 2 – No intervention has rank credible intervals $\leq 33\%$ of total distribution of comparators

15 **9.5.4.2 Clinical evidence profile for non-NMA outcomes (nutritional status, time to next**
 16 **pulmonary exacerbation, adverse events and quality of life)**

17 The summary clinical evidence profile tables are presented in Table 137 - Table 140.

1
2

Table 137: Summary clinical evidence profile: Comparison 1. Fluticasone versus placebo

Comparison 1. Fluticasone versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Fluticasone				
Time to first exacerbation Follow-up: 6 months	460 per 10001	483 per 1000 (342 to 645) ¹	HR 1.07 (0.68 to 1.68) ²	171 (Balfour-Lynn 2006)	⊕⊕⊕⊖ low ¹	
Growth (change in height) SDS (standard deviation) score Follow-up: 12 months	The mean growth (change in height) in the placebo groups was -0.01 SDS	The mean growth (change in height) in the fluticasone groups was 0.37 SDS lower (0.77 lower to 0.03 higher)		30 (De Boeck 2007)	⊕⊕⊕⊖ moderate ³	
Growth (change in height) in paediatric participants cm Follow-up: 8 months	The mean growth (change in height) in paediatric participants in the placebo groups was 3.5 cm	The mean growth (change in height) in paediatric participants in the fluticasone groups was 0.6 cm higher (0.46 lower to 1.66 higher)		80 (Balfour-Lynn 2006)	⊕⊕⊕⊖ moderate ³	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; HR: hazard ratio; MD: mean difference; SDS: standard deviation score

3
4
5

1 The quality of the evidence was downgraded by 2 as 95%CI crossed the null effect line, and it is very wide.
2 Calculated by the NGA technical team from percentage of participants in group with at least 1 exacerbation.
3 The quality of the evidence was downgraded by 1 because 95%CI crossed 1 default MID.

6
7

Table 138: Summary clinical evidence profile: Comparison 2. Prednisone/ Prednisolone versus placebo

Comparison 2. Prednisone/ Prednisolone versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Prednisone/ Prednisolone				
[2 mg prednisone] Absolute change in	The mean absolute change in weight in the placebo groups	The mean absolute change in weight in the prednisone groups was		25 (Greally 1994)	⊕⊖⊖⊖ very low ^{1,2}	

weight (kg) Follow-up: 12 weeks	was 0.01 kg	0.34 kg higher (2.32 lower to 3 higher)			
[1 mg prednisone] Weight (Kg) Boys at 18 Years of Age	The mean absolute weight in the placebo groups was 63.7 kg	The mean weight in the prednisone groups was 4.6 kg lower (9.69 lower to 0.49 higher)		55 (Lai 2000)	⊕⊕⊕⊕ low ³
[2 mg prednisone] Weight (Kg) Boys at 18 Years of Age	The mean absolute weight in the placebo groups was 63.7 kg	The mean weight in the prednisone groups was 6.7 kg lower (11.59 to 1.81 lower)		52 (Lai 2000)	⊕⊕⊕⊕ moderate ⁴
[1 mg prednisone] Weight (Kg) Girls at 18 Years of Age	The mean absolute weight in the placebo groups was 51.9 kg	The mean weight in the prednisone groups was 0 kg higher (7.62 lower to 3.02 higher)		43 (Lai 2000)	⊕⊕⊕⊕ very low ²
[2 mg prednisone] Weight (Kg) Girls at 18 Years of Age	The mean absolute weight in the placebo groups was 51.9 kg	The mean weight in the prednisone groups was 1.7 kg higher (3.37 lower to 6.77 higher)		46 (Lai 2000)	⊕⊕⊕⊕ very low ²
[1 mg prednisone] Height (cm) Boys at 18 Years of Age -	The mean absolute height in the placebo groups was 174.6 cm	The mean height in the prednisone groups was 3.9 cm lower (7.77 to 0.03 lower)		55 (Lai 2000)	⊕⊕⊕⊕ very low ³
[2 mg prednisone] Height (cm) Boys at 18 Years of Age	The mean absolute height in the placebo groups was 174.6 cm	The mean height in the prednisone groups was 4.1 cm lower (7.82 to 0.38 lower)		52 (Lai 2000)	⊕⊕⊕⊕ very low ³
[1 mg prednisone] Height at 18 Years of Age - Girls cm	The mean absolute height in the placebo groups was 160.3 cm	The mean height in the prednisone groups was 1 cm lower (4.54 lower to 2.54 higher)		43 (Lai 2000)	⊕⊕⊕⊕ very low ²
[2 mg prednisone] Height (cm) Girls at 18	The mean absolute height in the placebo groups was	The mean height in the prednisone groups was		46 (Lai 2000)	⊕⊕⊕⊕ very low ²

Years of Age	160.3 cm	0.5 cm lower (4.43 lower to 3.43 higher)				
[1 mg prednisone] Adverse effects - Cataracts Follow-up: 4 years	74 per 1000	32 per 1000 (8 to 119)	RR 0.43 (0.11 to 1.61)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[2mg prednisone] Adverse effects - Cataracts Follow-up: 3 years	74 per 1000	116 per 1000 (47 to 286)	RR 1.57 (0.64 to 3.88)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[1 mg prednisone] Adverse effects - Diabetes mellitus Follow-up: 4 years	11 per 1000	32 per 1000 (3 to 298)	RR 3 (0.32 to 28.33)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[2 mg prednisone] Adverse effects - Diabetes mellitus Follow-up: 3 years	11 per 1000	63 per 1000 (8 to 515)	RR 6.00 (0.74 to 48.89)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[1 mg prednisone] Adverse effects – Glycosuria Follow-up: 4 years	42 per 1000	63 per 1000 (19 to 217)	RR 1.5 (0.44 to 5.15)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[2 mg prednisone] Adverse events – Glycosuria Follow-up: 3 years	42 per 1000	105 per 1000 (34 to 324)	RR 2.5 (0.81 to 7.69)	190 (Eigen 1995)	⊕⊕⊕⊕ low ^{1,3}	
[1 mg prednisone] Adverse effects - Hyperglycaemia Follow-up: 4 years	21 per 1000	32 per 1000 (5 to 185)	RR 1.5 (0.26 to 8.78)	190 (Eigen 1995)	⊕⊕⊕⊕ very low ^{1,2}	
[2 mg prednisone]	21 per 1000	105 per 1000 (24 to 468)	RR 5 (1.13 to 22.8)	190 (Eigen 1995)	⊕⊕⊕⊕ low ^{1,3}	

Adverse effects - Hyperglycaemia			to 22.21)	1995)		
Follow-up: 3 years						
Mortality	42 per 1000	16 per 1000 (1 to 368)	RR 0.38 (0.02 to 8.83)	45 (Auberch 1985)	⊕⊕⊕⊖ low ^{5,6}	
Follow-up: 4 years						

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; kg: kilogrammes; MD: mean difference; mg: milligrams; RR: risk ratio

- 1 The quality of the evidence was downgraded by 1, as allocation concealment and blinding were unclear.
- 2 The quality of the evidence Evidence downgraded by 2 due to serious imprecision as 95% CI crossed 2 default MIDs.
- 3 The quality of the evidence Evidence downgraded by 1 due to serious imprecision as 95% CI crossed 1 default MID.
- 4 The quality of the evidence was upgraded by 1 as there is evidence of dose-response within study
- 5 Allocation concealment and blinding were unclear, but the quality of the evidence was not downgraded for this outcome
- 6 The quality of the evidence was downgraded by 2 as 95%CI crossed the null effect line, and it is very wide.

Table 139: Summary clinical evidence profile: Comparison 3. Azithromycin versus placebo

Comparison 3. Azithromycin versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Azithromycin				
Time to next exacerbation: time free of exacerbation Follow-up: mean 6 months	Study population		HR 0.59 (0.44 to 0.79)	445 (Saiman 2003, Saiman 2010)	⊕⊕⊕⊕ high	
	348 per 1000	223 per 1000 (172 to 287) ¹				
	Moderate					
	348 per 1000	223 per 1000 (172 to 287) ¹				
Time to next exacerbation Follow-up: 12 months	Study population		HR 0.37 (0.217 to 0.6299)	82 (Clement 2006)	⊕⊕⊕⊕ high	
	48 per 1000 ¹	18 per 1000 (11 to 30) ¹				
	Moderate					
	36 per 1000 ¹	13 per 1000 (8 to 23) ¹				
Mild adverse effects of antibiotic treatment - Hearing impairment Follow-up: 6 months	10 per 1000	12 per 1000 (1 to 181)	RR 1.13 (0.07 to 17.74)	185 (Saiman 2003)	⊕⊕⊖⊖ low ²	

Comparison 3. Azithromycin versus placebo						
Mild adverse effects of antibiotic treatment – Tinnitus Follow-up: 6 months	10 per 1000	12 per 1000 (1 to 181)	RR 1.13 (0.07 to 17.74)	185 (Saiman 2003)	⊕⊕⊕⊕ low ²	
Change in BMI z score Follow-up: 12 months	The mean change in BMI z score in the placebo groups was -0.12	The mean change in BMI z score in the azithromycin groups was 0.15 higher (0.03 lower to 0.33 higher)		82 (Clement 2006)	⊕⊕⊕⊕ moderate ³	
Change in weight (kg) Follow-up: 6 months	Not reported	The mean change in weight in the azithromycin groups was 0.62 higher (0.26 to 0.98 higher)		440 (Saiman 2003, Saiman 2010)	⊕⊕⊕⊕ moderate ³	
Quality of life: change in CFQ-R total score Scale from: 0 to 100 Follow-up: 6 months	The mean change in total quality of life score (CFQ-R) in the placebo groups was 0.1	The mean change in total quality of life score (CFQ-R) in the azithromycin groups was 1.6 higher (0.61 lower to 3.81 higher)		177 (Saiman 2003)	⊕⊕⊕⊕ high	
Quality of life: change in CFQ-R physical domain Scale from: 0 to 100 Follow-up: 6 months	The mean change in physical domain of CFQ-R score in the placebo groups was -1.9	The mean change in physical domain of CFQ-R score in the azithromycin groups was 2.7 higher (0.09 to 5.31 higher)		177 (Saiman 2003)	⊕⊕⊕⊕ high	
Quality of life: change in CFQ-R psychosocial domain Scale from: 0 to 100 Follow-up: 6 months	The mean change in psychosocial domain of CFQ-R score in the placebo groups was 1.2	The mean change in psychosocial domain of CFQ-R score in the azithromycin groups was 0.4 higher (3 lower to 3.8 higher)		177 (Saiman 2003)	⊕⊕⊕⊕ high	
Quality of life: Change in CFQ-R body image domain Scale from: 0 to 100	The mean change in body image domain of CFQ-R score in the placebo	The mean change in body image domain of CFQ-R score in the azithromycin groups was		177 (Saiman 2003)	⊕⊕⊕⊕ high	

Comparison 3. Azithromycin versus placebo

Follow-up: 6 months	groups was -0.1	3.2 higher (0.24 lower to 6.64 higher)			
---------------------	-----------------	--	--	--	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CFQ-R: cystic fibrosis questionnaire revised; CI: confidence interval; MD: mean difference; RR: risk ratio

1 Calculated by the NGA technical team from probability of remaining free from exacerbation.

2 The quality of the evidence Evidence downgraded by 2 due to serious imprecision as 95% CI crossed two default MIDs.

3 The quality of the evidence Evidence downgraded by 1 due to serious imprecision as 95% CI crossed one default MID.

Table 140: Summary clinical evidence profile: Comparison 4. Ibuprofen versus placebo

Comparison 4. Ibuprofen versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Ibuprofen				
Adverse effects: increase in abdominal pain Follow-up: 2 years	56 per 1000	14 per 1000 (2 to 124)	RR 0.26 (0.03 to 2.24)	142 (Lands 2007)	⊕⊕⊖ ⊖ low ¹	
Adverse effects: Increase in abdominal pain Follow-up: 4 years	163 per 1000	122 per 1000 (42 to 353)	RR 0.75 (0.26 to 2.17)	84 (Konstan 1995)	⊕⊕⊖ ⊖ very low ^{1,2}	
Adverse effects: Gastrointestinal bleeding Follow-up: 2 years	0 per 1000	0 per 1000 (0 to 0) ²	RR 3.08 (0.13 to 74.46)	142 (Lands 2007)	⊕⊕⊖ ⊖ low ¹	
Annual rate of change in % ideal body weight Follow-up: 4 years	The mean annual rate of change in % ideal body weight in the placebo groups was -0.94	The mean annual rate of change in % ideal body weight in the ibuprofen groups was 0.99 higher (0.17 to 1.81 higher)		84 (Konstan 1995)	⊕⊕⊖ ⊖ low ^{3,4}	
[Under 13 years at randomisation] Annual rate of change in % ideal body weight (by age)	The mean annual rate of change in % ideal body weight (by age) - under 13 years at	The mean annual rate of change in % ideal body weight (by age) in the ibuprofen groups was		49 (Konstan 1995)	⊕⊕⊖ ⊖ low ^{3,4}	

Comparison 4. Ibuprofen versus placebo					
– Follow-up: 4 years	randomisation in the placebo groups was -1.5	1.45 higher (0.33 to 2.57 higher)			
[13 years or older at randomisation] Annual rate of change in % ideal body weight (by age) Follow-up: 4 years	The mean annual rate of change in % ideal body weight (by age) - 13 years or older at randomisation in the placebo groups was -0.15	The mean annual rate of change in % ideal body weight (by age) in the ibuprofen groups was 0.34 higher (0.61 lower to 1.29 higher)		35 (Konstan 1995)	⊕⊕⊕ ⊖ very low ^{1,3}

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; MD: mean difference; RR: risk ratio

- 1 1 The quality of the evidence downgraded by 2 due to serious imprecision as 95% CI crossed 2 default MIDs.
- 2 2 Absolute effect not calculable as there are 0 events in control (placebo) arm.
- 3 3 The quality of the evidence was downgraded by 1 due to reporting bias.
- 4 4 The quality of the evidence downgraded by 1 due to serious imprecision as 95% CI crossed 1 default MID.

5 9.5.5 Economic evidence

6 No economic evaluations of immunomodulatory agents were identified in the literature
7 search conducted for this guideline. Full details of the search and economic article selection
8 flow chart can be found in Appendix E and F, respectively.

9 This area was prioritised for de novo economic modelling; consequently, a cost-utility model
10 was developed. The model uses a lifetime horizon based on the assumption that
11 immunomodulatory agents are given on a long-term basis.

12 The model takes the form of a state transition model to estimate transitions between 3 lung
13 function (FEV₁% predicted) strata. Transition probabilities between the three FEV₁% strata
14 and the number of exacerbations experienced each cycle were taken from the NMA.

15 Treatment related adverse effects and post lung transplant health states were also included
16 in the model to reflect the clinical pathway.

17 A series of deterministic sensitivity analyses were undertaken in order to test how sensitive
18 the results were to uncertainty in individual parameters. Probabilistic sensitivity analysis was
19 also conducted in the model to take account of the simultaneous effect of uncertainty relating
20 to model parameter values. The methods used to construct the model and the results of all
21 analyses are reported in Appendix K. Table 141 below presents the results from the base-
22 case, where it is clear azithromycin dominates the alternatives as it is most the effective and
23 least expensive option.

24 **Table 141: Base case results from the economic model**

Treatment	Total costs	Total QALYs	ICER
Macrolide (azithromycin)	£158,404	14.2	-
Oral corticosteroid (prednisolone)	£289,619	12.5	Dominated
NSAID (ibuprofen)	£291,035	12.3	Dominated
No treatment	£302,045	12.4	Dominated
Inhaled corticosteroid (fluticasone)	£411,046	11.1	Dominated

1 **9.5.6 Evidence statements**

2 **9.5.6.1 Evidence statements for NMA outcomes (FEV₁ % predicted and rate of pulmonary**
3 **exacerbations)**

4 See section 9.5.4.1.

5 **9.5.6.2 Evidence statements for non-NMA outcomes (nutritional status, time to next**
6 **pulmonary exacerbation, adverse events and quality of life)**

7 **9.5.6.2.1 Corticosteroids**

8 **Inhaled Beclamethasone**

9 No evidence was found for this treatment.

10 **Inhaled Budesonide**

11 No evidence was found for this treatment.

12 **Inhaled Fluticasone**

13 **Comparison 1. Fluticasone versus placebo**

14 **Time to next exacerbation**

15 Low quality evidence from 1 RCT with 171 children, young people and adults with cystic
16 fibrosis showed no clinically significant difference in the time to next exacerbation between
17 fluticasone and placebo at 6 months follow-up.

18 **Nutritional status**

19 No evidence was found for this important outcome.

20 **Quality of life**

21 No evidence was found for this important outcome.

22 **Adverse effects**

23 Moderate quality evidence from 1 RCT with 30 children with cystic fibrosis showed no
24 clinically significant difference in growth measured by change in height standard deviation
25 score between fluticasone and placebo over 12 months follow-up.

26 Moderate quality evidence from 1 RCT with 80 people with cystic fibrosis showed no
27 clinically significant difference in growth measured by change in height measured by
28 centimetres between fluticasone and placebo over 8 months follow-up.

29 **Mortality**

30 No evidence was found for this important outcome.

31 **9.5.6.2.2 Corticosteroids**

32 **IV Methylprednisolone**

33 No evidence was found for this treatment.

1 **Oral Prednisone/ Prednisolone**

2 **Comparison 2. Prednisone/ prednisolone versus placebo**

3 **Time to next exacerbation**

4 No evidence was found for this critical outcome.

5 **Nutritional status: weight and height**

6 Very low quality evidence from 1 RCT with 25 children and young people with cystic fibrosis
7 showed no clinically significant difference in weight measured in kilograms between 2 mg/kg
8 prednisolone and placebo at 12 weeks follow-up.

9 Low quality evidence from 1 observational study with 55 young people (males) with cystic
10 fibrosis showed no clinically significant difference in weight measured in kilograms between 1
11 mg/kg prednisolone and placebo at the age of 18.

12 Moderate quality evidence from 1 observational study with 52 young people (males) with
13 cystic fibrosis showed a clinically significant harmful effect of 2 mg/kg prednisolone in weight
14 measured in kilograms compared to placebo at the age of 18.

15 Very low quality evidence from 1 observational study with 43 young people (females) with
16 cystic fibrosis showed no clinically significant difference in weight measured in kilograms
17 between 1 mg/kg prednisolone and placebo at the age of 18.

18 Very low quality evidence from 1 observational study with 46 young people (females) with
19 cystic fibrosis showed no clinically significant difference in weight measured in kilograms
20 between 2 mg/kg prednisolone and placebo at the age of 18.

21 Very low quality evidence from 1 observational study with 52 young people (males) with
22 cystic fibrosis showed a clinically significant harmful effect of 1 mg/kg prednisolone in height
23 measured in centimetres compared to placebo at the age of 18.

24 Very low quality evidence from 1 observational study with 52 young people (males) with
25 cystic fibrosis showed a clinically significant harmful effect of 2 mg/kg prednisolone in height
26 measured in centimetres compared to placebo at the age of 18.

27 Very low quality evidence from 1 observational study with 43 young people (females) with
28 cystic fibrosis showed no clinically significant difference in height measured in centimetres
29 between 1 mg/kg prednisolone and placebo at the age of 18.

30 Very low quality evidence from 1 observational study with 46 young people (females) with
31 cystic fibrosis showed no clinically significant difference in height measured in centimetres
32 between 2 mg/kg prednisolone and placebo at the age of 18.

33 **Quality of life**

34 No evidence was found for this important outcome.

35 **Adverse effects**

36 Very low quality evidence from 1 RCT with 190 children with cystic fibrosis showed no
37 clinically significant difference in cataracts, diabetes mellitus, glycosuria and hyperglycaemia
38 between 1 mg/kg prednisone and placebo at 4 years follow-up.

39 Very low quality evidence from 1 RCT with 190 children with cystic fibrosis showed no
40 clinically significant difference in cataracts and diabetes mellitus between 2 mg/kg
41 prednisone and placebo at 3 years follow-up.

1 Low quality evidence from 1 RCT with 190 children with cystic fibrosis showed no clinically
2 significant difference in glycosuria between 2 mg/kg prednisone and placebo at 3 years
3 follow-up.

4 Low quality evidence from 1 RCT with 190 children with cystic fibrosis showed a clinically
5 significant harmful effect of 2 mg/kg prednisone compared with placebo for hyperglycaemia
6 at 3 years follow-up.

7 **Mortality**

8 Low quality evidence with 45 children with cystic fibrosis showed no clinically significant
9 difference in mortality between 2mg/kg prednisone and placebo at 4 years follow-up.

109.5.6.2.3 **Macrolide antibiotics**

11 **Azithromycin**

12 **Comparison 3. Azithromycin versus placebo**

13 **Time to next exacerbation**

14 High quality evidence from 2 RCT with 445 children and young people with cystic fibrosis
15 showed a clinically significant beneficial effect of azithromycin in the time to next
16 exacerbation compared to placebo at 6 months follow-up.

17 High quality evidence from 1 RCT with 82 children and young people with cystic fibrosis
18 showed a clinically significant beneficial effect of azithromycin in the time to next
19 exacerbation compared to placebo at 12 months follow-up.

20 **Nutritional status: BMI and weight**

21 Moderate quality evidence from 1 RCT with 82 children and young people with cystic fibrosis
22 showed no clinically significant difference in change in BMI z score from baseline between
23 azithromycin and placebo at 12 months follow-up.

24 Moderate quality evidence from 2 RCTs with 440 people with cystic fibrosis > 6 years
25 showed a clinically significant beneficial effect of azithromycin in weight change measured in
26 kilograms compared to placebo at 24 weeks follow-up.

27 **Quality of life**

28 High quality evidence from 1 RCT with 177 people with cystic fibrosis > 6 years showed no
29 clinically significant difference in change in quality of life (measure with CFQ-R total score,
30 and CFQ-R physical, psychosocial and body image domains) between azithromycin and
31 placebo at 6 months follow-up.

32 **Adverse effects**

33 Low quality evidence from 1 RCT with 185 people with cystic fibrosis > 6 years showed no
34 clinically significant difference in hearing impairment and tinnitus between azithromycin and
35 placebo at 6 months follow-up.

36 **Mortality**

37 No evidence was found for this important outcome.

19.5.6.2.4 NSAIDs

2 Ibuprofen

3 Comparison 4. Ibuprofen versus placebo

4 Time to next exacerbation

5 No evidence was found for this critical outcome.

6 Nutritional status: weight

7 Low quality evidence from 1 RCT with 84 people with cystic fibrosis aged 5 to 39 years
8 showed a clinically significant beneficial effect of ibuprofen in annual rate of change in
9 percent ideal body weight compared to placebo at 4 years follow-up. However, this clinically
10 beneficial significant effect was seen in children under 13 years only (n=49). Very low quality
11 evidence from this study showed no clinically significant difference in change in percent ideal
12 body weight in people with cystic fibrosis over 13 years (n=35) at 4 years follow-up.

13 Quality of life

14 No evidence was found for this important outcome.

15 Adverse effects

16 Low quality evidence from 1 RCT with 142 children with cystic fibrosis showed no clinically
17 significant difference in increase of abdominal pain between ibuprofen and placebo at 2
18 years follow-up.

19 Very low quality evidence from 1 RCT with 84 children, young people and adults with cystic
20 fibrosis showed no clinically significant difference in increase of abdominal pain between
21 ibuprofen and placebo at 4 years follow-up.

22 Low quality evidence from 1 RCT with 142 children with cystic fibrosis showed no clinically
23 significant difference in abdominal bleeding between ibuprofen and placebo at 2 years follow-
24 up.

25 Mortality

26 No evidence was found for this important outcome.

27 9.5.6.3 Monoclonal antibody

28 Omalizumab

29 No evidence was found for this treatment.

30 9.5.6.4 Economic evidence statements

31 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

32 The economic model found that azithromycin dominated (more effective and less expensive)
33 the remaining treatments in the model (NSAIDs, oral corticosteroids, inhaled corticosteroids
34 and “no treatment”). This result was also found in the extensive deterministic and
35 probabilistic sensitivity analysis that were undertaken.

1 **9.5.7 Evidence to recommendations**

2 **9.5.7.1 Relative value placed on the outcomes considered**

3 The aim of this review was to determine the clinical and cost effectiveness of
4 immunomodulatory agents in reducing pulmonary inflammation in children, young people and
5 adults with cystic fibrosis.

6 The guideline Committee identified FEV₁ % predicted, time to next exacerbation and adverse
7 events (particularly growth retardation in children) as critical outcomes for decision making;
8 whereas quality of life, nutritional status and mortality were rated as important outcomes.

9 **9.5.7.2 Consideration of clinical benefits and harms**

10 The Committee discussed the results of the evidence, and their experience in clinical
11 practice.

12 The Committee discussed the NMA results that found azithromycin had the best probability
13 of reducing exacerbations, and one of the worst for improving lung function. Based on their
14 clinical experience, the Committee agreed azithromycin can reduce exacerbations, but may
15 not necessarily improve lung function. They highlighted, however, that there is no evidence
16 that supports a direct link between lung function and clinical exacerbations, and the critical
17 outcome is to reduce the number of pulmonary exacerbations. They noted azithromycin does
18 not have such a problematic interaction profile compared to other alternative
19 immunomodulatory agents. They also noted azithromycin is usually offered as first line in
20 current practice, and they agreed to recommend it to people who are suffering a clinical
21 deterioration (as assessed by lung function), and to those who present recurrent pulmonary
22 exacerbations. They suggested that due to its pharmacokinetic profile, it can be administered
23 3 times per week, rather than daily. The Committee also discussed the duration of treatment,
24 as in practice it tends to be used for longer than the duration in studies. It was agreed that
25 treatment should be reviewed periodically to assess response.

26 The Committee agreed that oral corticosteroids can be considered if clinical deterioration
27 continues despite treatment with azithromycin, where all other treatments have been
28 maximised.

29 The Committee noted there was less evidence on fluticasone than the other treatments in the
30 NMA – it was tested in only 12 patients suggesting that more research on fluticasone is
31 needed to increase the confidence in the results. They noted that in practice, fluticasone
32 does not improve lung function to the extent the NMA inferred. In the absence of evidence-
33 base and empirical evidence to support its use, they agreed to not recommend the use of
34 inhaled corticosteroids.

35 The Committee also noted the lack of evidence for omalizumab, and that this is limited to
36 case reports.

37 The Committee acknowledged ibuprofen showed a beneficial effect in terms of lung function
38 and nutritional status. However, they were reluctant to recommend it widely, due to the high
39 dose and therapeutic drug monitoring required (which is not universally available), its
40 adverse effects profile and potential interaction with other drugs. Although the studies did not
41 show significant adverse events for ibuprofen, they emphasized longer follow-up trials are
42 needed to assess this. Also none of the studies reported on renal function, which is known to
43 be negatively affected by long-term ibuprofen use. In addition, the Committee noted
44 ibuprofen is not currently routinely used in clinical practice for the management of cystic
45 fibrosis in the UK. Nevertheless, they agreed not to write a “do not do” recommendation, as
46 they acknowledged ibuprofen may be suitable for some people (for example when
47 azithromycin is not deemed appropriate).

1 The Committee agreed it is important to assess tolerability and adverse effects in addition to
2 efficacy when making decisions about treatment.

3 **9.5.7.3 Consideration of economic benefits and harms**

4 The Committee stated that it was crucial the adverse effects of treatment were taken into
5 consideration when making their recommendations, as they may outweigh the benefits
6 related to lung function and/or exacerbations the agents can provide. As a result, the
7 economic modelling was used by the Committee as one of many ways to assess those
8 trade-offs.

9 NMAs were undertaken for this review question, to allow the treatments identified in the
10 review to be compared to a single comparator and enable the economic model to perform a
11 fully incremental analysis that compares all treatments simultaneously, to identify the most
12 cost-effective treatment. However, in the network, there were a lot of indirect comparisons
13 coming from a small number of head-to-head trials, and for most comparisons where direct
14 evidence was available, it came from a single trial. Consequently, the NMAs were not over-
15 interpreted by the Committee when making their recommendations.

16 From their clinical experience, azithromycin can reduce exacerbations, but not necessarily
17 improve lung function. Based on this, the Committee accepted the results from the NMA that
18 found azithromycin to have the best probability of reducing exacerbations and one of the
19 worst for improving lung function.

20 On the other hand, the Committee did not agree fluticasone improves lung function to the
21 extent the NMA inferred. Following this, the Committee noted that there was less evidence
22 on fluticasone than the other treatments in the NMA, suggesting that more evidence on
23 fluticasone was needed to increase their confidence in the results. The Committee also
24 questioned the inclusion of fluticasone in the economic model as it was no longer used as an
25 immunomodulatory agent in clinical practice. Subsequently, the Committee recognised that
26 the model provided sufficient evidence not to recommend fluticasone as an
27 immunomodulatory agent, as it was dominated (more expensive and less effective) by its
28 comparators in all analyses explored.

29 Conversely, the Committee agreed that azithromycin should be offered as the first line
30 treatment given that it dominated (less expensive and more effective) all alternatives in the
31 model and had the highest probability of being the most cost-effective agent in probabilistic
32 analysis. The Committee also noted that azithromycin is the first line treatment in current
33 clinical practice, particularly as it has a relatively small interactions profile.

34 The Committee advised that ibuprofen and oral corticosteroids were associated with more
35 serious treatment-related adverse effects in clinical practice than azithromycin. For this
36 reason, the Committee agreed that those agents would not be considered cost-effective
37 compared to azithromycin as they would be dominated (more expensive and less effective).
38 However, the Committee agreed they could not recommend against those agents if
39 azithromycin was no longer effective or contraindicated, given that clinical uncertainty was
40 not completely removed by the economic model. One reason for this was the absence of
41 exacerbation data for ibuprofen and the resulting ICER in the south-west quadrant (less
42 expensive and less effective) for ibuprofen compared to “no treatment” when the
43 exacerbation rates were equivalent. Given that there was evidence from the review that
44 ibuprofen improved lung function and clinical experience from the Committee that
45 immunomodulatory benefits were demonstrated by ibuprofen, the Committee concluded they
46 could not recommend against the use of ibuprofen.

47 Following this, the Committee noted that mucolytic use would be optimised if azithromycin
48 was no longer effective, before a second line immunomodulatory agent is considered. The
49 Committee also added that NSAIDs are rarely used as immunomodulatory agents in clinical
50 practice and concluded that a recommendation in favour of oral prednisolone would be

1 appropriate following no beneficial effect from azithromycin and mucolytics. This was also
2 supported by the economic model that found oral prednisolone to dominate NSAIDs and “no
3 treatment”, or, in other words, that oral prednisolone provides greater benefits at a lower
4 cost. A recommendation was also considered by the Committee to regularly review the
5 effectiveness, tolerability and side effects of immunomodulatory agents, to lead to more
6 timely identification to reduce the downstream costs to manage those events. However the
7 recommendation was subsequently removed as the Committee agreed that treatment-related
8 adverse effects should be monitored as part of good practice.

9 Overall, the results from the NMAs and subsequently, the model, were generally considered
10 to be in line with UK practice, which is why the Committee’s recommendations broadly follow
11 those results. Committee consensus and current UK practice played a large part in informing
12 their strong recommendations, where the evidence was weaker.

13 **9.5.7.4 Quality of evidence**

14 The quality of the evidence presented in this report ranged from very low to high as assessed
15 by GRADE. The main reasons that led to downgrading the quality of the evidence were:

- 16 • For the domain risk of bias, the studies were assigned the same risk of bias as in the
17 Cochrane reviews, and were not individually reviewed. The main biases that led to
18 downgrading the quality of the evidence were selection process, lack of blinding,
19 inadequate concealment, attrition, and reporting bias.
- 20 • Another reason that led to downgrading the quality of the evidence was the imprecision,
21 as confidence intervals crossed 1 or 2 MIDs. The Committee noted that some trials were
22 underpowered to detect a clinically important difference.

23 For the rate of exacerbations after short-term treatment inconsistency (heterogeneity) was
24 found to be very serious, and is likely to be due to substantial clinical heterogeneity in the
25 clinical history of patients that cannot be captured from the study reports. For other outcomes
26 no serious inconsistency was found, as most outcomes were reported by single studies,
27 though clinical heterogeneity may still have been an issue. This can lead to issues when
28 comparing treatments across different trials.

29 No issues were identified regarding the directness of the population (generalisability of the
30 results).

31 **9.5.7.5 Other considerations**

32 No equality issues were identified by the Committee for this review question.

33 The Committee agreed a research recommendation was not needed, as research is unlikely
34 to change clinical practice. In addition, the Committee were aware that there are large
35 ongoing trials in this area.

36 **9.5.7.6 Key conclusions**

37 The Committee concluded that azythromycin should be offered as long term treatment to
38 people with cystic fibrosis who are deteriorating or having pulmonary exacerbations. The
39 response to treatment should be assessed periodically, and treatment may be stopped if
40 there is no evidence of clinical benefit. Oral corticosteroids or NSIADs may be a suitable
41 alternative, but it is important to assess tolerability and side effects regularly. The use of
42 inhaled corticosteroids should not be considered as immunomodulatory treatment.

1 **9.5.8 Recommendations**

2 **92. For people with cystic fibrosis and deteriorating lung function or repeated**
3 **pulmonary exacerbations, offer long-term treatment with azithromycin at an**
4 **immunomodulatory dose^a.**

5 **93. For people who have continued deterioration in lung function, or continuing**
6 **pulmonary exacerbations while receiving long-term treatment with azithromycin,**
7 **stop azithromycin and consider oral corticosteroids at an immunomodulatory**
8 **dose.**

9 **94. Do not offer inhaled corticosteroids as an immunomodulatory treatment for**
10 **people with cystic fibrosis.**

a At the time of publication (October 2017), azithromycin did not have a UK marketing authorisation for this indication. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council's Prescribing guidance: prescribing unlicensed medicines for further information.

10 Other monitoring, assessment and management

10.1 Nutritional Interventions

Review question: What is the clinical and cost effectiveness of nutritional interventions in people with cystic fibrosis?

10.1.1 Introduction

People with cystic fibrosis are at risk of poor nutritional status which is a strong predictor of morbidity and mortality. Nutritional deficiency and malnutrition in people with cystic fibrosis is affected by both pancreatic and clinical status. Pancreatic insufficiency results in nutrient and fat soluble vitamin malabsorption, added to by the increased energy and nutrient requirement associated with chronic lung infections. Additionally, growth spurts in childhood (particularly in the first year and at adolescence) increase nutritional requirements where there is an additional risks in cystic fibrosis. Optimising nutritional status is therefore an important part of routine care for all people with cystic fibrosis, and early detection and monitoring of nutritional decline is necessary to empower potentially beneficial treatments and interventions. The strategies commonly used include dietary advice, oral supplements, enteral tube feeding, the use of appetite stimulants and behavioural interventions.

10.1.2 Description of clinical evidence

The aim of this review was to determine the clinical and cost-effectiveness of nutritional interventions for people with cystic fibrosis.

The interventions reviewed were:

- oral supplementary prescribed feeds
- enteral tube feeding
- appetite stimulants (cyproheptadine, megace)
- dietary advice, or educational interventions
- psychological and behavioural interventions

Interventions were considered “behavioural” rather than educational if they were facilitated by psychologists, psychotherapists, or psychological therapists in training.

Systematic reviews of randomised controlled trials (RCTs) and RCTs were prioritised. Systematic reviews were assessed for inclusion against the protocol, and if relevant, their quality was assessed using AMSTAR. High-quality systematic reviews were included in our review, and where possible, data and quality assessment was taken directly from the review. Individual studies were also retrieved for completeness and accuracy, and were checked for additional outcomes of interest. Low-quality systematic reviews were excluded from the review, but the list of included studies was checked to identify relevant trials.

Cross-over RCTs, RCTs with less than 10 participants, quasi-randomised trials or studies published before or during 1997 were excluded unless these studies were included in 1 of the 4 Cochrane systematic reviews that were included in this review.

Given that no evidence was found for enteral tube feeding or quality of life (a critical outcome) in the RCTs, cohort studies were assessed for inclusion and data were reported from these studies only in relation to this intervention or this critical outcome not covered by the RCTs.

1 For full details see review protocol in Appendix D.

2 Five Cochrane reviews were included in the review:

- 3 • Chinuck (2014) evaluated the effectiveness of appetite stimulants compared to placebo; 3
4 RCTs were included in this review (Eubanks 2002, Homnick 2004, Marchand 2000)
- 5 • Goldbeck (2014) evaluated the effectiveness of behavioural interventions, assessing 2
6 comparisons: behavioural intervention versus wait list control; and behavioural
7 management training plus nutritional intervention versus nutritional intervention alone; 4
8 RCTs were included in this review (Stark 1996, Stark 2009, Powers 2003)
- 9 • Morton (2015) evaluated the effectiveness of supplemental enteral tube feeding for 1
10 month or longer compared to no specific intervention; no studies were eligible for inclusion
11 in the review
- 12 • Savage (2014) evaluated the effectiveness of nutrition education compared to standard
13 treatment; 1 RCT was included in this review (Watson 2008)
- 14 • Smyth (2014) evaluated the effectiveness of oral calorie supplementation compared to
15 additional nutritional advice or no intervention; 2 RCTs (Hanning 1993, Poustie 2006) and
16 1 quasi-randomised trial (Kalnins 2005) were included in this review.

17 In addition, 1 RCT (Powers 2015) was included. It evaluated a behavioural intervention
18 compared to an educational intervention. Finally, 2 cohort studies (Bradley 2012, White
19 2013) were included. These studies evaluated the effectiveness of enteral tube feeding.

20 The size of the RCTs or cohort studies ranged from 9 to 102 participants. One study included
21 infants and children (Powers 2003), 3 studies included children (Marchand 2000, Stark 1996,
22 Powers 2015), 3 studies included children and young people (Hanning 1993, Poustie 2016,
23 Stark 2009), 4 studies included children, young people and adults (Bradley 2012, Eubanks
24 2002, Homnick 2004, Kalnins 2005), 1 study included people older than 16 (Watson 2008)
25 and 1 study included adults (White 2013).

26 Four studies (Hanning 1993, Homnick 2004, Stark 1996, Watson 2008) were considered
27 indirect in terms of the population because no inclusion criteria related to underweight were
28 specified, therefore the study population was unlikely to be representative of those who
29 would receive the interventions in clinical practice.

30 Nine studies (Bradley 2012, Eubanks 2002, Kalnins 2005, Marchand 2000, Poustie 2006,
31 Powers 2003, Stark 2009, Powers 2015, White 2013) were considered direct in terms of the
32 population because the authors specified some inclusion criteria which related to either
33 underweight, or reduced weight gain or body mass index (BMI), or pancreatic insufficiency,
34 or need for nutritional supplementation.

35 Three studies were conducted in the UK (Poustie 2016, Watson 2008, White 2013), 8 in the
36 USA (Bradley 2012, Eubanks 2002, Homnick 2004, Marchand 2000, Powers 2003, Powers
37 2015, Stark 1996, Stark 2009) and 2 in Canada (Hanning 1993, Kalnins 2005).

38 A summary of the studies included in this review is presented in Table 142. See study
39 selection flow chart in Appendix F, study evidence tables in Appendix G, list of excluded
40 studies in Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

41 10.1.3 Summary of included studies

42 A summary of the studies that were included in this review are presented in Table 142.

43 **Table 142: Summary of included studies**

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				

Study	Intervention/Comparison	Population	Outcomes	Comments
Chinuck 2014 Cochrane SR	Appetite stimulants versus placebo (Eubanks 2002, Homnick 2004, Marchand 2000)	Adults and children with CF	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight z score • Change in FEV₁ (% predicted) • Number of pulmonary exacerbations • Number of adverse effects 	
Goldbeck 2014 Cochrane SR	Comparison 1. Behavioural intervention versus usual care (Stark 1996) Comparison 2. Behavioural management training plus educational intervention versus educational intervention alone (Powers 2003, Stark 2009)	Children or adults with CF	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight z score • Change in BMI z score • Change in % ideal body weight • Change in weight % for age • Change in height (cm) • Change in height z score • Change in FEV₁ % predicted 	
Morton 2015 Cochrane SR	Supplemental enteral tube feeding for 1 month or longer versus no specific intervention (No studies were included)	People with CF of any age	No studies were identified for inclusion in this review	
Savage 2014 Cochrane SR	Nutrition education versus usual care (Watson 2008)	Individuals of all ages with CF or family members or both	<ul style="list-style-type: none"> • Change in weight • Change in FEV₁ % predicted 	
Smyth 2014 Cochrane SR	Comparison 1. Oral calorie supplementation versus usual care (Hanning 1993, Poustie 2006) Comparison 2. Oral calorie supplementation versus additional nutritional advice	People with CF	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight centile (percentile points) • Change in weight z score • Change in BMI (kg/m²) 	

Study	Intervention/Comparison	Population	Outcomes	Comments
	(Kalnins 2005)		<ul style="list-style-type: none"> • Change in BMI centile (percentile points) • Change in weight for height (percentage) • Change in height (cm) • Change in height centile (percentile points) • Change in height z score • Change in FEV₁ (% predicted) 	
Primary studies included in SRs				
Eubanks 2002 USA RCT	Intervention: Appetite stimulant <ul style="list-style-type: none"> • Megestrol acetate 10 mg/kg/day (adjusted at subsequent visits) Control: Placebo	N=17 participants (intervention: n=10; placebo: n=7) <ul style="list-style-type: none"> • Age: > 6 years • Inclusion criteria: pancreatic insufficiency, FEV₁>40% growth failure defined as no weight gain in the preceding 6 months 	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight z score • Change in FEV₁% • Number of pulmonary exacerbations • Number of adverse events: <ul style="list-style-type: none"> ○ Constipation ○ Decreased morning cortisol levels 	Included in Chinuck 2014 SR Duration: 6 months. After completion of the 6-month trial, the placebo group was offered MA for a further 6 months. Population was considered as direct.
Hanning 1993 Canada RCT	Intervention: oral calorie supplements Dietary supplements, drink powders, milk shakes, tinned puddings to achieve 25% of normal energy recommendations in addition to normal diet for 6 months Control: usual care	N=20 children and young people with CF (20 randomised, 16 studied) <ul style="list-style-type: none"> • Age: 7 to 15 years • Inclusion criteria: not reported 	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight as % expected for age and height • Change in height as % of expected for age • Change in FEV₁ % predicted 	Included in Smyth 2014 SR Follow-up: 6 months Population was considered as indirect.
Homnick 2004 USA	Intervention: Appetite stimulant Cyproheptadine	N=18 people with CF enrolled, 16 completed study	<ul style="list-style-type: none"> • Change in weight z score 	Included in Chinuck 2014 SR Follow-up: 3

Study	Intervention/Comparison	Population	Outcomes	Comments
RCT	hydrochloride 4mg 4 x daily Control: Placebo	(intervention: n=8; placebo: n=8) <ul style="list-style-type: none"> Age: ≥5 years Inclusion criteria: Ideal body weight for height <100% 	<ul style="list-style-type: none"> Change in height (cm) Change in BMI (kg/m²) Change in BMI percentile Change in % ideal body weight 	months Population was considered as indirect.
Kalnins 2005 Canada Quasi-randomised controlled trial	Intervention: Oral calorie supplementation <ul style="list-style-type: none"> High calorie drink to increase energy intake by 20% of predicted energy needs Control: Nutritional counselling <ul style="list-style-type: none"> Nutritional counselling to increase energy intake by 20% of predicted energy needs by eating high calorie foods. 	<ul style="list-style-type: none"> N=15 participants with CF were enrolled but 2 dropped out Participants were aged >10 years. Mean (SD) age on entry to trial: advice group: 16.4 years(6.7); supplement group: 19.5 years (11.3). < 90% ideal WFH or 5% reduction in ideal WFH over 3 months 	<ul style="list-style-type: none"> Change in weight (kg) Change in weight z score Change in weight for height (%) Change in % ideal body weight Change in height (cm) Change in height z score Change in FEV₁ (% predicted) 	Included in Smyth 2014 SR Interventions implemented for 3 months, follow-up: 3 and 6 months Population was considered as direct.
Marchand 2000 USA RCT	Intervention: Appetite stimulant <ul style="list-style-type: none"> Megasterol acetate 10 mg/kg/day for 12 weeks Control: Placebo	N=12 children with CF <ul style="list-style-type: none"> Age: mean age 7.4 years. Range: 21 months to 10.4 years Inclusion criteria were loss of weight or plateau in weight gain for more than 3 months, weight-for-height less than 85%, and a negative change in weight z score 	<ul style="list-style-type: none"> Change in weight z score Number of pulmonary exacerbations Fasting blood glucose levels Decreased morning cortisol levels 	Included in Chinuck 2014 SR Clinical assessment at week 0,6,12,24 and 36 Population was considered as direct.
Poustie 2006 UK RCT	Intervention 1: Oral calorie supplements Intervention 2: Routine dietary	N=102 children and young people aged 2 - 15 years with CF <ul style="list-style-type: none"> Children with at least 1 of following 	<ul style="list-style-type: none"> Change in weight (kg) Change in weight centile (percentile points) Change in BMI 	Included in Smyth 2014 SR Interventions implemented for 12 months. Outcomes at 3, 6 and 12 months

Study	Intervention/Comparison	Population	Outcomes	Comments
	advice (usual care)	criteria: BMI <25th centile but > 0.4th centile; or no increase in weight over the previous 3 months; or 5% decrease in weight from baseline over a period of < 6 months	(kg/m ²) <ul style="list-style-type: none"> • Change in BMI centile (percentile points) • Change in height (cm) • Change in height centile (percentile points) • Change in FEV₁ % predicted 	Population was considered as direct.
Powers 2003 USA RCT	Intervention 1: Behavioural management training plus educational intervention <ul style="list-style-type: none"> • Nutrition intervention with strategies for enhancing calorie intake • Behavioural management training for parents designed to encourage children to eat food consistent with CF dietary recommendations. Intervention 2: Educational intervention only	N=12 infants and children with CF <ul style="list-style-type: none"> • (intervention 1: n=7, intervention 2: n=5). • Age: Less than 3 years old. • Pancreatic insufficiency. 	<ul style="list-style-type: none"> • Change in weight (kg) • Change in % ideal body weight • Change in weight % for age • Change in height (cm) 	Included in Goldbeck 2014 SR Both groups received 8 sessions (45 to 60 minutes) over 1 year: Sessions 1 to 4 (3 months) intensive education Follow-up: 1 years Population was considered as direct.
Stark 1996 USA RCT	Intervention: Group behavioural intervention. <ul style="list-style-type: none"> • 7 weekly sessions - baseline assessment plus snack, breakfast, relaxation skills training, lunch, dinner and maintenance strategies targeted over following 7 	N=10 children with CF. <ul style="list-style-type: none"> • 1 withdrew from control group after randomisation. • Total sample n = 9 (intervention group: n=5, control group: n=4). • Age range: 5.3 years to 10.1 years; mean (SD) age: 7.3 	<ul style="list-style-type: none"> • Change in weight (kg) • Change in weight (z score) • Change in height (cm) • Change in FEV₁ % predicted 	Included in Goldbeck 2014 SR Duration of interventions: 6 weeks Population was considered as indirect.

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>sessions.</p> <p>Control: Usual care (Wait list control)</p> <ul style="list-style-type: none"> • Parent meeting and 7-day food diaries at times corresponding to baseline and last week of intervention 	<p>years (1.7).</p>		
Stark 2009 USA RCT	<p>Intervention 1: Behavioural intervention</p> <ul style="list-style-type: none"> • Behavioural intervention in group setting for change around nutrition an energy (Be-In-CHARGE!; n = 33) (available online at www.oup.com/us/pediatricpsych). <p>Intervention 2: Nutrition education</p> <ul style="list-style-type: none"> • Nutrition education in group setting 	<p>N=79 children and young people with CF</p> <ul style="list-style-type: none"> • Number randomised, n = 79 • Received the intervention, n=67 (behavioural intervention plus nutrition education: n = 33, nutrition education: n = 34) • Age: 4 to 12 years • With pancreatic insufficiency; and weight for age and height \leq 40th percentile. 	<ul style="list-style-type: none"> • Change in weight (kg) • Change in BMI z score change • Change in height (cm) • Change in height z score • FEV₁ change • Parent satisfaction (reported narratively) 	<p>Included in Goldbeck 2014 SR</p> <p>Duration of interventions: 9 weeks. Timing of sessions: 7 sessions (each 90 minutes): pre-treatment (session 1), 2 weeks later 5 weekly groups sessions (sessions 2 to 6), 2 weeks later post-treatment (session 7; follow up)</p> <p>Follow-up: up to 2 years</p> <p>Population was considered as direct.</p>
Watson 2008 UK RCT	<p>Intervention: Nutrition education</p> <ul style="list-style-type: none"> • General and disease-specific nutrition education ('Eat Well with CF') • Content: knowledge on general and disease-specific nutrition topics; self-management skills on goal setting in small incremental steps to establish new 	<p>N=74 people with CF older than 16 years of age</p> <ul style="list-style-type: none"> • participants were enrolled and stratified by disease severity into low or high risk disease. • Participants were randomly allocated into intervention (n = 37) and control (n = 37) group • 48 participants completed the study through 	<ul style="list-style-type: none"> • Change in FEV₁ (% predicted) • Change in weight • Quality of life (narrative reporting and p values only) 	<p>Included in Savage 2014 SR</p> <p>Duration of the intervention: 10 weeks. Outcomes measured at 6 and 12 months</p> <p>Population was considered as indirect.</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
	behaviours <ul style="list-style-type: none"> • Mode of delivery: written material; supplementary workshops (introductory, weeks 5 and 10) and weekly telephone calls delivered by a dietitian Control: Usual care	to 12-month follow-up assessment (23 in intervention group, 25 in control group) <ul style="list-style-type: none"> • Age: intervention group 26.4 (17.2 - 43.2) years; control group 24.2 (16.9 - 38.1) years 		
Additional primary studies				
Bradley 2012 United States Cohort study	Intervention: Gastrostomy Control: No gastrostomy (usual care)	N=40 people with CF (20 in the intervention group, 20 in the control group) <ul style="list-style-type: none"> • Age range: 2 to 20 years • Each child in the intervention group was pair-matched on age, sex, pancreatic status, BMI and lung function with a children from the control group • Children who had a gastrostomy for reasons other than nutritional supplementation were excluded 	<ul style="list-style-type: none"> • Change in weight z-score • Change in BMI z-score • Change in height z-score • Change in FEV₁ % predicted 	Outcomes measured at 6 and 12 months Population was considered as direct.
Powers 2015 USA RCT	Behavioural intervention <ul style="list-style-type: none"> • Individualized nutritional counselling targeting increased energy intake and behavioural child management skills Control:	N=78 children with CF and pancreatic insufficiency (intervention: n=36, control: n=42) <ul style="list-style-type: none"> • Age: 2 to 6 years • Confirmed pancreatic insufficiency; no restrictions in consuming a 	<ul style="list-style-type: none"> • Change in weight z score • Change in height z score • Number of adverse events (digestive system) 	Both treatments were delivered in person or telehealth (via telephone) Sessions occurred weekly for 8 weeks then monthly for 4 months (6 months). The control arm served as a behavioural placebo

Study	Intervention/Comparison	Population	Outcomes	Comments
	education and attention control treatment <ul style="list-style-type: none"> Education on general nutrition information and other topics (e.g. infection control and bicycle safety) 	high-fat diet.		controlling for attention and contact frequency. Participants then returned to standard care for 1 year. Follow-up: 18 months Population was considered as direct.
White 2013 UK Cohort study	Intervention: Enteral tube feeding <ul style="list-style-type: none"> Supplemental enteral tube feeding administered over 3 years Overnight enteral tube feed Control: Usual care	<ul style="list-style-type: none"> N=21 adults with CF (intervention: n=15, control: n=6) Mean (SD) age: intervention: 21.8 (3.6); control: 23.0 (5.7) All study participants fulfilled the criteria for commencement of enteral tube feeding (CF Trust, 2002): BMI<19 kg/m² and/or 5% acute weight loss over a 2 month period with a failure or oral nutritional supplements to adequately improve nutritional status. 	<ul style="list-style-type: none"> Change in weight (kg) Change in BMI (kg/m²) Change in FEV₁ (%) Change in IV treatment days 	Initially n=17 accepted enteral tube feeding. However 2 people died during the baseline year and subsequent analyses were conducted on the surviving participants. Control group: people who declined enteral tube feeding Follow-up: 1 year Population was considered as direct.

1 BMI: body mass index; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; RCT: randomised
2 controlled trial; SR: systematic review; WFH: weight for height

3 10.1.4 Clinical evidence profile

4 The summary clinical evidence profiles for this review question (nutrition interventions in
5 people with CF) are presented in Table 143 to Table 150.

6 **Table 143: Summary clinical evidence profile: Comparison 1.1 Oral calorie**
7 **supplementation versus usual care**

Comparison 1.1 Oral calorie supplementation versus usual care					
Outcomes	Illustrative comparative risks* (95% CI)	Relative effect	No of Participan	Quality of the	Comm ents

Comparison 1.1 Oral calorie supplementation versus usual care						
	Assumed risk	Corresponding risk	(95% CI)	ts (studies)	evidence (GRADE)	
	Usual care	Oral calorie supplementation				
Change in weight (kg) Follow-up: 3 months	The mean change in weight (kg) in the control group was 0.77	The mean change in weight (kg) oral calorie supplementation groups was 0.34 higher (0.07 lower to 0.75 higher)		99 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in weight (kg) Follow-up: 6 months	The mean change in weight (kg) in the usual care group was 1.33 in 1 study and 1.72 in the other study	The mean change in weight (kg) in the oral calorie supplementation groups was 0.47 higher (0.07 lower to 1.02 higher)		117 (Hanning 1993, Poustie 2006)	⊕⊕⊖⊖ low ^{1,2,3}	
Change in weight (kg) Follow-up: 1 year	The mean change in weight (kg) in the usual care group was 2.97	The mean change in weight (kg) in the oral calorie supplementation groups was 0.16 higher (0.68 lower to 1 higher)		102 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in height (cm) Follow-up: 3 months	The mean change in height (cm) in the usual care group was 1.68	The mean change in height (cm) in the oral calorie supplementation groups was 0.03 lower (0.36 lower to 0.3 higher)		99 (Poustie 2006)	⊕⊕⊕⊕ high	
Change in height (cm) Follow-up: 6 months	The mean change in height (cm) in the usual care group was 3.56	The mean change in height (cm) in the oral calorie supplementation groups was 0.47 lower (1.32 lower to 0.38 higher)		101 (Poustie 2006)	⊕⊕⊕⊕ high	
Change in height (cm) Follow-up: 1 year	The mean change in height (cm) in the usual care group was 5.85	The mean change in height (cm) in the oral calorie supplementation groups was 0.06 higher (0.5 lower to 0.62 higher)		102 (Poustie 2006)	⊕⊕⊕⊕ high	
Change in weight as % expected for age and height	The mean change in weight as % expected for age and height	The mean change in weight as % expected for age and height in the oral calorie		16 (Hanning 1993)	⊕⊖⊖⊖ very low ^{2,4,5}	

Comparison 1.1 Oral calorie supplementation versus usual care						
Follow-up: 6 months	in the usual care group was -2.7	supplementation groups was 3.3 higher (6.27 lower to 12.87 higher)				
Change in BMI (kg/m ²) Follow-up: 3 months	The mean change in BMI (kg/m ²) in the usual care group was 0.05	The mean change in BMI (kg/m ²) in the oral calorie supplementation groups was 0.14 higher (0.08 lower to 0.36 higher)		99 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in BMI (kg/m ²) Follow-up: 6 months	The mean change in BMI (kg/m ²) in the usual care group was 0.15	The mean change in BMI (kg/m ²) in the oral calorie supplementation groups was 0.24 higher (0.06 lower to 0.54 higher)		101 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in BMI (kg/m ²) Follow-up: 1 year	The mean change in BMI (kg/m ²) in the usual care group was 0.24	The mean change in BMI (kg/m ²) in the oral calorie supplementation groups was 0.08 higher (0.28 lower to 0.44 higher)		102 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in BMI (centile) Follow-up: 3 months	The mean change in BMI (centile) in the usual care group was -0.56	The mean change in BMI (centile) in the oral calorie supplementation groups was 3.28 higher (0.7 lower to 7.26 higher)		99 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in BMI (centile) Follow-up: 6 months	The mean change in BMI (centile) in the usual care group was -1.29	The mean change in BMI (centile) in the oral calorie supplementation groups was 5.75 higher (0.22 to 11.28 higher)		101 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in BMI (centile) Follow-up: 1 year	The mean change in BMI (centile) the usual care group was -2.32	The mean change in BMI (centile) in the oral calorie supplementation groups was 2.99 higher (2.69 lower to 8.67 higher)		102 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	
Change in weight (centile) Follow-up: 3 months	The mean change in weight (centile) - in the usual care	The mean change in weight (centile) in the oral calorie supplementation groups was		99 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹	

Comparison 1.1 Oral calorie supplementation versus usual care					
	group was 0.4	1.72 higher (0.59 lower to 4.03 higher)			
Change in weight (centile) Follow-up: 6 months	The mean change in weight (centile) in the usual care group was 0.63	The mean change in weight (centile) in the oral calorie supplementation groups was 2.12 higher (0.94 lower to 5.18 higher)		101 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹
Change in weight (centile) Follow-up: 1 year	The mean change in weight (centile) in the usual care group was -1	The mean change in weight (centile) in the oral calorie supplementation groups was 1.83 higher (1.77 lower to 5.43 higher)		102 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹
Change in height (centile) Follow-up: 3 months	The mean change in height (centile) - in the usual care group was 1.13	The mean change in height (centile) in the oral calorie supplementation groups was 0.56 lower (2.04 lower to 0.92 higher)		99 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹
Change in height (centile) Follow-up: 6 months	The mean change in height (centile) - in the usual care group was 1.98	The mean change in height (centile) in the oral calorie supplementation groups was 1.74 lower (4.4 lower to 0.92 higher)		101 (Poustie 2006)	⊕⊕⊕⊕ high
Change in height (centile) Follow-up: 1 year	The mean change in height (centile) in the usual care group was 1.18	The mean change in height (centile) in the oral calorie supplementation groups was 0.65 lower (3.11 lower to 1.81 higher)		102 (Poustie 2006)	⊕⊕⊕⊖ moderate ¹
Change in height as % of expected for age Follow-up: 6 months	The mean change in height as % of expected for age in the usual care group was 1.7	The mean change in height as % of expected for age in the oral calorie supplementation groups was 1.6 lower (21.54 lower to 18.34 higher)		16 (Hanning 1993)	⊕⊖⊖⊖ very low ^{2,4,5}
Adverse effects	No evidence was found				
Change in FEV ₁ % predicted	The mean change in FEV ₁ % predicted in	The mean change in FEV ₁ % predicted in the oral calorie		69 (Poustie 2006)	⊕⊕⊕⊖ moderate ⁶

Comparison 1.1 Oral calorie supplementation versus usual care						
Follow-up: 3 months	the usual care group was 5.37	supplementation groups was 7.92 lower (13.89 to 1.95 lower)				
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 months	The mean change in FEV ₁ % predicted in the usual care group was -4.3 in 1 study and 1.61 in the other study	The mean change in FEV ₁ % predicted in the oral calorie supplementation groups was 3.84 lower (9.63 lower to 1.94 higher)		86 (Hanning 1993, Poustie 2006)	⊕⊕⊕⊕ low ^{2,3,6}	
Change in FEV ₁ % predicted Follow-up: 1 year	The mean change in FEV ₁ % predicted in the usual care group was -1.5	The mean change in FEV ₁ % predicted in the oral calorie supplementation groups was 1.91 lower (8.57 lower to 4.75 higher)		70 (Poustie 2006)	⊕⊕⊕⊕ moderate ⁶	
Quality of life	No evidence was found					
Pulmonary exacerbations	No evidence was found					
Patient or carer satisfaction	No evidence was found					

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; cm: centimetres; FEV₁: forced expiratory volume in 1 second; kg: kilogrammes; kg/m²: kilogrammes per metre square; MD: mean difference

- 1 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 2 The quality of the evidence was downgraded by 1 because of high risk of bias in relation to the randomisation (the treated group appeared to be in better clinical condition at baseline in 1 study).
- 3 The inclusion criteria in the paper by Hanning et al. did not mention underweight therefore the population in the study is unlikely to be representative of people who would receive oral supplements in clinical practice; however the quality of the evidence was not downgraded because the inclusion criteria in the paper by Poustie et al. are likely to be representative of people who receive oral supplements in clinical practice
- 4 The quality of the evidence was downgraded by 1 because the inclusion criteria did not mention underweight therefore the population in the study is unlikely to be representative of people who would usually receive oral supplements
- 5 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 defaults MIDs
- 6 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID

Table 144: Summary clinical evidence profile: Comparison 1.2 Oral calorie supplementation versus nutrition advice

Comparison 1.2 Oral calorie supplementation versus nutrition advice						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Nutrition advice	Oral calorie supplementation				

Comparison 1.2 Oral calorie supplementation versus nutrition advice					
Change in weight (kg) Follow-up: 3 months	The mean change in weight (kg) in the nutrition advice group was 2.15	The mean change in weight (kg) in the oral calorie supplementation groups was 0.69 lower (3.3 lower to 1.92 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in weight for height (%) Follow-up: 3 months	The mean change in weight for height (%) in the nutrition advice group was 1.67	The mean change in weight for height (%) in the oral calorie supplementation groups was 0.96 lower (5.23 lower to 3.31 higher)		19 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in weight z score Follow-up: 3 months	The mean change in weight z score in the nutrition advice group was 0.1	The mean change in weight z score in the oral calorie supplementation groups was 0 higher (0.59 lower to 0.59 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in weight z score Follow-up: 6 months	The mean change in weight z score in the nutrition advice group was 0.2	The mean change in weight z score in the oral calorie supplementation groups was 0.3 lower (0.98 lower to 0.38 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in % ideal body weight Follow-up: 3 months	The mean change in % ideal body weight in the nutrition advice group was 1	The mean change in % ideal body weight in the oral calorie supplementation groups was 2 lower (10.59 lower to 6.59 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in % ideal body weight Follow-up: 6 months	The mean change in % ideal body weight in the nutrition advice group was 0	The mean change in % ideal body weight in the oral calorie supplementation groups was 3 lower (11.59 lower to 5.59 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}
Change in height (cm) Follow-up: 3 months	The mean change in height (cm) in the nutrition	The mean change in height (cm) in the oral calorie		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}

Comparison 1.2 Oral calorie supplementation versus nutrition advice						
	advice group was 2.55	supplementation groups was 0.38 lower (3.05 lower to 2.29 higher)				
Change in height z score Follow-up: 3 months	The mean change in height z score - in the nutrition advice group was 0.1	The mean change in height z score in the oral calorie supplementation groups was 0 higher (0.96 lower to 0.96 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}	
Change in height z score Follow-up: 6 months	The mean change in height z score - in the nutrition advice group was 0.2	The mean change in height z score in the oral calorie supplementation groups was 0.1 lower (1.07 lower to 0.87 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,2}	
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 months	The mean change in FEV ₁ % predicted in the nutrition advice group was 1.6	The mean change in FEV ₁ % predicted in the oral calorie supplementation groups was 8.2 lower (23.37 lower to 6.97 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,3}	
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 months	The mean change in FEV ₁ % predicted in the nutrition advice group was 4	The mean change in FEV ₁ % predicted in the oral calorie supplementation groups was 8 lower (26.96 lower to 10.96 higher)		13 (1 Kalnins 2005)	⊕⊖⊖⊖ very low ^{1,3}	
Quality of life	No evidence was found					
Pulmonary exacerbations	No evidence was found					
Adverse effects	No evidence was found					
Patient or carer satisfaction	No evidence was found					
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). Abbreviations: confidence interval; CF: cystic fibrosis; cm: centimetres; FEV₁: forced expiratory volume in 1 second; kg: kilogrammes; MD: mean difference</p>						

1 The quality of the evidence was downgraded by 2 because of unclear risk of bias in relation to randomisation, high risk of bias in relation to allocation concealment, and inability to make judgment in relation to other bias.
2 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDIs
3 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDIs

1
2

Table 145: Summary clinical evidence profile: Comparison 2. Enteral tube feeding versus usual care

Comparison 2. Enteral tube feeding versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual care	Enteral tube feeding				
Change in weight (kg) Follow-up: 1 year	The mean change in weight (kg) the usual care group was -0.3	The mean change in weight (kg) in the enteral tube feeding groups was 7.60 higher (4.74 to 10.46 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	
Change in weight (kg) Follow-up: 2 years	The mean change in weight (kg) the usual care group was -0.8	The mean change in weight (kg) in the enteral tube feeding groups was 9.10 higher (5.43 to 12.77 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	
Change in weight (kg) Follow-up: 3 years	The mean change in weight (kg) the usual care group was -0.1	The mean change in weight (kg) in the enteral tube feeding groups was 9.00 higher (5.21 to 12.79 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	
Change in weight z score - Scale from: -4 to 4. Follow-up: 6 months	The mean change in weight z score in the usual care group was 0.05	The mean change in weight z score in the enteral tube feeding groups was 0.62 higher (0.27 to 0.97 higher)		40 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,3}	
Change in weight z score - Scale from: -4 to 4. Follow-up: 1 years	The mean change in weight z score in the usual care group was 0.2	The mean change in weight z score in the enteral tube feeding groups was 0.44 higher (0.11 to 0.77 higher)		40 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,3}	
Change in height z-score Scale from: -	The mean change in height z-score in the	The mean change in height z-score in the enteral tube		40 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,3}	

Comparison 2. Enteral tube feeding versus usual care						
4 to 4. Follow-up: 6 months	usual care group was 0.3	feeding groups was 0.2 higher (0.19 lower to 0.59 higher)				
Change in height z-score Scale from: -4 to 4. Follow-up: 1 years	The mean change in height z-score in the usual care group was 0	The mean change in height z-score in the enteral tube feeding groups was 0.1 higher (0.29 lower to 0.49 higher)		40 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,3}	
Change in BMI z score Scale from: -4 to 4. Follow-up: 6 months	The mean change in BMI z score in the usual care group was 0.08	The mean change in BMI z score in the enteral tube feeding groups was 0.82 higher (0.48 to 1.16 higher)		40 (Bradley 2012)	⊕⊕⊕⊕ very low ²	
Change in BMI z score Scale from: -4 to 4. Follow-up: 1 years	The mean change in BMI z score in the usual care group was 0.39	The mean change in BMI z score in the enteral tube feeding groups was 0.39 higher (0.09 to 0.69 higher)		40 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,3}	
Change in BMI (kg/m ²) Follow-up: 1 years	The mean change in BMI in the usual care group was -0.2	The mean change in BMI (kg/m ²) in the enteral tube feeding groups was 2.90 higher (2.2 to 3.6 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	
Change in BMI (kg/m ²) Follow-up: 2 years	The mean change in BMI in the usual care group was -0.3	The mean change in BMI (kg/m ²) in the enteral tube feeding groups was 3.20 higher (2.33 to 4.07 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	
Change in BMI (kg/m ²) Follow-up: 3 years	The mean change in BMI in the usual care group was 0.8	The mean change in BMI (kg/m ²) in the enteral tube feeding groups was 2.50 higher (1.55 to 3.45)		21 (White 2013)	⊕⊕⊕⊕ very low ¹	

Comparison 2. Enteral tube feeding versus usual care					
Change in FEV ₁ %predicted Scale from: 0 to 100. Follow-up: 6 months	The mean change in FEV ₁ %predicted in the usual care group was 3.2	higher) The mean change in FEV ₁ %predicted in the enteral tube feeding groups was 4.5 lower (16.18 lower to 7.18 higher)		27 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,4}
Change in FEV ₁ %predicted Scale from: 0 to 100. Follow-up: 1 years	The mean change in FEV ₁ %predicted in the usual care group was 6.6	The mean change in FEV ₁ %predicted in the enteral tube feeding groups was 8.2 lower (20.5 lower to 4.1 higher)		27 (Bradley 2012)	⊕⊕⊕⊕ very low ^{2,5}
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1 years	The mean change in FEV ₁ % predicted in the usual care group was -5.3	The mean change in FEV ₁ % predicted in the enteral tube feeding groups was 10.60 higher (10.34 lower to 31.54 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,4}
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 2 years	The mean change in FEV ₁ % predicted in the usual care group was -8	The mean change in FEV ₁ % predicted in the enteral tube feeding groups was 12.20 higher (2.57 lower to 26.97 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,5}
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 3 years	The mean change in FEV ₁ % predicted in the usual care group was -11	The mean change in FEV ₁ % predicted in the enteral tube feeding groups was 12.20 higher (1.84 lower to 26.24 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,5}
Change in IV treatment days Follow-up: 1 years	The mean change in IV treatment days in the usual care group was 2.8	The mean change in IV treatment days in the enteral tube feeding groups was 17.90 higher (5.96 lower to 41.76 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,3}
Change in IV treatment days Follow-	The mean change in IV treatment	The mean change in IV treatment days		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,3}

Comparison 2. Enteral tube feeding versus usual care					
up: 2 years	days in the usual care group was -8	in the enteral tube feeding groups was 36.00 higher (5.06 to 66.94 higher)			
Change in IV treatment days Follow-up: 3 years	The mean change in IV treatment days in the usual care group was 7	The mean change in IV treatment days in the enteral tube feeding groups was 36.20 higher (6.29 lower to 78.69 higher)		21 (White 2013)	⊕⊕⊕⊕ very low ^{1,3}
Quality of life	No evidence available				
Patient or carer satisfaction	No evidence available				
Adverse events	No evidence available				
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: BMI: body mass index; confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; IV: intravenous; k/m²g: kilogrammes per square metre; MD: mean difference</p>					

- 1 The quality of the evidence was downgraded by 2 due to high risk of bias in relation to selection of the study population and comparability of the 2 groups
2
3 The quality of the evidence was downgraded by 1 because of high risk of bias in relation to comparability
4 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
5 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs
6 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID

7 **Table 146: Summary clinical evidence profile: Comparison 3. Appetite stimulants**
8 **versus placebo**

Comparison 3. Appetite stimulants versus placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Appetite stimulants				
Change in weight in kg. Follow-up: 3 months	The mean change in weight in kg. in the placebo group was 1.3 in 1 study, 1.1 in the other study	The mean change in weight in kg. in the appetite stimulants groups was 2.97 higher (0.94 to 4.99 higher)		33 (Eubanks 2002, Homnick 2004)	⊕⊕⊕⊕ low ¹	
Change in weight in kg.	The mean change in	The mean change in		17 (Eubanks	⊕⊕⊕⊕ low ²	

Comparison 3. Appetite stimulants versus placebo						
Follow-up: 6 months	weight in kg. in the placebo group was 1.5	weight in kg. in the appetite stimulants groups was 3.8 higher (1.27 to 6.33 higher)		2002)		
Change in weight z score Scale from: -4 to 4. Follow-up: 3 months	The mean change in weight z score in the 3 placebo groups was: 0.07, 0.04, -0.05	The mean change in weight z score in the appetite stimulants groups was 0.61 higher (0.29 to 0.93 higher)		40 (Eubanks 2002, Homnick 2004, Marchand 2000)	⊕⊕⊕⊖ low ³	
Change in weight z score Scale from: -4 to 4. Follow-up: 6 months	The mean change in weight z score in the placebo group was 0.02	The mean change in weight z score in the appetite stimulants groups was 0.74 higher (0.26 to 1.22 higher)		17 (Eubanks 2002)	⊕⊕⊕⊖ low ²	
Change in height (cm) Follow-up: 3 months	The mean change in height in the placebo group was 1	The mean change in height (cm) in the appetite stimulants groups was 0.2 higher (11.88 lower to 12.28 higher)		16 (Homnick 2004)	⊕⊖⊖⊖ very low ^{4,5,6}	
Change in BMI (kg/m ²) Follow-up: 3 months	The mean change in BMI in the placebo group was 0.29	The mean change in BMI (kg/m ²) in the appetite stimulants groups was 0.88 higher (0.76 lower to 2.52 higher)		16 (Homnick 2004)	⊕⊖⊖⊖ very low ^{4,5,7}	
Change in BMI percentile Follow-up: 3 months	The mean change in BMI percentile in the placebo group was 1.78	The mean change in BMI centile in the appetite stimulants groups was 11.1 higher (0.15 to 22.05 higher)		16 (Homnick 2004)	⊕⊖⊖⊖ very low ^{4,5,7}	
Change in % ideal body weight Follow-up: 3 months	The mean change in % ideal body weight in the placebo group was	The mean change in % ideal body weight in the appetite stimulants		16 (Homnick 2004)	⊕⊖⊖⊖ very low ^{4,5,7}	

Comparison 3. Appetite stimulants versus placebo						
	1.15	groups was 5.14 higher (0.2 to 10.08 higher)				
Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 3 months	The mean change in FEV ₁ % predicted in the placebo group was -3.7	The mean change in FEV ₁ % predicted in the appetite stimulants groups was 13.55 higher (1.88 lower to 28.98 higher)		17 (Eubaks 2002)	⊕⊕⊕⊕ very low ^{2,8}	
Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 6 months	The mean change in FEV ₁ % predicted in the placebo group was 0.83	The mean change in FEV ₁ % predicted - at 6 months in the appetite stimulants groups was 5.64 higher (4.43 lower to 15.71 higher)		17 (Eubaks 2002)	⊕⊕⊕⊕ very low ^{2,8}	
Quality of life	No evidence was found					
Number of pulmonary exacerbations Follow-up: 3 months	Study population		RR 1.67 (0.69 to 4)	12 (Marchand 2000)	⊕⊕⊕⊕ very low ^{6,9}	
	500 per 1000	835 per 1000 (345 to 1000)				
Adverse effects: constipation Follow-up: 6 months	-	-	RR 2.18 (0.1 to 46.92)	17 (Eubanks 2002)	⊕⊕⊕⊕ very low ^{2,6}	
Adverse effects: high blood glucose Follow-up: 3 months	Fasting blood glucose levels remained unchanged in both groups. Values not reported			12 (Marchand 2000)	⊕⊕⊕⊕ low ¹⁰	
Adverse effects: decreased morning cortisol levels <0.6mcg/dl Follow-up: 3 months	-	All participants in the intervention group had normal morning cortisol levels at baseline; at follow-up 4 out of the 6 participants in the intervention group had morning cortisol levels decreased to <0.6mcg/dl		12 (Marchand 2000)	⊕⊕⊕⊕ low ¹⁰	
Adverse effects:	-	-	RR 10.91	17 (Eubanks)	⊕⊕⊕⊕ very	

Comparison 3. Appetite stimulants versus placebo					
decreased morning cortisol levels <30nmol/L Follow-up: 6 months			(0.72 to 164.61)	2002)	low ^{2,6}
Patient or carer satisfaction	No evidence available				
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: BMI: body mass index; confidence interval; CF: cystic fibrosis; FEV ₁ : forced expiratory volume in 1 second; IV: intravenous; kg: kilogrammes; kg/m ² : kilogrammes per square metre; MD: mean difference; nmol/L: nanomoles per litre; RR: risk ratio					

- 1 The quality of the evidence was downgraded by 2 due to very serious risk of bias in relation to the evidence from the Eubanks 2002 paper and serious risk of bias in relation to the evidence from the Hornick 2004 paper
- 2 The quality of the evidence was downgraded by 2 due to unclear risk of bias in relation to allocation concealment, and high risk of bias in relation to incomplete outcome data and selective reporting.
- 3 The quality of the evidence was downgraded by 2 due to very serious risk of bias in relation to the evidence from the Eubanks 2002 paper, serious risk of bias in relation to the evidence from the Hornick 2004 paper, and very serious risk of bias in relation to the evidence from the Marchand 2000 paper.
- 4 The quality of the evidence was downgraded by 1 due to unclear risk of bias in relation to allocation concealment and high risk of bias in relation to selective reporting.
- 5 The evidence was downgraded by 1 because ideal body weight for height <100% was an inclusion criteria. However in clinical practice some people with ideal body weight for height under this cut-off may be considered with normal weight and therefore would not be the target population of appetite stimulants.
- 6 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs
- 7 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 8 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID
- 9 The quality of the evidence was downgraded by 2 due to unclear risk of bias in relation to random sequence generation and allocation concealment, and high risk of bias in relation to incomplete outcome data and selective reporting
- 10 The quality of the evidence was downgraded by 2 due to unclear risk of bias in relation to random sequence generation and allocation concealment, and high risk of bias in relation to incomplete outcome data, selective reporting, and bad reporting (relevant values not provided)
- a. Reversible decrease: 30+ days after treatment levels went back up to 270 +-6.9 nmol/L

Table 147: Summary clinical evidence profile: Comparison 4. Nutrition education versus usual care

Comparison 4. Nutrition education versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Standard treatment	Nutrition education				
Change in weight (kg) Follow-up: 6 months	The mean change in weight (kg) - in the standard treatment group was 0.8	The mean change in weight (kg) in the nutrition education groups was 0.4 lower (4.85 lower to 4.05 higher)		48 (Watson 2008)	⊕⊕⊕⊕ very low ^{1,2,3}	
Change in weight (kg)	The mean change in	The mean change in		48 (Watson	⊕⊕⊕⊕ low ^{1,2,4}	

Comparison 4. Nutrition education versus usual care						
Follow-up: 1 year	weight (kg) in the standard treatment group was 1.2	weight (kg) in the nutrition education groups was 0.4 lower (4.87 lower to 4.07 higher)		2008)		
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 months	The mean change in FEV ₁ % predicted in the standard treatment group was 0.81	The mean change in FEV ₁ % predicted in the nutrition education groups was 1.49 higher (8.84 lower to 11.82 higher)		48 (Watson 2008)	⊕⊖⊖⊖ very low ^{1,2,3}	
Change in FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 1 year	The mean change in FEV ₁ % predicted in the standard treatment group was -0.79	The mean change in FEV ₁ % predicted in the nutrition education groups was 0.99 higher (9.29 lower to 11.27 higher)		48 (Watson 2008)	⊕⊖⊖⊖ very low ^{1,2,5}	
Quality of life: CF-QOL, physical functioning Scale from: 0 to 100 Follow-up: 6 months		p-value= 0.05		48 (Watson 2008)	⊕⊕⊖⊖ low ^{2,6}	
Quality of life: CF-QOL, physical functioning Scale from: 0 to 100 Follow-up: 12 months		p-value= 0.61		48 (Watson 2008)	⊕⊕⊖⊖ low ^{2,6}	
Quality of life: CF-QOL, social functioning Scale from: 0 to 100 Follow-up: 6 months		p-value= 0.85		48 (Watson 2008)	⊕⊕⊖⊖ low ^{2,6}	
Quality of life: CF-QOL, social functioning Scale from: 0		p-value= 0.54		48 (Watson 2008)	⊕⊕⊖⊖ low ^{2,6}	

Comparison 4. Nutrition education versus usual care						
to 100						
Follow-up: 12 months						
Quality of life: treatment issues CF-QOL, Follow-up: 6 months		p-value= 0.74		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, treatment issues Scale from: 0 to 100 Follow-up: 12 months		p-value= 0.68		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, chest symptoms Follow-up: 6 months		p-value= 0.59		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, chest symptoms Scale from: 0 to 100 Follow-up: 12 months		p-value= 0.62		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, emotional responses Scale from: 0 to 100 Follow-up: 6 months		p-value= 0.45		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, emotional responses Scale from: 0 to 100 Follow-up: 12 months		p-value= 0.07		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, concerns for the future Scale from: 0 to 100 Follow-up: 6 months		p-value= 0.46		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life:		p-value= 0.03		48	⊕⊕⊕⊕	

Comparison 4. Nutrition education versus usual care						
CF-QOL, concerns for the future Scale from: 0 to 100 Follow-up: 12 months				(Watson 2008)	low ^{2,6}	
Quality of life: CF-QOL, interpersonal relationship Scale from: 0 to 100 Follow-up: 6 months		p-value= 0.75		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, interpersonal relationship Scale from: 0 to 100 Follow-up: 12 months		p-value= 0.64		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, body image Scale from: 0 to 100 Follow-up: 6 months		p-value=0.24		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, body image Scale from: 0 to 100 Follow-up: 12 months		p-value=0.59		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, career issues Scale from: 0 to 100 Follow-up: 6 months		p-value=0.15		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Quality of life: CF-QOL, career issues Scale from: 0 to 100 Follow-up: 12 months		p-value=0.28		48 (Watson 2008)	⊕⊕⊕⊕ low ^{2,6}	
Pulmonary exacerbations	No evidence was found					
Adverse	No evidence was found					

Comparison 4. Nutrition education versus usual care

effects	
Patient or carer satisfaction	No evidence was found

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; CF: cystic fibrosis; CFQOL: cystic fibrosis quality of life questionnaire; FEV₁: forced expiratory volume in 1 second; kg: kilogrammes; MD: mean difference

1 The quality of the evidence was not downgraded despite unclear risk of bias in relation to blinding and selective reporting, because objective measures are unlikely to be influenced by the lack of blinding

2 The quality of the evidence was downgraded by 1 because there were no inclusion criteria related to underweight, therefore the study population is unlikely to be representative of people who would receive this intervention in clinical practice

3 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs

4 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

5 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs

6 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to selective reporting and high risk of bias due to bad reporting (only p values and U test statistic provided)

Table 148: Summary clinical evidence profile: Comparison 5.1 Behavioural intervention versus usual care

Comparison 5.1 Behavioural intervention versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual care	Behavioural intervention				
Change in weight (kg) Follow-up: 6 weeks	The mean change in weight (kg) in the usual care group was 0	The mean change in weight (kg) in the behavioural intervention groups was 1.7 higher (4.02 lower to 7.42 higher)		9 (Stark 1996)	⊕⊕⊕⊕ very low ^{1,2,3}	
Change in height (cm) Follow-up: 6 weeks	The mean change in height (cm) in the usual care group was 1.3	The mean change in height (cm) in the behavioural intervention groups was 0.1 lower (16.75 lower to 16.55 higher)		9 (Stark 1996)	⊕⊕⊕⊕ very low ^{1,2,3}	
Change in weight z score Follow-up: 6 weeks	The mean change in weight z score in the usual care group was -0.05	The mean change in weight z score in the behavioural intervention groups was 0.5 higher (0.19 lower to 1.19 higher)		9 (Stark 1996)	⊕⊕⊕⊕ very low ^{1,2,4}	
Change in	The mean	The mean		9	⊕⊕⊕⊕	

Comparison 5.1 Behavioural intervention versus usual care					
FEV ₁ % predicted Scale from: 0 to 100 Follow-up: 6 weeks	change in FEV ₁ % predicted in the usual care group was 0.5	change in FEV ₁ % predicted in the behavioural intervention groups was 6.5 lower (28.09 lower to 15.09 higher)		(Stark 1996)	very low ^{1,2,5}
Quality of life	No evidence was found				
Pulmonary exacerbations	No evidence was found				
Adverse effects	No evidence was found				
Patient or carer satisfaction	No evidence was found				
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: CI: confidence interval; CF: cystic fibrosis; cm: centimetres; FEV ₁ : forced expiratory volume in 1 second; MD: mean difference					

- 1 The quality of the evidence was downgraded by 1 due to unclear risk of bias in relation to random sequence generation, allocation concealment and selective reporting. Cochrane rated the risk of bias for blinding as high however objective measures are unlikely to be influenced by the lack of blinding.
- 2 The quality of the evidence was downgraded by 1 because there were no inclusion criteria related to underweight or calorie intake therefore the study population is unlikely to be representative of people who would receive this intervention in clinical practice
- 3 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs
- 4 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 5 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs

10 **Table 149: Summary clinical evidence profile: Comparison 5.2 Behavioural**
11 **intervention versus education and attention control treatment**

Comparison 5.2 Behavioural intervention versus education and attention control treatment						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Educational intervention	Behavioural intervention				
Change in weight z score Follow-up: 6 months	The mean change in weight z score in the educational intervention group was 0.06	The mean change in weight z score in the intervention groups was 0.06 higher (0.1 lower to 0.22 higher)		78 (Powers 2015)	⊕⊕⊕⊖ moderate ^{1,2}	
Change in weight z score Follow-up: 18 months	The mean change in weight z score in the educational intervention group was 0.11	The mean change in weight z score in the behavioural intervention groups was 0.04 higher (0.2 lower to		78 (Powers 2015)	⊕⊕⊕⊕ high ¹	

Comparison 5.2 Behavioural intervention versus education and attention control treatment						
Change in height z score Follow-up: 18 months	The mean change in weight z score in the educational intervention group was -0.02	The mean change in height z score in the behavioural intervention groups was 0.11 higher (0.02 lower to 0.24 higher)		78 (Powers 2015)	⊕⊕⊕⊖ moderate ^{1,2}	
Change in FEV ₁	No evidence was found					
Quality of life	No evidence was found					
Pulmonary exacerbations	No evidence was found					
Adverse effects (digestive system) Follow-up: 6 months	Study population		RR 1.61 (1.14 to 2.27)	78 (Powers 2015)	⊕⊕⊕⊖ moderate ^{1,2}	
	500 per 1000	805 per 1000 (570 to 1000)				
	Moderate					
	500 per 1000	805 per 1000 (570 to 1000)				
Patient or carer satisfaction	No evidence was found					
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; MD: mean difference						

- 1 The quality of the evidence was not downgraded although there was unclear risk of bias in relation to allocation concealment and blinding, because objective measures are unlikely to be influenced by the lack of blinding.
2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

4 **Table 150: Summary clinical evidence profile: Comparison 5.3. Behavioural**
5 **management training + educational intervention versus educational**
6 **intervention alone**

Comparison 5.3. Behavioural management training + educational intervention versus educational intervention alone						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Educational intervention alone	Behavioural management training + nutritional intervention				
Change in weight (kg) Follow-up: 2 months	The mean change in weight (kg) in the control group was 0.92	The mean change in weight (kg) in the behavioural management training +		67 (Stark 2009)	⊕⊕⊕⊖ moderate ¹	

Comparison 5.3. Behavioural management training + educational intervention versus educational intervention alone						
		nutritional intervention groups was 0.55 higher (0 to 1.1 higher)				
Change in weight (kg) Follow-up: 1 year	The mean change in weight (kg) in the educational intervention group was 1.75	The mean change in weight (kg) in the behavioural management training + nutritional intervention groups was 0.43 lower (1.27 lower to 0.41 higher)		8 (Powers 2003)	⊕⊕⊕⊕ very low ^{2,3}	
Change in weight (kg) Follow-up: 2 years	The mean change in weight (kg) in the educational intervention group was 6.45	The mean change in weight (kg) in the behavioural management training + nutritional intervention groups was 0.52 higher (1.34 lower to 2.38 higher)		59 (Stark 2009)	⊕⊕⊕⊕ moderate ¹	
Change in BMI z score Follow-up: 2 months	The mean change in BMI z score in the educational intervention group was 0.18	The mean change in BMI z score in the behavioural management training + nutritional intervention groups was 0.2 higher (0.02 lower to 0.42 higher)		67 (Stark 2009)	⊕⊕⊕⊕ moderate ¹	
Change in BMI z score Follow-up: 2 years	The mean change in BMI z score in the educational intervention group was -0.22	The mean change in BMI z score in the behavioural management training + nutritional intervention groups was 0.35 higher (0 to 0.7 higher)		59 (Stark 2009)	⊕⊕⊕⊕ moderate ¹	
Change in %	The mean	The mean		7	⊕⊕⊕⊕	

Comparison 5.3. Behavioural management training + educational intervention versus educational intervention alone						
ideal body weight Follow-up: 1 year	change in % ideal body weight in the educational intervention group was 9.4	change in % ideal body weight in the behavioural management training + nutritional intervention groups was 0.91 lower (37.52 lower to 35.7 higher)		(Powers 2003)	very low ^{2,3}	
Change in weight % for age Follow-up: 1 year	The mean change in weight % for age in the educational intervention group was 4.8	The mean change in weight % for age in the behavioural management training + nutritional intervention groups was 0.6 lower (17.25 lower to 16.05 higher)		8 (Powers 2003)	⊕⊕⊕⊕ very low ^{2,3}	
Change in height (cm) Follow-up: 1 year	The mean change in height (cm) in the educational intervention group was 7.13	The mean change in height (cm) in the behavioural management training + nutritional intervention groups was 2.03 lower (4.87 lower to 0.81 higher)		(Powers 2003)	⊕⊕⊕⊕ very low ^{2,3}	
Change in height (cm) Follow-up: 2 years	The mean change in height (cm) in the educational intervention group was 13.54	The mean change in height (cm) in the behavioural management training + nutritional intervention groups was 0.2 lower (1.45 lower to 1.05 higher)		59 (Stark 2009)	⊕⊕⊕⊕ high	
Change in height z score Follow-up: 2 years	The mean change in height z score in the educational intervention group was 0.04	The mean change in height z score in the behavioural management training + nutritional intervention groups was 0.01 lower		59 (Stark 2009)	⊕⊕⊕⊕ moderate ¹	

Comparison 5.3. Behavioural management training + educational intervention versus educational intervention alone					
		(0.17 lower to 0.15 higher)			
Change in FEV ₁ % Scale from: 0 to 100 Follow-up: 2 years	The mean change in FEV ₁ in the educational intervention group was -5	The mean change in FEV ₁ in the behavioural management training + nutritional intervention groups was 5.16 higher (8.49 lower to 18.81 higher)		28 (Stark 2009)	⊕⊕⊕⊖ low ⁴
Quality of life	No evidence was found				
Adverse effects	No evidence was found				
Time to next exacerbation	No evidence was found				
Patient or carer satisfaction	Parents in both groups reported high ratings of satisfaction with treatment (>6 in a 7 point scale)			67 (Stark 2009)	⊕⊕⊕⊖ moderate ⁵
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: BMI: body mass index; CI: confidence interval; FEV ₁ : forced expiratory volume in 1 second; kg: kilogrammes; cm: centimetres; MD: mean difference					

- 1 *1 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID*
- 2 *2 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to random*
- 3 *sequence generation, allocation concealment and incomplete outcome data. Cochrane rated the risk of bias in*
- 4 *relation to blinding as high risk however objective measures are unlikely to be influenced by a lack of blinding.*
- 5 *3 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs*
- 6 *4 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs*
- 7 *5 The quality of the evidence was downgraded by 1 due to bad reporting (narrative reporting only)*

8 10.1.5 Economic evidence

9 No economic evaluations of nutritional interventions were identified in the literature search
10 conducted for this guideline. Full details of the search and economic article selection flow
11 chart can be found in Appendix E and F, respectively.

12 This review question was not prioritised for de novo economic modelling. To aid
13 consideration of cost-effectiveness relevant resource and cost use data are presented in
14 Appendix K.

15 10.1.6 Evidence statements

16 10.1.6.1 Oral calorie supplementation

17 10.1.6.1.1 Comparison 1.1. Oral calorie supplementation versus usual care

18 Indices of nutrition and growth: weight

19 Moderate quality evidence from 1 RCT with 99 children and young people with cystic fibrosis
20 2 to 15 years old showed no clinically significant difference in weight (measured as change in

1 kg and change in weight centile) in the group of participants receiving oral calorie
2 supplementation compared to the participants in the control group receiving usual care at 3
3 months follow-up.

4 Similarly, low quality evidence from 2 RCTs with 117 children and young people with cystic
5 fibrosis 2 to 15 years old showed no clinically significant difference in weight (measured as
6 change in kg) in the group of participants receiving oral calorie supplementation compared to
7 the participants in the control group receiving usual care at 6 months follow-up.

8 Similarly, moderate quality evidence from 1 RCT with 101 children and young people with
9 cystic fibrosis 2 to 15 years old showed no clinically significant difference in weight
10 (measured as change in weight centile) in the group of participants receiving oral calorie
11 supplementation compared to the participants in the control group receiving usual care at 6
12 months follow-up.

13 Very low quality evidence from 1 RCT with 16 children and young people with cystic fibrosis
14 7 to 15 years old showed no clinically significant difference in weight (measured as change in
15 percentage expected for age and height) in the group of participants receiving oral calorie
16 supplementation compared to the participants in the control group receiving usual care at 6
17 months follow-up.

18 Likewise, moderate quality evidence from 1 RCT with 102 children and young people with
19 cystic fibrosis 2 to 15 years old showed no clinically significant difference in weight
20 (measured as change in kg and change in weight centile) in the group of participants
21 receiving oral calorie supplementation compared to the participants in the control group
22 receiving usual care at 1 year follow-up.

23 **Indices of nutrition and growth: BMI**

24 Moderate quality evidence from 1 RCT with 99 children and young people with cystic fibrosis
25 2 to 15 years old showed no clinically significant difference in BMI (measured as change in
26 kg/m² and as change in BMI centile) in the group of participants receiving oral calorie
27 supplementation compared to the participants in the control group receiving usual care at 3
28 months follow-up.

29 Similarly, moderate quality evidence from 1 RCT with 101 children and young people with
30 cystic fibrosis 2 to 15 years old showed no clinically significant difference in BMI (measured
31 as change in kg/m² and as change in BMI centile) in the group of participants receiving oral
32 calorie supplementation compared to the participants in the control group receiving usual
33 care at 6 months follow-up.

34 Likewise, moderate quality evidence from 1 RCT with 102 children and young people with
35 cystic fibrosis 2 to 15 years old showed no clinically significant difference in BMI (measured
36 as change in kg/m² and as change in BMI centile) in the group of participants receiving oral
37 calorie supplementation compared to the participants in the control group receiving usual
38 care at 1 year follow-up.

39 **Indices of nutrition and growth: height**

40 High to moderate quality evidence from 1 RCT with 99 children and young people with cystic
41 fibrosis 2 to 15 years old showed no clinically significant difference in height (measured as
42 change in cm and as change in height centile) in the group of participants receiving oral
43 calorie supplementation compared to the participants in the control group receiving usual
44 care at 3 months follow-up.

45 Likewise, high quality evidence from 1 RCT with 101 children and young people with cystic
46 fibrosis 2 to 15 years old showed no clinically significant difference in height (measured as
47 change in cm and change in height centile) in the group of participants receiving oral calorie

1 supplementation compared to the participants in the control group receiving usual care at 6
2 months follow-up.

3 Very low quality evidence from 1 RCT with 16 children and young people with cystic fibrosis
4 7 to 15 years old showed no clinically significant difference in height (measured as change in
5 percentage expected for age) in the group of participants receiving oral calorie
6 supplementation compared to the participants in the control group receiving usual care at 6
7 months follow-up.

8 High to moderate quality evidence from 1 RCT with 102 children and young people with
9 cystic fibrosis 2 to 15 years old showed no clinically significant difference in height
10 (measured as change in cm and change in height centile) in the group of participants
11 receiving oral calorie supplementation compared to the participants in the control group
12 receiving usual care at 1 year follow-up.

13 **Lung function: FEV₁**

14 Moderate quality evidence from 1 RCT with 69 children and young people with cystic fibrosis
15 2 to 15 years old showed a clinically significant decrease in lung function (measured as
16 change in FEV₁ % predicted) in the group of participants receiving oral calorie
17 supplementation compared to the participants in the control group receiving usual care at 3
18 months follow-up.

19 However, low quality evidence from 2 RCTs with 86 children and young people with cystic
20 fibrosis 2 to 15 years old showed no clinically significant change in lung function (measured
21 as change in FEV₁ % predicted) in the group of participants receiving oral calorie
22 supplementation compared to the participants in the control group receiving usual care at 6
23 months follow-up.

24 Similarly, moderate quality evidence from 1 RCT with 70 children and young people with
25 cystic fibrosis 2 to 15 years old showed no clinically significant change in lung function
26 (measured as change in FEV₁ % predicted) in the group of participants receiving oral calorie
27 supplementation compared to the participants in the control group receiving usual care at 1
28 year follow-up.

29 **Quality of life**

30 No evidence was found for this critical outcome.

31 **Pulmonary exacerbations**

32 No evidence was found for this important outcome.

33 **Adverse effects**

34 No evidence was found for this important outcome.

35 **Patient or carer satisfaction**

36 No evidence was found for this important outcome.

37 **370.1.6.1.2 Comparison 1.2. Oral calorie supplementation versus nutritional advice**

38 **Indices of nutrition and growth: weight**

39 Very low quality evidence from 1 quasi-RCT with 13 people with CF >10 years old showed
40 no clinically significant difference in weight (measured as change in kg, change in Z-score,
41 change in percentage weight for height, change in percentage of ideal body weight) between
42 the group of participants receiving oral calorie supplementation for 3 months and the
43 participants receiving nutritional advice at 3 months follow-up.

1 Very low quality evidence from 1 quasi-RCT with 13 people with CF >10 years old showed
2 no clinically significant difference in weight (measured as change in z score and change in
3 percentage of ideal body weight) between the group of participants receiving oral calorie
4 supplementation for 3 months and the participants receiving nutritional advice at 6 months
5 follow-up.

6 **Indices of nutrition and growth: height**

7 Very low quality evidence from 1 quasi-RCT with 13 people with cystic fibrosis >10 years old
8 showed no clinically significant difference in height (measured as change in cm and change
9 in Z-score) between the group of participants receiving oral calorie supplementation for 3
10 months and the participants receiving nutritional advice at 3 months follow-up.

11 Very low quality evidence from 1 quasi-RCT with 13 people with cystic fibrosis >10 years old
12 showed no clinically significant difference in height (measured as change in Z-score)
13 between the group of participants receiving oral calorie supplementation for 3 months and
14 the participants receiving nutritional advice at 6 months follow-up.

15 **Lung function: FEV₁**

16 Very low quality evidence from 1 quasi-RCT with 13 people with cystic fibrosis >10 years old
17 showed a clinically significant decrease in lung function (measured as change in FEV₁ %
18 predicted) in the group of participants receiving oral calorie supplementation for 3 months
19 compared to the participants receiving nutritional advice at 3 and at 6 months follow-up.

20 **Quality of life**

21 No evidence was found for this critical outcome.

22 **Pulmonary exacerbations**

23 No evidence was found for this important outcome.

24 **Adverse effects**

25 No evidence was found for this important outcome.

26 **Patient or carer satisfaction**

27 No evidence was found for this important outcome.

28 **10.1.6.2 Enteral tube feeding**

29 **10.1.6.2.1 Comparison 2. Enteral tube feeding versus usual care**

30 **Indices of nutrition and growth: weight**

31 Very low quality evidence from 1 cohort study with 21 adults with cystic fibrosis showed a
32 clinically significant improvement in weight (measured as change in kg) in the group
33 receiving enteral tube feeding compared to those who received usual care at 1,2 and 3 years
34 follow-up.

35 Very low quality evidence from 1 cohort study with 40 people with cystic fibrosis aged 2 to 20
36 years showed a clinically significant improvement in weight (measured as change in z score)
37 in the group receiving gastrostomy compared to those who received usual care at 6 months
38 and 1 year follow-up.

39 **Indices of nutrition and growth: height**

40 Very low quality evidence from 1 cohort study with 40 people with cystic fibrosis aged 2 to 20
41 years showed no clinically significant increase in height (measured as change in z score)

1 between the group receiving gastrostomy and those who received usual care at 6 months
2 and 1 year follow-up.

3 **Indices of nutrition and growth: BMI**

4 Very low quality evidence from 1 cohort study with 21 adults with cystic fibrosis showed a
5 clinically significant improvement in BMI (measured as change in kg/m²) in the group
6 receiving enteral tube feeding compared to those who received usual care at 1,2 and 3 years
7 follow-up.

8 Very low quality evidence from 1 cohort study with 40 people with cystic fibrosis aged 2 to 20
9 years showed a clinically significant improvement in BMI (measured as change in z score) in
10 the group receiving gastrostomy compared to those who received usual care at 6 months
11 and 1 year follow-up.

12 **Lung function: FEV₁**

13 Very low quality evidence from 1 cohort study with 21 adults with cystic fibrosis showed no
14 clinically significant improvement in lung function (measured as change in FEV₁ % predicted)
15 between the group receiving enteral tube feeding and those who received usual care at 1,2
16 and 3 years follow-up.

17 Very low quality evidence from 1 cohort study with 27 people with cystic fibrosis aged 2 to 20
18 years showed no clinically significant difference in lung function (measured as change in
19 FEV₁ % predicted) between the group receiving gastrostomy and those who received usual
20 care at 6 months and 1 year follow-up.

21 **Quality of life**

22 No evidence was found for this critical outcome.

23 **Days on IV treatment (proxy outcome for pulmonary exacerbations)**

24 Very low quality evidence from 1 cohort study with 21 adults with cystic fibrosis showed no
25 clinically significant improvement in IV treatment (measured as change in number of days of
26 treatment) between the group receiving enteral tube feeding and those who received usual
27 care at 1 and 3 years follow-up. However, the same evidence found a clinically significant
28 increase in IV treatment (measured as change in number of days of treatment) in the group
29 receiving enteral tube feeding compared to those who received usual care at 2 years follow-
30 up.

31 **Adverse effects**

32 No evidence was found for this important outcome.

33 **Patient or carer satisfaction**

34 No evidence was found for this important outcome.

35 **10.1.6.3 Appetite stimulants**

36 **10.1.6.3.1 Comparison 3. Appetite stimulants versus placebo**

37 **Indices of nutrition and growth: weight**

38 Low quality evidence from 2 RCTs with 33 people with cystic fibrosis aged ≥5 years showed
39 no clinically significant difference in weight (measured as change in kg) between the group of
40 participants receiving an appetite stimulant and the group receiving placebo at 3 months
41 follow-up.

1 Low quality evidence from 3 RCTs with 40 people with cystic fibrosis aged ≥ 21 months
2 showed a clinically significant improvement in weight (measured as change in weight z
3 score) in the group of participants receiving an appetite stimulant as opposed to the group
4 receiving placebo at 3 months follow-up.

5 Very low quality evidence from 1 RCT with 16 people with cystic fibrosis aged ≥ 5 years
6 showed a clinically significant improvement in weight (measured as change in % ideal body
7 weight) in the group of participants receiving an appetite stimulant compared to the group
8 receiving placebo at 3 months follow-up.

9 Low quality evidence from 1 RCT with 17 people with cystic fibrosis older than 6 years
10 showed a clinically significant improvement in weight (measured as change in kg and as
11 change in weight z score) in the group of participants receiving an appetite stimulant as
12 opposed to the group receiving placebo at 6 months follow-up.

13 **Indices of nutrition and growth: height**

14 Very low quality evidence from 1 RCT with 16 people with cystic fibrosis aged ≥ 5 years
15 showed no clinically significant difference in height (measured as change in cm) between the
16 group of participants receiving an appetite stimulant and the group receiving placebo at 3
17 months follow-up.

18 **Indices of nutrition and growth: BMI**

19 Very low quality evidence from 1 RCT with 16 people with cystic fibrosis aged ≥ 5 years
20 showed no clinically significant difference in BMI (measured as change in kg/m²) between
21 the group of participants receiving an appetite stimulant and the group receiving placebo at 3
22 months follow-up.

23 However, very low quality evidence from 1 RCT with 16 people with cystic fibrosis aged ≥ 5
24 years showed a clinically significant improvement in BMI (measured as change in percentile)
25 in the group of participants receiving an appetite stimulant compared to the group receiving
26 placebo at 3 months follow-up.

27 **Lung function: FEV₁**

28 Very low quality evidence from 1 RCT with 17 people with cystic fibrosis older than 6 years
29 showed no clinically significant difference in lung function (measured as change in FEV₁ %
30 predicted) between the group of participants receiving an appetite stimulant and the group
31 receiving placebo at 3 and 6 months follow-up.

32 **Pulmonary exacerbations**

33 Very low quality evidence from 1 RCT with 12 children with cystic fibrosis aged 21 months to
34 10.4 years showed no clinically significant difference in the number of pulmonary
35 exacerbations between the group of participants receiving an appetite stimulant and the
36 group receiving placebo at 3 months follow-up.

37 **Adverse effects**

38 Very low quality evidence from 1 RCT with 17 people with cystic fibrosis older than 6 years
39 showed no clinically significant difference in the number of constipation events between the
40 group of participants receiving an appetite stimulant and the group receiving placebo at 6
41 months follow-up.

42 Low quality evidence from 1 RCT with 12 children with cystic fibrosis aged 21 months to 10.4
43 years showed that fasting blood glucose levels remained unchanged in both the group of
44 participants receiving an appetite stimulant and the group receiving placebo at 3 months
45 follow-up. Imprecision and clinical significance could not be calculated.

1 Low quality evidence from 1 RCT with 12 children aged 21 months to 10.4 years with cystic
2 fibrosis showed that all participants in the group receiving an appetite stimulant had normal
3 cortisol levels at baseline; at follow-up 4 out of the 6 participants in this same group had
4 morning cortisol levels decreased to <0.6 mcg/dl. The values for the control group were not
5 reported. Imprecision and clinical significance could not be calculated.

6 Very low quality evidence from 1 RCT with 17 people with cystic fibrosis older than 6 years
7 showed no clinically significant difference in the number of people with decreased morning
8 cortisol levels <30nmol/L between the group of participants receiving an appetite stimulant
9 and the group receiving placebo at 6 months follow-up. The study described that 7 out of 10
10 people receiving an appetite stimulant had decreased morning cortisol levels <30nmol/L at 6
11 months follow-up, however after completing a 1-month-long wean from the appetite
12 stimulant, levels increased to 270 +-6.9 nmol/L.

13 **Quality of life**

14 No evidence was found for this critical outcome.

15 **Patient or carer satisfaction**

16 No evidence was found for this important outcome.

17 **10.1.6.4 Nutrition education/dietary advice**

18 **10.1.6.4.1 Comparison 4. Nutrition education versus standard treatment**

19 **Indices of nutrition and growth: weight**

20 Low to very low quality evidence from 1 RCT with 48 people with cystic fibrosis older than 16
21 years showed no clinically significant difference in weight (measured as change in kg) between
22 the group of participants receiving a nutrition education intervention for 10 weeks and those
23 who received usual care at 6 and 12 months follow-up.

24 **Lung function: FEV₁**

25 Very low quality evidence from 1 RCT with 48 people with cystic fibrosis older than 16 years
26 showed no clinically significant difference in lung function (measured as change in FEV₁ %
27 predicted) between the group of participants receiving a nutrition education intervention for
28 10 weeks and those who received usual care at 6 and 12 months follow-up.

29 **Quality of life**

30 Low quality evidence from 1 RCT with 48 people with cystic fibrosis older than 16 years
31 showed the following differences in health-related quality of life measured with the CF-QOL
32 questionnaire: the control group scored significantly higher in physical functioning than the
33 intervention group at 6 months (p-value=0.05) and the control group scored significantly
34 higher in concerns for the future (meaning it had less concerns) than the intervention group
35 at 12 months follow-up. This outcome was reported narratively only and imprecision could
36 not be calculated.

37 **Time to next exacerbation**

38 No evidence was found for this important outcome

39 **Adverse effects**

40 No evidence was found for this important outcome

41 **Patient and carer satisfaction**

42 No evidence was found for this important outcome

1 10.1.6.5 Psychological and behavioural interventions

20.1.6.5.1 Comparison 5.1 Behavioural intervention versus usual care

3 Indices of nutrition and growth: weight

4 Very low quality evidence from 1 RCT with 9 children with cystic fibrosis aged 5 to 10 years
5 showed no clinically significant difference in weight (measured as change in kg and change
6 in z score) between the group of participants receiving a group behavioural intervention and
7 those who received usual care at 6 weeks follow-up.

8 Indices of nutrition and growth: height

9 Very low quality evidence from 1 RCT with 9 children with cystic fibrosis aged 5 to 10 years
10 showed no clinically significant difference in height (measured as change in cm) between the
11 group of participants receiving a group behavioural intervention and those who received
12 usual care at 6 weeks follow-up.

13 Lung function: FEV₁

14 Very low quality evidence from 1 RCT with 9 children with cystic fibrosis aged 5 to 10 years
15 showed no clinically significant difference in lung function (measured as change in FEV₁ %
16 predicted) between the group of participants receiving a group behavioural intervention and
17 those who received usual care at 6 weeks follow-up.

18 Quality of life

19 No evidence was found for this critical outcome

20 Time to next exacerbation

21 No evidence was found for this important outcome

22 Adverse effects

23 No evidence was found for this important outcome

24 Patient and carer satisfaction

25 No evidence was found for this important outcome

26 20.1.6.5.2 Comparison 5.2. Behavioural intervention versus education and attention control 27 treatment

28 Indices of nutrition and growth: weight

29 Moderate to high quality evidence from 1 RCT with 78 children with cystic fibrosis aged 2 to 6
30 years showed no clinically significant difference in weight (measured as change in z score)
31 between the group of participants receiving a behavioural intervention and those who
32 received an education and attention control treatment at 6 and 18 months follow-up.

33 Indices of nutrition and growth: height

34 Moderate quality evidence from 1 RCT with 78 children with cystic fibrosis aged 2 to 6 years
35 showed no clinically significant difference in height (measured as change in z score) between
36 the group of participants receiving a behavioural intervention and those who received an
37 education and attention control treatment at 18 months follow-up.

38 Lung function: FEV₁

39 No evidence was found for this critical outcome

1 **Quality of life**

2 No evidence was found for this critical outcome

3 **Time to next exacerbation**

4 No evidence was found for this important outcome

5 **Adverse effects**

6 Moderate quality evidence from 1 RCT with 78 children with cystic fibrosis aged 2 to 6 years
7 showed a clinically significant higher number of adverse effects to the digestive system in the
8 group of participants receiving a behavioural intervention compared to those who received an
9 education and attention control treatment at 6 months follow-up.

10 **Patient and carer satisfaction**

11 No evidence was found for this important outcome

12 **10.1.6.5.3 Comparison 5.3. Behavioural management training + educational intervention versus**
13 **educational intervention alone**

14 **Indices of nutrition and growth: weight**

15 Moderate quality evidence from 1 RCT with 67 children and young people with cystic fibrosis
16 aged 4 to 12 years showed no clinically significant difference in weight (measured as change
17 in kg) between the group receiving a combination of a behavioural and educational
18 intervention for 2 months and those who received the educational intervention alone for 2
19 months at 2 months follow-up.

20 Very low quality evidence from 1 RCT with 8 infants and children less than 3 years old with
21 cystic fibrosis showed no clinically significant difference in weight (measured as change in
22 kg, as change in % ideal body weight at as change in % weight for age) between the group
23 receiving a combination of a behavioural and educational intervention for 1 year and those
24 who received the educational intervention alone for 1 year at 1 year follow-up.

25 Moderate quality evidence from 1 RCT with 59 children and young people with cystic fibrosis
26 aged 4 to 12 years showed no clinically significant difference in weight (measured as change
27 in kg) between the group receiving a combination of a behavioural and educational
28 intervention for 2 months and those who received the educational intervention alone for 2
29 months at 2 years follow-up.

30 **Indices of nutrition and growth: BMI**

31 Moderate quality evidence from 1 RCT with 67 children and young people with cystic fibrosis
32 aged 4 to 12 years showed no clinically significant difference in BMI (measured as change in
33 z score) between the group receiving a combination of a behavioural and educational
34 intervention for 2 months and those who received the educational intervention alone for 2
35 months at 2 months follow-up.

36 Moderate quality evidence from 1 RCT with 59 children and young people with cystic fibrosis
37 aged 4 to 12 years showed no clinically significant difference in BMI (measured as change in
38 z score) between the group receiving a combination of a behavioural and educational
39 intervention for 2 months and those who received the educational intervention alone for 2
40 months at 2 years follow-up.

41 **Indices of nutrition and growth: height**

42 Very low quality evidence from 1 RCT with 7 infants and children less than 3 years old with
43 cystic fibrosis showed no clinically significant difference in height (measured as change in
44 cm) between the group receiving a combination of a behavioural and educational intervention

1 for 1 year and those who received the educational intervention alone for 1 year at 1 year
2 follow-up.

3 High to moderate quality evidence from 1 RCT with 59 children and young people with cystic
4 fibrosis aged 4 to 12 years showed no clinically significant difference in height (measured as
5 change in cm and change in z score) between the group receiving a combination of a
6 behavioural and educational intervention for 2 months and those who received the
7 educational intervention alone for 2 months at 2 years follow-up.

8 **Lung function: FEV₁**

9 Low quality evidence from 1 RCT with 28 children and young people with cystic fibrosis aged
10 4 to 12 years showed no clinically significant difference in lung function (measured as
11 change in FEV₁ % predicted) between the group of participants receiving a combination of a
12 behavioural and educational intervention for 2 months and those who received the
13 educational intervention alone for 2 months at 2 years follow-up.

14 **Quality of life**

15 No evidence was found for this critical outcome

16 **Time to next exacerbation**

17 No evidence was found for this important outcome

18 **Adverse effects**

19 No evidence was found for this important outcome

20 **Patient and carer satisfaction**

21 Moderate quality evidence from 1 RCT with 67 children and young people with cystic fibrosis
22 aged 4 to 12 years showed that parents in both groups (the group receiving a combination of
23 a behavioural and educational intervention for 2 months and the group receiving the
24 educational intervention only for 2 months) reported high ratings of satisfaction with
25 treatment (>6 in a 7 point scale) at 2 months follow-up. The outcome was reported narratively
26 only and the imprecision could not be assessed.

27 **10.1.6.6 Economic evidence statements**

28 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

29 **10.1.7 Evidence to recommendations**

30 **10.1.7.1 Relative value placed on the outcomes considered**

31 The aim of this review was to determine the clinical and cost effectiveness of nutritional
32 interventions in improving health outcomes for people with cystic fibrosis.

33 The Committee chose change in weight or height or body mass index (BMI) or z score or
34 other indices of nutrition or growth, lung function (FEV₁) and quality of life as critical
35 outcomes for decision making; whereas changes to body composition detected by
36 anthropometric measure, pulmonary exacerbations, patient and parent or carer satisfaction
37 and adverse effects (including diarrhoea, reduced appetite, abdominal bloating, episodes of
38 distal intestinal obstruction syndrome) were rated as important. For appetite stimulants, the
39 following adverse effects were also rated as important outcomes: high blood glucose and
40 adrenal insufficiency.

1 10.1.7.2 Consideration of clinical benefits and harms

2 People with cystic fibrosis often suffer from undernutrition due to faecal fat loss, increased
3 energy requirements caused by chronic infections and malabsorption due to pancreatic
4 insufficiency. It is well established that nutrition is important for lung function and overall
5 health, therefore, different nutritional interventions to improve the nutritional status and
6 growth of people with cystic fibrosis should be considered. Because nutrition is such an
7 important component of overall health and a considerable problem among people with cystic
8 fibrosis, the Committee agreed that dietitians should be an integral part of the
9 multidisciplinary team caring for the person with cystic fibrosis and review the patient
10 regularly. This should be from an individualised basis, considering a myriad of factors,
11 including current diet, bowel habit in relation to pancreatic enzyme use as well as family
12 circumstances, needs and capabilities before recommending any nutritional intervention.

13 If there are nutrition concerns, the Committee recommended based on their clinical
14 experience and expertise to encourage people to increase portion size and eat high-energy
15 foods in order to increase calorie intake and counterbalance increased energy requirements
16 and malabsorption.

17 After reviewing the evidence on the effectiveness of nutritional interventions, the Committee
18 concluded that enteral tube feeding and appetite stimulants can improve nutritional status
19 and growth, but the available evidence does not show routine oral calorie supplementation to
20 be effective.

21 Despite this, considering the invasiveness of some nutritional interventions the Committee
22 agreed that the first treatment to follow additional nutritional advice should be a trial of oral
23 calorie supplementation. If no improvement is achieved, either enteral tube feeding or for
24 adults, appetite stimulants, should be considered.

25 Enteral tube feeding is shown to be effective in improving nutrition and growth in people with
26 cystic fibrosis. The Committee agreed that the capacity and the capabilities of the patient and
27 the family should always be carefully considered before embarking on this.

28 The Committee looked at appetite stimulants as an alternative to enteral tube feeding. The
29 evidence shows that they are effective but the Committee discussed whether they can have
30 adverse effects such as hyperglycaemia and adrenal insufficiency. There was limited
31 evidence available on adverse effects; this evidence showed no clinically significant
32 difference in constipation and no difference in fasting blood glucose levels between
33 participants receiving appetite stimulants and those receiving placebo. According to the
34 evidence, some participants had decreased morning cortisol levels after receiving appetite
35 stimulants, however in one study values in the control group were not reported, while in the
36 other study there was no clinically significant difference with the control group, and values
37 increased after the intervention group stopped receiving appetite stimulants. The Committee
38 discussed that although many people with cystic fibrosis considering appetite stimulants
39 might already have diabetes, and in their clinical experience, adrenal insufficiency is not very
40 often observed, they agreed to recommend them only in adults, short-term and after all other
41 options had been fully explored. Moreover, possible adverse effects should be explained so
42 that an informed decision can be made.

43 The decision about these treatments should be based on the whole clinical picture as well as
44 the patient's preferences and capabilities.

45 Because the evidence showed that oral calorie supplements are not effective in improving
46 nutrition or growth in people in cystic fibrosis, the Committee agreed not to recommend them
47 as a routine intervention for the general population of people with cystic fibrosis. They
48 discussed whether to recommend them if there are nutrition concerns. They noted that out of
49 three studies on oral nutritional supplements, the population in two studies (Hanning 1993
50 and Kalnins 2005) was small (between 15 and 20 participants) and did not represent the

1 population that dietitians would actually consider offering nutrition interventions to, because
2 inclusion criteria were either unclear (Hanning 1993) or used relatively high thresholds for
3 weight (Kalnins 2005) to define the study populations. Only one study (Poustie 2006, 102
4 participants) showed no effectiveness of oral nutritional supplements in a population defined
5 by inclusion criteria that were similar to the thresholds for additional nutritional support
6 outlined in the CF Trust consensus document on nutritional management of cystic fibrosis.
7 The Committee agreed that supplements if effective would be preferable from a patient's
8 perspective to enteral tube feeding, which is an invasive technique, or to appetite stimulant
9 drugs which may be associated with adverse effects. Therefore, based on their clinical
10 experience and expertise they agreed that oral nutritional supplements should be considered
11 on a trial basis for people requiring additional nutrition who had not responded to dietary
12 advice, before considering more invasive interventions.

13 10.1.7.3 Consideration of economic benefits and harms

14 The Committee advised that oral supplements should not be routinely offered to all people
15 with cystic fibrosis, as dietary modifications are at least as effective and do not take away
16 from NHS resources. Moreover, the cost of oral supplementation could be substantial over
17 the longer term, especially if they are used to substitute rather than complement a healthy
18 diet. A single measure would cost approximately £4.36 (BNF August 2016, Calogen®
19 emulsion, 200ml), but the specific brand would depend on the person's deficiencies which
20 could require a more expensive preparation.

21 However, the Committee agreed that oral supplements should be used to complement a
22 person's diet for acute use, during periods of ill health, such as an exacerbation, when they
23 are nutritionally unwell and have the scope to benefit from oral supplementation when their
24 normal diet is insufficient. Following this, the Committee agreed that a research
25 recommendation to assess the clinical and cost-effectiveness of oral supplementation in
26 people with cystic fibrosis who are nutritionally unwell, would assess if the benefits from the
27 acute use of oral supplementation can justify the costs to reduce current uncertainty in this
28 area. However, a research recommendation was not prioritised, given that the findings would
29 not contradict a recommendation to consider a trial of oral nutritional supplements if dietary
30 modifications are not effective.

31 Conversely, in the longer term, the Committee stated that optimal nutrition can prevent a
32 decline in lung function, iterating that the studies included in the clinical evidence review
33 were too short to demonstrate the differences seen in clinical practice over several years. As
34 a result, longer term studies would be needed to justify the additional cost of interventions
35 compared to usual care when the aim is to maintain lung function.

36 The high upfront cost of tube feeding and the initial monitoring schedule was recognised by
37 the Committee, and they acknowledged that tube feeding is associated with adverse effects
38 (incurring a treatment cost and disutility) and can negatively impact social interactions during
39 meal times. Despite this, the Committee considered a role for tube feeding in the event of
40 failure of efficacy or intolerance of alternative interventions, where the benefits could
41 outweigh the costs. The Committee also added that the most appropriate type of tube
42 feeding would be determined through a discussion with the dietician, the person with cystic
43 fibrosis and their family, to ensure their quality of life is maximised.

44 The Committee advised that appetite stimulants are associated with severe side effects such
45 as high blood glucose and adrenaline insufficiency, and should not be offered as a first line
46 option as the expected cost to manage those complications could outweigh the benefits from
47 an increased appetite. Moreover, the cost of appetite stimulants could soon overtake the cost
48 of tube feeding when they are prescribed on a long-term basis (BNF NHS Drug Tariff price,
49 August 2016; cyproheptadine hydrochloride 4mg 4 times daily, £24.28/month; megace
50 480mg/day, £59.34/month).

1 Overall, the Committee agreed that appropriate dietary modifications should be considered,
2 before initiating oral supplements, tube feeding or appetite stimulants; hence, options
3 associated with the least cost and resource use would be considered first. The Committee
4 were reluctant to specify the second line intervention in their recommendations as this would
5 depend on the capabilities and preferences of the person with cystic fibrosis and their family
6 or carers. The Committee also added that there would need to be a decline in health, or
7 faltering growth, before an intervention more costly and invasive than advice is considered.

8 **10.1.7.4 Quality of evidence**

9 The quality of the evidence presented in this review range from very low to high as assessed
10 by GRADE.

11 For the domain risk of bias, the studies were assigned the same risk of bias as in the
12 Cochrane reviews, and were not individually reviewed. The main biases that lead to
13 downgrading the quality of the evidence included randomization, allocation concealment, and
14 reporting bias.

15 Sample sizes of the studies are relatively small and statistical power might therefore be too
16 low to show an effect. Another problem with the studies are that the intervention time and the
17 follow-up time are not necessarily adequate to detect an effect in some outcomes (for
18 example change in weight).

19 No serious issues were found regarding inconsistency (heterogeneity), as most comparisons
20 included only 1 study.

21 Some issues regarding indirectness were also identified. The Committee discussed that the
22 participants in some of the studies on nutrition interventions among people with cystic fibrosis
23 do not represent the population that dietitians would actually consider offering nutrition
24 interventions to, instead the studies might include all patients at a cystic fibrosis clinic, not
25 just the ones with faltering growth or undernutrition.

26 **10.1.7.5 Other considerations**

27 The Committee agreed that studies with longer follow-up are needed (over 18 months), as
28 the aim of nutritional interventions is to prevent lung function from deterioration in the long
29 term, rather than improving in the short term.

30 They also discussed that a possible explanation for interventions not proving to be effective
31 is the lack of adherence.

32 No equality issues were identified by the Committee for this review question.

33 The Committee agreed that a research recommendation was not a priority in this area,
34 although it was noted that there was a lack of studies about nutritional interventions and
35 children with faltering growth.

36 **10.1.7.6 Key conclusions**

37 The Committee concluded that a dietitian should be an integral part of the multidisciplinary
38 team caring for the person with cystic fibrosis. The evidence showed that enteral tube
39 feeding and appetite stimulants are effective in improving nutritional status and growth in
40 people with cystic fibrosis. However, because of the invasive nature of enteral tube feeding
41 and the concern for potential adverse effects of appetite stimulants, dietary modifications
42 through nutritional advice should always be considered as the first choice of treatment for a
43 person with cystic fibrosis with undernutrition or faltering growth. The Committee did not
44 recommend routine use of oral calorie supplements for people with cystic fibrosis because no
45 evidence was found to justify this. However they did recommend that if there are signs that

1 raise nutrition concerns, a trial of oral supplementation should be considered before
2 proceeding to more invasive approaches.

3 **10.1.8 Recommendations**

4 **95. The cystic fibrosis specialist dietitian should offer advice on the benefits of**
5 **optimal nutrition, and at the annual assessment, review the person's:**

- 6 • total nutritional intake, including energy intake (calories)
- 7 • estimated nutritional needs
- 8 • pancreatic exocrine replacement therapy, if appropriate (see Exocrine
9 pancreatic insufficiency).

10 **96. Encourage people to increase calorie intake by increasing portion size and eating**
11 **high-energy foods, if there is concern about their nutrition (including weight loss**
12 **and inadequate weight gain).**

13 **97. If increased portion size and high-energy foods are not effective, consider a trial**
14 **of oral nutritional supplements.**

15 **98. If attempts to increase calorie intake are not effective, consider:**

- 16 • supplementation with enteral tube feeding, or
- 17 • for adults, a short-term trial of an appetite stimulant.

18 **10.2 Exocrine pancreatic insufficiency**

19 **Review question: In people with cystic fibrosis, what is the most effective regimen of**
20 **pancreatic enzyme replacement therapy (PERT) in the treatment of exocrine**
21 **pancreatic insufficiency?**

22 **10.2.1 Introduction**

23 Cystic Fibrosis causes a number of gastrointestinal complications. Exocrine pancreatic
24 insufficiency (PI) is a common gastrointestinal complication which affects people with cystic
25 fibrosis.

26 Pancreatic insufficiency is caused by a progressive fibrotic process that begins in-utero.
27 Pancreatic cells are damaged by deposits of dehydrated pancreatic secretions and replaced
28 with fibrous scar tissue. The pancreas no longer functions effectively, and produces reducing
29 amounts of enzymes essential for digestion. Pancreatic exocrine insufficiency is the major
30 cause of maldigestion of dietary macronutrients including fat, protein and carbohydrate.
31 Untreated it will result in sub-optimal nutritional status, impaired growth and development,
32 deficiency of fat soluble vitamins, and symptoms of malabsorption such as steatorrhoea.

33 Pancreatic exocrine insufficiency is treated by the administration of pancreatic enzyme
34 replacement therapy (PERT) with all fat and protein containing meals, snacks and drinks.
35 PERT preparations contain variable concentrations of lipase, protease and amylase. The
36 preparation chosen is dependent on clinical and individual requirements. The aim of PERT is
37 to ensure adequate digestion of all nutrients thereby controlling the symptoms of
38 malabsorption and ensuring growth, development and the maintenance of normal nutritional
39 and fat soluble vitamin status.

40 An understanding of the signs and symptoms of pancreatic insufficiency are crucial to ensure
41 its effective management with PERT. Pancreatic enzyme requirements vary at different
42 times, influenced by respiratory health, changing nutritional requirements throughout

1 childhood, and progressing pancreatic insufficiency. The purpose of this evidence review is
2 to determine the optimal PERT regimen to manage pancreatic exocrine insufficiency in
3 people with cystic fibrosis.

4 **10.2.2 Description of clinical evidence**

5 The objectives of this review were to evaluate:

- 6 • The effectiveness of PERT given with an acid neutralising or suppressing agent in
7 comparison with PERT alone
- 8 • The effectiveness of a high dose of PERT in comparison with a low dose of PERT

9 In the treatment of exocrine pancreatic insufficiency.

10 We looked for systematic reviews of randomised controlled trial (RCTs) and RCTs (including
11 cross-over trials).

12 For full details see review protocol in Appendix D.

13 One Cochrane Review was identified for this review question (Somaraju 2014) which
14 evaluated the efficacy and safety of pancreatic enzyme replacement therapy in children and
15 adults with cystic fibrosis. However, most comparisons of interest were different from those
16 stated in our evidence review protocol as the Cochrane looked at enteric coated PERT
17 compared to non-enteric coated PERT and enteric coated microspheres compared to tablets.
18 The Committee agreed that both non-enteric coated and tablets were not commonly used in
19 clinical practice in the UK. It also did not included studies with less than 28 days follow-up.
20 The individual relevant studies have been identified for potential inclusion in our review.

21 For comparison 1, the effectiveness and safety of adding an acid supressing agents (an H2
22 receptor antagonists or a protein pump inhibitor) to PERT therapy, 4 RCTs (Durie 1980,
23 Heijerman 1991, Heijerman 1993 and Francisco 2002) were included.

24 For comparison 2, the effectiveness and safety of high dose versus low dose PERT, 4 RCTs
25 (Heijerman 1991 – same as for comparison 1 -, Brady 1991, Mitchell 1982 and Beker 1994),
26 were included.

27 3 studies were conducted in the USA (Brady 1991, Beker 1994, Francisco 2002), 2 in
28 Holland (Heijerman 1991, Heijerman 1993), 1 in Canada (Durie 1990) and 1 in New Zealand
29 (Mitchel 1982).

30 With regards to the population, 4 studies were done in a children's population (Beker 1994,
31 Brady 1991, Durie 1980, Mitchell 1982) and 2 with adults (Heijerman 1991, Heijerman 1993).
32 One study included both adults and children population (Francisco 2002).

33 All studies had cross-over design. In all of them the patients were allocated to at least 2
34 different interventions, with a variable follow-up (from 5 to 28 days). In all studies a stool
35 collection was performed for the last 3 days of each treatment period and analysed for fat.

36 Of the outcomes listed in the protocol, all studies reported on the critical reduction of
37 steatorrhoea and faecal fat. All seven studies measured faecal fat excretion (FFE), either as
38 grams of fat per kg per 24 hours (Brady 1991), as grams of fat lost in the stool per 24 hours
39 (Brady 1991, Beker 1994, Durie 1980) or as fat excreted as a percentage of dietary fat intake
40 (Brady 1991, Durie 1980, Heijerman 1991, Heijerman 1993). Three studies measured fat
41 absorption as a percentage of intake (Beker 1994, Francisco 2002, Mitchell 1982) (the
42 coefficient of fat absorption, CFA). One study reported on resolution of symptoms of
43 malabsorption (Mitchel 1982) and 1 study reported side effects of the treatment (Brady
44 1994).

45 No results were found for weight or BMI, health-related quality of life and patient satisfaction

1 A summary of the included studies is included in Table 151. See also study selection flow
2 chart in Appendix F, study evidence tables in Appendix G, list of excluded studies in
3 Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

4 10.2.3 Summary of included studies

5 A summary of the studies that were included in this review are presented in Table 151.

6 **Table 151: Summary of included studies**

Study	Intervention/ comparison	Population	Outcomes	Comments
Beker 1994 (USA) Cross- over RCT	High-dose: 1500u lipase per kg/body for meals & 750u lipase per kg/body for snacks. Low-dose: 500u lipase per kg/body for meals & 250u lipase per kg/body for snacks. Additional information: • Daily fat intake (g): 100g in both groups.	N=21 children and young people with CF Age (SD, range): 11.5; 5 to 28 years	Reduction of steatorrhoea and faecal fat • Faecal fat excretion (FFE) (g/24 h) • Fat absorption (CFA) (%) of intake) Adverse event • Constipation • Elevations in serum uric acid levels	
Brady 1991 (USA) Cross- over RCT	High-dose: median 12 (8 to 18) capsules per meal Low-dose: median 3 (2 to 5) capsules per meal. Additional information: Constituent enzymes per capsule: 7.020u of lipase. Daily fat intake (g) 94±6 in both groups.	N=9 children with CF Age (median, range): 9; 6 to 10 years	Reduction of steatorrhoea and faecal fat • Faecal fat excretion (FFE) (% of intake) • Faecal fat excretion (FFE) (g/kg/24h) • Faecal fat excretion (FFE) (g/24h)	
Durie 1980 (Canada) Cross- over RCT	Pancrealipase alone: 26 Pancrease capsules (Cotazym) per day , 6 per meal and 3 per snack Pancrealipase + Cimetidine (200 & 300 mg. tablets)	N=21 children and young people with CF Age: 10 to 17 years	Reduction of steatorrhoea and faecal fat • Faecal fat excretion (FFE) (g/24h) • Faecal fat excretion (FFE) (% of intake)	
Francisco 2002 (USA) Cross- over RCT	Pancrease (MT10 or MT16) + adjunct low-dose or high-dose Ranitidine. Children weighting ≤40 kg were given ranitidine 5 mg/kg or 10 mg/kg daily, divided into 2 equal doses 30 minutes before breakfast and dinner. Children weighting >40 kg and adults received 150 mg or 300 mg twice daily. Pancrease (MT10 or	N=22 people with CF 12 children and young people, age: 6 to 17 years; 10 adults, age: 18 to 36 years	Reduction of steatorrhoea and faecal fat • Fat absorption (CFA) (%)	

Study	Intervention/ comparison	Population	Outcomes	Comments
	MT16) + adjunct Omeprazole (adults only) 20 mg daily, 30 minutes before breakfast Pancrease (MT10 or MT16) + placebo Additional information: diet fat was kept constant			
Heijerman 1991 (Netherlands) Cross- over RCT	High-dose 4 capsules x 3 times per day Low-dose 2 capsules x 3 times per day. Additional information: Constituent enzymes per capsule 5000u lipase, 2900u lipase, 330u protease. Fat intake was not standardized	N=9 adults with CF Age (median, range): 29; 23 to 42 years	Resolution of symptoms of malabsorption • Faecal fat excretion (FFE) (% of intake)	Both high- dose and low-dose are low-dose and very low- dose in current practice: indirect evidence
Heijerman 1993 (Netherlands) Cross- over RCT	Pancrease 2 caps, 3 per day and Omeprazole placebo Pancrease 2 caps, 3 per day + Omeprazole 20 mg once in the morning Additional information: Constituent enzymes per capsule of Pancrease: 5000 units lipase, 2900 units amylase, 330 units protease per capsule	N=11 adults with cystic fibrosis Age: 20 to 42	Resolution of symptoms of malabsorption • Faecal fat excretion (FFE) (% of intake)	This dose is a very-low dose in current practice: indirect evidence
Mitchell 1982 (New Zealand) Cross- over RCT	High-dose 22 capsules/day Low-dose 11 capsules/ day Pancrease®. Additional information: Constituent enzymes per capsule 4,000 USNF lipase units; 25,000 USNF protease units; 20,000 amylase units. Fat intake was not standardized	N=12 children and young people with CF Age: (mean, SD): 9.6±2.1 years	Resolution of symptoms of malabsorption • Faecal fat excretion (FFE) (g/kg/day) • Faecal fat excretion (FFE) (g/day) • Fat absorption (CFA) (%) Resolution of symptoms of malabsorption • Stool frequency (bowel actions/ day) • Abdominal pain	

1
2
CF: cystic fibrosis; CFA: coefficient of fat absorption; FFE: faecal fat excretion; RCT: randomised controlled trial;
SD: standard deviation

1 10.2.4 Clinical evidence profile

2 The clinical evidence profiles for this review question are presented in Table 152 to Table
3 157.

4 10.2.4.1 Acid suppressing agents as adjuvant therapy to PERT

5 **Table 152: Summary clinical evidence profile: Comparison 1.1. PERT + cimetidine**
6 **versus PERT alone in children**

Comparison 1.1. PERT + cimetidine versus PERT alone in children						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	PERT alone	PERT + cimetidine				
Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 14 days	The mean faecal fat excretion (as % of intake) in the PERT alone group was 27.7	The mean faecal fat excretion (as % if intake) in the PERT + cimetidine group was 17.8 (p<0.01)		21 (Durie 1980) ^{1, 2}	⊕⊕⊕ ⊖ low ^{3,4}	
Faecal fat excretion (FFE) Measured as g/24 hours Follow-up: 14 days	The mean faecal fat excretion (as g/24 h) in the control PERT alone was 31.3	The mean faecal fat excretion (as g/24 h) in the PERT cimetidine group was 11 lower (18.577 to 3.423 lower)		21 (Durie 1980) ^{1, 2}	⊕⊕⊕ ⊖ low ^{5,6,7}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; FFE: faecal fat excretion; g: grams; MD: mean difference; PERT: pancreatic endocrine enzyme therapy						

7 *1 Cross-over trial*

8 *2 Treatment details: Cotazym 26 capsules/ day + Cimetadine 20 mg/kg/day or placebo*

9 *3 The quality of evidence was downgraded by 1 due to unclear randomization, concealment and single-blinding. The quality of the evidence was further downgraded by 1 due to the quality of the statistical analysis. Means are provided instead of medians, although it is not normally distributed.*

10 *4 Imprecision was not assessed, as it was considered not appropriate. See footnote 3.*

11 *5 The quality of evidence was downgraded by 1 due to unclear randomization, concealment and single-blinding.*

12 *6 The quality of the evidence was downgraded by 2 because method of measuring fat excreted is inaccurate, as it does not take into account fat intake.*

13 *7 The quality of the evidence was downgraded by 1 because the CI crossed 1 clinical MID*

14 **Table 153: Summary clinical evidence profile: Comparison 1.2. PERT + Ranitidine**
15 **versus PERT alone in children**

Comparison 1.2. PERT + Ranitidine versus PERT alone in children						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				

	PERT alone	PERT + ranitidine	CI)			
[PERT + low dose ranitidine] Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 12 days	The median fat absorption (as % of intake) in the PERT alone group was 80.37	The median fat absorption (as % of intake) in the PERT + low dose ranitidine group was 83.60 p=0.87*		12 (Francisco 2002) ^{1,2}	⊕⊕⊕⊕ high ^{3,4}	
[PERT + high dose ranitidine] Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 12 days	The median fat absorption (as % of intake) in the PERT alone group was 80.37	The median fat absorption (as % of intake) in the PERT + high dose ranitidine group was 80.91 p=1*		12 (Francisco 2002) ^{1,5}	⊕⊕⊕⊕ high ^{3,4}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFA: coefficient of fat absorption; CI: confidence interval; MD: mean difference; PERT: pancreatic endocrine enzyme therapy

* The paper provided raw data. Medians and p-values were calculated by the NGA technical team

1 Cross-over trial

2 Treatment details: Pancrease M10 or M16 + ranitidine or placebo. Children weighting ≤40 kg were given 5 mg/kg. Children weighting >40 kg received 150 mg. twice daily.

3 Imprecision cannot be calculated from medians.

4 Reporting bias not detected, but drugs were provided by the Pharmaceutical industry

5 Treatment details: Pancrease M10 or M16 + ranitidine or placebo. Children weighting ≤40 kg were given 10 mg/kg. Children weighting >40 kg received 300 mg. twice daily.

Table 154: Summary clinical evidence profile: Comparison 1.3. PERT + Omeprazole versus PERT alone in adults

Comparison 1.3. PERT + Omeprazole versus PERT alone in adults						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	PERT alone	PERT + omeprazole				
Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 12	The median fat absorption (as % of intake) in the PERT alone group was 88.59	The median fat absorption (as % of intake) in the PERT + omeprazole groups was 87.40 p≤0.05*		9 (Francisco 2002) ^{1,2}	⊕⊕⊕⊖ moderate ^{3,4}	

days					
[low-dose PERT + omeprazole or placebo] Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 4 weeks	The median faecal fat excretion (as % of intake) in the PERT alone group was 20	The median faecal fat excretion (as % of intake) in the PERT + omeprazole group was 14 p>0.05		9 (Heijerman 1991) ^{1,5}	⊕⊕⊕⊕ very low ^{6,7,8,9}
[high-dose PERT + omeprazole or placebo] Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 4 weeks	The median faecal fat excretion (as % of intake) in the PERT alone group was 18	The median faecal fat excretion (as % of intake) in the PERT + omeprazole group was 9 p<0.01		9 (Heijerman 1991) ^{1,10}	⊕⊕⊕⊕ very low ^{6,7,8,9}
Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 4 weeks	The median fat excretion (as % of intake) in the PERT alone group was 20	The median faecal fat excretion (as % of intake) in the PERT + omeprazole group was 17 p>0.05		11 (Heijerman 1993) ^{1,11}	⊕⊕⊕⊕ low ^{12,13}

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFA: coefficient of fat absorption; CI: confidence interval; FFE: faecal fat excretion; PERT: pancreatic endocrine enzyme therapy

* The paper provided raw data. Medians and p-values were calculated by the NGA technical team

1 Cross-over trial

2 Treatment details: Pancrease M10 or M16 + omeprazole 20 mg/day or placebo

3 Imprecision cannot be calculated from medians

4 Reporting bias not detected, but drugs were provided by the Pharmaceutical industry. Quality of evidence was downgraded by 1 due to small population (n=9).

5 Treatment details: PERT 2 capsules x 3 times per day + Omeprazole 20mg/day or placebo. Constituent enzymes per capsule 5000u lipase, 2900u lipase, 330u protease. Fat intake was not standardized.

6 The quality of the evidence was downgraded by 1 due to unclear randomization and concealment

7 The quality of the evidence was of evidence downgraded by 2 as this dosage is not used in current practice

8 Imprecision cannot be calculated from medians.

9 Reporting bias not detected. Evidence downgraded by 1 due to small sample size (n=9).

10 Treatment details: PERT 4 capsules x 3 times per day + Omeprazole 20mg/day or placebo. Constituent enzymes per capsule 5000u lipase, 2900u lipase, 330u protease. Fat intake was not standardized.

11 Treatment details: PERT 2 capsules x 3 times per day + Omeprazole 20mg/day or placebo. Constituent enzymes per capsule 5000u lipase, 2900u lipase, 330u protease. Fat intake was not standardized.

12 The quality of the evidence was of evidence downgraded by 2 as this dosage is not used in current practice
13 Imprecision cannot be calculated from medians

Table 155: Summary clinical evidence profile: Comparison 1.4. PERT + Ranitidine versus PERT alone in adults

Comparison 1.4. PERT + Ranitidine versus PERT alone in adults						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	PERT alone	PERT + ranitidine				
[PERT + low-dose ranitidine] Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 12 days	The median fat absorption (as % of intake) in the control group was 89.20	The median fat absorption (as % of intake) in the intervention group was 93.06 p=0.01*		10 (Francisco 2002) ^{1,2}	⊕⊕⊕⊕ high ^{3,4}	
[PERT + high-dose ranitidine] Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 12 days	The median fat absorption (as % of intake) in the control group was 88.59	The median fat absorption (as % of intake) in the intervention group was 88.92 p≤0.05*		9 (Francisco 2002) ^{1,5}	⊕⊕⊕⊖ moderate ^{3,4,6}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CFA: coefficient of fat absorption; CI: confidence interval; PERT: pancreatic endocrine enzyme therapy						

* The paper provided raw data. Medians and p-values were calculated by the NGA technical team

1 Cross-over study

2 Treatment details: Pancrease M10 or M16 + ranitidine 150 mg. twice daily or placebo

3 Imprecision cannot be calculated from medians.

4 Reporting bias not detected, but drugs were provided by the Pharmaceutical industry

5 Treatment details: Pancrease M10 or M16 + ranitidine 300 mg. twice daily or placebo

6 Reporting bias not detected. Evidence downgraded by 1 due to small sample size (n=9).

12 10.2.4.2 High-dose PERT versus low-dose of PERT

Table 156: Summary clinical evidence profile: Comparison 2.1. High dose PERT versus low dose PERT in children

Comparison 2.1. High dose PERT versus low dose PERT in children					
Outcomes	Illustrative comparative risks* (95% CI)	Relative effect	No of Participants	Quality of the evidence	Comments

	Assumed risk	Corresponding risk	(95% CI)	(studies)	(GRADE)	
	Control					
Faecal fat excretion (FFE) (g/kg/day) Follow-up: 14 days	The mean faecal fat excretion (g/kg/day) in the control group was 0.437	The mean faecal fat excretion (g/kg/day) in the intervention groups was 0.141 lower (0.253 to 0.029 lower)		9 (Brady 1991) ^{1,2}	⊕⊕⊕⊕ very low ^{3,4,5,6,a}	a. This method of measuring fat excreted is inaccurate, as it does not take into account fat intake.
Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 14 days	The mean faecal fat excretion (as % of intake) in the control group was 13	The mean faecal fat excretion (as % of intake) in the intervention groups was 0 higher (0 to 0 higher)		9 (Brady 1991) ^{1,2}	⊕⊕⊕⊕ very low ^{3,4,5,6}	
Faecal fat excretion (FFE) (g/day) Follow-up: 9 days	-	The mean faecal fat excretion (g/day) in the intervention groups was 5 lower (8.877 to 1.123 lower)		30 (Brady 1991, Beker 1994) ^{1,2,3}	⊕⊕⊕⊕ very low ^{4,5,7,a}	a. This method of measuring fat excreted is inaccurate, as it does not take into account fat intake.
Faecal fat excretion (FFE) (g/day) Follow-up: 4 weeks	The mean faecal fat excretion (g/day) in the control group was 11.5	The mean faecal fat excretion (g/day) in the intervention groups was 8.7		12 (Mitchell 1982) ^{2,8}	⊕⊕⊕⊕ very low ^{4,9,10,11,a}	The MD could not be calculated a. This method of measuring fat excreted is inaccurate, as it does not take into account fat intake.
Fat absorption	The mean fat	The mean fat absorption in		12 (Mitchell	⊕⊕⊕⊕ very	

(CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 4 weeks	absorption (as % of intake) in the control group was 85.4	the intervention (as % of intake) groups was 89.5		1982) ^{2,8}	low ^{4,9,11,12}	
Fat absorption (CFA) Measured as % of intake (consumed fat that is absorbed) Follow-up: 9 days	The mean fat absorption (as % of intake) in the control group was 86.2	The mean fat absorption (as % of intake) in the intervention groups was 91.2		21 (Beker 1994) ^{2,3}	⊕⊕⊕⊕ very low ^{4,12,13,14}	The MD could not be calculated
Stool frequency (bowel movements/day, self-report) Follow-up: 4 weeks	The mean stool frequency in the intervention group was 1.8	The mean stool frequency in the intervention groups was 0.1 lower (0.189 lower to 0.011 higher)		12 (Mitchell 1982) ^{2,8}	⊕⊕⊕⊕ very low ^{4,9,11}	
Abdominal pain self-report Follow-up: 4 weeks	Not reported	No differences were found between both groups	Not estimable ¹⁴	12 (Mitchell 1982) ^{2,8}	⊕⊕⊕⊕ very low ^{4,9,11,15}	
Adverse events (constipation, elevation in serum uric acid levels) self-report Follow-up: 9 days	None	None	Not estimable ¹⁴	21 (Beker 1994) ^{2,3}	⊕⊕⊕⊕ very low ^{4,13,14,15}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CFA: coefficient of fat absorption; CI: confidence interval; FFE: faecal fat excretion; g: grams; kg: kilogrammes; MD: mean difference; ns: not significant; PERT: pancreatic endocrine enzyme therapy; SEM: standard error of measurement

1 Cross-over trial

2 Treatment details: high-dose 12 (8 to 18) & low-dose 3 (2 to 5) capsules per meal. Constituent enzymes per capsule: 7.020u of lipase. Daily fat intake (g) 94±6 in both groups.

3 Treatment details: high-dose: 1500u lipase per kg/body for meals & 750u lipase per kg/body for snacks. Low-dose: 500u lipase per kg/body for meals & 250u lipase per kg/body for snacks. Daily fat intake (g): 100g in both groups.

4 The quality of the evidence was downgraded by 2 as these doses are not used in current practice. Low-dose is in fact very low dose, and high-dose is just low-dose

5 Imprecision could not be calculated, as SD was not available for the control group

6 Reporting bias not detected, although funding not reported. Evidence downgraded by 1 due to small sample (n=9)

1
2
3
4
5
6
7
8
9
10
11

7 The quality of the evidence was downgraded by 1 due to unclear randomization and concealment in both studies.
 8 Treatment details: high-dose 22 capsules/day & low-dose 11 capsules/ day Pancrease®. Constituent enzymes per capsule 4,000 USNF lipase units; 25,000 USNF protease units; 20,000 amylase units.
 9 The quality of the evidence was downgraded by 1 due to unclear randomization and concealment. It is unclear if blinding was done, but given the outcome this may not have an impact.
 10 The quality of the evidence was downgraded by 1 as the results are poorly reported: authors do not report p-value and MD cannot be calculated
 11 Reporting bias not detected, although Pancrealipase capsules were provided by Ethnor Pty Ltd.
 12 The quality of the evidence was downgraded by 2 due to the quality of the statistical analysis. Means are provided instead of medians, although it is not normally distributed, therefore differences cannot be calculated as it is not appropriate.
 13 The quality of the evidence was downgraded by 1 because it is an open-label study.
 14 Reporting bias not detected, although the study is partly funded by a grant from Johnson Pharmaceutical.
 15 Imprecision cannot be calculated.

Table 157: Summary clinical evidence profile: Comparison 2.2. High-dose PERT versus low-dose PERT in adults

Comparison 2.2. High-dose PERT versus low-dose PERT in adults						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control					
Faecal fat excretion (FFE) Measured as % of intake (consumed that is excreted) Follow-up: 14 days	The median faecal fat excretion (% of intake) in the control groups was 20 (12 to 44)	The median faecal fat excretion (% of intake) in the intervention groups was 18 (10 to 34); p>0.05		9 (Heijerman 1991) ^{1,2}	⊕⊕⊕⊕ very low ^{3,4,5,6}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; FFE: faecal fat excretion; PERT: pancreatic endocrine enzyme therapy						

18 1 Cross-over trial
 19 2 High-dose 4 capsules x 3 times per day & low-dose 2 capsules x 3 times per day. Constituent enzymes per capsule 5000u lipase, 2900u lipase, 330u protease. Fat intake was not standardized.
 20 3 Evidence downgraded by 1 due to unclear randomization and concealment.
 21 4 Evidence downgraded by 2 as these doses are not used in current practice. Low-dose is in fact very low dose, and high-dose is just low-dose
 22 5 Imprecision cannot be calculated from medians
 23 6 Reporting bias not detected. Evidence downgraded by 1 due to small sample size (n=9).
 24
 25

26 10.2.5 Economic evidence

27 No economic evaluations of interventions relevant to PERT were identified in the literature
 28 search conducted for this guideline. Full details of the search and economic article selection
 29 flow chart can be found in Appendix E and F, respectively.

30 This review question was not prioritised for de novo economic modelling. To aid
 31 consideration of cost-effectiveness, relevant resource and cost use data are presented in
 32 Appendix K.

1 **10.2.6 Evidence statements**

2 **10.2.6.1 Comparison 1. Acid suppressing agents as adjuvant therapy to PERT**

3 **10.2.6.1.1 Evidence for children**

4 **Reduction of steatorrhea and faecal fat excretion**

5 Low quality evidence from 1 cross-over trial with 21 children with cystic fibrosis showed that
6 the mean faecal fat excretion (measured as percentage of consumed fat that is excreted)
7 was lower when the children received Cimetidine (20 mg/kg/day) in addition to PERT
8 treatment than when they received PERT and placebo at 2 weeks follow-up. However, the
9 uncertainty around this could not be calculated.

10 Low quality evidence from 1 cross-over trial with 21 children with cystic fibrosis showed that
11 there was a clinically significant difference in the reduction of faecal fat excretion (measured
12 as g/ 24 hours) when the children received Cimetidine (20 mg/kg/day) in addition to PERT
13 than when they received PERT and placebo at 2 weeks follow-up. This method of measuring
14 fat excreted is inaccurate, as it does not take into account fat intake.

15 **Fat absorption**

16 High quality evidence from 1 cross-over trial with 12 children with cystic fibrosis showed no
17 differences in the median of fat absorption (measured as % of consumed fat that is
18 absorbed) between adding low-dose Ranitidine (5 mg/kg for children weighting under 40 kg
19 or 150 mg twice daily for children weighting over 40kg) or placebo to PERT treatment at 12
20 days follow-up. However, the uncertainty around this could not be calculated.

21 High quality evidence from 1 cross-over trial with 12 children with cystic fibrosis showed no
22 differences in the median of fat absorption (measured as % of consumed fat that is
23 absorbed) between adding high-dose Ranitidine (10 mg/kg for children weighting under 40
24 kg or 300 mg twice daily for children weighting over 40kg) or placebo to PERT treatment at
25 12 days follow-up. However, the uncertainty around this could not be calculated.

26 **Resolution of symptoms of malabsorption**

27 No evidence was found for this critical outcome in children.

28 **Weight**

29 No evidence was found for this critical outcome in children.

30 **Quality of life**

31 No evidence was found for this important outcome in children.

32 **Satisfaction**

33 No evidence was found for this important outcome in children.

34 **Adverse events**

35 No evidence was found for this important outcome in children.

10.2.6.1.2 Evidence for adults

2 Reduction of steatorrhea and faecal fat excretion

3 Very low quality evidence from 1 cross-over trial with 9 adults with cystic fibrosis showed no
4 difference in the median faecal fat excretion (measured as percentage of consumed fat that
5 is excreted) between adding Omeprazole (20 mg/day) or placebo to low-dose PERT
6 treatment at 4 weeks follow-up. However, the uncertainty around this could not be calculated.

7 Very low quality evidence from 1 cross-over trial with 9 adults with cystic fibrosis showed a
8 decrease in the median faecal fat excretion (measured as percentage of consumed fat that is
9 excreted) when the participants received Omeprazole (20 mg/day) in addition to high-dose
10 PERT treatment than when they received PERT and placebo at 4 weeks follow-up. However,
11 the uncertainty around this could not be calculated.

12 Low quality evidence from 1 cross-over trial with 11 adults with cystic fibrosis showed no
13 difference in the median faecal fat excretion (measured as percentage of consumed fat that
14 is excreted) between adding Omeprazole (20 mg/day) or placebo to PERT treatment at 4
15 weeks follow-up. However, the uncertainty around this could not be calculated.

16 Fat absorption

17 Moderate quality evidence from 1 cross-over trial with 9 adults with cystic fibrosis showed
18 that the fat absorption (measured as % of consumed fat that is absorbed) was higher when
19 the participants received omeprazole (20 mg/day) in addition to PERT treatment than when
20 they received PERT and placebo at 12 days follow-up. However, the uncertainty around this
21 could not be calculated.

22 High quality evidence from 1 cross-over trial with 10 adults with cystic fibrosis showed that
23 the fat absorption (measured as % of consumed fat that is absorbed) was higher when the
24 participants received low-dose Ranitidine (150 mg twice daily) in addition to PERT treatment
25 than when they received PERT and placebo at 12 days follow-up. However, the uncertainty
26 around this could not be calculated.

27 Moderate quality evidence from 1 cross-over trial with 10 adults with cystic fibrosis showed
28 that the fat absorption (measured as % of consumed fat that is absorbed) was higher when
29 the participants received high-dose Ranitidine (300 mg twice daily) in addition to PERT
30 treatment than when they received PERT and placebo. However, the uncertainty around this
31 could not be calculated.

32 Weight

33 No evidence was found for this critical outcome in adults.

34 Resolution of symptoms of malabsorption

35 No evidence was found for this critical outcome in adults.

36 Quality of life

37 No evidence was found for this important outcome in adults.

38 Satisfaction

39 No evidence was found for this important outcome in adults.

1 **Adverse events**

2 No evidence was found for this important outcome in adults.

3 **10.2.6.2 Comparison 2. High versus low dose of PERT**

4 **40.2.6.2.1 Evidence for children**

5 **Reduction of steatorrhea and faecal fat excretion**

6 Very low quality evidence from 1 cross-over trial with 9 children with cystic fibrosis showed
7 that the mean faecal fat excretion (measured as g/kg/day) was lower when the children were
8 allocated to the high-dose treatment arm at 2 weeks follow-up. However, the uncertainty
9 around this could not be calculated. This method of measuring fat excreted is inaccurate, as
10 it does not take into account fat intake.

11 Very low quality evidence from 1 cross-over trial with 9 children with cystic fibrosis showed
12 that the mean faecal fat excretion (measured as percentage of consumed fat that is
13 excreted) was lower when the children were allocated to the high-dose treatment arm at 2
14 weeks follow-up. However, the uncertainty around this outcome cannot be calculated.

15 Very low quality evidence from 2 cross-over trials with 30 children with cystic fibrosis showed
16 that the mean faecal fat excretion (measured as g/day) was lower when the children were
17 allocated to the high-dose treatment arm at 4 weeks follow-up at 4 weeks follow-up.
18 However, the uncertainty around this could not be calculated. This method of measuring fat
19 excreted is inaccurate, as it does not take into account fat intake.

20 Very low quality evidence from 1 cross-over trial with 12 children with cystic fibrosis showed
21 no difference in the faecal fat excretion (measured as g/day) between the high-dose and the
22 low-dose PERT treatment arms at 4 weeks follow-up. However, the uncertainty around this
23 could not be calculated. This method of measuring fat excreted is inaccurate, as it does not
24 take into account fat intake.

25 **Fat absorption**

26 Very low quality evidence from 2 cross-over trials with 33 children with cystic fibrosis showed
27 that the fat absorption (measured as % of consumed fat that is absorbed) was higher when
28 the children were allocated to the high-dose treatment arm at 9 days and at 4 weeks follow-
29 up. However, the uncertainty around this could not be calculated.

30 **Weight**

31 No evidence was found for this critical outcome.

32 **Resolution of symptoms of malabsorption**

33 Very low quality evidence from 1 cross-over trial with 12 children with cystic fibrosis showed
34 no clinically significant difference in stool frequency between the high and the low dose
35 PERT treatment arms at 4 weeks follow-up.

36 Very low quality evidence from 1 cross-over trial with 12 children with cystic fibrosis showed
37 no difference in the occurrence of abdominal pain between the high and the low dose PERT
38 treatment arms at 4 weeks follow-up. However the uncertainty around this outcome could not
39 be calculated.

1 **Adverse Events**

2 Very low quality evidence from 1 cross-over trial with 21 children with cystic fibrosis showed
3 no episodes of constipation or elevation in serum acid levels in both high and low dose
4 treatment arms at 9 days follow-up.

5 **10.2.6.2.2 Evidence for adults**

6 **Reduction of steatorrhea and faecal fat excretion**

7 Very low quality evidence from 1 cross-over trial with 9 adults with cystic fibrosis showed no
8 clinically significant difference in the reduction of faecal fat excretion (measured as
9 percentage of intake) between high dose and low dose PERT treatment arms at 2 weeks
10 follow-up.

11 **Fat absorption**

12 No evidence was found for this important outcome.

13 **Weight**

14 No evidence was found for this critical outcome.

15 **Resolution of symptoms of malabsorption**

16 No evidence was found for this critical outcome.

17 **Adverse events**

18 No evidence was found for this important outcome.

19 **10.2.6.3 Economic evidence statements**

20 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

21 **10.2.7 Evidence to recommendations**

22 **10.2.7.1 Relative value placed on the outcomes considered**

23 The aim of this review was to evaluate the effectiveness of PERT given with an acid
24 neutralising or suppressing agent in comparison with PERT alone and to evaluate the
25 effectiveness of a high dose of PERT in comparison with a low dose of PERT in the
26 treatment of exocrine pancreatic insufficiency.

27 The critical outcomes specified by the Committee were reduction of steatorrhea, using
28 measures such as faecal fat excretion (total or as a percentage of fat consumption) and the
29 coefficient of fat absorption (the percentage of consumed fat absorbed, based on the
30 difference between fat consumed and fat excreted in the stool); weight, BMI (also % weight
31 for height if available) and resolution of symptoms of malabsorption. Quality of life,
32 satisfaction and drug related side effects were rated as important outcomes.

33 **10.2.7.2 Consideration of clinical benefits and harms**

34 The Committee noted that pancreatic enzyme insufficiency is a common disorder among
35 people with cystic fibrosis that prevents digestion and absorption of nutrients, resulting in
36 nutrient malabsorption and other symptoms such as diarrhoea. These in turn can affect
37 quality of life and eventually result in malnutrition. Based on this, the Committee agreed to
38 recommend that children and young people with cystic fibrosis should be offered an annual

1 test for exocrine pancreatic insufficiency, and that all people with cystic fibrosis should be
2 offered a test if there are signs or symptoms suggestive of malabsorption. This
3 recommendation was based on their clinical knowledge and experience.

4 The Committee discussed the technique that should be used to assess exocrine pancreatic
5 insufficiency. They agreed direct pancreatic stimulation tests (such as the direct pancreatic
6 stimulation test) are invasive, uncomfortable for the person, and costly. Instead, they
7 recommended the use of non-invasive techniques, such as stool elastase estimation, as it is
8 a non-invasive easy to perform alternative. They noted there are other alternatives, such as
9 the analysis of the percentage of fat absorbed or excreted, but they agreed they are
10 uncomfortable for the person (as a 72-hours stool collection is needed), and they rely on the
11 person accurately recording food intake.

12 The Committee agreed that the use of PERT is well-established in clinical practice, as it is
13 known that PERT treatment is useful in overcoming enzyme deficiency in people with cystic
14 fibrosis. However, they noted there is uncertainty regarding the optimal doses of enzymes
15 needed.

16 The Committee acknowledged the evidence presented comparing different treatment
17 dosages, but they agreed it was of limited use due poor quality of the studies. They noted
18 that although normalisation of fat absorption and thus prevention of steatorrhea would
19 ideally be achieved with PERT, demonstrating this was in practice difficult, because as
20 indicated above, the measurement of stool fat entails stool collection and measurement of
21 dietary fat intake over a period of days in order to calculate the coefficient of fat absorption.
22 This approach is rarely done other than in a research context, being troublesome and
23 impractical.

24 The Committee acknowledged that the evidence suggested improved fat absorption with
25 higher doses of pancreatic enzyme replacement therapy, but that dosage was likely to vary
26 on an individual basis, since the severity of pancreatic insufficiency was not uniform in
27 people with cystic fibrosis. Optimal dose might differ depending on body size and on dietary
28 composition and intake. Infants and young children have a higher intake of fat proportionately
29 than older children, young people and adults.

30 Based on this, the Committee agreed to recommend to offer PERT to people with cystic
31 fibrosis with pancreatic insufficiency, and that the dose should be adjusted for each person in
32 order to minimise symptoms of malabsorption. This recommendation is consistent with
33 clinical practice. The Committee also noted this recommendation aligned with the CF Trust
34 Consensus recommendations (CF Trust, Nutritional Management of Cystic Fibrosis 2016).

35 The Committee agreed that evidence regarding the effectiveness of PERT dose and acid
36 suppression in relation to resolution of malabsorption symptoms, improvement in weight, and
37 improvement in patient satisfaction or health-related quality of life was either very limited and
38 of very low quality or completely lacking. They noted that the normal clinical approach to
39 determining individual need was an empirical one (for instance titrating the PERT dose in
40 terms of units of lipase against the amount of fat being ingested). A standard dose, related to
41 age in children, was usually given and adjustment then made based on the clinical response,
42 in terms of trying to achieve a normal bowel habit and the resolution of any malabsorptive
43 symptoms. They recommended that in people with confirmed pancreatic exocrine
44 insufficiency the dose was titrated against symptoms and regularly reviewed. High enzyme
45 concentration products would aid treatment optimisation where there was a higher dose
46 requirement.

47 The Committee noted that trials were of short duration, and therefore it was not possible to
48 assess whether prescribing high dose PERT treatment or adding an antacid had an impact
49 on weight. They also noted the dosages used are very low compared to those used in clinical
50 practice. In addition, most trials included a very small sample size, and were underpowered
51 to detect differences between treatment arms.

1 10.2.7.3 Consideration of economic benefits and harms

2 It was noted by the Committee that the dose-response may justify the additional cost of high-
3 concentration PERT over low-concentration PERT.

4 With regards to acid suppression, the Committee agreed the clinical evidence was too
5 uncertain to justify the costs for intermittent malabsorption. However, in people with cystic
6 fibrosis with persistent symptoms of malabsorption, the Committee believed the benefits from
7 acid suppression would justify the costs.

8 Overall, the Committee agreed clinical judgement is necessary to provide the most cost-
9 effective treatment as the optimal dose and concentration of enzymes is individualised based
10 on weight and drug adherence.

11 10.2.7.4 Quality of evidence

12 The quality of the evidence presented in this report ranged from very low to high as assessed
13 by GRADE. The main reasons that lead to downgrading the quality of the evidence were the
14 following.

15 For the domain risk of bias, the 6 trials included in the review were assessed as having
16 unclear or high risk of bias regarding randomization and concealment methods. All of the
17 trials were of cross-over design, but given the nature of the intervention it is not expected that
18 treatments have a carry-over effect.

19 The Committee noted that there were issues regarding indirectness of the intervention, as
20 the dosage used in some of the studies was much lower than current practice. In addition,
21 they noted that measuring fat excretion or fat absorption as grams per 24 hours is
22 inaccurate, as it does not take into account daily intake.

23 Most trials were underpowered to detect a difference between both treatment groups.

24 Finally, there were some concerns regarding the statistical analysis used and the type of
25 data reported. Due to this limitation most of the results could not be meta-analysed.

26 Imprecision could not be assessed when results were provided in medians.

27 10.2.7.5 Other considerations

28 No equality issues were identified by the Committee for this review question.

29 The Guideline Committee agreed that research recommendations were not prioritised for this
30 topic because clinical practice is already established and there are treatments which are
31 known (based on clinical expertise) to be useful in the treatment of pancreatic exocrine
32 insufficiency.

33 10.2.7.6 Key conclusions

34 The Committee concluded that people with cystic fibrosis with pancreatic insufficiency should
35 be offered oral pancreatic enzyme replacement therapy. Although the evidence showed
36 some indication that high-dose PERT treatment may have a beneficial impact in both faecal
37 fat excretion and fat absorption when compared to standard or low-dose treatment, without
38 an increase in side effects, the Committee agreed that the dose should be adjusted to
39 minimise symptoms of malabsorption.

40 In addition, the Committee noted that there was some indication that addition of acid
41 suppressants to PERT treatment may reduce faecal fat excretion, and therefore the use of
42 an acid suppressant agents can be considered in people with cystic fibrosis who have
43 persistent symptoms of malabsorption despite optimal PERT therapy.

1 10.2.8 Recommendations

2 **99. Test for exocrine pancreatic insufficiency in people with cystic fibrosis, using a**
3 **non-invasive technique such as stool elastase estimation. If the test result is**
4 **normal, repeat it:**

- 5 • annually in children and young people
- 6 • if symptoms or signs suggesting malabsorption occur.

7 **100. Offer oral pancreatic enzyme replacement therapy to people with exocrine**
8 **pancreatic insufficiency. Adjust the dose as needed to minimise any symptoms or**
9 **signs of malabsorption.**

10 **101. Consider an acid suppression agent (for example an H2 receptor antagonist or**
11 **a proton pump inhibitor) for people who have persistent symptoms or signs of**
12 **malabsorption despite optimal pancreatic enzyme replacement therapy.**

13 10.3 Distal ileal obstruction syndrome

14 **Review question: What are the effective strategies for treatment and secondary**
15 **prevention of distal ileal obstruction syndrome?**

16 10.3.1 Introduction

17 Cystic fibrosis (CF) can cause a number of gastro-intestinal problems. One gastro-intestinal
18 problem characteristic of cystic fibrosis is a blockage in the bowel known as distal intestinal
19 obstruction syndrome, or 'DIOS' for short. DIOS is a specific cystic fibrosis-related clinical
20 problem and therefore is best managed by clinical teams with experience of treating people
21 with cystic fibrosis. Approximately 7-8 % of people with cystic fibrosis will experience an
22 episode of DIOS in their lifetime. Symptoms of DIOS are abdominal pain, abdominal
23 distension, nausea and vomiting. Typical clinical examination findings include a distended
24 abdomen, a palpable mass in the right iliac fossa, and reduced or absent bowel sounds.
25 Radiological imaging, usually involving an abdominal x-ray, ultrasound scan or CT-scan,
26 helps to confirm the diagnosis. DIOS is much more common in those patients who have
27 exocrine pancreatic insufficiency. The pathophysiology is not fully understood, but there are
28 often multiple contributory factors including, dehydration, salt loss, rapid increase in
29 pancreatic enzyme dosage or poor compliance with pancreatic enzyme therapy, altered gut
30 motility and pH, and dietary factors.

31 A number of patients suffer recurrent episodes in their lifetime. This review will look at the
32 evidence for strategies both to prevent DIOS and for treating episodes of DIOS if they occur.

33 10.3.2 Description of clinical evidence

34 The aim of this review was to identify the effective strategies of primary treatment (acute
35 treatment) in those with a diagnosis of cystic fibrosis and DIOS. Additionally, this review
36 aimed to identify the effective strategies for the secondary prevention of DIOS.

37 Systematic reviews and randomised controlled trials (RCTs) that assessed the effectiveness
38 of primary and secondary treatments for DIOS were eligible for inclusion in this review.
39 However, no relevant systematic reviews or RCTs were identified. Hence, as per the
40 protocol, conference abstracts of RCTs and non-randomised comparative studies were also
41 considered although none was identified for inclusion and no evidence was available to
42 inform the review.

43 For full details see review protocol in Appendix D.

1 See also study selection flow chart in Appendix F, and list of excluded studies in Appendix H.

2 **10.3.3 Summary of included studies**

3 Not applicable, as no studies were identified for inclusion in this review.

4 **10.3.4 Clinical evidence profile**

5 Not applicable, as no studies were identified for inclusion in this review.

6 **10.3.5 Economic evidence**

7 No economic evaluations of interventions relevant to DIOS were identified in the literature
8 search conducted for this guideline. Full details of the search and economic article selection
9 flow chart can be found in Appendix E and F, respectively.

10 This review question was not prioritised for de novo economic modelling. To aid
11 consideration of cost-effectiveness, relevant resource and cost use data are presented in
12 Appendix K.

13 **10.3.6 Evidence statements**

14 Not applicable, as no clinical or economic studies were identified for inclusion in this review.

15 **10.3.7 Evidence to recommendations**

16 **10.3.7.1 Relative value placed on the outcomes considered**

17 The aim of this review was to identify the most effective strategies for the primary treatment
18 and secondary prevention of DIOS.

19 For the primary treatment, when a person presented with acute abdominal pain and other
20 symptoms due to DIOS, the Committee identified the following outcomes as critical:
21 reduction in clinical manifestations, adverse events from treatment and treatment failure.
22 Patient satisfaction and duration of hospital stay were rated as important outcomes.

23 For the secondary prevention of DIOS, the Committee identified the following outcomes as
24 critical: reduction in clinical manifestations, adverse events from treatment and recurrence of
25 DIOS. Patient satisfaction was rated as an important outcome.

26 **10.3.7.2 Consideration of clinical benefits and harms**

27 The Committee noted the absence of trials in the review. They agreed that there are
28 difficulties in undertaking randomised controlled trials of treatments for those presenting with
29 DIOS. In particular, the clinical presentation varies considerably, some having relatively
30 minor symptoms (for example abdominal pain) while others might have symptoms of
31 intestinal obstruction with vomiting. The approach to managing these patients would be
32 tailored to their individual difficulties. Given the lack of evidence, the list of recommendations
33 as part of this topic are derived from both current good practice and the Committee expert
34 opinion. In addition the Committee emphasized the variability in practice, and the need to
35 define a correct treatment. Treatment of an acute episode of DIOS

36 Although the diagnosis of DIOS was not part of this review, the Committee considered it was
37 important to make recommendations on the recognition of DIOS in people with cystic fibrosis.
38 Therefore, by consensus based on their clinical knowledge and experience they agreed
39 recommendations on this.

1 First, the Committee highlighted the importance of making a differential diagnosis of acute
2 abdominal pain in people with cystic fibrosis. Some causes of abdominal pain that may
3 resemble DIOS include constipation, appendicitis, intussusception, and cholecystitis. Their
4 correct identification is important in order to decide the adequate treatment approach.

5 Although the symptoms of DIOS are variable, they agreed that DIOS should be suspected in
6 people with cystic fibrosis who have an acute onset of peri-umbilical or right lower quadrant
7 abdominal pain, and any of the following signs or symptoms are present: a palpable faecal
8 mass in the right lower quadrant, faecal loading in the right lower quadrant on a plain
9 abdominal radiograph, especially if associated with small intestine air-fluid levels and clinical
10 features of partial or complete intestinal obstruction, such as vomiting (especially bilious) and
11 abdominal distension. In the absence of clinical or radiological features of DIOS, additional
12 imaging techniques, such as an abdominal ultrasound scan or an abdominal CT scan
13 considered.

14 They noted that people with cystic fibrosis presenting to non-cystic fibrosis clinical settings
15 might be assumed to have appendicitis or other suspected causes for intestinal obstruction
16 when in fact DIOS was more likely, and therefore be more likely to be subjected to
17 unnecessary and potentially harmful surgical intervention.

18 Although it is common for a distinction to be made between complete versus impending
19 DIOS the Committee choose not to make this distinction as it was more accurate to conceive
20 of the syndrome as consisting of a continuum, some presenting with acute pain and evidence
21 of stool retention in the distal ileum (either a palpable mass in the right iliac fossa or based
22 on radiological imaging) while others might have overt clinical and radiological evidence of
23 distal small intestinal obstruction. It was worth noting that some people with cystic fibrosis
24 have evidence of a palpable mass in the right lower quadrant of the abdomen for years
25 without developing the symptoms of DIOS.

26 As noted above, the Committee recognised the lack of randomised controlled trials for the
27 treatment of DIOS and the prevention of recurrent DIOS. However they agreed there are
28 certain basic principles governed currently accepted management.

29 The Committee discussed that important contributors to the development of DIOS were
30 dehydration of intestinal mucus due to the underlying defect in the cystic fibrosis
31 conductance regulator. Other possible contributors were the increased viscosity of luminal
32 contents and alterations in intestinal motility (delayed small intestinal transit) in cystic fibrosis.
33 Although DIOS can occur in people with cystic fibrosis who are pancreatic sufficient there is
34 speculation that fat malabsorption can increase the risk either by increasing luminal content
35 viscosity or by delaying intestinal transit (the ileal brake mechanism). For these reasons the
36 Committee agreed that adequate hydration and if necessary rehydration was advisable in
37 those presenting with DIOS and that encouraging ample oral fluid intake was advisable in
38 those who were experiencing recurrent episodes of DIOS. In the latter group it was also a
39 rational approach to review and attempt to optimise oral pancreatic enzyme replacement
40 therapy as a preventive measure.

41 The Committee noted that 2 approaches to medical therapy for DIOS were currently in
42 common use. Diatrizoate meglumine and diatrizoate sodium solution (Gastrografin) could be
43 given either orally or via an enteral tube. They considered that this was often the quickest
44 and most effective way of dealing with a blockage, and adverse effects, if any, were usually
45 minor. In addition, it is usually patient preference due to its ease to take and being more
46 tolerable. Based on this, the Committee agreed that a treatment regimen could be
47 considered as first-line treatment.

48 If this was unsuccessful, the Committee recommended a trial of treatment with stool
49 softeners such as osmotic laxatives containing polyethylene glycol (PEG) in an iso-osmotic
50 electrolyte solution. This could be taken orally or administered through an enteral tube.

1 Occasionally, if these measures proved unsuccessful in relieving the DIOS symptoms,
2 recourse to surgery to alleviate the obstruction could be considered. However they
3 emphasised surgery should only be considered as a last resource.

4 **Secondary prevention**

5 In addition to the potential value of enhanced oral fluid intake and optimisation of pancreatic
6 enzyme replacement, as discussed above, the Committee recommended trying a stool
7 softening agent such as osmotic laxatives containing polyethylene glycol (PEG) in an iso-
8 osmotic electrolyte solution. There was no clinical trial evidence for the effectiveness of such
9 agents in this setting, but the Committee considered that they made rational sense and were
10 readily tolerated and that if recurrent DIOS was a concern it was reasonable to give a trial of
11 prophylaxis with them. Osmotic laxatives as described above are commonly used in the long-
12 term management of chronic constipation and are free of serious adverse effects.

13 **10.3.7.3 Consideration of economic benefits and harms**

14 The Committee considered imaging studies to reduce the number of false positives as some
15 people with cystic fibrosis have potentially serious conditions such as appendicitis or
16 intussusception that might mimic the symptoms of DIOS. For this reason, the Committee
17 agreed imaging studies would be a cost-effective use of NHS resources as they would
18 minimise the foregone benefits and additional treatment costs associated with unidentified
19 cases.

20 The Committee also added that DIOS can be a serious and costly condition that negatively
21 impacts a person's health-related quality of life if it is incorrectly managed. Therefore, it
22 would be necessary to manage suspected DIOS in a specialist cystic fibrosis centre, with
23 supervision from specialists who have expertise in recognising and treating the condition and
24 its complications in order to minimise the number of incorrectly managed cases. As a result,
25 a recommendation was prioritised to ensure resource allocations in this area are maintained.

26 According to the Committee people with an acute onset of DIOS symptoms can respond to
27 an intravenous fluids to regain adequate hydration, or osmotic laxative containing
28 polyethylene glycol which is a relatively inexpensive treatment at a cost of less than £10 per
29 month. However, the Committee noted that the more expensive treatment, diatrizoate, is
30 often chosen as the first line treatment in current clinical practice based on experience that it
31 is the most effective first line treatment with the potential for fewer side effects. Moreover,
32 cheaper alternatives would overtake the cost of diatrizoate when they are required for longer
33 durations to achieve the same effects..

34 Following this, the Committee also agreed that the cost and distress incurred by a surgical
35 procedure would be overtaken by pharmacological treatments that require ongoing costs and
36 treatment burden. As a result, the Committee made a recommendation to consider surgery if
37 prolonged pharmacological treatment is not effective.

38 When making their recommendations, the Committee referred to previous guidance
39 (Colombo 2011) that recommends a stepwise approach to DIOS treatment, using the least
40 invasive options first and surgery only as a last resort. As a result, the Committee believed
41 guideline recommendations are not likely to represent a change in current practice and
42 resource use.

43 **10.3.7.4 Quality of evidence**

44 Not applicable, as no studies were identified for inclusion in this review.

1 10.3.7.5 Other considerations

2 The guideline committee were not aware of any published trial and it was their clinical
3 experience that the choice of a treatment over another was mainly empirically based. Current
4 good practice guidance (Colombo 2011) was acknowledged by the Guideline Committee,
5 which they agreed was largely in line with their clinical experience.

6 The Committee discussed potential equality issues. They noted adults may be more likely to
7 go into a local hospital if the live far from the specialist centre. However they agreed there
8 was no need to draft additional recommendations, as they had already recommended that
9 suspected DIOS should be managed in a specialist cystic fibrosis centre, with supervision
10 from specialists who have expertise in recognising and treating the condition and its
11 complications.

12 The Guideline Committee agreed that research recommendations were not prioritised for this
13 topic because clinical practice is already established and there are treatments which are
14 known (based on clinical expertise) to be useful in the acutely ill patient and for secondary
15 prevention.

16 10.3.7.6 Key conclusions

17 The Guideline Committee concluded that in the absence of relevant published evidence,
18 recommendations are based on their clinical experience, expert opinion and existing
19 guidance. They agreed treatment of DIOS should be provided in a cystic fibrosis centre. The
20 less invasive treatments should be the first choice for treating DIOS, as they are generally
21 effective in almost all patients with DIOS.

22 10.3.8 Recommendations

23 **102. Be aware that a variety of conditions can cause acute abdominal pain and**
24 **resemble distal ileal obstruction syndrome in people with cystic fibrosis, for**
25 **example:**

- 26 • constipation
- 27 • appendicitis
- 28 • intussusception
- 29 • cholecystitis.

30 **103. Suspect distal ileal obstruction syndrome in people with cystic fibrosis who**
31 **have an acute onset of peri-umbilical or right lower quadrant abdominal pain and**
32 **any of the following:**

- 33 • a palpable mass in the right lower quadrant
- 34 • faecal loading in the right lower quadrant on a plain abdominal
35 radiograph, especially if associated with small intestine air-fluid levels
- 36 • clinical features of partial or complete intestinal obstruction, such as
37 vomiting (especially bilious) and abdominal distension.

38 **104. For people who have an acute onset of peri-umbilical abdominal pain but no**
39 **other clinical or radiological features of distal ileal obstruction syndrome,**
40 **consider further investigation with an:**

- 41 • abdominal ultrasound scan or
- 42 • abdominal CT scan.

- 1 **105. Manage suspected distal ileal obstruction syndrome in a specialist cystic**
2 **fibrosis centre, with supervision from specialists who have expertise in**
3 **recognising and treating the condition and its complications.**
- 4 **106. Offer oral or intravenous fluids to ensure adequate hydration (and rehydration**
5 **if needed) for people with distal ileal obstruction syndrome.**
- 6 **107. Consider diatrizoate meglumine and diatrizoate sodium solution (orally or via**
7 **an enteral tube) as first-line treatment for distal ileal obstruction syndrome.**
- 8 **108. If diatrizoate meglumine and diatrizoate sodium solution is not effective,**
9 **consider an iso-osmotic polyethylene glycol and electrolyte (PEG) solution (orally**
10 **or via an enteral tube) as a second-line treatment.**
- 11 **109. Consider surgery as a last resort, if prolonged treatment with a PEG solution is**
12 **not effective.**
- 13 **110. To reduce the risk of distal ileal obstruction syndrome recurring:**
- 14 • encourage people to drink plenty of fluids
 - 15 • optimise pancreatic enzyme replacement therapy (see Exocrine
 - 16 pancreatic insufficiency).
 - 17 • consider advising regular treatment with a stool-softening agent such as
 - 18 lactulose or a PEG solution.

19

20 **10.4 Monitoring for liver disease**

21 **Review question 1. What is the diagnostic accuracy of tests to detect / strategies to**
22 **detect early and late cystic fibrosis liver disease?**

23 **Review question 2. What is the diagnostic and prognostic value of different strategies**
24 **to detect cystic fibrosis liver disease and predict progression (including progression**
25 **to cirrhosis and portal hypertension with (out) oesophageal varices)?**

26 **10.4.1 Introduction**

27 The estimated prevalence of liver involvement in cystic fibrosis is reported to be between
28 26% and 45%. Clinical presentation with hepatomegaly and/or splenomegaly usually occurs
29 within the first 10 years of age, and there may be progression from focal hepatic biliary
30 fibrosis to the development of cirrhosis and then portal hypertension with or without
31 oesophageal varices complications (Debray 2011).

32 Diagnosis of cystic fibrosis-related liver disease is typically made by clinical examination
33 (presence or absence of hepatosplenomegaly), biochemical tests, radiological methods (for
34 example ultrasound scanning), and where necessary histological assessment (for example
35 liver biopsy). Current UK practice uses clinical examination and biochemical liver testing to
36 determine a baseline and to monitor progression of disease thereafter annually. Ultrasound
37 is not usually offered on an annual basis to those without any signs of liver disease, but may
38 be performed at intervals. Early detection of hepatic injury and fibrosis enables early
39 treatment which may prevent further damage or even reverse very early damage.

40 There are currently no universally accepted definitions of liver disease (early, progressive,
41 late), and these vary according to tests performed.

1 More recently new definitions of liver disease have come into practice as a result of
2 recommendations that are based on results from monitoring tests performed at clinical
3 review; for example, clinical examination and routine liver function testing, with ultrasound
4 imaging only being used if necessary. Broadly cystic fibrosis-related liver disease is
5 diagnosed if certain criteria are met; for example, some have suggested diagnosis based on
6 2 of the following conditions being met on (at least 2) consecutive examinations spanning a
7 set period (for example 3 months to 1 year):

- 8 • hepatomegaly confirmed by ultrasound;
- 9 • 2 abnormal (>upper limit of normal) serum liver enzyme levels (eg alanine
10 aminotransferase, ALT; aspartate aminotransferase, AST; alkaline phosphatase, ALP,
11 gamma-glutamyl transferase, GGT);
- 12 • ultrasound abnormalities other than hepatomegaly (increased heterogeneous
13 echogenicity, nodularity, irregular margins).

14 If liver disease has not developed by adulthood, it may be that the patient will not develop it.
15 Therefore the Committee queried the value of using ultrasound at annual review in addition
16 to clinical examination and biochemical liver function tests when there was no indication of
17 liver disease from either test.

18 **10.4.2 Description of clinical evidence**

19 The aim of this review was to assess the diagnostic accuracy of different diagnostic
20 strategies to detect cystic fibrosis- related liver disease (including cirrhosis, portal
21 hypertension and oesophageal varices) defined by gold standard tests and to identify
22 whether any tests are useful in predicting the progression of cystic fibrosis-related liver
23 disease.

24 No “test and treat” randomised controlled trials (RCTs), were available. Thus, for the
25 diagnostic part of the review we looked for SRs of diagnostic studies, cross-sectional
26 diagnostic studies, and case-control studies. For the prognostic question we searched for
27 SRs of prognostic studies and cohort studies.

28 For full details see review protocol in Appendix D.

29 Eleven studies were included in this review (Fagundes 2004, Karlas 2012, Kitson 2013,
30 Lewindon 2011, Lindblad 1999, Mueller-Abt 2008, Patriquin 1999, Rath 2012, Rath 2013,
31 Sadler 2015, Witters 2009).

32 Studies were from Germany (Karlas 2012, Rath 2012, Rath 2013), Sweden (Lindblad 1999),
33 Belgium (Witters 2009), Australia (Kitson 2013, Lewindon 2011, Mueller-Abt 2008), Canada
34 (Patriquin 1999, Sadler 2015) and Brazil (Fagundes 2004).

35 With regards to study design, 7 were prospective cohorts (Fagundes 2004, Karlas 2012,
36 Lewindon 2011, Patriquin 1999, Rath 2012, Rath 2013, Sadler 2015), 3 were retrospective
37 cohorts (Lindblad 1999, Mueller-Abt 2008, Witters 2009) and 1 was a case control study
38 (Kitson 2013) with population sizes ranging from 30 to 280 participants.

39 With regards to the population, 3 studies recruited children and young people (Lewindon
40 2011, Mueller-Abt 2008, Patriquin 1999), 3 studies only recruited adults (Karlas 2012, Kitson
41 2013, Sadler 2015) and 5 studies recruited both children and adults (Fagundes 2004,
42 Lindblad 1999, Rath 2012, Rath 2013, Witters 2009), 1 of which reported children’s and
43 adults’ results separately (Rath 2012).

44 Evidence was available for the following, used as reference standards to detect cystic
45 fibrosis-related liver disease: clinical examination, biochemical testing, clinical examination
46 and biochemical testing in combination, ultrasound, published cystic fibrosis-related liver
47 disease definitions (incorporating clinical examination, biochemical testing and ultrasound),

1 and biopsy. No evidence was available for MRI or CT scanning used as the reference
2 standard.

3 Evidence was available for the following index tests: single biochemical tests (ALT, AST,
4 GGT, and ALP), formulas using biochemical and other parameters (APRI and Forn's score),
5 ultrasound, transient elastography (FibroScan®) and clinical examination and ultrasound (in
6 combination).

7 The definitions of cystic fibrosis-related liver disease varied according to the gold standard
8 used and mostly incorporated more severe disease (cirrhosis). In a few studies, either
9 separate data was available for or the reference standard was limited to definitions of cystic
10 fibrosis-related liver disease (without cirrhosis), cirrhosis or portal hypertension (with or
11 without oesophageal varices).

12 Diagnostic data (sensitivity, specificity, likelihood ratios) were available or calculated from
13 information available in the full text of the report for all comparisons. Area under receiver
14 operating characteristic curve (AUROC) data was less frequently presented in study reports.

15 Prognostic data for the development of cystic fibrosis-related liver disease and portal
16 hypertension were available from 3 studies (Kitson 2013, Lewindon 2011, Woodruff 2016).

17 A summary of the included studies is presented in Table 158. See study selection flow chart
18 in Appendix F, study evidence tables in Appendix G, list of excluded studies in Appendix H,
19 forest plots in Appendix I, and full GRADE profiles in Appendix J.

20 10.4.3 Summary of included studies

21 A summary of the 11 studies included in this review is presented in Table 158

22 **Table 158: Summary of studies**

Study	Index test	Reference standard	Population	Target condition
Fagundes 2004 (Brazil) Prospective cohort study	<ul style="list-style-type: none"> Williams ultrasound score: normal ultrasound results (score = 3) or abnormal (score > 3). 	<ul style="list-style-type: none"> Clinical and biochemical criteria. Abnormal clinical examination: <ul style="list-style-type: none"> the presence of a palpable spleen and/or hepatomegaly (presence of a palpable liver more than 2.5 cm below the right costal margin of firm consistency). Abnormal biochemistry: <ul style="list-style-type: none"> a significant and persistent increase of at least 1.5 times the upper limit of the reference range, of at least 2 of the enzymes aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (AP) or gamma-glutamyl 	N=70 infants, children and young people with CF Mean age (SD): 10.9 (6.4) Participants were followed prospectively at a CF outpatient clinic at a Brazilian university and who underwent clinical, biochemical and ultrasound examinations.	<ul style="list-style-type: none"> CFLD (includes cirrhosis)

Study	Index test	Reference standard	Population	Target condition
		transpeptidase (GGT), for a period of more than 6 months.		
Karlas 2012 (Germany) Case control study	<ul style="list-style-type: none"> • Transient elastography (TE): • AST/Platelets-Ratio-Index (APRI) • Forns' score was calculated according to the formula: score=7.811-3.131 x platelet count (109 /l)+0.781 x ln GGT (U/l) + 3.467 x ln age (years)-0.014xcholesterol (mg/dl) 	<ul style="list-style-type: none"> • Cystic fibrosis-related liver disease: at least 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-year period (Sokol 1999, Colombo 2002): <ul style="list-style-type: none"> ○ (1) Ultrasound confirmed hepatomegaly; ○ (2) elevated serum liver enzyme levels of ALT, AST, AP, or GGT; ○ (3) ultrasound abnormalities other than hepatomegaly (i.e., increased, heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). • Liver cirrhosis: distinct ultrasonographic signs (i.e. coarse nodularity, presence of portal hypertension and rarefaction of peripheral portal veins) and clinical signs (e.g. esophageal varices, splenomegaly) 	<p>N=55 adults with CF, n=14 with CFLD</p> <ul style="list-style-type: none"> • Mean age (SD): 31.9 (8.8) • Participants were prospectively investigated at presentation to the pulmonary outpatient clinic for clinical routine examinations. • Healthy probands and patients with alcoholic liver cirrhosis served as controls to establish test ranges 	<ul style="list-style-type: none"> • CFLD (includes cirrhosis) • Liver cirrhosis
Kitson 2013 Australia Case control study	<ul style="list-style-type: none"> • Transient elastography (TE): (FibroScan®) • AST/Platelets-Ratio-Index (APRI) performed at baseline 	<ul style="list-style-type: none"> • CFLD: according to established criteria if least two of the following conditions on consecutive examinations spanning a one-year period were present: <ul style="list-style-type: none"> ○ (i) Hepatomegaly and/or splenomegaly confirmed by ultrasound, ○ (ii) abnormal serum liver enzyme levels, consisting of elevation above the upper limit of normal of 2 of the following: ALT, AST, GGT, ○ (iii) ultrasound abnormalities other 	<p>N=50 adults with CF</p> <ul style="list-style-type: none"> • 25 with CFLD, mean age (SD): 30.5 (9.3) • 25 without CFLD, mean age (SD): 34.1 (9.8) • Participants were prospectively studied at a large CF referral centre in Australia. 	<ul style="list-style-type: none"> • CFLD (includes cirrhosis) • Portal Hypertension • Oesophageal varices

Study	Index test	Reference standard	Population	Target condition
		than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular margins; splenomegaly; presence of porto-systemic collateral veins; ascites).		
Lemaitre 2016 (France) Retrospective cohort study	<ul style="list-style-type: none"> • Transient Elastography: Results were expressed in kilopascal (kPa) using the Metavir scoring system • Index test: Biliary and Hepatic Magnetic Resonance Imaging 	<ul style="list-style-type: none"> • Liver function test or ultrasound: Details not reported 	<p>N=25 adults with CF</p> <ul style="list-style-type: none"> • Median age, years (range): 25 (18 to 43) • The sample included adult with CF, investigated by hepatobiliary MRI and by transient elastography for liver stiffness measurement (LSM) between July 2009 and July 2010 	<ul style="list-style-type: none"> • CFLD
Lewindon 2011 (Australia) Prospective cohort study	<ul style="list-style-type: none"> • Clinical examinations: Hepatomegaly with or without splenomegaly • Serum ALT levels performed at enrollment • Ultrasound images were obtained after fasting. Briefly, liver images were recorded as nodular edge, nodular, heterogeneous, or normal echogenicity with or without splenomegaly. Normal US was defined as normal echogenicity with no splenomegaly. US evidence of 	<ul style="list-style-type: none"> • Dual pass percutaneous liver biopsy with US guidance under general anesthesia For fibrosis scoring, the Scheuer F0-F4 staging system was used (F0 =no fibrosis, F4 = cirrhosis). Only sections with at least five portal tracts were deemed adequate for assessment. 	<p>N=40 children and young people with CF</p> <ul style="list-style-type: none"> • Age range: 2.38 to 18.73 years, median=10.64 years at enrollment) • Participants were attending a cystic fibrosis referral clinic with suspected cystic fibrosis liver disease defined as the following: <ul style="list-style-type: none"> • hepatomegaly (HM) with or without splenomegaly • a persistent (>6-month) elevation of serum alanine aminotransferase (ALT; level > 1.5 x upper limit of normal) 	<ul style="list-style-type: none"> • F1-F4 fibrosis • F2-F4 significant fibrosis

Study	Index test	Reference standard	Population	Target condition
	PHT included a nodular liver with splenomegaly.		<ul style="list-style-type: none"> abnormal liver US findings (abnormal echogenicity or a nodular edge) 	
Lindblad 1999 (Sweden) Prospective cohort study	<ul style="list-style-type: none"> Ultrasonography characterized as normal or pathological (increased and/or irregular echogenicity) Liver function test included serum activities of alanine transaminase (ALT), aspartate transaminase (AST), and g-glutamyltransferase (gGT) (with an upper reference level of ,0.8, ,0.8, and ,0.5 μkatal/l, respectively). Combined US and LFT 	<ul style="list-style-type: none"> Liver biopsy performed under general anesthesia in patients younger than 16 years and under local anesthesia in older patients. The biopsies were evaluated regarding fibrosis (normal; slight, enlarged portal zones; moderate, tendency towards septa formation; severe, bridging fibrosis; and cirrhosis, complete septa with regenerative noduli). Steatosis, bile duct proliferation, and inflammation were classified as absent, slight, moderate, or severe. A minimum of 4 portal zones were evaluated in each biopsy. Definitions: Biochemical liver disease (BLD) was defined as elevation above the upper reference level of any serum liver enzyme included in the LFT for at least 2 consecutive years in patients 4 years of age or older. A patient was thereafter classified as BLD even if LFT results were later normalized. Clinical liver disease was defined as multilobular cirrhosis (MLC) and always included clinical (hepato)splenomegaly with esophageal varices or signs of hypersplenism and biopsy-proven cirrhosis. All other patients were classified as having no liver disease (NLD). 	<p>N=41 children, young people and adults with CF</p> <ul style="list-style-type: none"> Median age: 19 years, range 5 to 43 years Participants were cared for at the Stockholm CF center and attended the center 2 or more times between 1976 and 1993 and who received biopsy in 1989-1993. 	<ul style="list-style-type: none"> Moderate or severe fibrosis and cirrhosis and/or moderate to severe steatosis Moderate or severe fibrosis and cirrhosis
Mueller-Abt 2008	<ul style="list-style-type: none"> US scans obtained after a 	Percutaneous liver biopsy using ultrasound	N= 30 infants, children and	<ul style="list-style-type: none"> Liver disease

Study	Index test	Reference standard	Population	Target condition
(Australia) Retrospective cohort study	4-hour fast in children under 2 years and a 6-hour fast in children over 2 years. There were 3 categories on the summary interpretation of findings: normal, indeterminate (suggestion of liver disease but no definite signs of cirrhosis) and cirrhosis. Increased hepatic echogenicity, heterogeneity and/or increased attenuation in the absence of nodularity of the liver surface were classified as indeterminate. Splenomegaly as an isolated finding was also regarded as indeterminate. All patients with nodularity of the liver surface were classified as cirrhosis.	guidance. Two samples, to limit sampling error, were obtained from the right lobe, at least 6 portal tracts were available for analysis. Scheuer-Score of 0 was regarded as normal, a score of 1–2 as mild to moderate reversible periportal changes and 3–4 was assessed as definite fibrosis/cirrhosis.	young people with CF. <ul style="list-style-type: none"> • Age range 11 months to 17 years. • Participants were attending the CF clinic with evidence of 2 out of 3 of the following criteria: biochemical liver disease (persistent elevation of ALT above upper limit of normal over a period of at least 6 months), clinical hepatomegaly or hepatosplenomegaly or sonographic evidence of liver disease 	(includes cirrhosis) (F1-F4 fibrosis score) <ul style="list-style-type: none"> • Cirrhosis (F3-F4 fibrosis score)
Patriquin 1999 (Canada) Prospective cohort study	<ul style="list-style-type: none"> • Liver function tests included aminotransferase (AST), alanine aminotransferase (ALT), gamma glutamyltransferase (GGT). 	<ul style="list-style-type: none"> • US scans were obtained without sedation after a 4-hour fast in children aged 2–6 years and after an 8-hour fast in patients older than 6 years. US included a survey of the entire abdomen as well as a detailed examination of liver architecture. • US signs were interpreted as follows: hypoechogenicity with prominent portal tracks as edema, hyperechogenicity as steatosis, hyperechogenicity with increased attenuation and nodules within or at 	N=195 children, young people and adults with CF <ul style="list-style-type: none"> • Mean age: 8.5 years, range 1 to 23 years. • Participants were attending a CF clinic who underwent abdominal US and a standard set of liver function tests over 1 year. 	<ul style="list-style-type: none"> • CFLD (includes cirrhosis)

Study	Index test	Reference standard	Population	Target condition
		the edge of the liver as cirrhosis. Signs of portal hypertension also were sought and Doppler US used to assess presence and direction of blood flow and detection of esophageal varices. US signs were compared with liver fu		
Rath 2012 (Germany) Prospective cohort study	<ul style="list-style-type: none"> • Transient elastography (TE): Liver stiffness by TE was evaluated using the same FibroScan® device in all patients. Non-invasive measurements were on the right lobe of the liver through the intercostal space at a depth of 25 and 65 mm from skin surface. In children below 15 kg of weight the FibroScan® S probe was used. For each patient, the stiffness value was calculated as the median of ten successful measurements. TE was considered valid if 10 successful measurements with a success rate $\geq 60\%$ and an interquartile range $\leq 30\%$ of the median were obtained. Results are expressed in kilopascal (kPa). Total examination time was approximately 5 minutes per patient. 	<ul style="list-style-type: none"> • Diagnosis of CFLD was established according to recent guidelines if least 2 of the following conditions on at least 2 consecutive examinations spanning a one-year period were present: <ul style="list-style-type: none"> ○ (i) Hepatomegaly (liver span >2 cm below the costal margin on the medioclavicular line) confirmed by ultrasound, ○ (ii) 2 abnormal serum liver enzyme levels (ALT, AST, γGT $>$ ULN), ○ (iii) ultrasound abnormalities other than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular margins). • Diagnosis of PHT was based on clinical and lab data combined with sonographic or endoscopic signs of PHT. 	<p>N=70 adults with CF;</p> <ul style="list-style-type: none"> • n=32 without CFLD, mean age (SD) 32.3 (9.3); • n=29 with CFLD, mean age (SD) 30.6 (8.6); • n=9 with LD and portal hypertension, mean age (SD) 21.4 (1.7). 	<ul style="list-style-type: none"> • CFLD (includes cirrhosis)
Rath 2013	<ul style="list-style-type: none"> • Alkaline 	<ul style="list-style-type: none"> • Diagnosis of CFLD was 	N=45 children,	<ul style="list-style-type: none"> • CFLD

Study	Index test	Reference standard	Population	Target condition
(Germany) Prospective cohort study	<p>phosphatase (ALP)</p> <ul style="list-style-type: none"> • AST/Platelets-Ratio-Index (APRI) • Transient elastography (TE): Liver stiffness by TE was evaluated using the same FibroScan® (Echosens, Paris, France) device in all patients. Non-invasive measurements were performed on the right lobe of the liver through the intercostal space at a depth of 25 and 65 mm from skin surface. In children below 15 kg of weight the FibroScan® S probe was used. For each patient, the stiffness value was calculated as the median of ten successful measurements. TE was considered valid if 10 successful measurements with a success rate $\geq 60\%$ and an interquartile range $\leq 30\%$ of the median were obtained. Results are expressed in kilopascal (kPa). Total examination time was approximately 5 minutes per patient. 	<p>established according to recent guidelines if least 2 of the following conditions on at least 2 consecutive examinations spanning a 1-year period were present:</p> <ul style="list-style-type: none"> ○ (i) Hepatomegaly (liver span >2 cm below the costal margin on the medioclavicular line) confirmed by ultrasound, ○ (ii) 2 abnormal serum liver enzyme levels (ALT, AST, γGT $>$ ULN), ○ (iii) ultrasound abnormalities other than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular margins). 	<p>young people and adults with CF;</p> <ul style="list-style-type: none"> • n=28 without CFLD; mean age (SD) 21.4 (11.8) (5 to 50 years) • and n=17 with CFLD; mean age (SD) 29 (10.8) (14 to 47) • Setting not confirmed. 	<p>(includes cirrhosis)</p>
Sadler 2015 (Canada)	<ul style="list-style-type: none"> • Liver stiffness measurement by transient 	<ul style="list-style-type: none"> • Diagnosis of CFLD was established according to previously published 	<p>N=127 adults with CF</p> <ul style="list-style-type: none"> • Median age, 	<ul style="list-style-type: none"> • CFLD (includes cirrhosis)

Study	Index test	Reference standard	Population	Target condition
Prospective cohort study	<p>elastography (TE) using FibroScan® probe.</p> <ul style="list-style-type: none"> • Aspartate aminotransferase to Platelets-Ratio-Index (APRI) was calculated as (AST/upper limit of normal for AST) x (100/platelets (x10⁹/L)). 	<p>criteria if least 2 of the following conditions were present:</p> <ul style="list-style-type: none"> ○ (i) Hepatomegaly and/or splenomegaly confirmed by ultrasonography, ○ (ii) abnormal liver biochemistry consisting of elevated levels of any 2 of ALT, AST, or GGT, ○ (iii) ultrasound abnormalities other than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular margins, splenomegaly presence). 	<p>years (interquartile range): 27 (22 to 37)</p> <ul style="list-style-type: none"> • Participants were attending a CF clinic of Calgary and Southern Alberta. 	
Witters 2009 (Belgium) Prospective cohort study	<ul style="list-style-type: none"> • Transient elastography (TE): (FibroScan®) – Liver disease was defined as a result above the age-related upper limit of normal liver stiffness • Ultrasound: Normal ultrasound results (Williams score = 3). Liver disease was defined as a Williams score of ≥ 4 (i.e. intermediate coarse to irregular liver parenchyma, liver edge nodularity and/or moderate to severe periportal fibrosis). 	<ul style="list-style-type: none"> • The North-American cystic fibrosis foundation (CFF) consensus workgroup Definition of CFLD: the presence of either clinical or biochemical liver disease. • Clinical liver disease was defined as the presence of hepatomegaly or splenomegaly • Biochemical liver disease was defined as the elevation of 2 of these tests: Liver tests (AST, ALT, alkaline phosphatase, bilirubin and gamma-GT). All CF patients from January 1996 to July 2007 were studied and patients with persistently elevated liver tests were identified (3–6 months, 1.5 times age-dependent upper limit of normal). 	<p>N=66 children, young people and adults with CF</p> <ul style="list-style-type: none"> • Mean age (SD): 13.6 (7.8) • Participants were followed up in CF clinic at a university hospital 	<ul style="list-style-type: none"> • Clinical CFLD (includes cirrhosis) • Biochemical CFLD (includes cirrhosis) • CFF guideline defined (clinical or biochemical) CFLD (includes cirrhosis)
Woodruff 2016 (USA) Prospective cohort study	<ul style="list-style-type: none"> • Monitoring strategy based on the assessment of liver function tests. 	<ul style="list-style-type: none"> • Not relevant (prognostic study) 	<p>N=298 children with CF identified</p> <ul style="list-style-type: none"> • Mean age at diagnosis (SD): 3.8 weeks (2.4 to 5.7) 	<ul style="list-style-type: none"> • CFLD

Study	Index test	Reference standard	Population	Target condition
	<ul style="list-style-type: none"> Median follow-up: 7.23 years 			

1 *Abbreviations: ALT: alkaline phosphatase; AST: alanine aminotransferase; APRI: Aspartate aminotransferase to*
 2 *Platelets-Ratio-Index; CF: cystic fibrosis; CFF: cystic fibrosis consensus; CFLD: cystic fibrosis liver disease; cm:*
 3 *centimetres; GGT: gamma-glutamyl transpeptidase; LFT: liver function test; MRI: magnetic resonance; SD:*
 4 *standard deviation; TE: transient elastography; PHT: portal hypertension*

5 10.4.4 Clinical evidence profile

6 Summary clinical evidence profile tables are not applicable to this review. Please see clinical
 7 evidence GRADE profiles in Appendix J.

8 10.4.5 Economic evidence

9 No economic evaluations of tests to detect related liver disease in people with cystic fibrosis
 10 were identified in the literature search conducted for this guideline. Full details of the search
 11 and economic article selection flow chart can be found in Appendix E and F, respectively. To
 12 aid consideration of cost-effectiveness relevant resource and cost use data are presented in
 13 Appendix K.

14 10.4.6 Evidence statements

15 10.4.6.1 Review question 1. What is the diagnostic accuracy of tests to detect/ strategies to 16 detect early and late cystic fibrosis-related liver disease?

17 10.4.6.1.1 Target condition: cystic-fibrosis liver disease (includes cirrhosis)

18 Clinical examination (hepatomegaly, splenomegaly)

19 Test 1. Diagnostic accuracy of clinical liver examination (versus biopsy)

20 High quality evidence from 1 cohort study of 40 children with cystic fibrosis found that clinical
 21 liver examination was not useful for ruling in (no serious imprecision) or for ruling out (no
 22 serious imprecision) cystic fibrosis-related liver disease (F1-F4 fibrosis).

23 Sensitivity was 68 (95% CI: 61-77)* and specificity was 33 (95% CI: 10-65)*.

24 Liver function blood tests (AST, ALT, GGT, Alkaline phosphatase, bilirubin, albumin, 25 platelets and clotting) and indices based on these tests (eg APRI, Forn's score, INR 26 ratio)

27 Test 2. Diagnostic accuracy of ALT, AST and GGT (versus ultrasound)

28 High quality evidence was available from 1 cohort study of 195 children, young people and
 29 adults with cystic fibrosis that examined the use of ALT, AST and GGT at unspecified
 30 thresholds for detecting cystic fibrosis-related liver disease found that:

- 31 • ALT was not useful for ruling in (no serious imprecision) or ruling out (no serious
 32 imprecision) cystic fibrosis-related liver disease
- 33 • AST was not useful for ruling in (serious imprecision) or ruling out (no serious
 34 imprecision) cystic fibrosis-related liver disease
- 35 • GGT was moderately useful for ruling in cystic fibrosis-related liver disease (serious
 36 imprecision), but was not useful for ruling out cystic fibrosis-related liver disease (no
 37 serious imprecision).

1 Sensitivities were 63.6 (95% CI: 34.4-86.0)*, 47.4 (95% CI: 33.4-60.6)* and 50.0 (95% CI:
2 22.0-75.1)* respectively and specificities were 79.0 (95% CI: 75.3-82.2)* and 87.9 (95% CI:
3 84.5-91.1)* and 90.4 (95% CI: 87.1-93.4)* respectively.

4 Note: Ultrasound signs were interpreted as follows: hypoechogenicity with prominent portal
5 tracts as oedema, hyperechogenicity as steatosis, hyperechogenicity with increased
6 attenuation and nodules within or at the edge of the liver as cirrhosis. Signs of portal
7 hypertension also were sought and Doppler ultrasound used to assess presence and
8 direction of blood flow and detection of oesophageal varices.

9 **Test 3. Diagnostic accuracy of ALT, AST and GGT (versus biopsy)**

10 Moderate quality evidence from 1 cohort study of 41 children, young people and adults with
11 cystic fibrosis found that liver function testing (ALT, AST and GGT at thresholds of 0.8, 0.8
12 and 0.5 μ kata respectively):

- 13 • was not useful for ruling in (no serious imprecision) or for ruling out (serious imprecision)
14 moderate or severe fibrosis/cirrhosis with steatosis;
- 15 • was not useful for ruling in, but was very useful (no serious imprecision) for ruling out
16 (serious imprecision) moderate or severe fibrosis/cirrhosis.

17 Sensitivities were 83 (95% CI: 68-94)* and 100 (95% CI: 78-100)* and specificities were 44
18 (95% CI: 26-58)* and 44 (95% CI: 33-44)* respectively.

19 **Test 4. Diagnostic accuracy of ALT testing (versus biopsy)**

20 Moderate quality evidence from 1 cohort study of 40 children and young people with cystic
21 fibrosis found that ALT testing was not useful for ruling in (very serious imprecision) or for
22 ruling out (no serious imprecision) cystic fibrosis-related liver disease (F1-F4 fibrosis).

23 Sensitivity was 30 (95% CI: 0-0.60)* and specificity was 98 (95% CI: 96-100)*.

24 **Test 5. Diagnostic accuracy of alkaline phosphatase (ALP) (versus practice guideline 25 cystic fibrosis-related liver disease definitions)**

26 Moderate quality evidence from 1 cohort study of 45 children, young people and adults with
27 cystic fibrosis found that use of ALP (age and gender specific cutoffs) was not useful for
28 ruling in (serious imprecision) or for ruling out (serious imprecision) cystic fibrosis-related
29 liver disease.

30 Sensitivity was 70.6 (95% CI: 49.5-85.5)* and specificity was 82.1 (95% CI: 69.3-91.2)*

31 Note: Practice guideline definitions included criteria for clinical, biochemical and ultrasound
32 testing.

33 **Test 6. Diagnostic accuracy of APRI (versus practice guideline cystic fibrosis-related 34 liver disease definitions)**

35 Low, moderate and high quality evidence from 3 cohort studies of children, young people and
36 adults with cystic fibrosis (n=45, 55 and 122) examined the use of APRI at various thresholds
37 to detect cystic fibrosis-related liver disease.

- 38 • APRI thresholds of 0.133 in children and adults and of 0.4 and 0.5 in adults were
39 moderately useful for ruling in (very serious imprecision), but not useful for ruling out (no
40 serious imprecision) cystic fibrosis-related liver disease
 - 41 ○ sensitivities were 47.1 (95% CI: 28.2-56.7)*, 50 (95% CI: 29-69)* and 50 (95% CI: 29-
42 68)* and specificities were 93.1 (95% CI: 82.0-98.7)*, 92 (95% CI: 88-95)* and 94 (95%
43 CI: 90-97)* respectively.

- 1 • An APRI threshold of 0.231 in adults was not useful for ruling in (no serious imprecision),
2 but was moderately useful for ruling out (very serious imprecision) cystic fibrosis-related
3 liver disease
4 ○ sensitivity was 85.7 (95% CI: 60-97.4)* and specificity 70.7 (95% CI: 62.0-74.7)*.

5 Note: Practice guideline definitions included criteria for clinical, biochemical and ultrasound
6 testing.

7 **Test 7. Diagnostic accuracy of Forn's score (versus practice guideline cystic fibrosis-**
8 **related liver disease definitions)**

9 Moderate quality evidence from 1 cohort study of 55 adults with cystic fibrosis found that use
10 of Forn's score at a threshold of 2.154 was not useful for ruling in (no serious imprecision)
11 but was moderately useful for ruling out (very serious imprecision) cystic fibrosis-related liver
12 disease.

13 Sensitivity was 92.9 (95% CI: 67.8-99.6)* and specificity was 61.0 (95% CI: 52.4-63.3)*.

14 Note: Practice guideline definitions included criteria for clinical, biochemical and ultrasound
15 testing.

16 **Imaging techniques**

17 **Test 8. Diagnostic accuracy of ultrasound (versus clinical cystic fibrosis-related liver**
18 **disease definition)**

19 Low quality evidence from 1 cohort study of 66 children, young people and adults with cystic
20 fibrosis found that use of ultrasound at a threshold of Williams score ≥ 4 was not a useful test
21 for ruling in (no serious imprecision) or for ruling out (no serious imprecision) cystic fibrosis-
22 related liver disease.

23 Sensitivity was 66.7 (95% CI: 25.0-93.9)* and specificity was 66.7 (95% CI: 62.5-69.4)*.

24 Note: Diagnosis of cystic fibrosis-related liver disease according to the presence or absence
25 of hepatomegaly or splenomegaly determined by clinical examination.

26 **Test 9. Diagnostic accuracy of ultrasound (versus biochemical definition)**

27 Low quality evidence from 1 cohort study of 66 children, young people and adults with cystic
28 fibrosis found that use of ultrasound at a threshold of Williams score ≥ 4 was not useful for
29 ruling in (no serious imprecision) or ruling out (no serious imprecision) cystic fibrosis-related
30 liver disease.

31 Sensitivity was 50.0 (95% CI: 14.3-85.6)* and specificity was 66.7 (95% CI: 63.1-70.2)*.

32 Note: Diagnosis of cystic fibrosis-related liver disease was defined as persistently elevated
33 results (3–6 months, 1.5 times age-dependent upper limit of normal) for 2 of these liver tests:
34 AST, ALT, alkaline phosphatase, bilirubin and gamma-GT.

35 **Test 10. Diagnostic accuracy of ultrasound (versus clinical and or biochemical**
36 **definition)**

37 Moderate quality evidence from 1 cohort study of 70 infants, children, young people and
38 adults with cystic fibrosis found that use that use of ultrasound at a threshold of Williams
39 score ≥ 4 was moderately useful for ruling in (very serious imprecision), but not for ruling out
40 (no serious imprecision) cystic fibrosis-related liver disease.

1 However moderate quality evidence from another cohort study of 66 children, young people
2 and adults with cystic fibrosis found that use of ultrasound at a threshold of Williams score ≥ 4
3 was not useful for ruling in (no serious imprecision) and for ruling out (no serious imprecision)
4 cystic fibrosis-related liver disease.

5 Sensitivities were 50.0 (95% CI: 22.0-75.1)* and 63.6 (95% CI: 33.6-87.0)* and specificities
6 were 91.7 (95% CI: 87.0-95.8)* and 70.9 (95% CI: 64.9-75.6)* respectively.

7 Note: Diagnosis of cystic fibrosis-related liver disease was defined using clinical and
8 biochemical criteria.

9 **Test 11. Diagnostic accuracy of ultrasound (versus biopsy)**

10 High, moderate and low quality evidence was available from 3 cohort studies of infants,
11 children, young people and adults with cystic fibrosis (n=30, 40 and 41) that examined
12 ultrasound imaging.

- 13 • Ultrasound imaging was not useful for ruling in (serious imprecision) or for ruling out
14 (serious imprecision) F1-F4 fibrosis in children
 - 15 ○ sensitivities were 81 (95% CI: 73-89)* and 65 (95% CI: 55-74)* and specificities were
16 44 (95% CI: 17-73)* and 57 (95% CI: 22-87)* respectively (high quality evidence from 2
17 studies).
- 18 • Ultrasound imaging was not useful for ruling in (no serious imprecision) or ruling out (no
19 serious imprecision) moderate or severe fibrosis/cirrhosis with steatosis
 - 20 ○ sensitivity was 70 (95% CI: 54-80)* and specificity was 78 (95% CI: 58-92)* (moderate
21 quality evidence from 1 study).
- 22 • Ultrasound testing was not useful for ruling in (no serious imprecision), but moderately
23 useful for ruling out (very serious imprecision) moderate or severe fibrosis/cirrhosis
24 without steatosis
 - 25 ○ sensitivity was 86 (95% CI: 61-97)* and specificity was 70 (95% CI: 58-76)* (low quality
26 evidence from 1 study).

27 **Test 12. Diagnostic accuracy of MRI (versus liver function tests or abnormal 28 ultrasound)**

29 Moderate quality evidence from 1 cohort study of 23 adults with cystic fibrosis found that use
30 of MRI to detect 1 abnormal sign was not useful to rule in (very serious imprecision) or to rule
31 out (no serious imprecision) cystic fibrosis-related liver disease.

32 Sensitivity 36.4 (95% CI: 14.7-51.1) and specificity was 83.3 (95% CI: 63.5-96.8)*.

33 Note: details of the diagnosis of cystic fibrosis-related liver disease according to liver function
34 tests or abnormal ultrasound were not reported in the study.

35 **Liver stiffness measurement (transient elastography or FibroScan®)**

36 **Test 13. Diagnostic accuracy of transient elastography (versus clinical cystic fibrosis- 37 related liver disease definition)**

38 Low quality evidence from 1 cohort study of 66 children, young people and adults with cystic
39 fibrosis found that use of transient elastography at a threshold of 5.63kPa for <12 years and
40 6.50kPa for ≥ 12 years was moderately useful for ruling in (serious imprecision) and for ruling
41 out (very serious imprecision) cystic fibrosis-related liver disease.

42 Sensitivity was 83.3 (95% CI: 38.7-99.1)* and specificity was 85.0 (95% CI: 80.5-86.6)*

1 Note: Diagnosis of cystic fibrosis-related liver disease according to the presence or absence
2 of hepatomegaly or splenomegaly determined by clinical examination.

3 **Test 14. Diagnostic accuracy of transient elastography (versus biochemical definition)**

4 Low quality evidence from 1 cohort study of 66 children, young people and adults with cystic
5 fibrosis found that use of transient elastography at a threshold of 5.63kPa for <12 years and
6 6.50kPa for ≥12 years was not a useful test for ruling in (serious imprecision) or ruling out
7 (serious imprecision) cystic fibrosis-related liver disease.

8 Sensitivity was 50.0 (95% CI: 14.5-85.3)* and specificity was 83.3 (95% CI: 79.8-86.9).

9 Note: Diagnosis of cystic fibrosis-related liver disease was defined as persistently elevated
10 results (3–6 months, 1.5 times age-dependent upper limit of normal) for 2 of these liver tests:
11 AST, ALT, alkaline phosphatase, bilirubin and gamma-GT.

12 **Test 15. Diagnostic accuracy of transient elastography (versus clinical and or**
13 **biochemical definition)**

14 Moderate quality evidence from 1 cohort study of 66 children, young people and adults with
15 cystic fibrosis found that use of transient elastography at a threshold of 5.63kPa for <12
16 years and 6.50kPa for ≥12 years was moderately useful for ruling in (very serious
17 imprecision) but not useful for ruling (serious imprecision).

18 Sensitivity was 63.6 (95% CI: 34.4-86.0)* and specificity was 87.3 (95% CI: 81.4-91.8)*.

19 Note: Diagnosis of cystic fibrosis-related liver disease was defined using clinical and
20 biochemical criteria.

21 **Test 16. Diagnostic accuracy of transient elastography (versus practice guideline**
22 **cystic fibrosis-related liver disease definition)**

23 High quality evidence from 1 cohort study (n=136; 75 children and 61 adults) found that use
24 of transient elastography at a threshold of 5.5kPa was not a useful test for ruling in (serious
25 imprecision) or ruling out (no serious imprecision) cystic fibrosis-related liver disease.

26 Sensitivities were 53.3 (95% CI 43.2 to 61.2) and 55.2 (95% CI 40.7 to 66.8), and
27 specificities were 76.7 (95% CI 61.4 to 88.4) and 78.1 (95% CI 60.0 to 88.7) in children and
28 adults respectively.

29 Note: Diagnosis of cystic fibrosis-related liver disease was established according to
30 published guidelines (Debray 2011) if least 2 of the following conditions on at least 2
31 consecutive examinations spanning a 1-year period were present: (i) Hepatomegaly (liver
32 span >2 cm below the costal margin on the medioclavicular line) confirmed by ultrasound, (ii)
33 2 abnormal serum liver enzyme levels (ALT, AST, γGT > ULN), (iii) ultrasound abnormalities
34 other than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular
35 margins).

36 **Test 17. Diagnostic accuracy of transient elastography (versus practice guideline**
37 **cystic fibrosis-related liver disease definitions)**

38 Low, moderate and high quality evidence from 3 cohort studies and 1 case control study of
39 children, young people and adults with cystic fibrosis (n=45, 49, 127 and 50) was available
40 that examined the use of transient elastography at various thresholds to detect cystic
41 fibrosis-related liver disease as defined by practice guidelines.

- 42 • At a threshold of 3.7 transient elastography was not useful for ruling in (no serious
43 imprecision) or for ruling out (very serious imprecision) cystic fibrosis-related liver disease
44 in a population of adults

- 1 ○ sensitivity was 89 (95% CI: 66-98)* and specificity was 37 (95% CI: 33-38)*.
- 2 ● At a threshold of 5.3 kPa transient elastography was not useful for ruling in (serious
3 imprecision) or for ruling out (serious imprecision) cystic fibrosis-related liver disease in a
4 population of adults
- 5 ○ sensitivity was 67 (95% CI: 43-85)* and specificity was 83 (95% CI: 79-86)*.
- 6 ● At a threshold of 5.9kPa transient elastography was very useful (very serious imprecision)
7 for ruling in but not useful for ruling out (no serious imprecision) cystic fibrosis-related liver
8 disease in a population of adults
- 9 ○ sensitivity was 42.9 (95% CI: 2s.6-49.6)* and specificity was 97.1 (95% CI: 89.0-99.8)*.
- 10 ● At a threshold of 6.0 kPa, transient elastography was moderately useful (very serious
11 imprecision) for ruling in but not useful for ruling out (no serious imprecision) cystic
12 fibrosis-related liver disease in a population of adults
- 13 ○ sensitivity was 56 (95% CI: 34-75)* and specificity was 91 (95% CI: 87-94)*.
- 14 ● At a threshold of 6.3kPa, transient elastography was very useful for ruling in (very serious
15 imprecision, but at least moderately useful) and moderately useful (serious imprecision)
16 for ruling out cystic fibrosis-related liver disease in a population of children, young people
17 and adults
- 18 ○ sensitivity was 82.4 (95% CI: 64.2-85.3)* and specificity was 98.2 (95% CI: 87.4-100)*.
- 19 ● At a threshold of 6.8kPa, transient elastography was moderately useful for ruling in (very
20 very serious imprecision) but not useful for ruling out (serious imprecision) cystic fibrosis-
21 related liver disease in a population of adults
- 22 ○ sensitivity was 76 (95% CI: 61.6-82.5)* and specificity was 92 (95% CI: 77.6-98.5)*.
- 23 ● Note: Practice guideline definitions included criteria for clinical, biochemical and
24 ultrasound testing.

25 **Test 18. Diagnostic accuracy of transient elastography (versus liver function tests or**
26 **abnormal ultrasound)**

27 Very low quality evidence from 1 cohort study of 23 adults with cystic fibrosis found that use
28 of transient elastography to detect F2-F4 fibrosis was not useful to rule in (serious
29 imprecision) or to rule out (very serious imprecision) cystic fibrosis-related liver disease.
30 Sensitivity was 75 (95% CI: 24.2-98.6)* and specificity was 84.2 (95% CI: 73.5-69.2)*.

31 Note: details of the diagnosis of cystic fibrosis-related liver disease according to liver function
32 tests or abnormal ultrasound were not reported in the study.

33 **Combination of tests**

34 **Test 19. Diagnostic accuracy of ALT, AST and GGT + ultrasound (versus biopsy)**

35 Moderate and low quality evidence from 1 cohort study of 41 children, young people and
36 adults with cystic fibrosis found that combined use of liver function testing (ALT, AST and
37 GGT at thresholds of 0.8, 0.8 and 0.5 μ kata/ respectively) and ultrasound imaging

- 38 ● was not useful for ruling in (serious imprecision) or ruling out (no serious imprecision)
39 moderate or severe fibrosis/cirrhosis with steatosis
- 40 ○ sensitivity was 65 (95% CI: 50-76)* and specificity was 78 (95% CI: 58-92)*.
- 41 ● was not useful for ruling in (serious imprecision) but were moderately useful (very serious
42 imprecision) for ruling out in moderate or severe fibrosis/cirrhosis without steatosis
- 43 ○ sensitivity was 86 (95% CI: 62-97)* and specificity was 74 (95% CI: 62-80)*.

1 **Test 20. Diagnostic accuracy of clinical liver examination + ALT + ultrasound imaging**
2 **(versus biopsy)**

- 3 High and moderate quality evidence from 1 cohort study of 40 children with cystic fibrosis
4 found that combined use of clinical liver examination, ALT testing and ultrasound imaging:
- 5 • was not useful for ruling in (serious imprecision) or for ruling out (very serious imprecision)
6 cystic fibrosis-related liver disease (F1-F4 fibrosis)
 - 7 ○ sensitivity was 97 (95% CI: 85-100)* and specificity was 13 (95% CI: 4-15)*.
 - 8 • was not useful for ruling in (serious imprecision) or for ruling out (very serious imprecision)
9 cystic fibrosis-related liver disease (F2-F4 significant fibrosis)
 - 10 ○ sensitivity was 82 (95% CI: 62-95)* and specificity was 48 (95% CI: 33-57)*.

11 **110.4.6.1.2 Target condition: cirrhosis only**

12 **Clinical examination (hepatomegaly, splenomegaly)**

13 No evidence was found.

14 **Liver function blood tests (AST, ALT, GGT, Alkaline phosphatase, bilirubin, albumin,**
15 **platelets and clotting) and indices based on these tests (eg APRI, Forn's score, INR**
16 **ratio)**

17 **Test 1. Diagnostic accuracy of APRI (versus clinical examination + ultrasound)**

18 Low quality evidence from 1 cohort study of 14 adults with cystic fibrosis and cystic fibrosis-
19 related liver disease found that APRI at a threshold of 0.334 was moderately useful for ruling
20 in (very serious imprecision) and moderately useful for ruling out (very serious imprecision)
21 cirrhosis.

22 Sensitivity was 83.3.7 (95% CI: 45.0-98.5) and specificity was 87.5 (95% CI: 58.8-98.9).

23 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
24 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
25 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
26 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
27 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
28 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and
29 rarefication of peripheral portal veins) and clinical signs (for example esophageal varices,
30 splenomegaly).

31 **Test 2. Diagnostic accuracy of Forns score (versus clinical examination + ultrasound)**

32 Low quality evidence from 1 cohort study of 14 adults with cystic fibrosis and cystic fibrosis-
33 related liver disease found that Forns's score at a threshold of 4.059 was very useful for ruling
34 in (very serious imprecision) and not useful for ruling out (no serious imprecision) cirrhosis.

35 Sensitivity was 66.7 (95% CI: 30.1-75.0) and specificity was 94.1 (95% CI: 68.3-100).

36 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
37 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
38 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
39 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
40 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
41 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and
42 rarefication of peripheral portal veins) and clinical signs (for example esophageal varices,
43 splenomegaly).

1 **Imaging techniques**

2 **Test 3. Diagnostic accuracy of ultrasound (versus biopsy)**

3 Moderate quality evidence from 1 cohort study of 30 infants, children and young people with
4 cystic fibrosis found that ultrasound was a moderately useful test to rule in (very serious
5 imprecision) but not useful to rule out (no serious imprecision) fibrosis.

6 Sensitivity was 0.57 (95% CI: 0.36-0.64)* and specificity was 0.94 (95% CI: 0.75-1.00)*.

7 **Liver stiffness measurement (transient elastography or FibroScan®)**

8 **Test 4. Diagnostic accuracy of transient elastography (versus clinical examination +**
9 **ultrasound)**

10 Low quality evidence from 1 cohort study of 14 adults with cystic fibrosis and cystic fibrosis-
11 related liver disease found that transient elastography at a threshold of 4.4kPa was not
12 useful to rule in (serious imprecision), but was very useful to rule out (very serious
13 imprecision) cirrhosis.

14 Sensitivity was 92.3 (95% CI: 56.2-100)* and specificity was 75 (95% CI: 45.7-81.2)*.

15 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
16 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
17 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
18 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
19 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
20 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and
21 rarefaction of peripheral portal veins) and clinical signs (for example esophageal varices,
22 splenomegaly).

23 **Combination of tests**

24 No evidence was found.

25 **10.4.6.2 Target condition: portal hypertension**

26 **Clinical examination (hepatomegaly, splenomegaly)**

27 No evidence was found.

28 **Liver function blood tests (AST, ALT, GGT, Alkaline phosphatase, bilirubin, albumin,**
29 **platelets and clotting) and indices based on these tests (eg APRI, Forns score, INR**
30 **ratio)**

31 **Test 1. Diagnostic accuracy of APRI (versus clinical examination)**

32 Low quality evidence was available from 1 case control study of 50 adults with cystic fibrosis
33 (25 with cystic fibrosis-related liver disease and 25 without cystic fibrosis-related liver
34 disease) found that APRI at a threshold of 0.49 was very useful for ruling in (very serious
35 imprecision) and moderately useful for ruling out (very serious imprecision) portal
36 hypertension in the whole adult population and in those with cystic fibrosis-related liver
37 disease.

38 Sensitivities were 87.5 (95% CI: 52.0-99.3)* and 87.5 (95% CI: 54.8-98.9)* respectively and
39 specificities were 92.9 (95% CI: 86.1-95.1)*and 94.1 (95% CI: 78.7-99.5)*respectively.

1 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
2 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
3 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
4 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
5 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
6 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and
7 rarefaction of peripheral portal veins) and clinical signs (for example esophageal varices,
8 splenomegaly). Portal hypertension: platelet count <140x10⁹/L, splenomegaly, presence of
9 porto-systemic collateral veins, portal diameter >13mm, or ascites.

10 **Test 2. Diagnostic accuracy of Forns score (versus clinical examination)**

11 Low quality evidence was available from 1 case control study of 50 adults with cystic fibrosis
12 (25 with cystic fibrosis-related liver disease and 25 without cystic fibrosis-related liver
13 disease) found that Forns' score at a threshold of 0.68 was moderately useful for ruling in
14 (serious imprecision) and moderately useful for ruling out (very serious imprecision) portal
15 hypertension in the whole adult population and in those with cystic fibrosis-related liver
16 disease.

17 Sensitivities were 87.5 (95% CI: 53.2-99.3)*and 87.5 (95% CI: 50.7-99.3)* respectively and
18 specificities were 82.4 (95% CI: 66.2-87.9)*and 85.7 (95% CI: 78.7-88.0)* respectively.

19 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
20 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
21 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
22 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
23 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
24 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and
25 rarefaction of peripheral portal veins) and clinical signs (for example esophageal varices,
26 splenomegaly). Portal hypertension: platelet count <140x10⁹/L, splenomegaly, presence of
27 porto-systemic collateral veins, portal diameter >13mm, or ascites.

28 **Imaging techniques**

29 No evidence was found.

30 **Liver stiffness measurement (transient elastography or FibroScan®)**

31 **Test 3. Diagnostic accuracy of transient elastography (versus clinical examination)**

32 Low quality evidence was available from 1 case control study of 50 adults with cystic fibrosis
33 (25 with cystic fibrosis-related liver disease and 25 without cystic fibrosis-related liver
34 disease) found that transient elastography at a threshold of 8.9kPa was moderately useful
35 (very serious imprecision) for ruling in portal hypertension in the whole adult population but
36 was not useful (serious imprecision) in those with cystic fibrosis-related liver disease. This
37 threshold was moderately useful to rule out (very serious imprecision) portal hypertension in
38 the whole adult population and in those with cystic fibrosis-related liver disease.

39 Sensitivities were 87.5 (95% CI: 51.4-99.3)*and 87.5 (95% CI: 52.9-99.3)* respectively and
40 specificities were 90.5 (95% CI: 83.6-92.7)* and 76.5 (95% CI: 60.2-82.0)* respectively.

41 Note: Diagnosis of cystic fibrosis-related liver disease (Sokol 1999, Colombo 2002) if at least
42 2 of the following conditions present on at least 2 consecutive examinations spanning a 1-
43 year period: (1) Ultrasound confirmed hepatomegaly;(2) elevated serum liver enzyme levels
44 of ALT, AST, AP, or GGT;(3) ultrasound abnormalities other than hepatomegaly (increased,
45 heterogeneous echogenicity, nodularity, irregular margins, splenomegaly). Liver cirrhosis:
46 distinct ultrasonographic signs (coarse nodularity, presence of portal hypertension and

1 rarefication of peripheral portal veins) and clinical signs (for example esophageal varices,
2 splenomegaly). Portal hypertension: platelet count $<140 \times 10^9/L$, splenomegaly, presence of
3 porto-systemic collateral veins, portal diameter $>13\text{mm}$, or ascites.

4 **Test 4. Diagnostic accuracy of transient elastography (versus biochemical and**
5 **imaging)**

6 High quality evidence was available from 1 cohort study of 70 adults with cystic fibrosis found
7 that transient elastography at a threshold of 11.5kPa was very useful for ruling in (serious
8 imprecision, but at least moderately useful) but not for ruling out (no serious imprecision)
9 portal hypertension.

10 Sensitivity was 66.7 (95% CI: 36.2-77.2)* and specificity was 98.4 (95% CI: 93.9-99.9)*.

11 Note: Diagnosis of cystic fibrosis-related liver disease was established according to
12 published guidelines (Debray 2011) if least 2 of the following conditions on at least 2
13 consecutive examinations spanning a 1-year period were present: (i) Hepatomegaly (liver
14 span $>2\text{ cm}$ below the costal margin on the medioclavicular line) confirmed by ultrasound, (ii)
15 2 abnormal serum liver enzyme levels (ALT, AST, $\gamma\text{GT} > \text{ULN}$), (iii) ultrasound abnormalities
16 other than hepatomegaly (increased, heterogeneous echogenicity, nodularity, irregular
17 margins). Diagnosis of portal hypertension was based on clinical and lab data combined with
18 sonographic or endoscopic signs of portal hypertension (defined splenomegaly, increased
19 portal vein pressure in duplex Doppler sonography, platelet count $150,000/\text{mm}^3$,
20 oesophageal varices or other signs of portal hypertension on
21 oesophagogastrroduodenoscopy.

22 **Combination of tests**

23 No evidence was found.

24 **240.4.6.2.1 Target condition: oesophageal varices**

25 **Clinical examination (hepatomegaly, splenomegaly)**

26 No evidence was found.

27 **Liver function blood tests (AST, ALT, GGT, Alkaline phosphatase, bilirubin, albumin,**
28 **platelets and clotting) and indices based on these tests (eg APRI, Forn's score, INR**
29 **ratio)**

30 **Test 1. Diagnostic accuracy of APRI (versus published definition)**

31 Low quality evidence was available from 1 case control study of 23 adults with cystic fibrosis
32 (13 with cystic fibrosis-related liver disease, and 10 without cystic fibrosis-related liver
33 disease) found that APRI at a threshold of 0.49 was very useful for ruling in (very serious
34 imprecision) and very useful for ruling out (very serious imprecision) oesophageal varices in
35 the whole adult population and in those with cystic fibrosis-related liver disease.

36 Sensitivities were 100 (95% CI: 60.0-100)* and 100 (95% CI: 62.9-100)* respectively and
37 specificities were 94.1(95% CI: 80.0-94.1)* and 93.3(95% CI: 63.7-93.3)* respectively.

38 **Test 2. Diagnostic accuracy of Forn's score (versus published definition)**

39 Very low quality evidence was available from 1 case control study of 23 adults with cystic
40 fibrosis (13 with cystic fibrosis-related liver disease, and 10 without cystic fibrosis-related liver
41 disease) found that Forn's score at a threshold of 0.68 was moderately useful for ruling in

1 (serious imprecision) and very useful for ruling out (very serious imprecision) oesophageal
2 varices in the whole adults population and in those with cystic fibrosis-related liver disease.

3 Sensitivities were 100 (95% CI: 62.9-100)* and 100 (95% CI: 58.9-100)* respectively and
4 specificities were 85.7 (95% CI: 53.9-85.7)* and 88.2 (95% CI: 73.7-88.2)* respectively.

5 **Imaging techniques**

6 No evidence was found.

7 **Liver stiffness measurement (transient elastography or FibroScan®)**

8 **Test 3. Diagnostic accuracy of transient elastography (versus published definition)**

9 Very low quality evidence was available from 1 case control study of 23 adults with cystic
10 fibrosis (13 with cystic fibrosis-related liver disease, and 10 without cystic fibrosis-related liver
11 disease) found that transient elastography at a threshold of 8.9kPa was not useful to rule in
12 (no serious imprecision), but was very useful to rule out cirrhosis (very serious imprecision)
13 oesophageal varices in adults.

14 Sensitivity was 100 (95% CI: 57.8-100)* and specificity was 76.5 (95% CI: 61.6-76.5)*.

15 **Combination of tests**

16 No evidence was found.

17 **10.4.6.3 Review question 2. What is the diagnostic and prognostic value of different strategies 18 to detect cystic fibrosis-related liver disease and predict progression (including 19 progression to cirrhosis and portal hypertension without oesophageal varices)?**

20 **10.4.6.3.1 Strategy 1. Clinical examination (hepatomegaly, splenomegaly)**

21 **Prognosis of liver disease**

22 No evidence was identified.

23 **Prognosis of cirrhosis**

24 No evidence was identified.

25 **Prognosis of portal hypertension**

26 No evidence was identified.

27 **10.4.6.3.2 Strategy 2. Liver function blood tests (AST, ALT, GGT, Alkaline phosphatase, bilirubin, 28 albumin, platelets and clotting) and indices based on these tests (eg APRI, Forns 29 score, INR ratio)**

30 **Prognosis of liver disease**

31 High quality evidence from 1 prospective cohort study of 298 children with cystic fibrosis
32 described a significant association between:

- 33 • an elevated AST ≥ 1.5 ULN
- 34 • an elevated GGTP ≥ 1.5 ULN

35 and the risk of developing cystic fibrosis-related liver disease at a median of 7 years follow-
36 up.

1 However, no significant association was described for the following:

- 2 • an elevated AST \geq 2.0 ULN
3 • an elevated ALT \geq 1.5 ULN
4 • an elevated ALT \geq 2.0 ULN
5 • an elevated GGTP \geq 2.0 ULN.

6 **Prognosis of cirrhosis**

7 No evidence was identified.

8 **Prognosis of portal hypertension**

9 No evidence was identified.

10 **100.4.6.3.3 Strategy 3. Imaging techniques**

11 **Prognosis of liver disease**

12 No evidence was identified.

13 **Prognosis of cirrhosis**

14 No evidence was identified.

15 **Prognosis of portal hypertension**

16 No evidence was identified.

17 **170.4.6.3.4 Strategy 4. Liver stiffness measurement - transient elastography (FibroScan®)**

18 **Prognosis of liver disease**

19 No evidence was identified.

20 **Prognosis of cirrhosis**

21 No evidence was identified.

22 **Prognosis of portal hypertension**

23 No evidence was identified.

24 **240.4.6.3.5 Strategy 5. Combination of different strategies**

25 **Prognosis of liver disease**

26 **Combination of transient elastography and biopsy**

27 Low quality evidence from 1 case control study of 50 adults with cystic fibrosis set in a cystic
28 fibrosis referral centre described a significant association between increasing liver stiffness
29 and increasing risk of developing cystic fibrosis-related liver disease (adjOR: 2.74 (95% CI
30 1.53-4.89, $p=0.001$).

31 **Prognosis of cirrhosis**

32 No evidence was identified

1 **Prognosis of portal hypertension**

2 **Combination of transient elastography and biopsy**

3 High quality evidence from 1 cohort study of 40 children and young people with cystic fibrosis
4 set in a cystic fibrosis clinic in a city hospital described a significant association between
5 increasing fibrosis stage on biopsy and increasing risk of developing portal hypertension in
6 adults from birth (adjHR: 3.9 (p<0.001, no 95% CI given) and from the time of biopsy (adjHR:
7 3.4 (p<0.002, no 95% CI given).

8 **10.4.6.4 Economic evidence statements**

9 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

10 **10.4.7 Evidence to recommendations**

11 **10.4.7.1 Relative value placed on the outcomes considered**

12 The aim of this review was to assess the diagnostic accuracy of different diagnostic
13 strategies to detect cystic fibrosis-related liver disease (including cirrhosis, portal
14 hypertension and oesophageal varices) defined by gold standard tests and to identify
15 whether any tests are useful in predicting the progression of cystic fibrosis-related liver
16 disease.

17 As sensitivity and specificity reflect patient outcomes these were considered critical
18 outcomes for this review. Consequences for people with cystic fibrosis following diagnosis
19 were also considered by the Committee. Likelihood ratios were also considered to be critical
20 diagnostic outcomes because they provide information about a test's usefulness in assisting
21 the healthcare professional to make a diagnosis. AUROC diagnostic data (which inform the
22 best thresholds for defining disease) were regarded as important but not critical to the
23 review.

24 Sensitivity was used to evaluate imprecision for the detection of cystic fibrosis-related liver
25 disease (including cirrhosis). Early accurate identification of disease might be important
26 because there was a possibility that early use of ursodeoxycholic acid treatment might
27 prevent liver disease progression. Specificity was used to evaluate imprecision for the
28 detection of portal hypertension or oesophageal varices (because accurate identification of
29 those without disease would avoid unnecessary referral to specialist hepatology centres and
30 unnecessary investigation for this complication of advanced cystic fibrosis-related liver
31 disease.

32 Adjusted hazard ratios (adjHRs) and odds ratios (ORs) were regarded as critical prognostic
33 outcomes for assessing the risk/probability of developing clinically significant liver disease
34 (cirrhosis, portal hypertension, oesophageal varices) in the future based on current available
35 information. Estimates that were not adjusted for confounders were excluded from the
36 review.

37 **10.4.7.2 Consideration of clinical benefits and harms**

38 The Committee agreed that their priority was early detection and treatment of liver disease in
39 patients with cystic fibrosis as this may prevent further damage and, in some cases, may be
40 reversible. They agreed that irreversible liver disease would be under liver specialist
41 management and was out with the scope of this guideline.

42 The Committee discussed the importance of reducing the likelihood of a false negative
43 diagnosis which could result in the withholding of ursodeoxycholic acid treatment which was
44 potentially effective. The Committee agreed that the impact of a false negative diagnosis

1 would be mitigated by the slow progression of disease and that the subsequent annual
2 review would provide a further opportunity for identification. They noted that a false positive
3 result might lead to unnecessary treatment and have a negative psychological impact.

4 It was noted that while neonates with cystic fibrosis can demonstrate transient liver
5 dysfunction (manifested by a temporary derangement of liver function tests), any perceived
6 harm of starting unnecessary treatment is likely to be offset by the benefit of preventing
7 irreversible liver damage.

8 The Committee also discussed the relative benefits and harms associated with the different
9 diagnostic tests. Although the gold standard test for identification of liver pathology in cystic
10 fibrosis is histological examination of biopsy samples, this is rarely performed in practice as it
11 is invasive, may miss focal lesions and has associated risks for example with respect to
12 general anaesthesia or infection. CT scanning and MRI scanning are also used as gold
13 standards because of the high definition images that are obtained, but the evidence review
14 found no evidence for CT or MRI scanning as the reference standard. The Committee noted
15 that although these investigations are not invasive, they are expensive and not routinely
16 performed. The Committee also noted that there would be exposure to radiation with CT
17 scans (which the patient may also need to receive for other systemic investigations). Overall,
18 the Committee did not find evidence to support the use of liver biopsy, MRI and CT scans for
19 the detection of liver disease in people with cystic fibrosis and did not make
20 recommendations regarding their use.

21 The Committee considered that interpretation of ultrasound images could be subjective in
22 some respects, but that transient elastography produces a objective score that might be
23 more reliable in detecting liver disease. The Committee noted that unlike transient
24 elastography (which uses sound waves to estimate liver stiffness), images from ultrasound
25 provide information of the staging of disease progression occurring in the liver including
26 portal hypertension when Doppler is used. Instead of an image, FibroScan® produces sound
27 waves that limit the results that can be interpreted. Overall, the Committee agreed there was
28 little evidence to support the use of FibroScan® for early stage detection of liver disease, and
29 agreed that because treatment with ursodeoxycholic acid would have already been initiated,
30 the value of adding this test to a diagnostic regimen was diminished.

31 The Committee considered whether a baseline ultrasound might be performed before the
32 patient's fifth birthday, and thereafter only if abnormal clinical assessments and abnormal
33 liver function tests. However, the Committee questioned the value of performing a baseline
34 ultrasound if it is not used as a reference subsequently.

35 The Committee agreed that routine repeat ultrasounds may not be necessary in adults with
36 previous normal scans because the development of (new) liver disease after late
37 adolescence is very unusual. The value of annual review in adults is more to monitor the
38 progression of liver disease, rather than detect (new) liver disease.

39 Children with persistently elevated liver function tests or significant abnormalities on the
40 ultrasound of liver and spleen need to be referred to a specialist paediatric liver unit. In
41 children with cystic fibrosis the objective of monitoring would be to detect early evidence of
42 liver disease, to look for evidence of disease progression and especially progression to
43 chronic liver disease and its complications including portal hypertension. Clinical examination
44 is routinely performed when people with cystic fibrosis attend for clinic appointments. Clinical
45 examination would be unlikely to detect early disease but with progression might reveal
46 important signs such as hepatosplenomegaly. The Committee therefore recommended that a
47 clinical assessment for evidence of liver disease should be conducted annually for people
48 with cystic fibrosis.

49 It is currently common practice to perform blood test investigations to look for evidence of
50 liver disease. The Committee recommended that blood liver function tests should be
51 performed in people with cystic fibrosis every year.

1 The Committee recognised that ultrasound can detect evidence of liver disease (for example
2 changes in liver echogenicity, as well as advanced changes suggestive of portal
3 hypertension) and also the presence of gallstones (which sometimes occur in cystic fibrosis).
4 Transient elastography does not reveal these specific changes although it can detect
5 increased liver stiffness suggestive of liver fibrosis and progressive liver disease. Ultrasound
6 scanning is commonly used to monitor cystic fibrosis-related liver disease. The Committee
7 agreed that this was not necessary if there was no evidence of liver disease, but that a liver
8 ultrasound scan should be performed if the blood liver function tests were abnormal.

9 FibroScan® gives measurement of liver stiffness. If liver stiffness is present, the person is
10 usually referred to a specialist liver service. The lay members on the Committee pointed out
11 that from a patient perspective, FibroScan® is very quick and easy to do and may not require
12 further referral/referral time wait. However, the Committee felt the added value of
13 FibroScan® was unclear and it was unclear what decisions should follow an abnormal
14 FibroScan® result.

15 10.4.7.3 Consideration of economic benefits and harms

16 The economic harms associated with an incorrect diagnosis are much smaller for early stage
17 than late stage liver disease. If the majority of people with cystic fibrosis are incorrectly
18 diagnosed as not having liver disease (false negatives) are likely to be picked up at their next
19 annual review, the question arises as to whether additional monitoring between the annual
20 reviews is cost-effective. On the other hand, the Committee noted the cost of monitoring for
21 liver disease would be relatively insignificant when compared with the downstream costs
22 associated with liver cirrhosis and subsequent portal hypertension that could require
23 management with a liver transplant. However, the Committee agreed such cases would be
24 rare and concluded that more frequent assessments would not be a cost-effective use of
25 resources.

26 The Committee noted that if abnormal liver function blood tests are the first indication of liver
27 disease, there are potential cost savings to the NHS if those tests can replace ultrasound
28 scans at the annual review, especially in adults who are unlikely to develop liver disease
29 without prior suspicion. Following this, the Committee agreed that a recommendation in
30 favour of liver function blood tests, as the first assessment, was warranted.

31 However, the Committee agreed that the results from an ultrasound scan could lead to a
32 change in the management strategy when, for example, cirrhosis and portal hypertension
33 (with Doppler) are detected. Given that an ultrasound can add additional information to an
34 abnormal liver function blood test, a recommendation in that subgroup was considered as a
35 cost-effective use of ultrasound scans.

36 10.4.7.4 Quality of evidence

37 Most of the evidence was derived from cohort studies and outcomes presented within these
38 studies were often downgraded to moderate or low for serious or very serious imprecision of
39 sensitivity/specificity estimates. Evidence for detection of portal hypertension was largely
40 derived from a single included case control study, was low quality at best and downgraded to
41 very low quality for imprecision of specificity for some outcomes. The settings in either cystic
42 fibrosis or liver clinics were similar in all the studies.

43 The Committee was concerned that the included studies were heterogeneous in terms of
44 their population and the reference standards used. The Committee noted that a small body of
45 evidence was included overall and that the studies provided no evidence regarding early
46 cystic fibrosis-related liver disease as specified in the review question. The studies provided
47 evidence for cystic fibrosis-related liver disease which included those with cirrhosis and in
48 some studies those with clinically significant liver disease states (portal hypertension with/out
49 oesophageal varices) as well although this was not always specified clearly. There was also

1 some evidence that related to cirrhosis alone and to portal hypertension with or without
2 oesophageal varices, but no studies with participants with early disease were retrieved.
3 Although the point estimates for likelihood ratios for several comparisons indicated that the
4 index test would be moderately or very useful for ruling in or ruling out disease, the 95% CIs
5 for these point estimates were mostly wide and there was uncertainty as to whether tests
6 would be useful at all. Two exceptions were:

- 7 • High quality evidence from 1 cohort study of children and adults found that transient
8 elastography at a threshold of 6.3kPa was at least moderately useful to rule in cystic
9 fibrosis-related liver disease. The definition of cystic fibrosis-related liver disease was
10 based on practice guidelines requiring biochemical and ultrasound testing.
- 11 • A second cohort study provided high quality evidence that transient elastography at a
12 threshold of 11.5kPa was at least moderately useful to rule in portal hypertension.

13 There was no evidence that examined the use of MRI or CT scanning as reference tests nor
14 evidence that examined the use of ultrasound in addition to clinical and biochemical testing in
15 adults with no history of cystic fibrosis-related liver disease.

16 There were very little data available regarding tests that were useful to predict the
17 development of clinically significant liver disease. One study provided low quality evidence of
18 an association between increasing liver stiffness detected by transient elastography and the
19 probability of developing cystic fibrosis-related liver disease in adulthood and a second study
20 provided high quality evidence of an association between increasing fibrosis detected by
21 biopsy and the later development of portal hypertension. The paucity of prognostic outcome
22 data did not permit determination of a “best” reference standard from the evidence review. As
23 such, liver disease was defined in accordance with each reference standard.

24 10.4.7.5 Other considerations

25 No equality issues were identified by the Committee for this review question.

26 The Committee discussed the need for a research recommendation in this topic. They noted
27 FibroScan® gives measurement of liver stiffness, and this can be useful to select who should
28 referred for further specialist service. From the patient perspective, FibroScan® is very quick
29 and easy to do and may not require further referral/referral time wait. Based on this, the
30 Committee agreed it would be important to assess the added value of performing
31 FibroScan® in people with cystic fibrosis, as good quality evidence is lacking for this
32 population. However, the committee agreed this area was not a priority overall for a research
33 recommendation.

34 10.4.7.6 Key conclusions

35 The Committee chose not to make a recommendation regarding performance of clinical
36 examination at other appointments for monitoring leaving this to clinical judgement.

37 The Committee agreed that generally it was good practice to repeat liver function tests that
38 were abnormal to confirm their accuracy, although this would depend on the clinical context
39 and so they did not make a recommendation about this as this is part of routine clinical
40 practice and judgement.

41 The Committee decided that if there is evidence of chronic progressive liver disease based
42 on clinical assessment, liver function tests or the findings of an ultrasound scan the patient
43 should be referred to a liver specialist.

44 Alternatively, if investigations for liver disease (such as a clinical examination, liver function
45 tests, ultrasound imaging or FibroScan®) are persistently normal, then the cessation of
46 treatment with ursodeoxycholic acid should be considered. Monitoring for future liver damage
47 should continue in line with the annual monitoring schedule.

1 10.4.8 Recommendations

2 111. Perform a clinical assessment and liver function blood tests at the annual
3 review for people with cystic fibrosis.

4 112. Refer people with cystic fibrosis to a liver specialist if there is any of the
5 following:

- 6 • chronic progressive liver disease, based on clinical assessment, liver
7 function blood tests or the findings on a liver ultrasound scan
- 8 • liver failure, based on clinical assessment and liver function tests
- 9 • portal hypertension, haematemesis, splenomegaly or findings on a liver
10 ultrasound scan.

11 10.5 Ursodeoxycholic acid

12 **Review question: What is the effectiveness of ursodeoxycholic acid for preventing**
13 **liver disease progression in people with cystic fibrosis?**

14 10.5.1 Introduction

15 People with cystic fibrosis (CF) can develop cystic fibrosis related liver disease (CFRLD) as a
16 complication of their disease. Monitoring for the early detection of CFRLD is therefore part of
17 the overall routine care for all people with cystic fibrosis. CFRLD is a common complication in
18 cystic fibrosis, with clinical presentation of hepatomegaly or splenomegaly usually around 10
19 years of age (see Complications review). The early detection and monitoring of CFRLD is
20 therefore important in order to initiate and monitor potentially beneficial treatments.

21 Ursodeoxycholic acid (UDCA) is a commonly prescribed drug in people with cystic fibrosis. It
22 is a bile acid analogue which may act to improve biliary flow in people with cystic fibrosis
23 related disease. The current practice and indications for prescribing this drug varies widely.
24 Although UDCA is an inexpensive and well tolerated drug, it represents an additional
25 treatment burden for people with cystic fibrosis and it is within this context that this review of
26 effectiveness has been conducted.

27 10.5.2 Description of clinical evidence

28 The objective of this review was to assess the clinical and cost effectiveness of UDCA for
29 preventing liver disease progression in people with cystic fibrosis.

30 We looked for systematic reviews of randomised controlled trials (RCTs), RCTs (including
31 cross-over trials). Comparative cohort studies were not considered for inclusion, as enough
32 evidence was retrieved from RCTs.

33 For full details see review protocol in Appendix D.

34 One Cochrane systematic review was identified for this review question (Cheng 2014) which
35 included evidence from 3 RCTs (Colombo 1996, Merli 1994 and O'Brien 1992). Where
36 possible, data and risk of bias assessment were extracted directly from the Cochrane
37 systematic reviews. Individual studies were retrieved for completeness and accuracy, and
38 were also checked for additional outcomes of interest.

39 All the RCTs included had either 6 or 12 months follow-up. No RCTs from the time of
40 publication of the Cochrane systematic review and this review were found.

1 One RCT had cross-over study design (Merli 1994), and was conducted with 51 participants
2 allocated to UDCA for 6 months, a wash out period of 1 month and then placebo for 6
3 months.

4 The trials were conducted in Italy and Ireland, and the population ranged from 12 to 55
5 participants. All the trials included children, young people and adults with cystic fibrosis. Two
6 of the trials included people with liver disease (Colombo 1996, O'Brian 1992), and 1 included
7 people with and without liver disease (Merli 1994).

8 Where change in hepatocellular enzyme or bilirubin was not reported in the RCT, the final
9 value of the enzyme or bilirubin was reported.

10 A summary of the included studies is presented in Table 159. See also study selection flow
11 chart in Appendix F, study evidence tables in Appendix G, list of excluded studies in
12 Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

13 10.5.3 Summary of included studies

14 A summary of the studies that were included in this review are presented in Table 159.

15 **Table 159: Summary included studies**

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Cheng 2014 Cochrane systematic review	Comparison: UDCA vs placebo (Colombo 1996, Merli 1994, O'Brien 1992)	People with CF diagnosed by sweat test and clinical symptoms.	<ul style="list-style-type: none"> Lack of normalisation of liver enzymes (ALT, AST and GGT) 	
Primary studies included in the SR				
Colombo 1996 (Italy) RCT	UDCA <ul style="list-style-type: none"> 6 months at a daily dose of 1 to 3 300 mg capsules Placebo	N=55 children, young people and adults with CF and chronic liver disease with persistent alterations of serum liver enzymes. Median age (range): 13.8 years (4 to 22 years)	<ul style="list-style-type: none"> Percentage change in hepatocellular enzymes: ALT, AST and GGT. Liver failure Liver transplantation 	
Merli 1994 (Italy) Cross-over trial	UDCA <ul style="list-style-type: none"> dose of 12 mg/kg per day Placebo	N=51 children, young people and adults with CF. Median age (range): 14 years (8 to 32) Participants with CFRLD and no CFRLD included.	<ul style="list-style-type: none"> Change in hepatocellular enzymes (AST, ALT, GGT) No development of liver disease 	
O'Brien 1992 (Ireland) RCT	UDCA <ul style="list-style-type: none"> 20 mg/kg per day 	N=12 children, young people and adults with CF and liver disease	<ul style="list-style-type: none"> Change in hepatocellular enzymes (AST, ALT, GGT) 	

Study	Intervention/Comparison	Population	Outcomes	Comments
	No treatment	based on hepatomegaly and/ or splenomegaly. Median age (range): 17 (12 to 20) in the intervention group; 17.5 (14 to 25 in the control group)	<ul style="list-style-type: none"> Final bilirubin value 	

1 ALT: alanine aminotransferase; AST: aspartate aminotransferase; CF: cystic fibrosis; GGT, gammaglutamyl
2 transferase; mg/ kg: milligrams per kilogram; OR: odds ratio; SR: systematic review; UDCA: ursodeoxycholic acid

3 10.5.4 Clinical evidence profile

4 The summary clinical evidence profiles for this review question (UDCA) are presented in,
5 Table 160.

6 **Table 160: Summary clinical evidence profile: Comparison 1. UDCA versus Placebo**
7 **or control**

Comparison 1. UDCA versus Placebo or control						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo/control	UDCA				
Lack of normalisation of AST Follow-up: 6 months	Study population		RR 1.51 (0.83 to 2.78)	14 (Merli 1994, O'Brien 1992) ¹	⊕⊕⊕⊖ moderate ²	
	625 per 1000	944 per 1000 (519 to 1000)				
	Moderate					
	750 per 1000	1000 per 1000 (622 to 1000)				
Lack of normalisation of ALT Follow-up: 6 months	Study population		RR 0.69 (0.27 to 1.74)	12 (Merli 1994, O'Brien 1992) ¹	⊕⊕⊕⊖ moderate ²	
	750 per 1000	518 per 1000 (203 to 1000)				
	Moderate					
	833 per 1000	575 per 1000 (225 to 1000)				
Lack of normalisation of GGT Follow-up: 6 months	Study population		RR 0.6 (0.16 to 2.29)	10 (Merli 1994, O'Brien 1992) ¹	⊕⊕⊖⊖ low ³	
	500 per 1000	300 per 1000 (80 to 1000)				
	Moderate					
	333 per 1000	200 per 1000 (53 to 763)				
Final bilirubin value (umol/l) Follow-up: 6 months	The mean final bilirubin value in the control groups was 9.2	The mean final bilirubin value in the UDCA groups was 4 higher (3.72 lower to 11.72 higher)		12 (O'Brien 1992)	⊕⊕⊖⊖ low ³	

Comparison 1. UDCA versus Placebo or control						
Percentage change in AST Follow-up: 12 months	The mean percentage change in ALT in the control groups was -17 %	The mean percentage change in ALT in the UDCA groups was 13 lower (29.35 lower to 3.35 higher)		27 (Colombo 1996)	⊕⊕⊕⊕ low ^{2,7}	
Percentage change in ALT Follow-up: 12 months	The mean percentage change in ALT in the control groups was -17 %	The mean percentage change in ALT in the UDCA groups was 13 lower (29.35 lower to 3.35 higher)		27 (Colombo 1996)	⊕⊕⊕⊕ low ^{2,4}	
Percentage change in GGT Follow-up: 12 months	The mean percentage change in GGT in the control groups was -15 %	The mean percentage change in GGT in the UDCA groups was 11 lower (36.74 lower to 14.74 higher)		27 (Colombo 1996)	⊕⊕⊕⊕ low ^{2,4}	
No development of liver disease Follow-up: 6 months	None of the participants developed liver disease.	None of the participants developed liver disease.	Not calculable ⁵	11 (Merli 1994) ¹	⊕⊕⊕⊕ high	
Liver failure (jaundice) Follow-up: 12 months	1/15	0/13	Not calculable ⁶	28 (Colombo 1996)	⊕⊕⊕⊕ moderate ⁴	
Liver transplantation Follow-up: 12 months	-	1 person in the UDCA group was withdrawn to receive transplantation	Not applicable	28 (Colombo 1996)	⊕⊕⊕⊕ moderate ⁴	Outcome reported narratively
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CFLD: ALT: alanine aminotransferase; AST: aminotransferase; cystic fibrosis liver disease; CI: confidence interval; GGT: gamma glutamyltransferase; MD: mean difference; RR: risk ratio; UDCA: ursodeoxycholic acid						

- 1 Merli (1994) used a cross-over study design
- 2 The quality of the evidence was downgraded by 1 because the 95% CI crosses 1 default MID.
- 3 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs.
- 4 The quality of the evidence was downgraded by 1 due to lack of allocation concealment reporting.
- 5 RR not calculable - no development of liver disease in 11/11 participants who did not have CF related liver disease at entry in this cross-over trial.
- 6 Not calculable - 0 events in placebo arm.

1 10.5.5 Economic evidence

2 No economic evaluations of interventions relevant to UDCA were identified in the literature
3 search conducted for this guideline. Full details of the search and economic article selection
4 flow chart can be found in Appendix E and F, respectively.

5 This review question was not prioritised for de novo economic modelling. To aid
6 considerations of cost-effectiveness the cost of UDCA over the course of 1 week and a
7 month of continued has been estimated for all preparations.

8 Drug acquisition costs are taken from the NHS Electronic Drug Tariff October 2015. For an
9 indication of primary biliary cirrhosis the BNF recommends a dose of 12–16 mg/kg daily in 3
10 divided doses for 3 months, then 12–16 mg/kg once daily at bedtime. Conversely in the
11 studies by Merli 1994 and O'Brien 1992, participants received 12mg/kg daily and 20mg/kg
12 daily, respectively.

13 For illustrative purposes, the costs in Table 161 are based 600mg daily as this was
14 considered to be representative of the typical person with cystic fibrosis in clinical practice.

15 **Table 161: Acquisition cost of UDCA**

UDCA preparation	Quantity	Basic price	Cost per unit	Cost per day	Cost per week	Cost per month
150mg tablets	60	£19.02	£0.32	£1.27	£8.88	£39.82
300mg tablets	60	£38.86	£0.65	£1.30	£9.07	£40.67
250mg capsules	60	£25.29	£0.51	£1.21	£8.50	£38.12
Oral suspension sugar free 250mg/5ml	250ml	£26.98	£0.45a	£1.08b	£7.55	£33.89

16 (z) *b a unit equals 5ml; b 2.4x 5ml*

17 10.5.6 Evidence statements

18 10.5.6.1 Comparison 1. UDCA versus placebo or control

19 Change of hepatocellular enzymes or bilirubin level

20 Moderate quality evidence from 2 RCTs with 14 children, young people and adults with cystic
21 fibrosis showed no clinically significant difference in lack of normalisation of AST between the
22 group of participants receiving UDCA (12 or 20 mg/kg/day or) and those in the control group
23 at 6 months follow-up.

24 Moderate quality evidence from 2 RCTs with 12 children, young people and adults with cystic
25 fibrosis showed no clinically significant difference in lack of normalisation of ALT between the
26 group of participants receiving UDCA (12 or 20 mg/kg/day or) and those in the control group
27 at 6 months follow-up.

28 Low quality evidence from 2 RCTs with 10 children, young people and adults with cystic
29 fibrosis showed no clinically significant difference in lack of normalisation of GGT between
30 the group of participants receiving UDCA (12 or 20 mg/kg/day or) and those in the control
31 group at 6 months follow-up.

32 Low quality evidence from 1 RCT with 12 children, young people and adults with cystic
33 fibrosis showed no clinically significant difference in final bilirubin values (umol/l.) between
34 the group of participants receiving UDCA (20 mg/kg/day) and those in the control group at 6
35 months follow-up.

- 1 Low quality evidence from 1 RCT with 27 children, young people and adults with cystic
2 fibrosis showed no clinically significant difference in percentage change in AST, ALT and
3 GGT between the group of participants receiving UDCA (1 to 3 300 mg. capsules per day)
4 and those in the control group at 12 months follow-up.
- 5 **No development of liver disease**
- 6 High quality evidence from 1 cross-over trial with 11 participants found that participants who
7 did not have cystic fibrosis related liver disease who received both UDCA and placebo did
8 not develop liver disease at 6 months follow up.
- 9 **Liver failure**
- 10 Moderate quality evidence from 1 RCT with 28 participants reported that 1 participant taking
11 UDCA had liver failure due to jaundice and no liver failure was found in placebo arm during
12 the follow-up period of 12 months. There was no difference between UDCA and placebo.
- 13 **Liver transplantation**
- 14 Moderate quality evidence from 1 RCT with 28 participants reported that 1 participant taking
15 UDCA required liver transplantation which was performed successfully and no liver
16 transplantation was required in the placebo arm during the follow-up period of 12 months.
17 There was no difference between UDCA and placebo.
- 18 A Cochrane systematic review of RCTs reported that none of the participant in 2 of the trials
19 (Merli 1994 and O'Brien 1992) required liver transplantation (with personal communication
20 with authors).
- 21 **Liver related mortality**
- 22 No evidence was found for this important outcome. However, a Cochrane systematic review
23 of RCTs reported there was no deaths in 2 of the trials (Merli 1994 and O'Brien 1992, with
24 personal communication with authors).
- 25 **Development of portal hypertension**
- 26 No evidence was found for this critical outcome. However, a Cochrane SR of RCTs reported
27 that no development of portal hypertension occurred amongst participants in 2 RCTs (Merli
28 1994 and O'Brien 1992, with personal communication with authors).
- 29 **Quality of life**
- 30 No evidence was found for this important outcome.
- 31 **10.5.6.2 Economic evidence statements**
- 32 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.
- 33 **10.5.7 Evidence to recommendations**
- 34 **10.5.7.1 Relative value placed on the outcomes considered**
- 35 The objective of this review was to assess the clinical and cost effectiveness of UDCA for
36 preventing liver disease progression in people with cystic fibrosis.
- 37 The Committee agreed the following as critical outcomes for consideration in decision
38 making: change of hepatocellular enzyme levels, progression to liver failure and

1 development of portal hypertension indicated by an enlarged spleen (increased by at least
2 15%), development of varices or ultrasound evidence of portal hypertension. Health related
3 quality of life, liver transplantation, liver related mortality and no development of liver disease
4 were rated as important outcomes.

5 **10.5.7.2 Consideration of clinical benefits and harms**

6 The Committee discussed the use of UDCA for primary prevention and secondary
7 prophylaxis.

8 With regards to the use of UDCA in people with cystic fibrosis without liver disease, the
9 Committee noted most of the participants in the trials had pre-existing evidence of liver
10 disease. Therefore there was a lack of evidence to support the use of UDCA as primary
11 prophylaxis aimed at preventing the onset of liver disease.

12 The Committee noted that infants with cystic fibrosis who have meconium ileus (about 10-
13 15%) were at increased risk of developing liver disease. However, in the absence of
14 evidence of evidence showing a benefit, they did not recommend the use of UDCA even in
15 these infants.

16 The Committee also discussed when UDCA should be given to prevent progression of liver
17 disease, and the duration of administration. They noted the available trials were small and
18 the duration of follow-up was no more than a year, and consequently it was not surprising
19 that there was no evidence for clinical benefit with UDCA in terms of preventing development
20 of portal hypertension, liver failure or mortality.

21 The evidence did not show differences either in terms of normalisation of hepatobiliary
22 enzymes (including AST, ALT and GGT), and final bilirubin values at 6 months, or in
23 percentage change in hepatobiliary enzymes (including AST, ALT and GGT) at 12 months
24 between UDCA and placebo. However, the Committee noted that signs of advanced liver
25 disease, including portal hypertension were present at baseline for most of the participants.
26 The fact that the liver disease had already progressed to this degree before starting UDCA
27 might have led to an underestimate the potential benefit effect of UDCA in preventing
28 disease progression.

29 The Committee therefore recommended that if the liver function tests were abnormal,
30 treatment with UDCA could be considered. The optimal duration of treatment was unknown
31 and so they recommended that clinicians should think about discontinuing it if the liver
32 function tests returned to normal levels and there was no clinical or ultrasound scan evidence
33 of liver disease. If, subsequently, monitoring showed a recurrence of liver function test
34 abnormality then the clinician would again consider giving UDCA. The Committee also
35 agreed that if the abnormal liver function tests did not resolve with UDCA and remained
36 persistently abnormal, the clinician should consider referring to a liver specialist.

37 The Committee did not made specific recommendations on the degree of abnormality of the
38 various liver function tests used that should be considered abnormal or on the advisability of
39 repeating blood tests when considering UDCA. They recognised that this was a complex
40 area that required clinical expertise and judgement. A minor elevation of AST for example
41 might not be significant whereas elevation of the serum bilirubin would be a reason for
42 concern.

43 The Committee believed that UDCA was considered safe for use, although in higher dosages
44 it might cause diarrhoea.

45 **10.5.7.3 Consideration of economic benefits and harms**

46 The preparations of UDCA are comparable and relatively inexpensive in the short-term at a
47 cost of approximately £1.08 to £1.27 a day (600mg daily) for capsules and oral suspension,

1 respectively. However, the costs of prolonged use could be significant, and without knowing
2 the benefits of UDCA in people with cystic fibrosis without liver disease, we cannot know if
3 UDCA is a cost-effective prophylactic. Consequently, the Committee agreed not to
4 recommend UDCA in people with cystic fibrosis without evidence of liver disease.

5 Following this, the Committee added that portal hypertension and liver disease were reported
6 at baseline for many participants in the studies included in the clinical evidence review. As a
7 result, they stated that this could potentially underestimate the cost-effectiveness of UDCA to
8 prevent the initiation of liver disease.

9 The Committee agreed that although UDCA is relatively inexpensive and it has a good safety
10 profile, this does not justify the continued prescription of these drugs, without evidence of
11 effectiveness. As a result, the Committee made a recommendation to reduce unnecessary
12 treatment costs and treatment burden, by stopping UDCA if the annual clinical assessments
13 show no evidence of liver disease.

14 **10.5.7.4 Quality of evidence**

15 The quality of the evidence presented in this report ranged from low to high as assessed by
16 GRADE.

17 The risk of bias was 1 of the reasons that lead to downgrading the quality of the evidence, in
18 particular lack of allocation concealment.

19 Imprecision also lead to downgrading the quality of the evidence, as the confidence interval
20 crossed 1 or 2 clinical or default MIDs (minimal important difference).

21 Inconsistency was not an issue, as most outcomes were reported by a single study.

22 The evidence was not downgraded for indirectness, but the Committee noted that some of
23 the participants had signs of liver disease at enrolment, and therefore evidence on
24 preventing development of liver disease is lacking.

25 **10.5.7.5 Other considerations**

26 No equality issues were identified by the Committee for this review question.

27 The Committee agreed to draft a research recommendation for this topic. They noted liver
28 disease is the third most common cause of mortality in people with cystic fibrosis, and
29 around 10 to 30% of people with cystic fibrosis will develop CFRD (See complications
30 review). Children with meconium ileus are at an increased risk of liver disease, and starting
31 treatment with UDCA from diagnosis may reduce this risk. UDCA appears safe, and is well
32 tolerated and cheap. Routine use could increase people's overall quality of life and reduce
33 the need for subsequent treatment for liver disease, but more research is needed into the
34 effectiveness and safety of this treatment.

35 **10.5.7.6 Key conclusions**

36 The Guideline Committee concluded that in the absence of evidence, they would not
37 recommend using UDCA as primary prophylaxis.

38 The Committee also discussed that the available evidence did not show that UDCA was
39 beneficial in achieving normalization of hepatobiliary enzymes in people with cystic fibrosis
40 and pre-existing liver disease. They noted the limitations of the evidence, as most of the
41 participants in the studies had signs of advanced liver disease at baseline (therefore the fact
42 that the liver disease had already progressed to this degree before starting UDCA might
43 have led to an underestimate the potential benefit effect of UDCA in preventing disease
44 progression). They also agreed longer studies would be needed to assess whether UDCA is
45 effective in terms of preventing development of portal hypertension, liver failure or mortality.

1 Moreover, the Committee noted liver disease is a serious complication of cystic fibrosis, and
2 it is very important to delay its progression. Based on all this, the Committee agreed UDCA
3 could be considered if liver function tests were abnormal. They also agreed to monitor for
4 response to treatment, and to refer to specialist advice if abnormal results are persistent.

5 10.5.8 Recommendations

6 **113. If liver function blood tests are abnormal, perform a liver ultrasound scan and**
7 **consider ursodeoxycholic acid treatment.**

8 **114. Think about stopping ursodeoxycholic acid if liver function blood tests return**
9 **to normal and clinical assessment and liver ultrasound scan show no liver**
10 **disease.**

11 **115. If ursodeoxycholic acid is stopped, monitor for re-emergence of liver disease**
12 **using clinical assessment and liver function blood tests.**

13 **116. Think about referring people with cystic fibrosis to a liver specialist if the liver**
14 **function blood test results are persistently abnormal despite treatment with**
15 **ursodeoxycholic acid.**

16 10.5.9 Research recommendations

17 **4. Should all children with meconium ileus receive ursodeoxycholic acid from**
18 **diagnosis?**

19 **Table 162: Research recommendation justification**

Research question	Should all children with meconium ileus receive ursodeoxycholic acid from diagnosis?
Why this is needed	
Importance to 'patients' or the population	Liver disease is the third commonest cause of death in cystic fibrosis. Meconium ileus is a risk factor for the development of cystic fibrosis related liver disease. It is possible that ursodeoxycholic acid (UDCA) decreases the incidence or severity of cystic fibrosis related liver disease. UDCA is relatively inexpensive and well tolerated, and if it decreased the incidence or severity of cystic fibrosis related liver disease this could improve quality of life and decrease mortality.
Relevance to NICE guidance	If UDCA changed the incidence or severity of cystic fibrosis related liver disease, this would change the recommendations for monitoring and treatment. <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline.
Relevance to the NHS	Financial cost of regular therapy and treatment burden to the patient. Cost savings if decreased incidence and severity of liver disease
National priorities	No document identified.
Current evidence base	UDCA can improve deranged biochemical tests of liver function, but it is unclear whether if given to high risk groups it is able to alter the incidence or severity of disease.
Equality	This intervention is of particular relevance to infants born with meconium ileus who are at increased risk of cystic fibrosis related liver disease.
Feasibility	The proposed research can be carried out within a realistic timescale and at an acceptable cost. There are no ethical or technical issues.

Research question	Should all children with meconium ileus receive ursodeoxycholic acid from diagnosis?
Other comments	None

1

2

Table 163: Research recommendation statements

Criterion	Explanation
Population	Children with cystic fibrosis and meconium ileus
Intervention	<ul style="list-style-type: none"> • Ursodeoxycholic acid (UDCA)
Comparators	<ul style="list-style-type: none"> • Usual care • Placebo
Outcomes	<ul style="list-style-type: none"> • Health related quality of life (CF-QOL, CFQR) • Change of hepatocellular enzymes or bilirubin level • Liver failure • Liver transplantation • Liver related mortality • Development of cystic fibrosis related-disease (enlarged spleen, development of varices, ultrasound evidence of portal hypertension) • No development of cystic fibrosis liver disease • Resource use • Unit costs
Study design	Randomised control trial
Timeframe	Two years of randomisation and 5 year of follow up, providing recruitment numbers are sufficient to achieve population numbers of sufficient size to answer the research question.

3 **10.6 Monitoring for cystic fibrosis related diabetes**

4 **10.6.1 Introduction**

5 Diabetes mellitus is a common complication of cystic fibrosis (CF). It differs from both type 1
6 and 2 diabetes mellitus encountered in people without cystic fibrosis, and therefore is
7 referred to as CF-related diabetes or 'CFRD'. It is caused by a combination of slow
8 progressive loss of the insulin-producing β -cells in the pancreas and the development of end-
9 organ resistance to insulin. The prevalence of CFRD increases with age. Although symptoms
10 of CFRD can include weight loss, thirst and increased urinary frequency, its onset can be
11 insidious. Without treatment, uncontrolled CFRD can lead to a number of significant
12 complications, including more frequent respiratory exacerbations, an accelerated loss of lung
13 function, and a decline in nutritional status. People with cystic fibrosis are therefore regularly
14 screened for CFRD, so treatment can be started early to prevent any complications. The
15 clinical management of CFRD differs from other forms of diabetes mellitus, in both dietary
16 advice and the use of agents to control blood sugar levels.

17 **10.6.2 Description of clinical evidence**

18 The aim of this review was to determine what criteria should be used to determine the need
19 for insulin therapy to achieve optimal patient outcomes.

20 As the Committee anticipated a lack of evidence to respond to this question, we also aimed
21 to determine what thresholds of glucose dysregulation are associated with more rapid
22 progression of lung disease (as a proxy measure to determine which criteria should be used
23 to instigate treatment with insulin in people with cystic fibrosis).

1 We searched for systematic reviews, test-and-treat randomized controlled trials (RCTs) and
2 prospective and retrospective comparative cohort studies. Conference abstracts were also
3 considered in the absence of full published RCTs.

4 For full details see review protocol in Appendix D.

5 No studies were identified.

6 See study selection flow chart in Appendix F, and list of excluded studies in Appendix H.

7 **10.6.3 Summary of included studies**

8 **Review question 1. What criteria should be used to determine the need for insulin** 9 **therapy to achieve optimal patient outcomes?**

10 No studies were identified for this question.

11 **Review question 2. What thresholds of glucose dysregulation are associated with** 12 **more rapid progression of lung disease?**

13 No studies were identified for this question.

14 **10.6.4 Clinical evidence profile**

15 Not applicable, as no evidence was found for this review.

16 **10.6.5 Economic evidence**

17 No economic evaluations of strategies to monitor for the onset of CFRD were identified in the
18 literature search conducted for this guideline. Full details of the search and economic article
19 selection flow chart can be found in Appendix E and F, respectively.

20 This review question was not prioritised for de novo economic modelling. To aid
21 consideration of cost-effectiveness relevant resource and cost use data on the OGTT and
22 CGM are presented in Appendix K.

23 **10.6.6 Evidence statements**

24 Not applicable, as no clinical or economic evidence was found for this review.

25 **10.6.7 Evidence to recommendations**

26 **10.6.7.1 Relative value placed on the outcomes considered**

27 The aim of this review was to determine what criteria should be used to determine the need
28 for insulin therapy to achieve optimal patient outcomes.

29 The Committee chose change in lung function (forced expiratory volume in 1 second (FEV₁),
30 forced vital capacity (FVC) and lung clearance index (LCI)), pulmonary exacerbations and
31 body mass index (BMI) (z-scores for children) as critical outcomes for decision making;
32 whereas adverse events and patient acceptability or satisfaction were rated as important
33 outcomes.

34 **10.6.7.2 Consideration of clinical benefits and harms**

35 The Committee recognised that diabetes is a very common complication in people with cystic
36 fibrosis, and it is important to identify it without delay. They highlighted early treatment with

1 insulin is very important, as it may improve lung function and weight. However they
2 recognised the lack of evidence in relation to these matters, and they were aware of a lack of
3 consensus in clinical practice.

4 The Committee emphasised that glucose may fluctuate in people with cystic fibrosis, and the
5 tests usually used to diagnose diabetes in the general population may not be suitable for the
6 detection of cystic fibrosis related diabetes. For this reason, they agreed that single point
7 tests such as HbA1c or fasting plasma glucose measurement when used alone are not
8 useful for the identification of cystic fibrosis related diabetes. This is also supported by a
9 previous Health Technology Assessment (HTA 2012) and by the American Diabetes
10 Association (ADA 2010).

11 The Committee discussed at length the use of the oral glucose tolerance test (OGTT). They
12 noted this test is commonly used to monitor for cystic fibrosis related diabetes in current
13 practice, and is the test of choice for many professionals. The use of this test is also
14 recommended by the American Diabetes Association [consensus recommendation] (ADA
15 2010) and the European Cystic Fibrosis Foundation (Alan 2014). However, they also noted
16 the OGTT may not be an ideal test, given that as mentioned above, glucose dysregulation in
17 cystic fibrosis related diabetes is a dynamic and variable process. Based on this, they agreed
18 OGTT may not be completely reliable, and if used and found abnormal it should be backed
19 up with dynamic testing so that glucose regulation is assessed over several days. The
20 Committee discussed the important fact that hyperglycaemia associated with glucose
21 dysregulation in cystic fibrosis related diabetes can have a negative effect on clinical
22 outcomes, even before the usual diagnostic criteria for diabetes are met.

23 The Committee discussed the potential benefit of continuous glucose monitoring (CGM), in
24 that it would more reliably detect glucose dysregulation, as it monitors the blood sugar level
25 continuously for several days rather than relying on a single or few glucose measurements.
26 This test is increasingly used in clinical practice for the detection of the onset of diabetes in
27 people with cystic fibrosis. However, they acknowledged there is currently no evidence to
28 require the routine use of this test. In addition, some drawbacks were also noted. This test is
29 more difficult to perform and to interpret (compared to the OGTT) and it has to be
30 administered by trained staff, making it difficult to generalise its use to all cystic fibrosis
31 clinics.

32 The Committee agreed monitoring for cystic fibrosis related diabetes should be done
33 annually and should start from the age of 10, as recommended by the American Diabetes
34 Association (ADA 2010) and the US Cystic Fibrosis Foundation (Moran 2010). This is also
35 consistent with current practice. Apart from this process of annual monitoring, investigation
36 should be undertaken in a number of specific contexts. Clearly if people with cystic fibrosis
37 had symptoms or signs of diabetes such as polydipsia and polyuria they should be
38 appropriately investigated as recommended in existing NICE diabetes guidelines for children,
39 young people and adults. In addition, the committee recommended consideration be given to
40 investigating those with cystic fibrosis who, despite optimising their pulmonary management,
41 have unexplained weight loss or an unexplained deterioration in lung function as measured
42 by pulmonary spirometry, or frequent pulmonary exacerbations, or lethargy and malaise.
43 They also recommended monitoring for people taking long-term systemic corticosteroids or
44 enteral tube feeding, in whom glucose dysregulation may occur, and that this should be done
45 using continuous glucose monitoring or serial glucose monitoring.

46 With regards to monitoring diabetes during pregnancy, the Committee referred to the current
47 NICE guidelines on Diabetes in Pregnancy (NG3 2015), but they noted they do not
48 specifically address monitoring of cystic fibrosis related diabetes in pregnancy. The
49 Committee noted that insulin therapy is often needed in pregnant women with cystic fibrosis,
50 as the threshold for commencing insulin is lower than in those without cystic fibrosis. They
51 recommended that either continuous glucose monitoring or an oral glucose tolerance test
52 should be performed at the end of the first and second trimesters of pregnancy, this timing

1 being consistent with recommendations from the US Cystic Fibrosis Foundation (Moran
2 2010).

3 10.6.7.3 Consideration of economic benefits and harms

4 The Committee stated that people with cystic fibrosis require their own clinic space when
5 attending for an OGTT to reduce their risk of cross-infection, incurring a larger cost (£50
6 than OGTTs undertaken in a shared clinic space (NICE NG3 2015, 2-sample OGTT:
7 £22.06). The Committee also added that in cases with an abnormal OGTT result, further
8 dynamic tests such as serial blood sugar monitoring or CGM would be performed after a few
9 days to obtain a complete picture of glucose levels.

10 Given that single-point tests can result in more false positives and negatives than two-, or
11 three- point tests, the Committee agreed that their recommendations should favour dynamic
12 testing to reduce the number of people with cystic fibrosis that incur the downstream costs
13 from an incorrect diagnosis and ordered their recommendation on the type of test to that
14 effect. However, when OGTT or HbA1c are performed, the Committee agreed normal results
15 have a higher specificity and sensitivity than abnormal results; hence, abnormal results
16 should be followed by CGM or serial blood glucose monitoring to outweigh the potential costs
17 of an incorrect diagnosis.

18 Unlike CGM in confirmed diabetes, the duration of CGM to monitor for its onset would be up
19 to 5 to 7 days. For this reason, the Committee stated that CGM systems would be shared
20 across the clinic when used to monitor for the onset of CFRD, reducing the cost per patient
21 from approximately £3,500/ year (NICE NG17 2015) to £200/ year (including attendances
22 and delays to receive and return the system). Following this, the Committee questioned if
23 CGM was the most expensive investigation, as the cost of OGTT could overtake the cost of
24 CGM if several visits were required, and more so in children who require extra staff time to fit
25 a cannula for their OGTT.

26 However, the Committee noted that the majority of clinics rely on OGTT, or HbA1C to
27 monitor for CFRD, as they do not have access to CGM systems, or the expertise on how to
28 use and interpret them appropriately in this population. The Committee also noted that each
29 centre would require several CGM systems to ensure enough people were monitored by the
30 centre each week. As a result, recommending CGM would lead a change in current practice
31 and would incur upfront training costs and capital costs to implement.

32 Overall, the Committee agreed they could not change current practice from the OGTT to
33 CGM because there was not enough evidence to support the large injection of resources. As
34 a result, the Committee made a recommendation in favour of dynamic testing to allow
35 centres to decide which strategy (CGM, serial glucose testing over several days, OGTT, or
36 HbA1C) is cost-effective for them to provide. The Committee also prioritised a research
37 recommendation to identify the best strategy to diagnose CFRD in people with cystic fibrosis,
38 to mitigate current uncertainty in this area.

39 The Committee also outlined the populations at high risk of CFRD in their recommendations
40 that may require more frequent monitoring. Those populations were prioritised as their
41 condition could escalate much quicker between the annual reviews, leading to more intense
42 and costly management.

43 10.6.7.4 Quality of evidence

44 Not applicable, as no evidence was found for this review.

1 10.6.7.5 Other considerations

2 The Committee discussed potential equality issues. They noted cystic fibrosis related
3 diabetes is common in females. However they agreed care is the same for both sexes, and
4 therefore there was no need to draft additional recommendations.

5 Given the lack of evidence, and the uncertainty in relation to what is the best screening
6 strategy to detect the onset of diabetes in people with cystic fibrosis, the Committee agreed a
7 research recommendation was useful. They highlighted it would be important to look at
8 OGTT and CGM. This would provide guidance on when treatment with insulin should be
9 initiated in people with cystic fibrosis.

10 10.6.7.6 Key conclusions

11 The guideline Committee concluded that early detection of CFRD is very important. They
12 noted OGTT is often used in clinical practice but they agreed it is an unreliable test, and if
13 used it should be backed up with dynamic testing over several days. They suggested CGM
14 could be used instead as it monitors the blood sugar level continuously. Children should be
15 monitored for the onset of CFRD from the age of 10 years. Monitoring should be done as
16 part of the annual assessment, and if there are symptoms suggestive of diabetes.

17 10.6.8 Recommendations

18 117. Diagnose cystic-fibrosis-related diabetes using one of the following:

- 19
- 20 • continuous glucose monitoring (CGM)
 - 21 • serial glucose testing over several days
 - 22 • oral glucose tolerance testing (OGTT) – if OGTT is abnormal perform
CGM or serial glucose testing over several days to confirm the diagnosis
 - 23 • HbA1c – if HbA1c is abnormal perform CGM or serial glucose testing
24 over several days to confirm the diagnosis).

25 118. Test for cystic-fibrosis-related diabetes (as detailed in the recommendation on 26 diagnosing cystic-fibrosis-related diabetes) in people with cystic fibrosis annually 27 from 10 years of age.

28 119. Test for cystic-fibrosis-related diabetes at the end of the first and second 29 trimesters of pregnancy, using CGM or OGTT.

30 120. Test for cystic-fibrosis-related diabetes in people with cystic fibrosis who are 31 taking long-term systemic corticosteroids or receiving enteral tube feeding, using 32 CGM or serial glucose monitoring.

33 121. Think about testing for cystic-fibrosis-related diabetes in people who still have 34 any of the following despite optimised cystic fibrosis treatment:

- 35
- 36 • unexplained weight loss
 - 37 • a deterioration in lung function as measured by spirometry
 - 38 • frequent pulmonary exacerbations
 - excessive tiredness.

39 10.6.9 Research recommendations

40 5. What is the most clinical and cost effective screening strategy to detect diabetes 41 in people with cystic fibrosis?

1

Table 164: Research recommendation justification

Research question	What is the most clinical and cost effective screening strategy to detect diabetes in people with cystic fibrosis?
Why this is needed	
Importance to 'patients' or the population	<p>People with cystic fibrosis are at high risk of developing cystic fibrosis related diabetes (CFRD). The onset can be insidious but without institution of therapy is associated with accelerated progression of lung disease. A strategy for the early identification of CFRD will allow treatment to be commenced in a timely manner and help prevent the clinical decline associated with this condition.</p> <p>Currently there are 10,800 people with cystic fibrosis, of whom 3,759 are age >10 years screened for CFRD in 2015. Further 1,192 were not screened, but could have been eligible.</p>
Relevance to NICE guidance	<p>High: the research is essential to inform future updates of key recommendations in the guideline.</p> <p>The guideline committee spent some time debating the optimal screening strategy for CFRD. They agreed it was important for clinical care but there are differences in clinical practice between cystic fibrosis centres. They found no suitable evidence to guide their decision.</p>
Relevance to the NHS	It may help prevent both the progression of lung disease and the development of long term complications of diabetes in this population, both of which incur increased utilisation of healthcare resources
National priorities	None
Current evidence base	A search was undertaken to find evidence to inform the development of this current guideline; no suitable evidence base was identified.
Equality	No equality issues
Feasibility	No equality issues
Other comments	None

2

Table 165: Research recommendation statements

Criterion	Explanation
Population	People with cystic fibrosis age >10 years
Intervention	Continuous Glucose Monitoring (CGM), followed up with insulin treatment (Possible sub-groups according to different levels of glucose dysregulation)
Comparators	Current method (OGTT), followed up with insulin treatment
Outcomes	<ul style="list-style-type: none"> • Change in lung function (measured with FEV₁, FVC, LCI) • Time to next pulmonary exacerbation • Change in weight or BMI • Adverse events (for example hypoglycaemic episodes due to insulin therapy) • Patient acceptability/ satisfaction (with insulin therapy) • Resource use • Unit costs
Study design	Test-and-treat randomised controlled trial (RCT)
Timeframe	1 year

1 10.7 Monitoring for low bone mineral density

2 **Review question: What is the most effective strategy to monitor for the identification**
3 **of reduced bone mineral density in people with CF?**

4 10.7.1 Introduction

5 Improved survival in people with cystic fibrosis has contributed to the development of co-
6 morbidities. Low bone mineral density is a common co-morbidity affecting people with CF. It
7 is, however, more frequently seen in adolescents and adults. Low bone mineral density can
8 lead to the development of osteoporosis which results in the increased risk of fragility
9 fractures.

10 The aetiology of low bone mineral density in people with cystic fibrosis is multifactorial.
11 Contributory risk factors include poor nutritional status, delayed puberty, cystic fibrosis
12 transmembrane conductance regulator (CFTR) dysfunction, physical inactivity, deficiencies
13 of calcium, vitamins D and K, recurrent respiratory exacerbations, corticosteroid treatment
14 and hypogonadism. Medical and nutritional interventions should aim to optimise these
15 multiple factors that affect bone health.

16 An important part of standard cystic fibrosis care is the prevention and recognition of low
17 bone mineral density. Current bone health consensus committees recommend the
18 monitoring of bone health using dual energy x-ray absorptiometry (DXA) scans, although the
19 age to commence scanning and the frequency of scanning is variable and dependant on the
20 presence of risk factors.

21 10.7.2 Description of clinical evidence

22 The objective of this review was to determine the most effective strategy to monitor for the
23 identification of reduced mineral density in people with CF.

24 We looked for systematic reviews of test and treat RCTs, systematic reviews of cohort
25 studies and prospective and retrospective cohort studies that looked at regular DXA scans or
26 peripheral quantitative computed tomography (pQCT).

27 For full details see review protocol in Appendix D.

28 We identified 6 observational studies (Baker 2016, Bhudhikanok 1998, Brenckmann 2003,
29 Haworth 2002, Papaioannou 2008, Schulze 2006).

30 Three studies were conducted in the USA (Baker 2016, Bhudhikanok 1998, Schulze 2006); 2
31 in Canada (Brenckmann 2003, Papaioannou 2008) and 1 in the UK (Haworth 2002) and the
32 follow-up ranged from 1 to 4 years.

33 One study included young people (Schulze 2006); 3 included adults (Baker 2016,
34 Brenckmann 2003, Papaioannou 2008); 1 included young people and adults (Haworth 2002)
35 and 1 included children, young people and adults (Bhudhikanok 1998).

36 Population size ranged from 18 to 63 people.

37 With regards to the test, all studies used DXA scans and 1 study also used QCT (Haworth
38 2002).

39 Studies reported on change BMD at the lumbar spine (Baker 2016, Brenckmann 2003,
40 Bhudhikanok 1998, Haworth 2002, Papaioannou 2008, Schulze 2006); at the proximal femur
41 (Papaioannou 2008); at the hip (Brenckmann 2003, Haworth 2002); at the femoral neck
42 (Bhudhikanok 1998, Haworth 2002) and change in whole body BMD (Brenckmann 2003,
43 Bhudhikanok 1998, Papaioannou 2008, Schulze 2006). Five studies used z-scores (Baker

1 2016, Bhudhikanok 1998, Brenckmann 2003, Papaioannou 2008, Schulze 2006), 1 used t-
2 scores for adults (Papaioannou 2008); and 1 study used mg/ml or g/cm² (Haworth 2002).
3 Only 1 study (Baker 2016) used a multivariate model adjusting for age, gender, fat-free mass
4 index and height.

5 10.7.3 Summary of included studies

6 A summary of the studies that were included in this review are presented in Table 166.

7 **Table 166: Summary of included studies**

Study	Population	Prognostic indicator	Follow-up	Outcomes	Comments
Baker 2016 USA Retrospective observational study	N=63 adults with CF Mean age (SD): 31.7 (8.0) years (18 to 57) 50.9% male	<ul style="list-style-type: none"> • Low BMD, defined as z-score ≤ -1 • Very low BMD, defined as z-score ≤ -2 • Standard dual-energy X-ray absorptiometry (DXA) on a QDR4500A model (Hologic Inc, Bedford, MA) 	2 years (data available for n=39)	<ul style="list-style-type: none"> • Change in posterior-anterior spine BMD • Multivariate model adjusting for: age, gender, fat-free mass index and height 	Subjects that did not have a repeat PA spine DXA were significantly younger and tended to have lower baseline PA spine z-scores.
Brenckmann 2003 Canada Retrospective observational study	N=40 adults with CF Mean age (SD): 28.7 (8.4) years (19 to 52)	<ul style="list-style-type: none"> • Hip BMD • Lumbar spine • Total BMD • DXA scan. No details given. 	1 year (data available for n=27)	<ul style="list-style-type: none"> • Change in hip BMD • Change in lumbar spine • Change in total BMD 	Subgroup analysis was not possible as sample size was too small. N=21 participants were receiving oral or IV corticosteroids
Bhudhikanok 1998 USA Prospective observational study	N=47 children, young people and adults with CF Mean age, range: 20.6 years (8.4 to 48.5 years) 30 female, 19 male <ul style="list-style-type: none"> • Boy and young males n=9 • Adult males n=6 • Girls and 	<ul style="list-style-type: none"> • Lumbar spine BMD (QCT) • Femoral neck BMD (DXA) • Whole body BMD (DXA) Dual-energy X-ray (DXA) absorptiometry (QDR 100W, Hologic Corporation, Waltham, Mass)	Mean 17 months (11 to 25 months) (data available for n=41)	<ul style="list-style-type: none"> • Change in lumbar spine BMD • Change in femoral neck BMD • Change in whole body BMD 	

Study	Population	Prognostic indicator	Follow-up	Outcomes	Comments
	young females n=11 • Adult females n=15				
Haworth 2002 UK Prospective observational study	N=114 young people and adults with CF Mean (SD) age: 25.1 (6.9) years (15 to 49) • Young cohort ≤ 24 years n=55 • Adult cohort ≥ 25 years n=59	<ul style="list-style-type: none"> Lumbar spine BMD Femoral neck BMD Total hip BMD QCT and DXA (reported in a previous study) 	12 to 13 months	<ul style="list-style-type: none"> Change in lumbar spine BMD Change in femoral neck BMD Change in total hip BMD 	
Papaioannou 2008 Canada Retrospective observational study	N=49 adults with CF Mean age (SD): 25.2 (9.4) years 42.9% male	<ul style="list-style-type: none"> Lumbar spine BMD Proximal femur BMD Whole body BMD Standard dual-energy X-ray absorptiometry (DXA) on a QDR4500A model (Hologic Inc, Bedford, MA, USA) 	Mean (SD): 4.03 (1.45) years (data available for n=10)	<ul style="list-style-type: none"> Lumbar spine BMD Proximal femur BMD Whole body BMD 	
Schulze 2006 USA Prospective observational study	N=18 prepubertal and pubertal girls and young females Age range: 7.6 to 17.9 years	<ul style="list-style-type: none"> Lumbar spine BMD Whole body BMD Standard dual-energy X-ray absorptiometry (DXA) on a QDR4500A model (Hologic Inc, Bedford, MA, USA) 	Mean time: 2.13±1.14 years (range 1.6 to 4.10)	<ul style="list-style-type: none"> Change in lumbar spine BMD. Change in whole body BMD. 	Although the study reported data stratified by pubertal status, this was not reported in the review as sample size was very low (range 2 to 7) Significant differences were found by pubertal stage at baseline.

10.7.41 Clinical evidence profile

2 A summary of the results in presented in Table 167.

3 **Table 167: Prognostic indicators for bone mineral density**

Study	Prognostic indicator	Effect size	Quality
Baker 2016	<ul style="list-style-type: none"> • Low baseline BMD, defined as z-score ≤ -1 • Very low baseline BMD, defined as z-score ≤ -2 • 	Change in posterior-anterior spine BMD at 2 years <ul style="list-style-type: none"> • z-score ≤ 1 at baseline not significantly and independently associated with greater BMD loss (p-value = 0.81) • z-score ≤ 2 at baseline not significantly and independently associated with greater BMD loss (p-value = 0.47) 	Low
Brenckmnn 2003	<ul style="list-style-type: none"> • Baseline mean (SD) z-scores: • Left hip BMD: -0.9 (1.1) • Right hip BMD: -1.0 (1.1) • Lumbar spine BMD: -1.1 (1.3) • Baseline mean (SD) gm/cm²: • Total BMD: 1.2 (0.1) • 	BMD annual change: <ul style="list-style-type: none"> • Change in hip BMD: <ul style="list-style-type: none"> ◦ Left hip: -3.01% (95% CI -4.76 to -1.26) ◦ Right hip: -3.06% (95% CI -4.69 to -1.43) • Change in lumbar spine BMD: -0.86% (95% CI -2.46 to 0.75) • Change in total body BMD (n=21): 0.0% (SD 1.4%) 	Very low
Bhudhikanok 1998	<ul style="list-style-type: none"> • Baseline lumbar spine BMD (z-score for age and sex) • Males <ul style="list-style-type: none"> ◦ Children and young people < 18 years: -1.0\pm0.9 ◦ Adults \geq 18 years: -2.5\pm1.4 • Females <ul style="list-style-type: none"> ◦ Children and young people < 18 years: -1.5\pm1.5 ◦ Adults \geq 18 years: -1.9\pm1.6 • 	Change in lumbar spine BMD (z-score for age and sex) at mean 17 months follow-up <ul style="list-style-type: none"> • Males <ul style="list-style-type: none"> ◦ Children and young people < 18 years: -0.2\pm0.5; ns ◦ Adults \geq 18 years: 0.1\pm0.2; ns • Females <ul style="list-style-type: none"> ◦ Children and young people < 18 years: -0.6\pm0.8; p<0.05 ◦ Adults \geq 18 years: 0.1\pm0.3; ns 	Very low
	<ul style="list-style-type: none"> • Baseline femoral neck BMD (z-score for age and sex) 	Change in femoral neck BMD (z-score for age and sex) at mean 17 months follow-up	

Study	Prognostic indicator	Effect size	Quality
	<ul style="list-style-type: none"> • Males <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.8±0.6 ○ Adults ≥ 18 years: -2.5±0.8 • Females <ul style="list-style-type: none"> ○ Children and young people < 18 years: -2.0±1.6 ○ Adults ≥ 18 years: -2.2±1.6 • • Baseline whole body BMD (z-score for age and sex) • Males <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.3±0.5 ○ Adults ≥ 18 years: -2.0±1.2 • Females <ul style="list-style-type: none"> ○ Children and young people < 18 years: -1.3±1.2 ○ Adults ≥ 18 years: -1.3±1.2 • 	<ul style="list-style-type: none"> • Males <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.2±0.4; ns ○ Adults ≥ 18 years: -0.2±0.4; ns • Females <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.3±0.8; ns ○ Adults ≥ 18 years: 0.1±0.4; ns <p>Change in femoral neck BMD (z-score for age and sex) at mean 17 months follow-up</p> <ul style="list-style-type: none"> • Males <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.6±0.4; p<0.005 ○ Adults ≥ 18 years: 0.1±0.3; ns • Females <ul style="list-style-type: none"> ○ Children and young people < 18 years: -0.4±0.3; p<0.005 ○ Adults ≥ 18 years: -0.0±0.2; ns 	
Haworth 2002	<ul style="list-style-type: none"> • Baseline lumbar spine BMD (mg/ml by CQT): • Young cohort ≤ 24 years: 176.1 (166.9 to 185.2) • Adult cohort ≥ 25 years: 170.8 (161.3 to 180.3) • Baseline lumbar spine BMD (g/cm² by DXA): • Young cohort ≤ 24 years: 0.918 (0.8882 to 0.953) • Adult cohort ≥ 25 years: 0.942 (0.909 to 0.975) • Baseline femoral neck BMD (g/cm² by DXA): • Young cohort ≤ 24 years: 0.839 (0.801 to 0.877) • Adult cohort ≥ 25 years: 0.781 (0.756 to 0.819) • Baseline total hip BMD (g/cm² by DXA): • Young cohort ≤ 24 years: 0.917 (0.880 to 0.953) • Adult cohort ≥ 25 years: 0.881 (0.844 to 0.918) 	<ul style="list-style-type: none"> • Annual change in lumbar spine BMD (mg/ml by CQT): • Young cohort ≤ 24 years: -1.7% (-4.4 to 1.1) • Adult cohort ≥ 25 years: 0.7% (-1.5 to 2.8) • Annual change in lumbar spine BMD (g/cm² by DXA): • Young cohort ≤ 24 years: -0.9% (-2.0 to 0.2) • Adult cohort ≥ 25 years: -0.0% (-1.3 to 1.2) • Annual change in femoral neck BMD (g/cm² by DXA): • Young cohort ≤ 24 years: -2.5% (-3.8 to -1.2); p-value <0.001 • Adult cohort ≥ 25 years: -1.9% (-2.9 to -0.8); p-value <0.001 • Annual change in total hip BMD (g/cm² by DXA): • Young cohort ≤ 24 years: -2.2% (-3.3 to -1.0); p-value <0.001 • Adult cohort ≥ 25 years: -1.5% (-2.4 to -0.6); p=0.001 	Very low

Study	Prognostic indicator	Effect size	Quality
Papaioannou 2008	<ul style="list-style-type: none"> • Baseline mean (SD) T-score/ Z-score: • Lumbar spine BMD: -0.80 (1.10) • Proximal femur BMD: -0.57 (0.97) • Whole body BMD: -0.71 (1.11) 	Overall rate of bone loss at mean 4.3 years follow-up: <ul style="list-style-type: none"> • Lumbar spine BMD: -0.73% • Proximal femur BMD: -1.93% • Whole body BMD: -0.40% 	Very low
Schulze 2006	Baseline mean±SD gender and age matched z-scores: <ul style="list-style-type: none"> • Lumbar spine BMD: -0.40±1.13 • Whole body BMD: -0.29±1.01 	Change between baseline and follow-up (mean ±SD follow-up 2.13±1.16; range 1.06 to 4.10 years) <ul style="list-style-type: none"> • Lumbar spine BMD: -0.46±0.94 • Whole body BMD: -0.45±1.16 	Very low

1 BMD: bone mineral density; DXA: dual-energy X-ray absorptiometry; ns: not significant; SD: standard deviation; QCT: quantitative computed tomography

2

1 10.7.5 Economic evidence

2 No economic evaluations of strategies to identify reduced BMD were identified in the
3 literature search conducted for this guideline. Full details of the search economic article
4 selection flow chart can be found in Appendix E and F, respectively.

5 Diagnostic procedures, including imaging, will not be considered cost-effective if there is not
6 an effective treatment for the condition being diagnosed, or if the person's management is
7 not changed by the results. In other words, if the procedure does not add any additional
8 information to a clinical assessment and does not change the management strategy, those
9 procedures should not be recommended. However, to fully address the cost-effectiveness of
10 strategies (DXA scans and peripheral quantitative computed tomography (pQCT)) to detect
11 reduced BMD would require a model that included the management of reduced BMD which
12 lies outside the scope of this guideline. Moreover, the studies included in the clinical
13 evidence review compared the sites DXA scans could measure, as opposed to the accuracy
14 of DXA scans compared to another imaging procedure. As a result, if the resource use for
15 each site is equivalent, it is evident that the most accurate site should be measured and the
16 economic focus should be on the frequency those scans are performed.

17 According to the Committee, approximately one third of adults with cystic fibrosis have
18 reduced BMD which can lead to osteoporosis and predispose them to bone fractures. Those
19 complications can incur high treatment costs and negatively impact a person's quality of life
20 due to their reduced mobility and ability to perform their usual activities. For these reasons,
21 there are potential cost savings to the NHS if imaging can better identify bone disease
22 related to cystic fibrosis early in its course, resulting in more timely management and the
23 possible prevention of complications associated with reduced BMD.

24 To assess the risk of reduced BMD in people with cystic fibrosis, the Committee advised that
25 most people would receive regular DXA scans, but some may undergo a pQCT scan which
26 is more costly. The cost of those scans, according to NHS Reference Costs 2015, are
27 reported in Table 168.

28 **Table 168: Cost of imaging**

Currency code	Currency description	National average cost
Dexa Scan		
RD50Z	DXA scan	£59
Computerised tomography scan		
RD20A	one area, without contrast, 19 years and over	£93
RD20B	one area, without contrast, between 6 and 18 years	£93
RD20C	one area, without contrast, 5 years and under	£113
RD21A	one area, with post contrast only, 19 years and over	£104
RD21B	one area, with post contrast only, between 6 and 18 years	£111
RD21C	one area, with post contrast only, 5 years and under	£156
RD22Z	one area, with pre and post contrast	£113
RD23Z	two areas, without contrast	£121
RD24Z	two areas, with contrast	£122
RD25Z	three areas, without contrast	£107
RD26Z	three areas, with contrast	£124
RD27Z	more than three areas	£136
RD28Z	complex	£122

1 The Committee advised that it would be difficult to accurately judge a person's BMD prior to
2 a scan, based on clinical assessment of their risk factors. Therefore, if scans are necessary
3 to determine BMD, the Committee should consider when those scans are performed as their
4 frequency could have significant resource implications.

5 The Committee also advised that bone health is strongly linked to the severity of lung
6 disease, body weight and oral corticosteroids use. As a result, the Committee should
7 consider if there are populations which should undergo more frequent monitoring because
8 they have a higher risk of reduced BMD compared to the general population, whose
9 downstream costs would escalate much quicker if were are undetected. Finally, if pQCT
10 scans are more costly than DXA scans, the Committee should consider when pQCT scans
11 can provide additional information to a DXA scan to justify their additional cost.

12 **10.7.6 Evidence statements**

13 **10.7.6.1 Evidence in children and young people**

14 Very low quality evidence from 1 longitudinal study with 18 prepubertal and pubertal girls and
15 young females with CF (age range 7.6 to 17.9 years) suggested that:

- 16 • a baseline z-score of -0.40 in lumbar spine BMD (measured by DXA scan) was associated
17 with an overall change in BMD at the spine of -0.07 at 2.13 years follow-up
- 18 • a baseline z-score of -0.29 in whole body BMD (measured by DXA scan) was associated
19 with an overall change in whole body BMD of -0.17 at 2.13 years follow-up.

20 Very low quality evidence from 1 prospective longitudinal study with 20 children and young
21 people with CF (9 males, 11 females; age not reported) suggested that:

- 22 • a baseline z-score of -1.0 BMD at the lumbar spine (measured by DXA scan) was not
23 significantly associated with a decrease in BMD at the lumbar spine at 17 months follow-
24 up in boys and young males
- 25 • a baseline z-score of -1.5 BMD at the lumbar spine (measured by DXA scan) was
26 significantly associated with a decrease in BMD at the lumbar spine at 17 months follow-
27 up in girls and young females
- 28 • a baseline z-score of -0.8 BMD at the femoral neck (measured by DXA scan) was not
29 significantly associated with a decrease in BMD at the femoral neck at 17 months follow-
30 up in boys and young males
- 31 • a baseline z-score of -2.0 BMD at the femoral neck (measured by DXA scan) was not
32 significantly associated with a decrease in BMD at the femoral neck at 17 months follow-
33 up in girls and young females
- 34 • a baseline z-score of -0.3 in whole body BMD (measured by DXA scan) was significantly
35 associated with a decrease in whole body BMD at 17 months follow-up in boys and young
36 males
- 37 • a baseline z-score of -1.3 in whole body BMD (measured by DXA scan) was significantly
38 associated with a decrease in whole body BMD at 17 months follow-up in girls and young
39 females.

40 **10.7.6.2 Evidence in young people and young adults**

41 Very low quality evidence from 1 longitudinal study with 55 young people and young adults
42 with CF (age range 15 to 24 years) suggested that:

- 43 • baseline lumbar spine BMD (measured by CQT or DXA) was not significantly associated
44 with a decrease in lumbar spine BMD at 1 year follow-up
- 45 • baseline femoral neck BMD (measured by DXA) was significantly associated with a
46 decrease in femoral neck BMD at 1 year follow-up

- 1 • baseline total hip BMD (measured by DXA) was significantly associated with a decrease
2 in total hip BMD at 1 year follow-up.

3 10.7.6.3 Evidence in adults

4 Moderate quality evidence from 1 longitudinal study with 63 adults with CF (age range 18 to
5 57 years) suggested that low and very posterior-anterior spine BMD (defined as z-score \leq -1
6 and z-score \leq -2 and measured by DXA scan) were not significantly associated with greater
7 loss of posterior-anterior spine bone mineral density at 2 years follow-up.

8 Low quality evidence from 1 longitudinal study with 40 adults with CF (age range 18 to 52
9 years) suggested that:

- 10 • a baseline z-score of -0.9 BMD at the left hip (measured by DXA scan) was significantly
11 associated with a decrease in BMD at the left hip at 1 year follow-up
12 • a baseline z-score of -1.0 BMD at the right hip (measured by DXA scan) was significantly
13 associated with a decrease in BMD at the right hip at 1 year follow-up
14 • a baseline z-score of -1.1 BMD at the lumbar spine (measured by DXA scan) was not
15 significantly associated with a decrease in BMD at the lumbar spine at 1 year follow-up
16 • the change in total bone mineral density (measured by DXA scan) from baseline was
17 negligible at 1 year follow-up.

18 Very low quality evidence from 1 prospective longitudinal study with 21 adults with CF (6
19 males, 15 females; age not reported) suggested that:

- 20 • a baseline z-score of -2.5 BMD at the lumbar spine (measured by DXA scan) was not
21 significantly associated with a decrease in BMD at the lumbar spine at 17 months follow-
22 up in male adults
23 • a baseline z-score of -1.9 BMD at the lumbar spine (measured by DXA scan) was not
24 significantly associated with a decrease in BMD at the lumbar spine at 17 months follow-
25 up in adult females
26 • a baseline z-score of -2.5 BMD at the femoral neck (measured by DXA scan) was not
27 significantly associated with a decrease in BMD at the femoral neck at 17 months follow-
28 up in male adults
29 • a baseline z-score of -2.2 BMD at the femoral neck (measured by DXA scan) was not
30 significantly associated with a decrease in BMD at the femoral neck at 17 months follow-
31 up in adult females
32 • a baseline z-score of -2.0 in whole body BMD (measured by DXA scan) was not
33 significantly associated with a decrease in whole body BMD at 17 months follow-up in
34 male adults
35 • a baseline z-score of -1.3 in whole body BMD (measured by DXA scan) was not
36 significantly associated with a decrease in whole body BMD at 17 months follow-up in
37 adult females.

38 Very low quality evidence from 1 longitudinal study with 59 adults with CF (age \geq 25 years)
39 suggested that:

- 40 • baseline lumbar spine BMD (measured by CQT or DXA) was not significantly associated
41 with a decrease in lumbar spine BMD at 1 year follow-up
42 • baseline femoral neck BMD (measured by DXA) was significantly associated with a
43 decrease in femoral neck BMD at 1 year follow-up
44 • baseline total hip BMD (measured by DXA) was significantly associated with a decrease
45 in total hip BMD at 1 year follow-up.

46 Very low quality from 1 longitudinal study with 49 adults with CF (mean age 25.2 years)
47 suggested that:

- 1 • a baseline t-score/ z-score of -0.80 in lumbar spine BMD (measured by DXA scan) was
2 associated with an overall rate of bone loss at lumbar spine was -0.73% at 4.3 years
3 follow-up
- 4 • a baseline t-score/ z-score of -0.57 in proximal femur BMD (measured by DXA scan) was
5 associated with an overall rate of bone loss at proximal femur was -1.93% at 4.3 years
6 follow-up
- 7 • a baseline t-score/ z-score of -0.71 in whole body BMD (measured by DXA scan) was
8 associated with an overall rate whole body loss was -0.40% at 4.3 years follow-up.

9 **10.7.6.4 Economic evidence statement**

10 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

11 **10.7.7 Evidence to recommendations**

12 **10.7.7.1 Relative value placed on the outcomes considered**

13 The objective of this review was to determine the most effective strategy to monitor for the
14 identification of reduced mineral density in people with CF.

15 The Committee chose change in bone mineral density as critical outcome for decision
16 making; whereas number of fractures, and quality of life were rated as important outcomes.

17 **10.7.7.2 Consideration of clinical benefits and harms**

18 The Committee acknowledged that most CF Centres currently follow the European CF bone
19 mineralisation guidelines (Sermet-Gaudelus 2011). These guidelines recommend that a first
20 bone density scan should be done around the age of 8 to 10 years, and repeated every 1 - 5
21 years depending on severity (according to BMD t- or z-scores) and risk factors.

22 The Committee discussed the usefulness of conducting regular monitoring for bone mineral
23 density in children, as these tests are difficult to interpret in this population. Likewise, the
24 Committee also discussed whether routine DXA scans should be offered to all adults. They
25 noted that it is unlikely that people over 18 years can build bone mass, as this is mostly done
26 during teenage years, therefore there is little that can be done.

27 Careful liaison with a bone specialist should occur before commencing children or adults on
28 bisphosphonate therapy. This should include the optimal dose of treatment, duration and
29 monitoring of long-term effects. Patients should be counselled about the side-effects of
30 bisphosphates before commencing therapy. Careful consideration should also be given to
31 the use of bisphosphates in women of child-bearing age. Vitamin D deficiency and ensuring
32 an optimal calcium intake need to also be addressed prior to commencing bisphosphonates.
33 It is also important to minimise risk factors for low BMD e.g. improving nutritional status and
34 encouraging weight bearing exercise.

35 Even though it was not specifically addressed in the evidence, the Committee agreed that it
36 is recognised that certain groups across all age groups may actually benefit from monitoring,
37 as they are at a higher risk of reduced bone mineral density. These groups may include
38 people who present with under-nutrition, those using corticosteroids (either frequently or
39 long-term use), those with a history of low impact fracture, those who present exacerbations
40 that require intravenous antibiotics on regular basis, those with poor lung function (low FEV₁),
41 post- menopausal women and following transplants.

42 The Committee noted the available evidence did not provide any clarity in relation to which
43 body sites should be scanned, thus they agreed that a z-score of -2.0 standard deviations at
44 any body site would be considered abnormal and they recommended to seek expert advice.

1 The Committee agreed z-scores should be used in children, whereas z-scores or t-scores
2 can be used in adults (although t-scores are more useful in post-menopausal women and
3 men over 50 years).

4 10.7.7.3 Consideration of economic benefits and harms

5 Once presented with the clinical evidence and an outline of the resources scans entail, the
6 Committee questioned the value of monitoring for reduced BMD in current clinical practice,
7 stating that the absence of evidence does not mean current clinical practice should be
8 recommended when the opportunity cost of those resources is high.

9 The Committee noted that scans are useful to compare a person's BMD to the general
10 population as the results can infer if they are at a higher risk of fracture and subsequently,
11 potentially reduce the number of undiagnosed fractures and downstream costs they entail.
12 Despite this, the Committee agreed that management is rarely changed based on the results
13 obtained from a scan as people with cystic fibrosis should always be encouraged to optimise
14 calcium and vitamin D and participate in exercise. Moreover, clinicians are reluctant to
15 prescribe bisphosphonates to maintain BMD in young adults due to their negative effects on
16 fertility, iterating that a low BMD finding from a scan is unlikely to initiate such treatment. For
17 these reasons alone, it was evident to the Committee that the use of scans in current clinical
18 practice was not a cost-effective use of resources.

19 The Committee also added that DXA scans can overestimate reductions in BMD and if
20 people with reduced BMD are aware of their higher risk of fracture, they may stop exercising,
21 resulting in a counterproductive effect. Furthermore, even though the cost of one DXA scan
22 is not substantial, performing a DXA scan every 3 years in everyone with cystic fibrosis under
23 current practice has a substantial resource and cost use when the total is considered.

24 An alternative to a DXA scan is a pQCT scan, but the Committee advised that the use of
25 pQCT scans to monitor for BMD remains a research area as their interpretation is not fully
26 understood by clinicians in clinical practice. It was also noted that pQCT scans can emit
27 radiation and are more expensive to perform than DXA scans due to the extra time and
28 equipment they require.

29 Overall, the Committee agreed that monitoring for reduced BMD would not be considered
30 cost-effective in all cases and subsequently identified populations deemed to be at a higher
31 risk of reduced BMD who could benefit from a DXA scan.

32 In those populations, the Committee agreed scans could be repeated on an annual basis if
33 the BMD SDS score is less than -2.00 to know if treatment to prevent reduced BMD should
34 to be escalated or stopped, or if specialist advice should be sought regarding
35 bisphosphonate treatment.

36 The Committee considered performing a DXA scan after the transition from paediatric to
37 adult services for a baseline BMD measure and then as necessary. However, the Committee
38 noted that weight can change BMD quite markedly questioning the value of a baseline scan
39 that could soon be outdated. As a result, a recommendation was only prioritised to monitor
40 BMD in high risk populations. The Committee advised that this would lead to a large change
41 in current practice and result in cost savings, but would not compromise on health benefit as
42 current practice was not a cost-effective use of resources.

43 10.7.7.4 Quality of evidence

44 Prospective and retrospective observational studies were included in the review. The quality
45 of the evidence, as assessed per individual studies using a critical appraisal tool for the
46 evaluation of the quality of prognosis studies in systematic reviews (Hayden et al. 2006) was
47 very low. The main sources of bias in the studies were:

- 48
- poorly reported sample selection, with unclear inclusion and exclusion criteria;

- 1 • the study sample did not fully represent the population of interest with regard to key
2 characteristics,
3 • important potential confounders were not appropriately accounted for.
- 4 Due to all these limitations, the Committee considered that the usefulness of the data
5 reported in the studies is very limited.

6 10.7.7.5 Other considerations

7 Due to the scarcity and low quality of the evidence, the recommendations were mainly based
8 on Committee members' clinical experience, and consensus on good clinical practice.

9 The Committee acknowledge there is a lack of longitudinal studies to evaluate the impact of
10 low bone mineral density on fractures and quality of life. However, it is known that people
11 with CF are an increased risk of developing osteopenia and osteoporosis (See chapter on
12 complications of cystic fibrosis). They were also aware of cross-sectional evidence showing a
13 correlation between low bone mineral density and decreased lung function and poor
14 nutritional status. However, they noted that although the main concern is the risk of fractures,
15 the incidence of fractures in this population is not particularly high.

16 As discussed previously, the Committee agreed the absence of evidence showing a benefit
17 in conducting regular scans and the fact that people receive treatment anyway, justifies the
18 change in practice. Moreover conducting routine exams can create anxiety. However, they
19 also discussed that people with CF and their carers may feel concerned about a change in
20 practice. They suggested it is important to talk to the families and discuss their concerns.

21 No equality issues were identified by the Committee for this review question.

22 The Committee noted that evidence regarding the impact of low bone mineral density in
23 fractures was mostly conducted in post-menopausal women. They agreed it would be helpful
24 if a study could be conducted looking at people with CF and highlighted the importance in
25 assessing fractures and quality of life. However, it was agreed that this was not a priority
26 research recommendation for the guideline as a whole.

27 10.7.7.6 Key conclusions

28 The Committee concluded that there is no need to perform routine bone density scans to all
29 children and adults with CF. DXA scans should only be considered in children and adults
30 who are at risk of low bone mineral density. This is because there is little value in routine
31 monitoring, as children and young people get treated to help accrue bone mass in spite of
32 the results of the test.

33 10.7.8 Recommendations

34 **122. Consider dual energy X-ray absorptiometry (DXA) bone density scans for**
35 **people with cystic fibrosis who have factors that put them at high risk of low bone**
36 **mineral density, such as:**

- 37 • frequent or long-term oral corticosteroid use
38 • frequent intravenous antibiotic use
39 • severe lung disease
40 • undernutrition
41 • previous low-impact fractures
42 • previous transplants
43 • post menopause.

1 **123. Seek specialist advice for people with a bone mineral density standard**
2 **deviation below -2.0 (Z score) or -2.5 (T score).**

3 **10.8 Exercise**

4 **Review question: What is the effectiveness of programmes of exercise in the**
5 **management of cystic fibrosis?**

6 **10.8.1 Introduction**

7 Physical Exercise programmes are widely recommended as a routine part of the
8 management of cystic fibrosis. In addition to the benefits found by the general population,
9 exercise participation is advised in cystic fibrosis to help maintain and slow the decline in
10 respiratory function, facilitate airway clearance techniques, help improve bone mineral
11 density and to maintain and/or increase muscle strength, flexibility and posture.

12 Improving and maintaining cardiovascular fitness is very important for people with cystic
13 fibrosis. Regular education, monitoring and assessment of fitness from the point of diagnosis
14 allow for programmes to be tailored to the individual through the spectrum disease to meet
15 specific needs

16 People with cystic fibrosis can experience a number of barriers to full engagement in regular
17 exercise programmes due to physical limitations such as fatigue, frequent exacerbations and
18 the treatment burden of medications, nebulisers and physiotherapy. Environmental and
19 socioeconomic factors can also limit participation in exercise such as access to appropriate
20 facilities and support both in the community and in the hospital

21 Exercise education and prescription has been a routine part of clinical care in cystic fibrosis
22 for many years however there is extensive variability in content, supervision and frequency of
23 the intervention. This review question aims to evaluate if specific programmes of physical
24 exercise lead to improve outcomes and benefits for people with cystic fibrosis, taking into
25 consideration a variety of activity interventions, including supervised and unsupervised
26 programmes.

27 **10.8.2 Description of clinical evidence**

28 The aim of this review was to determine the effectiveness of different exercise programmes
29 in improving health outcomes for people with cystic fibrosis.

30 The interventions reviewed were: aerobic exercise programme; strength resistance training;
31 high intensity interval training; habitual physical activity; inspiratory muscle training
32 (performed at maximal inspiratory effort of at least 80%) or any combination of these
33 interventions.

34 We looked for systematic reviews of randomized controlled trials (RCTs) and RCTs.
35 Systematic reviews were assessed for inclusion against the protocol, and if relevant, their
36 quality was assessed using AMSTAR. High-quality systematic reviews were included in our
37 review, and where possible data and quality assessment were taken directly from the review.
38 Individual studies were retrieved for completeness and accuracy, and were also checked for
39 additional outcomes. Low-quality SR were excluded from our review, but the list of included
40 studies was checked to identify relevant trials.

41 Given that no evidence was found for high intensity interval training or time to next
42 exacerbation (a critical outcome) in the RCTs, cohort studies were assessed for inclusion
43 and data were reported from these studies only in relation to this intervention or this critical
44 outcome not covered by the RCTs.

1 Although there were no RCTs on habitual physical activity, cohort studies relating to this
2 intervention were not assessed because the Committee thought that the information from
3 RCTs on long term exercise programmes was sufficient to formulate recommendations. Only
4 1 study on habitual physical activity was included (Cox 2016) because it looked at the
5 outcome need for hospitalization, which was considered appropriate as a proxy outcome for
6 time to next exacerbation.

7 For full details see review protocol in Appendix D.

8 Three Cochrane reviews were identified in the search (Cox 2013, Houston 2013, Radtke
9 2015).

10 Two reviews were included in the review:

- 11 • Radtke (2015) evaluated the effectiveness of physical exercise training compared to no
12 training. 10 RCTs were included from this review (Hebestreit 2010, Hommerding 2015,
13 Klijn 2004, Kriemler 2013, Moorcroft 2004, Rovedder 2014, Santana-Sosa 2012, Santana-
14 Sosa 2014, Schneiderman-Walker 2000, Selvadurai 2002)
- 15 • Houston (2013) evaluated the effectiveness of inspiratory muscle training compared to no
16 training. 1 RCT was included from this review (Enright 2004)

17 One review was excluded:

- 18 • Cox (2013) evaluated interventions for promoting physical activity in people with cystic
19 fibrosis. No additional trials were included from this review, as they were included in
20 Radtke (2015).

21 In addition, 3 RCTs (Beaudoin 2016, Orenstein 2004, Schindel 2015) and 2 cohort studies
22 (Cox 2016, Gruber 2014) were included.

23 The size of the RCTs or cohort studies ranged from 14 to 67 participants. 5 studies included
24 adults (Beaudoin 2016, Cox 2016, Enright 2004, Gruber 2014, Moorcroft 2004), 1 young
25 people aged ≥ 16 years and adults (Rovedder 2014), 2 young people aged >12 years and
26 adults (Hebestreit 2010, Kriemler 2013), 6 children and young people (Hommerding 2015,
27 Klijn 2004, Orenstein 2004, Santana-Sosa 2012, Santana-Sosa 2014, Selvadurai 2002), 2
28 children, young people and adults (Schindel 2015; Schneiderman-Walker 2000)

29 2 studies were conducted in the UK (Enright 2004, Moorcroft 2004), 3 in Brazil (Hommerding
30 2015, Rovedder 2014, Schindel 2015), 2 in Germany (Hebestreit 2010, Gruber 2014), 2 in
31 Spain (Santana-Sosa 2012, Santana-Sosa 2014), 2 in Canada (Beaudoin 2016,
32 Schneiderman-Walker 2000), 2 in Australia (Cox 2016, Selvadurai 2002), 1 in the
33 Netherlands (Klijn 2004), 1 in Switzerland (Kriemler 2013), 1 in the USA (Orenstein 2004).

34 6 studies assessed a supervised exercise programme (Gruber 2014, Klijn 2004, Orenstein
35 2004, Santana-Sosa 2014, Santana-Sosa 2012, Selvadurai 2002). 8 studies assessed a
36 partially supervised or unsupervised exercise programme (Beaudoin 2016, Hebestreit 2010,
37 Hommerding 2015, Kriemler 2013, Moorcroft 2004, Rovedder 2014, Schindel 2015,
38 Schneiderman-Walker 2000). 1 study assessed supervised inspiratory muscle training
39 (Enright 2004). 1 study assessed habitual physical activity (Cox 2016)

40 1 study compared a supervised aerobic exercise programme for about 19 days to no
41 exercise programme (Selvadurai 2002). 3 studies compared an unsupervised aerobic
42 exercise programme to no exercise programme (Hommerding 2015, Kriemler 2013,
43 Schneiderman Walker 2000). Depending on the study, training lasted 3 months
44 (Hommerding 2015); 6 months (Kriemler 2013) or 3 years (Schneiderman-Walker 2000).

45 2 studies compared a supervised strength resistance / anaerobic training programme to no
46 exercise programme (Selvadurai 2002, Klijn 2004). Training lasted either about 19 days
47 (Selvadurai 2002) or 12 weeks (Klijn 2004). 1 study compared an unsupervised strength

- 1 resistance / anaerobic training programme for 6 months to no exercise programme (Kriemler
2 2013).
- 3 2 studies compared a supervised strength / anaerobic training programme to a supervised
4 aerobic programme (Selvadurai 2002, Orenstein 2004). Training lasted either about 19 days
5 (Selvadurai 2002) or 1 year (Orenstein 2004). One study compared an unsupervised
6 strength / anaerobic training programme for 6 months to an unsupervised aerobic
7 programme for 6 months (Kriemler 2013).
- 8 1 study compared compared a supervised high-intensity interval training for 6 weeks to
9 standard aerobic and anaerobic exercise programme for 6 weeks (Gruber 2014).
- 10 1study compared a supervised combined aerobic and anaerobic training programme for 8
11 weeks to no exercise programme (Santana-Sosa 2012).
- 12 5 studies compared an unsupervised combined aerobic and anaerobic training programme to
13 no exercise programme (Beaudoin 2016, Hebestreit 2010, Moorcroft 2004, Rovedder 2014,
14 Schindel 2015). Depending on the study, duration of training was 12 weeks (Beaudoin 2016),
15 3 months (Rovedder 2014 and Schindel 2015), 6 months (Hebestreit 2010), or 1 year
16 (Moorcroft 2004).
- 17 1 study compared a supervised combined inspiratory muscle training, resistance and aerobic
18 training for 8 weeks to no exercise programme (Santana-Sosa 2014).
- 19 1 study compared supervised inspiratory muscle training at 80% of maximal effort for 8
20 weeks to usual care (Enright 2004).
- 21 1 study compared a higher amount or longer duration of habitual physical activity to a smaller
22 amount or shorter duration (Cox 2016).
- 23 A summary of the included studies is presented in Table 169. See also study selection flow
24 chart in Appendix F, study evidence tables in Appendix G, list of excluded studies in
25 Appendix H, forest plots in Appendix I, and full GRADE profiles in Appendix J.

26 10.8.3 Summary of included studies

27 A summary of the studies that were included in this review are presented in Table 169.

28 **Table 169: Summary of included studies**

Study	Intervention/Comparison	Population	Outcomes	Comments
Cochrane systematic reviews				
Radtke 2015 Cochrane SR	Comparison 1: Aerobic exercise programme versus no defined exercise programme (Selvadurai 2002, Hommerding 2015, Kriemler 2013, Schneiderman-Walker 2000) Comparison 2: Strength resistance/ anaerobic training versus no defined exercise programme (Selvadurai 2002, Klijn 2004, Kriemler 2013) Comparison 3: Combined aerobic and strength	People with cystic fibrosis of any age and any degree of disease severity	Comparison 1: Aerobic training versus no physical training • FEV ₁ % predicted • FVC % predicted • VO ₂ peak • BMI • Not reported: • Time to next exacerbation • Quality of life • Adverse events • Preference	Santana-Sosa 2014 looks at combined IMT + resistance + aerobic

Study	Intervention/Comparison	Population	Outcomes	Comments
	resistance training versus no defined exercise programme (Hebestreit 2010, Moorcroft 2004, Rovedder 2014, Santana-Sosa 2012, Santana-Sosa 2014)		<p>Comparison 2: Anaerobic training versus no physical training</p> <ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted • VO₂ peak • QoL (CFQ) • BMI <p>Not reported:</p> <ul style="list-style-type: none"> • Time to next exacerbation • Adverse events • Preference <p>Comparison 3: Combined aerobic and anaerobic training versus no physical training</p> <ul style="list-style-type: none"> • FEV₁ % predicted • FVC % predicted • VO₂ peak • QoL (CFQ-R) • BMI • Adverse events <p>Not reported:</p> <ul style="list-style-type: none"> • Time to next exacerbation • Preference 	
Houston 2013 Cochrane SR	Comparison 1. Inspiratory muscle training (80% maximal effort) versus control (Enright 2004)	People with cystic fibrosis of any age	<p>Comparison 1. Inspiratory muscle training (80% maximal effort) versus control</p> <ul style="list-style-type: none"> • FEV₁ (I) • FVC (I) • Not reported: • VO₂ • Time to next exacerbation • Body composition • Preference • Quality of life • Adverse events 	Only data for IMT80% was included
Primary studies included in the SRs				
Enright 2004 (UK) RCT	<p>Intervention: IMT at 80% of "maximal inspiratory effort"</p> <ul style="list-style-type: none"> • direct supervision at home by designated training • IMT is incremental maximal effort with progressively 	<p>N=19 adults with cystic fibrosis</p> <p>Age, mean (SD): 22 (4.2) years</p>	<ul style="list-style-type: none"> • Change in FEV₁ (I) • Change in FVC (I) 	Included in Houston 2013 SR

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>shorter rest periods</p> <ul style="list-style-type: none"> • 3 times a week for 8 weeks <p>Control: no training</p>	<ul style="list-style-type: none"> • IMT group: n=9 • Control group: n=10 		
Hebestreit 2010 (Germany) RCT	<p>Intervention: endurance-type and strengthening exercises</p> <ul style="list-style-type: none"> • Unsupervised programme • Participants agreed to increase their vigorous physical activities by a minimum of 3x 60 min per week in the first 6 months of the study. An individual exercise plan was devised for participants; activity counselling was stopped after the first 6 months and participants were encouraged to maintain or further increase their physical activity level <p>Control:</p> <ul style="list-style-type: none"> • Participants told to keep their activity level constant during the first 12 months of the study. During the second year (period from 12 - 24 months) they were free to change their activity behaviour 	<p>N=38 people with cystic fibrosis >12 years</p> <ul style="list-style-type: none"> • Exercise group n=23; mean (SD) age: 19.5 (6.4) years • Control group n=15; mean (SD) age: 19.4 (5.3) years 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in FVC % predicted • Change in VO₂ peak • Change in BMI 	Included in Radtke 2015 SR and Cox 2013 SR Data for > 6 months was not considered, as training lasted 6 months
Hommerding 2015 (Brazil) RCT	<p>Intervention: aerobic exercise programme</p> <ul style="list-style-type: none"> • Unsupervised programme • Included jogging, swimming, walking, ball games and stretching exercises. • based on verbal and written guidelines • twice a week for at least 20 min for 3 months • participants received telephone calls every 2 weeks and instructions were provided by 1 of the authors <p>Control: usual care</p> <ul style="list-style-type: none"> • participants were instructed about aerobic exercises once at baseline according to the CF centre routine 	<p>N=34 children and young people with cystic fibrosis</p> <ul style="list-style-type: none"> • Exercise group n=17; mean (SD) age 13.4 (2.8) years • Control group n=17; mean (SD) age 12.7 (3.3) years 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in FVC % predicted • Change in VO₂ peak 	Included in Radtke 2015 SR
Klijn 2004 (Netherlands) RCT	<p>Intervention: anaerobic training</p> <ul style="list-style-type: none"> • Supervised programme • 2 days per week for 30 to 45 min • 12 weeks 	<p>N=20 children and young people with cystic fibrosis with stable disease</p>	<ul style="list-style-type: none"> • Change in VO₂ peak • Change in quality of life 	Included in Radtke 2015 SR and Cox 2013 SR 3

Study	Intervention/Comparison	Population	Outcomes	Comments
	<p>Control: normal daily activities</p> <ul style="list-style-type: none"> • 12 weeks 	<ul style="list-style-type: none"> • Intervention group n=11; mean (SD) age 13.6 (1.3) years • Control group n=9; mean (SD) age 14.2 (2.1) years 		<p>participants dropped out; 1 withdrew from the training group for practical reasons</p>
<p>Kriemler 2013 (Switzerland) RCT</p>	<p>Intervention 1: aerobic training</p> <ul style="list-style-type: none"> • Unsupervised programme • 24-months: 6-month intervention and long-term follow-up period • 3 sessions per week of 30 to 45 minutes for 6 months and received support which was stopped thereafter <p>Intervention 2: strength training</p> <ul style="list-style-type: none"> • Unsupervised programme • 24-months: 6-month intervention and long-term follow-up period • 3 sessions per week of 30 to 45 minutes for the first 6 months and received support which was stopped thereafter <p>Control: no programme</p> <ul style="list-style-type: none"> • Participants in the control group were told to keep their activity level constant. • Free access to a fitness centre for 1 year after the first study year 	<p>N=39 participants with cystic fibrosis >12 years</p> <ul style="list-style-type: none"> • Aerobic training group (n=17): mean (95% CI) age 23.8 (21.5 to 26.5) years • Strength training group (n=12): mean (95% CI) age 19.0 (16.0 to 22.0) years • Control group (n=10): mean (95% CI) age 20.3 (17.0 to 23.6) years. • A separate control group from a parallel study (Hebestreit 2010) was added due to an unusual deterioration of physical health in the control group in this study (n=15), mean (95% CI) age 19.5 (16.8 to 22.2) years 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in FVC % predicted • Change in VO₂ peak • Change in BMI 	<p>Included in Radtke 2015 SR Long-term exercise study.</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
Moorcroft 2004 (UK) RCT	<p>Intervention: aerobic exercise</p> <ul style="list-style-type: none"> Unsupervised programme exercise based on individual preferences general aerobic exercises for lower body and weight training for upper body) for 12 months 3 times per week <p>Control: usual activities</p> <ul style="list-style-type: none"> Continue with usual activities 	<p>N=51 adults with cystic fibrosis</p> <ul style="list-style-type: none"> Exercise group (n=30): mean (SD) age 23.5 (6.4) years. Control group (n=18): 23.6 (5.5) years. 	<ul style="list-style-type: none"> Change in BMI 	<p>Included in Radtke 2015 SR</p> <p>Long-term study over 1 year. 42 completed the study</p>
Rovedder 2014 (Brazil) RCT	<p>Intervention: aerobic and muscle strengthening exercises</p> <ul style="list-style-type: none"> unsupervised programme 3-month home-based exercise programme. printed guidance advised to perform the programme on a daily basis weekly telephone contacts were <p>Control: no programme</p> <ul style="list-style-type: none"> standard follow-up from a physiotherapist without any specific exercise instructions 	<p>N=41 people with cystic fibrosis ≥ 16 years</p> <ul style="list-style-type: none"> Exercise group n=19; mean (SD) age 23.8 (8.3) years. Control group n=22; mean (SD) age 25.4 (6.9) years. 	<ul style="list-style-type: none"> Change in FEV₁ % predicted Change in FVC % predicted Change in Quality of life (CFQ-R) 	<p>Included in Radtke 2015 SR</p>
Santana-Sosa 2012 (Spain) RCT	<p>Intervention: endurance and strengthening exercises</p> <ul style="list-style-type: none"> supervised programme 8-week intrahospital programme followed by a 4-week detraining period 3 times per week same chest physiotherapy <p>Control: no programme</p> <ul style="list-style-type: none"> Participants were instructed on the positive effects of regular physical activity 	<p>N=22 children and young people with cystic fibrosis.</p> <ul style="list-style-type: none"> Exercise group n=11; mean (SEM, range) age 11 years (3 years, 5 - 15 years) Control group n=11; mean (SEM, range) age 10.0 years (2 years, 6 to 14 years) 	<ul style="list-style-type: none"> Change in Quality of life (CFQ-R children's and parents' report) Adverse events 	<p>Included in Radtke 2015 SR</p>
Santana-Sosa 2014 (Spain) RCT	<p>Intervention: aerobic + anaerobic + IMT</p> <ul style="list-style-type: none"> supervised programme 8-week programme followed by a 4-week detraining period whole body aerobic and weight training 3 times per week 	<p>N=20 children and young people with cystic fibrosis</p> <p>Age: 6 to 17 years</p> <ul style="list-style-type: none"> Exercise group n=10; mean (SEM) 	<ul style="list-style-type: none"> Change in FEV₁ (l) Change in FVC (l) Change in Quality of life (CFQ-R) Change in weight (kg) 	<p>Included in Radtke 2015 SR</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
	<ul style="list-style-type: none"> plus 2 daily IMT sessions same chest physiotherapy <p>Control: low intensity IMT</p>	<p>age 11.1 (1.1) years.</p> <ul style="list-style-type: none"> Control group n=10; mean (SEM) age 10.1 (1.1) years. 	<ul style="list-style-type: none"> Adverse events 	
Schneiderman-Walker 2000 (Canada) RCT	<p>Intervention 1: aerobic programme</p> <ul style="list-style-type: none"> unsupervised programme home programme Minimum of 20 min 3 times per week for 3 years <p>Control: maintained regular activity</p>	<p>N=65 people with cystic fibrosis</p> <p>Age: 7 to 19 years</p> <ul style="list-style-type: none"> Exercise group (n=30): mean (SD) age 13.4 (3.9) years. Control group (n=35): mean (SD) age 13.3 (3.6) years. 	<ul style="list-style-type: none"> Change in FEV₁ % predicted Change in FVC % predicted 	<p>Included in Radtke 2015 SR and Cox 2013 SR</p> <p>Long-term study</p> <p>2 groups similar at baseline.</p> <p>7 dropouts.</p>
Selvadurai 2002 (Australia) RCT	<p>Intervention 1: aerobic training</p> <ul style="list-style-type: none"> supervised programme 30 min, 5 times per week Training during hospital admission; mean (SD) duration of admission: 18.6 (3.9) days <p>Intervention 2: resistance training</p> <ul style="list-style-type: none"> supervised programme 30 min, 5 times per week Training during hospital admission; mean (SD) duration of admission: 18.8 (4.1) days <p>Control: no specific training</p>	<p>N=66 children and young people with cystic fibrosis aged 8 to 16 years admitted to hospital due to a pulmonary exacerbation</p> <ul style="list-style-type: none"> Aerobic training group (n=22): mean (SD) age 13.2 (2.0) years Resistance training group (n=22): mean (SD) age 13.1 (2.1) years Control group (n=22): mean (SD) age 13.2 (2.0) years 	<ul style="list-style-type: none"> Change in FEV₁ % predicted Change in FVC % predicted Change in VO₂ peak 	<p>Included in Radtke 2015 SR and Cox 2013 SR</p> <p>No dropouts.</p>
Additional primary studies				
Beaudoin	Intervention: Combined aerobic	N= 14 adults	• Change in	18 adults

Study	Intervention/Comparison	Population	Outcomes	Comments
2016 (Canada) RCT	<p>and resistance training programme</p> <ul style="list-style-type: none"> Unsupervised programme Both aerobic and resistance training: 3 times per week for 12 weeks Aerobic training: 20 to 40 min; resistance training: 5 to 7 exercises for a progressively increasing number of sets and repetitions <p>Control: no specific training</p>	<p>with cystic fibrosis aged ≥ 18 years with glucose abnormality;</p> <ul style="list-style-type: none"> Exercise group (n=8): mean age 31.9; age range 24 to 41 Control group (n=6): mean age 35.5; age range 22 to 57 	<p>FEV₁% predicted</p> <ul style="list-style-type: none"> Change in quality of life (measured with CFQ-R) Change in FVC % predicted Change in VO₂ peak Change in BMI Change in weight 	<p>were recruited; 17 were randomized; 2 dropped out because of pulmonary exacerbations; 1 was excluded because he was noncompliant</p> <ul style="list-style-type: none">
Cox 2016 (Australia) Prospective cohort study	<p>Comparison 1.</p> <ul style="list-style-type: none"> Intervention: ≥ 30 minutes daily of habitual moderate-vigorous physical activity Control: < 30 minutes daily of habitual moderate-vigorous physical activity <p>Comparison 2.</p> <ul style="list-style-type: none"> Intervention 2: ≥ 30 minutes daily of habitual moderate-vigorous physical activity accumulated in bouts of > 10 minutes Control 2: < 30 minutes or ≥ 30 minutes of habitual moderate-vigorous physical activity not accumulated in bouts of > 10 minutes 	<ul style="list-style-type: none"> N=61 adults with cystic fibrosis aged ≥ 18 years Intervention: n=33 Control: n=28 	<ul style="list-style-type: none"> Need for hospitalization, n (%) during 12 months 	<p>Physical activity was measured over 5-7 days using a portable multi-sensor armband; moderate PA intensity was classified as ≥ 4.8 metabolic equivalents. People were considered to have reached physical activity in bouts of at least 10 minutes duration if said bouts were recorded on any 1 day in the monitoring period.</p>

Study	Intervention/Comparison	Population	Outcomes	Comments
				65 adults were recruited; 4 were excluded because they wore the armband for insufficient time at baseline
Gruber 2014 (Germany) Cohort study	<p>Intervention 1: interval-training</p> <ul style="list-style-type: none"> • supervised programme • interval-training treadmill program • high intensity bouts alternated with active recovery phases • 5 times weekly for 6 weeks <p>Intervention 2: Standard aerobic + anaerobic exercise program</p> <ul style="list-style-type: none"> • supervised programme • 5 times per week for 45 minutes, for 6 weeks • activities based on participants' fitness level • including prolonged endurance exercise in terms of walking or Nordic-Walking complemented by ball games, stretching, balance training, and resistance training 	<p>N=43 rehabilitation clinic cystic fibrosis adult inpatients with FEV₁<40% predicted Age: overall age not reported</p> <ul style="list-style-type: none"> • Interval training group: n=20. Mean (SD) age: 26.4 (7.5) • Standard exercise group: n=23. Mean (SD) age: 26.3 (9.9) 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in VC % predicted • Change in VO₂ (ml/kg/min) • Change in BMI • 	<p>Patients who were unable to participate in standard exercise programme (Subjects who de-saturated (SpO₂ < 90%) at very low power (≤0.3 W/kg) or had a SpO₂ ≤ 90% at rest) were allocated to high intensity interval training Not stated explicitly if prospective or retrospective, seems prospective</p>
Orenstein 2004 (USA) RCT	<p>Intervention 1: upper-body strength training</p> <ul style="list-style-type: none"> • supervised programme • individually tailored to the participants' characteristics • participants were given an upper-body-only-weight-resistance machine 	<p>N=67 children and young people with cystic fibrosis Age: 8 to 18 years</p> <ul style="list-style-type: none"> • Exercise group: n=26) • Control 	<ul style="list-style-type: none"> • Change in FEV₁ % predicted • Change in VO₂ 	

Study	Intervention/Comparison	Population	Outcomes	Comments
	<ul style="list-style-type: none"> at least 3 times per week, for 1 year home visits once a week for the first 8 weeks, followed by monthly visits instructed to perform biceps curls, lateral pull-downs, and military and bench presses instructed to keep their heart rate <55% of their maximum <p>Intervention 2: aerobic training</p> <ul style="list-style-type: none"> supervised programme each child was given a stair-stepping machine at least 3 times per week, for 1 year home visits once a week for the first 8 weeks, followed by monthly visits instructed to exercise 5 minutes per session, gradually increasing to 30 minutes taught to gradually increase their target heart rate to 70% of their maximum heart rate 	group: n=30		
Schindel 2015 (Brazil) RCT	<p>Intervention: aerobic exercise and stretching</p> <ul style="list-style-type: none"> supervised programme 3-month-long exercise programme instruction handbook calendar where patients marked the days they performed exercise at least 3 times per week for a minimum of 20 minutes and perform each stretch 2 times for 20 seconds each phone calls from the researcher every 2 weeks <p>Control: usual care</p> <ul style="list-style-type: none"> Verbal orientations to perform exercise and stretching 	<p>N=34 people with cystic fibrosis</p> <p>Age range: 7 to 20 years</p> <ul style="list-style-type: none"> Exercise group: n=17. Mean (SD) age: 13.6 (2.8) years Control group: n=17. Mean (SD) age: 12.9 (3.9) years 	<ul style="list-style-type: none"> Change in FEV₁ % predicted Change in FVC % predicted 	

1
2
3
BMI: body mass index; CF: cystic fibrosis; FVC: forced vital capacity; FEV 1: forced expiratory volume in 1 second; IMT: inspiratory muscle training; RCT: randomised controlled trial; SR: systematic review; VC: vital capacity

4 10.8.4 Clinical evidence profile

5 The clinical evidence profiles for this review question (exercise in people with cystic fibrosis)
6 are presented in Table 170 to Table 177.

1
2

Table 170: Summary clinical evidence profile: Comparison 1. Aerobic exercise training programme versus no exercise programme

Comparison 1. Aerobic exercise training programme compared to no exercise programme for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise programme	Aerobic exercise training programme				
[Supervised programme] Change in FEV ₁ % predicted - Scale from: 0 to 100. Follow-up: at hospital discharge, mean 18.7 days	The mean change in FEV ₁ % predicted in the control group was 4.51	The mean change in FEV ₁ % predicted in the aerobic exercise training programme groups was 2.03 higher (2.31 lower to 6.37 higher)		44 (Selvadurai 2002)	⊕⊕⊕⊕ low ^{1,2}	
[Unsupervised programme] Change in FEV ₁ % predicted - Scale from: 0 to 100. Follow-up: 3 months	The mean change in FEV ₁ % predicted in the control group was 1 in 1 study, - 7.92 in the other study	The mean change in FEV ₁ % predicted in the aerobic exercise training programme groups was MD 5.23 higher (10.06 lower to 20.52 higher)		58 (Hommerding 2015, Kriemler 2013)	⊕⊕⊕⊕ very low ^{3,4,5}	
[Unsupervised programme] Change in FEV ₁ % predicted - Scale from: 0 to 100. Follow-up: 6 months	The mean change in FEV ₁ % predicted in the control group was - 11	The mean change in FEV ₁ % predicted in the aerobic exercise training programme groups was 17.17 higher (8.59 to 25.75 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ low ⁶	
[Unsupervised programme] Change in FEV ₁ % predicted - Scale from: 0 to 100. Follow-up: 3 years	The mean change in FEV ₁ % predicted in the control group was - 3.47	The mean change in FEV ₁ % predicted - unsupervised programme in the intervention groups was 2.01 higher (0.06 lower to 4.08 higher)		65 (Schneiderman-Walker 2000)	⊕⊕⊕⊕ moderate ⁷	
[Supervised programme] Change in FVC % predicted - Scale from: 0 to 100.	The mean change in FVC ₁ % predicted in the control group was	The mean change in FVC % predicted in the aerobic exercise training programme		44 (Selvadurai 2002)	⊕⊕⊕⊕ very low ^{1,8}	

Comparison 1. Aerobic exercise training programme compared to no exercise programme for cystic fibrosis						
Follow-up: at hospital discharge, mean 18.7 days	2.28	groups was 0.06 higher (2.55 lower to 2.67 higher)				
[Unsupervised programme] Change in FVC % predicted - Scale from: 0 to 100. Follow-up: 3 months	The mean change in FVC1% predicted in the control group was 2 in 1 study. - 5.57 in the other study	The mean change in FVC % predicted in the aerobic exercise training programme groups was 3.99 higher (6.62 lower to 14.61 higher)		58 (Hommerding 2015, Kriemler 2013)	⊕⊕⊕⊕ very low3,8,9	
[Unsupervised programme] Change in FVC % predicted - Scale from: 0 to 100. Follow-up: 6 months	The mean change in FVC1% predicted in the control group was - 7.85	The mean change in FVC % predicted in the aerobic exercise training programme groups was 12.51 higher (5.9 to 19.12 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ low6	
[Unsupervised programme] Change in FVC % predicted - Scale from: 0 to 100. Follow-up: 3 years	The mean change in FVC1% predicted in the control group was - 2.42	The mean change in FVC % in the aerobic exercise training programme in the intervention groups was 2.17 higher (0.47 to 3.87 higher)		65 (Schneiderman-Walker 2000)	⊕⊕⊕⊕ low7,10	
[Supervised programme] Change in VO ₂ peak - ml/min per kg BW Follow-up: at hospital discharge, mean 18.7 days	The mean change in VO ₂ in the control group was - 1.22	The mean change in VO ₂ peak in the aerobic exercise training programme groups was 8.53 higher (4.85 to 12.21 higher)		44 (Selvadurai 2002)	⊕⊕⊕⊕ moderate 1	
[Unsupervised programme] Change in VO ₂ peak - ml/min per kg BW Follow-up: 3 months	The mean change in VO ₂ in the control group was 2.3 in 1 study, -2.45 in the other study.	The mean change in VO ₂ peak in the aerobic exercise training programme groups was MD 3.76 higher (6.89 lower to 14.41 higher)		59 (Hommerding 2015, Kriemler 2013)	⊕⊕⊕⊕ very low8,11,12	
[Unsupervised programme] Change in VO ₂	The mean change in VO ₂ in the	The mean change in VO ₂ peak in the		25 (Kriemler 2013)	⊕⊕⊕⊕ low6	

Comparison 1. Aerobic exercise training programme compared to no exercise programme for cystic fibrosis						
peak - ml/min per kg BW Follow-up: 6 months	control group was - 11.48	aerobic exercise training programme groups was 18.33 higher (8.95 to 27.71 higher)				
Time to next exacerbation	No evidence was found					
[Unsupervised programme] Change in BMI - kg/m2 Follow-up: 3 months	The mean change in BMI in the control group was - 0.3	The mean change in BMI in the aerobic exercise training programme groups was 0.3 higher (0.13 lower to 0.73 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ very low6,10	
[Unsupervised programme] Change in BMI - Kg/ m2 Follow-up: 6 months	The mean change in BMI in the control group was - 0.4	The mean change in BMI in the aerobic exercise training programme groups was 0.4 higher (0 to 0.8 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ very low6,10	
[Supervised programme] Change in BMI -	No evidence was found					
Quality of life	No evidence was found					
Preference for training programme	No evidence was found					
Adverse events	No evidence was found					
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; FEV ₁ : forced expiratory volume in 1 second; FVC: forced vital capacity; kg: kilogrammes MD: mean difference; min: minute; ml: millilitres; VO ₂ max/ peak: maximal oxygen consumption						

- 1 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to random sequence generation, blinding of participants and personnel and blinding of outcome assessment.
- 2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 because of unclear risk of bias in relation to allocation concealment, blinding of participants and personnel and blinding of outcome assessment in 1 study; high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of blinding of personnel, unclear risk of other bias (due to the deterioration of physical health in the control group) in the other study
- 4 The quality of the evidence was downgraded by 2 due to very serious heterogeneity (chi-squared p<0.1, I-squared inconsistency statistic of 90%) and no plausible explanation was found with sensitivity or subgroup analysis.
- 5 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs
- 6 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of bias in relation to blinding of participants and personnel, and unclear risk of other bias (due to the deterioration of physical health in the control group)

7 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to allocation concealment, blinding of participants and personnel, incomplete outcome data and other bias (exclusion criteria were not stated)

8 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs

9 The quality of the evidence was downgraded by 2 due to very serious heterogeneity (chi-squared $p < 0.1$, I-squared inconsistency statistic of 84%) and no plausible explanation was found with sensitivity or subgroup analysis.

10 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

11 The quality of the evidence was downgraded by 2 because of unclear risk of bias in relation to allocation concealment, blinding of participants and personnel, blinding of outcome assessment and other bias (the mean peak heart rate reached during the exercise test is indicative of submaximal effort, which is likely to underestimate the true VO_2 peak of the study participants) in 1 study; high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of blinding of personnel, unclear risk of other bias (due to the deterioration of physical health in the control group) in the other study

12 The quality of the evidence was downgraded by 2 due to very serious heterogeneity (chi-squared $p < 0.1$, I-squared inconsistency statistic of 75%) and no plausible explanation was found with sensitivity or subgroup analysis.

Table 171: Summary clinical evidence profile: Comparison 2.1. Strength resistance/ anaerobic training programme versus no exercise programme

Comparison 2.1. Strength resistance/ anaerobic training programme compared to no exercise programme for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise programme	Strength resistance/ anaerobic training programme				
[Supervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: at hospital discharge, mean 18.7 days	The mean change in FEV ₁ % predicted in the control group was 4.51	The mean change in FEV ₁ % predicted in the strength resistance/ anaerobic training groups was 5.58 higher (1.34 to 9.82 higher)		44 (Selvadurai 2002)	⊕⊕⊖ ⊖ low1,2	
[Unsupervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: mean 3 months	The mean change in FEV ₁ % predicted in the control group was -7.92	The mean change in FEV ₁ % predicted in the strength resistance/ anaerobic training groups was 11.11 higher (5.16 to 17.06 higher)		21 (Kriemler 2013)	⊕⊕⊖ ⊖ low3	
[Unsupervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100.	The mean change in FEV ₁ % predicted in the control group was	The mean change in FEV ₁ % predicted in the strength resistance/ anaerobic		21 (Kriemler 2013)	⊕⊕⊖ ⊖ low3	

Comparison 2.1. Strength resistance/ anaerobic training programme compared to no exercise programme for cystic fibrosis						
Follow-up: mean 6 months	-11	training groups was 19.51 higher (10.57 to 28.45 higher)				
[Supervised programme] Change in FVC % predicted - Scale from: 0 to 100. Follow-up: at hospital discharge, mean 18.7 days	The mean change in FVC % predicted in the control group was 2.28	The mean change in FVC % predicted in the strength resistance/ anaerobic training groups was 0.17 higher (2.31 lower to 2.65 higher)		44 (Selvadurai 2002)	⊕⊕⊕ ⊖ very low ^{1,4}	
[Unsupervised programme] Change in FVC % predicted - Scale from: 0 to 100. Follow-up: mean 3 months	The mean change in FVC % predicted in the control group was -5.57	The mean change in FVC % in the strength resistance/ anaerobic training groups was 7.37 higher (1.89 to 12.85 higher)		21 (Kriemler 2013)	⊕⊕⊕ ⊖ very low ^{3,5}	
[Unsupervised programme] Change in FVC % predicted Scale from: 0 to 100. Follow-up: mean 6 months	The mean change in FVC % predicted in the control group was -7.85	The mean change in FVC % predicted in the strength resistance/ anaerobic training groups was 14.05 higher (7.16 to 20.94 higher)		21 (Kriemler 2013)	⊕⊕⊕ ⊖ low ³	
[Supervised programme] Change in VO ₂ peak ml/min per kg body weight Follow-up: at hospital discharge, mean 18.7 days	The mean change in VO ₂ peak in the control group was -1.22	The mean change in VO ₂ peak in the strength resistance/ anaerobic training groups was 1.95 higher (1.61 lower to 5.51 higher)		44 (Selvadurai 2002)	⊕⊕⊕ ⊖ low ^{1,5}	
[pooled supervised and unsupervised] Change in VO ₂ peak at 3 months ml/min per kg body weight	The mean change in VO ₂ peak in the control group was -1.84 in 1 study. 2.45 in the other	The mean change in VO ₂ peak in the strength resistance/ anaerobic training groups was 6.36 higher		41 (Klijn 2004, Kriemler 2013)	⊕⊕⊕ ⊖ very low ^{5,6}	

Comparison 2.1. Strength resistance/ anaerobic training programme compared to no exercise programme for cystic fibrosis						
Follow-up: mean 3 months	study	(1.22 to 11.49 higher)				
[Unsupervised programme] Change in VO ₂ peak ml/min per kg body weight. Scale from: 0 to 100. Follow-up: mean 3 months	The mean change in VO ₂ peak in the control group was -1.84	The mean change in VO ₂ peak in the strength resistance/ anaerobic training groups was 9.34 higher (1.66 to 17.02 higher)		21 (Kriemler 2013)	⊕⊕⊕ ⊖ very low ^{3,5}	
[Supervised programme] Change in VO ₂ peak ml/min per kg body weight. Scale from: 0 to 100. Follow-up: mean 3 months	The mean change in VO ₂ peak in the control group was -2.45	The mean change in VO ₂ peak in the strength resistance/ anaerobic training groups was 3.95 higher (2.95 lower to 10.85 higher)		20 (Klijn 2004)	⊕⊕⊕ ⊖ low ^{5,7}	
[Unsupervised programme] Change in VO ₂ peak ml/min per kg body weight Follow-up: mean 6 months	The mean change in VO ₂ peak in the control group was -11.48	The mean change in VO ₂ peak in the strength resistance/ anaerobic training groups was 17.7 higher (5.98 to 29.42 higher)		18 (Kriemler 2013)	⊕⊕⊕ ⊖ very low ^{3,5}	
Time to next exacerbation	No evidence was found					
[Unsupervised programme] Change in BMI Follow-up: mean 3 months	The mean change in BMI in the control group was -0.3	The mean change in BMI in the strength resistance/ anaerobic training groups was 0.5 higher (0.07 to 0.93 higher)		25 (Kriemler 2013)	⊕⊕⊕ ⊖ very low ^{3,5}	
[Unsupervised programme] Change in BMI Follow-up: mean 6 months	The mean change in BMI in the control group was -0.4	The mean change in BMI in the strength resistance/ anaerobic training groups was 0.7 higher (0.27 to 1.13 higher)		25 (Kriemler 2013)	⊕⊕⊕ ⊖ low ³	

Comparison 2.1. Strength resistance/ anaerobic training programme compared to no exercise programme for cystic fibrosis					
[Supervised programme] Change in BMI	No evidence was found				
[Unsupervised programme] Change in quality of life	No evidence was found				
[Supervised programme] Change in quality of life CFQ - physical function domain. Scale from: 0 to 100. Follow-up: mean 3 months	The mean change in quality of life in the control group was 87.1	The mean change in health-related quality of life in the strength resistance/ anaerobic training groups was 1.3 higher (11.55 lower to 14.15 higher)		20 (Kijn 2004)	⊕⊕⊕ ⊖ very low3,8
Preference for training programme	No evidence was found				
Adverse events	No evidence was found				
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; FEV ₁ : forced expiratory volume in 1 second; FVC: forced vital capacity; kg: kilogrammes MD: mean difference; min: minute; ml: millilitres; VO ₂ max/ peak: maximal oxygen consumption					

- 1 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to random sequence generation, blinding of participants and personnel and blinding of outcome assessment.
- 2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of bias in relation to blinding of participants and personnel, and unclear risk of other bias (due to the deterioration of physical health in the control group)
- 4 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs
- 5 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 6 The quality of the evidence was downgraded by 2 because of: high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of bias in relation to blinding of participants and personnel, and unclear risk of other bias (due to the deterioration of physical health in the control group) in 1 study; unclear risk of bias in relation to random sequence generation, blinding of participants and personnel, blinding of outcome assessment, other bias (exclusion criteria were not reported) in the other study.
- 7 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to random sequence generation (described as randomised but no details given), blinding of participants and personnel, blinding of outcome assessment (the primary researcher was blinded but their role in the study is unclear), other bias (exclusion criteria were not reported)
- 8 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs

19 **Table 172: Summary clinical evidence profile: Comparison 2.2. Strength/ anaerobic**
20 **training programme versus aerobic training programme**

Comparison 2.2. Strength/ anaerobic training compared to aerobic training for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				

Comparison 2.2. Strength/ anaerobic training compared to aerobic training for cystic fibrosis						
	Aerobic training	Strength/ anaerobic training)	
[Supervised programme] Change in FEV ₁ % predicted at hospital discharge Scale from: 0 to 100. Follow-up: at hospital discharge, mean 18.7 days	The mean change in FEV ₁ % predicted in the aerobic training group was 6.54	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 3.55 higher (0.94 lower to 8.04 higher)		44 (Selvadurai 2002)	⊕⊕⊕⊕ low1,2	
[Unsupervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 3 months	The mean change in FEV ₁ % predicted in the aerobic training group was 4.89	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 1.7 lower (7.67 lower to 4.27 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ very low2,3	
[Unsupervised exercise] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 6 months	The mean change in FEV ₁ % predicted in the aerobic training group was 6.17	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 2.34 higher (6.33 lower to 11.01 higher)		26 (Kriemler 2013)	⊕⊕⊕⊕ very low3,4	
[Supervised exercise] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 6 months	The mean change in FEV ₁ % predicted in the aerobic training group was -2.57	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 1.66 lower (11.24 lower to 7.92 higher)		56 (Orenstein 2004)	⊕⊕⊕⊕ very low4,5	
[Pooled results for supervised and unsupervised programmes] Change in FEV ₁ % predicted Scale from: 0 to	The mean change in FEV ₁ % predicted in the aerobic training group was 6.17 in 1 study and -	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 0.54 higher (5.89 lower to 6.97 higher)		82 (Kriemler 2013, Orenstein 2004)	⊕⊕⊕⊕ very low4,6	

Comparison 2.2. Strength/ anaerobic training compared to aerobic training for cystic fibrosis						
100 Follow-up: 6 months	2.57 in the other study					
[Supervised exercise] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 12 months	The mean change in FEV ₁ % predicted in the aerobic training group was -1.19	The mean change in FEV ₁ % predicted in the strength/ anaerobic training groups was 0.3 higher (9.21 lower to 9.81 higher)		53 (Orenstein 2004)	⊕⊕⊕⊕ very low ^{4,5}	
[Supervised programme] Change in FVC % predicted Scale from: 0 to 100. Follow-up: at hospital discharge, mean 18.7 days	The mean change in FVC % predicted in the aerobic training group was 2.34	The mean change in FVC % predicted in the strength/ anaerobic training groups was 0.11 higher (2.49 lower to 2.71 higher)		44 (Selvadurai 2002)	⊕⊕⊕⊕ very low ^{1,7}	
[Unsupervised programme] Change in FVC % predicted Scale from: 0 to 100 Follow-up: 3 months.	The mean change in FVC % predicted in the aerobic training group was 3.67	The mean change in FVC % predicted in the strength/ anaerobic training groups was 1.87 lower (7.33 lower to 3.59 higher)		25 (Kriemler 2013)	⊕⊕⊕⊕ very low ³	
[Unsupervised programme] Change in FVC % predicted Scale from: 0 to 100. Follow-up: 6 months	The mean change in FVC % predicted in the aerobic training group was 4.66	The mean change in FVC % predicted in the strength/ anaerobic training groups was 1.54 higher (5.12 lower to 8.2 higher)		26 (Kriemler 2013)	⊕⊕⊕⊕ very low ^{3,7}	
[Supervised programme] Change in VO ₂ peak Follow-up: at hospital discharge, mean 18.7 days	The mean change in VO ₂ peak in the aerobic training group was 7.31	The mean change in VO ₂ peak in the strength/ anaerobic training groups was 6.58 lower (10.18 to 2.98 lower)		44 (Selvadurai 2002)	⊕⊕⊕⊕ low ^{1,8}	
[Unsupervised programme] Change in VO ₂	The mean change in VO ₂ peak in	The mean change in VO ₂ peak in the		26 (Kriemler 2013)	⊕⊕⊕⊕ very low ^{3,7}	

Comparison 2.2. Strength/ anaerobic training compared to aerobic training for cystic fibrosis						
peak Follow-up: 3 months	the aerobic training group was 7.26	strength/ anaerobic training groups was 0.24 higher (6.1 lower to 6.58 higher)				
[Unsupervised exercise] Change in VO ₂ max Follow-up: 6 months	The mean change in VO ₂ peak in the aerobic training group was 6.85	The mean change in VO ₂ max in the strength/ anaerobic training groups was 0.63 lower (10.94 lower to 9.68 higher)		26 (Kriemler 2013)	⊕⊕⊕⊕ very low ^{3,7}	
[Supervised exercise] Change in VO ₂ max Follow-up: 6 months	The mean change in VO ₂ peak in the aerobic training group was -1.91	The mean change in VO ₂ max in the strength/ anaerobic training groups was 0.25 lower (3.35 lower to 2.85 higher)		56 (Orenstein 2004)	⊕⊕⊕⊕ very low ^{5,8}	
[Pooled results for supervised and unsupervised programmes] Change in VO ₂ max Follow-up: 6 months	The mean change in VO ₂ max in the aerobic training group was 6.85 in 1 study and - 1.91 in the other study	The mean change in VO ₂ max in the strength/ anaerobic training groups was -0.28 lower (3.25 lower to 2.69 higher)		82 (Kriemler 2013, Orenstein 2004)	⊕⊕⊕⊕ low ⁶	
[Supervised exercise] Change in VO ₂ max Follow-up: 12 months	The mean change in VO ₂ peak in the aerobic training group was -0.91	The mean change in VO ₂ max in the strength/ anaerobic training groups was 0.82 lower (4.32 lower to 2.68 higher)		53 (Orenstein 2004)	⊕⊕⊕⊕ very low ^{5,8}	
[Unsupervised programme] Change in BMI Follow-up: 3 months	The mean change in BMI in the aerobic training group was 0	The mean change in BMI in the strength/ anaerobic training groups was 0.2 higher (0.23 lower to 0.63 higher)		30 (Kriemler 2013)	⊕⊕⊕⊕ very low ^{3,8}	
[Unsupervised programme]	The mean change in	The mean change in BMI		30 (Kriemler	⊕⊕⊕⊕ very	

Comparison 2.2. Strength/ anaerobic training compared to aerobic training for cystic fibrosis					
Change in BMI Follow-up: 6 months	BMI in the aerobic training group was 0	in the strength/ anaerobic training groups was 0.3 higher (0.1 lower to 0.7 higher)		2013)	low3,8
[Supervised programme] Change in BMI -	No evidence available				
Quality of life	No evidence available				
Preference for training programme	No evidence available				
Adverse events	No evidence available				
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).					
Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; FEV ₁ : forced expiratory volume in 1 second; FVC: forced vital capacity; kg: kilogrammes MD: mean difference; min: minute; ml: millilitres; VO ₂ max/ peak: maximal oxygen consumption					

- 1 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to random sequence generation, blinding of participants and personnel and blinding of outcome assessment.
- 2 The quality of the evidence was downgraded by 1 because the 95% 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to random sequence generation and allocation concealment, unclear risk of bias in relation to blinding of participants and personnel, and unclear risk of other bias (due to the deterioration of physical health in the control group)
- 4 The quality of the evidence was downgraded by 2 because the 95% 95% CI crossed 2 clinical MIDs
- 5 The quality of the evidence was downgraded by 2 due to high risk of bias in relation to blinding of participants and personnel and unclear risk of bias in relation to random sequence generation and allocation concealment.
- 6 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to random sequence generation and allocation concealment in 1 study, and unclear risk of bias in relation to the same domains in the other study; high risk of bias in relation to blinding of participants and personnel in 1 study and unclear risk of bias in relation to the same domains in the other study; and unclear risk of other bias in 1 study (due to the deterioration of physical health in the control group)
- 7 The quality of the evidence was downgraded by 2 because the 95% 95% CI crossed 2 default MIDs
- 8 The quality of the evidence was downgraded by 1 because the 95% 95% CI crossed 1 default MID

17 **Table 173: Summary clinical evidence profile: Comparison 3. High-intensity interval**
18 **training versus standard aerobic and anaerobic exercise programme**

Comparison 3. High intensity interval training programme compared to standard combined aerobic and anaerobic exercise programme for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Standard combined aerobic and anaerobic exercise programme	High intensity interval training programme				
[Unsupervised programme] Change in	No evidence was found					

Comparison 3. High intensity interval training programme compared to standard combined aerobic and anaerobic exercise programme for cystic fibrosis						
FEV ₁ %						
[Supervised programme] Change in FEV ₁ % predicted - Scale from: 0 to 100. Follow-up: 6 weeks	The mean change in FEV ₁ % predicted in the control group was 2.8	The mean change in FEV ₁ % predicted in the high intensity interval training groups was 3.9 lower (7.61 to 0.19 lower)		43 (Gruber 2014)	⊕⊕⊕ ⊖ very low _{1,2}	
[Unsupervised programme] Change in vital capacity (VC) % predicted	No evidence was found					
[Supervised programme] Change in vital capacity (VC) % predicted Scale from: 0 to 100. Follow-up: 6 weeks	The mean change in VC % predicted in the standard exercise group was 3.9	The mean change in vital capacity (VC) % predicted in the high intensity interval training groups was 5.1 lower (11.05 lower to 0.85 higher)		43 (Gruber 2014)	⊕⊕⊕ ⊖ very low _{1,3}	
[Unsupervised programme] Change in VO ₂ peak	No evidence was found					
[Supervised programme] Change in VO ₂ peak Follow-up: 6 weeks	The mean change in VO ₂ peak in the control standard exercise was 3.3	The mean change in VO ₂ peak in the high intensity interval training groups was 0.8 lower (4.59 lower to 2.99 higher)		43 (Gruber 2014)	⊕⊕⊕ ⊖ very low _{1,3}	
Time to next exacerbation	No evidence was found					
[Unsupervised programme] BMI	No evidence was found					
[Supervised programme] BMI Follow-up: 6 weeks	The mean change in BMI in the standard exercise group was 0.4	The mean BMI in the high intensity interval training groups was 0 higher (1.34 lower to 1.34 higher)		44 (Gruber 2014)	⊕⊕⊕ ⊖ very low _{1,4}	
Quality of life	No evidence was found					
Preference for training programme	No evidence was found					
Adverse events	No evidence was found					

Comparison 3. High intensity interval training programme compared to standard combined aerobic and anaerobic exercise programme for cystic fibrosis

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; VC: vital capacity; kg: kilogrammes MD: mean difference; min: minute; ml: millilitres; VO₂ max/ peak: maximal oxygen consumption

1 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to the selection of the participants for each group and the comparability of the groups

2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID

3 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

4 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs

Table 174: Summary clinical evidence profile: Comparison 4. Inspiratory muscle training (IMT) at 80% of maximal effort versus usual care

Comparison 4. Inspiratory muscle training (80% of maximal effort) programme compared to usual care for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual care	Inspiratory muscle training (IMT) (80% of maximal effort) programme				
Change in FEV ₁ % (litres) Scale from: 0 to 100. Follow-up: 2 to 6 months	The mean change in FEV ₁ (litres) in the control group was 2	The mean change in FEV ₁ % (litres) in the IMT groups was 0 higher (0.9 lower to 0.9 higher)		19 (Enright 2004)	⊕⊕⊕⊕ low1	
Change in FVC (litres) Scale from: 0 to 100. Follow-up: 2 to 6 months	The mean change in FEV ₁ (litres) in the control group was 3	The mean change in FVC (litres) in the IMT groups was 0.1 higher (0.9 lower to 1.1 higher)		19 (Enright 2004)	⊕⊕⊕⊕ very low1,2	
VO ₂ peak	No evidence was found					
Time to next exacerbation	No evidence was found					
Body composition	No evidence was found					
Quality of life	No evidence was found					
Preference for training	No evidence was found					

Comparison 4. Inspiratory muscle training (80% of maximal effort) programme compared to usual care for cystic fibrosis

programme	
Adverse events	No evidence was found
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; IMT: inspiratory muscle training; MD: mean difference</p>	

1 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to blinding (performance bias and detection bias), and unclear risk of bias in relation to random sequence generation, allocation concealment, incomplete outcome data, selective reporting, and other bias.

2 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs

Table 175: Summary clinical evidence profile: Comparison 5. Combined aerobic and anaerobic training programme versus no exercise programme

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise programme	Combined aerobic and anaerobic training programme				
[Unsupervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 3 months	The mean change in FEV ₁ % predicted in the control group was -0.5 in 1 study; -2 in another study; 2.7 in the third study.	The mean change in FEV ₁ % predicted in the combined aerobic and anaerobic training groups was 4.27 lower (9.63 lower to 1.09 higher)		89 (Beaudoin 2016, Rovedder 2014, Schindel 2015)	⊕⊕⊖ ⊖ low1,2	
[Unsupervised programme] Change in FEV ₁ % predicted Scale from: 0 to 100. Follow-up: 3-6 months	The mean change in FEV ₁ % predicted in the control group was -4.1	The mean change in FEV ₁ % predicted in the combined aerobic and anaerobic training groups was 2 higher (5.31 lower to 9.31 higher)		35 (Hebestreit 2010)	⊕⊖⊖ ⊖ very low3,4	
[Unsupervised programme] Change in FVC % predicted	The mean change in FVC % predicted in	The mean change in FVC % predicted in the combined		89 (Beaudoin 2016, Rovedder	⊕⊕⊖ ⊖ low1,5	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
Scale from: 0 to 100. Follow-up: 3 months	the control group was: -3.68 in 1 study; -3.5 in 1 study; 1.8 in 1 study	aerobic and anaerobic training groups was 1.47 lower (6.21 lower to 3.27 higher)		2014, Schindel 2015)		
[Unsupervised programme] Change in FVC % predicted Scale from: 0 to 100. Follow-up: 3-6 months	The mean change in FVC % predicted in the control group was not calculable	The mean change in FVC % predicted in the combined aerobic and anaerobic training groups was 0.5 higher (4.3 lower to 5.3 higher)		35 (Hebestreit 2010)	⊕⊕⊕ ⊖ very low3,6	
[Unsupervised programme] Change in VO ₂ peak Follow-up: 3 months	The mean change in VO ₂ peak in the control group was 2.37	The mean change in VO ₂ peak in the combined aerobic and anaerobic training groups was 2.13 lower (7.06 lower to 2.80 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low6,7	
[Unsupervised programme] Change in VO ₂ peak Follow-up: 3-6 months	The mean change in VO ₂ peak in the control group was not calculable	The mean change in VO ₂ peak in the combined aerobic and anaerobic training groups was 2.04 higher (0.08 to 4 higher)		38 (Hebestreit 2010)	⊕⊕⊕ ⊖ low3	
[Unsupervised programme] Change in weight (kg) Follow-up: 3 months	The mean change in weight (kg) in the control groups was 0.07	The mean change in weight (kg) in the combined aerobic and anaerobic training groups was 0.27 lower (12.95 lower to 12.41 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low6,7	
[Unsupervised programme] Change in BMI Follow-up: 3 months	The mean change in BMI in the control groups was -0.15	The mean change in BMI in the combined aerobic and anaerobic training groups		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low6,7	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
		was 0.06 higher (2.68 lower to 2.80 higher)				
[Unsupervised programme] Change in BMI Follow-up: 3 to 6 months	The mean change in BMI in the control groups was not calculable	The mean change in BMI in the combined aerobic and anaerobic training groups was 0.4 higher (0.17 lower to 0.97 higher)		35 (Hebestreit 2010)	⊕⊕⊕ ⊖ very low 3,5	
[Unsupervised programme] Change in BMI Follow-up: 12 months	The mean change in BMI in the control groups was not calculable	The mean change in BMI in the combined aerobic and anaerobic training groups was 0.54 higher (0.09 lower to 1.17 higher)		48 (Moorcroft 2004)	⊕⊕⊕ ⊖ very low 5,8	
[Unsupervised programme] Change in QOL: CFQ-R physical - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R physical in the control group was 2.4 (-1.0 to 13)	Not calculable.	P=0.742	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate 9,10	
[Unsupervised programme] Change in QOL: CFQ-R physical Follow-up: 3 months	The mean change in CFQ-R physical in the control group was 6.92	The mean change in CFQ-R physical in the combined aerobic and anaerobic training in the intervention groups was 0.60 higher (17.56 lower to 18.76 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low 4,7	
[Unsupervised programme] Change in QOL: CFQ-R body image Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R body image in the control group was 3.0 (-2 to 11)	Not calculable.	P=0.915	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate 9,10	
[Unsupervised programme] Change in	The mean change in CFQ-R body	The mean change in CFQ-R body image in		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
QOL: CFQ-R (body image) Follow-up: 3 months	image in the control group was 1.87	the combined aerobic and anaerobic training groups was 6.03 lower (18.89 lower to 6.83 higher)			low2,7	
[Unsupervised programme] Change in QOL: CFQ-R digestive - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R digestive in the control group was -0.5 (0 to 0)	Not calculable.	P=0.953	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	
[Unsupervised programme] Change in QOL: CFQ-R digestive Follow-up: 3 months	The mean change in CFQ-R digestive in the control group was -9.25	The mean change in CFQ-R in the combined aerobic and anaerobic training groups was 14.80 higher (0.43 to 29.17 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low2,7	
[Unsupervised programme] Change in QOL: CFQ-R respiratory - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R respiratory in the control group was -4.7 (-1 to 7)	Not calculable.	P=0.925	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	
[Unsupervised programme] Change in QOL: CFQ-R respiratory Follow-up: 3 months	The mean change in CFQ-R respiratory in the control group was 4.63	The mean change in CFQ-R respiratory in the combined aerobic and anaerobic training groups was 4.63 lower (16.88 lower to 7.62 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low2,7	
[Unsupervised programme] Change in QOL: CFQ-R emotional - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R emotional in the control group was -4.3 (-13 to 6)	Not calculable.	P=0.458	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
[Unsupervised programme] Change in QOL: CFQ-R emotional Supervised programme Scale from: 0 to 100. Follow-up: 3 months	The mean change in CFQ-R emotional in the control group was 1.11	The mean change in CFQ-R emotional in the combined aerobic and anaerobic training groups was 7.78 lower (18.65 lower to 3.09 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low ^{2,7}	
[Unsupervised programme] Change in QOL: CFQ-R social - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R social in the control group was -1.7 (5 to 11)	Not calculable.	P=0.953	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate ^{9,10}	
[Unsupervised programme] Change in QOL: CFQ-R social Scale from: 0 to 100. Follow-up: 3 months	The mean change in CFQ-R social in the control group was 2.79	The mean change in CFQ-R social in the combined aerobic and anaerobic training groups was 5.29 lower (18.10 lower to 7.52 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low ^{2,7}	
[Unsupervised programme] Change in QOL: (CFQ-R eating disturbances - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R eating disturbances in the control group was -2.0 (-11 to 0)	Not calculable.	P=0.913	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate ^{9,10}	
[Unsupervised programme] Change in QOL: CFQ-R eating disturbances Follow-up: 3 months	The mean change in CFQ-R food in the control group was 0	The mean change in CFQ-R eating disturbances in the combined aerobic and anaerobic training groups was MD -1.39 (4.91 lower to 2.13 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ low ⁷	
[Unsupervised programme] Change in	The median (IQR) CFQ-R treatment in	Not calculable.	P=0.850	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
QOL: CFQ-R treatment at 3 months - u	the control group was -2.0 (-11 to 0)				e9,10	
[Unsupervised programme] Change in QOL: CFQ-R treatment Scale from: 0 to 100. Follow-up: 3 months	The mean change in CFQ-R treatment in the control group was 0	The mean change in CFQ-R treatment in the combined aerobic and anaerobic training groups was 5.56 lower (26.03 lower to 14.91 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low4,7	
[Unsupervised programme] Change in QOL: CFQ-R vitality Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R vitality in the control group was 2.6 (-8 to 10)	Not calculable.	P=0.579	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	
[Unsupervised programme] Change in QOL: CFQ-R vitality Follow-up: 3 months	The mean change in CFQ-R vitality in the control group was 0	The mean change in CFQ-R vitality in the combined aerobic and anaerobic training in the intervention groups was 3.13 higher (13.45 lower to 19.71 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low4,7	
[Unsupervised programme] Change in QOL: CFQ-R health - Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R health in the control group was -3.0	Not calculable.	P=0.382	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	
[Unsupervised programme] Change in QOL: CFQ-R health Scale from: 0 to 100. Follow-up: 3 months	The mean change in CFQ-R health in the control group was 16.68	The mean change in CFQ-R health at 3 months in the combined aerobic and anaerobic training groups was 5.57 lower (21.75 lower to 10.61 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low4,7	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis						
[Unsupervised programme] Change in QOL: CFQ-R weight Scale from: 0 to 100. Follow-up: 3 months	The median (IQR) CFQ-R weight in the control group was 4.6 (0 to 33)	Not calculable.	P=0.410	41 (Rovedder 2014)	⊕⊕⊕ ⊖ moderate9,10	
[Unsupervised programme] Change in QOL: CFQ-R weight Follow-up: 3 months	The mean change in CFQ-R weight in the control group was 0	The mean change in CFQ-R weight in the combined aerobic and anaerobic training groups was 8.34 lower (36.73 lower to 20.05 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low4,7	
[Unsupervised programme] Change in QOL: CFQ-R role limitations Follow-up: 3 months	The mean change in CFQ-R role limitations in the control group was - 5.56	The mean change in CFQ-R role limitations in the combined aerobic and anaerobic training groups was 4.52 higher (13.37 lower to 22.41 higher)		14 (Beaudoin 2016)	⊕⊕⊕ ⊖ very low4,7	
[Supervised programme] Change in QOL: CFQ-R children's Scale from: 0 to 100. Follow-up: 2 months	In the control group the median pre-intervention was 696 (495 to 741); the median post-intervention was 719 (550 to 734)	Not calculable.	P=0.257	22 (Santana-Sosa 2012)	⊕⊕⊕ ⊖ low10,11	
[Supervised programme] Change in QOL: CFQ-R parents' Scale from: 0 to 100. Follow-up: 2 months	In the control group the median pre-intervention was 896 (688 to 1011); the median post-intervention was 889 (811 to 973)	Not calculable.	P=0.143	22 (Santana-Sosa 2012)	⊕⊕⊕ ⊖ low10,11	
Preference for training programme	No evidence available					
[Supervised programme] Adverse events	No data reported for the control	Not calculable. No adverse events occurred		22 (Santana-Sosa)	⊕⊕⊕ ⊖ low10,11	

Comparison 5. Combined aerobic and anaerobic training programme compared to no exercise programme for cystic fibrosis

Follow-up: 2 months	group	during exercise training		2012)	1	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: BMI: body mass index; CI: confidence interval; CF: cystic fibrosis; CFQ-R: cystic fibrosis questionnaire revised; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; kg: kilogrammes MD: mean difference; min: minute; ml: millilitres; VO₂ max/ peak: maximal oxygen consumption</p>						

- 1 The quality of the evidence was downgraded by 1 because of unclear risk of bias in relation to the allocation concealment and blinding of participants and personnel across the 3 studies; high risk of bias in relation to incomplete outcome data and unclear risk of bias in relation to blinding of outcome assessors and selective reporting in 1 study
- 2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 clinical MID
- 3 The quality of the evidence was downgraded by 2 because of high risk of bias for the random sequence generation and allocation concealment domains and unclear risk of bias for the blinding, outcome assessment and reporting domains
- 4 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 clinical MIDs
- 5 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 6 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs
- 7 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to incomplete outcome data, unclear risk of bias in relation to allocation concealment, selective reporting, blinding of participants and personnel and outcome assessors
- 8 The quality of the evidence was downgraded by 2 due to unclear risk of bias for the random sequence generation, allocation concealment, blinding and incomplete outcome data domains
- 9 The quality of the evidence was downgraded by 1 because of unclear risk of bias for the domains allocation concealment and blinding
- 10 Imprecision could not be calculated, as results are provided as medians
- 11 The quality of the evidence was downgraded by 2 because of high risk of bias for incomplete outcome data, and unclear risk of bias for random sequence generation, allocation concealment and blinding

Table 176: Summary clinical evidence profile: Comparison 6. Combined inspiratory muscle training (IMT), resistance and aerobic training

Comparison 6. Combined inspiratory muscle training resistance and aerobic training compared to no exercise programme for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No exercise programme	Combined inspiratory muscle training (IMT) resistance and aerobic training				
[Unsupervised programme] Change in FEV ₁ (litres)	No evidence was found.					
[Supervised programme] Change in FEV ₁ (litres) Scale from: 0 to 100. Follow-up: 2 months	The mean change in FEV ₁ (litres) in the control group was 0.02	The mean change in FEV ₁ (litres) in the combined inspiratory muscle training (IMT) resistance and aerobic		20 (Santana-Sosa 2014)	⊕⊕⊖⊖ low1	

Comparison 6. Combined inspiratory muscle training resistance and aerobic training compared to no exercise programme for cystic fibrosis						
		training group was 0.07 higher (0.54 lower to 0.68 higher)				
[Unsupervised programme] Change in FVC	No evidence was found.					
Change in FVC (litres) Supervised programme Scale from: 0 to 100. Follow-up: 2 months	The mean change FVC (litres) in the control group was -0.05	The mean change in forced vital capacity (litres) in the combined inspiratory muscle training (IMT) resistance and aerobic training group was 0.16 higher (0.68 lower to 1 higher)		20 (Santana-Sosa 2014)	⊕⊕⊕⊕ very low ^{1,2}	
Change in VO ₂ peak	No evidence was found.					
Time to next exacerbation	No evidence was found.					
[Unsupervised programme] Change in weight	No evidence was found.					
[Supervised programme] Change in weight (kg) Follow-up: 2 months	The mean change weight in the control group was 0.9	The mean change in weight at 2 months in the intervention group was 0.50 higher (10.51 lower to 11.51 higher)		20 (Santana-Sosa 2014)	⊕⊕⊕⊕ very low ^{1,2}	
[Unsupervised programme] Change in QOL (CFQ-R) Follow-up: 2 months	No evidence was found.					
[Supervised programme] Change in QOL (CFQ-R) Scale from: 0 to 100. Follow-up: 2 months	Not reported	The median QOL (CFQ-R) at 2 months in the intervention group was 688 (609 to 791); whereas the median pre-intervention: 629 (505 to 701) (p=0.071)		20 (Santana-Sosa 2014)	⊕⊕⊕⊕ low ^{1,3}	

Comparison 6. Combined inspiratory muscle training resistance and aerobic training compared to no exercise programme for cystic fibrosis

Preference for training programme	No evidence was found.					
[Unsupervised programme] Adverse events	No evidence was found.					
[Supervised programme] Adverse events Follow-up: 2 months	Not reported	No adverse events occurred during exercise training.	Not estimable	20 (Santana-Sosa 2014)	⊕⊕⊕⊖ low1,3	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval; CF: cystic fibrosis; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; kg: kilogrammes MD: mean difference; VO₂ max/ peak: maximal oxygen consumption</p>						

- 1 *1 The quality of the evidence was downgraded by 2 due to high risk of bias for outcome reporting, and unclear*
2 *risk of bias for randomization, allocation concealment and blinding*
3 *2 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs*
4 *3 Imprecision could not be calculated, as data was reported narratively only*

5 **Table 177: Summary clinical evidence profile: Comparison 7. Physical activity:**
6 **higher amount or longer duration versus lower amount or shorter duration**

Comparison 7. Physical activity: higher amount or longer duration compared to physical activity for lower amount or shorter duration for cystic fibrosis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Physical activity for lower amount or shorter duration	Physical activity for higher amount or longer duration				
Lung function: FEV ₁ % predicted	No evidence available					
Lung function: FVC% predicted	No evidence available					
VO ₂ peak	No evidence available					
Body composition	No evidence available					
Quality of life	No evidence available					
Preference for training programme	No evidence available					
Adverse events	No evidence available					
[≥30 mins daily versus < 30 mins] Need for hospitalization Follow-up: 12	679 per 1000	482 per 1000 (312 to 746)	RR 0.71 (0.46 to 1.1)	61 (Cox 2016)	⊕⊖⊖⊖ very low1,2	

Comparison 7. Physical activity: higher amount or longer duration compared to physical activity for lower amount or shorter duration for cystic fibrosis

months						
[≥30 mins for ≥10 mins bouts daily versus lower amount or shorter duration]	650 per 1000	383 per 1000 (208 to 689)	RR 0.59 (0.32 to 1.06)	61 (Cox 2016)	⊕⊖⊖⊖ very low ^{1,2}	
Need for hospitalization						
Follow-up: 12 months						

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; RR: risk ratio

1 The quality of the evidence was downgraded by 2 due to high risk of bias in relation to the selection of the study population and the comparability of the 2 groups

2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

4 **10.8.5 Economic evidence**

5 No economic evaluations of exercise programmes were identified in the literature search
6 conducted for this guideline. Full details of the search and economic article selection flow
7 chart can be found in Appendix E and F, respectively.

8 This review question was not prioritised for de novo economic modelling. However, the
9 exercise programmes under consideration vary in the resources and costs required, for
10 example habitual exercise programmes could be incorporated into daily activities, whereas
11 programmes monitored and facilitated by exercise psychologists, therapy technical
12 instructors or physiotherapists can entail high staff costs, especially if they are performed
13 regularly.

14 According to NHS Reference Costs 2015/16 the cost per attendance with a sport and
15 exercise therapist is £94 (WF02A, Non-Admitted Face to Face Attendance, Follow-up, Non-
16 consultant led, 325, Sport and Exercise Medicine) and the cost per attendance with a
17 physiotherapist is £45 (WF02A, Non-Admitted Face to Face Attendance, Follow-up, Non-
18 consultant led, 650, Physiotherapy). Consequently, supervised programmes that incur high
19 staff costs will need to provide additional benefits, in relation to unsupervised programmes to
20 be considered cost-effective.

21 **10.8.6 Evidence statements**

22 **10.8.6.1 Aerobic exercise programmes**

23 **10.8.6.1.1 Comparison 1. Aerobic exercise training programme versus no exercise programme**

24 **Lung function: FEV₁**

25 Low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
26 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
27 difference in lung function (measured as change in FEV₁ % predicted) between the group of
28 participants attending a *supervised* hospital training programme – consisting of aerobic
29 exercise – and those who received usual care at hospital discharge (mean follow-up ≈ 19
30 days).

1 Very low quality evidence from 2 RCTs showed conflicting results in relation to the difference
2 in lung function (measured as change in FEV₁ % predicted) between the group of
3 participants receiving an *unsupervised* training programme – consisting of aerobic exercise –
4 and the participants in the control group at 3 months follow-up. One RCT with 34 children
5 and young people with cystic fibrosis showed no clinically significant difference between the
6 groups. One RCT with 24 participants with cystic fibrosis aged >12 years showed a clinically
7 significant improvement in the group of participants receiving an *unsupervised* training
8 programme –consisting of aerobic exercise- compared to the participants in the control
9 group.

10 Likewise, low quality evidence from 1 RCT with 25 people with cystic fibrosis >12 years
11 showed a clinically significant improvement in lung function (measured as change in FEV₁ %
12 predicted) in the group of participants receiving an *unsupervised* training programme –
13 consisting of aerobic exercise – compared to the participants in the control group at 6
14 months follow-up.

15 However, moderate quality evidence from 1 RCT with 65 people with cystic fibrosis showed
16 no clinically significant difference in lung function (measured as change in FEV₁ % predicted)
17 between the group of participants receiving an *unsupervised* home exercise programme –
18 consisting of aerobic training – and the participants in the control group at 3 years follow-up.

19 **Lung function: FVC**

20 Low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
21 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
22 difference in forced vital capacity (measured as change in FVC % predicted) between the
23 group of participants attending a *supervised* hospital training programme – consisting of
24 aerobic exercise – and those who received usual care at hospital discharge (mean follow-up
25 ≈ 19 days).

26 Very low quality evidence from 2 RCTs showed conflicting results in relation to the difference
27 in forced vital capacity (measured as change in FVC % predicted) between the group of
28 participants receiving an *unsupervised* training programme – consisting of aerobic exercise –
29 and the participants in the control group at 3 months follow-up. One RCT with 34 children
30 and young people with cystic fibrosis showed no clinically significant difference between the
31 groups. One RCT with 24 participants with cystic fibrosis aged >12 years showed a clinically
32 significant improvement in the group of participants receiving an *unsupervised* training
33 programme –consisting of aerobic exercise- compared to the participants in the control
34 group.

35 Likewise, low quality evidence from 1 RCT with 25 people with cystic fibrosis >12 years
36 showed a clinically significant improvement in forced vital capacity (measured as change in
37 FVC % predicted) in the group of participants receiving an *unsupervised* training programme
38 – consisting of aerobic exercise – compared to the participants in the control group at 6
39 months follow-up.

40 Low quality evidence from 1 RCT with 65 people with cystic fibrosis showed a clinically
41 significant improvement in forced vital capacity (measured as change in FVC % predicted) in
42 the group of participants receiving an *unsupervised* home exercise programme – consisting
43 of aerobic training – compared to the participants in the control group at 3 years follow-up.

44 **VO₂ max**

45 Moderate quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
46 admitted to hospital due to a pulmonary exacerbation showed a clinically significant
47 improvement in the maximum volume of oxygen (measured as change in VO₂ max) in the
48 group of participants attending a *supervised* hospital training programme – consisting of

1 aerobic exercise – compared those who received usual care at hospital discharge (mean
2 follow-up ≈ 19 days).

3 Very low quality evidence from 2 RCTs showed conflicting results in relation to the difference
4 in the maximum volume of oxygen (measured as change in VO₂ max) between the group of
5 participants receiving an *unsupervised* training programme – consisting of aerobic exercise –
6 and the participants in the control group at 3 months follow-up. One RCT with 34 children
7 and young people with cystic fibrosis showed no clinically significant difference between the
8 groups. One RCT with 25 participants with cystic fibrosis aged >12 years showed a clinically
9 significant improvement in the group of participants receiving an *unsupervised* training
10 programme –consisting of aerobic exercise- compared to the participants in the control
11 group.

12 However, low quality evidence from 1 RCT with 25 people with cystic fibrosis >12 years
13 showed a clinically significant improvement in the maximum volume of oxygen (measured as
14 change in VO₂ max) in the group of participants receiving an *unsupervised* training
15 programme – consisting of aerobic exercise – compared to the participants in the control
16 group at 6 months follow-up.

17 **Time to next exacerbation**

18 No evidence was found for this critical outcome.

19 **Body composition**

20 Very low quality evidence from 1 RCT with 25 people with cystic fibrosis >12 years showed
21 no clinically significant difference in weight (measured as change in BMI) between the
22 participants receiving an *unsupervised* training programme – consisting of aerobic exercise
23 – and the participants in the control group at 3 and 6 months follow-up.

24 No evidence was found for *supervised* training programmes.

25 **Quality of life**

26 No evidence was found for this critical outcome.

27 **Preference for training programme**

28 No evidence was found for this important outcome.

29 **Adverse events**

30 No evidence was found for this important outcome.

31 **10.8.6.2 Strength resistance training/ anaerobic training**

32 **10.8.6.2.1 Comparison 2.1. Strength resistance / anaerobic training programme versus no 33 exercise programme**

34 **Lung function: FEV₁**

35 Low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
36 admitted to hospital due to a pulmonary exacerbation showed a clinically significant
37 improvement in lung function (measured as change in FEV₁ % predicted) in the group of
38 participants attending a *supervised* hospital training programme – consisting of anaerobic
39 exercise – compare to those who received usual care at hospital discharge (mean follow-up
40 ≈ 19 days).

1 Low quality evidence from 1 RCT with 21 people with cystic fibrosis >12 years showed a
2 clinically significant improvement in lung function (measured as change in FEV₁ % predicted)
3 in the group of participants attending an *unsupervised* anaerobic programme – consisting of
4 strength training sessions – compared to the control group at 3 and 6 months follow-up.

5 **Lung function: FVC**

6 Very low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
7 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
8 difference in forced vital capacity (measured as change in FVC % predicted) between the
9 group of participants attending a *supervised* hospital training programme – consisting of
10 anaerobic exercise – and those who received usual care at hospital discharge (mean follow-
11 up ≈ 19 days).

12 However, low to very low quality evidence from 1 RCT with 21 people with cystic fibrosis >12
13 years showed a clinically significant improvement in forced vital capacity (measured as
14 change in FVC % predicted) in the group of participants attending an *unsupervised*
15 anaerobic programme – consisting of strength training sessions – compared to the control
16 group at 3 and 6 months follow-up.

17 **VO₂ max**

18 Very low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
19 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
20 difference in maximum volume of oxygen (measured as change in VO₂ max) between the
21 group of participants attending a *supervised* hospital training programme – consisting of
22 anaerobic exercise – and those who received usual care at hospital discharge (mean follow-
23 up ≈ 19 days).

24 Low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis with
25 stable disease showed no clinically significant difference in the maximum volume of oxygen
26 (measured as change in VO₂ max) between the participants attending a supervised
27 anaerobic training programme and the control group at 3 months follow-up. However, very
28 low quality evidence from 1 RCT with 21 people with cystic fibrosis >12 years showed a
29 clinically significant improvement between the group of participants attending an
30 *unsupervised* anaerobic programme – consisting of strength training sessions – and the
31 control group at the same follow-up. Very low quality evidence from the pooled results of
32 both *supervised and unsupervised* training programmes showed a clinically significant
33 beneficial effect in VO₂ max in the group of participants attending the training programme at
34 3 months follow-up-.

35 Very low quality evidence from 1 RCT with 18 people with cystic fibrosis >12 years showed a
36 clinically significant improvement in the maximum volume of oxygen (measured as change in
37 VO₂ max) in the group of participants attending an *unsupervised* anaerobic programme –
38 consisting of strength training sessions – compared to the control group at 6 months follow-
39 up.

40 **Time to next exacerbation**

41 No evidence was found for this critical outcome.

42 **Body composition: BMI**

43 Low quality evidence from 1 RCT with 25 people with cystic fibrosis >12 years showed a
44 clinically significant improvement in weight (measured as change in BMI) in the group of
45 participants attending an *unsupervised* anaerobic programme – consisting of strength
46 training sessions – compared to the control group at 3 and 6 months follow-up.

1 **Quality of life**

2 Very low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
3 with stable disease showed no clinically significant difference in health related quality of life
4 (measured with the CFQ-R tool, physical domain) between the participants attending a
5 supervised anaerobic training programme and the control group at 3 months follow-up.

6 No evidence was found for *supervised* training programmes.

7 **Preference for training programme**

8 No evidence was found for this important outcome.

9 **Adverse events**

10 No evidence was found for this important outcome.

11 **110.8.6.2.2 Comparison 2.2. Strength/ anaerobic training programme versus aerobic training**
12 **programme**

13 **Lung function: FEV₁**

14 Low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
15 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
16 difference in lung function (measured as change in FEV₁ % predicted) between the group of
17 participants attending a *supervised* hospital anaerobic exercise training programme and
18 those attending a *supervised* hospital aerobic exercise training programme at hospital
19 discharge (mean follow-up ≈ 19 days).

20 Very low quality evidence from 1 RCT with 26 people with cystic fibrosis >12 years showed
21 no clinically significant difference in lung function (measured as change in FEV₁ % predicted)
22 between the group of participants attending an *unsupervised* anaerobic programme –
23 consisting of strength training sessions –and those attending an *unsupervised* aerobic
24 programme at 3 and 6 months follow-up. Likewise, very low quality evidence from 1 RCT
25 with 56 children and young people with cystic fibrosis showed no clinically significant
26 difference in lung function (measured as change in FEV₁ % predicted) between the group of
27 participants attending a *supervised* anaerobic training programme - consisting of upper-body
28 strength regime – and the participants attending an aerobic training programme at 6 months
29 follow-up. Very low quality evidence from pooled results of both *supervised and*
30 *unsupervised* training programmes showed no clinically significant difference in lung function
31 (measured as change in FEV₁ % predicted) between the group of participants attending an
32 anaerobic training programme and those attending an aerobic training programme at 6
33 months follow-up.

34 Very low quality evidence from 1 RCT with 56 children and young people with cystic fibrosis
35 showed no clinically significant difference in lung function (measured as change in FEV₁ %
36 predicted) between the group of participants attending a *supervised* anaerobic training
37 programme - consisting of upper-body strength regime – and the participants attending an
38 aerobic training programme at 12 months follow-up.

39 **Lung function: FVC**

40 Very low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
41 admitted to hospital due to a pulmonary exacerbation showed no clinically significant
42 difference in lung function (measured as change in FVC % predicted) between the group of
43 participants attending a *supervised* hospital anaerobic exercise training programme and
44 those attending a *supervised* hospital aerobic exercise training programme at hospital
45 discharge (mean follow-up ≈ 19 days).

1 Very low quality evidence from 1 RCT with 26 people with cystic fibrosis >12 years showed
2 no clinically significant difference in lung function (measured as change in FVC % predicted)
3 between the group of participants attending an *unsupervised* anaerobic programme –
4 consisting of strength training sessions –and those attending an *unsupervised* aerobic
5 programme at 3 and 6 months follow-up.

6 **VO₂ max**

7 Low quality evidence from 1 RCT with 44 children and young people with cystic fibrosis
8 admitted to hospital due to a pulmonary exacerbation showed a clinically significant lower
9 improvement in the maximum volume of oxygen (measured as change in VO₂ max) in the
10 group of participants attending a *supervised* hospital anaerobic exercise training programme
11 compared to those attending a *supervised* hospital aerobic exercise training programme at
12 hospital discharge (mean follow-up ≈ 19 days).

13 Very low quality evidence from 1 RCT with 26 people with cystic fibrosis >12 years showed
14 no clinically significant difference in the maximum volume of oxygen (measured as change in
15 VO₂ max) between the group of participants attending an *unsupervised* anaerobic
16 programme – consisting of strength training sessions –and those attending an *unsupervised*
17 aerobic programme at 3 months follow-up.

18 Very low quality evidence from 1 RCT with 26 people with cystic fibrosis >12 years showed
19 no clinically significant difference in the maximum volume of oxygen (measured as change in
20 VO₂ max) between the group of participants attending an *unsupervised* anaerobic
21 programme – consisting of strength training sessions –and those attending an *unsupervised*
22 aerobic programme at 6 months follow-up. Likewise, very low quality evidence from 1 RCT
23 with 56 children and young people with cystic fibrosis showed no clinically significant
24 difference in the maximum volume of oxygen (measured as change in VO₂ max) between the
25 group of participants attending a *supervised* anaerobic training programme - consisting of
26 upper-body strength regime – and the participants attending a *supervised* aerobic training
27 programme at 6 months follow-up. Low quality evidence from the pooled results of both
28 *supervised* and *unsupervised* training programmes showed no clinically significant difference
29 in in the maximum volume of oxygen (measured as change in VO₂ max) between the group
30 of participants attending an anaerobic training programme and those attending an aerobic
31 training programme at 6 months follow-up.

32 Very low quality evidence from 1 RCT with 56 children and young people with cystic fibrosis
33 showed no clinically significant difference in the maximum volume of oxygen (measured as
34 change in VO₂ max) between the group of participants attending a *supervised* anaerobic
35 training programme - consisting of upper-body strength regime – and the participants
36 attending a *supervised* aerobic training programme at 12 months follow-up.

37 **Time to next exacerbation**

38 No evidence was found for this critical outcome.

39 **Body composition**

40 Very low quality evidence from 1 RCT with 30 people with cystic fibrosis >12 years showed
41 no clinically significant difference in BMI between the group of participants attending an
42 *unsupervised* anaerobic programme – consisting of strength training sessions –and those
43 attending an *unsupervised* aerobic programme at 3 and 6 months follow-up.

44 No evidence was found for *supervised* exercise programmes.

45 **Quality of life**

46 No evidence was found for this critical outcome.

1 **Preference for training programme**

2 No evidence was found for this important outcome.

3 **Adverse events**

4 No evidence was found for this important outcome.

5 **10.8.6.3 High intensity interval training**

6 **10.8.6.3.1 Comparison 3. High intensity interval training versus standard aerobic and anaerobic exercise programme**

7
8 **Lung function: FEV₁**

9 Very low quality evidence from 1 cohort study with 43 adult inpatients with cystic fibrosis with
10 severe disease (FEV₁ <40% predicted) showed no clinically significant difference in lung
11 function (measured as change in FEV₁ % predicted) between the group of participants
12 performing supervised high intensity interval training and the participants performing a
13 supervised standardised exercise programme at 6 weeks follow-up.

14 No evidence was found for *unsupervised* training programmes.

15 **Lung function: vital capacity (VC)**

16 Very low quality evidence from 1 cohort study with 43 adult inpatients with cystic fibrosis with
17 severe disease (FEV₁ <40% predicted) showed no clinically significant difference in vital
18 capacity (measured as change in VC % predicted) between the group of participants
19 performing supervised high intensity interval training and the participants performing a
20 supervised standardised exercise programme at 6 weeks follow-up.

21 No evidence was found for *unsupervised* training programmes.

22 **VO₂ max**

23 Very low quality evidence from 1 cohort study with 43 adult inpatients with cystic fibrosis with
24 severe disease (FEV₁ <40% predicted) showed no clinically significant difference in the
25 maximum volume of oxygen (VO₂ max) between the group of participants performing
26 supervised high intensity interval training and the participants performing a supervised
27 standardised exercise programme at 6 weeks follow-up.

28 No evidence was found for *unsupervised* training programmes.

29 **Time to next exacerbation**

30 No evidence was found for this critical outcome.

31 **Body composition: BMI**

32 Very low quality evidence from 1 cohort study with 43 adult inpatients with cystic fibrosis with
33 severe disease (FEV₁ <40% predicted) showed no clinically significant difference in BMI
34 between the group of participants performing supervised high intensity interval training and
35 the participants performing a supervised standardised exercise programme at 6 weeks
36 follow-up.

37 No evidence was found for *unsupervised* training programmes.

- 1 **Quality of life**
- 2 No evidence was found for this critical outcome.
- 3 **Preference for training programme**
- 4 No evidence was found for this important outcome.
- 5 **Adverse events**
- 6 No evidence was found for this important outcome.
- 7 **10.8.6.4 Inspiratory muscle training**
- 8 **10.8.6.4.1 Comparison 4. Inspiratory muscle training (IMT) at 80% of maximal effort versus usual**
- 9 **care**
- 10 **Lung function: FEV₁**
- 11 Low quality evidence from 1 RCT with 19 adults with cystic fibrosis showed no clinically
- 12 significant difference in lung function (measured as change in FEV₁ % predicted) between
- 13 the group of participants receiving IMT at home at 80% of maximal inspiratory effort and the
- 14 participants receiving usual care at 2 to 6 months follow-up.
- 15 **Lung function: FVC**
- 16 Low quality evidence from 1 RCT with 19 adults with cystic fibrosis showed no clinically
- 17 significant difference in lung function (measured as change in FVC % predicted) between the
- 18 group of participants receiving IMT at home at 80% of maximal inspiratory effort and the
- 19 participants receiving usual care at 2 to 6 months follow-up.
- 20 **VO₂ max**
- 21 No evidence was found for this important outcome.
- 22 **Time to next exacerbation**
- 23 No evidence was found for this critical outcome.
- 24 **Body composition**
- 25 No evidence was found for this important outcome.
- 26 **Quality of life**
- 27 No evidence was found for this critical outcome.
- 28 **Preference for training programme**
- 29 No evidence was found for this important outcome.
- 30 **Adverse events**
- 31 No evidence was found for this important outcome.

1 10.8.6.5 Combined programmes

20.8.6.5.1 Comparison 5. Combined aerobic and anaerobic training versus no exercise programme

4 Lung function: FEV₁

5 Low quality evidence from 3 RCTs with 89 people with cystic fibrosis aged ≥7 years showed
6 no clinically significant difference in lung function (measured as change in FEV₁ % predicted)
7 between the participants attending an *unsupervised* training programme - consisting of a
8 combination of aerobic and anaerobic exercises - and the participants in the control group at
9 3 months follow-up.

10 Likewise, very low quality evidence from another RCT with 35 people with cystic fibrosis >12
11 years showed no clinically significant difference in change in FEV₁ % predicted between the
12 participants attending an *unsupervised* training programme - consisting of a combination of
13 endurance-type and strengthening exercises - and the control group at 3 to 6 months follow-
14 up.

15 No evidence was found for *supervised* training programmes.

16 Lung function: FVC

17 Likewise, low quality evidence from 3 RCTs with 89 people with cystic fibrosis >7 years old
18 showed no clinically significant difference in forced vital capacity (measured as change in
19 FVC % predicted) between the participants attending an *unsupervised* training programme -
20 consisting of a combination of aerobic and anaerobic or resistance training exercises - and
21 the participants in the control group at 3 months follow-up.

22 Likewise, very low quality evidence from another RCT with 35 people with cystic fibrosis >12
23 years showed no clinically significant difference in change in FVC % predicted between the
24 participants attending an *unsupervised* training programme - consisting in a combination of
25 endurance-type and strengthening exercises - and the control group at 3 to 6 months follow-
26 up.

27 No evidence was found for *supervised* training programmes.

28 VO₂ max

29 Very low quality evidence from 1 RCT with 14 adults with cystic fibrosis showed no clinically
30 significant difference in maximum volume of oxygen (measured as change in VO₂ max)
31 between the participants attending an *unsupervised* training programme - consisting of a
32 combination of aerobic and resistance training exercise- and the participants in the control
33 group at 3 months follow-up.

34 Low quality evidence from 1 RCT with 38 people with cystic fibrosis >12 years showed no
35 clinically significant difference in the maximum volume of oxygen (measured as change in
36 VO₂ max) between the participants attending an *unsupervised* training programme -
37 consisting in a combination of endurance-type and strengthening exercises - and the
38 participants in the control group at 3 to 6 months follow-up.

39 No evidence was found for *supervised* training programmes.

40 Time to next exacerbation

41 No evidence was found for this critical outcome.

1 **Body composition: weight and BMI**

2 Very low quality evidence from 1 RCT with 14 adults with cystic fibrosis showed no clinically
3 significant difference in weight (measured as change in kg) between the participants
4 attending an *unsupervised* training programme - consisting of a combination of aerobic and
5 resistance training exercise- and the participants in the control group at 3 months follow-up.

6 Very low quality evidence from 1 RCT with 14 adults with cystic fibrosis showed no clinically
7 significant difference in weight (measured as change in BMI) between the participants
8 attending an *unsupervised* training programme - consisting of a combination of aerobic and
9 resistance training exercise- and the participants in the control group at 3 months follow-up.

10 Very low quality evidence from 1 RCT with 35 people with cystic fibrosis >12 years showed
11 no clinically significant difference in weight (measured at change in BMI) between the
12 participants attending an *unsupervised* training programme - consisting of a combination of
13 endurance-type and strengthening exercises - and the participants in the control group at 3
14 to 6 months follow-up.

15 Likewise, very low quality evidence from another RCT with 48 adults with cystic fibrosis
16 showed no clinically significant difference between the participants attending an
17 *unsupervised* training programme – consisting of general aerobic exercises and weight
18 training – and the participants in the control group at 12 months follow-up.

19 No evidence was found for *supervised* training programmes.

20 **Quality of life**

21 Low to very low quality evidence from 1 RCT with 14 adults with cystic fibrosis showed no
22 clinically significant difference in the following quality of life domains (measured as change in
23 the scores obtained with CFQ-R questionnaire) between the participants attending an
24 *unsupervised* training programme - consisting of a combination of aerobic and resistance
25 training exercise- and the participants in the control group at 3 months follow-up: physical
26 functioning, vitality, emotional state, eating disturbances, treatment burden, health
27 perception, social limitations, body image, role limitations, weight problems, respiratory
28 symptoms. The same evidence showed a clinically significant improvement in the quality of
29 life domain digestion symptoms among the participants attending the training programme
30 compared to the control group at 3 months follow-up.

31 Low quality evidence from 1 RCT with 22 children and young people with cystic fibrosis
32 showed no significant difference in quality of life (measured with CFQ-R children's and
33 parents' scales) between the participants attending a *supervised* intra-hospital exercise
34 programme – consisting of endurance and strengthening exercises - and the control group at
35 2 months follow-up. The clinical significance of these outcomes could not be calculated.

36 Likewise, moderate quality evidence from 1 RCT with 41 participants with cystic fibrosis >16
37 years showed no significant difference in quality of life (measured with CFQ-R questionnaire,
38 all domains) between the participants attending an *unsupervised* home-based exercise
39 programme – consisting of aerobic and muscle strengthening exercises – and the control
40 group at 3 months follow-up. The clinical significance of these outcomes could not be
41 calculated.

42 **Preference for training programme**

43 No evidence was found for this important outcome.

1 **Adverse events**

2 Low quality evidence from 1 RCT with 22 children with cystic fibrosis showed that none of the
3 participants attending a *supervised* training programme - consisting of endurance and
4 strengthening exercises – experienced an adverse event at 2 months follow-up.

5 No evidence was found for *unsupervised* training programmes.

6 **6.8.6.5.2 Comparison 6. Combined inspiratory muscle training, resistance and aerobic training
7 versus no exercise programme**

8 **Lung function: FEV₁**

9 Low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
10 showed no clinically significant difference in lung function (measured as change in FEV₁
11 litres) between the participants attending a *supervised* training programme - consisting of a
12 combination of inspiratory muscle training, resistance and aerobic training - and the
13 participants in the control group at 2 months follow-up.

14 No evidence was found for *unsupervised* training programmes.

15 **Lung function: FVC**

16 Very low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
17 showed no clinically significant difference in forced vital capacity (measured as change in
18 FVC litres) between the participants attending a *supervised* training programme - consisting
19 of a combination of inspiratory muscle training, resistance and aerobic training - and the
20 participants in the control group at 2 months follow-up.

21 No evidence was found for *unsupervised* training programmes.

22 **VO₂ max**

23 No evidence was found for this important outcome.

24 **Time to next exacerbation**

25 No evidence was found for this critical outcome.

26 **Body composition: weight**

27 Very low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
28 showed no clinically significant difference in weight change (measured in kg) between the
29 participants attending a *supervised* training programme - consisting of a combination of
30 inspiratory muscle training, resistance and aerobic training - and the participants in the
31 control group at 2 months follow-up.

32 No evidence was found for *unsupervised* training programmes.

33 **Quality of life**

34 Low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
35 showed no significant difference in quality of life (measured with CFQ-R questionnaire)
36 between the participants attending a *supervised* training programme – consisting of a
37 combination of inspiratory muscle training, resistance and aerobic training - at 2 months
38 follow-up. The clinical significance of this outcome could not be calculated.

39 No evidence was found for *unsupervised* training programmes.

- 1 **Preference for training programme**
- 2 No evidence was found for this important outcome.
- 3 **Adverse events**
- 4 Low quality evidence from 1 RCT with 20 children and young people with cystic fibrosis
5 showed that none of the participants attending a *supervised* training programme - consisting
6 of a combination of inspiratory muscle training, resistance and aerobic training – experienced
7 an adverse event at 2 months follow-up.
- 8 No evidence was found for *unsupervised* training programmes.
- 9 **10.8.6.6 Habitual physical activity**
- 10 **10.8.6.6.1 Comparison 7. Physical activity: higher amount or longer duration versus lower
11 amount or shorter duration**
- 12 **Lung function: FEV₁**
- 13 No evidence was found for this critical outcome.
- 14 **Lung function: FVC**
- 15 No evidence was found for this important outcome.
- 16 **VO₂ max**
- 17 No evidence was found for this important outcome.
- 18 **Need of hospitalization (proxy outcome for time to next exacerbation)**
- 19 Very low quality evidence from a cohort study with 61 adults with cystic fibrosis showed no
20 clinically significant difference in need for hospitalization between the group of people doing
21 at least 30 minutes daily of moderate-vigorous physical activity compared to those doing less
22 physical activity at 12 months follow-up. The same evidence showed that there was no
23 clinically significant difference in need for hospitalization between the group that did at least
24 30 minutes of moderate-vigorous physical activity per day accumulated in bouts of more than
25 10 minutes and those who did less than 30 minutes daily or did more than 30 minutes but in
26 shorter bouts at 12 months follow-up.
- 27 **Body composition**
- 28 No evidence was found for this important outcome.
- 29 **Quality of life**
- 30 No evidence was found for this critical outcome.
- 31 **Preference for training programme**
- 32 No evidence was found for this important outcome.
- 33 **Adverse events**
- 34 No evidence was found for this important outcome.
- 35 **10.8.6.7 Economic evidence statements**
- 36 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

1 10.8.7 Evidence to recommendations

2 10.8.7.1 Relative value placed on the outcomes considered

3 The aim of this review was to determine the effectiveness of different exercise programmes
4 in improving health outcomes for people with cystic fibrosis.

5 The Committee chose lung function (FEV₁), quality of life and time to next exacerbation as
6 critical outcomes for decision making; whereas forced vital capacity (FVC), maximum volume
7 of oxygen (VO₂), body composition, preference for training programme and adverse events
8 were rated as important.

9 10.8.7.2 Consideration of clinical benefits and harms

10 The Committee reviewed the evidence on the effectiveness of aerobic exercise programmes
11 compared to no exercise programme. The evidence was of moderate to very low quality. The
12 evidence showed that these programmes were mostly effective in relation to FVC %
13 predicted and VO₂ max; results were mixed in relation to FEV₁ % predicted; the evidence
14 showed no benefits in relation to BMI. No evidence was found in relation to the other
15 outcomes, including adverse events. The Committee concluded that the evidence showed
16 some benefit of aerobic exercise programmes and there was no evidence of harm.

17 The Committee also reviewed the evidence on programmes of strength resistance or
18 anaerobic training compared to no exercise programme. The evidence was of low to very low
19 quality. The evidence showed that these programmes were mostly effective in relation to
20 FEV₁% predicted, VO₂ max and body composition; results were mixed in relation to FVC%
21 predicted; the evidence showed no benefits in relation to quality of life. No evidence was
22 found in relation to the other outcomes of interest, including adverse events. The Committee
23 concluded that the evidence showed some benefit of strength resistance or anaerobic
24 exercise programmes and there was no evidence of harm.

25 The Committee also reviewed the evidence on programmes of combined aerobic and
26 anaerobic training compared to no exercise programme. The evidence was of moderate to
27 very low quality and showed no benefits of training in relation to FEV₁% predicted, FVC%
28 predicted, VO₂ max, BMI, quality of life. This kind of training did not lead to an increase in
29 adverse events.

30 Overall, the Committee noted that the evidence showed some benefits for lung function and
31 no harm of either aerobic or strength resistance or anaerobic training. In addition to these
32 benefits for lung function, the Committee agreed that people with cystic fibrosis would also
33 gain the same kind of fitness benefits from exercise that would be gained by the general
34 population. Therefore, the Committee decided to recommend to advise people with cystic
35 fibrosis of the benefits of regular exercise in relation to lung function in addition to the
36 expected fitness benefits.

37 The Committee reviewed the evidence on the effectiveness of a strength resistance or
38 anaerobic training programme compared to an aerobic training programme. The evidence
39 was of very low quality and showed no differences in relation to FEV₁ % predicted and VO₂
40 max. No evidence was found in relation to the other outcomes. Therefore, the Committee
41 decided not to recommend 1 type of exercise over the other. Rather it decided to recommend
42 to offer all people with cystic fibrosis individualised exercise programmes, taking into account
43 the capability and preferences of the individual. The Committee noted that taking the
44 individual's preferences into account is key to increase the likelihood of adherence to a
45 training programme.

46 The Committee reviewed the evidence on high intensity interval training compared to a
47 standard exercise programme. The evidence was of very low quality and was based on a
48 population of inpatients with FEV₁ <40% predicted. The evidence showed no benefits in

1 relation to FEV₁% predicted, VC% predicted, VO₂ max, or BMI. No evidence was found in
2 relation to the other outcomes, including adverse events.

3 The Committee reviewed the evidence on inspiratory muscle training (IMT) at 80% of
4 maximal effort compared to usual care. The evidence was of low quality and showed no
5 benefits in relation to FEV₁% predicted and FVC% predicted. No evidence was found in
6 relation to the other outcomes, including adverse events.

7 The Committee reviewed the evidence on a programme of combined inspiratory muscle
8 training, resistance and aerobic training compared to no exercise programme. The evidence
9 was of low to very low quality. The evidence showed no benefits in relation to FEV₁, FVC,
10 weight, or quality of life. This kind of training did not lead to an increase in adverse events.

11 Given that the evidence on the 3 aforementioned training programmes did not show benefits
12 or harm and was of low to very low quality, the Committee chose not to make a
13 recommendation specific to these kinds of training.

14 The Committee agreed that they could not draw a clear conclusion from the evidence on
15 whether supervised programmes were more effective than unsupervised programmes,
16 therefore it decided not to make a recommendation on this. It recommended however to
17 regularly monitor exercise programmes at clinic visits so that they can be adapted as
18 necessary; it noted that regular exercise testing is an important part of monitoring.

19 The Committee noted that there was no evidence on what intensity or frequency of exercise
20 would be the most effective, therefore decided not to make a recommendation specific to
21 this.

22 10.8.7.3 Consideration of economic benefits and harms

23 People with cystic fibrosis are encouraged to participate in activity and exercise that is
24 available to the general population, leading to no additional resource or cost use. However,
25 the Committee noted that there may be individuals who would benefit from more specialised
26 advice.

27 According to the Committee, exercise programmes for individuals with cystic fibrosis should
28 be developed along with support of physiotherapy teams that specialise in cystic fibrosis who
29 are able to provide regular monitoring and support. This enables exercise to complement
30 existing airway clearance techniques and treatment regimens, to maximise and support
31 adherence. In turn, this will improve treatment effects which the Committee believed to
32 subsequently outweigh the cost to develop and maintain those programmes.

33 From the clinical review, there was no strong evidence supervised programmes were more
34 effective than unsupervised programmes, or no programme. In light of this, the Committee
35 agreed that not all people with cystic fibrosis require supervised programmes. However,
36 there may be times when supervision is important such as to help teach technique. In those
37 cases, such as strength and resistance training, supervision is key to ensure the exercise is
38 performed correctly to minimise injuries and maximise benefits. The Committee stated that
39 such initial costs would be negligible compared to the potential downstream costs from
40 potential injuries and inactivity.

41 Given that not every person would benefit from supervised programmes, the Committee
42 agreed that physiotherapists must consider the opportunity cost of their time to supervise
43 individuals. For those reasons, the Committee agreed not to recommend supervised
44 programmes in their recommendation given that the level of supervision would be
45 individualised.

46 The Committee stated that patient preference is paramount to the success and sustainability
47 of a programme; hence, despite higher costs, a programme could be considered cost-
48 effective if that programme provides them with greater benefits than a cheaper programme.

1 However, the Committee noted that freely available activities would be trialled first.
2 Following this, the Committee did not want to specify the type or duration of exercise as this
3 should be tailored to the individual's preferences and capabilities.

4 The Committee agreed that offering all people with cystic fibrosis individualised exercise
5 programmes would not lead to a change in clinical practice as physiotherapists and dietitians
6 regularly review participation in exercise at each review. Therefore, recommendations to
7 offer exercise programmes were prioritised to enforce the importance of exercise. Moreover,
8 the Committee advised the cost to create an individualised programme would be negligible
9 compared to the benefits sustained exercise can provide.

10 Following this, the Committee highlighted the importance of maintaining an exercise
11 programme, even during inpatient care, to prevent any additional deteriorations in their
12 health that could lead to additional costs, such as longer hospital stays. As a result, the
13 Committee prioritised a recommendation to offer inpatients the opportunity to exercise. The
14 Committee agreed that inpatients usually have poorer health than outpatients and in their
15 experience are more at risk of adverse events. Therefore supervised exercise would be more
16 appropriate for some inpatients, to prevent the downstream costs from unsupervised
17 exercise that is potentially unsafe or ineffective.

18 The Committee agreed that an assessment upon hospital admission by a physiotherapist,
19 and any subsequent supervision would not deviate from current clinical practice. However,
20 the Committee was concerned that not all hospitals have sufficient exercise facilities for
21 people with cystic fibrosis, or space to store exercise equipment. Their solution was not to
22 build a "second" gym, but to provide a schedule that promotes cross-infection control
23 measures, for people with cystic fibrosis to access the facilities. Achieving those schedules
24 may incur additional cleaning and equipment, and reduce the time facilities are available for
25 patient use. For these reasons, hospitals should consider if their strategies to prevent cross-
26 infection using existing exercise facilities, outweighs the cost to provide additional space and
27 equipment devoted to people with cystic fibrosis.

28 Overall, the Committee advised that their recommendations were within the remit of
29 specialist cystic fibrosis physiotherapy teams, but noted that staffing levels and exercise
30 facilities during episodes of inpatient care need to be prioritised for exercise programmes to
31 be continued.

32 **10.8.7.4 Quality of evidence**

33 The quality of the evidence presented in this review ranged from very low to moderate as
34 assessed by GRADE.

35 For the domain risk of bias, the studies were assigned the same risk of bias as in the
36 Cochrane reviews, and were not individually reviewed. The main biases that led to
37 downgrading the quality of the evidence were randomization, allocation concealment,
38 attrition, and reporting bias. It is also important to note that it was not possible to blind
39 participants to the exercise intervention in randomised controlled trials thus increasing the
40 risk of performance bias.

41 In studies on unsupervised programmes it was not possible to independently assess if the
42 participants actually performed the exercise programmes as prescribed.

43 Another reason that led to downgrading the quality of the evidence was the imprecision, as
44 confidence intervals crossed 1 or 2 MIDs. The Committee noted that many trials were
45 underpowered to detect a clinically important difference.

46 No serious issues were found regarding inconsistency (heterogeneity), as most outcomes
47 were reported by a single study. Where heterogeneity was identified, and sensitivity or sub-

1 group analysis did not explain the source of inconsistency, the results were explained to the
2 Committee.

3 No serious issues were found regarding indirectness of the population or the interventions.

4 **10.8.7.5 Other considerations**

5 The Committee noted the NHS service specifications for cystic fibrosis do not cover exercise
6 in detail, although they mention that inpatients should have access to facilities for exercise.

7 No equality issues were identified by the Committee for this review question.

8 The Committee felt a research recommendation was not needed, as there is enough
9 available evidence to show that regular exercise is beneficial for people with cystic fibrosis.

10 **10.8.7.6 Key conclusions**

11 The Committee concluded that regular exercise is especially beneficial for people with cystic
12 fibrosis due to the benefits to lung function in addition to other fitness benefits. Exercise
13 programmes should be individualised to take into account individual circumstances and
14 preferences, given that there is no evidence that a specific programme may be better than
15 other programmes. There is no evidence to indicate whether a supervised exercise
16 programme may be better than an unsupervised programme, however the programme
17 should be regularly monitored and adapted if necessary. Inpatient stays should be used as
18 opportunities to promote regular exercise and appropriate supervision should be offered if
19 necessary depending on individual circumstances. Moreover, access to appropriate exercise
20 facilities should be guaranteed in the inpatient setting while taking into account local infection
21 control guidelines.

22 **10.8.8 Recommendations**

23 **124. Advise people with cystic fibrosis and their family members or carers (as**
24 **appropriate) that regular exercise improves both lung function and overall fitness.**

25 **125. Offer people with cystic fibrosis an individualised exercise programme, taking**
26 **into account their capability and preferences.**

27 **126. Regularly review exercise programmes to monitor the person's progress and**
28 **ensure that the programme continues to be appropriate for their needs.**

29 **127. Provide people with cystic fibrosis who are having inpatient care with:**

- 30 • an assessment of their exercise capacity
- 31 • the facilities and support to continue their exercise programme (as
- 32 appropriate), taking into account the need to prevent cross-infection (see
- 33 Prevention of cross infection) and local infection control guidelines.

34 **10.9 Psychological assessment**

35 **Review question: What strategies are effective at identifying people with cystic**
36 **fibrosis for the presence of a psychological and behavioural problem?**

37 **10.9.1 Introduction**

38 The emotional impact of a diagnosis of cystic fibrosis is significant and without appropriate
39 psychological guidance people with cystic fibrosis and their families and carers can find the

1 impact extends to their interpersonal relationships, health and quality of life. It is known that
2 physical health outcomes can be improved if psychological distress is prevented or reduced.

3 Clinical psychologists with expert knowledge of cystic fibrosis, can offer strategies to identify
4 psychological and behavioural problems, and interventions to prevent mental health
5 symptoms from developing into intractable mental health diagnoses.

6 These clinical psychologists, as members of the core multi-disciplinary team, will be aware of
7 any likely impact of cystic fibrosis treatments on emotional functioning and the key triggers
8 for potential distress.

9 Likely psychological and behavioural issues are: adherence, procedural-anxiety and phobias,
10 engagement with health care team, adjustment to diagnosis or deteriorated function, self-
11 esteem problems, anger management, relationship difficulties, sleep problems, body
12 image/eating issues, medical trauma, substance misuse, generalised anxiety disorders, low
13 mood and depression.

14 **10.9.2 Description of clinical evidence**

15 The aim of this review was to determine which assessment strategies are effective at
16 identifying psychological, behavioural and adherence problems in children, young people and
17 adults with cystic fibrosis.

18 The Committee identified the following tools as relevant for this review (See Table 178 for full
19 a description of the tools).

- 20 • Generalised Anxiety Disorder 7-item scale (GAD-7)
- 21 • Patient Health Questionnaire 2-item scale (PHQ-2)
- 22 • Patient Health Questionnaire 9-item primary care scale (PHQ-9)
- 23 • Hospital Anxiety and Depression Scale (HADS)
- 24 • Paediatric Index of Emotional Distress (PI-ED)
- 25 • Centre for Epidemiologic Studies Depression Scale (CES-D)
- 26 • Eating Disorders Examination (EDE)
- 27 • Child Eating Disorders Examination (CEDE, ChEDE)
- 28 • Eating Attitudes Test (EAT)
- 29 • Child Eating Attitude Test (ChEAT)

30 For the diagnostic accuracy data, the following reference standards were considered.

31 For psychological problems (including anxiety, depression, mood disorders, emotional
32 distress, adjustment disorders and eating disorders), the reference standard diagnoses
33 according to the Diagnostic and Statistical Manual of Mental Disorders (DSM) or International
34 Classification of Diseases (ICD). Studies that did not clearly state the comparator to be DSM
35 or ICD diagnosis of depression, or that did not provide sufficient diagnostic accuracy data,
36 were excluded.

37 For adherence, electronic monitoring was considered the gold standard. Alternatively,
38 pharmacy collection records were also considered a reference standard.

39 For other behavioural problems listed in the protocol (such as school phobia), the reference
40 standard was considered as reported by the study.

41 We also included validation studies that looked at the validity and reliability of the tools.

42 We looked for systematic reviews of diagnostic studies, and prospective and retrospective
43 cohort studies.

1 For this review, quality appraisal of the evidence has been conducted by study.

2 For full details see review protocol in Appendix D.

3 **Table 178: Description of tools assessed**

Tool name	Key features
Centre for Epidemiologic Studies Depression Scale (CES-D)	<ul style="list-style-type: none"> • The CES-D is a self-report measure and screens for depression and depressive disorders. It measures symptoms defined by the American Psychiatric Association's Diagnostic and Statistical Manual (DSM-V) for a major depressive episode. • Designed for use with the general adult population. • The questionnaire is comprised of 20 items and measures symptoms of depression in nine different groups as defined by the DSM-V: sadness (dysphoria), loss of interest (anhedonia), appetite, sleep, thinking/concentration, guilt (worthlessness), tired (fatigue), movement (agitation) and suicidal ideation. • The response values for each item ask the individual to indicate the frequency of symptoms in the past week.
Child eating attitudes test (ChEAT)	<ul style="list-style-type: none"> • The ChEAT is a modified version of the Eating Attitudes Test (EAT) by Garner and Garfinkle (1979) and provides an assessment of attitudes towards eating and dietary behaviours. • The ChEAT is a self-report measure for children aged between 8 to 14 years. • The questionnaire comprises 26 items that are rated on a 6-point scale with responses: 'always', 'very often', 'often', 'sometimes', 'rarely' and 'never'. • The items of the ChEAT are mostly scored and interpreted according to total score. However, factor analysis supported three factors – subscales derived from these factors are: dieting, restricting and purging and food preoccupation.
Child eating disorders examination (CEDE, ChEDE)	<ul style="list-style-type: none"> • The ChEDE was adapted from the Eating Disorder Examination (EDE) for recommended use with children and young people aged 8 to 14 years to assess psychopathology associated with the diagnosis of an eating disorder. • The measure is a semi-structured interview, with 2 main administrative modifications compared with the EDE. Ideas about weight and shape are assessed via a sort task rather than through questioning, and some of the questions assessing actual behaviour in adults, were reformulated to assess intent in children and young people. • The interview provides an overall global score and a score for 4 subscales: restraint, eating concern, shape concern and weight concern.
Eating attitudes test (EAT)	<ul style="list-style-type: none"> • The EAT is a self-report measure designed for adolescents and adults which screens for symptoms and concerns characteristic of eating disorders. • The questionnaire was originally developed in 1979 and called EAT-40, containing 40 questions. The EAT-26 is used more commonly nowadays, which is a refined version of the questionnaire developed following factor analysis. • There are three parts to the 26-item version: part A asks for the age, weight and other physical attributes of the respondent, part B screens for the respondent's attitude towards their weight, height and shape and part C asks about behavioural tendencies of the respondent over the past 6 months. • Total scores are interpreted in relation to BMI norms for the individual's age. Items are also divided to produce scores for 2 areas: dieting, bulimia and food preoccupation and oral control.
Eating disorders examination (EDE)	<ul style="list-style-type: none"> • The EDE is a semi-structured interview for use with adults to assess psychopathology associated with the diagnosis of an eating disorder, including range and severity of eating disorder features. • It is described as an assessment and diagnostic tool which provides operationally defined eating disorder diagnoses. • Questions address the frequency in which an individual engages in behaviours

Tool name	Key features
	<p>indicative of an eating disorder over a 28-day period.</p> <ul style="list-style-type: none"> • The interview provides an overall global score and a score for 4 subscales: restraint, eating concern, shape concern and weight concern. • The Eating Disorders Examination Questionnaire (EDE-Q) has been developed from the EDE, which provides a 41-item self-report questionnaire that retains the 4 subscales and global score.
Generalised Anxiety Disorder 7-item scale (GAD-7)	<ul style="list-style-type: none"> • The GAD-7 is a self-report questionnaire which provides a screening measure for generalised anxiety disorder (GAD). • The questionnaire has 7 items which measure the severity of symptoms associated with GAD, according to DSM-IV criteria. The individual is asked to consider how often, in the past 2 weeks, they were bothered by each symptom. • Items are scored on a 4 point scale with responses: 'not at all', 'several days', 'more than half the days' and 'nearly every day'.
Hospital Anxiety and Depression scale (HADS)	<ul style="list-style-type: none"> • The HADS is a self-rating scale for use with adults that screens for anxiety and depression in both hospital and community settings. • The measure assesses symptom severity of anxiety and depression in people with illness and the general population. • The questionnaire contains 14 items, 7 of which assess anxiety, and the remaining 7, depression.
Patient Health Questionnaire 2-item scale (PHQ-2)	<ul style="list-style-type: none"> • The PHQ-2 is a self-report screening tool for depression, which enquires about the frequency of depressed mood and anhedonia over the past 2 weeks. • For use with adolescents and adults. • The questionnaire contains only 2 items, which are the first two questions from the PHQ-9. • It is recommended that individuals who screen positive should be evaluated further with the PHQ-9 to determine whether they meet criteria for depressive disorder.
Patient Health Questionnaire 9-item scale (PHQ-9)	<ul style="list-style-type: none"> • The PHQ-9 is a self-report tool for screening, diagnosing, monitoring and measuring the severity of depression. • For use with both adolescents and adults. • It contains 9 items, which incorporate DSM-IV depression diagnostic criteria. • Individuals are required to rate how frequently over the past 2 weeks they are bothered by the 9 statements presented. • Items are scored on a 4 point scale with responses: 'not at all', 'several days', 'more than half the days' and 'nearly every day'.
Paediatric Index of Emotional Distress (PI-ED)	<ul style="list-style-type: none"> • The PI-ED is based on the HADS and is a self-rating scale which screens for emotional distress in children and young people aged 8 to 16 years. • The measure is designed for use with both children that have physical health problems and children in a normal population. • It contains 14 questions which ask about symptoms of anxiety and depression.

- 1 Four studies were identified for inclusion in this review.
- 2 One study (Shearer 2014) aimed to evaluate the CEDE for the assessment of eating
- 3 disorders in children with cystic fibrosis. The study was conducted in the UK and included 55
- 4 children and young people.
- 5 Three studies (Daniels 2011, Siracusa 2015, White 2014) aimed to determine what
- 6 assessments are effective in measuring adherence to treatment in children, young people
- 7 and adults with cystic fibrosis. These studies were conducted in the UK and the USA.
- 8 Sample sizes ranged from 12 to 250.
- 9 None of the studies looked at tools assessing anxiety, depression, mood and emotional
- 10 distress or adjustment disorders.

1 A summary of the included studies is presented in Table 159 and Table 180. See also study
2 selection flow chart in Appendix F, study evidence tables in Appendix G, and list of excluded
3 studies in Appendix H.

4 **10.9.3 Summary of included studies and results**

5 A summary of the included studies and results for this review is presented in Table 159 and
6 Table 180.

7

1 **Table 179: Summary of included studies and results for psychological problems**

Study	Psychological disorder	Tool assessed	Participants	Results	Comments
Shearer 2004 (UK)	Eating disorder	CEDE	N=55 children and young people with CF not undergoing psychological therapy Age: 11 to 17 years	Reliability • Inter-rater reliability = 0.69 to 1 Validity Not reported	Overall quality: moderate

2 *CF: cystic fibrosis; CEDE: Child eating disorders examination; UK: United Kingdom*

3 **Table 180: Summary of included studies and results for adherence to treatment**

Study	Treatment	Method assessed	Participants	Results	Comments
Daniels 2011 (UK)	Nebulizers	Index: • Self-report • Clinician-report Gold standard: • Electronic monitoring (I-neb nebulizer system)	N=78 adults with CF on nebulizer therapy. Median age (IQR): 26 (21 to 31) years	Adherence according to self-report: • Median (IQR) = 80% (57.5% to 95%) of treatment prescribed Adherence according to electronic monitoring: • Median (IQR) = 36% (5% to 84.8%) of treatment prescribed Clinician agreement: • ICC = 0.95 (95% CI 0.44 to 0.66) • Agreement between clinician report and electronic monitoring: • ICC dietitian = 0.36 (95% CI 0.11 to 0.55) • ICC liaison/ home nurse = 0.36 (95% CI 0.15 to 0.54) • ICC physician = 0.42 (95% CI 0.21 to 0.59) • ICC ward nurse = 0.34 (95% CI 0.11 to 0.54) • ICC pharmacist = 0.28 (95% CI 0.07 to 0.47) • ICC physiotherapist = 0.54 (95% CI 0.36	• Overall quality: moderate • Extreme inaccuracy was observed for individual participants by clinicians and self-report adherence.

Study	Treatment	Method assessed	Participants	Results	Comments
Siracusa 2015 (USA)	Ivacaftor	Index: <ul style="list-style-type: none"> • Self-report • Pharmacy refill history Gold standard: <ul style="list-style-type: none"> • Electronic monitoring • Mean monitoring duration: 118 days (SD=35) 	N=12 children, young people and adults with CF previously prescribed Ivacaftor Mean age (range): 20.8 years (6 to 48 years)	to 0.68) Adherence rates according to (mean, SD, range): <ul style="list-style-type: none"> • self-report: 100% (14% to 100%) • pharmacy refill history: 84% (31) (13% to 124%) • electronic monitoring: 61% (28) (4% to 99%) Electronic monitoring versus self-report <ul style="list-style-type: none"> • $r_s=0.40$; $p=0.22$ • $ICC=0.14$; $p=0.23$ Electronic monitoring versus pharmacy refill history <ul style="list-style-type: none"> • $r_s=0.26$; $p=0.42$ • $ICC=-0.26$; $p=0.14$ 	Overall quality: low Individuals demonstrated wide variability in regards to the different measures of adherence.
White 2014 (UK)	<ul style="list-style-type: none"> • Aerosol to open air • Aerosol to thin mucus • Inhaler • PERT • Oral nutritional supplements • Oral antibiotics • Nebulised antibiotics 	Index: <ul style="list-style-type: none"> • Self-report (CFQ-R) Index: <ul style="list-style-type: none"> • Pharmacy script data collection 	N=250 young people and adults with CF Mean (SD) age: 29.7 (9.2)	Correlation between pharmacy script collection and self-report: <ul style="list-style-type: none"> • Aerosol to open air: $r=0.34$; $p<0.005$ • Aerosol to thin mucus: $r=0.51$; $p<0.001$ • Inhaler: $r=0.51$; $p<0.001$ • PERT: $r=0.45$; $p<0.001$ • Oral nutritional supplements: $r=0.51$; $p<0.001$ • Oral antibiotics: $r=0.46$; $p<0.001$ • Nebulised antibiotics: $r=0.55$; $p<0.001$ • Total: $r=0.61$; $p<0.001$ 	Overall quality: cannot be assessed Conference abstract

1 CI: confidence interval; CF: cystic fibrosis; CFQ-R: cystic fibrosis questionnaire revised; ICC: intraclass correlation coefficient; IQR: inter-quartile range; PERT: pancreatic exocrine replacement therapy; SD: standard deviation

3

1 **10.9.4 Clinical evidence profile**

2 See summary of results in Table 159 and Table 180 in Summary of included studies and
3 results.

4 **10.9.5 Economic evidence**

5 No economic evaluations of psychological and behavioural assessments were identified in
6 the literature search conducted for this guideline and this review question was not prioritised
7 for de novo economic modelling.

8 Full details of the search and economic article selection flow chart can be found in Appendix
9 E and F, respectively.

10 **10.9.6 Evidence statements**

11 **10.9.6.1 Psychological disorders**

12 **10.9.6.1.1 Anxiety**

13 **Generalised Anxiety Disorder 7-item scale (GAD-7)**

14 No evidence was found for this tool.

15 **Hospital Anxiety and Depression scale (HADS)**

16 No evidence was found for this tool.

17 **10.9.6.1.2 Depression, mood disorders and emotional distress**

18 **Centre for Epidemiologic Studies Depression Scale (CES-D)**

19 No evidence was found for this tool.

20 **Hospital Anxiety and Depression scale (HADS)**

21 No evidence was found for this tool.

22 **Patient Health Questionnaire 2-item scale (PHQ-2)**

23 No evidence was found for this tool.

24 **Patient Health Questionnaire 9-item scale (PHQ-9)**

25 No evidence was found for this tool.

26 **Paediatric Index of Emotional Distress (PI-ED)**

27 No evidence was found for this tool.

28 **10.9.6.1.3 Adjustment disorders**

29 No evidence was found for this disorder.

10.9.6.1.4 **Eating disorders and feeding issues**

2 **Eating attitudes test (EAT)**

3 No evidence was found for this tool.

4 **Child eating attitudes test (ChEAT)**

5 No evidence was found for this tool.

6 **Eating disorders examination (EDE)**

7 No evidence was found for this tool.

8 **Child eating disorders examination (CEDE)**

9 One study reported on the usefulness of the CEDE scale in a population of 55 children and
10 young people with cystic fibrosis not receiving psychological treatment. Inter-rater reliability
11 ranged from 0.69 to 1. No measures of validity were reported. This overall quality of this
12 study was moderate.

13 10.9.6.2 **Non adherence to treatment**

14 One study reported on the usefulness of self-report and clinician-report as measures of
15 adherence in a population of 78 adults with cystic fibrosis on nebulizer therapy:

- 16 • there was an overestimation of adherence by the participants. No measures of reliability
17 were reported;
- 18 • there was an overestimation of adherence by the clinicians. The intra-class correlation
19 agreement ranged between 0.28 and 0.54.

20 No measures of validity were reported. It is important to note that extreme inaccuracy was
21 observed for individual participants by clinicians and self-report adherence. This overall
22 quality of this study was moderate.

23 One study reported on the usefulness of self-report and pharmacy refill history as measures
24 of adherence in a population of 12 children, young people and adults with cystic fibrosis
25 receiving Ivacaftor:

- 26 • there was an overestimation of adherence by the participants. There was no statistically
27 significant correlation between overall electronic-monitoring adherence and self-report
28 (intra-class correlation coefficient was 0.14);
- 29 • there was an overestimation of adherence by the pharmacy records. There was also no
30 statistically significant correlation between overall electronic-monitoring adherence and
31 pharmacy refill history (intra-class correlation coefficient was 0.26).
- 32 • No measures of validity were reported. It is important to note that individuals
33 demonstrated wide variability in regards to the different measures of adherence. The
34 overall quality of this study was low, mainly because it was underpowered.

35 One study reported on the usefulness of self-report (using the CFQ-R tool) as a measure of
36 adherence to treatment in a population of 250 young people and adults with cystic fibrosis:

- 37 • there was a statistically significant correlation between pharmacy script collection and self-
38 report adherence to aerosol to open air ($r=0.34$);
- 39 • there was a statistically significant correlation between pharmacy script collection and self-
40 report adherence to aerosol to thin mucus ($r=0.51$);
- 41 • there was a statistically significant correlation between pharmacy script collection and self-
42 report adherence to inhalers ($r=0.51$);

- 1 • there was a statistically significant correlation between pharmacy script collection and self-
2 report adherence to PERT ($r=0.45$);
3 • there was a statistically significant correlation between pharmacy script collection and self-
4 report adherence to oral nutritional supplements ($r=0.51$);
5 • there was a statistically significant correlation between pharmacy script collection and self-
6 report adherence to oral antibiotics ($r=0.51$).
7 • there was a statistically significant correlation between pharmacy script collection and self-
8 report adherence to nebulised antibiotics ($r=0.55$).
9 The overall quality of this study could not be assessed, as the information was extracted from
10 a conference abstract. Full publication not available.

11 10.9.6.3 Economic evidence statements

12 No evidence on cost-effectiveness in people with cystic fibrosis was available for this review.

13 10.9.7 Evidence to recommendations

14 10.9.7.1 Relative value placed on the outcomes considered

15 The aim of this review was to determine which assessment strategies are effective at
16 identifying psychological, behavioural and adherence problems in children, young people and
17 adults with cystic fibrosis.

18 Sensitivity and specificity of the tools were prioritised as critical outcomes for decision
19 making. Whereas positive likelihood ratio, negative likelihood ratio, AUROC, and reliability or
20 validity were rated as important outcomes.

21 10.9.7.2 Consideration of clinical benefits and harms

22 The Committee agreed the limited available evidence was not helpful in guiding them to
23 make recommendations.

24 The Committee discussed the issue of adherence at length. They noted adherence is an
25 overarching issue in this guideline. It should not actually fall under the category of
26 psychological or behavioural disorder or problem. To reflect this, results for adherence were
27 presented separately in this review.

28 The Committee stressed that adherence problems are common in people with chronic
29 conditions and those with a number of concurrent treatments, and are not particularly specific
30 to cystic fibrosis. They agreed that the overarching principles from the NICE guidance on
31 Medicines adherence: involving patients in decisions about prescribed medicines and
32 supporting adherence [CG76] is applicable to people with cystic fibrosis.

33 The results from this review showed that there is poor correlation between what participants
34 or professionals report, and what is actually taken. Likewise, there was poor correlation
35 between what it is collected and what it is actually taken. According to the Committee, these
36 results are consistent with clinical practice. They noted that even following transplant,
37 adherence is still an issue. An increase in adherence is normally seen immediately after
38 transplantation, but it eventually goes back to the adherence levels from before surgery. In
39 some instances, adherence problems may overlap with a psychological problem. An
40 example is in people with eating disorders not taking medication to manipulate weight.

41 The Committee felt no specific recommendations could be made regarding assessment
42 tools, as no evidence was found. They highlighted there are no available tools specific to
43 people with cystic fibrosis. They noted that, although assessment tools are helpful to assess

1 the severity of the psychological or behavioural disorder, in practice psychologists are able to
2 intervene without a formal diagnosis.

3 Following this discussion, the Committee agreed a psychologist with expertise in cystic
4 fibrosis should be an integral member of the multi-disciplinary team. They agreed people with
5 cystic fibrosis should be seen by the psychologist as part of the annual review. However, all
6 members in the multidisciplinary team should be aware of how to identify psychological and
7 behavioural problems.

8 The annual review should be individualised, depending on the circumstances of the person.
9 But as a general guidance it should cover aspects such as general mental health and quality
10 of life, behavioural problems impacting on health outcomes, adherence to treatment, school
11 attendance, friendship and social life. If a severe problem is identified as part of the annual
12 screening, the person should be referred to a mental health practitioner. This is already
13 consistent with current practice. Detailed guidance on identification and management for
14 specific mental health conditions can be found at other NICE guidelines (Common mental
15 health problems: identification and pathways to care [CG123], Depression in children and
16 young people [CG28], Depression in adults [CG90], Generalised anxiety disorder and panic
17 disorder in adults [CG113], Eating disorders in over 8s: management [CG9]).

18 10.9.7.3 Consideration of economic benefits and harms

19 Effective psychological and behavioural assessments will lead to improved identification and
20 management. Therefore there are potentially important indirect resource implications.
21 However, the Committee stated that psychological and behavioural assessment tools are not
22 validated for people with cystic fibrosis who differ to the general population as they have a
23 multi-system disorder which requires a lifetime of frequent and multiple daily treatments that
24 negatively impacts their daily life.

25 If treatment regimens are not followed, treatment effectiveness and hence cost-effectiveness
26 will be reduced. This is especially the case if unused treatments are followed by another
27 supply, or if the dose is increased because the former was thought to be too low. The
28 Committee highlighted that some people with cystic fibrosis have high cost contracts with
29 drugs such as Ivacaftor. They noted that such contracts should only be offered to people with
30 cystic fibrosis who are treatment compliant to ensure the benefits from treatment can
31 outweigh the high cost of treatment.

32 The Committee stated that in times of austerity psychologists are often reliant on the
33 remaining multidisciplinary team to complete psychological and behavioural assessments on
34 their behalf. The Committee added that the number of people with cystic fibrosis with
35 psychological and behavioural problems is growing as there is little time for psychologists to
36 take preventative measures. In addition, psychologists could be at risk of missing emerging
37 problems in people who have not been referred if the times for identification are limited.
38 Ideally, the Committee advised that psychologists should be available to see people with
39 cystic fibrosis at each clinic. This will allow for quicker identification and prevention and
40 minimise the decrements in quality of life and treatment costs associated with psychological
41 and behavioural problems. To reflect current clinical practice, the Committee prioritised a
42 recommendation to suggest the minimum level of interaction psychologists should have with
43 people with cystic fibrosis as an outpatient or inpatient and their families and carers, to
44 prevent geographical variation.

45 10.9.7.4 Quality of evidence

46 The studies included in the review aimed to establish the reliability and validity of
47 assessment tools, and the following were considered as the main criteria for assessing the
48 quality of each study, as reported by Jerosch-Herold (2005):

- 49 • Sample size

- 1 • Sampling methodology
- 2 • Blinding of raters
- 3 • Statistical analysis

4 Main reason of bias in the included studies were little information given on whether
5 observer/tester were appropriately trained or certified.

6 One study was only available as conference abstract, and the quality could not be assessed.

7 In 1 study extreme inaccuracy was observed for individual participants by clinicians and self-
8 report adherence. In another study individuals demonstrated wide variability in regards to the
9 different measures of adherence.

10 **10.9.7.5 Other considerations**

11 The Committee discussed whether the cystic fibrosis psychologist should be available to
12 meet people with cystic fibrosis and their families or carers at every cystic fibrosis clinic if
13 necessary, but they highlighted it is not happening in current practice. This is particularly the
14 case in adult clinics, as the number of psychologists have remained the same, whereas the
15 number of adult people with cystic fibrosis has increased. While this may lead to a resource
16 impact, it was noted that early identification and intervention can help prevent more serious
17 issues in the future.

18 The Committee stated that when a psychologist is introduced at diagnosis stage as a natural
19 member of the team (for example a nurse) then people with cystic fibrosis and their carers
20 are more accepting of their role.

21 The Committee suggested some strategies effective at identifying the presence of
22 psychological or behavioural problems.

- 23 • Informal but frequent “screening” by the psychologist and multidisciplinary team members
24 by regularly asking questions about emotional wellbeing as well as physical health.
- 25 • The team clinical psychologist supports the rest of the team in this by encouraging it,
26 helping staff if they feel out of depth with the answers they get, providing training on
27 delivery of basic emotional support.
- 28 • Team clinical psychologist presence at patient case discussions of outpatients, inpatients,
29 complex case meetings and annual review reports.
- 30 • Annual assessment by clinical psychologist, using face to face professional assessment
31 and standardised measures of psychological wellbeing and mental health functioning.
- 32 • Information to people with cystic fibrosis regarding psychology services and support
33 available and easy access through self-referral.
- 34 • Increased access to clinical psychology and review of vulnerable groups and people at
35 particular complicated stages of cystic fibrosis (for example newly diagnosed, mental
36 health problems in parents/carers, substance misuse, pre- and post-transition, end stage
37 illness, pregnancy and assisted conception, secondary diagnosis, referral for transplant).
- 38 • Provide support for people with cystic fibrosis and training for staff around sensitive issues
39 that may be difficult to discuss. In particular, anxieties relating to transplant decisions and
40 people’s wishes for end-of-life care.

41 The Committee discussed potential equality issues. They noted psychological problems may
42 be more likely in people from lower socio-economic groups. However they agreed care is
43 available to all people, and therefore there was no need to draft additional recommendations.

44 The Committee agreed there was a lack of evidence to determine the best objective
45 measures of adherence. Difficulties with full adherence to all components of the cystic
46 fibrosis daily treatment regimen were well documented and directly resulted in poor health

1 outcomes. Studies indicated that reports of rates of adherence varied and so subjective
2 measures were unreliable. There needed to be further development of objective measures of
3 medication and other treatment usage so data on accurate adherence could be captured.
4 Understanding the occasions when people with cystic fibrosis do and don't follow prescribed
5 recommendations and which types of treatment are more likely to be taken than others would
6 help adapt prescribed treatments to support increased likelihood of compliance, leading to
7 better health outcomes. The area was not prioritised for a research recommendation but the
8 Committee wanted it noted.

9 The Committee agreed a research recommendation around psychological assessment. As
10 noted previously, there are no validated tools to assess psychological and behavioural
11 problems in people with cystic fibrosis and the Committee thought it would be useful to
12 validate generic measures (for example for depression and anxiety). They noted that people
13 with a long term physical health condition are more likely to present with psychological and
14 mental health difficulties than people without a physical health condition. NHSE
15 recommendations are that prevention of psychological problems is the most cost-effective
16 service provision. This means that all people with cystic fibrosis must be routinely and
17 regularly assessed not just for their physical health status but also their mental health status.
18 People with cystic fibrosis would benefit, therefore, in having a routine screen which would
19 indicate those who require further psychological intervention. This would allow early
20 intervention by a team psychologist to enable maintenance of good quality of life, prevention
21 of the development of mental health disorders and improvement in health outcomes as a
22 result of improved wellbeing.

23 **10.9.7.6 Key conclusions**

24 The Committee concluded that the cystic fibrosis psychologist should be an integral member
25 of the multidisciplinary team and should be available to people with cystic fibrosis and their
26 families or carers. A psychological and behavioural assessment should be part of the annual
27 review. People should be referred to a mental health practitioner if a serious issue is
28 identified.

29 **10.9.8 Recommendations**

30 **128. A specialist cystic fibrosis clinical psychologist (see Multidisciplinary team)**
31 **should be available to see people with cystic fibrosis at outpatient clinic visits and**
32 **during inpatient admissions.**

33 **129. The specialist cystic fibrosis clinical psychologist should assess the needs of**
34 **the family members or carers (as appropriate) of people with cystic fibrosis.**

35 **130. At the annual review, the specialist clinical psychologist should include**
36 **assessments of:**

- 37 • general mental health and wellbeing
- 38 • quality of life
- 39 • any factors that are making treatment adherence difficult
- 40 • psychosocial indicators
- 41 • behaviours that affect health outcomes.

42 **131. If a serious mental health issue is identified at the annual review or at any**
43 **assessment performed by the cystic fibrosis clinical psychologist, refer the**
44 **person with cystic fibrosis to a mental health practitioner. For more guidance on**
45 **treating depression, anxiety or panic disorder in adults, see the NICE guidelines**

1 on depression in adults and generalised anxiety disorder and panic disorder in
2 adults.

3 **10.9.9 Research recommendations**

4 **6. What is the most effective measure of psychological functioning to use screening**
5 **test for thresholds of concern for people with cystic fibrosis?**

6 **Table 181: Research recommendations rationale**

Research question	What is the best routine measure of psychological functioning to be used as a method of screening for thresholds of concern for people with cystic fibrosis?
Why this is needed	
Importance to 'patients' or the population	People with a long term physical health condition are more likely to present with psychological and mental health difficulties than people without a physical health condition. NHSE recommendations are that prevention of psychological problems is the most cost-effective service provision. This means that all people with cystic fibrosis must be routinely and regularly assessed not just for their physical health status but also their mental health status. People with cystic fibrosis would benefit therefore in having a routine screen which would indicate those who require further psychological intervention. This would allow early intervention by a team psychologist to enable maintenance of good quality of life, prevention of the development of mental health disorders and improvement in health outcomes as a result of improved wellbeing.
Relevance to NICE guidance	High: There is a need to understand the particular psychological and mental health needs of people with cystic fibrosis. Mental health difficulties decrease physical health outcomes and so understanding the extent of psychological needs of the cystic fibrosis population especially as an expanding population with diagnosis now in infancy and longer lifespan, NICE guidance might have to change to accommodate the psychological health as well as physical health needs. Recommendations to treat psychological health could reduce drug costs by improving attitude to health management and improving likelihood of being able to contribute economically to the workforce.
Relevance to the NHS	Early intervention to prevent psychological and mental health diagnosis will save NHS money.
National priorities	Yes: Future in Mind, NHSE, 2015; Five Year Forward View for Mental health, NHSE, 2016
Current evidence base	Research questions the evidence base is sparse. Though a few papers have moderate quality, the research has not been done.
Equality	Yes, it is anecdotally reported that those children and adults with cystic fibrosis who are most likely to suffer from mental health difficulties and be less likely to manage a complicated daily treatment regimen are over-represented in the most vulnerable societal groups.
Feasibility	Yes, though it would need to be a large scale study but potential use of the national cystic fibrosis registry for this research question
Other comments	Highlighted as an area for research development by Guideline Committee, no existing literature in this area, of benefit for clinical assessment, direction of appropriate treatment resource, research and national registry data.

7 **Table 182: Research recommendation statements**

Criterion	Explanation
Population	Children and young people with cystic fibrosis aged 6 years and over, and their parents or carers. Adults with cystic fibrosis.

Criterion	Explanation
Index test	Administration of existing outcome and screening measures of psychological wellbeing and mental health.
Reference standard	For diagnostic: <ul style="list-style-type: none"> • Diagnosis statistical manual (DSM) or International Classification of diseases (ICD) diagnosis for anxiety, depression, mood disorders, emotional distress and eating disorders
Reference population	For validation: <ul style="list-style-type: none"> • Other physical health condition population • Nominated well participants (for example siblings, schools, colleges)
Outcomes	<ul style="list-style-type: none"> • Diagnostic outcomes: • Sensitivity • Specificity • Positive likelihood ratio • Negative likelihood ratio • Psychometric properties • Reliability • Validity
Study design	Diagnostic cohort study (to conduct logistical regression analyses to look for predictors of wellbeing, co-morbidities, agreement between measures and concordance between psychological symptoms and physical symptoms)
Timeframe	Two years

11 Prevention of cross infection

2 Review questions:

- 3 • What is the effectiveness of cohorting on the basis of pathogen status versus not
- 4 cohorting on the basis of pathogen status in reducing transmission of CF
- 5 pathogens?
- 6 • What is the effectiveness of different models of segregating patients in reducing
- 7 transmission of CF pathogens?
- 8 • What is the effectiveness of individual protective equipment in reducing
- 9 transmission of CF pathogens?
- 10 • What is the effectiveness of the combination of cohorting, segregating and
- 11 protective equipment in reducing transmission of CF pathogens?

12 11.1 Introduction

13 Measures to reduce the risk of cross infection with transmissible pathogens are widely
14 accepted as good/optimum clinical practice in the care of people with cystic fibrosis. Prior to
15 the 1990's it was not unusual for people with cystic fibrosis to attend social events together
16 and share hospital waiting areas. However, landmark evidence of the transmission of
17 bacterial lung infection between people treated at the same hospital and the emergence of
18 new bacteria, has led to care providers adopting increasingly stringent infection control
19 strategies.

20 Segregation in single rooms on a ward or in a clinic, cohorting clinics by pathogen status,
21 discouraging social contact and the use of personal protective equipment are all strategies
22 employed in isolation or combination in cystic fibrosis care in the UK. Despite
23 acknowledgement that it is essential to use these measures to reduce the risk of cross
24 infection, people with cystic fibrosis, families and carers can express feelings of anxiety and
25 social isolation as a result, especially where such measures are employed in a varied or
26 inconsistent manner.

27 This review will therefore look at the evidence for the effectiveness of cohorting by pathogen
28 status; the different models of segregation and personal protective equipment and the
29 effectiveness of the combination of cohorting, segregating and protective equipment in
30 reducing the transmission of cystic fibrosis pathogens.

31 11.2 Description of clinical evidence

32 The aim of this review was to determine the effectiveness of the different strategies (such as
33 cohorting, segregation, or protective equipment) in reducing the transmission of cystic
34 fibrosis pathogens.

35 The interventions that are reviewed are either cohort segregation by time (for example by
36 clinic schedule), cohort segregation by location (for example separate clinics or separate
37 wards), individual segregation by location (for example patients in separate rooms with en-
38 suite facilities), use of protective equipment, or any combination of these interventions.

39 One single literature search was run for the review questions. We looked for systematic
40 reviews, RCTs and prospective and retrospective comparative cohort studies that were
41 conducted in Western countries. Studies based on registry and audit data from the UK were
42 also eligible for inclusion. Conference abstracts of RCTs were considered if RCTs were
43 unavailable. However, given that only 1 RCT and no cohort studies were identified for
44 inclusion, before-and-after implementation studies were also considered eligible for inclusion
45 where no data for critical outcomes was available from higher quality studies. Evidence from

1 questionnaires conducted within cross sectional studies was also considered (for example for
2 quality of life or patient satisfaction).

3 For full details see review protocol in Appendix D.

4 In total 16 studies were included in this review. There was 1 RCT (Hayes 2010), 3 surveys
5 (Griffiths 2004, Russo 2006, Waine 2007) and 12 retrospective before-and-after studies
6 (Chen 2001, France 2008, Frederiksen 1999, Griffiths 2005, Griffiths 2012, Hoiby &
7 Pedersen 1989, Jones 2005, Lee 2004, McKay 2009, Savant 2014, Thomassen 1986,
8 Whiteford 1995)

9 Depending on the study, interventions were implemented in the following setting: either
10 outpatient, inpatient or mixed (both outpatient and inpatient) settings. For the purpose of
11 structuring the review, interventions implemented in the same setting are grouped together.

12 In the outpatient setting, the following comparisons were assessed:

- 13 • Comparison 1. Cohort segregation by clinic times versus no cohort segregation (2 studies:
14 Hayes 2010, McKay 2009)
- 15 • Comparison 2. Cohort segregation by location versus no cohort segregation (1 study: Lee
16 2004)
- 17 • Comparison 3. Combination of protective equipment + individual segregation versus
18 incomplete protective equipment + incomplete individual segregation (1 study: Savant
19 2014).

20 In the inpatient setting, the following comparisons are assessed:

- 21 • Comparison 4. Cohort segregation by location versus no cohort segregation (2 studies:
22 Chen 2001, Thomassen 1986)
- 23 • Comparison 5. Individual segregation by location versus usual care (1 study: Russo
24 2006).

25 When interventions were implemented both in the inpatient and outpatient setting, the
26 following comparisons were assessed:

- 27 • Comparison 6. Cohort segregation versus no cohort segregation (7 studies: France 2008,
28 Frederiksen 1999, Griffiths 2005, Griffiths 2012, Hoiby & Pedersen 1989, Jones 2005,
29 Whiteford 1995)
- 30 • Comparison 7. Complete cohort segregation versus incomplete cohort segregation (1
31 study: France 2008)
- 32 • Comparison 8. Individual segregation versus usual care (1 study: Waine 2007)
- 33 • Comparison 9. Cohort segregation + individual segregation versus cohort segregation (2
34 studies: Chen 2001, France 2008)
- 35 • Comparison 10. Cohort segregation + individual segregation + protective equipment
36 versus usual care. (1 study: Chen 2001)
- 37 • Comparison 11. Cohort segregation + individual segregation versus usual care (1 study:
38 Griffiths 2004).

39 The reported size of the studies ranged from 39 to 232 participants with cystic fibrosis,
40 however for some studies the total size was not reported. 3 studies included adults (France
41 2008, Jones 2005, Waine 2007), 2 included children and young people and their parents
42 (Griffiths 2004, Russo 2006), 2 included infants and children (Hayes 2010, McKay 2009), 1
43 included infants, children and young people (Whiteford 1995), 1 included infants, children,
44 young people and adults (Savant 2014). 4 included people receiving paediatric care,
45 however the age range was not reported (Griffiths 2005, Griffiths 2012, Lee 2004,
46 Thomassen 1986). In 3 studies age of the study population was not reported (Chen 2001,
47 Frederiksen 1999, Hoiby & Pedersen 1989).

1 6 studies were conducted in the UK (France 2008, Jones 2005, Lee 2004, Russo 2006,
2 Waine 2007, Whiteford 1995), 4 in the USA (Chen 2001, Hayes 2010, Savant 2014,
3 Thomassen 1986), 2 in Denmark (Frederiksen 1999, Hoiby & Pedersen 1989), 4 in Australia
4 (Griffiths 2004, Griffiths 2005, Griffiths 2012, McKay 2009)

5 A summary of the included studies is presented in Table 159. See study selection flow chart
6 in Appendix F, study evidence tables in Appendix G, list of excluded studies in Appendix H,
7 forest plots in Appendix I, and full GRADE profiles in Appendix J.

8 11.3 Summary of included studies

9 A summary of the studies that were included in this review are presented in Table 183.

10 **Table 183: Summary of included studies**

Study	Intervention/ Comparison	Population	Outcomes	Comments
OUTPATIENT SETTINGS				
Cohorting into different pathogens by clinic times				
Hayes 2010 USA RCT	<p>Intervention: Cohort segregation</p> <ul style="list-style-type: none"> Segregated clinics free of patients with PA held on a separate day in the same clinic space used for mixed clinics; Large clinics and waiting rooms, and hygienic precautions. <p>Comparison: No cohort segregation</p> <ul style="list-style-type: none"> Mixed clinics that included PA positive patients; Large clinics and waiting rooms, and hygienic precautions. 	<p>Infants and children with cystic fibrosis. N=39 intervention group: n=21 comparison group: n=18 Age: not reported</p>	<p>Incidence of PA infection over 10 years</p>	<p>Study dates 1996-2005</p>
McKay 2009 Australia Retrospective before and after study	<p>Intervention: Cohort segregation by age.</p> <ul style="list-style-type: none"> Outpatients clinics were designated by colour as "red" (children 5 and under who were PA-free), "blue" (primary school age or children under 5 already colonised with PA) or "green" (secondary school age). Additional infection measures (for example removal of toys from the waiting room and hand cleansing). <p>Comparison: No cohort segregation</p> <ul style="list-style-type: none"> One all age (0-18) clinic Free mixing of patients in waiting area 	<p>Infants and children with cystic fibrosis. N=Between 72 and 90 were seen in each year of the study The results of 2837 sputum cultures were analysed for the study Age: ≤5 years</p>	<p>Prevalence of culture results (MRSA, non-mucoid PA, mucoid PA) Staff compliance</p>	<p>Segregation policy introduced in April and May 2003 and outcome data for 1999-2002 versus 2004-2007 The paper also mentions that all inpatients were treated in single rooms or in rooms shared with children without cystic fibrosis. However it is unclear if this</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
				policy started in 2003.
Cohorting into different pathogens by location				
Lee 2004 UK Retrospective before and after study	<p>Intervention: Cohort segregation.</p> <ul style="list-style-type: none"> Separate clinics for patients chronically infected with PA and uninfected patients. Improved hygienic measures in a purpose-built cystic fibrosis centre <p>Comparison: No cohort segregation</p> <ul style="list-style-type: none"> Various management strategies over the years to reduce the prevalence of chronic PA, including regular microbiological monitoring 	<p>People with cystic fibrosis receiving paediatric care.</p> <p>N=232 people 1990: n=966 patient months (mean age: 7.73) 2000: n=1803 patient months; (mean age: 9.42)</p>	<p>Incidence of PA infection</p> <p>Prevalence of chronic PA infection</p> <p>Prevalence of intermittent PA infection</p>	<p>Patients receiving full-time care at the Leeds Paediatric CF Centre. Separate clinics at the Leeds Regional CF Unit.</p> <p>Intervention implemented in 1991, outcome data provided for 1990 and 2000.</p> <p>Incidence reported narratively. Prevalence data calculated based on patient months</p>
Combination strategies				
Savant 2014 USA Retrospective before and after study	<p>Intervention: Protective equipment and individual segregation</p> <ul style="list-style-type: none"> Contact precautions for all patients in the outpatient clinic, regardless of respiratory tract culture results: gowning and gloving and hand hygiene by all providers; requesting that all patients use hand gel and mask when entering the facility or when outside of the exam room Abolishing the designated communal area for taking vital sign measurements and converting to exam rooms. Re-enforcement of the "no-waiting" room policy (Immediate placement within the examination room) Education of patients and families; cleaning rooms thoroughly. <p>Comparison: Incomplete use of</p>	<p>People with cystic fibrosis receiving paediatric care.</p> <p>N= ranged from 126 to 177 during the study years. Age range: 0 to 21.</p>	<p>Prevalence (% of patients cultured each quarter with positive tract cultures) of PA infections</p> <p>Prevalence (% of patients cultured each quarter with positive tract cultures) of MRSA infections</p>	<p>Mean no. of respiratory tract cultures per quarter = 169</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>protective equipment and incomplete individual separation</p> <ul style="list-style-type: none"> Any patient with respiratory tract cultures revealing a multi-resistant pathogen had a flag placed on the chart to indicate the need for contact precautions. However a consistent process for the use of this indicator was not systematic or routine. Vital signs were performed in a common station in the hallway close to the exam rooms, without specific cleaning between patients. “No-waiting” room policy 			
INPATIENT SETTINGS				
Cohorting into different pathogens by location				
<p>Chen 2001 USA Retrospective before and after study</p>	<p>Intervention 1. Cohort segregation.</p> <ul style="list-style-type: none"> Cohorting of hospitalized patients with cystic fibrosis on the basis of <i>B cepacia</i> colonization status <p>Comparison 1: No cohort segregation No details</p>	<p>People with cystic fibrosis. N not reported Age not reported.</p>	<p>Incidence of <i>B cepacia</i> complex infection</p>	<p>Intervention introduced in early 1990, incidence data for 1988, 1989, 1990 and 1991. We compare 1989 and 1991 (the closest pre-intervention full year and the closest post-intervention full year)</p>
<p>Thomassen 1986 USA Retrospective before and after study</p>	<p>Intervention: Incomplete cohort segregation</p> <ul style="list-style-type: none"> All patients with <i>Pseudomonas cepacia</i> recovered from sputum or throat culture were admitted to 1 floor of the hospital - other patients with cystic fibrosis were not admitted to this floor. The patients on this ward were not permitted to visit other inpatient wards but were not isolated in any other way. They had free access to the elevators, hospital cafeteria, and other common areas. No special precautions were taken to totally avoid chance meetings of the 2 groups of patients in the radiology department, outpatient areas, 	<p>People with cystic fibrosis receiving paediatric care admissions pre-segregation: n=453 admissions post-segregation: n=389 Age: not reported</p>	<p>Incidence of hospital-associated colonisation of <i>P. cepacia</i></p>	<p>Intervention introduced in August 1983. Data on 1 year and 5 months before the intervention compared to 1 year and 5 months afterwards Serotyping</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>pulmonary function laboratory, or other hospital areas.</p> <ul style="list-style-type: none"> • Equipment in pulmonary function lab was sterilized or changed between patients. Handwashing was emphasized. • Masks or other isolation equipment were not used by patients or hospital personnel. • Summer camp facility was reserved for patients free of P. cepacia, another camp site was provided for colonized patients. <p>Comparison: No cohort segregation in hospital</p> <ul style="list-style-type: none"> • Basic infection control procedures (no details) • No segregation in camp facility 			
Individual segregation				
Russo 2006 UK Survey. Questionnaire with both closed-ended and open-ended questions	<p>Intervention: individual segregation.</p> <ul style="list-style-type: none"> • Policy of segregation requiring all patients to remain in their individual rooms for the duration of their hospital stay <p>Comparison: Usual care</p>	<p>Children and young people with cystic fibrosis and their parents N= 192 parents, 101 people with cystic fibrosis. Mean age of eligible participants: 13 (range 10 to 17)</p>	<p>Patient and carer satisfaction: % of parents and children who supported segregated treatment</p>	<p>Views were elicited in preparation for the process of implementing a formal policy of segregation</p>
MIXED POPULATIONS (BOTH OUTPATIENT AND INPATIENT SETTINGS)				
Cohorting into different pathogens				
France 2008 UK Retrospective before and after study	<p>Intervention 2. Complete cohort segregation.</p> <ul style="list-style-type: none"> • Patients with Burkholderia species infection were cohorted into separate wards to non-Bcc infected patients and each inpatient has their own single room. • Patients with Burkholderia species infection have also continued to attend a different outpatient clinic. • Isolation policy for patients with B. species not yet implemented <p>Intervention 1: Incomplete cohort segregation.</p>	<p>Adults with cystic fibrosis. N not reported. Age not reported.</p>	<p>Incidence of Burkholderia species infection</p>	<p>Incomplete inpatient and complete outpatient A lot of graphic reporting, some numbers are mentioned, see below: Intervention 1 versus comparison 1: Intervention 1 implemented in 1991,</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<ul style="list-style-type: none"> Patients with Burkholderia species infection were admitted to inpatient beds on the opposite side of the corridor to non-Burkholderia species infected patients. There was continued patient mixing within a day-room facility on the ward and within areas such as the radiology department. Patients with Burkholderia species infection attended separate outpatient clinics to other cystic fibrosis patients. <p>Comparison 1. No infection control measures to prevent <i>B cepacia</i> complex cross-infection No details</p>			<p>range of incidence of Burkholderia species infection for 1983-1990, incidence 1992.</p> <p>Intervention 2 versus intervention 1: Intervention 2 implemented from November 1993 onwards. Incidence 1992 and incidence "following complete cohort segregation"</p>
Frederiksen 1999 Denmark Retrospective before and after study	<p>Intervention: Cohort segregation:</p> <ul style="list-style-type: none"> The cystic fibrosis centre was reconstructed, separating the wards and the outpatient clinic Patients with PA in their sputum were separated from patients without PA in the wards, in the outpatient clinic, and during social events. <p>Comparison: No cohort segregation</p> <ul style="list-style-type: none"> The wards with inpatients receiving iv treatment were near the outpatient clinic visited by all cystic fibrosis patients Cystic fibrosis patients were not segregated according to presence or absence of PA in their sputum 	People with cystic fibrosis. 1974: n=107. Median age: 9.0 1995: n=256. Median age: 18.5	Incidence of intermittent PA infection Incidence of chronic PA infection	Cohort segregation introduced in 1981. Incidence given for each year from 1974 and 1995, we compare 1980 and 1982.
Griffiths 2005 and 2012 Australia Retrospective before and after study	<p>Intervention: Cohort segregation</p> <ul style="list-style-type: none"> Cohorts: PA (negative, positive non-epidemic, epidemic PA (AES-1)) Separation of cohorts was maintained at outpatient visits and during hospital admissions. Standard infection control measures were reinforced, and education seminars were 	People with cystic fibrosis receiving paediatric care. 1999: n=325 Sputum producers: n=153 2002: n=291 Sputum	Prevalence of AES-1 (PA epidemic strain)	Cohort segregation in 2000, outcome data for 1999 and 2002

Study	Intervention/ Comparison	Population	Outcomes	Comments
	<p>arranged for staff and families.</p> <p>Comparison: No cohort segregation.</p> <ul style="list-style-type: none"> • No segregation based on PA. • Standard infection control measures, <i>B cepacia</i> complex and MRSA strict individual segregation 	<p>producers: n=149</p> <p>Age range: from <10 to ≥16 years, exact age range not reported.</p>		
<p>Hoiby & Pedersen 1989</p> <p>Denmark</p> <p>Retrospective before and after study</p>	<p>Intervention 2: Cohort segregation (by multiply resistant PA strain)</p> <ul style="list-style-type: none"> • Three groups: Patients with multiply resistant PA strain; cohort with normally sensitive strains of chronic PA infection; PA negative patients • Improved hygiene precautions <p>Intervention 1. Cohort segregation (by PA infection, but not multiply resistant PA strain)</p> <ul style="list-style-type: none"> • Two cohorts, cohort with chronic PA infection separated from PA negative patients • Segregated from each other in different wards and seen on different days in the outpatient clinic. 	<p>People with cystic fibrosis. N (range): 54 to 226 (between 1970 and 1987). Subgroup of people with PA infection in 1983: n=119 Age: Not reported</p>	<p>Incidence of multiply resistant PA strain</p> <p>Prevalence of multiply resistant PA strain</p>	<p>Intervention 2 (compared to intervention 1) was implemented in April 1983. Incidence and prevalence of multiply resistant PA is provided for Jan, Feb, Mar, Apr, May, June 1983</p>
<p>Jones 2005</p> <p>UK</p> <p>Retrospective before and after study</p>	<p>Intervention 1: Incomplete cohort segregation.</p> <ul style="list-style-type: none"> • Patients without chronic PA infection attended outpatient clinic appointments on a different day than other patients with cystic fibrosis. • Inpatients without chronic PA infection were housed on the same cystic fibrosis ward as patients with chronic PA infection, but in rooms with en-suite facilities, and were advised not to socialise with other patients on the ward. However there was still some social mixing between patients on the ward. <p>Comparison: No cohort segregation.</p> <ul style="list-style-type: none"> • Purpose-built facilities: all inpatients had their own bedroom, although only 2 of 11 rooms had en-suite facilities. • Treatments with door closed • Practice of strict hygiene. 	<p>People with cystic fibrosis who attend an adult centre. 1999: n= 216 2000: n=221 2001: n=228 Age not reported</p>	<p>Prevalence of chronic PA infection</p> <p>Prevalence of transmissible PA infection</p> <p>Prevalence of chronic infection with transmissible PA strain</p>	<p>Outpatient policy of segregation instituted in 2000. Outcome data provided for 1999, 2000, 2001, 2002 and 2003. We compare 1999 to 2001</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
	Rooms are cleaned between patients, equipment not shared between patients, hand hygiene for staff.			
Whiteford 1995 UK Retrospective Before and after study	<p>Intervention. Cohort segregation.</p> <ul style="list-style-type: none"> • Children with <i>B cepacia</i> were admitted to a separate ward • Children with <i>B cepacia</i> were moved to a different waiting area and given appointment times at the end of the clinic. • Recommendations to parents to avoid close physical contact outside the hospital <p>Comparison: no cohort segregation</p> <ul style="list-style-type: none"> • No cohort segregation for children with <i>B cepacia</i> • Children with cystic fibrosis needing inpatient care were admitted to 1 ward where they had complete freedom to play together and socialise • Each child had an individual peak flow meter and nebuliser for use throughout the admission • Children colonised with PA had their chest physiotherapy separately from children without PA 	Infants, children and young people with cystic fibrosis. N=115. Mean (range) age: 7.6 years (0.6 to 15.8 year)	Incidence of <i>B cepacia</i> infection	Intervention introduced in June 1992; number of cases given for each month of 1992. , 114 at risk in Jan 1992
Individual segregation				
Waine 2007 UK Survey	<p>Intervention: Individual separation Not mixing with others with cystic fibrosis</p> <p>Comparison: no separation Mixing with others with cystic fibrosis</p>	Adults with cystic fibrosis. N=94 respondents (184 invited to participate) Mean (SD) age: 27.2 (8.5) years	Patient satisfaction % of those who mixed with others who said that if they avoided others with cystic fibrosis, their quality of life would be: significantly or greatly affected / not suffer	All patients attending a clinic appointment/inpatients at the West Midlands Adult CF Centre were offered a questionnaire . Questionnaires were posted to 8 patients colonized with the <i>B cepacia</i> complex who attended a separate clinic.
Combination of strategies				

Study	Intervention/ Comparison	Population	Outcomes	Comments
Chen 2001 USA Retrospective before and after study	<p>Intervention 2: Individual segregation in addition to cohort segregation</p> <ul style="list-style-type: none"> Hospitalized non-colonized patients were prohibited from sharing rooms, then this policy was expanded so that all patients with cystic fibrosis irrespective of <i>B cepacia</i> colonization were in separate rooms. Separate waiting rooms were established for outpatients. <p>Intervention 1. Cohort segregation.</p> <ul style="list-style-type: none"> Cohorting of hospitalized patients with cystic fibrosis on the basis of <i>B cepacia</i> colonization status <p>Intervention 3: Cohort segregation combined with individual segregation and with protective equipment.</p> <ul style="list-style-type: none"> Cohorting of hospitalized patients on the basis of <i>B cepacia</i> colonization status. Furthermore inpatients colonized with <i>B cepacia</i> were placed in contact isolation and were required to wear mask and gloves when out of their rooms. In the outpatient setting, patients infected with <i>B cepacia</i> were restricted to separate clinic days during which no non-colonized patients were seen. Patients were required to wear masks while in the waiting room. A hospital-wide educational program regarding infection control measures was introduced Particular attention was given to disinfection of clinic rooms and equipment <p>Comparison 3: Usual care No details</p>	People with cystic fibrosis. N not reported. Age not reported.	Annual incidence of <i>B cepacia</i> complex infection. Prevalence of <i>B cepacia</i> complex infection	Intervention 2 versus intervention 1: Progressive isolation measures in mid-1996, 1997, 1998, and prevalence data for 1992 and 1999 Intervention 3 versus comparison 3: Measures introduced in early 1997, annual incidence given for 1996, 1997 and "since time of implementation"
France 2008 UK	<ul style="list-style-type: none"> Intervention 3. Cohort segregation combined with individual segregation. 	Adults with cystic fibrosis. N not reported.	Prevalence of Burkholderia species	Policy of isolation in 2000

Study	Intervention/ Comparison	Population	Outcomes	Comments
Retrospective before and after study	<ul style="list-style-type: none"> • A policy of isolation was introduced for patients infected with all Burkholderia species. This policy involves patients not having any contact with other patients, either at an inpatient or outpatient level. • Patients being admitted to single rooms during admissions and attending outpatient appointments and being immediately isolated within their own clinic room. • Patients with Burkholderia species infection were cohorted into separate wards to non-Bcc infected patients and attended a different outpatient clinic. <p>Intervention 2. Cohort segregation.</p> <ul style="list-style-type: none"> • Patients with Burkholderia species infection were cohorted into separate wards to non-Bcc infected patients and each inpatient had their own single room. • Patients with Burkholderia species infection attended a different outpatient clinic. • Isolation policy for patients with Burkholderia species not yet implemented 	Age not reported.	Prevalence of transmissible <i>B cepacia</i> complex infection.	Peak prevalence of Burkholderia species infection 1994 and prevalence 2005. Prevalence of B cepacia complex transmissible infections for 1993/94 and for 2005.
Griffiths 2004 Australia Survey	<p>Intervention: Combination of cohort and individual segregation.</p> <ul style="list-style-type: none"> • Cohort segregation was based on five separate groups: PA positive; epidemic strain PA; <i>B cepacia</i>; MRSA; and PA negative. • Inpatients were nursed in separate sections and attended physiotherapy sessions at different times. • Children infected with epidemic strain PA, MRSA or BC were isolated from each other and all other patients • Those within the other groups were allowed to mix within their cohort groups. <p>Comparison: Usual care</p>	<p>People with cystic fibrosis receiving paediatric care and their parents N= 190 questionnaires were completed (291 people were sent the questionnaire) 114 parents alone 75 parents completed the questionnaire together with a person with cystic fibrosis aged >=12 years</p>	<p>Patient and carer satisfaction: Children's and parents' overall response to segregation measures: positive / negative / unsure</p>	<p>Survey was conducted 2 years after the introduction of the intervention. Royal Children's Hospital, Melbourne, not clear if both inpatient and outpatient</p>

Study	Intervention/ Comparison	Population	Outcomes	Comments
		1 questionnaire completed by child only. Mean age: unclear		

1 PA: *P aeruginosa*; B cepacia: *B cepacia*; MRSA: *Meticillin-resistant S aureus*

2 11.4 Clinical evidence profile

3 The summary clinical evidence profiles for this review question (cross-infection control) are
4 presented in Table 184 to Table 194.

5 11.4.1 Outpatient care

6 **Table 184: Summary clinical evidence profile: Comparison 1. Cohort segregation by**
7 **clinic times versus no cohort segregation**

Outpatient care: Comparison 1. Cohort segregation by clinic times versus no cohort segregation						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No cohort segregation	Cohort segregation into different pathogens by clinic times				
10-year incidence of <i>P aeruginosa</i> infections. Follow-up: 10 years	778 per 1000	622 per 1000 (404 to 941)	RR 0.8 (0.52 to 1.21)	39 (Hayes 2010)	⊕⊕⊖⊖ low ^{1,2}	
4-year prevalence of MRSA (percentages) Follow-up: 4 years	1%	1.3%	ns	≈2,837 sputum cultures (McKay 2009) ⁴	⊕⊖⊖⊖ very low ³	
4-year prevalence of non-mucoid <i>P aeruginosa</i> (percentages) Follow-up: 4 years	22.3%	22.7%	ns	≈2,837 sputum cultures (McKay 2009) ⁴	⊕⊖⊖⊖ very low ³	
4-year prevalence of mucoid <i>P aeruginosa</i> (percentages) Follow-up: 4 years	5.9%	1.0%	P=0.001	≈2,837 sputum cultures (McKay 2009) ⁴	⊕⊖⊖⊖ very low ³	

Outpatient care: Comparison 1. Cohort segregation by clinic times versus no cohort segregation

Staff compliance (percentages) Follow-up: 4 years	-	Adherence to the "coloured" clinic booking scheme: % of children attending the red clinic who were 5 and under: 2004: 96.8%; 2005: 97.5%; 2006: 94.4%; 2007: 95.9%	Not estimable	N not reported (McKay 2009) ⁴	⊕⊖⊖⊖ very low ³	Narrative data
--	---	--	---------------	--	-------------------------------	----------------

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; MRSA: methicillin-resistant *S aureus*; ns: not significant; RR: risk ratio

1 The quality of the evidence was downgraded by 1 due to unclear randomization, allocation concealment, blinding, incomplete data outcome and selective reporting

2 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID

3 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection, comparability between groups and outcome reporting

4 Intervention group: data for the period 2004 to 2007; comparison group: data for the period 1999 to 2002. Intervention introduced in 2003.

Table 185: Summary clinical evidence profile: Comparison 2. Cohort segregation by location versus no cohort segregation

Outpatient care: Comparison 2. Cohort segregation by location versus no cohort segregation

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No cohort segregation	Cohort segregation into different pathogens by location				
Annual incidence of new growths of <i>P aeruginosa</i> . Follow-up: 9 years	-	The annual incidence of new growths of <i>P aeruginosa</i> , while fluctuating, showed no downward trend, despite segregation	Not estimable	232 (Lee 2004) ³	⊕⊖⊖⊖ very low ^{1,2}	Narrative data
Yearly prevalence of chronic <i>P aeruginosa</i> infection Follow-up: 9 years	245 per 1000	181 per 1000 (154 to 210)	OR 0.68 (0.56 to 0.82)	2769 patient months (Lee 2004) ³	⊕⊖⊖⊖ very low ^{1,4}	
Yearly prevalence of	262 per 1000	574 per 1000 (528 to 620)	OR 3.8 (3.15 to	2769 patient	⊕⊖⊖⊖ very low ¹	

Outpatient care: Comparison 2. Cohort segregation by location versus no cohort segregation

intermittent <i>P aeruginosa</i> infection Follow-up: 9 years			4.59)	months (Lee 2004) ³		
--	--	--	-------	--------------------------------	--	--

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; OR: odds ratio

- 1 *1 The quality of the evidence was downgraded by 2 because high risk of bias in relation to sample selection, comparability between groups, and outcome assessment and reporting*
- 2 *2 Imprecision cannot be calculated with the data provided*
- 3 *3 Intervention group: data from 2000; comparison group: data from 1990. Intervention implemented in 1991.*
- 4 *4 The quality of the evidence was downgraded by 1 as the 95% CI crossed 1 default MID*

Table 186: Summary clinical evidence profile: Comparison 3. Combination of protective equipment + individual segregation versus incomplete protective equipment + incomplete individual segregation

Outpatient care: Comparison 3. Combination of protective equipment + individual segregation versus incomplete protective equipment + incomplete individual segregation

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Incomplete protective equipment + incomplete individual segregation	Protective equipment + individual segregation				
4-month prevalence of <i>P aeruginosa</i> infections (percentages). Follow-up: 5 years	The 4-month prevalence of <i>P aeruginosa</i> infections in the control group was: 29.79%	The 4-month prevalence of <i>P aeruginosa</i> infections in the intervention group was: 21.78%	p<0.0001	N varied from 126 to 177 during the study years (Savant 2014) ³	⊕⊕⊕⊖ very low ^{1,2}	
4-month prevalence of MRSA infections (percentages). Follow-up: 5 years	The 4-month prevalence of MRSA infections in the control group was: 10.76%	The 4-month prevalence of MRSA infections in the intervention group was: 8.68%	p=0.008	N varied from 126 to 177 during the study years (Savant 2014) ³	⊕⊕⊕⊖ very low ^{1,2}	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval; MRSA: methicillin-resistant *S aureus*

- 9 *1 The quality of the evidence was downgraded by 2 because of high risk bias in relation to sample selection, comparability between groups and outcome assessment.*
- 10 *2 Imprecision cannot be assessed with the reported data.*
- 11

1 3 Intervention group: mean data for the period 2008 to 2012; comparison group: mean data for the period 2005 to
2 2007. Intervention implemented in 2007.

3 11.4.2 Inpatient care

4 **Table 187: Summary clinical evidence profile: Comparison 4. Cohort segregation by**
5 **location versus no cohort segregation**

Inpatient care: Comparison 4. Cohort segregation by location versus no cohort segregation						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No cohort segregation	Cohort segregation into different pathogens by location				
Annual incidence of <i>B cepacia</i> complex (percentages) Follow-up: 1 year	The annual incidence of <i>B cepacia</i> complex infections in the control group was: 5.8%	The annual incidence of <i>B cepacia</i> complex infections in the intervention group was: 3.7%	Not estimable	N not reported (Chen 2001) ³	⊕⊕⊕⊕ very low ^{1,2}	
5-month incidence of hospital-associated colonisation of <i>P. Cepacia</i> Follow-up: 5 months	78 per 1000	26 per 1000 (10 to 61)	OR 0.31 (0.12 to 0.77)	N of people unclear. N of admissions: ≈842 (Thomassen 1986) ⁵	⊕⊕⊕⊕ very low ⁴	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval; OR: odds ratio						

6 1 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection,
7 comparability between groups and outcome assessment
8 2 Imprecision cannot be calculated with the data reported
9 3 Intervention group: data from 1991; comparison group: data from 1989. Intervention implemented in early 1990.
10 4 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to the comparability
11 between groups and outcome assessment
12 5 Intervention group: data for the period 1 Aug 1983 to 31 Dec 1984; comparison group: data for the period 1 Mar
13 1982 to 31 Jul 1983. Intervention introduced in August 1983.

14 **Table 188: Summary clinical evidence profile: Comparison 5. Individual segregation**
15 **by location versus usual care**

Inpatient care: Comparison 5. Individual segregation by location versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual	Individual				

Inpatient care: Comparison 5. Individual segregation by location versus usual care						
	care	segregation				
Patient satisfaction (narrative)		92% of children supported segregated treatment	na	101 (Russo 2006)	⊕⊕⊕⊕ very low ^{1,2}	Narrative results only
Parents' satisfaction (narrative)		91% of parents supported segregated treatment	na	192 (Russo 2006)	⊕⊕⊕⊕ very low ^{1,2}	Narrative results only

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: CI: confidence interval

- 1 *The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection, the comparability between groups and outcome assessment.*
 2 *The imprecision cannot be calculated with the data reported*

4 11.4.3 Combined inpatient and outpatient care

5 **Table 189: Summary clinical evidence profile: Comparison 6. Cohort segregation**
 6 **versus no cohort segregation**

Inpatient/ outpatient care: Comparison 6. Cohort segregation versus no cohort segregation						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Cohort segregation into pathogens				
Monthly incidence of multiply resistant <i>P aeruginosa</i> strain Follow-up: 1 month	206 per 1000	65 per 1000 (25 to 161)	OR 0.27 (0.1 to 0.74)	119 (Hoiby & Pedersen 1989) ²	⊕⊕⊕⊕ very low ¹	
Annual incidence of intermittent <i>P aeruginosa</i> Follow-up: 1 year	333 per 1000	225 per 1000 (99 to 433)	OR 0.58 (0.22 to 1.53)	Total N unclear (Frederiksen 1999) ⁵	⊕⊕⊕⊕ very low ^{3,4}	
Annual incidence of chronic <i>P aeruginosa</i> Follow-up: 1 year	200 per 1000	101 per 1000 (41 to 229)	OR 0.45 (0.17 to 1.19)	Total N unclear (Frederiksen 1999) ⁵	⊕⊕⊕⊕ very low ^{3,6}	
6-month incidence <i>B cepacia</i> Follow-up: 6 months	46 per 1000	11 per 1000 (1 to 87)	OR 0.23 (0.03 to 1.97)	115 (Whiteford 1995) ⁸	⊕⊕⊕⊕ very low ^{4,7}	
Annual incidence of <i>Burkholderia</i> species infection (percentages) Follow-up: 1 year	3-5%	16.3%	Not estimable	N not reported (France) ¹¹	⊕⊕⊕⊕ very low ^{9,10}	
Monthly prevalence of	328 per 1000	332 per 1000 (226 to 462)	OR 1.02 (0.60 to	119 (Hoiby	⊕⊕⊕⊕ very	

Inpatient/ outpatient care: Comparison 6. Cohort segregation versus no cohort segregation						
multiple resistant <i>P aeruginosa</i> strain (percentages) Follow-up: 1 month			176)	1989) ²	low ^{1,4}	
Prevalence of AES-1 (<i>P aeruginosa</i> epidemic strain) Follow-up: 2 years	-	-	aRR 0.64 (0.47 to 0.87)	Total N unclear (Griffiths 2005) ¹²	⊕⊕⊕⊕ very low ⁶	
Annual prevalence of chronic <i>P aeruginosa</i> infection Follow-up: 1 year	722 per 1000	807 per 1000 (728 to 867)	OR 1.61 (1.03 to 2.51)	Total N unclear (Jones 2005) ¹³	⊕⊕⊕⊕ very low ⁶	
Annual prevalence of transmissible <i>P aeruginosa</i> infection Follow-up: 1 year	130 per 1000	154 per 1000 (96 to 237)	OR 1.22 (0.71 to 2.08)	Total N unclear (Jones 2005) ¹³	⊕⊕⊕⊕ very low ⁴	
Annual prevalence of chronic infection with transmissible <i>P aeruginosa</i> strain (percentages) Follow-up: 1 year	13%	15.4%	Not estimable	Total N unclear (Jones 2005) ¹³	⊕⊕⊕⊕ very low ¹⁰	

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

Abbreviations: adjRR: adjusted risk ratio; ASUSP-1: Australian epidemic strain, type 1; CI: confidence interval; MRSA: methicillin-resistant *S aureus*; OR: odds ratio

- 1 The quality of the evidence was downgraded by 1 because of high risk of bias in relation to comparability of the groups, and outcome reporting
- 2 Intervention group: data from May 1983; comparison group: data from March 1983. Intervention implemented in April 1983.
- 3 The quality of the evidence was downgraded by 1 because of high risk of bias in relation to comparability between groups, and outcome assessment
- 4 The quality of the evidence was downgraded by 2 because the 95% CI crossed 2 default MIDs
- 5 Intervention group: data from 1982; comparison group: data from 1980. Intervention implemented in 1981
- 6 The quality of the evidence was downgraded by 1 because the 95% CI crossed 1 default MID
- 7 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to the comparability between groups, outcome assessment and unclear sample selection
- 8 Intervention group: data from December 1992; comparison group: data from May 1992. Intervention implemented in June 1992.
- 9 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection, comparability between groups and outcome assessment
- 10 Imprecision cannot be calculated with the data reported
- 11 Intervention group: data from 1992; comparison group: data from 1983-1990. Intervention implemented in November 1991. Intervention was incomplete cohort segregation.
- 12 Intervention group: data from 2002; comparison group: data from 1999. Intervention implemented in January 2000.
- 13 Intervention group: data from 2001; comparison group: data from 1999. Intervention implemented in 2000.

1
2

Table 190: Summary clinical evidence profile: Comparison 7. Complete cohort segregation versus incomplete cohort segregation

Inpatient/ outpatient care: Comparison 7. Complete cohort segregation versus incomplete cohort segregation						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Incomplete cohort segregation ³	Complete cohort segregation ³				
Annual incidence of Burkholderia species (percentages). Follow-up: unclear	The annual incidence of Burkholderia species for the control group was: 16.3%	The annual incidence of Burkholderia species for the intervention group was : < 3% (for all but 1 year)	Not estimable	N not reported (France 2008) ³	very low ^{1,2}	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>Abbreviations: CI: confidence interval</p>						

3
4
5
6
7
8
9

1 The quality of the evidence was downgraded by 2 because high risk of bias in relation to sample selection, the comparability between the groups and the outcome reporting and assessment.
2 Imprecision cannot be calculated with the data reported
3 Intervention group: data after 1993; comparison group: data from 1992. Intervention implemented in November 1993.

Table 191: Summary clinical evidence profile: Comparison 8. Individual segregation versus usual care

Inpatient/ outpatient care: Comparison 8. Individual segregation versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual care	Individual segregation				
Patient satisfaction (narrative results)	23.3% of patients said that their quality of life would suffer a 'significant amount' or 'a great deal' if they were to begin avoiding others	62.5% of patients said that their quality of life did not suffer as a result.	Not estimable	94 (Waive 2007)	very low ^{1,2}	Narrative results only
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p>						

Inpatient/ outpatient care: Comparison 8. Individual segregation versus usual care

Abbreviations: CI: confidence interval

1 The quality of the evidence was downgraded by 2 because high risk of bias in relation to sample selection, the comparability between the groups and the outcome reporting and assessment.
2 Imprecision cannot be calculated with the data reported

Table 192: Summary clinical evidence profile: Comparison 8. Individual segregation versus usual care

Inpatient/ outpatient care: Comparison 8. Individual segregation versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Cohort segregation	Cohort segregation + individual segregation				
Yearly prevalence of <i>B cepacia</i> complex infection (percentages) Follow-up: 1 year	The annual incidence of <i>B cepacia</i> complex infections in the control group was: 15%	The annual incidence of <i>B cepacia</i> complex infections in the intervention group was: 7%	Not estimable	N not reported (Chen 2001)3	very low1,2	
Yearly prevalence of <i>Burkholderia</i> species (percentages) Follow-up: 1 year	The annual incidence of <i>Burkholderia</i> species for the control group was: 31.2%	The annual incidence of <i>Burkholderia</i> species for the intervention group was : 9.3%	Not estimable	N not reported (France 2008)5	very low2,4	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval						

6 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection, comparability between groups and outcome assessment
7
8 Imprecision cannot be calculated with the data reported
9 Intervention group: data from 1999; comparison group: data from 1992. Intervention introduced in 1996..
10 The quality of the evidence was downgraded by 2 because high risk of bias in relation to sample selection, the comparability between the groups and the outcome reporting and assessment.
11
12 Intervention group: data from 2005; comparison group: data from 1994. Intervention implemented in 2000.

Table 193: Summary clinical evidence profile: Comparison 10. Cohort segregation + individual segregation + protective equipment versus usual care

Inpatient/ outpatient care: Comparison 10. Cohort segregation + individual segregation + protective equipment versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				

Inpatient/ outpatient care: Comparison 10. Cohort segregation + individual segregation + protective equipment versus usual care						
	Usual care	Cohort segregation + individual segregation + protective equipment				
Annual incidence of <i>B cepacia</i> complex infection (percentages) Follow-up: 1 year	The annual incidence of <i>B cepacia</i> complex infections in the control group was: 8.8%	The annual incidence of <i>B cepacia</i> complex infections in the intervention group was: < 1%	Not estimable	N not reported (Chen 2001) ³	Very low ^{1,2}	
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval						

- 1 The quality of the evidence was downgraded by 2 because of high risk of bias in relation to sample selection, comparability between groups and outcome assessment
 2 Imprecision cannot be calculated with the data reported
 3 Intervention group: data after implementation; comparison group: data from 1996. Intervention implemented in early 1997.

6 **Table 194: Summary clinical evidence profile: Comparison 11. Cohort segregation + individual segregation versus usual care**

Inpatient/ outpatient care: Comparison 11. Cohort segregation + individual segregation versus usual care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Cohort segregation into pathogens				
Patient satisfaction	-	Positive: 63%: Negative: 12%: Unsure: 25% (p<0.001)	Not estimable	190 (Griffiths 2004)	⊕⊖⊖⊖⊖ very low ^{1,2}	Narrative results only
Carer satisfaction	-	Positive: 85%: Negative: 4%: Unsure: 11% (p<0.001)	Not estimable	190 (Griffiths 2004)	⊕⊖⊖⊖⊖ very low ^{1,2}	Narrative results only
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
Abbreviations: CI: confidence interval						


- 8 1 The quality of the evidence was downgraded by 1 because of high risk of bias in relation to sample selection and outcome reporting
 9 2 Imprecision cannot be calculated with the data reported
 10

11.5 Economic evidence

No economic evaluations of strategies to prevent cross-infection were identified in the literature search conducted for this guideline. Full details of the search and economic article selection flow chart can be found in Appendix E and F, respectively.

This area was prioritised for de novo economic modelling, subsequently a cost-utility model was developed. The decision analytic model took the form of a decision tree (Figure 15) where the outcome (probability of transmissible pathogen) is associated with a treatment cost and utility value. The methods used to construct the model and the results of all analyses are reported in Appendix K. A summary of cost-effectiveness estimates are provided in Table 195 for ease of reference.

Figure 15: Decision tree



11

Table 195: Summary of cost-effectiveness estimates

Strategy	Infection with transmissible pathogen			
	Intermittent BCC	Intermittent PA	Chronic PA	Super infection with chronic PA
Cohort segregation	Cost-effectiveness uncertain (dominant to dominated)	Not Cost-Effective (ICER £239,634 to dominated)	Cost-effective (dominant)	NC
Protective equipment	Cost-effective (dominant)	Cost-effective (dominant)	NC	NC
Single inpatient rooms versus beds in shared rooms	Cost-effective (dominant)	NC	NC	NC

Strategy	Infection with transmissible pathogen			
	Intermittent BCC	Intermittent PA	Chronic PA	Super infection with chronic PA
Incomplete cohort segregation including en suite bathroom facilities versus no cohort segregation including shared bathroom facilities	NC	Cost-effective if en suite facilities cost less than an additional £21,600/clinic /year	NC	Cost-effectiveness uncertain

1 BCC, *B cepacia complex*; ICER, incremental cost-effectiveness ratio; NC, not calculable; PA, *P. aeruginosa*

2 11.6 Evidence statements

3 11.6.1 Outpatient care

4 11.6.1.1 Cohorting into different pathogens by clinic times

5 11.6.1.1.1 Comparison 1. Cohort segregation by clinic times versus no cohort segregation

6 Incidence of patients infected with transmissible pathogens

7 Low quality evidence from 1 RCT with 39 infants and children with cystic fibrosis showed no
8 clinically significant difference in the incidence of *P aeruginosa* infections between cohorting
9 patients into different pathogens by clinic times and no cohort segregation at 10 years follow-
10 up.

11 Prevalence of patients infected with transmissible pathogens

12 Very low quality evidence from 1 observational study with ≈2,837 sputum cultures showed no
13 significant difference in the prevalence of MRSA and non-mucoid *P aeruginosa* infections
14 among infants and children with cystic fibrosis after segregation measures (consisting in
15 segregating patients by age using a booking system) were put in place during the 4-year
16 follow-up. However, a significant lower prevalence of mucoid *P aeruginosa* was observed
17 during the same study period. The uncertainty for these outcomes could not be calculated.

18 Quality of life

19 No evidence was found.

20 Emotional function including anxiety and depression

21 No evidence was found.

22 Carer satisfaction

23 No evidence was found.

24 Patient satisfaction

25 No evidence was found.

- 1 **Staff experience**
- 2 No evidence was found.
- 3 **Staff and patient compliance**
- 4 Very low quality evidence from 1 observational study with staff looking after infants and
5 children with cystic fibrosis, showed that the staff adherence to the segregation programme
6 (using a clinic booking system) was over 90% during the 4-year follow-up period. This result
7 is provided narratively. The total number of people included in the study was not reported.
- 8 **11.6.1.2 Cohorting into different pathogens by location**
- 9 **11.6.1.2.1 Comparison 2. Cohort segregation by location versus no cohort segregation**
- 10 **Incidence of patients infected with transmissible pathogens**
- 11 Very low quality evidence from 1 observational study with people with 232 people with cystic
12 fibrosis receiving paediatric care suggested that the annual incidence of new growths of *P*
13 *aeruginosa*, while fluctuating, showed no downward trend 9 years after segregation
14 measures (consisting of separate clinics) were put in place, compared to previous usual
15 care. This outcome was reported narratively only.
- 16 **Prevalence of patients infected with transmissible pathogens**
- 17 Very low quality evidence from 1 observational study with 2,769 patient months showed a
18 clinically significant lower yearly prevalence of chronic *P aeruginosa* infections among people
19 with cystic fibrosis receiving paediatric care 9 years after segregation measures (consisting
20 of separate clinics) were put in place, compared to previous usual care. However, a clinically
21 significant higher yearly prevalence of intermittent *P aeruginosa* infections was observed.
- 22 **Quality of life**
- 23 No evidence was found.
- 24 **Emotional function including anxiety and depression**
- 25 No evidence was found.
- 26 **Carer satisfaction**
- 27 No evidence was found.
- 28 **Patient satisfaction**
- 29 No evidence was found.
- 30 **Staff experience**
- 31 No evidence was found.
- 32 **Staff and patient compliance**
- 33 No evidence was found

- 1 **11.6.1.3 Individual segregation**
- 2 No studies have been identified.
- 3 **11.6.1.4 Protective equipment**
- 4 No studies have been identified.
- 5 **11.6.1.5 Combination of strategies**
- 6 **11.6.1.5.1 Comparison 3. Combination of protective equipment + individual segregation versus**
- 7 **incomplete protective equipment + incomplete individual segregation**
- 8 **Incidence of patients infected with transmissible pathogens**
- 9 No evidence was found.
- 10 **Prevalence of patients infected with transmissible pathogens**
- 11 Very low quality evidence from 1 observational study with people with cystic fibrosis receiving
- 12 paediatric care and aged 0 to 21 showed a significantly lower prevalence of *P aeruginosa*
- 13 and MRSA infections calculated as average prevalence for 4 month intervals over a 5 year
- 14 period after an infection prevention and control policy (consisting in a combination of
- 15 individual segregation and protective equipment) was put in place, compared to the previous
- 16 strategy (that consisted in a combination of incomplete protective equipment + incomplete
- 17 individual segregation). The uncertainty for these outcomes could not be calculated. The total
- 18 number of people included in the study was not reported.
- 19 **Quality of life**
- 20 No evidence was found.
- 21 **Emotional function including anxiety and depression**
- 22 No evidence was found.
- 23 **Carer satisfaction**
- 24 No evidence was found.
- 25 **Patient satisfaction**
- 26 No evidence was found.
- 27 **Staff experience**
- 28 No evidence was found.
- 29 **Staff and patient compliance**
- 30 No evidence was found.

1 **11.6.2 Inpatient care**

2 **11.6.2.1 Cohorting into pathogen by location**

3 **11.6.2.1.1 Comparison 4. Cohort segregation by location versus no cohort segregation**

4 **Incidence of patients infected with transmissible pathogens**

5 Very low quality evidence from 1 observational study with people with cystic fibrosis showed
6 a lower annual incidence of *B cepacia complex* infection over 1 year after infection control
7 measures (consisting in cohorting hospitalised patients on the basis of *B cepacia*
8 colonisation status) were put in place, compared to the previous situation (no cohorting
9 segregation). The significance and the uncertainty of this result could not be calculated. The
10 age range and the total number of people in the study was not reported.

11 Very low quality evidence from 1 observational study with people with cystic fibrosis receiving
12 paediatric care showed a clinically significant lower incidence of hospital-associated
13 colonisation of *P cepacia* over 5 months after precautionary measures were implemented
14 (consisting in admitting all patients with *P cepacia* to the same ward), compared to the
15 previous situation (where only basic infection prevention measures were in place). The total
16 number of people in the study was not reported

17 **Prevalence of patients infected with transmissible pathogens**

18 No evidence was found.

19 **Quality of life**

20 No evidence was found.

21 **Emotional function including anxiety and depression**

22 No evidence was found.

23 **Carer satisfaction**

24 No evidence was found.

25 **Patient satisfaction**

26 No evidence was found.

27 **Staff experience**

28 No evidence was found.

29 **Staff and patient compliance**

30 No evidence was found.

31 **11.6.2.1.2 Comparison 5. Individual segregation by location versus usual care.**

32 **Incidence of patients infected with transmissible pathogens**

33 No evidence was found.

- 1 **Prevalence of patients infected with transmissible pathogens**
- 2 No evidence was found.
- 3 **Quality of life**
- 4 No evidence was found.
- 5 **Emotional function including anxiety and depression**
- 6 No evidence was found.
- 7 **Patient and carer satisfaction**
- 8 Very low quality from a cross-sectional study with 101 children and young people with cystic
9 fibrosis and their parents suggested that a high percentage of children and parents support
10 having segregated treatment (92% and 91% respectively).
- 11 **Staff experience**
- 12 No evidence was found.
- 13 **Staff and patient compliance**
- 14 No evidence was found.
- 15 **11.6.2.2 Inpatient room with ensuite facilities**
- 16 No studies have been identified.
- 17 **11.6.2.3 Inpatient recreational facilities**
- 18 No studies have been identified.
- 19 **11.6.3 Combined inpatient and outpatient care**
- 20 **11.6.3.1 Cohorting into pathogens**
- 21 **11.6.3.1.1 Comparison 6. Cohort segregation versus no cohort segregation**
- 22 **Incidence of patients infected with transmissible pathogens**
- 23 Very low quality evidence from 1 observational study with 119 people with cystic fibrosis
24 showed a clinically significant lower monthly incidence of multiply resistant *P aeruginosa*
25 strain 1 month after control measures (consisting in segregating patients based on PA
26 strains), compared to the previous policy (where patients were separated on the basis of the
27 PA status). The age of the people included in the study was not reported.
- 28 Very low quality evidence from 1 observational study with people with cystic fibrosis showed
29 no clinically significant difference in the annual incidence of intermittent and chronic *P*
30 *aeruginosa* 1 year after cohort isolation was introduced, compared to previous usual care.
31 The age range and the total number of people in the study was not reported.
- 32 Very low quality evidence from 1 observational study with 115 infants, children and young
33 people with cystic fibrosis showed no clinically significant difference in the incidence of *B*
34 *cepacia* during the 6-month study period after segregation measures were introduced,

1 compared to previous usual care. The total number of people included in the study was not
2 reported.

3 Very low evidence from 1 observational study with adults with cystic fibrosis showed a higher
4 annual incidence of *Burkholderia* species infections after changing infection control practices
5 (consisting in partially segregating patients with *Burkholderia* species infection), compared to
6 previous usual care. The significance and the uncertainty of this result could not be
7 calculated. The total number of people in the study was not reported.

8 **Prevalence of patients infected with transmissible pathogens**

9 Very low quality evidence from 1 observational study with 119 people with cystic fibrosis
10 showed no clinically significant difference in monthly prevalence of multiply resistant *P*
11 *aeruginosa* strain 1 month after control measures (consisting in segregating patients based
12 on PA strains), compared to the previous policy (where patients were separated on the basis
13 of the PA status). The age of the people included in the study was not reported.

14 Very low quality evidence from 1 observational study with people with cystic fibrosis
15 receiving paediatric care who were able to produce sputum showed a clinically significant
16 lower prevalence of *P aeruginosa* epidemic strain 2 years after cohort segregation measures
17 based on PA status were put in place, compare to previous usual care. The total number of
18 people included in the study was unclear.

19 Very low quality evidence from 1 observational study with adults with cystic fibrosis showed a
20 clinically significant higher annual prevalence of chronic *P aeruginosa* infection 1 year after
21 simple segregation measures were put in place, compared to previous usual care. However,
22 no clinically significant differences were found in the annual prevalence of transmissible *P*
23 *aeruginosa* infection. In addition, a slightly higher annual prevalence of chronic infection with
24 transmissible *P aeruginosa* strain was also observed, but the significance and the uncertainty
25 of this outcome could not be calculated. The total number of people included in the study
26 was unclear.

27 **Quality of life**

28 No evidence was found.

29 **Emotional function including anxiety and depression**

30 No evidence was found.

31 **Patient and carer satisfaction**

32 **No evidence was found.**

33 **Staff experience**

34 No evidence was found.

35 **Staff and patient compliance**

36 No evidence was found

11.6.3.1.2 Comparison 7. Complete cohort segregation versus incomplete cohort segregation

2 Incidence of patients infected with transmissible pathogens

3 Very low evidence from 1 observational study with adults with cystic fibrosis showed a lower
4 annual incidence of *Burkholderia* species infections after changing infection control practices
5 (consisting in cohorting patients with *Burkholderia* species infection), compared to the
6 previous situation (partial cohort segregation). The significance and the uncertainty of this
7 result could not be calculated. The follow-up and the total number of people included in the
8 study were unclear.

9 Prevalence of patients infected with transmissible pathogens

10 Quality of life

11 No evidence was found.

12 Emotional function including anxiety and depression

13 No evidence was found.

14 Carer satisfaction

15 No evidence was found.

16 Patient satisfaction

17 No evidence was found.

18 Staff experience

19 No evidence was found.

20 Staff and patient compliance

21 No evidence was found.

22 11.6.3.1.3 Comparison 8. Individual segregation versus usual care

23 Incidence of patients infected with transmissible pathogens

24 No evidence was found.

25 Prevalence of patients infected with transmissible pathogens

26 No evidence was found.

27 Quality of life

28 No evidence was found.

29 Emotional function including anxiety and depression

30 No evidence was found.

- 1 **Carer satisfaction**
- 2 No evidence was found.
- 3 **Patient satisfaction**
- 4 Very low quality evidence from 1 observational study with 94 adults with cystic fibrosis
5 indicated that the majority of the people (62.5%) who did not mix with others with cystic
6 fibrosis felt that their quality of life did not suffer as a result of this prevention control strategy;
7 whereas almost a quarter of the people (23.3%) who mixed with others with cystic fibrosis
8 were concerned that their quality of life would suffer a 'significant amount' or 'a great deal' if
9 they were to begin avoiding others. These results are provided narratively only.
- 10 **Staff experience**
- 11 No evidence was found.
- 12 **Staff and patient compliance**
- 13 No evidence was found.
- 14 **11.6.3.2 Protective equipment**
- 15 No studies have been identified.
- 16 **11.6.3.3 Combination of strategies**
- 17 **11.6.3.3.1 Comparison 9. Cohort segregation + individual segregation versus cohort segregation**
- 18 **Incidence of patients infected with transmissible pathogens**
- 19 No evidence was found.
- 20 **Prevalence of patients infected with transmissible pathogens**
- 21 Very low quality evidence from 1 observational study with people with cystic fibrosis showed
22 a lower annual prevalence of *B cepacia complex* infection 3 years after infection control
23 measures (consisting in cohorting hospitalised patients on the basis of *B cepacia*
24 colonisation status and individual isolation) were put in place, compared to the previous
25 situation (cohort segregation on the basis of *B cepacia* colonisation status only). The
26 significance and the uncertainty of this result could not be calculated. The age and the total
27 number of people included in the study was unclear.
- 28 Very low evidence from 1 observational study with adults with cystic fibrosis showed a lower
29 annual prevalence of *Burkholderia* species infections 5 years after changing infection control
30 practices (consisting in isolating patients with *Burkholderia* species infection in addition to
31 cohort segregation), compared to the previous situation (cohorting patients with *Burkholderia*
32 species infection). The significance and the uncertainty of this result could not be calculated.
33 The total number of people included in the study was unclear.
- 34 **Quality of life**
- 35 No evidence was found.
- 36 **Emotional function including anxiety and depression**
- 37 No evidence was found.

- 1 **Carer satisfaction**
- 2 No evidence was found.
- 3 **Patient satisfaction**
- 4 No evidence was found.
- 5 **Staff experience**
- 6 No evidence was found.
- 7 **Staff and patient compliance**
- 8 No evidence was found.
- 9 **1.6.3.3.2** **Comparison 10. Cohort segregation + individual segregation + protective equipment**
10 **versus usual care**
- 11 **Incidence of patients infected with transmissible pathogens**
- 12 Very low quality evidence from 1 observational study with people with cystic fibrosis showed
13 a lower annual incidence of *B cepacia complex* infection after a combination of infection
14 control measures (consisting in cohorting hospitalised patients on the basis of *B cepacia*
15 colonisation status, in addition to isolation and use of protective equipment) were put in
16 place, compared to the previous situation (usual care). The significance and the uncertainty
17 of this result could not be calculated. The follow-up, age and total number of people included
18 in the study were not reported.
- 19 **Prevalence of patients infected with transmissible pathogens**
- 20 No evidence was found.
- 21 **Quality of life**
- 22 No evidence was found.
- 23 **Emotional function including anxiety and depression**
- 24 No evidence was found.
- 25 **Carer satisfaction**
- 26 No evidence was found.
- 27 **Patient satisfaction**
- 28 No evidence was found.
- 29 **Staff experience**
- 30 No evidence was found.
- 31 **Staff and patient compliance**
- 32 No evidence was found.

11.6.3.3.3 Comparison 11. Cohort segregation + individual segregation versus usual care

2 Incidence of patients infected with transmissible pathogens

3 No evidence was found

4 Prevalence of patients infected with transmissible pathogens

5 No evidence was found.

6 Quality of life

7 No evidence was found.

8 Emotional function including anxiety and depression

9 No evidence was found.

10 Patient and carer satisfaction

11 Very low quality evidence from 190 people with cystic fibrosis receiving paediatric care or
12 their parents or carers indicated that both patients and carers showed an overall positive
13 response to segregation measures. This results are provided narratively only.

14 Staff experience

15 No evidence was found.

16 Staff and patient compliance

17 No evidence was found.

18 11.6.4 Economic evidence statements

19 No published evidence on cost-effectiveness in people with cystic fibrosis was available for
20 this review.

21 The economic model found that cohort segregation according to chronic *P.aeruginosa* was
22 cost-effective compared to no cohort segregation (3 studies).

23 The economic model found that cohort segregation according to intermittent *P.aeruginosa*
24 was not cost-effective compared to no cohort segregation (2 studies).

25 The economic model found that cohort segregation according to intermittent *B.cepacia*
26 complex was cost-effective compared to no cohort segregation in 1 study and cost-ineffective
27 in 1 study.

28 The economic model found that the addition of protective equipment was cost-effective to
29 prevent cross-infection with intermittent *B.cepacia* complex (1 study) and intermittent
30 *P.aeruginosa* (1 study).

31 The economic model found that individual inpatient segregation (single inpatient rooms)
32 according to intermittent *B.cepacia* complex would be cost-effective compared to no
33 individual inpatient segregation (shared ward) (2 studies).

1 11.7 Evidence to recommendations

2 11.7.1 Relative value placed on the outcomes considered

3 The aim of this review was to determine the effectiveness of the different strategies (such as
4 cohorting, segregation, or protective equipment) in reducing the transmission of cystic
5 fibrosis pathogens.

6 The Committee identified incidence and prevalence of patients infected with transmissible
7 pathogens as critical outcomes for decision making, whereas quality of life, emotional
8 function, patient satisfaction, staff experience and staff and patient compliance were rated as
9 important outcomes.

10 11.7.2 Consideration of clinical benefits and harms

11 The Committee noted that there were existing NICE guidelines on preventing and controlling
12 infection, therefore they incorporated a reference to these guidelines and focussed their
13 recommendations on cross-infection concerns specific to people with cystic fibrosis.

14 The Committee noted that the evidence showed mixed results with regards to whether
15 incidence and prevalence of people infected with transmissible pathogens decreased after
16 cohort segregation was implemented across the inpatient and outpatient setting or in the
17 outpatient setting alone. There was some evidence that the incidence of *B cepacia* complex
18 and of hospital-associated *P cepacia* infections decreased after cohort segregation was
19 implemented in the inpatient setting alone. Moreover, there was some evidence that
20 prevalence of *B cepacia* complex infection decreased after individual segregation was added
21 to cohort segregation across the inpatient and outpatient setting. The Committee noted that
22 feelings of isolation may be a potential disadvantage of individual or cohort segregation,
23 however there was some evidence that people with cystic fibrosis and carers supported both
24 cohort and individual segregation.

25 The Committee noted that infection control would have to be implemented both in the
26 inpatient and outpatient setting at the same time, therefore the Committee recommended
27 that arrangements to prevent cross-infection among people with cystic fibrosis should be
28 based on a local infection control strategy that covers both outpatient and inpatient care. This
29 strategy would cover interventions such as: cleaning of rooms and equipment; closing the
30 doors of rooms in the hospital or the outpatient clinic; ventilation in gyms to ensure that the
31 next person does not breathe the same air as the previous person. Moreover, the Committee
32 recommended that arrangements to prevent cross-infection should also be based on a
33 microbiological surveillance programme.

34 The Committee noted that segregation interventions would be ineffective if people did not
35 adhere to segregation rules or if they met with each other outside of the health care setting.
36 Therefore a recommendation was prioritised to inform people with cystic fibrosis, their family
37 members or carers and staff involved in their care about the risk of cross-infection and how
38 to avoid it.

39 The Committee noted that individual segregation is likely to be more effective than cohort
40 segregation because cross-infection can occur between people with cystic fibrosis who are
41 designated as having the same pathogen but actually have different strains which may have
42 differing virulence characteristics. For example, superinfection with different, more virulent
43 epidemic strains of *P aeruginosa*, potentially as a result of cross-infection, can occur in
44 people who already have *P aeruginosa*; similarly, patients with less virulent species of *B*
45 *cepacia* complex, for example *Burkholderia multivorans*, could become infected with the
46 more virulent *Burkholderia cenocepacia* if cohorting is based on *B cepacia* complex status
47 without species differentiation or if patients have as yet undetected infection. If people were
48 to be cohorted based on different pathogen strains, then cohort interventions would become

1 complex due to a relatively high number of different cohorts which may increase as
2 understanding of individual pathogen virulence develops. Moreover, any information on a
3 person's infection status refers to the last sputum culture that was performed. Therefore,
4 after acquisition of an infection there would be a time interval until the next sputum culture,
5 during which a person would be cohorted together with people who are free of infection.
6 Therefore, the Committee concluded that separating all people with cystic fibrosis
7 (regardless of their infection status) from each other was bound to be more effective as
8 opposed to separating them into groups based on infection status. The Committee also
9 noted that the inpatient setting is the setting where people would be more likely to come into
10 close contact with each other because they may spend multiple days or weeks in the
11 hospital. Therefore, the Committee prioritised a recommendation to give people with cystic
12 fibrosis individual rooms with en-suite facilities. This recommendation was consistent with the
13 NHS service specifications for cystic fibrosis. The Committee noted that in practice people
14 with cystic fibrosis are sometimes segregated into individual rooms but have contact in
15 communal areas, which is to be avoided. Therefore, the Committee recommended to help
16 inpatients with cystic fibrosis plan their attendance to avoid contact with each other, for
17 example when they use hospital restaurants, schools and recreation areas, such as the gym;
18 and diagnostic and treatment facilities. The Committee noted that a timetable could be used
19 for school and gym attendance. Moreover, for diagnostic and treatment procedures there
20 could be a computerised system suggesting who should enter specific areas at what time.
21 Finally, systems should be in place to ensure that if someone with cystic fibrosis comes to
22 the hospital, contact with the inpatients is prevented.

23 The Committee noted that when it comes to the outpatient setting, contact between people
24 with cystic fibrosis is mostly casual, as the chances of close contact are considerably lower
25 compared to the inpatient setting. Moreover, in the outpatient setting it would be more
26 difficult to ensure individual segregation because the whole patient journey would need to be
27 taken into account and because it is more common that people attend for the use of
28 diagnostic or treatment facilities without a prior booking. The Committee also noted the lack
29 of evidence assessing individual segregation as an intervention on its own in the outpatient
30 setting alone. Considering these issues on the one hand and the aforementioned advantages
31 of individual segregation on the other hand, the Committee concluded that in the outpatient
32 setting the decision on whether to segregate patients into cohorts or individually should be
33 left to the individual clinics, therefore it did not specify how people should be separated, and
34 it only recommended to separate people, for example by managing communal waiting areas
35 and attendance at diagnostic and treatment facilities. With regards to these points, the
36 Committee noted that there were multiple options to separate people; for example, placing
37 the person directly into a room, where the multidisciplinary team can go to conduct the visit;
38 or using a communal waiting area for people with a specific pathogen on 1 day of the week
39 and for people with a different pathogen on a different day of the week.

40 Although it did not specify whether to implement individual or cohort segregation in the
41 outpatient setting, the Committee agreed to prioritise a recommendation to separate people
42 with transmissible or chronic *P aeruginosa* or *B cepacia* complex infection from people who
43 do not have these infections during outpatient clinics.

44 The Committee noted that it would be difficult to separate people with intermittent *P*
45 *aeruginosa* infection from those who do not have this infection because intermittent infection
46 is harder to diagnose given that tests with positive results would alternate with tests with
47 negative results. Moreover some evidence showed a higher prevalence of intermittent *P*
48 *aeruginosa* infection after cohort segregation in the outpatient setting; however a lower
49 prevalence of chronic *P aeruginosa* infection was observed in the same setting so these
50 trends may be due to treatment strategies rather than to cohort segregation (because
51 treatment may prevent the intermittent pathogen from becoming chronic). With regards to
52 this point, the Committee agreed that changes in prevalence and incidence may be due not
53 only to cross-infection but also to other factors, such as acquisition of infection from
54 environmental sources and advances in treatment; moreover, prevalence also relates to

1 survival. The evidence also showed no clinically significant difference in the incidence of
2 intermittent *P aeruginosa* after cohort segregation across the inpatient and outpatient setting.
3 Given the lack of evidence supporting segregation and considering the difficulties involved in
4 diagnosing intermittent *P aeruginosa*, the Committee decided not to be too prescriptive and
5 recommended to consider separating people with intermittent *P aeruginosa* from people who
6 do not have this infection during outpatient clinics.

7 There was some evidence that prevalence of people infected with transmissible pathogens
8 decreased after a combination of individual segregation with protective equipment was
9 implemented in the outpatient setting alone and incidence of infection decreased after a
10 combination of cohort and individual segregation and protective equipment was implemented
11 across the inpatient and outpatient settings. However there was no evidence on the use of
12 protective equipment alone. The Committee noted that the use of protective equipment is
13 difficult to implement, as people's motivation is limited because of stigma and because
14 protective equipment protects other people but not themselves, therefore they tend to take it
15 off. Therefore the Committee decided not to make a recommendation on protective
16 equipment.

17 **11.7.3 Consideration of economic benefits and harms**

18 **11.7.3.1 Cohort segregation**

19 ***P aeruginosa***

20 The economic model showed that cohort segregation was cost-effective to reduce the
21 transmission of chronic *P aeruginosa* compared to no cohort segregation, but was not cost-
22 effective to reduce the transmission of intermittent *P aeruginosa*. This difference was driven
23 by Lee 2004 who found that a fall in the number of cases with chronic infection was
24 associated with a rise in those classified as intermittent. Following this, the Committee noted
25 that it is difficult to define an intermittent infection, adding that the distinction is even more
26 difficult in children who cannot produce sputum.

27 Based on the evidence presented to them and their clinical expertise, the Committee agreed
28 they could justify a recommendation to cohort people with transmissible *P aeruginosa* and
29 chronic *P aeruginosa* as most clinics implement this as a cost-effective measure.

30 The Committee also wanted to enable clinics that had the ability to differentiate pathogens
31 for intermittent *P aeruginosa* to do so, although they agreed that the findings from Lee 2004
32 were limited. This was because Lee 2004 did not genotype the participants' *P aeruginosa*
33 isolates, to infer if their strains were transmissible and potentially acquired from the clinic.
34 Given that the findings from Lee 2004 can produce an unreliable estimate of cost-
35 effectiveness and cohort segregation depends on the ability of the clinic to identify
36 intermittent *P.aeruginosa*, the Committee reduced the strength of their recommendation to
37 "consider cohorting people with cystic fibrosis who have intermittent pseudomonas
38 aeruginosa" to reflect the weaker evidence compared to chronic *P.aeruginosa*.

39 The Committee discussed the potential harms of seeing less people with cystic fibrosis using
40 cohorted clinics and advised that waiting times could increase in the short term, but in the
41 longer term, the number of people with a transmissible pathogen would decrease,
42 subsequently increasing the number of people that could be seen during "usual" clinics. The
43 Committee also referred to their recommendations on service delivery, where people with
44 cystic fibrosis and their family members or carers can contact the MDT when they have
45 urgent enquiries at any time, to minimise delays in identification and management.

46 The Committee stated that genotyping is currently used by cystic fibrosis clinics to keep a
47 track on the number of epidemic strains and prompt further investigation when there is a
48 significant increase in numbers. Cystic fibrosis clinics currently follow CF Trust laboratory

standards and infection control guidelines that advise surveillance should be performed on all new isolates, and annually on those from people infected with *P aeruginosa*. Therefore, genotyping all isolates of *P aeruginosa* in order to provide more up to date information for accurate clinic segregation, as a means to reduce cross-infections with transmissible pathogens could not be achieved with existing laboratory resources. The Committee advised that it would be relatively easy to set up in-house polymerase chain reaction (PCR) methods for detecting the presence or otherwise of known epidemic strains, but this would require dedicated project budget and recruitment of technical and scientific staff to deliver this work. Moreover, to achieve the required level of discriminatory power, more complex typing methodologies such as variable number of tandem repeats (VNTR) typing or whole genome sequencing (WGS) would be necessary, but would raise costs and workloads even further. Overall, the Committee considered a recommendation was needed to ensure microbiological surveillance and local infection control strategies were in place. The Committee were reluctant to specify how such arrangements should be achieved as this would depend on the expertise and prevalence of pathogens within each clinic.

In addition, a recommendation in favour of regular, detailed genotyping was not prioritised by the Committee as there is insufficient evidence to enable full understanding of the results, especially as there are data suggesting significant intra-patient (and intra-isolate) variability when detailed methods are used referring to the study by Darch 2015. For these reasons, the Committee wanted to make a research recommendation to define optimal microbiological methods and their role to inform infection control strategies, to assess if the benefits of regular genotyping could justify the additional resources. However, this was later deprioritised as it was considered a public health issue for local areas to devise their own policies.

***B cepacia* complex**

One study (Whitford 1995) found cohort segregation according to *B cepacia* complex to be cost-effective compared to no cohort segregation. Conversely, the second study (France 2008) found no cohort segregation to dominate cohort segregation according to *B cepacia* complex, as no cohort segregation was less expensive and more effective than cohort segregation. However, given that complete cohort segregation with the addition of single inpatient rooms, dominated (less expensive and more effective) incomplete cohort segregation, the Committee agreed that people infected with *B cepacia* complex also should be at outpatient clinics and inpatient care. Following this, the Committee noted that the species of *B cepacia* complex must be determined otherwise those with *cenocercaria* could be cohorted with *multivorans* which would be inappropriate and hazardous. The Committee also noted that the prevalence of *B cepacia* complex is decreasing (circa 3%) according to UK registry data, indicating that current infection control measures to cohort these people with cystic fibrosis are working.

11.7.3.2 Individual segregation

The Committee advised that unlike cohort segregation, individual segregation acts as a preventative measure as there is less need to worry about the specific pathogen. The Committee also added that individual segregation is a strategy currently applied in cystic fibrosis clinics, to reduce the risk of cross-infection with transmissible pathogens.

The Committee agreed that the additional cost of single rooms compared to beds in shared rooms was best reflected by NHS Estates 2005 (an additional £10.61/day in 2016 prices) and potentially overestimated when an additional 25% was assumed.

The clinical evidence was not meta-analysed as the studies were too heterogeneous. Consequently, the Committee considered the cost-effective results produced by each of the studies and concluded that France 2008 was the study most reflective of UK clinical practice today. Based on France 2008, the dominant strategy (less expensive and more effective

1 strategy) was to admit people with cystic fibrosis to single rooms when an additional
2 £10.61/bed/day or 25%/bed/day compared to beds in shared rooms was assumed. As a
3 result, the Committee prioritised a recommendation for all inpatients to have their own single
4 room as the economic model provided additional justifications for current practice to be
5 followed.

6 The Committee also considered the acceptable additional cost given the additional QALY
7 gain, based on the study by Jones 2005 who compared incomplete cohort segregation to no
8 cohort segregation. Incomplete cohort segregation provided inpatient rooms with en-suite
9 facilities, whereas only 2 of the 11 rooms in no cohort segregation strategy included en-suite
10 facilities. Incomplete cohort segregation also meant people without chronic *P aeruginosa*
11 attended a different outpatient clinic appointments to other people with cystic fibrosis. The
12 Committee acknowledged that the incidence of superinfection by transmissible strains
13 among trial participants (Jones 2005) already infected with chronic *P aeruginosa* was
14 sporadic, potentially resulting in a dominated (more expensive and less effective) strategy.

15 However, incomplete cohort segregation produced more QALYs than no cohort segregation,
16 with a fall in the incidence of intermittent *P aeruginosa* from 9.7% to 0% each year, following
17 the new strategy. Based on this benefit, the Committee believed that incomplete cohort
18 segregation could be cost-effective. The Committee also added that the majority of single
19 rooms include en-suite facilities and referred to the NHS Commissioning document on
20 service specifications that states “Every CF patient will be in their own room, with en-suite
21 facilities to minimise the risk of cross infection and to enable them to continue life as normally
22 as possible”. Given that the majority of centres follow this, the Committee agreed a
23 recommendation in favour of en-suite facilities that would not lead to substantial increase
24 from current resource use.

25 11.7.3.3 Protective equipment

26 The new strategies implemented by Savant 2014 and Chen 2001 that included protective
27 equipment to prevent intermittent infections were cost-effective (dominant). However, it is
28 important to note that the effect of protective equipment alone could not be disaggregated
29 into a separate effect measure from the combination of strategies applied. Following this, the
30 Committee agreed that a dominant result was unlikely to be driven by the protective
31 equipment as Chen 2001 also cohorted their participants according to *B cepacia* complex
32 infection and Savant 2014 applied a “no-waiting” room policy, which the Committee believed
33 to be the true cost-effective strategies.

34 Despite the low cost of protective equipment, the Committee agreed, based on their
35 knowledge and expertise, that there was not enough evidence to support the use of
36 protective equipment as compliance outside of a trial setting would be low, due to the
37 negative impact the equipment has on social interaction.

38 11.7.3.4 Other considerations

39 The Committee agreed that communal areas in cystic fibrosis clinics promote patient contact
40 and increase the risk of infection from transmissible pathogens, moreover the opportunity
41 cost of the space would be of greater value in other areas of the hospital. For these reasons,
42 the Committee made a recommendation to manage the use of communal waiting areas in
43 outpatient clinics. The Committee also considered the benefits from closing clinic room doors
44 as this is a very simple, quick, and costless way to prevent transmissions that can be
45 overlooked, or stopped by reluctant parents; however, a recommendation was not prioritised.

46 11.7.4 Quality of evidence

47 All the evidence included in the review was of very low quality as assessed by GRADE.

1 One RCT was included. This study had unclear risk of selection bias, performance bias,
2 detection bias, attrition bias, and outcome reporting bias.

3 Most included studies were retrospective uncontrolled before-after studies. In some of these
4 studies, the intervention and comparison groups were drawn from years or time intervals that
5 were considerably distant in time, thus increasing the likelihood that the intervention group
6 may have been exposed to additional interventions affecting incidence or prevalence.
7 Moreover, in some studies the frequency of cultures was unclear or only annual. Therefore,
8 people that were supposedly “at risk” when the intervention started may have actually had
9 the infection. Consequently, there was a high risk of selection bias. Most studies did not
10 control for any factor, therefore there was also a high risk of comparability bias. Furthermore,
11 many studies looked at changes in prevalence and incidence without making a distinction
12 between transmissible strains and unique strains; most studies did not use genotyping in
13 order to understand whether changes in incidence or prevalence were related to cross-
14 infection. Therefore, there was high risk of bias in relation to the outcomes of incidence and
15 prevalence.

16 Three surveys were included. Only 1 study assessed comparability between respondents
17 and non-respondents and all three studies had high risk of selection bias. None of the three
18 studies controlled the analysis for possible confounders, therefore there was high risk of
19 comparability bias.

20 **11.7.5 Other considerations**

21 There was some evidence that staff adherence to cohort segregation was high. This
22 indicated that cohort segregation was feasible.

23 No equality issues were identified by the Committee for this review question.

24 The Committee discussed the need for a research recommendation in this area, but they
25 agreed it was not needed, as they noted it is more for local areas to devise their own policies
26 and public health issue. They also acknowledged there is a NICE guideline on infection.

27 **11.7.6 Key conclusions**

28 The Committee concluded that in the inpatient setting people with cystic fibrosis should avoid
29 contact with each other irrespective of their infection status. However in the outpatient
30 setting, where the chances of close contact are lower, individual clinics should decide
31 whether to implement cohort segregation or individual segregation. The minimum
32 requirement in outpatient clinics is that people with transmissible or chronic *P aeruginosa* or
33 *B cepacia* complex infection are kept separate from people who do not have these infections.

34 **11.8 Recommendations**

35 **132. For recommendations on preventing and controlling infection, see the NICE**
36 **guidelines on infection control in primary and community care, healthcare-**
37 **associated infections and the NICE quality standard on infection prevention and**
38 **control.**

39 **133. To prevent cross-infection among people with cystic fibrosis, use**
40 **microbiological surveillance and a local infection control strategy that covers**
41 **outpatient and inpatient care.**

42 **134. Inform people with cystic fibrosis, their family members or carers (as**
43 **appropriate) and staff involved in their care about the risk of cross-infection and**
44 **how to avoid it.**

- 1 **135. Separate people with cystic fibrosis who have transmissible or chronic**
2 **pseudomonas aeruginosa or Burkholderia cepacia complex infection, for example**
3 **during outpatient clinics.**
- 4 **136. Consider separating people with cystic fibrosis who have intermittent**
5 **pseudomonas aeruginosa, for example during outpatient clinics.**
- 6 **137. All specialist cystic fibrosis clinics should be organised to prevent cross-**
7 **infection. Separate people at outpatient clinics, for example by managing:**
8 • use of communal waiting areas
9 • attendance at diagnostic and treatment facilities.
- 10 **138. Help people with cystic fibrosis plan their inpatient attendance to avoid contact**
11 **with each other, for example when they use:**
12 • hospital restaurants, schools and recreation areas
13 • diagnostic and treatment facilities.
- 14 **139. During inpatient care, give people with cystic fibrosis individual rooms with en-**
15 **suite facilities.**
- 16
17
18

12 References

- 1 **Aaron et al., 2012**
- 2
- 3 Aaron, S. D., Vandemheen, K. L., Freitag, A., Pedder, L., Cameron, W., Lavoie, A., Paterson,
4 N., Wilcox, P., Rabin, H., Tullis, E., Morrison, N., Ratjen, F., Treatment of *Aspergillus*
5 *fumigatus* in patients with cystic fibrosis: a randomized, placebo-controlled pilot study, *PLoS*
6 *ONE*, 7, e36077, 2012
- 7 **Ahmed & Mukherjee, 2016**
- 8 Ahmed, M. I., Mukherjee, S., Treatment for chronic methicillin-sensitive *S aureus* pulmonary
9 infection in people with cystic fibrosis, *Cochrane Database of Systematic Reviews*, 3,
10 CD011581, 2016
- 11 **Aitken et al., 2012**
- 12 Aitken, M. L., Bellon, G., De Boeck, K., Flume, P. A., Fox, H. G., Geller, D. E., Haarman, E.
13 G., Hebestreit, H. U., Lapey, A., Schou, I. M., Zuckerman, J. B., Charlton, B., C. F.
14 Investigators, Long-term inhaled dry powder mannitol in cystic fibrosis: an international
15 randomized study, *American Journal of Respiratory & Critical Care Medicine*, 185, 645-52,
16 2012
- 17 **Al-Yateem, 2012**
- 18 Al-Yateem, N., Child to adult: transitional care for young adults with cystic fibrosis, *British*
19 *Journal of Nursing*, 21, 850-4, 2012
- 20 **Al-Yateem, 2013**
- 21 Al-Yateem, N., Guidelines for the transition from child to adult cystic fibrosis care, *Nursing*
22 *Children and Young People*, 25, 29-34, 2013
- 23 **Amin et al., 2010**
- 24 Amin, R., Subbarao, P., Jabar, A., Balkovec, S., Jensen, R., Kerrigan, S., Gustafsson, P.,
25 Ratjen, F., Hypertonic saline improves the LCI in paediatric patients with CF with normal lung
26 function, *Thorax*, 65, 379-83, 2010
- 27 **Amin et al., 2011**
- 28 Amin, R., Subbarao, P., Lou, W., Jabar, A., Balkovec, S., Jensen, R., Kerrigan, S.,
29 Gustafsson, P., Ratjen, F., The effect of dornase alfa on ventilation inhomogeneity in patients
30 with cystic fibrosis, *European Respiratory Journal*, 37, 806-812, 2011
- 31 **Angst & Deatrck, 1996**
- 32 Angst, D. B., Deatrck, J. A., Involvement in health care decisions: parents and children with
33 chronic illness, *Journal of Family Nursing*, 2, 174-194, 1996
- 34 **Anyanwu et al., 2001**
- 35 Anyanwu, A., McGuire, A., Rogers, C., Murday, A., Assessment of quality of life in lung
36 transplantation using a simple generic tool, *Thorax*, 56, 218-222, 2001
- 37 **Anyanwu et al., 2002**
- 38 Anyanwu, A. C., McGuire, A., Rogers, C. A., Murday, A. J., An economic evaluation of lung
39 transplantation, *Journal of Thoracic and Cardiovascular Surgery*, 123, 411-8; discussion 418-
40 20, 2002

- 1 Assael et al., 2013
- 2 Assael, B. M., Pressler, T., Bilton, D., Fayon, M., Fischer, R., Chiron, R., LaRosa, M., Knoop,
3 C., McElvaney, N., Lewis, S. A., Bresnik, M., Montgomery, A. B., Oermann, C. M., Azli,
4 Active Comparator Study Group, Inhaled aztreonam lysine vs. inhaled tobramycin in cystic
5 fibrosis: a comparative efficacy trial, *Journal of Cystic Fibrosis*, 12, 130-40, 2013
- 6 Auerbach et al., 1985
- 7 Auerbach, H. S., Williams, M., Kirkpatrick, J. A., Colten, H. R., Alternate-day prednisone
8 reduces morbidity and improves pulmonary function in cystic fibrosis, *Lancet*, 2, 686-8, 1985
- 9 Bagnasco et al., 2013
- 10 Bagnasco, A., Petralia, P., Furnari, S., Ghio, S., Calza, S., Sasso, L., Paediatric nurses'
11 perception of the child-family dyad's autonomy in managing a chronic disease situation: the
12 experience of an Italian paediatric department, *Journal of Preventive Medicine & Hygiene*,
13 54, 124-9, 2013
- 14 Baker et al., 2016
- 15 Baker, J. F., Putman, M. S., Herlyn, K., Tillotson, A. P., Finkelstein, J. S., Merkel, P. A., Body
16 composition, lung function, and prevalent and progressive bone deficits among adults with
17 cystic fibrosis, *Joint, Bone, Spine: Revue du Rhumatisme*, 83, 207-11, 2016
- 18 Balaguer & Gonzalez de Dios, 2015
- 19 Balaguer, Albert, Gonzalez de Dios, Javier, Home versus hospital intravenous antibiotic
20 therapy for cystic fibrosis, *Cochrane Database of Systematic Reviews*, 12, CD00191,7 2015
- 21 Balfour-Lynn & Welch, 2016
- 22 Balfour-Lynn, I. M., Welch, Karen, Inhaled corticosteroids for cystic fibrosis, *Cochrane*
23 *Database of Systematic Reviews*, 8, CD001915, 2016
- 24 **Balfour-Lynn et al., 1997**
- 25 Balfour-Lynn, I. M., Klein, N. J., Dinwiddie, R., Randomised controlled trial of inhaled
26 corticosteroids (fluticasone propionate) in cystic fibrosis, *Archives of Disease in Childhood*,
27 77, 124-30, 1997
- 28 Balfour-Lynn et al., 2006
- 29 Balfour-Lynn, I. M., Lees, B., Hall, P., Phillips, G., Khan, M., Flather, M., Elborn, J. S., Cf
30 Wise Investigators, Multicenter randomized controlled trial of withdrawal of inhaled
31 corticosteroids in cystic fibrosis, *American Journal of Respiratory & Critical Care Medicine*,
32 173, 1356-62, 2006
- 33 **Ballmann & vonderHardt, 1998**
- 34 Ballmann, M., von der Hardt, H., Hypertonic saline and recombinant human DNase: a
35 randomised cross-over pilot study in patients with cystic fibrosis, 22nd European Cystic
36 Fibrosis Conference, 1998, [Abstract]
- 37 **Ballmann & von der Hardt, 2002**
- 38 Ballmann, M., von der Hardt, H., Hypertonic saline and recombinant human DNase: a
39 randomised cross-over pilot study in patients with cystic fibrosis, *Journal of Cystic Fibrosis*, 1,
40 35-7, 2002
- 41 **Barker et al., 2012**

- 1 Barker, D., Driscoll, K., Modi, A., Light, M., Quittner, A., Supporting cystic fibrosis disease
2 management during adolescence: The role of family and friends, *Child: care, health and*
3 *development*, 38, 497-504, 2012
- 4 **Beardsmore et al., 1994**
- 5 Beardsmore, C.S., Thompson, J.R., Williams, A., McArdle, E.K., Gregory, G.A., Weaver, L.T.,
6 Simpson, H., Pulmonary function in infants with cystic fibrosis: the effect of antibiotic
7 treatment, *Archives of Disease in Childhood*, 71, 133-137, 1994
- 8 **Beaudoin et al., 2016**
- 9 Beaudoin, N., Bouvet, G. F., Coriati, A., Rabasa-Lhoret, R., Berthiaume, Y., Combined
10 Exercise Training Improves Glycemic Control in Adult With Cystic Fibrosis, *Medicine and*
11 *Science in Sports and Exercise*, 2016
- 12 **Begley, 2013**
- 13 Begley, T., Transition to adult care for young people with long-term conditions, *British Journal*
14 *of Nursing*, 22, 506, 508-11, 2013
- 15 **Beker et al., 1994**
- 16 Beker, L. T., Fink, R. J., Shamsa, F. H., Chaney, H. R., Kluff, J., Evans, E., Schidlow, D. V.,
17 Comparison of weight-based dosages of enteric-coated microtablet enzyme preparations in
18 patients with cystic fibrosis, *Journal of Pediatric Gastroenterology & Nutrition*, 19, 191-7,
19 1994
- 20 **Bell et al., 2011**
- 21 Bell, S. C., Bye, P. T., Cooper, P. J., Martin, A. J., McKay, K. O., Robinson, P. J., Ryan, G.
22 F., Sims, G. C., Cystic fibrosis in Australia, 2009: results from a data registry, *Medical*
23 *Journal of Australia*, 195, 396-400, 2011
- 24 **Beresford & Sloper, 2003**
- 25 Beresford, B. A., Sloper, P., Chronically ill adolescents' experiences of communicating with
26 doctors: a qualitative study, *Journal of Adolescent Health*, 33, 172-9, 2003
- 27 **Bhudhikanok et al., 1998**
- 28 Bhudhikanok, G. S., Wang, M. C., Marcus, R., Harkins, A., Moss, R. B., Bachrach, L. K.,
29 Bone acquisition and loss in children and adults with cystic fibrosis: a longitudinal study,
30 *Journal of Pediatrics*, 133, 18-27, 1998
- 31 **Bilton et al., 2011**
- 32 Bilton, D., Robinson, P., Cooper, P., Gallagher, C. G., Kolbe, J., Fox, H., Jaques, A.,
33 Charlton, B., C. F. Study Investigators, Inhaled dry powder mannitol in cystic fibrosis: an
34 efficacy and safety study, *European Respiratory Journal*, 38, 1071-80, 2011
- 35 **Blumer et al., 2005**
- 36 Blumer, J. L., Saiman, L., Konstan, M. W., Melnick, D., The efficacy and safety of
37 meropenem and tobramycin vs ceftazidime and tobramycin in the treatment of acute
38 pulmonary exacerbations in patients with cystic fibrosis, *Chest*, 128, 2336-46, 2005
- 39 **Bond et al., 2007**
- 40 Bond, Mary, Mealing, Stuart, Anderson, Rob, Elston, Julian, Weiner, Graeme, Taylor, Rod
41 S., Hoyle, Martin, Liu, Zulian, Price, Alison, Stein, Ken, Peninsula Technology Assessment
42 Group (PenTAG), Peninsula Medical School, Universities of Exeter and Plymouth, The

- 1 Effectiveness and Cost-Effectiveness of Cochlear Implants for Severe to Profound Deafness
2 in Children and Adults: a Systematic Review and Economic Model, 2007, Plymouth:
3 Peninsula Technology Assessment Group (PenTAG)
- 4 **Bosworth & Nielson, 1997**
- 5 Bosworth, D. G., Nielson, D. W., Effectiveness of home versus hospital care in the routine
6 treatment of cystic fibrosis, *Pediatric Pulmonology*, 24, 42-7, 1997
- 7 **Bradley et al., 2010**
- 8 Bradley, J., Blume, S., Stafford, M., Balp, M., Elborn, S., Quality of life and health utility in
9 patients with cystic fibrosis, *European Respiratory Journal*, 36, 2652, 2010
- 10 **Bradley et al., 2012**
- 11 Bradley, G. M., Carson, K. A., Leonard, A. R., Mogayzel, P. J., Jr., Oliva-Hemker, M.,
12 Nutritional outcomes following gastrostomy in children with cystic fibrosis, *Pediatric*
13 *Pulmonology*, 47, 743-8, 2012
- 14 **Brady et al., 1991**
- 15 Brady, M. S., Rickard, K., Yu, P. L., Eigen, H., Effectiveness and safety of small vs. large
16 doses of enteric coated pancreatic enzymes in reducing steatorrhea in children with cystic
17 fibrosis: a prospective randomized study, *Pediatric Pulmonology*, 10, 79-85, 1991
- 18 **Braggion et al., 1995**
- 19 Braggion, C., Cappelletti, L. M., Cornacchia, M., Zanolla, L., Mastella, G., Short-term effects
20 of three chest physiotherapy regimens in patients hospitalized for pulmonary exacerbations
21 of cystic fibrosis: a cross-over randomized study, *Pediatric Pulmonology*, 19, 16-22, 1995
- 22 **Braithwaite et al., 2011**
- 23 Braithwaite, M., Philip, J., Tranberg, H., Finlayson, F., Gold, M., Kotsimbos, T., Wilson, J.,
24 End of life care in CF: patients, families and staff experiences and unmet needs, *Journal of*
25 *Cystic Fibrosis*, 10, 253-7, 2011
- 26 **Brenckmann et al., 2003**
- 27 Brenckmann, C., Papaioannou, A., Freitag, A., Hennessey, R., Hansen, S., Ioannidis, G.,
28 Webber, C., Adachi, J., Osteoporosis in Canadian adult cystic fibrosis patients: a descriptive
29 study, *BMC Musculoskeletal Disorders*, 4, 13, 2003
- 30 **Briesacher et al., 2011**
- 31 Briesacher, B. A., Quittner, A. L., Fouayzi, H., Zhang, J., Swensen, A., Nationwide trends in
32 the medical care costs of privately insured patients with cystic fibrosis (CF), 2001-2007,
33 *Pediatric Pulmonology*, 46, 770-6, 2011
- 34 **Brumfield & Lansbury, 2004**
- 35 Brumfield, K., Lansbury, G., Experiences of adolescents with cystic fibrosis during their
36 transition from paediatric to adult health care: a qualitative study of young Australian adults,
37 *Disability & Rehabilitation*, 26, 223-34, 2004
- 38 **Chatfield et al., 1991**
- 39 Chatfield, S., Owen, G., Ryley, H. C., Williams, J., Alfaham, M., Goodchild, M. C., Weller, P.,
40 Neonatal screening for cystic fibrosis in Wales and the West Midlands: clinical assessment
41 after five years of screening, *Archives of Disease in Childhood*, 66, 29-33, 1991

- 1 **Chavasse et al., 2004**
- 2 Chavasse, R. J., Francis, J., Balfour-Lynn, I., Rosenthal, M., Bush, A., Serum vitamin D
3 levels in children with cystic fibrosis, *Pediatric Pulmonology*, 38, 119-22, 2004
- 4 **Chen et al., 2001**
- 5 Chen, J. S., Witzmann, K. A., Spilker, T., Fink, R. J., LiPuma, J. J., Endemicity and inter-city
6 spread of *B cepacia* genomovar III in cystic fibrosis, *Journal of Pediatrics*, 139, 643-9, 2001
- 7 **Cheng et al., 2014**
- 8 Cheng, Katharine, Ashby, Deborah, Smyth, Rosalind L., Ursodeoxycholic acid for cystic
9 fibrosis-related liver disease, *Cochrane Database of Systematic Reviews*, 2014
- 10 **Cheng et al., 2015**
- 11 Cheng, K., Ashby, D., Smyth, R. L., Oral steroids for long-term use in cystic fibrosis,
12 *Cochrane Database of Systematic Reviews*, 12, CD000407, 2015
- 13 **Chinuck et al., 2014**
- 14 Chinuck, R., Dewar, J., Baldwin, D. R., Hendron, E., Appetite stimulants for people with
15 cystic fibrosis, *Cochrane Database of Systematic Reviews*, 7, CD008190, 2014
- 16 **Christopher et al., 1999**
- 17 Christopher, F., Chase, D., Stein, K., Milne, R., rhDNase therapy for the treatment of cystic
18 fibrosis patients with mild to moderate lung disease, *Journal of Clinical Pharmacy &*
19 *Therapeutics*, 24, 415-26, 1999
- 20 **Chuchalin et al., 2007**
- 21 Chuchalin, A., Csiszer, E., Gyurkovics, K., Bartnicka, M. T., Sands, D., Kapranov, N., Varoli,
22 G., Monici Preti, P. A., Mazurek, H., A formulation of aerosolized tobramycin (Bramitob) in
23 the treatment of patients with cystic fibrosis and *P aeruginosa* infection: a double-blind,
24 placebo-controlled, multicenter study, *Paediatric Drugs*, 9 Suppl 1, 21-31, 2007
- 25 **Clement et al., 2006**
- 26 Clement, A., Tamalet, A., Leroux, E., Ravilly, S., Fauroux, B., Jais, J. P., Long term effects of
27 azithromycin in patients with cystic fibrosis: A double blind, placebo controlled trial, *Thorax*,
28 61, 895-902, 2006
- 29 **Coates et al., 2007**
- 30 Coates, Nicola, Gregory, Maggie, Skirton, Heather, Gaff, Clara, Patch, Christine, Clarke,
31 Angus, Parsons, Evelyn, Family communication about cystic fibrosis from the mother's
32 perspective: An exploratory study, *Journal of Research in Nursing*, 12, 619-634, 2007
- 33 **Colombo et al., 1996**
- 34 Colombo, C., Battezzati, P. M., Podda, M., Bettinardi, N., Giunta, A., Ursodeoxycholic acid
35 for liver disease associated with cystic fibrosis: a double-blind multicenter trial. The Italian
36 Group for the Study of Ursodeoxycholic Acid in Cystic Fibrosis, *Hepatology*, 23, 1484-90,
37 1996
- 38 **Conrad et al., 2015**
- 39 Conrad, C., Lymp, J., Thompson, V., Dunn, C., Davies, Z., Chatfield, B., Nichols, D., Clancy,
40 J., Vender, R., Egan, M. E., Quittell, L., Michelson, P., Antony, V., Spahr, J., Rubenstein, R.
41 C., Moss, R. B., Herzenberg, L. A., Goss, C. H., Tirouvanziam, R., Long-term treatment with

- 1 oral N-acetylcysteine: affects lung function but not sputum inflammation in cystic fibrosis
2 subjects. A phase II randomized placebo-controlled trial, *Journal of Cystic Fibrosis*, 14, 219-
3 27, 2015
- 4 **Conway et al., 1997**
- 5 Conway, S. P., Pond, M. N., Watson, A., Etherington, C., Robey, H. L., Goldman, M. H.,
6 Intravenous colistin sulphomethate in acute respiratory exacerbations in adult patients with
7 cystic fibrosis, *Thorax*, 52, 987-93, 1997
- 8 **Corry et al., 2008**
- 9 Corry, J., Poon, W., McPhee, N., Milner, A. D., Cruickshank, D., Porceddu, S. V., Rischin, D.,
10 Peters, L. J., Randomized study of percutaneous endoscopic gastrostomy versus
11 nasogastric tubes for enteral feeding in head and neck cancer patients treated with
12 (chemo)radiation, *Journal of Medical Imaging and Radiation Oncology*, 52, 503-510, 2008
- 13 **Cox et al., 2016**
- 14 Cox, N. S., Alison, J. A., Button, B. M., Wilson, J. W., Morton, J. M., Holland, A. E., Physical
15 activity participation by adults with cystic fibrosis: An observational study, *Respirology*, 21,
16 511-8, 2016
- 17 **Curtis et al., 2016**
- 18 Curtis, L., Burns, A., Personal Social Services Research Unit, University of Kent, Unit costs
19 of health and social care 2016, 2016, Canterbury: Personal Social Services Research Unit,
20 University of Kent
- 21 **Cystic Fibrosis Trust, 2015**
- 22 Cystic Fibrosis Trust, UK CF Registry, Data on intestinal obstruction disaggregated by history
23 of meconium ileus in relation to the UK, year 2015, [online; accessed 11 January 2017]
- 24 **Cystic Fibrosis Trust, 2016**
- 25 Cystic Fibrosis Trust, UK Cystic Fibrosis Registry 2015 Annual Data Report, London: Cystic
26 Fibrosis Trust, 2016
- 27 **Cystic Fibrosis Trust, 2011**
- 28 Cystic Fibrosis Trust, Standards for the Clinical Care of Children and Adults with Cystic
29 Fibrosis in the UK. London: Cystic Fibrosis Trust, 2011
- 30 **Cystic Fibrosis Trust, 2015a**
- 31 Cystic Fibrosis Trust, UK CF Registry, BMI and BMI percentile in relation to the UK, year
32 2015. [online; accessed 23 November 2016]
- 33 **Daniels et al., 2011**
- 34 Daniels, T., Goodacre, L., Sutton, C., Pollard, K., Conway, S., Peckham, D., Accurate
35 assessment of adherence: self-report and clinician report vs electronic monitoring of
36 nebulizers, *Chest*, 140, 425-32, 2011
- 37 **Darbee et al., 2005**
- 38 Darbee, J. C., Kanga, J. F., Ohtake, P. J., Physiologic evidence for high-frequency chest wall
39 oscillation and positive expiratory pressure breathing in hospitalized subjects with cystic
40 fibrosis, *Physical Therapy*, 85, 1278-89, 2005
- 41 **Darch et al., 2015**

- 1 Darch, S. E., McNally, Alan, Harrison, Freya, Corander, Jukka Barr, Helen L., Paszkiewicz,
2 Konrad, Holden, Stephen, Fogarty, Andrew, Crusz, Shanika A., Diggle, Stephen P.,
3 Recombination is a key driver of genomic and phenotypic diversity in a *P aeruginosa*
4 population during cystic fibrosis infection, *Scientific Reports*, 5, 7649, 2015
- 5 **D'Auria et al., 2000**
- 6 D'Auria, J. P., Christian, B. J., Henderson, Z. G., Haynes, B., The company they keep: the
7 influence of peer relationships on adjustment to cystic fibrosis during adolescence, *Journal of*
8 *Pediatric Nursing*, 15, 175-182, 2000
- 9 **DeBoeck et al., 1989**
- 10 De Boeck, K., Smet, M., Eggermont, E., Treatment of *Pseudomonas* lung infection in cystic
11 fibrosis with piperacillin plus tobramycin versus ceftazidime monotherapy: preliminary
12 communication, *Pediatric Pulmonology*, 7, 171-3, 1989
- 13 **DeBoeck et al., 2007**
- 14 De Boeck, K., De Baets, F., Malfroot, A., Desager, K., Mouchet, F., Proesmans, M., Do
15 inhaled corticosteroids impair long-term growth in prepubertal cystic fibrosis patients?,
16 *European Journal of Pediatrics*, 166, 23-8, 2007
- 17 **Debray et al., 2011**
- 18 Debray, D., Kelly, D., Houwen, R., Strandvik, B., Colombo, C., Best practice guidance for the
19 diagnosis and management of cystic fibrosis-associated liver disease, *Journal of Cystic*
20 *Fibrosis*, 10, Suppl 2, S29-36, 2011
- 21
- 22 **Dellon et al., 2012**
- 23 Dellon, E. P., Sawicki, G. S., Shores, M. D., Wolfe, J., Hanson, L. C., Physician practices for
24 communicating with patients with cystic fibrosis about the use of noninvasive and invasive
25 mechanical ventilation, *Chest*, 141, 1010-7, 2012
- 26 **Dentice et al., 2016**
- 27 Dentice, R. L., Elkins, M. R., Middleton, P. G., Bishop, J. R., Wark, P. A., Dorahy, D. J.,
28 Harmer, C. J., Hu, H., Bye, P. T., A randomised trial of hypertonic saline during
29 hospitalisation for exacerbation of cystic fibrosis, *Thorax*, 71, 141-7, 2016
- 30 **Department of Health, 2010**
- 31 Department of Health, *Healthcare Premises Cost Guides (HPCGs) 2010*, 2010
- 32 **Department of Health, 2016**
- 33 Department of Health, *NHS reference costs 2015 to 2016*, 2016
- 34 **DerSimonian & Laird, 1986**
- 35 DerSimonian, R., Laird, N., Meta-analysis in clinical trials, *Controlled Clinical Trials*, 7, 177-
36 88, 1986
- 37 **Diabetes.co.uk, 2012**
- 38 Cost of diabetes, 2012
- 39 **Dias et al., 2016**

- 1 Dias, Sofia, Welton, Nicky J., Sutton, Alex J., Ades, A. E., NICE, NICE DSU Technical
2 support document 2: a generalised linear modelling framework for pairwise and network
3 meta-analysis of randomised controlled trials, 2016
- 4 **Dixon-Woods et al., 2005**
- 5 Dixon-Woods, M., Agarwal, S., Jones, D., Young, B., Sutton, A., Synthesising qualitative and
6 quantitative evidence: a review of possible methods, *Journal of Health Services & Research*
7 *Policy*, 10, 45-53, 2005
- 8 **Dolan, 1997**
- 9 Dolan, P., Modeling valuations for EuroQol health states, *Medical care*, 35, 1095-108, 1997
- 10 **Donati et al., 1987**
- 11 Donati, M. A., Guenette, G., Auerbach, H., Prospective controlled study of home and hospital
12 therapy of cystic fibrosis pulmonary disease, *Journal of Pediatrics*, 111, 28-33, 1987
- 13 **Dupuis et al., 2011**
- 14 Dupuis, F., Duhamel, F., Gendron, S., Transitioning care of an adolescent with cystic fibrosis:
15 development of systemic hypothesis between parents, adolescents, and health care
16 professionals, *Journal of Family Nursing*, 17, 291-311, 2011
- 17 **Durie et al., 1980**
- 18 Durie, P. R., Bell, L., Linton, W., Corey, M. L., Forstner, G. G., Effect of cimetidine and
19 sodium bicarbonate on pancreatic replacement therapy in cystic fibrosis, *Gut*, 21, 778-86,
20 1980
- 21 **Eigen et al., 1995**
- 22 Eigen, H., Rosenstein, B. J., FitzSimmons, S., Schidlow, D. V., A multicenter study of
23 alternate-day prednisone therapy in patients with cystic fibrosis. Cystic Fibrosis Foundation
24 Prednisone Trial Group, *Journal of Pediatrics*, 126, 515-23, 1995
- 25 **Elborn et al., 1992**
- 26 Elborn, S., Colville, A., Cordon, S., Hiller, E. J., Shale, D., A comparison of intravenous
27 ceftazidime and aztreonam in the treatment of respiratory exacerbations in cystic fibrosis,
28 11th International Cystic Fibrosis Congress, 1992, [Abstract]
- 29 **Elkins et al., 2006**
- 30 Elkins, M. R., Robinson, M., Rose, B. R., Harbour, C., Moriarty, C. P., Marks, G. B.,
31 Belousova, E. G., Xuan, W., Bye, P. T., National Hypertonic Saline in Cystic Fibrosis Study,
32 Group, A controlled trial of long-term inhaled hypertonic saline in patients with cystic fibrosis,
33 *New England Journal of Medicine*, 354, 229-40, 2006
- 34 **Elliott et al., 2005**
- 35 Elliott, R. A., Thornton, J., Webb, A. K., Dodd, M., Tully, M. P., Comparing costs of home-
36 versus hospital-based treatment of infections in adults in a specialist cystic fibrosis center,
37 *International Journal of Technology Assessment in Health Care*, 21, 506-10, 2005
- 38 **Elphick & Southern, 2014**
- 39 Elphick, H. E., Southern, K. W., Antifungal therapies for allergic bronchopulmonary
40 aspergillosis in people with cystic fibrosis, *Cochrane Database of Systematic Reviews*, 11,
41 CD002204, 2014

- 1 **Enright et al., 2004**
- 2 Enright, S., Chatham, K., Ionescu, A. A., Unnithan, V. B., Shale, D. J., Inspiratory muscle
3 training improves lung function and exercise capacity in adults with cystic fibrosis, *Chest*,
4 126, 405-11, 2004
- 5 **Equi et al., 2002**
- 6 Equi, A., Balfour-Lynn, I. M., Bush, A., Rosenthal, M., Long term azithromycin in children with
7 cystic fibrosis: A randomised, placebo-controlled crossover trial, *Lancet*, 360, 978-984, 2002
- 8 **Ericsson et al., 2013**
- 9 Ericsson, A., Pollock, R. F., Hunt, B., Valentine, W. J., Evaluation of the cost-utility of insulin
10 degludec vs insulin glargine in Sweden, *Journal of Medical Economics*, 16, 1442-52, 2013
- 11 **Eriksen & Ingebretsen, 2006**
- 12 Eriksen, B. O., Ingebretsen, O. C., The progression of chronic kidney disease: a 10-year
13 population-based study of the effects of gender and age, *Kidney international*, 69, 375-82,
14 2006
- 15 **Esmond et al., 2006**
- 16 Esmond, G., Butler, M., McCormack, A. M., Comparison of hospital and home intravenous
17 antibiotic therapy in adults with cystic fibrosis, *Journal of Clinical Nursing*, 15, 52-60, 2006
- 18 **Etherington et al., 2008**
- 19 Etherington, C., Hall, M., Conway, S., Peckham, D., Denton, M., Clinical impact of reducing
20 routine susceptibility testing in chronic *P aeruginosa* infections in cystic fibrosis, *Journal of*
21 *Antimicrobial Chemotherapy*, 61, 425-7, 2008
- 22 **Eubanks et al., 2002**
- 23 Eubanks, V., Koppersmith, N., Wooldridge, N., Clancy, J. P., Lyrene, R., Arani, R. B., Lee, J.,
24 Moldawer, L., Atchison, J., Sorscher, E. J., Makris, C. M., Effects of megestrol acetate on
25 weight gain, body composition, and pulmonary function in patients with cystic fibrosis,
26 *Journal of Pediatrics*, 140, 439-44, 2002
- 27 **European Medicines Agency, 2012**
- 28 European Medicines Agency, Report of the workshop on endpoints for cystic fibrosis clinical
29 trials, 2012, London: European Medicines Agency
- 30 **Fagundes et al., 2004**
- 31 Fagundes, E. D. T., Silva, R. A. P., Roquete, M. L. V., Penna, F. J., Reis, F. J. C., Goulart, E.
32 M. A., Duque, C. G., Validation of the Williams ultrasound scoring system for the diagnosis of
33 liver disease in cystic fibrosis, *Jornal de Pediatria*, 80, 380-386, 2004
- 34 **Fair et al., 2000**
- 35 Fair, A., Griffiths, K., Osman, L. M., Attitudes to fertility issues among adults with cystic
36 fibrosis in Scotland. The Collaborative Group of Scottish Adult CF Centres, *Thorax*, 55, 672-
37 7, 2000
- 38 **Farmer et al., 2009**
- 39 Farmer, A. J., Wade, A. N., French, D. P., Simon, J., Yudkin, P., Gray, A., Craven, A.,
40 Goyder, L., Holman, R. R., Mant, D., Kinmonth, A. L., Neil, H. A., Blood glucose self-

- 1 monitoring in type 2 diabetes: a randomised controlled trial, Health Technology Assessment
2 (Winchester, England), 13, iii-iv, ix-xi, 1-50, 2009
- 3 **Filigno et al., 2012**
- 4 Filigno, S. S., Brannon, E. E., Chamberlin, L. A., Sullivan, S. M., Barnett, K. A., Powers, S.
5 W., Qualitative analysis of parent experiences with achieving cystic fibrosis nutrition
6 recommendations, Journal of Cystic Fibrosis, 11, 125-30, 2012
- 7 **Finkelstein et al., 1992**
- 8 Finkelstein, S. M., Wielinski, C. L., Kujawa, S. J., Loewenson, R., Warwick, W. J., The impact
9 of home monitoring and daily diary recording on patient status in cystic fibrosis, Pediatric
10 Pulmonology, 12, 3-10, 1992
- 11 **Flume et al., 2016**
- 12 Flume, P. A., Clancy, J. P., Retsch-Bogart, G. Z., Tullis, D. E., Bresnik, M., Derchak, P. A.,
13 Lewis, S. A., Ramsey, B. W., Continuous alternating inhaled antibiotics for chronic
14 pseudomonal infection in cystic fibrosis, Journal of Cystic Fibrosis, 15, 809-815, 2016
- 15 **Forest Laboratories UK, 2011**
- 16 Forest Laboratories UK, Colistimethate sodium powder for inhalation for the treatment of
17 Pseudomonas lung infection in cystic fibrosis – Forest submission to NICE.
18 COLO/DPI/02/05., 2011, Kent: Forest Laboratories UK
- 19 **Forest Laboratories UK, 2011a**
- 20 Forest Laboratories UK, A randomised, open label study to compare the efficacy and safety
21 of a dry powder formulation of inhaled colistimethate sodium and nebulised TNSFI
22 (tobramycin nebuliser solution for inhalation, TOBI®) in cystic fibrosis patients with *P*
23 *aeruginosa* lung infection. Final protocol no: COLO/DPI/02/06., 2011, Kent: Forest
24 Laboratories UK
- 25 **France et al., 2008**
- 26 France, M. W., Dodd, M. E., Govan, J. R., Doherty, C. J., Webb, A. K., Jones, A. M., The
27 changing epidemiology of Burkholderia species infection at an adult cystic fibrosis centre,
28 Journal of Cystic Fibrosis, 7, 368-72, 2008
- 29 **Francisco et al., 2002**
- 30 Francisco, M. P., Wagner, M. H., Sherman, J. M., Theriaque, D., Bowser, E., Novak, D. A.,
31 Ranitidine and omeprazole as adjuvant therapy to pancrelipase to improve fat absorption in
32 patients with cystic fibrosis, Journal of Pediatric Gastroenterology & Nutrition, 35, 79-83,
33 2002
- 34 **Frederiksen et al., 1999**
- 35 Frederiksen, B., Koch, C., Hoiby, N., Changing epidemiology of *P aeruginosa* infection in
36 Danish cystic fibrosis patients (1974-1995), Pediatric Pulmonology, 28, 159-66, 1999
- 37 **Fuchs et al., 1994**
- 38 Fuchs, H. J., Borowitz, D. S., Christiansen, D. H., Morris, E. M., Nash, M. L., Ramsey, B. W.,
39 Rosenstein, B. J., Smith, A. L., Wohl, M. E., Effect of aerosolized recombinant human DNase
40 on exacerbations of respiratory symptoms and on pulmonary function in patients with cystic
41 fibrosis, New England Journal of Medicine, 331, 637-642, 1994
- 42 **Gabriel et al., 1991**

- 1 Gabriel, S. E., Jaakkimainen, L., Bombardier, C., Risk for serious gastrointestinal
2 complications related to use of nonsteroidal anti-inflammatory drugs. A meta-analysis,
3 *Annals of Internal Medicine*, 115, 787-96, 1991
- 4 **Galeva et al., 2013**
- 5 Galeva, I., Konstan, M. W., Higgins, M., Angyalosi, G., Brockhaus, F., Piggott, S., Thomas, K.,
6 Chuchalin, A. G., Tobramycin inhalation powder manufactured by improved process in cystic
7 fibrosis: the randomized EDIT trial, *Current Medical Research and Opinion*, 29, 947-956,
8 2013
- 9 **Garcia Rodriguez & Jick, 1994**
- 10 Garcia Rodriguez, L. A., Jick, H., Risk of upper gastrointestinal bleeding and perforation
11 associated with individual non-steroidal anti-inflammatory drugs, *Lancet*, 343, 769-72, 1994
- 12 **Gold et al., 1985**
- 13 Gold, R., Overmeyer, A., Knie, B., Fleming, P. C., Levison, H., Controlled trial of ceftazidime
14 vs. ticarcillin and tobramycin in the treatment of acute respiratory exacerbations in patients
15 with cystic fibrosis, *Pediatric Infectious Disease*, 4, 172-7, 1985
- 16 **Goldbeck et al., 2014**
- 17 Goldbeck, L., Fidika, A., Herle, M., Quittner, A. L., Psychological interventions for individuals
18 with cystic fibrosis and their families, *Cochrane Database of Systematic Reviews*, 6,
19 CD003148, 2014
- 20 **Greally et al., 1994**
- 21 Greally, P., Hussain, M. J., Vergani, D., Price, J. F., Interleukin-1 alpha, soluble interleukin-2
22 receptor, and IgG concentrations in cystic fibrosis treated with prednisolone, *Archives of*
23 *Disease in Childhood*, 71, 35-9, 1994
- 24 **Grieve et al., 2003**
- 25 Grieve, R., Thompson, S., Normand, C., Suri, R., Bush, A., Wallis, C., A cost-effectiveness
26 analysis of rhDNase in children with cystic fibrosis, *International Journal of Technology*
27 *Assessment in Health Care*, 19, 71-9, 2003
- 28 **Griffiths et al., 2004**
- 29 Griffiths, A. L., Armstrong, D., Carzino, R., Robinson, P., Cystic fibrosis patients and families
30 support cross-infection measures, *European Respiratory Journal*, 24, 449-452, 2004
- 31 **Griffiths et al., 2005**
- 32 Griffiths, A. L., Jansen, K., Carlin, J. B., Grimwood, K., Carzino, R., Robinson, P. J., Massie,
33 J., Armstrong, D. S., Effects of segregation on an epidemic *P aeruginosa* strain in a cystic
34 fibrosis clinic, *American Journal of Respiratory & Critical Care Medicine*, 171, 1020-5, 2005
- 35 **Griffiths et al., 2012**
- 36 Griffiths, A. L., Wurzel, D. F., Robinson, P. J., Carzino, R., Massie, J., Australian epidemic
37 strain pseudomonas (AES-1) declines further in a cohort segregated cystic fibrosis clinic,
38 *Journal of Cystic Fibrosis*, 11, 49-52, 2012
- 39 **Grimaldi et al., 2015**
- 40 Grimaldi, C., Bremont, F., Berlioz-Baudoin, M., Brouard, J., Corvol, H., Couderc, L., Lezmi,
41 G., Pin, I., Petit, I., Reix, P., Remus, N., Schweitzer, C., Thumerelle, C., Dubus, J. C., Sweat

- 1 test practice in pediatric pulmonology after introduction of cystic fibrosis newborn screening,
2 European Journal of Pediatrics, 174, 1613-20, 2015
- 3 **Grob, 2008**
- 4 Grob, R., Is my sick child healthy? Is my healthy child sick?: changing parental experiences
5 of cystic fibrosis in the age of expanded newborn screening, Social Science & Medicine, 67,
6 1056-64, 2008
- 7 **Grossoehme et al., 2014**
- 8 Grossoehme, D. H., Filigno, S. S., Bishop, M., Parent routines for managing cystic fibrosis in
9 children, Journal of Clinical Psychology in Medical Settings, 21, 125-135, 2014
- 10 **Gruber et al., 2014**
- 11 Gruber, W., Orenstein, D. M., Braumann, K. M., Beneke, R., Interval exercise training in
12 cystic fibrosis - Effects on exercise capacity in severely affected adults, Journal of Cystic
13 Fibrosis, 13, 86-91, 2014
- 14 **Grzincich et al., 2008**
- 15 Grzincich, G. L., Longo, F., Faverzani, S., Chetta, A., Spaggiari, C., Pisi, G., Short term
16 effects of high frequency chest compression (HFCC) and positive expiratory pressure (PEP)
17 in adults with cystic fibrosis, European Respiratory Society Annual Congress, Berlin,
18 Germany, October, 2008, [Abstract]
- 19 **Gupta et al., 2012**
- 20 Gupta, S., Ahmed, F., Lodha, R., Gupta, Y. K., Kabra, S. K., Comparison of effects of 3 and
21 7% hypertonic saline nebulization on lung function in children with cystic fibrosis: a double-
22 blind randomized, controlled trial, Journal of Tropical Pediatrics, 58, 375-81, 2012
- 23 **Gutthann et al., 1997**
- 24 Gutthann, S. P., Garcia Rodriguez, L. A., Raiford, D. S., Individual nonsteroidal
25 antiinflammatory drugs and other risk factors for upper gastrointestinal bleeding and
26 perforation, Epidemiology (Cambridge, Mass.), 8, 18-24, 1997
- 27 **Hanning et al., 1993**
- 28 Hanning, R. M., Blimkie, C. J., Bar-Or, O., Lands, L. C., Moss, L. A., Wilson, W. M.,
29 Relationships among nutritional status and skeletal and respiratory muscle function in cystic
30 fibrosis: does early dietary supplementation make a difference?, American Journal of Clinical
31 Nutrition, 57, 580-7, 1993
- 32 **Haworth et al., 2002**
- 33 Haworth, C. S., Selby, P. L., Horrocks, A. W., Mawer, E. B., Adams, J. E., Webb, A. K., A
34 prospective study of change in bone mineral density over one year in adults with cystic
35 fibrosis, Thorax, 57, 719-23, 2002
- 36 **Hayden et al., 2013**
- 37 Hayden, J. A., van der Windt, D. A., Cartwright, J. L., Cote, P., Bombardier, C., Assessing
38 bias in studies of prognostic factors, Annals of Internal Medicine, 158, 280-6, 2013
- 39 **Hayes et al., 2010**
- 40 Hayes, D., Jr., West, S. E., Rock, M. J., Li, Z., Splaingard, M. L., Farrell, P. M., *P aeruginosa*
41 in children with cystic fibrosis diagnosed through newborn screening: assessment of clinic
42 exposures and microbial genotypes, Pediatric Pulmonology, 45, 708-16, 2010

- 1 **Health Technology Assessment, 2015**
- 2 Health Technology Assessment, Mannitol dry powder for inhalation for the treatment of cystic
3 fibrosis (Project record), Health Technology Assessment (Winchester, England), 2015
- 4 **Hebestreit et al., 2010**
- 5 Hebestreit, H., Kieser, S., Junge, S., Ballmann, M., Hebestreit, A., Schindler, C., Schenk, T.,
6 Posselt, H. G., Kriemler, S., Long-term effects of a partially supervised conditioning
7 programme in cystic fibrosis, *European Respiratory Journal*, 35, 578-83, 2010
- 8 **Heijerman et al., 1991**
- 9 Heijerman, H. G., Lamers, C. B., Bakker, W., Omeprazole enhances the efficacy of
10 pancreatin (pancrease) in cystic fibrosis, *Annals of Internal Medicine*, 114, 200-1, 1991
- 11 **Heijerman et al., 1993**
- 12 Heijerman, H. G., Lamers, C. B., Bakker, W., Dijkman, J. H., Improvement of fecal fat
13 excretion after addition of omeprazole to pancrease in cystic fibrosis is related to residual
14 exocrine function of the pancreas, *Digestive Diseases & Sciences*, 38, 1-6, 1993
- 15 **Heltshe et al., 2014**
- 16 Heltshe, S. L., Borowitz, D. S., Leung, D. H., Ramsey, B., Mayer-Hamblett, N., Early attained
17 weight and length predict growth faltering better than velocity measures in infants with CF,
18 *Journal of Cystic Fibrosis*, 13, 723-9, 2014
- 19 **Henry et al., 1996**
- 20 Henry, D., Lim, L. L., Garcia Rodriguez, L. A., Perez Gutthann, S., Carson, J. L., Griffin, M.,
21 Savage, R., Logan, R., Moride, Y., Hawkey, C., Hill, S., Fries, J. T., Variability in risk of
22 gastrointestinal complications with individual non-steroidal anti-inflammatory drugs: results of
23 a collaborative meta-analysis, *BMJ*, 312, 1563-1566, 1996
- 24 **Hilliard et al., 2014**
- 25 Hilliard, M. E., Hahn, A., Ridge, A. K., Eakin, M. N., Riekert, K. A., User Preferences and
26 Design Recommendations for an mHealth App to Promote Cystic Fibrosis Self-Management,
27 *JMIR MHealth and UHealth*, 2, e44, 2014
- 28 **Hodgkinson & Lester, 2002**
- 29 Hodgkinson, R., Lester, H., Stresses and coping strategies of mothers living with a child with
30 cystic fibrosis: implications for nursing professionals, *Journal of Advanced Nursing*, 39, 377-
31 83, 2002
- 32 **Hodson et al., 2002**
- 33 Hodson, M. E., Gallagher, C. G., Govan, J. R., A randomised clinical trial of nebulised
34 tobramycin or colistin in cystic fibrosis, *European Respiratory Journal*, 20, 658-64, 2002
- 35 **Hoiby & Pedersen, 1989**
- 36 Hoiby, N., Pedersen, S. S., Estimated risk of cross-infection with *P aeruginosa* in Danish
37 cystic fibrosis patients, *Acta Paediatrica Scandinavica*, 78, 395-404, 1989
- 38 **Hommerding et al., 2015**
- 39 Hommerding, P. X., Baptista, R. R., Makarewicz, G. T., Schindel, C. S., Donadio, M. V. F.,
40 Pinto, L. A., Marostica, P. J. C., Effects of an educational intervention of physical activity for

- 1 children and adolescents with cystic fibrosis: A randomized controlled trial, *Respiratory Care*,
2 60, 81-87, 2015
- 3 **Homnick et al., 1998**
- 4 Homnick, D. N., Anderson, K., Marks, J. H., Comparison of the flutter device to standard
5 chest physiotherapy in hospitalized patients with cystic fibrosis: a pilot study, *Chest*, 114,
6 993-7, 1998
- 7 **Homnick et al., 2004**
- 8 Homnick, D. N., Homnick, B. D., Reeves, A. J., Marks, J. H., Pimentel, R. S., Bonnema, S.
9 K., Cyproheptadine is an effective appetite stimulant in cystic fibrosis, *Pediatric Pulmonology*,
10 38, 129-34, 2004
- 11 **Horsley et al., 2016**
- 12 Horsley, A., Jones, A. M., Lord, R., Antibiotic treatment for *B cepacia* complex in people with
13 cystic fibrosis experiencing a pulmonary exacerbation, *Cochrane Database of Systematic*
14 *Reviews*, 1, CD009529, 2016
- 15 **Houston et al., 2013**
- 16 Houston, B. W., Mills, N., Solis-Moya, A., Inspiratory muscle training for cystic fibrosis,
17 *Cochrane Database of Systematic Reviews*, 11, CD006112, 2013
- 18 **Hubert et al., 2004**
- 19 Hubert, D., Fajac, I., Bienvenu, T., Desmazes-Dufeu, N., Ellaffi, M., Dall'ava-Santucci, J.,
20 Dusser, D., Diagnosis of cystic fibrosis in adults with diffuse bronchiectasis, *Journal of Cystic*
21 *Fibrosis*, 3, 15-22, 2004
- 22 **Hummelinck & Pollock, 2006**
- 23 Hummelinck, A., Pollock, K., Parents' information needs about the treatment of their
24 chronically ill child: A qualitative study, *Patient Education and Counseling*, 62, 228-234, 2006
- 25 **Hurley et al., 2015**
- 26 Hurley, M. N., Prayle, A. P., Flume, P., Intravenous antibiotics for pulmonary exacerbations
27 in people with cystic fibrosis, *Cochrane Database of Systematic Reviews*, 7, CD009730,
28 2015
- 29 **Iles & Lowton, 2010**
- 30 Iles, N., Lowton, K., What is the perceived nature of parental care and support for young
31 people with cystic fibrosis as they enter adult health services?, *Health and Social Care in the*
32 *Community*, 18, 21-29, 2010
- 33 **Iles et al., 2003**
- 34 Iles, R., Legh-Smith, J., Drummond, M., Prevost, A., Vowler, S., Economic evaluation of
35 Tobramycin nebuliser solution in cystic fibrosis, *Journal of Cystic Fibrosis*, 2, 120-8, 2003
- 36 **Jaques et al., 2008**
- 37 Jaques, A., Daviskas, E., Turton, J. A., McKay, K., Cooper, P., Stirling, R. G., Robertson, C.
38 F., Bye, P. T., Lesouef, P. N., Shadbolt, B., Anderson, S. D., Charlton, B., Inhaled mannitol
39 improves lung function in cystic fibrosis, *Chest*, 133, 1388-96, 2008
- 40 **Jensen et al., 1987**

- 1 Jensen, T., Pedersen, S. S., Garne, S., Heilmann, C., Hoiby, N., Koch, C., Colistin inhalation
2 therapy in cystic fibrosis patients with chronic *P aeruginosa* lung infection, Journal of
3 Antimicrobial Chemotherapy, 19, 831-8, 1987
- 4 **Jessup & Parkinson, 2010**
- 5 Jessup, M., Parkinson, C., "All at sea": the experience of living with cystic fibrosis, Qualitative
6 Health Research, 20, 352-64, 2010
- 7 **Jessup et al., 2016**
- 8 Jessup, M., Douglas, T., Priddis, L., Branch-Smith, C., Shields, L., Arest, C. F., Parental
9 Experience of Information and Education Processes Following Diagnosis of Their Infant With
10 Cystic Fibrosis Via Newborn Screening, Journal of Pediatric Nursing, 31, e233-41, 2016
- 11 **Johannesson et al., 1998**
- 12 Johannesson, M., Carlson, M., Brucefors, A. B., Hjelte, L., Cystic fibrosis through a female
13 perspective: psychosocial issues and information concerning puberty and motherhood,
14 Patient Education & Counseling, 34, 115-23, 1998
- 15 **Jones et al., 2005**
- 16 Jones, A. M., Dodd, M. E., Govan, J. R., Doherty, C. J., Smith, C. M., Isalska, B. J., Webb, A.
17 K., Prospective surveillance for *P aeruginosa* cross-infection at a cystic fibrosis center,
18 American Journal of Respiratory & Critical Care Medicine, 171, 257-60, 2005
- 19 **Kalnins et al., 2005**
- 20 Kalnins, D., Corey, M., Ellis, L., Pencharz, P. B., Tullis, E., Durie, P. R., Failure of
21 conventional strategies to improve nutritional status in malnourished adolescents and adults
22 with cystic fibrosis, Journal of Pediatrics, 147, 399-401, 2005
- 23 **Karlas et al., 2012**
- 24 Karlas, T., Neuschulz, M., Oltmanns, A., Guttler, A., Petroff, D., Wirtz, H., Mainz, J. G.,
25 Mossner, J., Berg, T., Troltsch, M., Keim, V., Wiegand, J., Non-invasive evaluation of cystic
26 fibrosis related liver disease in adults with ARFI, transient elastography and different fibrosis
27 scores, PLoS ONE, 7, e42139, 2012
- 28 **Kazmerski et al., 2016**
- 29 Kazmerski, T. M., Borrero, S., Tuchman, L. K., Weiner, D. J., Pilewski, J. M., Orenstein, D.
30 M., Miller, E., Provider and Patient Attitudes Regarding Sexual Health in Young Women With
31 Cystic Fibrosis, Pediatrics, 137, 2016
- 32 **Kind et al., 1999**
- 33 Kind, P., Hardman, G., Macran, S., Centre for Health Economics, University of York, UK
34 Population Norms for EQ-5D, 1999, York: University of York
- 35 **Kirk & Milnes, 2016**
- 36 Kirk, S., Milnes, L., An exploration of how young people and parents use online support in
37 the context of living with cystic fibrosis, Health Expectations, 19, 309-21, 2016
- 38 **Kitson et al., 2013**
- 39 Kitson, M. T., Kemp, W. W., Iser, D. M., Paul, E., Wilson, J. W., Roberts, S. K., Utility of
40 transient elastography in the non-invasive evaluation of cystic fibrosis liver disease, Liver
41 International, 33, 698-705, 2013

- 1 **Klijn et al., 2004**
- 2 Klijn, P. H. C., Oudshoorn, A., Van Der Ent, C. K., Van Der Net, J., Kimpen, J. L., Helders, P.
3 J. M., Effects of anaerobic training in children with cystic fibrosis: A randomized controlled
4 study, *Chest*, 125, 1299-1305, 2004
- 5 **Konstan et al., 1995**
- 6 Konstan, M.W., Byard, P.J., Hoppel, C.L., Davis, P.B., Effect of high-dose ibuprofen in
7 patients with cystic fibrosis, *New England Journal of Medicine*, 332, 848-854, 1995
- 8 **Konstan et al., 2011**
- 9 Konstan, M. W., Geller, D. E., Minic, P., Brockhaus, F., Zhang, J., Angyalosi, G., Tobramycin
10 inhalation powder for *P aeruginosa* infection in cystic fibrosis: the EVOLVE trial, *Pediatric*
11 *Pulmonology*, 46, 230-8, 2011
- 12 **Konstan et al., 2011a**
- 13 Konstan, M.W., Flume, P.A., Kappler, M., Chiron, R., Higgins, M., Brockhaus, F., Zhang, J.,
14 Angyalosi, G., He, E., Geller, D.E., Safety, efficacy and convenience of tobramycin inhalation
15 powder in cystic fibrosis patients: The EAGER trial, *Journal of Cystic Fibrosis*, 10, 54-61,
16 2011
- 17 **Konstan et al., 2012**
- 18 Konstan, Michael W., Wagener, Jeffrey S., VanDevanter, Donald R., Pasta, David J., Yegin,
19 Ashley, Rasouliyan, Lawrence, Morgan, Wayne J., Risk factors for rate of decline in FEV₁ in
20 adults with cystic fibrosis, *Journal of Cystic Fibrosis*, 11, 405-411, 2012
- 21 **Kriemler et al., 2013**
- 22 Kriemler, S., Kieser, S., Junge, S., Ballmann, M., Hebestreit, A., Schindler, C., Stussi, C.,
23 Hebestreit, H., Effect of supervised training on FEV₁ in cystic fibrosis: a randomised
24 controlled trial, *Journal of Cystic Fibrosis*, 12, 714-20, 2013
- 25 **Lai & Shoff, 2008**
- 26 Lai, H. J., Shoff, S. M., Classification of malnutrition in cystic fibrosis: implications for
27 evaluating and benchmarking clinical practice performance, *American Journal of Clinical*
28 *Nutrition*, 88, 161-6, 2008
- 29 **Lai et al., 2000**
- 30 Lai, H. C., FitzSimmons, S. C., Allen, D. B., Kosorok, M. R., Rosenstein, B. J., Campbell, P.
31 W., Farrell, P. M., Risk of persistent growth impairment after alternate-day prednisone
32 treatment in children with cystic fibrosis, *New England Journal of Medicine*, 342, 851-9, 2000
- 33 **Lands & Stanojevic, 2016**
- 34 Lands, L. C., Stanojevic, S., Oral non-steroidal anti-inflammatory drug therapy for lung
35 disease in cystic fibrosis, *Cochrane Database of Systematic Reviews*, 4, CD001505, 2016
- 36 **Lands et al., 2007**
- 37 Lands, L.C., Milner, R., Cantin, A.M., Manson, D., Corey, M., High-Dose Ibuprofen in Cystic
38 Fibrosis: Canadian Safety and Effectiveness Trial, *Journal of Pediatrics*, 151, 249-254, 2007
- 39 **Lang et al., 2005**
- 40 Lang, L., Duff, A. J., Brownlee, K. G., Introducing the need for lung transplantation in children
41 with cystic fibrosis: parental experiences, *Journal of Cystic Fibrosis*, 4, 259-62, 2005

- 1 **Langton Hewer & Smyth, 2014**
- 2 Langton Hewer, S. C., Smyth, A. R., Antibiotic strategies for eradicating *P aeruginosa* in
3 people with cystic fibrosis, Cochrane Database of Systematic Reviews, 11, CD004197, 2014
- 4 **Laube et al., 1996**
- 5 Laube, B. L., Auci, R. M., Shields, D. E., Christiansen, D. H., Lucas, M. K., Fuchs, H. J.,
6 Rosenstein, B. J., Effect of rhDNase on airflow obstruction and mucociliary clearance in
7 cystic fibrosis, American Journal of Respiratory & Critical Care Medicine, 153, 752-60, 1996
- 8 **Lee et al., 2004**
- 9 Lee, T. W., Brownlee, K. G., Denton, M., Littlewood, J. M., Conway, S. P., Reduction in
10 prevalence of chronic *P aeruginosa* infection at a regional pediatric cystic fibrosis center,
11 Pediatric Pulmonology, 37, 104-10, 2004
- 12 **Lee et al., 2011**
- 13 Lee, Joyce M., Rhee, Kirsten, O'Grady, Michael J., Basu, Anirban, Winn, Aaron, John, Priya,
14 Meltzer, David O., Kollman, Craig, Laffel, Lori M., Lawrence, Jean M., Tamborlane, William
15 V., Wysocki, Tim, Xing, Dongyuan, Huang, Elbert S., Health Utilities for Children and Adults
16 with Type 1 Diabetes, Medical Care, 49, 924-931, 2011
- 17 **Lemaitre et al., 2016**
- 18 Lemaitre, C., Dominique, S., Billoud, E., Eliezer, M., Montialoux, H., Quillard, M., Riachi, G.,
19 Koning, E., Morisse-Pradier, H., Savoye, G., Savoye-Collet, C., Gorla, O., Relevance of 3D
20 Cholangiography and Transient Elastography to Assess Cystic Fibrosis-Associated Liver
21 Disease?, Canadian Respiratory Journal, 2016, 4592702, 2016
- 22 **Lenoir et al., 2007**
- 23 Lenoir, G., Antypkin, Y. G., Miano, A., Moretti, P., Zanda, M., Varoli, G., Monici Preti, P. A.,
24 Aryayev, N. L., Efficacy, safety, and local pharmacokinetics of highly concentrated nebulized
25 tobramycin in patients with cystic fibrosis colonized with *P aeruginosa*, Paediatric Drugs, 9
26 (Suppl 1), 11-20, 2007
- 27 **Lesko & Mitchell, 1995**
- 28 Lesko, S.M., Mitchell, A.A., An assessment of the safety of pediatric ibuprofen. A practitioner-
29 based randomized clinical trial, JAMA, 273, 929-933, 1995
- 30 **Lewin et al., 2015**
- 31 Lewin, Simon, Glenton, Claire, Munthe-Kaas, Heather, Carlsen, Benedicte, Colvin,
32 Christopher J., Gülmezoglu, Metin, Noyes, Jane, Booth, Andrew, Garside, Ruth, Rashidian,
33 Arash, Using Qualitative Evidence in Decision Making for Health and Social Interventions: An
34 Approach to Assess Confidence in Findings from Qualitative Evidence Syntheses (GRADE-
35 CERQual), PLoS Medicine, 12, e1001895, 2015
- 36 **Lewindon et al., 2011**
- 37 Lewindon, P. J., Shepherd, R. W., Walsh, M. J., Greer, R. M., Williamson, R., Pereira, T. N.,
38 Frawley, K., Bell, S. C., Smith, J. L., Ramm, G. A., Importance of hepatic fibrosis in cystic
39 fibrosis and the predictive value of liver biopsy, Hepatology, 53, 193-201, 2011
- 40 **Lewis et al., 2015**

- 1 Lewis, C., Blackman, S. M., Nelson, A., Oberdorfer, E., Wells, D., Dunitz, J., Thomas, W.,
2 Moran, A., Diabetes-related mortality in adults with cystic fibrosis. Role of genotype and sex,
3 American Journal of Respiratory & Critical Care Medicine, 191, 194-200, 2015
- 4 **Lindblad et al., 1999**
- 5 Lindblad, A., Glaumann, H., Strandvik, B., Natural history of liver disease in cystic fibrosis,
6 Hepatology, 30, 1151-8, 1999
- 7 **Lo et al., 2015**
- 8 Lo, D. K., Hurley, M. N., Muhlebach, M. S., Smyth, A. R., Interventions for the eradication of
9 meticillin-resistant *S aureus* (MRSA) in people with cystic fibrosis, Cochrane Database of
10 Systematic Reviews, 2, CD009650, 2015
- 11 **Lowson et al., 2011**
- 12 Lowson, K., Kelly, J., Bending, M., Whitehead, S., Wright, D., Lowson, P., Duffy, S.,
13 Trueman, P., Saxby, R., West, P., York Health Economics Consortium, University of York,
14 Cost-Effectiveness of Hospital Design: Options to Improve Patient Safety and Wellbeing ,
15 2011, York: University of York
- 16 **Lucidi et al., 2009**
- 17 Lucidi, V., Alghisi, F., Raia, V., Russo, B., Valmarana, L., Valmarana, R., Coruzzo, A.,
18 Beschi, S., Dester, S., Rinaldi, D., Maglieri, M., Guidotti, M. L., Ravaioli, E., Pesola, M., De
19 Alessandri, A., Padoan, R., Grynzich, L., Ratclif, L., Repetto, T., Ambroni, M., Provenzano,
20 E., Tozzi, A. E., Colombo, C., Growth assessment of paediatric patients with CF comparing
21 different auxologic indicators: A multicentre Italian study, Journal of Pediatric
22 Gastroenterology & Nutrition, 49, 335-42, 2009
- 23 **Lucidi et al., 2011**
- 24 Lucidi, V., Alghisi, F., Dall'Oglio, L., D'Apice, M. R., Monti, L., De Angelis, P., Gambardella,
25 S., Angioni, A., Novelli, G., The etiology of acute recurrent pancreatitis in children: a
26 challenge for pediatricians, Pancreas, 40, 517-21, 2011
- 27 **Maben et al., 2015**
- 28 Maben, J., Griffiths, P., Penfold, C., Simon, M., Pizzo, E., Anderson, J., Robert, G., Hughes,
29 J., Murrells, T., Brearley, S., Barlow, J., Evaluating a major innovation in hospital design:
30 workforce implications and impact on patient and staff experiences of all single room hospital
31 accommodation, 2015, In: Health Services and Delivery Research, Vol. 3, No. 3
- 32 **Macdonald & Greggans, 2010**
- 33 Macdonald, K., Greggans, A., 'Cool friends': an evaluation of a community befriending
34 programme for young people with cystic fibrosis, Journal of Clinical Nursing, 19, 2406-14,
35 2010
- 36 **Macfarlane et al., 1985**
- 37 Macfarlane, P. I., Hughes, D. M., Landau, L. I., Olinsky, A., The role of piperacillin therapy in
38 pulmonary exacerbations of cystic fibrosis: a controlled study, Pediatric Pulmonology, 1, 249-
39 55, 1985
- 40 **Maes et al., 2011**
- 41 Maes, I. H., Joore, M. A., Cima, R. F., Vlaeyen, J. W., Anteunis, L. J., Assessment of health
42 state in patients with tinnitus: a comparison of the EQ-5D and HUI mark III, Ear and Hearing,
43 32, 428-35, 2011

- 1 **Mainz et al., 2016**
- 2 Mainz, J. G., Schumacher, U., Schadlich, K., Hentschel, J., Koitschev, C., Koitschev, A.,
3 Riethmuller, J., Prenzel, F., Sommerburg, O., Wiedemann, B., Staab, D., Gleiber, W.,
4 Fischer, R., Beck, J. F., Arnold, C., Cooperators,, Sino nasal inhalation of isotonic versus
5 hypertonic saline (6.0%) in CF patients with chronic rhinosinusitis - Results of a multicenter,
6 prospective, randomized, double-blind, controlled trial, *Journal of Cystic Fibrosis*, 4, 4, 2016
- 7 **Mantel & Haenszel, 1959**
- 8 Mantel, N., Haenszel, W., Statistical aspects of the analysis of data from retrospective
9 studies of disease, *Journal of the National Cancer Institute*, 22, 719-48, 1959
- 10 **Marchand et al., 2000**
- 11 Marchand, V., Baker, S. S., Stark, T. J., Baker, R. D., Randomized, double-blind, placebo-
12 controlled pilot trial of megestrol acetate in malnourished children with cystic fibrosis, *Journal*
13 of *Pediatric Gastroenterology & Nutrition*, 31, 264-9, 2000
- 14 **Master et al., 2001**
- 15 Master, V., Roberts, G.W., Coulthard, K.P., Baghurst, P.A., Martin, A., Roberts, M.E.,
16 Onishko, C.R., Martin, A.J., Linke, R.J., Holmes, M., Jarvinen, A., Kennedy, D.,
17 Colebatch, K.A., Hansman, D., Parsons, D.W., Efficacy of once-daily tobramycin
18 monotherapy for acute pulmonary exacerbations of cystic fibrosis: a preliminary study,
19 *Pediatric Pulmonology*, 31, 367-376, 2001
- 20 **McCarty et al., 1988**
- 21 McCarty, J. M., Tilden, S. J., Black, P., Craft, J. C., Blumer, J., Waring, W., Halsey, N. A.,
22 Comparison of piperacillin alone versus piperacillin plus tobramycin for treatment of
23 respiratory infections in children with cystic fibrosis, *Pediatric Pulmonology*, 4, 201-4, 1988
- 24 **McCoy et al., 1996**
- 25 McCoy, K., Hamilton, S., Johnson, C., Effects of 12-week administration of dornase alfa in
26 patients with advanced cystic fibrosis lung disease. Pulmozyme Study Group, *Chest*, 110,
27 889-95, 1996
- 28 **McCoy et al., 2008**
- 29 McCoy, K. S., Quittner, A. L., Oermann, C. M., Gibson, R. L., Retsch-Bogart, G. Z.,
30 Montgomery, A. B., Inhaled aztreonam lysine for chronic airway *P aeruginosa* in cystic
31 fibrosis, *American Journal of Respiratory & Critical Care Medicine*, 178, 921-8, 2008
- 32 **Mcllwaine et al., 2001**
- 33 Mcllwaine, P. M., Wong, L. T., Peacock, D., Davidson, A. G., Long-term comparative trial of
34 positive expiratory pressure versus oscillating positive expiratory pressure (flutter)
35 physiotherapy in the treatment of cystic fibrosis, *Journal of Pediatrics*, 138, 845-50, 2001
- 36 **Mcllwaine et al., 2013**
- 37 Mcllwaine, M. P., Alarie, N., Davidson, G. F., Lands, L. C., Ratjen, F., Milner, R., Owen, B.,
38 Agnew, J. L., Long-term multicentre randomised controlled study of high frequency chest wall
39 oscillation versus positive expiratory pressure mask in cystic fibrosis, *Thorax*, 68, 746-51,
40 2013
- 41 **Mcllwaine et al., 2014**

- 1 Mcllwaine, M. P., Richmond, M., Agnew, J. L., Alarie, N., Lands, L., Chilvers, M., Ratjen, F.,
2 Cost-effectiveness of performing positive expiratory pressure versus high frequency chest
3 wall oscillation, *Journal of Cystic Fibrosis*, 13, S11, 2014
- 4 **Mcllwaine et al., 2015**
- 5 Mcllwaine, M, Button, B, Dwan, K, Positive expiratory pressure physiotherapy for airway
6 clearance in people with cystic fibrosis, *Cochrane Database of Systematic Reviews*,
7 CD003147, 2015
- 8 **McIntyre, 1999**
- 9 McIntyre, A. M., Dornase alpha and survival of patients with cystic fibrosis, *Hospital Medicine*
10 (London), 60, 736-9, 1999
- 11 **McKay et al., 2009**
- 12 McKay, K. O., Cooper, P. J., van Asperen, P. P., Segregation of children with CF diagnosed
13 via newborn screening and acquisition of *P aeruginosa*, *Journal of Cystic Fibrosis*, 8, 400-4,
14 2009
- 15 **McNamara et al., 1997**
- 16 McNamara, R. L., Lima, J. A., Whelton, P. K., Powe, N. R., Echocardiographic identification
17 of cardiovascular sources of emboli to guide clinical management of stroke: a cost-
18 effectiveness analysis, *Annals of Internal Medicine*, 127, 775-87, 1997
- 19 **Medicines Complete, 2017**
- 20 BNF 72 and BNF for Children (September 2016 – March 2017), 2017, London:
21 Pharmaceutical Press
- 22 **Menzin et al., 1996**
- 23 Menzin, J., Oster, G., Davies, L., Drummond, M. F., Greiner, W., Lucioni, C., Merot, J. L.,
24 Rossi, F., vd Schulenburg, J. G., Souetre, E., A multinational economic evaluation of
25 rhDNase in the treatment of cystic fibrosis, *International Journal of Technology Assessment*
26 *in Health Care*, 12, 52-61, 1996
- 27 **Merli et al., 1994**
- 28 Merli, M., Bertasi, S., Servi, R., Diamanti, S., Martino, F., De Santis, A., Goffredo, F.,
29 Quattrucci, S., Antonelli, M., Angelico, M., Effect of a medium dose of ursodeoxycholic acid
30 with or without taurine supplementation on the nutritional status of patients with cystic
31 fibrosis: a randomized, placebo-controlled, crossover trial, *Journal of Pediatric*
32 *Gastroenterology & Nutrition*, 19, 198-203, 1994
- 33 **Miller et al., 2009**
- 34 Miller, A. R., Condin, C. J., McKellin, W. H., Shaw, N., Klassen, A. F., Sheps, S., Continuity
35 of care for children with complex chronic health conditions: parents' perspectives, *BMC*
36 *Health Services Research*, 9, 242, 2009
- 37 **Minasian et al., 2010**
- 38 Minasian, C., Wallis, C., Metcalfe, C., Bush, A., Comparison of inhaled mannitol, daily
39 rhDNase and a combination of both in children with cystic fibrosis: a randomised trial,
40 *Thorax*, 65, 51-6, 2010
- 41 **Mitchell et al., 1982**

- 1 Mitchell, E. A., Qusted, C., Marks, R. E., Pinnock, R. E., Elliott, R. B., Comparative trial of
2 viokase, pancreatin and Pancrease pancrelipase (enteric coated beads) in the treatment of
3 malabsorption in cystic fibrosis, *Australian Paediatric Journal*, 18, 114-7, 1982
- 4 **Moen et al., 2011**
- 5 Moen, I. E., Nilsson, K., Andersson, A., Fagerland, M. W., Fluge, G., Hollsing, A., Gilljam, M.,
6 Mared, L., Pressler, T., Santi, H., Storrosten, O. T., Hjelte, L., Dietary intake and nutritional
7 status in a Scandinavian adult cystic fibrosis-population compared with recommendations,
8 *Food and Nutrition Research*, 55 (no pagination), 2011
- 9 **Moodie et al., 2014**
- 10 Moodie, M., Lal, A., Vidmar, S., Armstrong, D. S., Byrnes, C. A., Carlin, J. B., Cheney, J.,
11 Cooper, P. J., Grimwood, K., Robertson, C. F., Tiddens, H. A., Wainwright, C. E.,
12 Australasian Cystic Fibrosis Bronchoalveolar Lavage Study, Investigators, Costs of
13 bronchoalveolar lavage-directed therapy in the first 5 years of life for children with cystic
14 fibrosis, *Journal of Pediatrics*, 165, 564-569.e5, 2014
- 15 **Moola & Norman, 2011**
- 16 Moola, F. J., Norman, M. E., 'Down the rabbit hole': enhancing the transition process for
17 youth with cystic fibrosis and congenital heart disease by re-imagining the future and time,
18 *Child: Care, Health & Development*, 37, 841-51, 2011
- 19 **Moorcroft et al., 2004**
- 20 Moorcroft, A. J., Dodd, M. E., Morris, J., Webb, A. K., Individualised unsupervised exercise
21 training in adults with cystic fibrosis: a 1 year randomised controlled trial, *Thorax*, 59, 1074-
22 80, 2004
- 23 **Moran et al., 2010**
- 24 Moran, A., Brunzell, Carol, Cohen, Richard C., Katz, Marcia, Marshall, Bruce C., Onady,
25 Gary, Robinson, Karen A., Sabadosa, Kathryn A., Stecenko, Arlene, Slovis, Bonnie, The
26 CFRD Guidelines Committee, Clinical Care Guidelines for Cystic Fibrosis-Related Diabetes:
27 A position statement of the American Diabetes Association and a clinical practice guideline of
28 the Cystic Fibrosis Foundation, endorsed by the Pediatric Endocrine Society, *Diabetes Care*,
29 33, 2697-2708, 2010
- 30
- 31 **Moran et al., 2009**
- 32 Moran, A., Dunitz, J., Nathan, B., Saeed, A., Holme, B., Thomas, W., Cystic fibrosis-related
33 diabetes: current trends in prevalence, incidence, and mortality, *Diabetes Care*, 32, 1626-31,
34 2009
- 35 **Moran et al., 2014**
- 36 Moran, Fidelma, Bradley, Judy M., Piper, Amanda J., Non-invasive ventilation for cystic
37 fibrosis, *Cochrane Database of Systematic Reviews*, 4, CD00276, 2013
- 38 **Morrison & Agnew, 2014**
- 39 Morrison, L., Agnew, J., Oscillating devices for airway clearance in people with cystic fibrosis,
40 *Cochrane Database of Systematic Reviews*, 7, CD006842, 2014
- 41 **Morton & Wolfe, 2015**

- 1 Morton, A., Wolfe, S., Enteral tube feeding for cystic fibrosis, Cochrane Database of
2 Systematic Reviews, 4, CD001198, 2015
- 3 **Mueller-Abt et al., 2008**
- 4 Mueller-Abt, P. R., Frawley, K. J., Greer, R. M., Lewindon, P. J., Comparison of ultrasound
5 and biopsy findings in children with cystic fibrosis related liver disease, Journal of Cystic
6 Fibrosis, 7, 215-21, 2008
- 7 **Murphy et al., 2004**
- 8 Murphy, T. D., Anbar, R. D., Lester, L. A., Nasr, S. Z., Nickerson, B., VanDevanter, D. R.,
9 Colin, A. A., Treatment with tobramycin solution for inhalation reduces hospitalizations in
10 young CF subjects with mild lung disease, Pediatric Pulmonology, 38, 314-20, 2004
- 11 **National Horizon Scanning Centre, 2008**
- 12 National Horizon Scanning Centre, National Institute for Health Research, Mannitol dry
13 powder for inhalation (Bronchitol) for cystic fibrosis at risk for bronchiectasis, 2008,
14 Southampton: NIHR
- 15 **National Horizon Scanning Centre, 2008a**
- 16 National Horizon Scanning Centre, National Institute for Health Research, Transient
17 elastography (FibroScan) for evaluating liver fibrosis, 2008, Southampton: NIHR
- 18 **National Institute for Health and Care Excellence, 2013**
- 19 National Institute for Health and Care Excellence, Acute kidney injury: prevention, detection
20 and management, 2013
- 21 **National Institute for Health and Clinical Excellence, 2011**
- 22 National Institute for Health and Clinical Excellence, Common mental health problems:
23 identification and pathways to care, 2011
- 24 **National Institute for Health and Clinical Excellence, 2009**
- 25 National Institute for Health and Clinical Excellence, Depression in adults: recognition and
26 management, 2009
- 27 **National Institute for Clinical Excellence, 2004**
- 28 National Institute for Clinical Excellence, Eating disorders in over 8s: management, 2004
- 29 **National Institute for Clinical Excellence, 2005**
- 30 National Institute for Clinical Excellence, Depression in children and young people:
31 identification and management, 2005
- 32 **National Institute for Health and Care Excellence , 2013a**
- 33 National Institute for Health and Care Excellence, Generalised anxiety disorder and panic
34 disorder in adults: management, 2013
- 35 **National Institute for Health and Clinical Excellence, 2012**
- 36 National Institute for Health and Clinical Excellence, Mannitol dry powder for inhalation for
37 treating cystic fibrosis. Single technology appraisal, 2012
- 38 **National Institute for Health and Clinical Excellence, 2009a**

- 1 National Institute for Health and Clinical Excellence, Medicines adherence: involving patients
2 in decisions about prescribed medicines and supporting adherence, 2009
- 3 **National Institute for Health and Clinical Excellence, 2012a**
- 4 National Institute for Health and Clinical Excellence, NICE guidelines manual, 2012
- 5 **National Institute for Health and Care Excellence, 2014**
- 6 National Institute for Health and Care Excellence, NICE guidelines manual, 2014
- 7 **National Institute for Health and Care Excellence, 2015**
- 8 National Institute for Health and Care Excellence, Diabetes in pregnancy: management from
9 preconception to the postnatal period, 2015
- 10 **National Institute for Health and Care Excellence, 2015a**
- 11 National Institute for Health and Care Excellence, Type 1 diabetes in adults: diagnosis and
12 management, 2015
- 13 **National Institute for Health and Care Excellence, 2016**
- 14 National Institute for Health and Care Excellence, Integrated sensor-augmented pump
15 therapy systems for managing blood glucose levels in type 1 diabetes, 2016
- 16 **National Institute for Health and Care Excellence, 2016a**
- 17 National Institute for Health and Care Excellence, Cirrhosis in over 16s: assessment and
18 management, 2016
- 19 **National Institute for Health and Care Excellence, 2016b**
- 20 National Institute for Health and Care Excellence, NICE TA submission by Vertex.
21 (Lumacaftor and ivacaftor combination therapy for treating CF homozygous for the F508del
22 mutation [ID786]), 2016
- 23 **National Institute for Health and Clinical Excellence, 2012b**
- 24 National Institute for Health and Clinical Excellence, Patient experience in adult NHS
25 services: improving the experience of care for people using adult NHS services, 2012
- 26 **National Institute for Health and Care Excellence, 2016c**
- 27 National Institute for Health and Care Excellence, Transition from children's to adults'
28 services for young people using health or social care services, 2016
- 29 **National Institute for Health and Clinical Excellence, 2009b**
- 30 National Institute for Health and Clinical Excellence, Type 2 diabetes: The management of
31 type 2 diabetes, 2009
- 32 **Newbold et al., 2005**
- 33 Newbold, M. E., Tullis, E., Corey, M., Ross, B., Brooks, D., The Flutter Device versus the
34 PEP Mask in the treatment of adults with cystic fibrosis, *Physiotherapy Canada*, 57, 199-207,
35 2005
- 36 **NHS, 2015**
- 37 NHS, NHS Supply Chain 2015, 2015, Leeds: NHS
- 38 **NHS, 2015a**

- 1 NHS, National Diabetes Audit (NDA) Mortality Analysis 2012-2013, 2015, Leeds: NHS Digital
- 2 **NHS Business Services Authority,**
- 3 NHS Business Services Authority, NHS Electronic Drug Tariff, Newcastle upon Tyne: NHS
- 4 Business Services Authority
- 5 **NHS England,**
- 6 NHS England, Service Specifications: Cystic Fibrosis – Adults, Leeds: NHS
- 7 **NHS England,**
- 8 NHS England, Service Specifications: Cystic Fibrosis – Children, Leeds: NHS
- 9 **NHS Estates, 2005**
- 10 NHS Estates, Ward layouts with single rooms and space for flexibility, 2005, London: The
- 11 Stationery Office
- 12 **Nolan, Sarah et al., 2015**
- 13 Nolan Sarah, J., Thornton, Judith, Murray Clare, S., Dwyer, Tiffany, Inhaled mannitol for
- 14 cystic fibrosis, Cochrane Database of Systematic Reviews, 10, CD008649, 2015
- 15 **O'Brien et al., 1992**
- 16 O'Brien, S., Fitzgerald, M. X., Hegarty, J. E., A controlled trial of ursodeoxycholic acid
- 17 treatment in cystic fibrosis-related liver disease, European Journal of Gastroenterology and
- 18 Hepatology, 4, 857-863, 1992
- 19 **Oermann et al., 2001**
- 20 Oermann, C. M., Sockrider, M. M., Giles, D., Sontag, M. K., Accurso, F. J., Castile, R. G.,
- 21 Comparison of high-frequency chest wall oscillation and oscillating positive expiratory
- 22 pressure in the home management of cystic fibrosis: a pilot study, Pediatric Pulmonology, 32,
- 23 372-7, 2001
- 24 **Ooi et al., 2012**
- 25 Ooi, C. Y., Dupuis, A., Ellis, L., Jarvi, K., Martin, S., Gonska, T., Dorfman, R., Kortan, P.,
- 26 Solomon, M., Tullis, E., Durie, P. R., Comparing the American and European diagnostic
- 27 guidelines for cystic fibrosis: same disease, different language?, Thorax, 67, 618-24, 2012
- 28 **Orenstein et al., 2004**
- 29 Orenstein, D. M., Hovell, M. F., Mulvihill, M., Keating, K. K., Hofstetter, C. R., Kelsey, S.,
- 30 Morris, K., Nixon, P. A., Strength vs aerobic training in children with cystic fibrosis: a
- 31 randomized controlled trial, Chest, 126, 1204-14, 2004
- 32 **Padman et al., 1999**
- 33 Padman, R., Geouque, D. M., Engelhardt, M. T., Effects of the flutter device on pulmonary
- 34 function studies among pediatric cystic fibrosis patients, Delaware Medical Journal, 71, 13-8,
- 35 1999
- 36 **Papaioannou et al., 2008**
- 37 Papaioannou, A., Kennedy, C. C., Freitag, A., O'Neill, J., Pui, M., Ioannidis, G., Webber, C.,
- 38 Pathak, A., Hansen, S., Hennessey, R., Adachi, J. D., Longitudinal analysis of vertebral
- 39 fracture and BMD in a Canadian cohort of adult cystic fibrosis patients, BMC Musculoskeletal
- 40 Disorders, 9, 125, 2008

- 1 **Patriquin et al., 1999**
- 2 Patriquin, H., Lenaerts, C., Smith, L., Perreault, G., Grignon, A., Filiatrault, D., Boisvert, J.,
3 Roy, C. C., Rasquin-Weber, A., Liver disease in children with cystic fibrosis: US-biochemical
4 comparison in 195 patients, *Radiology*, 211, 229-32, 1999
- 5 **Philips et al., 2004**
- 6 Philips, Z., Ginnelly, L., Sculpher, M., Claxton, K., Golder, S., Riemsma, R., Woolacoot, N.,
7 Glanville, J., Review of guidelines for good practice in decision-analytic modelling in Health
8 Technology Assessment, *Health Technology Assessment (Winchester, England)*, 8, iii-iv, ix-
9 xi, 1-158, 2004
- 10 **Pignone et al., 2006**
- 11 Pignone, M., Earnshaw, S., Tice, J. A., Pletcher, M. J., Aspirin, statins, or both drugs for the
12 primary prevention of coronary heart disease events in men: a cost-utility analysis, *Annals of*
13 *Internal Medicine*, 144, 326-36, 2006
- 14 **Placidi et al., 2006**
- 15 Placidi, G., Cornacchia, M., Polese, G., Zanolla, L., Assael, B. M., Braggion, C., Chest
16 physiotherapy with positive airway pressure: a pilot study of short-term effects on sputum
17 clearance in patients with cystic fibrosis and severe airway obstruction, *Respiratory Care*, 51,
18 1145-53, 2006
- 19 **Pollock et al., 2010**
- 20 Pollock, R. F., Valentine, W. J., Goodall, G., Brandle, M., Evaluating the cost-effectiveness of
21 self-monitoring of blood glucose in type 2 diabetes patients on oral anti-diabetic agents,
22 *Swiss Medical Weekly*, 140, w13103, 2010
- 23 **Poustie et al., 2006**
- 24 Poustie, V. J., Russell, J. E., Watling, R. M., Ashby, D., Smyth, R. L., Calico Trial
25 Collaborative Group, Oral protein energy supplements for children with cystic fibrosis:
26 CALICO multicentre randomised controlled trial, *BMJ*, 332, 632-6, 2006
- 27 **Powers et al., 2003**
- 28 Powers, S. W., Byars, K. C., Mitchell, M. J., Patton, S. R., Schindler, T., Zeller, M. H., A
29 randomized pilot study of behavioural treatment to increase calorie intake in toddlers with
30 cystic fibrosis, *Children's Health Care*, 32, 297-311, 2003
- 31 **Powers et al., 2015**
- 32 Powers, S. W., Stark, L. J., Chamberlin, L. A., Filigno, S. S., Sullivan, S. M., Lemanek, K. L.,
33 Butcher, J. L., Driscoll, K. A., Daines, C. L., Brody, A. S., Schindler, T., Konstan, M. W.,
34 McCoy, K. S., Nasr, S. Z., Castile, R. G., Acton, J. D., Wooldridge, J. L., Ksenich, R. A.,
35 Szczeniak, R. D., Rausch, J. R., Stallings, V. A., Zemel, B. S., Clancy, J. P., Behavioral and
36 nutritional treatment for preschool-aged children with cystic fibrosis: a randomized clinical
37 trial, *JAMA Pediatrics*, 169, e150636, 2015
- 38 **Proesmans et al., 2013**
- 39 Proesmans, M., Vermeulen, F., Boulanger, L., Verhaegen, J., De Boeck, K., Comparison of
40 two treatment regimens for eradication of *P aeruginosa* infection in children with cystic
41 fibrosis, *Journal of Cystic Fibrosis*, 12, 29-34, 2013
- 42 **Quan et al., 2001**

- 1 Quan, J. M., Tiddens, H. A., Sy, J. P., McKenzie, S. G., Montgomery, M. D., Robinson, P. J.,
2 Wohl, M. E., Konstan, M. W., Pulmozyme Early Intervention Trial Study, Group, A two-year
3 randomized, placebo-controlled trial of dornase alfa in young patients with cystic fibrosis with
4 mild lung function abnormalities, *Journal of Pediatrics*, 139, 813-20, 2001
- 5 **Quon et al., 2011**
- 6 Quon, B.S., Mayer-Hamblett, N., Aitken, M.L., Smyth, A.R., Goss, C.H., Risk factors for
7 chronic kidney disease in adults with cystic fibrosis, *American Journal of Respiratory and
8 Critical Care Medicine*, 184, 1147-1152, 2011
- 9 **Qvist et al., 2016**
- 10 Qvist, Tavs, Taylor-Robinson, David, Waldmann, Elisabeth, Olesen, Hanne Vebert, Hansen,
11 Christine Rønne, Mathiesen, Inger Hee, Høiby, Niels, Katzenstein, Terese L., Smyth,
12 Rosalind L., Diggle, Peter J., Pressler, Tania, Comparing the harmful effects of
13 nontuberculous mycobacteria and Gram negative bacteria on lung function in patients with
14 cystic fibrosis, *Journal of Cystic Fibrosis*, 15, 380-385, 2016
- 15 **Radtke et al., 2015**
- 16 Radtke, T., Nolan, S. J., Hebestreit, H., Kriemler, S., Physical exercise training for cystic
17 fibrosis [Update of Cochrane Database Systematic Reviews. 2008; (1): CD002768],
18 *Cochrane Database of Systematic Reviews*, 6, CD002768, 2015
- 19 **Ramsey et al., 1993**
- 20 Ramsey, B. W., Dorkin, H. L., Eisenberg, J. D., Gibson, R. L., Harwood, I. R., Kravitz, R. M.,
21 Schidlow, D. V., Wilmott, R. W., Astley, S. J., McBurnie, M. A., et al., Efficacy of aerosolized
22 tobramycin in patients with cystic fibrosis, *New England Journal of Medicine*, 328, 1740-6,
23 1993
- 24 **Ramsey et al., 1993a**
- 25 Ramsey, B. W., Astley, S. J., Aitken, M. L., Burke, W., Colin, A. A., Dorkin, H. L., Eisenberg,
26 J. D., Gibson, R. L., Harwood, I. R., Schidlow, D. V., et al., Efficacy and safety of short-term
27 administration of aerosolized recombinant human deoxyribonuclease in patients with cystic
28 fibrosis, *American Review of Respiratory Disease*, 148, 145-51, 1993
- 29 **Ramsey et al., 1999**
- 30 Ramsey, B. W., Pepe, M. S., Quan, J. M., Otto, K. L., Montgomery, A. B., Williams-Warren,
31 J., Vasiljev, K. M., Borowitz, D., Bowman, C. M., Marshall, B. C., Marshall, S., Smith, A. L.,
32 Intermittent administration of inhaled tobramycin in patients with cystic fibrosis. Cystic
33 Fibrosis Inhaled Tobramycin Study Group, *New England Journal of Medicine*, 340, 23-30,
34 1999
- 35 **Rana et al., 2014**
- 36 Rana, M., Wong-See, D., Katz, T., Gaskin, K., Whitehead, B., Jaffe, A., Coakley, J.,
37 Lochhead, A., Fat-soluble vitamin deficiency in children and adolescents with cystic fibrosis,
38 *Journal of Clinical Pathology*, 67, 605-8, 2014
- 39 **Ranasinha et al., 1993**
- 40 Ranasinha, C., Assoufi, B., Shak, S., Christiansen, D., Fuchs, H., Empey, D., Geddes, D.,
41 Hodson, M., Efficacy and safety of short-term administration of aerosolised recombinant
42 human DNase I in adults with stable stage cystic fibrosis, *Lancet*, 342, 199-202, 1993
- 43 **Rath et al., 2012**

- 1 Rath, T., Menendez, K. M., Kugler, M., Hage, L., Wenzel, C., Schulz, R., Graf, J., Nahrlich,
2 L., Roeb, E., Roderfeld, M., TIMP-1/-2 and transient elastography allow non invasive
3 diagnosis of cystic fibrosis associated liver disease, *Digestive & Liver Disease*, 44, 780-7,
4 2012
- 5 **Rath et al., 2013**
- 6 Rath, T., Hage, L., Kugler, M., Menendez Menendez, K., Zachoval, R., Naehrlich, L., Schulz,
7 R., Roderfeld, M., Roeb, E., Serum Proteome Profiling Identifies Novel and Powerful Markers
8 of Cystic Fibrosis Liver Disease, *PLoS ONE*, 8, 2013
- 9 **Ratjen et al., 1985**
- 10 Ratjen, F., Wonne, R., Posselt, H. G., Stover, B., Hofmann, D., Bender, S. W., A double-blind
11 placebo controlled trial with oral ambroxol and N-acetylcysteine for mucolytic treatment in
12 cystic fibrosis, *European Journal of Pediatrics*, 144, 374-8, 1985
- 13 **Regan & Bhatt, 2016**
- 14 Regan, K. H., Bhatt, J., Eradication therapy for *B cepacia* complex in people with cystic
15 fibrosis, *Cochrane Database of Systematic Reviews*, 11, CD009876, 2016
- 16 **Remington et al., 2016**
- 17 Remington, T., Jahnke, N., Harkensee, C., Oral anti-pseudomonal antibiotics for cystic
18 fibrosis, *Cochrane Database of Systematic Reviews* *Cochrane Database Syst Rev*, 7,
19 CD005405, 2016
- 20 **Retsch-Bogart et al., 2009**
- 21 Retsch-Bogart, G. Z., Quittner, A. L., Gibson, R. L., Oermann, C. M., McCoy, K. S.,
22 Montgomery, A. B., Cooper, P. J., Efficacy and safety of inhaled aztreonam lysine for airway
23 pseudomonas in cystic fibrosis, *Chest*, 135, 1223-32, 2009
- 24 **Richard et al., 1997**
- 25 Richard, D. A., Nousia-Arvanitakis, S., Sollich, V., Hampel, B. J., Sommerauer, B., Schaad,
26 U. B., Oral ciprofloxacin vs. intravenous ceftazidime plus tobramycin in pediatric cystic
27 fibrosis patients: comparison of antipseudomonas efficacy and assessment of safety with
28 ultrasonography and magnetic resonance imaging. Cystic Fibrosis Study Group, *Pediatric*
29 *Infectious Disease Journal*, 16, 572-8, 1997
- 30 **Riemsma et al., 2011**
- 31 Riemsma, R., Maiwenn, J. Al, Armstrong, Nigel, Misso, Kate, Allen, Alexander, Manning,
32 Nathan, Tusabe, Doreen Allen, Severans, Johan L., Kleijnen, Jos, Kleijnen Systematic
33 Reviews Ltd., Mannitol dry powder for inhalation for the treatment of cystic fibrosis: A Single
34 Technology Appraisal, 2011, York: Kleijnen Systematic Reviews Ltd
- 35 **Riethmueller et al., 2002**
- 36 Riethmueller, J., Busch, A., Damm, V., Ziebach, R., Stern, M., Home and hospital antibiotic
37 treatment prove similarly effective in cystic fibrosis, *Infection*, 30, 387-91, 2002
- 38 **Robinson et al., 2012**
- 39 Robinson, P., Schechter, M. S., Sly, P. D., Winfield, K., Smith, J., Brennan, S., Shinkai, M.,
40 Henke, M. O., Rubin, B. K., Clarithromycin therapy for patients with Cystic Fibrosis: A
41 randomized controlled trial, *Pediatric Pulmonology*, 47, 551-557, 2012
- 42 **Roehrer et al., 2013**

- 1 Roehrer, E., Cummings, E., Beggs, S., Turner, P., Hauser, J., Micallef, N., Ellis, L., Reid, D.,
2 Pilot evaluation of web enabled symptom monitoring in cystic fibrosis, *Informatics for Health
3 & Social Care*, 38, 354-65, 2013
- 4 **Rogers et al., 2012**
- 5 Rogers, Chris, Thomas, Helen, Taylor, Rhiannon, Cromwell, David, Parameshwar, Jayan,
6 Banner, Nicholas R., UKCTA Steering Group, Royal College of Surgeons of England, UK
7 Cardiothoracic Transplant Audit in patients who received a transplant between 1st July 1995
8 and 31st March 2012. Annual Report, 2012, London: Royal College of Surgeons of England
- 9 **Rosenfeld et al., 2012**
- 10 Rosenfeld, M., Ratjen, F., Brumback, L., Daniel, S., Rowbotham, R., McNamara, S.,
11 Johnson, R., Kronmal, R., Davis, S. D., Isis Study Group, Inhaled hypertonic saline in infants
12 and children younger than 6 years with cystic fibrosis: the ISIS randomized controlled trial,
13 *JAMA*, 307, 2269-77, 2012
- 14 **Rovedder et al., 2014**
- 15 Rovedder, P. M., Flores, J., Ziegler, B., Casarotto, F., Jaques, P., Barreto, S. S., et al.,
16 Exercise programme in patients with cystic fibrosis: a randomized controlled trial, *Respiratory
17 Medicine*, 108, 1134-40, 2014
- 18 **Royal College of Paediatrics and Child Health, 2017**
- 19 Royal College of Paediatrics and Child Health, School age charts and resources 2017, 2017,
20 London: Royal College of Paediatrics and Child Health
- 21 **Russell et al., 1996**
- 22 Russell, M. T., Reinbold, J., Maltby, H. J., Transferring to adult health care: experiences of
23 adolescents with cystic fibrosis, *Journal of Pediatric Nursing*, 11, 262-268, 1996
- 24 **Russo et al., 2006**
- 25 Russo, K., Donnelly, M., Reid, A. J., Segregation--the perspectives of young patients and
26 their parents, *Journal of Cystic Fibrosis*, 5, 93-9, 2006
- 27 **Ryan et al., 2011**
- 28 Ryan, G., Singh, M., Dwan, K., Inhaled antibiotics for long-term therapy in cystic fibrosis,
29 *Cochrane Database of Systematic Reviews*, 3, CD001021, 2011
- 30 **Ryan et al., 2012**
- 31 Ryan, G., Jahnke, N., Remington, T., Inhaled antibiotics for pulmonary exacerbations in
32 cystic fibrosis, *Cochrane Database of Systematic Reviews*, 12, CD008319, 2012
- 33 **Sadler et al., 2015**
- 34 Sadler, M. D., Crotty, P., Fatovich, L., Wilson, S., Rabin, H. R., Myers, R. P., Noninvasive
35 methods, including transient elastography, for the detection of liver disease in adults with
36 cystic fibrosis, *Canadian Journal of Gastroenterology & Hepatology*, 29, 139-44, 2015
- 37 **Saiman et al., 2003**
- 38 Saiman, L., Marshall, B. C., Mayer-Hamblett, N., Burns, J. L., Quittner, A. L., Cibene, D. A.,
39 Coquillotte, S., Fieberg, A. Y., Accurso, F. J., Campbell, P. W., 3rd, Macrolide Study, Group,
40 Azithromycin in patients with cystic fibrosis chronically infected with *P aeruginosa*: a
41 randomized controlled trial, *JAMA*, 290, 1749-56, 2003

- 1 **Saiman et al., 2010**
- 2 Saiman, L., Anstead, M., Mayer-Hamblett, N., Lands, L. C., Kloster, M., Hocevar-Trnka, J.,
3 Goss, C. H., Rose, L. M., Burns, J. L., Marshall, B. C., Ratjen, F., A. Z. Azithromycin Study
4 Group, Effect of azithromycin on pulmonary function in patients with cystic fibrosis uninfected
5 with *P aeruginosa*: a randomized controlled trial, *JAMA*, 303, 1707-15, 2010
- 6 **Salh et al., 1992**
- 7 Salh, B., Bilton, D., Dodd, M., Abbot, J., Webb, K., A comparison of aztreonam and
8 ceftazidime in the treatment of respiratory infections in adults with cystic fibrosis,
9 *Scandinavian Journal of Infectious Diseases*, 24, 215-8, 1992
- 10 **Sanders et al., 2015**
- 11 Sanders, D. B., Li, Z., Brody, A. S., Chest computed tomography predicts the frequency of
12 pulmonary exacerbations in children with cystic fibrosis, *Annals of the American Thoracic*
13 *Society*, 12, 64-69, 2015
- 14 **Santana-Sosa et al., 2012**
- 15 Santana-Sosa, E., Groeneveld, I. F., Gonzalez-Saiz, L., Lopez-Mojares, L. M., Villa-Asensi,
16 J. R., Barrio Gonzalez, M. I., Fleck, S. J., Perez, M., Lucia, A., Intrahospital weight and
17 aerobic training in children with cystic fibrosis: a randomized controlled trial, *Medicine &*
18 *Science in Sports & Exercise*, 44, 2-11, 2012
- 19 **Santana-Sosa et al., 2014**
- 20 Santana-Sosa, E., Gonzalez-Saiz, L., Groeneveld, I. F., Villa-Asensi, J. R., Barrio Gomez de
21 Aguero, M. I., Fleck, S. J., Lopez-Mojares, L. M., Perez, M., Lucia, A., Benefits of combining
22 inspiratory muscle with 'whole muscle' training in children with cystic fibrosis: a randomised
23 controlled trial, *British Journal of Sports Medicine*, 48, 1513-7, 2014
- 24 **Savage et al., 2014**
- 25 Savage, E., Beirne, P. V., Ni Chroinin, M., Duff, A., Fitzgerald, T., Farrell, D., Self-
26 management education for cystic fibrosis, *Cochrane Database of Systematic Reviews*, 9,
27 CD007641, 2014
- 28 **Savant et al., 2014**
- 29 Savant, A. P., O'Malley, C., Bichl, S., McColley, S. A., Improved patient safety through
30 reduced airway infection rates in a paediatric cystic fibrosis programme after a quality
31 improvement effort to enhance infection prevention and control measures, *BMJ Quality &*
32 *Safety*, 23, i73-i80, 2014
- 33 **Schaad et al., 1987**
- 34 Schaad, U. B., Wedgwood-Krucko, J., Suter, S., Kraemer, R., Efficacy of inhaled amikacin as
35 adjunct to intravenous combination therapy (ceftazidime and amikacin) in cystic fibrosis,
36 *Journal of Pediatrics*, 111, 599-605, 1987
- 37 **Schaad et al., 1989**
- 38 Schaad, U. B., Wedgwood-Krucko, J., Guenin, K., Buehlmann, U., Kraemer, R.,
39 Antipseudomonal therapy in cystic fibrosis: aztreonam and amikacin versus ceftazidime and
40 amikacin administered intravenously followed by oral ciprofloxacin, *European Journal of*
41 *Clinical Microbiology & Infectious Diseases*, 8, 858-65, 1989
- 42 **Schechter et al., 2015**

- 1 Schechter, M. S., Trueman, D., Farquharson, R., Higuchi, K., Daines, C. L., Inhaled
2 Aztreonam Lysine versus Inhaled Tobramycin in Cystic Fibrosis. An Economic Evaluation,
3 *Annals of the American Thoracic Society*, 12, 1030-8, 2015
- 4 **Schindel et al., 2015**
- 5 Schindel, C. S., Hommerding, P. X., Melo, D. A. S., Baptista, R. R., Marostica, P. J. C.,
6 Donadio, M. V. F., Physical exercise recommendations improve postural changes found in
7 children and adolescents with cystic fibrosis: A randomized controlled trial, *Journal of*
8 *Pediatrics*, 166, 710-716, 2015
- 9 **Schneider et al., 2006**
- 10 Schneider, V., Levesque, L. E., Zhang, B., Hutchinson, T., Brophy, J. M., Association of
11 selective and conventional nonsteroidal antiinflammatory drugs with acute renal failure: A
12 population-based, nested case-control analysis, *American Journal of Epidemiology*, 164,
13 881-9, 2006
- 14 **Schneiderman-Walker et al., 2000**
- 15 Schneiderman-Walker, J., Pollock, S. L., Corey, M., Wilkes, D. D., Canny, G. J., Pedder, L.,
16 Reisman, J. J., A randomized controlled trial of a 3-year home exercise program in cystic
17 fibrosis, *Journal of Pediatrics*, 136, 304-10, 2000
- 18 **Schulze et al., 2006**
- 19 Schulze, K. J., Cutchins, C., Rosenstein, B. J., Germain-Lee, E. L., O'Brien, K. O., Calcium
20 acquisition rates do not support age-appropriate gains in total body bone mineral content in
21 prepuberty and late puberty in girls with cystic fibrosis, *Osteoporosis International*, 17, 731-
22 40, 2006
- 23 **Schuster et al., 2013**
- 24 Schuster, A., Haliburn, C., Doring, G., Goldman, M. H., Freedom Study, Group, Safety,
25 efficacy and convenience of colistimethate sodium dry powder for inhalation (Colobreathe
26 DPI) in patients with cystic fibrosis: a randomised study, *Thorax*, 68, 344-50, 2013
- 27 **Seear & Wensley, 1997**
- 28 Seear, M., Wensley, D., Chronic cough and wheeze in children: do they all have asthma?,
29 *European Respiratory Journal*, 10, 342-345, 1997
- 30 **Selvadurai et al., 2002**
- 31 Selvadurai, H. C., Blimkie, C. J., Meyers, N., Mellis, C. M., Cooper, P. J., Van Asperen, P. P.,
32 Randomized controlled study of in-hospital exercise training programs in children with cystic
33 fibrosis, *Pediatric Pulmonology*, 33, 194-200, 2002
- 34 **Sermet-Gaudelus et al., 2011**
- 35 Sermet-Gaudelus, I., Bianchi, M. L., Garabedian, M., Aris, R. M., Morton, A., Hardin, D. S.,
36 Elkin, S. L., Compston, J. E., Conway, S. P., Castanet, M., Wolfe, S., Haworth, C. S.,
37 European cystic fibrosis bone mineralisation guidelines, *Journal of Cystic Fibrosis*, 10, Suppl
38 2, S16-23, 2011
- 39 **Shah et al., 1995**
- 40 Shah, P. I., Bush, A., Canny, G. J., Colin, A. A., Fuchs, H. J., Geddes, D. M., Johnson, C. A.,
41 Light, M. C., Scott, S. F., Tullis, D. E., et al., Recombinant human DNase I in cystic fibrosis
42 patients with severe pulmonary disease: a short-term, double-blind study followed by six
43 months open-label treatment, *European Respiratory Journal*, 8, 954-8, 1995

- 1 **Shah et al., 1996**
- 2 Shah, P. L., Scott, S. F., Knight, R. A., Marriott, C., Ranasinha, C., Hodson, M. E., In vivo
3 effects of recombinant human DNase I on sputum in patients with cystic fibrosis, *Thorax*, 51,
4 119-25, 1996
- 5 **Shearer & Bryon, 2004**
- 6 Shearer, J. E., Bryon, M., The nature and prevalence of eating disorders and eating
7 disturbance in adolescents with cystic fibrosis, *Journal of the Royal Society of Medicine*, 97
8 (Suppl 44), 36-42, 2004
- 9 **Sheldon et al., 1993**
- 10 Sheldon, C. D., Assoufi, B. K., Hodson, M. E., Regular three monthly oral ciprofloxacin in
11 adult cystic fibrosis patients infected with *P aeruginosa*, *Respiratory Medicine*, 87, 587-93,
12 1993
- 13 **Silverstein et al., 2000**
- 14 Silverstein, F. E., Faich, G., Goldstein, J. L., Simon, L. S., Pincus, T., Whelton, A., Makuch,
15 R., Eisen, G., Agrawal, N. M., Stenson, W. F., Burr, A. M., Zhao, W. W., Kent, J. D.,
16 Lefkowitz, J. B., Verburg, K. M., Geis, G. S., Gastrointestinal toxicity with celecoxib vs
17 nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis: the CLASS
18 study: A randomized controlled trial. Celecoxib Long-term Arthritis Safety Study, *JAMA*, 284,
19 1247-55, 2000
- 20 **Siracusa et al., 2014**
- 21 Siracusa, C., Clancy, J. P., Drotar, D., Electronic monitoring reveals highly variable
22 adherence patterns in patients prescribed ivacaftor, *Pediatric Pulmonology*, 49, 440, 2014
- 23 **Smyth et al., 2014**
- 24 Smyth, A. R., Bell, Scott C., Bojcin, Snezana, Bryon, Mandy, Duff, Alistair, Flume, Patrick,
25 Kashirskaya, Nataliya, Munck, Anne, Ratjen, Felix, Schwarzenberg, Sarah Jane, Sermet-
26 Gaudelus, Isabelle, Southern, Kevin W., Taccetti, Giovanni, Ullrich, Gerald, Wolfe, Sue,
27 European Cystic Fibrosis Society Standards of Care: Best Practice guidelinesmoran
28 , *Journal of Cystic Fibrosis* ,13, Suppl 1, S23-S42, 2014
- 29
- 30 **Smyth & Rayner, 2014a**
- 31 Smyth, R. L., Rayner, O., Oral calorie supplements for cystic fibrosis, *Cochrane Database of*
32 *Systematic Reviews*, 11, CD000406, 2014
- 33 **Smyth & Walters, 2014**
- 34 Smyth, A. R., Walters, S., Prophylactic anti-staphylococcal antibiotics for cystic fibrosis,
35 *Cochrane Database of Systematic Reviews*, 11, CD001912, 2014
- 36 **Solem et al., 2014**
- 37 Solem, C. T., Vera Llonch, M., Liu, S., Botteman, M. F., Lasch, K., Rodriguez, S.,
38 Castiglione, B., Responsiveness Of The Eq-5d Index And Visual Analog Scale To Changes
39 In Lung Function In Patients With Cystic Fibrosis, *Value in Health*, 17, A178, 2014
- 40 **Solem et al., 2014a**

- 1 Solem, C. T., Vera-Llonch, M., Liu, S., Botteman, M., Lin, F. J., Castiglione, B., Impact of
2 pulmonary exacerbations on EQ-5D measures in patients with cystic fibrosis, *Value in*
3 *Health*, 17 (7), A535, 2014
- 4 **Somerville et al., 2013**
- 5 Somerville, R., Lackson, A., Zhou, S., Fletcher, C., Fitzpatrick, P., Non-pulmonary chronic
6 diseases in adults with cystic fibrosis: analysis of data from the Cystic Fibrosis Registry, *Irish*
7 *Medical Journal*, 106, 166-8, 2013
- 8 **Sordelli et al., 1994**
- 9 Sordelli, D.O., Macri, C.N., Maillie, A.J., Cerquetti, M.C., A preliminary study on the effect of
10 anti-inflammatory treatment in cystic fibrosis patients with *P aeruginosa* lung infection,
11 *International Journal of Immunopathology and Pharmacology*, 7, 109-117, 1994
- 12 **South Devon Healthcare NHS Foundation Trust, 2015**
- 13 South Devon Healthcare NHS Foundation Trust, Private patient and overseas visitor price
14 list, 2015, Torquay: South Devon Healthcare NHS Foundation Trust
- 15 **Southampton Children's Hospital, 2014**
- 16 Southampton Children's Hospital, Transition to adult care: ready steady go, from
17 <http://www.uhs.nhs.uk/Ourservices/Childhealth/TransitiontoadultcareReadySteadyGo/Transitiontoadultcare.aspx>, 2014
- 18
- 19 **Southern et al., 2012**
- 20 Southern, Kevin W., Barker, Pierre M., SolisMoya, Arturo, Patel, Latifa, Macrolide antibiotics
21 for cystic fibrosis, *Cochrane Database of Systematic Reviews*, 11, CD002203, 2012
- 22 **Stark et al., 1996**
- 23 Stark, L. J., Mulvihill, M. M., Powers, S. W., Jelalian, E., Keating, K., Creveling, S., Byrnes-
24 Collins, B., Harwood, I., Passero, M. A., Light, M., Miller, D. L., Hovell, M. F., Behavioral
25 intervention to improve calorie intake of children with cystic fibrosis: treatment versus wait list
26 control, *Journal of Pediatric Gastroenterology & Nutrition*, 22, 240-53, 1996
- 27 **Stark et al., 2009**
- 28 Stark, L. J., Quittner, A. L., Powers, S. W., Opiari-Arrigan, L., Bean, J. A., Duggan, C.,
29 Stallings, V. A., Randomized clinical trial of behavioral intervention and nutrition education to
30 improve caloric intake and weight in children with cystic fibrosis, *Archives of Pediatrics &*
31 *Adolescent Medicine*, 163, 915-21, 2009
- 32 **Stephenson et al., 2013**
- 33 Stephenson, A. L., Mannik, L. A., Walsh, S., Brotherwood, M., Robert, R., Darling, P. B.,
34 Nisenbaum, R., Moerman, J., Stanojevic, S., Longitudinal trends in nutritional status and the
35 relation between lung function and BMI in cystic fibrosis: a population-based cohort study,
36 *American Journal of Clinical Nutrition*, 97, 872-7, 2013
- 37 **Stevens, 2011**
- 38 Stevens, J. W., A note on dealing with missing standard errors in meta-analyses of
39 continuous outcome measures in WinBUGS, *Pharmaceutical Statistics*, 10, 374-8, 2011
- 40 **Stutman et al., 2002**

- 1 Stutman, H. R., Lieberman, J. M., Nussbaum, E., Marks, M. I., Antibiotic prophylaxis in
2 infants and young children with cystic fibrosis: a randomized controlled trial, *Journal of*
3 *Pediatrics*, 140, 299-305, 2002
- 4 **Suri et al., 2001**
- 5 Suri, R., Metcalfe, C., Lees, B., Grieve, R., Flather, M., Normand, C., Thompson, S., Bush,
6 A., Wallis, C., Comparison of hypertonic saline and alternate-day or daily recombinant
7 human deoxyribonuclease in children with cystic fibrosis: a randomised trial, *Lancet*, 358,
8 1316-21, 2001
- 9 **Suri et al., 2002**
- 10 Suri, R., Grieve, R., Normand, C., Metcalfe, C., Thompson, S., Wallis, C., Bush, A., Effects of
11 hypertonic saline, alternate day and daily rhDNase on healthcare use, costs and outcomes in
12 children with cystic fibrosis, *Thorax*, 57, 841-6, 2002
- 13 **Suri et al., 2002a**
- 14 Suri, R., Wallis, C., Bush, A., Thompson, S., Normand, C., Flather, M., Grieve, R.,
15 Metcalfe, C., Lees, B., A comparative study of hypertonic saline, daily and alternate-day
16 rhDNase in children with cystic fibrosis, *Health Technology Assessment (Winchester,*
17 *England)*, 6, 2002
- 18 **Suri et al., 2002b**
- 19 Suri, R., Marshall, L.J., Wallis, C., Metcalfe, C., Bush, A., Shute, J.K., Effects of recombinant
20 human DNase and hypertonic saline on airway inflammation in children with cystic fibrosis,
21 *American Journal of Respiratory and Critical Care Medicine*, 166, 352-355, 2002
- 22 **Taccetti et al., 2012**
- 23 Taccetti, G., Bianchini, E., Cariani, L., Buzzetti, R., Costantini, D., Trevisan, F., Zavataro, L.,
24 Campana, S., Italian Group for, *P aeruginosa* Eradication in Cystic Fibrosis, Early antibiotic
25 treatment for *P aeruginosa* eradication in patients with cystic fibrosis: a randomised
26 multicentre study comparing two different protocols, *Thorax*, 67, 853-9, 2012
- 27 **Tajima et al., 2010**
- 28 Tajima, R., Kondo, M., Kai, H., Saito, C., Okada, M., Takahashi, H., Doi, M., Tsuruoka, S.,
29 Yamagata, K., Measurement of health-related quality of life in patients with chronic kidney
30 disease in Japan with EuroQol (EQ-5D), *Clinical & Experimental Nephrology*, 14, 340-8,
31 2010
- 32 **Tannenbaum et al., 2005**
- 33 Tannenbaum, E., Prasad, S.A., Main, E., Scrase, E., 374 Long term effects of positive expiratory
34 pressure (PEP) or oscillatory positive pressure (RC cornet®) on FEV₁ and perceived health
35 in children with CF, *Journal of Cystic Fibrosis*, 4, S100, 2005
- 36 **Tappenden et al., 2013**
- 37 Tappenden, P., Harnan, S., Uttley, L., Mildred, M., Carroll, C., Cantrell, A., Colistimethate
38 sodium powder and tobramycin powder for inhalation for the treatment of chronic *P*
39 *aeruginosa* lung infection in cystic fibrosis: systematic review and economic model, *Health*
40 *Technology Assessment (Winchester, England)*, 17, v-xvii, 2013
- 41 **Tappenden et al., 2014**

- 1 Tappenden, P., Harnan, S., Uttley, L., Mildred, M., Walshaw, M., Taylor, C., Brownlee, K.,
2 The cost effectiveness of dry powder antibiotics for the treatment of *P aeruginosa* in patients
3 with cystic fibrosis, *Pharmacoeconomics*, 32, 159-72, 2014
- 4 **The Cochrane Collaboration, 2011**
- 5 The Cochrane Collaboration, The Cochrane Collaboration's tool for assessing risk of bias,
6 2011
- 7 **Thomas et al., 2006**
- 8 Thomas, C., Mitchell, P., O'Rourke, P., Wainwright, C., Quality-of-life in children and
9 adolescents with cystic fibrosis managed in both regional outreach and cystic fibrosis center
10 settings in Queensland, *Journal of Pediatrics*, 148, 508-516, 2006
- 11 **Thomas et al., 2008**
- 12 Thomas, C. L., O'Rourke, P. K., Wainwright, C. E., Clinical outcomes of Queensland children
13 with cystic fibrosis: a comparison between tertiary centre and outreach services, *Medical*
14 *Journal of Australia*, 188, 135-9, 2008
- 15 **Thomassen et al., 1986**
- 16 Thomassen, M. J., Demko, C. A., Doershuk, C. F., Stern, R. C., Klinger, J. D., *Pseudomonas*
17 *cepacia*: decrease in colonization in patients with cystic fibrosis, *American Review of*
18 *Respiratory Disease*, 134, 669-71, 1986
- 19 **Thornton et al., 2005**
- 20 Thornton, J., Elliott, R. A., Tully, M. P., Dodd, M., Webb, A. K., Clinical and economic choices
21 in the treatment of respiratory infections in cystic fibrosis: comparing hospital and home care,
22 *Journal of Cystic Fibrosis*, 4, 239-47, 2005
- 23 **Tierney et al., 2013**
- 24 Tierney, S., Deaton, C., Jones, A., Oxley, H., Biesty, J., Kirk, S., Liminality and transfer to
25 adult services: a qualitative investigation involving young people with cystic fibrosis,
26 *International Journal of Nursing Studies*, 50, 738-46, 2013
- 27 **Tipping et al., 2010**
- 28 Tipping, C. J., Scholes, R. L., Cox, N. S., A qualitative study of physiotherapy education for
29 parents of toddlers with cystic fibrosis, *Journal of Cystic Fibrosis*, 9, 205-11, 2010
- 30 **Tluczek et al., 2006**
- 31 Tluczek, A., Kosciak, R. L., Modaff, P., Pfeil, D., Rock, M. J., Farrell, P. M., Lifchez, C.,
32 Freeman, M. E., Gershan, W., Zaleski, C., Sullivan, B., Newborn screening for cystic fibrosis:
33 parents' preferences regarding counseling at the time of infants' sweat test, *Journal of*
34 *Genetic Counseling*, 15, 277-91, 2006
- 35 **Tluczek et al., 2009**
- 36 Tluczek, A., Orland, K. M., Nick, S. W., Brown, R. L., Newborn screening: an appeal for
37 improved parent education, *Journal of Perinatal & Neonatal Nursing*, 23, 326-34, 2009
- 38 **Trapnell et al., 2012**
- 39 Trapnell, B. C., McColley, S. A., Kissner, D. G., Rolfe, M. W., Rosen, J. M., McKevitt, M.,
40 Moorehead, L., Montgomery, A. B., Geller, D. E., Phase, F. T. I. Study Group,
41 Fosfomycin/tobramycin for inhalation in patients with cystic fibrosis with *pseudomonas* airway
42 infection, *American Journal of Respiratory & Critical Care Medicine*, 185, 171-8, 2012

- 1 **Tuchman et al., 2008**
- 2 Tuchman, L. K., Slap, G. B., Britto, M. T., Transition to adult care: experiences and
3 expectations of adolescents with a chronic illness, *Child: Care, Health & Development*, 34,
4 557-63, 2008
- 5 **Tunis & Minshall, 2010**
- 6 Tunis, S. L., Minshall, M. E., Self-monitoring of blood glucose (SMBG) for type 2 diabetes
7 patients treated with oral anti-diabetes drugs and with a recent history of monitoring: cost-
8 effectiveness in the US, *Current Medical Research and Opinion*, 26, 151-62, 2010
- 9 **Tunis et al., 2010**
- 10 Tunis, S. L., Sauriol, L., Minshall, M. E., Cost effectiveness of insulin glargine plus oral
11 antidiabetes drugs compared with premixed insulin alone in patients with type 2 diabetes
12 mellitus in Canada, *Applied Health Economics and Health Policy*, 8, 267-80, 2010
- 13 **Turner et al., 2012**
- 14 Turner, Rebecca M., Davey, Jonathan, Clarke, Mike J., Thompson, Simon G., Higgins, Julian
15 P. T., Predicting the extent of heterogeneity in meta-analysis, using empirical data from the
16 Cochrane Database of Systematic Reviews, *International Journal of Epidemiology*, 41, 818-
17 827, 2012
- 18 **Van Gestel et al., 2010**
- 19 Van Gestel, Aukje, Severens, Johan L., Webers, Carroll A. B., Beckers, Henny J. M.,
20 Jansonius, Nomdo M., Schouten, Jan S. A. G., Modeling Complex Treatment Strategies:
21 Construction and Validation of a Discrete Event Simulation Model for Glaucoma, *Value in*
22 *Health*, 13, 358-367, 2010
- 23 **Van Koolwijk et al., 2002**
- 24 Van Koolwijk, L. M. E., Uiterwaal, C. S. P. M., Van der Laag, J., Hoekstra, J. H., Gulmans, V.
25 A. M., Van der Ent, C. K., Treatment of children with cystic fibrosis: Central, local or both?,
26 *Acta Paediatrica*, 91, 972-977, 2002
- 27 **Van Staa et al., 2011**
- 28 Van Staa, A. L., Jedeloo, S., van Meeteren, J., Latour, J. M., Crossing the transition chasm:
29 experiences and recommendations for improving transitional care of young adults, parents
30 and providers, *Child: Care, Health & Development*, 37, 821-32, 2011
- 31 **Van Valkenhoef et al., 2016**
- 32 Van Valkenhoef, Gert, Dias, Sofia, Ades, A. E., Welton, Nicky J., Automated generation of
33 node-splitting models for assessment of inconsistency in network meta-analysis, *Research*
34 *Synthesis Methods*, 7, 80-93, 2016
- 35 **Van Winden et al., 1998**
- 36 Van Winden, C. M., Visser, A., Hop, W., Sterk, P. J., Beckers, S., de Jongste, J. C., Effects
37 of flutter and PEP mask physiotherapy on symptoms and lung function in children with cystic
38 fibrosis, *European Respiratory Journal*, 12, 143-7, 1998
- 39 **Vieni et al., 2013**
- 40 Vieni, G., Faraci, S., Collura, M., Lombardo, M., Traverso, G., Cristadoro, S., Termini, L.,
41 Lucanto, M. C., Furnari, M. L., Trimarchi, G., Triglia, M. R., Costa, S., Pellegrino, S.,

- 1 Magazzu, G., Stunting is an independent predictor of mortality in patients with cystic fibrosis,
2 Clinical Nutrition, 32, 382-5, 2013
- 3 **Waine et al., 2007**
- 4 Waine, D. J., Whitehouse, J., Honeybourne, D., Cross-infection in cystic fibrosis: the
5 knowledge and behaviour of adult patients, Journal of Cystic Fibrosis, 6, 262-6, 2007
- 6 **Wainwright et al., 2011**
- 7 Wainwright, C. E., Vidmar, S., Armstrong, D. S., Byrnes, C. A., Carlin, J. B., Cheney, J.,
8 Cooper, P. J., Grimwood, K., Moodie, M., Robertson, C. F., Tiddens, H. A., Acfbal Study
9 Investigators, Effect of bronchoalveolar lavage-directed therapy on *P aeruginosa* infection
10 and structural lung injury in children with cystic fibrosis: a randomized trial, JAMA, 306, 163-
11 71, 2011
- 12 **Wainwright et al., 2011a**
- 13 Wainwright, C.E., Quittner, A.L., Geller, D.E., Nakamura, C., Wooldridge, J.L., Gibson, R.L.,
14 Lewis, S., Montgomery, A.B., Aztreonam for inhalation solution (AZLI) in patients with cystic
15 fibrosis, mild lung impairment, and *P aeruginosa*, Journal of Cystic Fibrosis, 10, 234-242,
16 2011
- 17 **Walters et al., 1994**
- 18 Walters, S., Britton, J., Hodson, M. E., Hospital care for adults with cystic fibrosis: an
19 overview and comparison between special cystic fibrosis clinics and general clinics using a
20 patient questionnaire, Thorax, 49, 300-6, 1994
- 21 **Wark & McDonald, 2009**
- 22 Wark, Peter, McDonald, Vanessa M., Nebulised hypertonic saline for cystic fibrosis,
23 Cochrane Database of Systematic Reviews, 2, CD001506, 2009
- 24 **Wark & McDonald, 2010**
- 25 Wark, Peter, McDonald, Vanessa M., Nebulised hypertonic saline for cystic fibrosis,
26 Cochrane Database of Systematic Reviews, 6, CD001506, 2010
- 27 **Warnock et al., 2013**
- 28 Warnock, L., Gates, A., van der Schans, C. P., Chest physiotherapy compared to no chest
29 physiotherapy for cystic fibrosis, Cochrane Database of Systematic Reviews, 9, CD001401,
30 2013
- 31 **Warwick et al., 2004**
- 32 Warwick, W. J., Wielinski, C. L., Hansen, L. G., Comparison of expectorated sputum after
33 manual chest physical therapy and high-frequency chest compression, Biomedical
34 Instrumentation & Technology, 38, 470-5, 2004
- 35 **Waters & Ratjen, 2014**
- 36 Waters, V., Ratjen, F., Antibiotic treatment for nontuberculous mycobacteria lung infection in
37 people with cystic fibrosis, Cochrane Database of Systematic Reviews, 12, CD010004, 2014
- 38 **Waters & Ratjen, 2016**
- 39 Waters, Valerie, Ratjen, Felix, Antibiotic treatment for nontuberculous mycobacteria lung
40 infection in people with cystic fibrosis, Cochrane Database of Systematic Reviews, 12,
41 CD010004, 2016

- 1 **Watson et al., 2008**
- 2 Watson, H., Bilton, D., Truby, H., A randomized controlled trial of a new behavioral home-
3 based nutrition education program, "Eat Well with CF," in adults with cystic fibrosis, *Journal*
4 of the American Dietetic Association, 108, 847-52, 2008
- 5 **Watts et al., 2009**
- 6 Watts, K. D., Seshadri, R., Sullivan, C., McColley, S. A., Increased prevalence of risk factors
7 for morbidity and mortality in the US Hispanic CF population, *Pediatric Pulmonology*, 44,
8 594-601, 2009
- 9 **Waugh et al., 2012**
- 10 Waugh, N., Royle, P., Craigie, I., Ho, V., Pandit, L., Ewings, P., Adler, A., Helms, P.,
11 Sheldon, C., Screening for cystic fibrosis-related diabetes: a systematic review, *Health*
12 Technology Assessment (Winchester, England), 16, 24, iii-iv, 1-179, 2012
- 13
- 14 **Weaver et al., 1994**
- 15 Weaver, L.T., Green, M.R., Nicholson, K., Mills, J., Heeley, M.E., Kuzemko, J.A., Austin, S.,
16 Gregory, G.A., Dux, A.E., Davis, J.A., Prognosis in cystic fibrosis treated with continuous
17 flucloxacillin from the neonatal period, *Archives of Disease in Childhood*, 70, 84-89, 1994
- 18 **Wells et al., 2008**
- 19 Wells, G. A., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., Tugwell, P., Ottawa
20 Hospital Research Institute, The Newcastle-Ottawa Scale (NOS) for assessing the quality of
21 nonrandomised studies in meta-analyses, 2008, Ottawa: Ottawa Hospital Research Institute
- 22 **Wesley et al., 1988**
- 23 Wesley, A. W., Qusted, C., Edgar, B. W., Lennon, D. R., A double-blind comparison of
24 ceftazidime with tobramycin and ticarcillin in the treatment of exacerbations of pseudomonas
25 chest infection in children with cystic fibrosis, 10th International Cystic Fibrosis Congress,
26 1988, [Abstract]
- 27 **White et al., 2013**
- 28 White, H., Morton, A. M., Conway, S. P., Peckham, D. G., Enteral tube feeding in adults with
29 cystic fibrosis; patient choice and impact on long term outcomes, *Journal of Cystic Fibrosis*,
30 12, 616-22, 2013
- 31 **White et al., 2014**
- 32 White, H., Denman, S., Shaw, N., Pollard, K., Peckham, D., Do longitudinal measures of
33 clinical variation correlate with adherence in cystic fibrosis, *Pediatric Pulmonology*, 49, 437,
34 2014
- 35 **Whiteford et al., 1995**
- 36 Whiteford, M. L., Wilkinson, J. D., McColl, J. H., Conlon, F. M., Michie, J. R., Evans, T. J.,
37 Paton, J. Y., Outcome of Burkholderia (Pseudomonas) cepacia colonisation in children with
38 cystic fibrosis following a hospital outbreak, *Thorax*, 50, 1194-8, 1995
- 39 **Whiting et al., 2014**
- 40 Whiting, P., Al, M., Burgers, L., Westwood, M., Ryder, S., Hoogendoorn, M., Armstrong, N.,
41 Allen, A., Severens, H., Kleijnen, J., Ivacaftor for the treatment of patients with cystic fibrosis

- 1 and the G551D mutation: a systematic review and cost-effectiveness analysis, Health
2 Technology Assessment (Winchester, England), 18, 1-106, 2014
- 3 **Whyte et al., 1995**
- 4 Whyte, D. A., Baggaley, S., Rutter, C., Chronic illness in childhood: A comparative study of
5 family support across four diagnostic groups, *Physiotherapy*, 81, 515-520, 1995
- 6 **Widerman, 2002**
- 7 Widerman, E., Communicating a diagnosis of cystic fibrosis to an adult: what physicians
8 need to know, *Behavioral Medicine*, 28, 45-52, 2002
- 9 **Widerman, 2003**
- 10 Widerman, E., Knowledge, interests and educational needs of adults diagnosed with cystic
11 fibrosis after age 18, *Journal of Cystic Fibrosis*, 2, 97-104, 2003
- 12 **Widerman, 2004**
- 13 Widerman, E., The experience of receiving a diagnosis of cystic fibrosis after age 20:
14 Implications for social work, *Social Work in Health Care*, 39, 415-433, 2004
- 15 **Wiedemann et al., 2001**
- 16 Wiedemann, B., Steinkamp, G., Sens, B., Stern, M., German Cystic Fibrosis Quality
17 Assurance, Group, The German cystic fibrosis quality assurance project: clinical features in
18 children and adults, *European Respiratory Journal*, 17, 1187-94, 2001
- 19 **Wilcock et al., 2015**
- 20 Wilcock, M. J., Ruddick, A., Gyi, K. M., Hodson, M. E., Renal diseases in adults with cystic
21 fibrosis: a 40 year single centre experience, *Journal of Nephrology*, 28, 585-91, 2015
- 22 **Wilkinson et al., 2008**
- 23 Wilkinson, O. M., Duncan-Skingle, F., Pryor, J. A., Hodson, M. E., A feasibility study of home
24 telemedicine for patients with cystic fibrosis awaiting transplantation, *Journal of Telemedicine
25 & Telecare*, 14, 182-5, 2008
- 26 **Wilmott et al., 1996**
- 27 Wilmott, R. W., Amin, R. S., Colin, A. A., DeVault, A., Dozor, A. J., Eigen, H., Johnson, C.,
28 Lester, L. A., McCoy, K., McKean, L. P., Moss, R., Nash, M. L., Jue, C. P., Regelman, W.,
29 Stokes, D. C., Fuchs, H. J., Aerosolized recombinant human DNase in hospitalized cystic
30 fibrosis patients with acute pulmonary exacerbations, *American Journal of Respiratory &
31 Critical Care Medicine*, 153, 1914-7, 1996
- 32 **Witters et al., 2009**
- 33 Witters, P., De Boeck, K., Dupont, L., Proesmans, M., Vermeulen, F., Servaes, R., Verslype,
34 C., Laleman, W., Nevens, F., Hoffman, I., Cassiman, D., Non-invasive liver elastography
35 (Fibroscan) for detection of cystic fibrosis-associated liver disease, *Journal of Cystic Fibrosis*,
36 8, 392-9, 2009
- 37 **Woestenbenk et al., 2013**
- 38 Woestenbenk, J. W., Castelijns, S. J. A. M., van der Ent, C. K., Houwen, R. H. J., Nutritional
39 intervention in patients with Cystic Fibrosis: A systematic review., *Journal of Cystic Fibrosis*,
40 12, 102-115, 2013
- 41 **Wolter et al., 1997**

- 1 Wolter, J. M., Bowler, S. D., Nolan, P. J., McCormack, J. G., Home intravenous therapy in
2 cystic fibrosis: a prospective randomized trial examining clinical, quality of life and cost
3 aspects, *European Respiratory Journal*, 10, 896-900, 1997
- 4 **Wolter et al., 2002**
- 5 Wolter, J., Seeney, S., Bell, S., Bowler, S., Masel, P., McCormack, J., Effect of long term
6 treatment with azithromycin on disease parameters in cystic fibrosis: a randomised trial,
7 *Thorax*, 57, 212-6, 2002
- 8 **Woodruff et al., 2016**
- 9 Woodruff, S. A., Sontag, M. K., Accurso, F. J., Sokol, R. J., Narkewicz, M. R., Prevalence of
10 elevated liver enzymes in children with cystic fibrosis diagnosed by newborn screen, *Journal*
11 *of Cystic Fibrosis*, no pagination, 2016
- 12 **Yang et al., 2016**
- 13 Yang, C., Chilvers, M., Montgomery, M., Nolan, S. J., Dornase alfa for cystic fibrosis,
14 *Cochrane Database of Systematic Reviews*, 4, CD001127, 2016
- 15 **Yi et al., 2003**
- 16 Yi, M. S., Britto, M. T., Wilmott, R. W., Kotagal, U. R., Eckman, M. H., Nielson, D. W.,
17 Kociela, V. L., Tsevat, J., Health values of adolescents with cystic fibrosis, *Journal of*
18 *Pediatrics*, 142, 133-40, 2003
- 19 **Young et al., 2008**
- 20 Young, A. C., Wilson, J. W., Kotsimbos, T. C., Naughton, M. T., Randomised placebo
21 controlled trial of non-invasive ventilation for hypercapnia in cystic fibrosis, *Thorax*, 63, 72-7,
22 2008
- 23 **Zhang et al., 2010**
- 24 Zhang, Z., Shoff, S. M., Lai, H. J., Incorporating genetic potential when evaluating stature in
25 children with cystic fibrosis, *Journal of Cystic Fibrosis*, 9, 135-42, 2010
- 26 **Zhang et al., 2013**
- 27 Zhang, Z., Lindstrom, M. J., Lai, H. J., Pubertal height velocity and associations with
28 prepubertal and adult heights in cystic fibrosis, *Journal of Pediatrics*, 163, 376-82, 2013
- 29

1 13 Glossary and abbreviations

Term	Definition
Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.
ACT	Airway clearance techniques help to clear mucous from the lungs reducing the risk of infection and improve lung function.
Allocation concealment	The process used to prevent advance knowledge of group assignment in a randomised controlled trial (RCT). The allocation process should be impervious to any influence by the individual making the allocation, by being administered by someone who is not responsible for recruiting participants.
Alveolus (Alveoli)	The specialised part of the lung where oxygen enters the blood and carbon dioxide can leave.
Antimicrobial prophylaxis	Antimicrobial drugs administered to those without symptoms or positive cultures with the intention of preventing future infection/colonisation.
Applicability	How well the results of a study or NICE evidence review can answer a clinical question or be applied to the population being considered.
APRI	AST to platelet ratio index (APRI) is a liver fibrosis test.
Arm (of a clinical study)	Subsection of individuals within a study who receive one particular intervention, for example placebo arm.
Association	Statistical relationship between 2 or more events, characteristics or other variables. The relationship may or may not be causal.
Attrition bias	Systematic differences between comparison groups in withdrawals or exclusion of participants from a study.
Available case analysis	Analysis of data that is available for participants at the end of follow-up.
BAL	Bronchoalveolar lavage. The bronchi and alveoli are washed via bronchoscope with a small amount of fluid which is then collected for analysis.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable) with which subsequent results are compared.
Before-and-after study	A study that investigates the effects of an intervention by measuring particular characteristics of a population both before and after taking the intervention, and assessing any change that occurs.
Bias	Influences on a study that can make the results look better or worse than they really are. (Bias can even make it look as if a treatment works when it does not.) Bias can occur by chance, deliberately or as a result of systematic errors in the design and execution of a study. It can also occur at different stages in the research process, for example during the collection, analysis, interpretation, publication or review of research data. For examples see selection bias, performance bias, information bias, confounding factor and publication bias.
Bilirubin	A substance formed in liver by the breakdown of haemoglobin and excreted in bile.
Bronchus (Bronchi)	Small airways in the lung.
Cardiopulmonary exercise testing	Measurement of the function of the heart and lungs at rest and during exercise.
Carer (caregiver)	Someone who looks after family, partners or friends in need of help because they are ill, frail or have a disability.
Case series	Report of a number of cases of a given disease, usually covering the course of the disease and the response to treatment. There is no comparison (control) group of patients.
Case-control study	A study to find out the cause(s) of a disease or condition. This is done by

Term	Definition
	comparing a group of patients who have the disease or condition (cases) with a group of people who do not have it (controls) but who are otherwise as similar as possible (in characteristics thought to be unrelated to the causes of the disease or condition). This means the researcher can look for aspects of their lives that differ to see if they may cause the condition. For example, a group of people with lung cancer might be compared with a group of people the same age that do not have lung cancer. The researcher could compare how long both groups had been exposed to tobacco smoke. Such studies are retrospective because they look back in time from the outcome to the possible causes of a disease or condition.
CFRD	Cystic fibrosis related diabetes.
Chronic pulmonary infection	Presence of pathogens on culture (colonisation) in the absence of worsening clinical symptoms/signs of respiratory disease. Antimicrobials may be administered to suppress or eradicate such pathogens with the intention of reducing future acute pulmonary exacerbations.
Cirrhosis	Progressive fibrous tissue overgrowth in an organ.
Clinical audit	A systematic process for setting and monitoring standards of clinical care. Whereas 'guidelines' define what the best clinical practice should be, 'audit' investigates whether best practice is being carried out. Clinical audit can be described as a cycle or spiral. Within the cycle there are stages that follow a systematic process of establishing best practice, measuring care against specific criteria, taking action to improve care and monitoring to sustain improvement. The spiral suggests that as the process continues, each cycle aspires to a higher level of quality.
Clinical effectiveness	How well a specific test or treatment works when used in the 'real world' (for example when used by a doctor with a patient at home), rather than in a carefully controlled clinical trial. Trials that assess clinical effectiveness are sometimes called management trials. Clinical effectiveness is not the same as efficacy.
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinician	A healthcare professional who provides patient care; for example a doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence based medicine databases including the Cochrane Database of Systematic Reviews (reviews of randomised controlled trials prepared by the Cochrane Collaboration).
Cohort study	A study with 2 or more groups of people – cohorts – with similar characteristics. One group receives a treatment, is exposed to a risk factor or has a particular symptom and the other group does not. The study follows their progress over time and records what happens.
Cohorting (cross-infection)	Grouping patients with positive cultures of the same pathogen(s).
Colonisation	Presence of pathogens on culture without signs of infection. See chronic pulmonary infection.
Community care	Care provided by community nurses, health visitors or school nurses of the region who look after people with CF and will administer treatments such as home IV antibiotics. It can include palliative services.
Comorbidity	A disease or condition that someone has in addition to the health problem being studied or treated.
Concealment of allocation	The process used to ensure that the person deciding to enter a participant into a randomised controlled trial does not know the comparison group into which that individual will be allocated. This is distinct from blinding and is aimed at preventing selection bias. Some attempts at concealing allocation are more prone to manipulation than others, and the method of allocation concealment is used as an assessment of the quality of a trial.

Term	Definition
Confidence interval (CI)	There is always some uncertainty in research. This is because a small group of patients is studied to predict the effects of a treatment on the wider population. The confidence interval is a way of expressing how certain we are about the findings from a study, using statistics. It gives a range of results that is likely to include the 'true' value for the population. The CI is usually stated as '95% CI', which means that the range of values has a 95 in 100 chance of including the 'true' value. For example, a study may state that "based on our sample findings, we are 95% certain that the 'true' population blood pressure is not higher than 150 and not lower than 110". In such a case the 95% CI would be 110 to 150. A wide confidence interval indicates a lack of certainty about the true effect of the test or treatment – often because a small group of patients has been studied. A narrow confidence interval indicates a more precise estimate (for example if a large number of patients have been studied).
Confounding factor	Something that influences a study and can result in misleading findings if it is not understood or appropriately dealt with. For example, a study of heart disease may look at a group of people who exercise regularly and a group who do not exercise. If the ages of the people in the 2 groups are different, then any difference in heart disease rates between the 2 groups could be because of age rather than exercise. Therefore age is a confounding factor.
Consensus methods	Techniques used to reach agreement on a particular issue. Consensus methods may be used to develop NICE guidance if there is not enough good quality research evidence to give a clear answer to a question. Formal consensus methods include Delphi and nominal group techniques.
Continuous outcome	Data with a potentially infinite number of possible values within a given range. Height, weight and blood pressure are examples of continuous variables.
Control group	A group of people in a study who do not receive the treatment or test being studied. Instead, they may receive the standard treatment (sometimes called 'usual care') or a dummy treatment (placebo). The results for the control group are compared with those for a group receiving the treatment being tested. The aim is to check for any differences. Ideally, the people in the control group should be as similar as possible to those in the treatment group, to make it as easy as possible to detect any effects due to the treatment.
Corticosteroids	Anti-inflammatory medicines.
Cost–benefit analysis (CBA)	Cost-benefit analysis is one of the tools used to carry out an economic evaluation. The costs and benefits are measured using the same monetary units (for example UK pounds) to see whether the benefits exceed the costs.
Cost–consequence analysis (CCA)	Cost–consequence analysis is one of the tools used to carry out an economic evaluation. This compares the costs (such as treatment and hospital care) and the consequences (such as health outcomes) of a test or treatment with a suitable alternative. Unlike cost–benefit analysis or cost-effectiveness analysis, it does not attempt to summarise outcomes in a single measure (like the quality adjusted life year) or in financial terms. Instead, outcomes are shown in their natural units (some of which may be monetary) and it is left to decision-makers to determine whether, overall, the treatment is worth carrying out
Cost-effectiveness analysis (CEA)	Cost-effectiveness analysis is one of the tools used to carry out an economic evaluation. The benefits are expressed in non-monetary terms related to health, such as symptom-free days, heart attacks avoided, deaths avoided or life years gained (that is, the number of years by which life is extended as a result of the intervention).
Cost-effectiveness model	An explicit mathematical framework, which is used to represent clinical decision problems and incorporate evidence from a variety of sources in

Term	Definition
	order to estimate the costs and health outcomes.
Cost–utility analysis (CUA)	Cost–utility analysis is one of the tools used to carry out an economic evaluation. The benefits are assessed in terms of both quality and duration of life, and expressed as quality adjusted life years (QALYs).
COX proportional hazard model	In survival analysis, a statistical model that asserts that the effect of the study factors (for example the intervention of interest) on the hazard rate (the risk of occurrence of an event) in the study population is multiplicative and does not change over time.
Credible interval (CrI)	The Bayesian equivalent of a confidence interval.
CT scan	Computerised tomography scan; computer-processed combinations of x-ray images producing cross-sectional images.
Decision analysis	An explicit quantitative approach to decision-making under uncertainty based on evidence from research. This evidence is translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Dichotomous outcomes	Outcome that can take 1 of 2 possible values, such as dead/alive, smoker/non-smoker, present/not present (also called binary data).
DIOS	Distal Intestinal Obstruction Syndrome. A blockage of the gut which occurs in older children and adults with cystic fibrosis.
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Dominance	A health economics term. When comparing tests or treatments, an option that is both less effective and costs more is said to be 'dominated' by the alternative.
Drop-out	A participant who withdraws from a trial before the end.
Economic evaluation	An economic evaluation is used to assess the cost effectiveness of healthcare interventions (that is, to compare the costs and benefits of a healthcare intervention to assess whether it is worth doing). The aim of an economic evaluation is to maximise the level of benefits – health effects – relative to the resources available. It should be used to inform and support the decision-making process; it is not supposed to replace the judgement of healthcare professionals. There are several types of economic evaluation: cost–benefit analysis, cost consequence analysis, cost-effectiveness analysis, cost-minimisation analysis and cost–utility analysis. They use similar methods to define and evaluate costs, but differ in the way they estimate the benefits of a particular drug, programme or intervention.
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	A measure that shows the magnitude of the outcome in 1 group compared with that in a control group. For example, if the absolute risk reduction is shown to be 5% and it is the outcome of interest, the effect size is 5%. The effect size is usually tested, using statistics, to find out how likely it is that the effect is a result of the treatment and has not just happened incidentally.
Effectiveness	How beneficial a test or treatment is under usual or everyday conditions, compared with doing nothing or opting for another type of care.
Efficacy	How beneficial a test, treatment or public health intervention is under ideal conditions (for example in a laboratory), compared with doing nothing or opting for another type of care.
Enteric coated	Covered with a coating which protects against acid in the stomach.
Epidemiological study	The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (for example

Term	Definition
	infection, diet) and interventions.
EQ-5D (EuroQol 5 dimensions)	A standardised instrument used to measure health-related quality-of-life. It provides a single index value for health status.
Equivalence study	A trial designed to determine whether the response to 2 or more treatments differs by an amount that is clinically unimportant. This is usually demonstrated by showing that the true treatment difference is likely to lie between a lower and an upper equivalence level of clinically acceptable differences.
Eradication regimen, antibiotic	An antibiotic regimen aimed at eliminating a specific pulmonary pathogen such as <i>S aureus</i> or <i>P aeruginosa</i> in people with cystic fibrosis
Evidence	Information on which a decision or guidance is based. Evidence is obtained from a range of sources including randomised controlled trials, observational studies, expert opinion (of clinical professionals or patients).
Exclusion criteria (clinical study)	Criteria that define who is not eligible to participate in a clinical study.
Exclusion criteria (literature review)	Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.
Extended dominance	If Option A is both more clinically effective than Option B and has a lower cost per unit of effect when both are compared with a do-nothing alternative, then Option A is said to have extended dominance over Option B. Option A is therefore more cost effective and should be preferred, other things remaining equal.
Extrapolation	An assumption that the results of studies of a specific population will also hold true for another population with similar characteristics.
Fibroscan	See Transient elastography
Fixed-effect model	In meta-analysis, a model that calculates a pooled effect estimate using the assumption that all observed variation between studies is caused by the play of chance. Studies are assumed to be measuring the same overall effect.
Follow-up	Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been assessed in order to observe changes in health status or health-related variables.
Forest plot	A graphical representation of the individual results of each study included in a meta-analysis together with the combined meta-analysis result. The plot also allows readers to see the heterogeneity among the results of the studies. The results of individual studies are shown as squares centred on each study's point estimate. A horizontal line runs through each square to show each study's confidence interval. The overall estimate from the meta-analysis and its confidence interval are shown at the bottom, represented as a diamond. The centre of the diamond represents the pooled point estimate, and its horizontal tips represent the confidence interval.
Forn's score	A non-invasive marker of liver fibrosis.
Generalisability	The extent to which the results of a study hold true for groups that did not participate in the research. See also external validity.
Gold standard	A method, procedure or measurement that is widely accepted as being the best available to test for or treat a disease.
GRADE, GRADE profile	A system developed by the GRADE Working Group to address the shortcomings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE profile.
<i>Haemophilus influenzae</i>	A bacterium which is a common cause of respiratory infection in cystic fibrosis.

Term	Definition
Haemoptysis	Coughing up blood.
Harms	Adverse effects of an intervention.
Hazard ratio	A hazard is the rate at which events happen, so that the probability of an event happening in a short time interval is the length of time multiplied by the hazard. Although the hazard may vary with time, the assumption in proportional hazard models for survival analysis is that the hazard in one group is a constant proportion of the hazard in the other group. This proportion is the hazard ratio.
Health economics	Study or analysis of the cost of using and distributing healthcare resources.
Health related quality of life (HRQoL)	A measure of the effects of an illness to see how it affects someone's day-to-day life.
Heterogeneity	The term is used in meta-analyses and systematic reviews to describe when the results of a test or treatment (or estimates of its effect) differ.
HFCWO	High frequency chest wall oscillation; technique aimed at improving airway clearance
Home care (e.g. hospital at home)	Giving care at home instead of a hospital, provided by the relevant cystic fibrosis specialist (such as a specialist nurse, dietician or psychologist).
Immunomodulatory dose	The use of a drug such as azithromycin prescribed at a lower dose than the minimum inhibitory dose.
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.
Inclusion criteria (literature review)	Explicit criteria used to decide which studies should be considered as potential sources of evidence.
Incremental cost	The extra cost linked to using one test or treatment rather than another. Or the additional cost of doing a test or providing a treatment more frequently.
Incremental cost effectiveness ratio (ICER)	The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another
Incremental net benefit (INB)	The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the INB is calculated as: (£20,000×QALYs gained) minus incremental cost.
Indirectness	The available evidence is different to the review question being addressed, in terms of population, intervention, comparison and outcome (PICO).
Intention-to-treat analysis (ITT)	An assessment of the people taking part in a clinical trial, based on the group they were initially (and randomly) allocated to. This is regardless of whether or not they dropped out, fully complied with the treatment or switched to an alternative treatment. Intention-to-treat analyses are often used to assess clinical effectiveness because they mirror actual practice: that is, not everyone complies with treatment and the treatment people receive may be changed according to how they respond to it.
Intervention	In medical terms this could be a drug treatment, surgical procedure, diagnostic or psychological therapy. Examples of public health interventions could include action to help someone to be physically active or to eat a more healthy diet.
Kappa statistic	A statistical measure of inter-rater agreement that takes into account the agreement occurring by chance.
LCI	Lung Clearance Index. A measure of lung function.
Length of stay	The total number of days a participant stays in hospital.

Term	Definition
Licence	See 'Product licence'.
Life years gained	Mean average years of life gained per person as a result of the intervention compared with an alternative intervention.
Likelihood ratio	The likelihood ratio combines information about the sensitivity and specificity. It tells you how much a positive or negative result changes the likelihood that a patient would have the disease. The likelihood ratio of a positive test result (LR+) is sensitivity divided by (1 minus specificity).
Loss to follow-up	Patients who have withdrawn from a clinical trial at the point of follow-up.
Malabsorption	A failure to absorb nutrients from the intestine. In cystic fibrosis this is due to the common occurrence of exocrine pancreatic insufficiency, so that there is a deficiency or absence of the pancreatic enzymes necessary to digest complex carbohydrates, proteins and fats (maldigestion) resulting in an inability to absorb these and other nutrients
Maldigestion	See malabsorption.
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
MDT	Multi-disciplinary team. A patient care team comprised of healthcare professionals of various specialties.
Mean	An average value, calculated by adding all the observations and dividing by the number of observations.
Mean difference	In meta-analysis, a method used to combine measures on continuous scales (such as weight), where the mean, standard deviation and sample size in each group are known. The weight given to the difference in means from each study (for example how much influence each study has on the overall results of the meta-analysis) is determined by the precision of its estimate of effect.
Meconium ileus	An obstruction of the small intestine at birth due to inspicated material in the gut lumen in infants with cystic fibrosis.
Meconium Ileus Equivalent	See DIOS
Median	The value of the observation that comes half-way when the observations are ranked in order.
Meta-analysis	A method often used in systematic reviews. Results from several studies of the same test or treatment are combined to estimate the overall effect of the treatment.
Microspheres	Enzyme granules contained within a pancreatin capsule.
Minimal important difference (MID)	Thresholds for clinical importance, which represent minimal important differences for benefit or for harm; for example the threshold at which drug A is less effective than drug B by an amount that is clinically important to patients.
MRI	Magnetic Resonance Imaging. Diagnostic imaging using magnetic fields and radio waves.
Mucoactive agent	See mucolytic agent.
Mucolytic agent	Drug affecting the viscosity of mucus, typically administered with the intention of making the removal of mucus through coughing easier.
Multivariate model	A statistical model for analysis of the relationship between 2 or more predictor (independent) variables and the outcome (dependent) variable.
Nebuliser	A small machine which converts liquid medication to a fine mist which can be breathed in to work directly in the lungs.
Net monetary benefit (NMB)	The value (usually in monetary terms) of an intervention net of its cost. The NMB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the NMB

Term	Definition
	is calculated as: (£20,000×QALYs gained) minus cost.
Network meta-analysis	Meta-analysis in which multiple treatments (that is, 3 or more) are being compared using both direct comparisons of interventions within randomised controlled trials and indirect comparisons across trials based on a common comparator.
Number needed to treat (NNT)	The average number of patients who need to be treated to get a positive outcome. For example, if the NNT is 4, then 4 patients would have to be treated to ensure 1 of them gets better. The closer the NNT is to 1, the better the treatment. For example, if you give a stroke prevention drug to 20 people before 1 stroke is prevented, the number needed to treat is 20.
Observational study	Individuals or groups are observed or certain factors are measured. No attempt is made to affect the outcome. For example, an observational study of a disease or treatment would allow 'nature' or usual medical care to take its course. Changes or differences in 1 characteristic (for example whether or not people received a specific treatment or intervention) are studied without intervening. There is a greater risk of selection bias than in experimental studies.
Odds ratio (OR)	Odds are a way to represent how likely it is that something will happen (the probability). An odds ratio compares the probability of something in 1 group with the probability of the same thing in another. An odds ratio of 1 between 2 groups would show that the probability of the event (for example a person developing a disease or a treatment working) is the same for both. An odds ratio greater than 1 means the event is more likely in the first group. An odds ratio less than 1 means that the event is less likely in the first group. Sometimes probability can be compared across more than 2 groups - in this case, 1 of the groups is chosen as the 'reference category', and the odds ratio is calculated for each group compared with the reference category. For example, to compare the risk of dying from lung cancer for non-smokers, occasional smokers and regular smokers, non-smokers could be used as the reference category. Odds ratios would be worked out for occasional smokers compared with non-smokers and for regular smokers compared with non-smokers. See also confidence interval, relative risk, risk ratio.
Odds ratio (OR)	Odds are a way to represent how likely it is that something will happen (the probability). An odds ratio compares the probability of something in 1 group with the probability of the same thing in another. An odds ratio of 1 between 2 groups would show that the probability of the event (for example a person developing a disease, or a treatment working) is the same for both. An odds ratio greater than 1 means the event is more likely in the first group. An odds ratio less than 1 means that the event is less likely in the first group. Sometimes probability can be compared across more than 2 groups – in this case, 1 of the groups is chosen as the 'reference category' and the odds ratio is calculated for each group compared with the reference category. For example, to compare the risk of dying from lung cancer for non-smokers, occasional smokers and regular smokers, non-smokers could be used as the reference category. Odds ratios would be worked out for occasional smokers compared with non-smokers and for regular smokers compared with non-smokers.
Opportunity cost	The loss of other healthcare programmes displaced by investment in or introduction of another intervention. This may be best measured by the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.
Outcome	The impact that a test, treatment, policy, programme or other intervention has on a person, group or population. Outcomes from interventions to improve the public's health could include changes in knowledge and behaviour related to health, societal changes (for example a reduction in crime rates) and a change in people's health and wellbeing or health status. In clinical terms, outcomes could include the number of patients

Term	Definition
	who fully recover from an illness or the number of hospital admissions, and an improvement or deterioration in someone's health, functional ability, symptoms or situation. Researchers should decide what outcomes to measure before a study begins.
Outreach care	A model of care in which the specialist multidisciplinary cystic fibrosis team provide outpatient clinics in local hospitals.
P value	The p value is a statistical measure that indicates whether or not an effect is statistically significant. For example, if a study comparing 2 treatments found that 1 seems more effective than the other, the p value is the probability of obtaining these results by chance. By convention, if the p value is below 0.05 (that is, there is less than a 5% probability that the results occurred by chance) it is considered that there probably is a real difference between treatments. If the p value is 0.001 or less (less than a 1% probability that the results occurred by chance), the result is seen as highly significant. If the p value shows that there is likely to be a difference between treatments, the confidence interval describes how big the difference in effect might be.
Pancreatin	An extract of animal pancreas; the general name for all pancreatic enzymes.
PEP	Positive expiratory pressure; a technique aimed at improving airway clearance.
Performance bias	Systematic differences between intervention groups in care provided apart from the intervention being evaluated. Blinding of study participants (both the recipients and providers of care) is used to protect against performance bias.
PERT	Pancreatic enzyme replacement therapy.
Placebo	A fake (or dummy) treatment given to participants in the control group of a clinical trial. It is indistinguishable from the actual treatment (which is given to participants in the experimental group). The aim is to determine what effect the experimental treatment has had – over and above any placebo effect caused because someone has received (or thinks they have received) care or attention.
Placebo effect	A beneficial (or adverse) effect produced by a placebo and not due to any property of the placebo itself.
Post-hoc analysis	Statistical analyses that are not specified in the trial protocol and are generally suggested by the data.
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Primary care	Healthcare delivered outside hospitals. It includes a range of services provided by GPs, nurses, health visitors, midwives and other healthcare professionals and allied health professionals such as dentists, pharmacists and opticians.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Product licence	An authorisation from the MHRA to market a medicinal product.
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good prognosis is associated with low rate of undesirable outcomes; poor prognosis is associated with a high rate of undesirable outcomes.
Prophylactic antibiotics	Antibiotics used for the prevention of infection complications.
Prospective study	A research study in which the health or other characteristic of participants is monitored (or 'followed up') for a period of time, with events recorded as they happen. This contrasts with retrospective studies.
<i>Pseudomonas</i>	A bacterial infection which affects the lungs.

Term	Definition
<i>aeruginosa</i>	
Publication bias	Publication bias occurs when researchers publish the results of studies showing that a treatment works well and don't publish those showing it did not have any effect. If this happens, analysis of the published results will not give an accurate idea of how well the treatment works. This type of bias can be assessed by a funnel plot.
Pulmonary exacerbation	The sudden or recent worsening of clinical symptoms or signs. This is frequently caused by a respiratory infection.
Pulmonary infection	The sudden or recent respiratory infection. In people with cystic fibrosis, this can be diagnosed based on new or worsening symptoms or signs, or by identifying pathogens in respiratory secretion samples.
Pyrexia	A fever.
Sepsis	A whole-body inflammation caused by an infection.
Shared-care (Network CF Clinic)	When a local hospital cares for people with cystic fibrosis, with oversight, support and direct involvement from members of a specialist cystic fibrosis centre.
Specialist centre	This is a centre for the diagnosis and management of cystic fibrosis, and working with a multidisciplinary team approach. Cystic fibrosis specialist centres are commissioned by NHS England.
Spirometry	A lung function test measuring the volume and/or speed of air that can be inhaled and exhaled.
Stakeholder	An organisation with an interest in a topic that NICE is developing a NICE guideline or piece of public health guidance on. Organisations that register as stakeholders can comment on the draft scope and the draft guidance. Stakeholders may be:
Standard deviation (SD)	A measure of the spread or dispersion of a set of observations, calculated as the average difference from the mean value in the sample.
<i>Staphylococcus aureus</i>	A bacterial infection that can affect the lungs.
Steatorrhoea	Abnormal amount of fat in faeces due to malabsorption.
Subgroup analysis	An analysis in which the intervention effect is evaluated in a defined subset of the participants in a trial, or in complementary subsets.
Suppression regimen, antibiotic	An antibiotic regimen aimed at reducing the level of infection with a specific pulmonary pathogen such as <i>S aureus</i> or <i>P aeruginosa</i> in people with cystic fibrosis
Systematic review (SR)	A review in which evidence from scientific studies has been identified, appraised and synthesised in a methodical way according to predetermined criteria. It may include a meta-analysis.
Telemedicine	Telemedicine refers to the provision of remote clinical services, via real-time two-way communication between the patient and the healthcare provider, using electronic audio and visual means.
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Transient elastography	An ultrasound technique to measure tissue stiffness.
Treatment allocation	Assigning a participant to a particular arm of a trial.
UDCA	Ursodeoxycholic acid; drug administered with the intention of preventing progression of liver disease.
Ultrasound	Imaging technique using ultrasound (high frequency sound waves).
Univariate	Analysis which separately explores each variable in a data set.
Utility	In health economics, a 'utility' is the measure of the preference or value that an individual or society places upon a particular health state. It is generally a number between 0 (representing death) and 1 (perfect health). The most widely used measure of benefit in cost–utility analysis is the

Term	Definition
	quality adjusted life year (QALY), but other measures include disability adjusted life years (DALYs) and healthy year equivalents (HYEs).

1