

Otitis media with effusion in under 12s

Supplement 3

NICE guideline tbc

Supplement 3: Other non-surgical interventions section - evidence from 2008 guideline

March 2023

Draft for Consultation

*These supplements were developed by
NICE*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2023 All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

| | |
|---|----------|
| Other non-surgical interventions section: evidence from 2008 guideline | 5 |
|---|----------|

1 **Other non-surgical interventions section:** 2 **evidence from 2008 guideline**

3

4 This supplement relates to recommendation 1.5.5 from the Other non-surgical
5 interventions section of the Otitis media with effusion in under 12s guideline, for
6 further details see the rationale and impact sections for that recommendation as well
7 as the Committee's discussion of the evidence section in Evidence review G:
8 Antibiotics.

9 The text below is extracted from the 2008 NICE guideline [CG60] Surgical
10 management of otitis media with effusion in children:

11

12 **3.3.4 Effectiveness of non-surgical interventions**

13 **Clinical question**

14 What is the effectiveness of various non-surgical interventions in children with OME?

15 Good-quality systematic reviews were available for evaluating the effectiveness of
16 steroids, antihistamines and/or decongestants, antibiotics and autoinflation, but not
17 for homeopathy or hearing aids. The two studies included for hearing aids are
18 surveys assessing the compliance and subjective improvement with their use, while
19 the single study included for homeopathy is a poorly conducted pilot RCT. No
20 published studies were identified for acupuncture, cranial osteopathy, dietary
21 modification, immunostimulants, massage or probiotics.

22 **Homeopathy**

23 **Description of included studies**

24 A pilot RCT¹ was carried out at two general practice (GP) centres in the UK to
25 determine whether homeopathic treatment of children with glue ear is more effective
26 than standard GP care. The study population comprised children aged 18 months to
27 8 years with a positive diagnosis of OME by the patient's GP, hearing loss > 20 dB
28 and an abnormal tympanogram. Children were randomised to the homeopathy or

1 standard GP care group, but the process of randomisation was not described and
2 there was no concealment of allocation or blinding of the participants. The standard
3 care involved 'watchful waiting' with autoinflation and in some cases a course of low-
4 dose antibiotics for 4–6 weeks. The sample size was small and the two groups were
5 not compared for baseline demographic characteristics except the age range.
6 Audiometry and tympanometry were conducted during the 1 year follow-up in this
7 study, and results were given without intention-to-treat analysis. [EL = 1–]

8 **Review findings**

9 A total of 33 children were randomised to either the homeopathic care group (n = 17)
10 or the standard care group (n = 16). Children in the two groups had similar age
11 ranges but there was a significant difference with regard to their initial hearing loss.
12 After 12 months of follow-up, a higher proportion of children in the homeopathic care
13 group had normal tympanograms and audiometric improvement (hearing loss < 20
14 dB), but the difference reached statistical significance only for improvement in
15 tympanograms (76.4% versus 31.3%; P = 0.01). Children in the intervention group
16 also had fewer courses of antibiotics in 12 months and fewer referrals to specialists,
17 but again the difference was not statistically significant.

18 **Evidence summary**

19 Results from a pilot trial show some improvement in tympanogram in children treated
20 with homeopathy after 12 months of follow-up compared with standard care, but
21 there was no benefit for the other outcomes.

22 **Translation from evidence to recommendations**

23 Homoeopathy, cranial osteopathy, acupuncture, dietary modification, massage,
24 immunostimulants and probiotics, although of potential interest as treatments, all lack
25 a published evidence base for effectiveness in OME.

26 **Recommendations on effectiveness of non-surgical interventions**

27 The following treatments are not recommended for the management of OME:

- 28 • antibiotics
- 29 • topical or systemic antihistamines
- 30 • topical or systemic decongestants

- 1 • topical or systemic steroids
- 2 • homeopathy
- 3 • osteopathy
- 4 • acupuncture
- 5 • dietary modification, including probiotics
- 6 • immunostimulants
- 7 • massage.

8

9 References

- 10 1) Harrison H. A randomized comparison of homoeopathic and standard care for
11 the treatment of glue ear in children. *Complementary Therapies in Medicine*
12 1999;7:132–5.