

Hip fracture: management

Consultation on research recommendations 18/01/16 to 15/02/17

Comments forms with attachments such as research articles, letters or leaflets cannot be accepted.

| ID | Туре | Stakeholder | Research rec. number | Should rec stand? (Yes/No) | Comments Please insert each new comment in a new row | Developer's response Please respond to each comment |
|----|------|---|----------------------------|----------------------------------|---|---|
| 1 | SH | The Royal College of Anaesthetists and the Association of Anaesthetists of Great Britain & Ireland – joint response | 1 | | No comment | |
| 2 | SH | Royal College of Nursing | 1 | Yes | This imaging recommendation may be a developmental recommendation for the way forward but may not be achievable in all acute trusts. Imaging is dependent on availability and variation in demand from other departments, but is an important component of diagnosing a hip fracture quickly to initiate treatment. | Thank you for your comment. We decided to retain this research recommendation based on feedback of its importance. |
| 3 | SH | | 2 | yes | Butthere are a couple of studies looking at this (notably REGAIN in the US). Cohort studies have repeatedly shown that it doesn't make a difference what type of anaesthesia is given. There are more pressing questions around for instance: Blood pressure management Perioperative transfusion Safety of tranexamic acid Utility of regional blocks (current evidence remains mixed) | Thank you for your responses. Although we can look to remove old research recommendations, we are not able to consider amendments to them as this area was not part of the scope of this guideline update. |

Comments received in the course of consultations carried out by NICE are published in the interests of openness and transparency, and to promote understanding of how recommendations are developed. The comments are published as a record of the submissions that NICE has received, and are not endorsed by NICE, its officers or advisory committees



Hip fracture: management

Consultation on research recommendations 18/01/16 to 15/02/17

Comments forms with attachments such as research articles, letters or leaflets cannot be accepted.

| ID | Туре | Stakeholder | Research rec. number | Should rec stand? (Yes/No) | Comments Please insert each new comment in a new row | Developer's response Please respond to each comment |
|----|------|---|----------------------------|----------------------------------|--|---|
| 3 | SH | The Royal College of Anaesthetists and the Association of Anaesthetists of Great Britain & Ireland – joint response | 2 | | Although this is from 2011, so no comments should be made, the research statement now looks dated. More specific studies should focus on: the control of blood pressure during hip fracture surgery. avoidance of delirium, by using anaesthesia techniques that do not include sedation focused risk assessment and use of HDU/ITU | Thank you for your comment. We decided to retain this research recommendation based on feedback of its importance. Although we can look to remove old research recommendations, we are not able to consider amendments to them as this area was not part of the scope of this guideline update. |
| 4 | SH | Royal College of Nursing | 2 | Yes | There remains a wide variation in practice and to ensure high quality patient care this needs further research. | Thank you for your comment. We decided to retain this research recommendation based on feedback of its importance |
| 5 | SH | The Royal College of Anaesthetists and the Association of Anaesthetists of Great Britain & Ireland – joint response | 4 | | No comment | |
| | SH | Royal College of Nursing | 4 | Yes | Further research is needed in this field but should physiotherapy and-Occupational Therapy be separate considerations for research? | Thank you for your comment. We decided to retain this research recommendation based on feedback of its importance Although we can look to remove old research recommendations, we are not able to consider amendments to |

Comments received in the course of consultations carried out by NICE are published in the interests of openness and transparency, and to promote understanding of how recommendations are developed. The comments are published as a record of the submissions that NICE has received, and are not endorsed by NICE, its officers or advisory committees



Hip fracture: management

Consultation on research recommendations 18/01/16 to 15/02/17

Comments forms with attachments such as research articles, letters or leaflets cannot be accepted.

| ID | Туре | Stakeholder | Research rec. number | Should rec stand? (Yes/No) | Comments Please insert each new comment in a new row | Developer's response Please respond to each comment |
|----|------|---|----------------------------|----------------------------------|--|--|
| | | | | | | them as this area was not part of the scope of this guideline update. |
| 6 | SH | The Royal College of Anaesthetists and the Association of Anaesthetists of Great Britain & Ireland – joint response | 5 | | No comment | |
| 7 | SH | Royal College of Nursing | 5 | Yes | Early supported discharge requires research to advance current thinking and practices. | Thank you for your comment. We decided to retain this research recommendation based on feedback of its importance |

Comments received in the course of consultations carried out by NICE are published in the interests of openness and transparency, and to promote understanding of how recommendations are developed. The comments are published as a record of the submissions that NICE has received, and are not endorsed by NICE, its officers or advisory committees