Appendix 23b: 2009 Access and engagement – secondary subgroup analyses

| Table 1: Studies included in the secondary subgroup analyses | 2 |
|---|---|
| Table 2: ACT versus standard care – 9 months data (REACT) | 2 |
| Table 3: ACT versus standard care – 18 months data (REACT) | 3 |
| Table 4: Crisis intervention team versus standard care – 8 weeks data (Crisis RCT) | 4 |
| Table 5: Crisis intervention team versus standard care – 6 months data (Crisis RCT) | 5 |

Table 1: Studies included in the secondary subgroup analyses

| Treatment | versus Comparator |
|--------------------------|---|
| | Standard care |
| ACT | REACT (Killaspy et al., 2006) |
| | |
| Crisis intervention team | North Islington Crisis RCT (Johnson <i>et al.</i> , 2005) |
| | |

Table 2: ACT versus standard care - primary and secondary outcomes at 9 months (REACT)

| | Whole group | | | BME subgroup | | |
|-----------------------|---|----------------|-----|--------------|---------|----|
| Outcome | ACT | Control | | ACT | Control | |
| Consent to | | | | | | |
| treatment (N) | | | | | | |
| Yes | 90 | 78 | | 37 | 26 | |
| No | 24 | 27 | | 14 | 10 | |
| No response | 13 | 19 | | 4 | 9 | |
| Lost to follow up | | | | | | |
| Yes (any reason) | 6 | 16 | * | 5 | 6 | |
| No | 121 | 108 | * | 50 | 39 | |
| Number of times | | | | | | |
| seen by MH | | | | | | |
| worker | | | | | | |
| Median | 31.00 | 10.00 | ** | 35.00 | 10.50 | ** |
| N | 124 | 119 | | 53 | 44 | |
| Time since last | | | | | | |
| seen by MH | | | | | | |
| worker (days) | | | | | | |
| Median | 5.00 | 15.00 | ** | 5.00 | 14.50 | ** |
| N | 124 | 119 | | 53 | 53 | |
| Inpatient days | | | | | | |
| Median | 63.50 | 45.00 | | 101.00 | 52.50 | |
| N | 124 | 119 | | 53 | 44 | |
| No. new | | | | | | |
| admissions | | | | | | |
| Mean | 0.52 | 0.53 | | 0.57 | 0.39 | |
| SD | 0.70 | 0.93 | | 0.72 | 0.87 | |
| Median | 0.00 | 0.00 | | 0.00 | 0.00 | |
| N | 124 | 119 | | 53 | 44 | |
| No. days/admission | | | | | | |
| Mean | 33.57 | 23.99 | | 39.62 | 22.50 | |
| SD | 56.22 | 50.37 | | 61.40 | 52.62 | |
| Median | 0.00 | 0.00 | | 0.00 | 0.00 | |
| N | 124 | 119 | | 53 | 44 | |
| Detentions under | 124 | 117 | | | | |
| MHA | | | | | | |
| Mean | 0.32 | 0.31 | | 0.36 | 0.26 | |
| SD | 0.548 | 0.61 | | 0.54 | 0.56 | |
| Median | 0.00 | 0.00 | | 0.00 | 0.00 | |
| N | 100 | 94 | | 39 | 35 | |
| 37 (4 : 1: (: | ::::::::::::::::::::::::::::::::::::::: | lastrona ACT a | 1 . | 1 O.O.E | | |

Note: * indicates a significant difference between ACT and control group, p<0.05.

^{**} indicated a significant difference between ACT and control group, p<0.01.

Table 3: ACT versus standard care - primary and secondary outcomes at 18 months (REACT)

| | | le group | | | ıbgroup | |
|--|--------|----------|----|--------|---------|---|
| Outcome | ACT | Control | | ACT | Control | |
| Lost to follow up (N) | | | | | | |
| Yes (any reason) | 5 | 15 | * | 3 | 4 | |
| No | 122 | 109 | * | 52 | 41 | |
| EAS | | | | | | |
| Mean | 9.07 | 8.04 | * | 8.57 | 8.32 | |
| SD | 3.35 | 3.76 | | 3.51 | 3.26 | |
| Median | 9.00 | 8.00 | | 8.00 | 8.00 | |
| N | 124 | 119 | | 53 | 44 | |
| Number of times seen by MH worker | | | | | | |
| Median | 31.00 | 12.00 | ** | 36.00 | 9.50 | • |
| N | 124 | 119 | | 53 | 44 | |
| Time since last seen by MH worker (days) | | | | | | |
| Median | 5.00 | 16.00 | ** | 7.00 | 15.50 | • |
| N | 124 | 119 | | 53 | 44 | |
| Inpatient days | | | | | | |
| Mean | 162.19 | 143.82 | | 198.04 | 155.32 | |
| SD | 161.55 | 140.62 | | 163.10 | 156.65 | |
| Median | 120.00 | 130.00 | | 152.00 | 137.00 | |
| N | 124 | 119 | | 53 | 44 | |
| No. new admissions | | | | | | |
| Mean | 1.00 | 1.12 | | 1.08 | 0.86 | |
| SD | 1.14 | 1.43 | | 1.17 | 1.32 | |
| Median | 1.00 | 1.00 | | 1.00 | 0.00 | |
| No. | 124 | 119 | | 53 | 44 | |
| days/admission | | | | | | |
| Mean | 64.79 | 52.00 | | 78.02 | 48.05 | |
| SD | 91.01 | 66.17 | | 105.34 | 67.45 | |
| Median | 27.00 | 27.00 | | 40.00 | 0.00 | |
| N | 124 | 119 | | 53 | 44 | |
| Detentions under MHA | | | | | | |
| Mean | 0.66 | 0.70 | | 0.79 | 0.59 | |
| SD | 0.785 | 0.98 | | 0.79 | 1.00 | |
| Median | 0.00 | 0.00 | | 1.00 | 0.00 | |
| N | 124 | 119 | | 53 | 44 | |

^{**} indicated a significant difference between ACT and control group, p<0.01.

Table 4: Crisis intervention team versus standard care – primary and secondary outcomes at 8 weeks (Crisis RCT)

| | Whole group | | | BME subgroup | | | |
|-------------------------------|-------------|---------|----|--------------|---------|----|--|
| Outcome | Crisis team | Control | | Crisis team | Control | | |
| Compliance with treatment (N) | | | | | | | |
| Fully cooperative | 59 | 57 | | 10 | 14 | | |
| Moderately | | | | 10 | | | |
| cooperative | 40 | 35 | | 6 | 10 | | |
| Moderately uncooperative | 16 | 13 | | 3 | 4 | | |
| Very | | | | | | | |
| uncooperative | 4 | 5 | | 2 | 1 | | |
| Not receiving a | | | | | | | |
| service | | | | | | | |
| n/N | 18/135 | 13/125 | | 7/27 | 3/32 | | |
| Number of | · | | | | | | |
| contact with MH | | | | | | | |
| workers | | | | | | | |
| Median | 14.00 | 8.00 | ** | 11.00 | 8.50 | ^ | |
| N | 135 | 125 | | 27 | 32 | | |
| Number of GP | | | | | | | |
| contacts | | | | | | | |
| (including those | | | | | | | |
| with no contact) | | | | | | | |
| Median | 1.00 | 0.00 | ** | 1.00 | 0.00 | * | |
| N | 135 | 125 | | 27 | 32 | | |
| Inpatient admission | | | | | | | |
| Yes | 29 | 74 | ** | 5 | 19 | ** | |
| No | 106 | 51 | ** | 22 | 13 | ** | |
| Number of admission since | | | | | | | |
| crisis | | | | | | | |
| Median | 0.00 | 1.00 | ** | 0.00 | 1.00 | * | |
| N | 135 | 135 | | 27 | 32 | | |
| Total inpatient | | | | | | | |
| days | | | | | | | |
| Median | 0.00 | 5.00 | ** | 0.00 | 18.50 | * | |
| N | 135 | 125 | | 27 | 32 | | |
| Subject to MHA | | | | | | | |
| Assessment | 100 | 00 | | 20 | 22 | | |
| No Yes | 109 | 92 | | 20 | 23 | | |
| Yes | 24 | 32 | | 7 | 9 | | |

Note: * indicates a significant difference between crisis and control group, p<0.05.

^{**} indicated a significant difference between crisis and control group, p<0.01.

[^] Lack of significance is likely to be due to loss of power within the BME subgroup.

Table 5: Crisis intervention team versus standard care – primary and secondary outcomes at 6 months (Crisis RCT)

| | Whole group | | | BME subgroup | | | |
|---------------------------|-------------|---------|----|--------------|---------|----|--|
| Outcome | Crisis team | Control | | Crisis team | Control | | |
| Compliance with | | | | | | | |
| treatment (N) | | | | | | | |
| Fully cooperative | 51 | 57 | | 8 | 19 | | |
| Moderately | 1.0 | | | | | | |
| cooperative | 46 | 32 | | 9 | 7 | | |
| Moderately | 10 | 0 | | 2 | 1 | | |
| uncooperative | 12 | 8 | | 2 | 1 | | |
| Very | 2 | 2 | | 1 | 1 | | |
| uncooperative | ۷ | ۷ | | 1 | 1 | | |
| Not receiving a | | | | | | | |
| service | | | | | | | |
| n/N | 24/135 | 20/125 | | 7/27 | 3/32 | | |
| Number of | | | | | | | |
| contact with MH | | | | | | | |
| workers | | | | | | | |
| Median | 25.00 | 16.00 | ** | 32.00 | 20.50 | ^ | |
| N | 135 | 125 | | 27 | 32 | | |
| Number of GP | | | | | | | |
| contacts | | | | | | | |
| Median | 4.00 | 3.00 | | 3.50 | 2.00 | | |
| N | 83 | 72 | | 16 | 17 | | |
| Number of GP | | | | | | | |
| contacts | | | | | | | |
| (including those | | | | | | | |
| with no contact) | 2.00 | 2.00 | * | 2.00 | 1.00 | ^ | |
| Median | 3.00 | 2.00 | | 2.00 | 1.00 | /\ | |
| N | 101 | 107 | | 21 | 27 | | |
| Inpatient admission | | | | | | | |
| | 20 | 0.4 | ** | 11 | 20 | ^ | |
| Yes | 39 | 84 | ** | 11 | 20 | ^ | |
| No | 96 | 41 | | 20 | 12 | | |
| Number of admission since | | | | | | | |
| crisis | | | | | | | |
| Median | 0.00 | 1.00 | ** | 0.00 | 1.00 | * | |
| N | 134 | 124 | | 27 | 32 | | |
| | 134 | 124 | | | 32 | | |
| Total inpatient days | | | | | | | |
| Median | 0.00 | 11.00 | ** | 0.00 | 23.50 | * | |
| N | 135 | 125 | | 27 | 32 | | |
| Total number of | 155 | 120 | | | 32 | | |
| days on section | | | | | | | |
| Mean | 9.90 | 13.04 | | 15.59 | 11.78 | | |
| SD | 31.51 | 32.95 | | 32.80 | 26.07 | | |
| Median | 0.00 | 0.00 | | 0.00 | 0.00 | | |
| | 134 | 125 | | 27 | 32 | | |
| N | 134 | 125 | | 21 | 32 | | |

Note: * indicates a significant difference between crisis and control group, p<0.05.

^{**} indicated a significant difference between crisis and control group, p<0.01.

[^] Lack of significance is likely to be due to loss of power within the BME subgroup.