

# Obesity prevention

Clinical guideline

Published: 13 December 2006

[nice.org.uk/guidance/cg43](https://www.nice.org.uk/guidance/cg43)

## Your responsibility

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals and practitioners are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or the people using their service. It is not mandatory to apply the recommendations, and the guideline does not override the responsibility to make decisions appropriate to the circumstances of the individual, in consultation with them and their families and carers or guardian.

Local commissioners and providers of healthcare have a responsibility to enable the guideline to be applied when individual professionals and people using services wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with complying with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

## Contents

Overview .....	5
Who is it for? .....	5
Introduction .....	6
Rationale for public health guidance.....	6
Working with people to prevent and manage overweight and obesity: the issues .....	8
Working with children and young adults.....	8
Person-centred care: principles for health professionals .....	10
Key priorities for implementation .....	11
Public health .....	11
1 Guidance .....	13
1.1 Public health recommendations .....	13
1.2 Clinical recommendations .....	27
2 Notes on the scope of the guidance .....	28
3 Implementation .....	30
3.1 The NHS.....	30
3.2 Other audiences and settings.....	30
4 Research recommendations .....	31
4.1 What are the most effective interventions to prevent or manage obesity in children and adults in the UK?.....	31
4.2 How does the effectiveness of interventions to prevent or manage obesity vary by population group, setting and source of delivery?.....	31
4.3 What is the cost effectiveness of interventions to prevent or manage obesity in children and adults in the UK? .....	32
4.4 What elements make an intervention effective and sustainable, and what training do staff need?.....	32
4.5 Evaluation and monitoring.....	33
5 Other versions of this guideline.....	35
5.1 Full guideline.....	35
6 Related NICE guidance.....	36

7 Updating the guideline .....	37
Appendix A: The Guidance Development Groups .....	38
Public health.....	38
Clinical management .....	39
Guidance Development Group co-optees .....	40
National Institute for Health and Clinical Excellence .....	41
Obesity Collaborating Centre – Cardiff.....	42
Obesity Collaborating Centre – University of Teesside.....	43
External Collaborators – University of York.....	43
National Collaborating Centre for Primary Care (NCC-PC).....	43
Appendix B: The Guideline Review Panel.....	45
Appendix C: The algorithms.....	46
Appendix D: Existing guidance on diet, physical activity and preventing obesity.....	47
Changes after publication.....	48
About this guideline .....	50

This guideline replaces TA22, TA31 and TA46.

This guideline is partially replaced by PH53, CG189 and NG7.

This guideline is the basis of QS111 and QS94.

This guideline should be read in conjunction with PH13.

## Overview

This guideline covers preventing children, young people and adults becoming overweight or obese. It outlines how the NHS, local authorities, early years' settings, schools and workplaces can increase physical activity levels and make dietary improvements among their target populations.

The clinical management of obesity is covered by NICE's guideline on [obesity: identification, assessment and management](#).

NICE has also produced guidelines on [preventing excess weight gain](#) and [weight management: lifestyle services for overweight or obese adults](#).

## Who is it for?

- Commissioners and providers
- Employers
- Local authorities
- Primary care trusts
- Healthcare professionals
- Head teachers and chairs of governors
- Children, young people and adults, and their families and carers

## Introduction

This is the first national guidance on the prevention of overweight and obesity in adults and children in England and Wales. The guidance aims to:

- stem the rising prevalence of obesity and diseases associated with it
- increase the effectiveness of interventions to prevent overweight and obesity
- improve the care provided to adults and children at risk of overweight and obesity.

The recommendations are based on the best available evidence of effectiveness, including cost effectiveness. The advice on the prevention of overweight and obesity applies in both NHS and non-NHS settings.

The guidance supports the implementation of the 'Choosing health' White Paper in England, 'Designed for life' in Wales, the revised GP contract and the existing national service frameworks (NSFs). It also supports the joint Department of Health, Department for Education and Skills and Department for Culture, Media and Sport target to halt the rise in obesity among children under 11 by 2010, and similar initiatives in Wales.

### *Rationale for public health guidance*

Public health and clinical audiences share the same need for evidence-based, cost-effective solutions to the challenges in their day-to-day practice, as well as to inform policies and strategies to improve health. Complementary clinical and public health guidance are essential to address the hazy divisions between prevention and management of obesity.

The 2004 Wanless report 'Securing good health for the whole population' stressed that a substantial change will be needed to produce the reductions in preventable diseases such as obesity that will lead to the greatest reductions in future healthcare costs. In addition to recommending a more effective delivery framework for health services providers, the report proposed an enhanced role for schools, local authorities and other public sector agencies, employers, and private and voluntary sector providers in developing opportunities for people to secure better health.

It is unlikely that the problem of obesity can be addressed through primary care management alone. More than half the adult population are overweight or obese and a large proportion will need help with weight management. Although there is no simple solution, the most effective strategies

for prevention and management share similar approaches. The clinical management of obesity cannot be viewed in isolation from the environment in which people live.

NICE continues to recognise the importance of an integrated approach to the prevention, identification, assessment and management of obesity, as shown in the [obesity](#) pathway.

## Working with people to prevent and manage overweight and obesity: the issues

Preventing and managing overweight and obesity are complex problems, with no easy answers. This guidance offers practical recommendations based on the evidence. But staff working directly with the public also need to be aware of the many factors that could be affecting a person's ability to stay at a healthy weight or succeed in losing weight.

- People choose whether or not to change their lifestyle or agree to treatment. Assessing their readiness to make changes affects decisions on when or how to offer any intervention.
- Barriers to lifestyle change should be explored. Possible barriers include:
  - lack of knowledge about buying and cooking food, and how diet and exercise affect health
  - the cost and availability of healthy foods and opportunities for exercise
  - safety concerns, for example about cycling
  - lack of time
  - personal tastes
  - the views of family and community members
  - low levels of fitness, or disabilities
  - low self-esteem and lack of assertiveness.
- Advice needs to be tailored for different groups. This is particularly important for people from black and minority ethnic groups, vulnerable groups (such as those on low incomes) and people at life stages with increased risk for weight gain (such as during and after pregnancy, at the menopause or when stopping smoking).

### *Working with children and young adults*

- Treating children for overweight or obesity may stigmatise them and put them at risk of bullying, which in turn can aggravate problem eating. Confidentiality and building self-esteem are particularly important if help is offered at school.



- Interventions to help children eat a healthy diet and be physically active should develop a positive body image and build self-esteem.

## Person-centred care: principles for health professionals

When working with people to prevent or manage overweight and obesity, health professionals should follow the usual principles of person-centred care.

Advice, treatment and care should take into account people's needs and preferences. People should have the opportunity to make informed decisions about their care and treatment, in partnership with their health professionals.

Good communication between health professionals and people is essential. It should be supported by evidence-based written information tailored to the person's needs. Advice, treatment and care, and the information people are given about it, should be non-discriminatory and culturally appropriate. It should also be accessible to people with additional needs such as physical, sensory or learning disabilities, and to people who do not speak or read English.

For older children who are overweight or obese, a balance needs to be found between the importance of involving parents and the right of the child to be cared for independently.

If people do not have the capacity to make decisions, healthcare professionals should follow the [Department of Health's advice on consent](#) and the [code of practice that accompanies the Mental Capacity Act](#). In Wales, healthcare professionals should follow [advice on consent from the Welsh Government](#).

If the person is under 16, healthcare professionals should follow the guidelines in the Department of Health's [Seeking consent: working with children](#).

.

## Key priorities for implementation

The prevention and management of obesity should be a priority for all, because of the considerable health benefits of maintaining a healthy weight and the health risks associated with overweight and obesity.

### *Public health*

#### NHS

- Managers and health professionals in all primary care settings should ensure that preventing and managing obesity is a priority, at both strategic and delivery levels. Dedicated resources should be allocated for action.

#### Local authorities and partners

- Local authorities should work with local partners, such as industry and voluntary organisations, to create and manage more safe spaces for incidental and planned physical activity, addressing as a priority any concerns about safety, crime and inclusion, by:
  - providing facilities and schemes such as cycling and walking routes, cycle parking, area maps and safe play areas
  - making streets cleaner and safer, through measures such as traffic calming, congestion charging, pedestrian crossings, cycle routes, lighting and walking schemes
  - ensuring buildings and spaces are designed to encourage people to be more physically active (for example, through positioning and signing of stairs, entrances and walkways)
  - considering in particular people who require tailored information and support, especially inactive, vulnerable groups.

#### Early years settings

- Nurseries and other childcare facilities should:
  - minimise sedentary activities during play time, and provide regular opportunities for enjoyable active play and structured physical activity sessions
  - implement Department for Education and Skills, Food Standards Agency and Caroline Walker Trust<sup>[1]</sup> guidance on food procurement and healthy catering.

## Schools

- Head teachers and chairs of governors, in collaboration with parents and pupils, should assess the whole school environment and ensure that the ethos of all school policies helps children and young people to maintain a healthy weight, eat a healthy diet and be physically active, in line with existing standards and guidance. This includes policies relating to building layout and recreational spaces, catering (including vending machines) and the food and drink children bring into school, the taught curriculum (including PE), school travel plans and provision for cycling, and policies relating to the National Healthy Schools Programme and extended schools.

## Workplaces

- Workplaces should provide opportunities for staff to eat a healthy diet and be physically active, through:
  - active and continuous promotion of healthy choices in restaurants, hospitality, vending machines and shops for staff and clients, in line with existing Food Standards Agency guidance
  - working practices and policies, such as active travel policies for staff and visitors
  - a supportive physical environment, such as improvements to stairwells and providing showers and secure cycle parking
  - recreational opportunities, such as supporting out-of-hours social activities, lunchtime walks and use of local leisure facilities.

## Self-help, commercial and community settings

- Primary care organisations and local authorities should recommend to patients, or consider endorsing, self-help, commercial and community weight management programmes only if they follow best practice (see recommendation 1.1.7.1 for details of best practice standards).

---

<sup>[1]</sup> [Caroline Walker Trust](#)

## 1 Guidance

The following guidance is based on the best available evidence. The [full guideline](#) gives details of the methods and the evidence used to develop the guidance (see section 5 for details).

In the recommendations, 'children' refers to anyone younger than 18 years. 'Young people' is used when referring to teenagers at the older end of this age group.

Staff who advise people on diet, weight and activity – both inside and outside the NHS – need appropriate training, experience and enthusiasm to motivate people to change. Some will need general training (for example, in health promotion), while those who provide interventions for obesity (such as dietary treatment and physical training) will need more specialised training. In the recommendations, the term 'specific' is used if the training will be in addition to staff's basic training. The term 'relevant' is used for training that could be part of basic professional training or in addition to it.

### 1.1 *Public health recommendations*

The public health recommendations are divided according to their key audiences and the settings they apply to:

- the public
- the NHS
- local authorities and partners in the community
- early years settings
- schools
- workplaces
- self-help, commercial and community programmes.

Some of the recommendations are at a strategic level (primarily for those involved in planning and management of service provision and policies), and others are at delivery level (for individual staff, teams and team managers).

Section 3 has information about the status of NICE guidance in different settings, and links to tools to help with implementing the recommendations and meeting training needs. In many cases, implementation will involve organisations working in partnership.

### 1.1.1 Recommendations for the public

This section has been replaced by [Maintaining a healthy weight and preventing excess weight gain among adults and children](#) (2015) NICE guideline NG7.

### 1.1.2 The NHS

The following recommendations are made specifically for health professionals and managers in the NHS, but may also be relevant to health professionals in other organisations. Recommendations in other sections may also be relevant for NHS health professionals working with local authorities and other organisations.

These recommendations are for:

- senior managers, GPs, commissioners of care and directors of public health
- staff in primary and secondary care, particularly those providing interventions, including public health practitioners, nurses, behavioural psychologists, physiotherapists, GPs, pharmacists, trained counsellors, registered dietitians, public health nutritionists and specifically trained exercise specialists.

With specific training, staff such as pharmacy assistants or support staff in general practices may also be able to give advice and support.

Implementing these recommendations will contribute to the English target to halt the annual rise in obesity in children younger than 11 years by 2010, and similar initiatives in Wales. Recommendations can be delivered through local strategic partnerships and other local agreements and partnerships.

Section 3 has links to tools to help with implementing the recommendations and meeting training needs. In many cases, implementation will involve organisations working together in partnership.

Primary care staff should engage with target communities, consult on how and where to deliver interventions and form key partnerships and ensure that interventions are person centred.

Tailoring advice to address potential barriers (such as cost, personal tastes, availability, time, views of family and community members) is particularly important for people from black and minority ethnic groups, people in vulnerable groups (such as those on low incomes) and people at life stages with increased risk for weight gain (such as during and after pregnancy, menopause or smoking cessation). Many of the recommendations below also highlight the need to provide ongoing support – this can be in person, or by phone, mail or internet as appropriate.

### ***Overarching recommendation***

1.1.2.1 Managers and health professionals in all primary care settings should ensure that preventing and managing obesity is a priority at both strategic and delivery levels. Dedicated resources should be allocated for action.

### ***Strategy: for senior managers and budget holders***

1.1.2.2 In their role as employers, NHS organisations should set an example in developing public health policies to prevent and manage obesity by following existing guidance and (in England) the local obesity strategy. In particular:

- on-site catering should promote healthy food and drink choices (for example by signs, posters, pricing and positioning of products)
- there should be policies, facilities and information that promote physical activity, for example, through travel plans, by providing showers and secure cycle parking and by using signposting and improved décor to encourage stair use.

1.1.2.3 All primary care settings should ensure that systems are in place to implement the local obesity strategy. This should enable health professionals with specific training, including public health practitioners working singly and as part of multidisciplinary teams, to provide interventions to prevent and manage obesity.

1.1.2.4 All primary care settings should:

- address the training needs of staff involved in preventing and managing obesity
- allocate adequate time and space for staff to take action
- enhance opportunities for health professionals to engage with a range of organisations and to develop multidisciplinary teams.

1.1.2.5 Local health agencies should identify appropriate health professionals and ensure that they receive training in:

- the health benefits and the potential effectiveness of interventions to prevent obesity, increase activity levels and improve diet (and reduce energy intake)
- the best practice approaches in delivering such interventions, including tailoring support to meet people's needs over the long term
- the use of motivational and counselling techniques.

Training will need to address barriers to health professionals providing support and advice, particularly concerns about the effectiveness of interventions, people's receptiveness and ability to change and the impact of advice on relationships with patients.

### ***Delivery: for all health professionals***

1.1.2.6 Interventions to increase physical activity should focus on activities that fit easily into people's everyday life (such as walking), should be tailored to people's individual preferences and circumstances and should aim to improve people's belief in their ability to change (for example, by verbal persuasion, modelling exercise behaviour and discussing positive effects). Ongoing support (including appropriate written materials) should be given in person or by phone, mail or internet.

1.1.2.7 Interventions to improve diet (and reduce energy intake) should be multicomponent (for example, including dietary modification, targeted advice, family involvement and goal setting), be tailored to the individual and provide ongoing support.

1.1.2.8 Interventions may include promotional, awareness-raising activities, but these should be part of a long-term, multicomponent intervention rather than one-off activities (and should be accompanied by targeted follow-up with different population groups).

1.1.2.9 Health professionals should discuss weight, diet and activity with people at times when weight gain is more likely, such as during and after pregnancy, the menopause and while stopping smoking.



1.1.2.10 All actions aimed at preventing excess weight gain and improving diet (including reducing energy intake) and activity levels in children and young people should actively involve parents and carers.

***Delivery: for health professionals in primary care***

1.1.2.11 All interventions to support smoking cessation should:

- ensure people are given information on services that provide advice on prevention and management of obesity if appropriate
- give people who are concerned about their weight general advice on long-term weight management, in particular encouraging increased physical activity.

***Delivery: for health professionals in broader community settings***

The recommendations in this section are for health professionals working in broader community settings, including healthy living centres and Sure Start programmes.

1.1.2.12 All community programmes to prevent obesity, increase activity levels and improve diet (including reducing energy intake) should address the concerns of local people from the outset. Concerns might include the availability of services and the cost of changing behaviour, the expectation that healthier foods do not taste as good, dangers associated with walking and cycling and confusion over mixed messages in the media about weight, diet and activity.

1.1.2.13 Health professionals should work with shops, supermarkets, restaurants, cafes and voluntary community services to promote healthy eating choices that are consistent with existing good practice guidance and to provide supporting information.

1.1.2.14 Health professionals should support and promote community schemes and facilities that improve access to physical activity, such as walking or cycling routes, combined with tailored information, based on an audit of local needs.

1.1.2.15 Health professionals should support and promote behavioural change programmes along with tailored advice to help people who are motivated to change become more active, for example by walking or cycling instead of driving or taking the bus.

1.1.2.16 Families of children and young people identified as being at high risk of obesity – such as children with at least one obese parent – should be offered ongoing support from an appropriately trained health professional. Individual as well as family-based interventions should be considered, depending on the age and maturity of the child.

***Delivery: for health professionals working with preschool, childcare and family settings***

1.1.2.17 Any programme to prevent obesity in preschool, childcare or family settings should incorporate a range of components (rather than focusing on parental education alone), such as:

- diet – interactive cookery demonstrations, videos and group discussions on practical issues such as meal planning and shopping for food and drink
- physical activity – interactive demonstrations, videos and group discussions on practical issues such as ideas for activities, opportunities for active play, safety and local facilities.

1.1.2.18 Family programmes to prevent obesity, improve diet (and reduce energy intake) and/or increase physical activity levels should provide ongoing, tailored support and incorporate a range of behaviour change techniques (see section 1.2.4). Programmes should have a clear aim to improve weight management.

***Delivery: for health professionals working with workplaces***

1.1.2.19 Health professionals such as occupational health staff and public health practitioners should establish partnerships with local businesses and support the implementation of workplace programmes to prevent and manage obesity.

### **1.1.3 Local authorities and partners in the community**

The environment in which people live may influence their ability to maintain a healthy weight – this includes access to safe spaces to be active and to an affordable, healthy diet. Planning decisions may therefore have an impact on the health of the local population. Fundamental concerns about safety, transport links and services need to be addressed. Effective interventions often require multidisciplinary teams and the support of a broad range of organisations.

These recommendations apply to:

- senior managers and budget holders in local authorities and community partnerships, who manage, plan and commission services such as transport, sports and leisure and open spaces (not just those with an explicit public health role)
- staff providing specific community-based interventions.

Implementation of these recommendations is likely to contribute to local area agreements and other local agreements and targets. The need to work in partnership should be reflected in the integrated regional strategies and reviewed regularly.

Recommendations that refer to the planning of buildings, and stair use in particular, should be implemented in the context of existing building regulations and policies, particularly in relation to access for disabled people.

Section 3 has links to tools to help with implementing the recommendations, meeting training needs, evaluating the impact of action and working in partnership with other organisations.

### ***Overarching recommendation***

1.1.3.1 As part of their roles in regulation, enforcement and promoting wellbeing, local authorities, primary care trusts (PCTs) or local health boards and local strategic partnerships should ensure that preventing and managing obesity is a priority for action – at both strategic and delivery levels – through community interventions, policies and objectives. Dedicated resources should be allocated for action.

### ***Strategy: for senior managers and budget holders***

1.1.3.2 Local authorities should set an example in developing policies to prevent obesity in their role as employers, by following existing guidance and (in England) the local obesity strategy.

- On-site catering should promote healthy food and drink choices (for example by signs, posters, pricing and positioning of products).
- Physical activity should be promoted, for example through travel plans, by providing showers and secure cycle parking and using signposting and improved décor to encourage stair use.

1.1.3.3 Local authorities (including planning, transport and leisure services) should engage with the local community, to identify environmental barriers to physical activity and healthy eating. This should involve:

- an audit, with the full range of partners including PCTs or local health boards, residents, businesses and institutions
- assessing (ideally by doing a health impact assessment) the effect of their policies on the ability of their communities to be physically active and eat a healthy diet; the needs of subgroups should be considered because barriers may vary by, for example, age, gender, social status, ethnicity, religion and whether an individual has a disability.

Barriers identified in this way should be addressed.

1.1.3.4 Local authorities should work with local partners, such as industry and voluntary organisations, to create and manage more safe spaces for incidental and planned physical activity, addressing as a priority any concerns about safety, crime and inclusion, by:

- providing facilities such as cycling and walking routes, cycle parking, area maps and safe play areas
- making streets cleaner and safer, through measures such as traffic calming, congestion charging, pedestrian crossings, cycle routes, lighting and walking schemes
- ensuring buildings and spaces are designed to encourage people to be more physically active (for example, through positioning and signing of stairs, entrances and walkways)
- considering in particular people who require tailored information and support, especially inactive, vulnerable groups.

1.1.3.5 Local authorities should facilitate links between health professionals and other organisations to ensure that local public policies improve access to healthy foods and opportunities for physical activity.

### ***Delivery: specific interventions***

1.1.3.6 Local authorities and transport authorities should provide tailored advice such as personalised travel plans to increase active travel among people who are motivated to change.

- 1.1.3.7 Local authorities, through local strategic partnerships, should encourage all local shops, supermarkets and caterers to promote healthy food and drink, for example by signs, posters, pricing and positioning of products, in line with existing guidance and (in England) with the local obesity strategy.
- 1.1.3.8 All community programmes to prevent obesity, increase activity levels and improve diet (and reduce energy intake) should address the concerns of local people. Concerns might include the availability of services and the cost of changing behaviour, the expectation that healthier foods do not taste as good, dangers associated with walking and cycling and confusion over mixed messages in the media about weight, diet and activity.
- 1.1.3.9 Community-based interventions should include awareness-raising promotional activities, but these should be part of a longer-term, multicomponent intervention rather than one-off activities.

#### 1.1.4 Early years settings

The preschool years (ages 2–5) are a key time for shaping lifelong attitudes and behaviours, and childcare providers can create opportunities for children to be active and develop healthy eating habits, and can act as positive role models.

These recommendations apply to:

- directors of children's services
- children and young people's strategic partnerships
- staff, including senior management, in childcare and other early years settings
- children's trusts, children's centres, Healthy Start and Sure Start teams
- trainers working with childcare staff, including home-based childminders and nannies.

Implementing these recommendations will contribute to meeting the target to halt the annual rise in obesity in children younger than 11 years by 2010 and to implementing the England and Wales National Service Frameworks for children, young people and maternity services (the Children's NSFs for England and Wales), and 'Every child matters' and similar initiatives in Wales.

Section 3 has links to tools to help with implementing the recommendations and meeting training needs.

### ***For all settings***

- 1.1.4.1 All nurseries and childcare facilities should ensure that preventing excess weight gain and improving children's diet and activity levels are priorities.
- 1.1.4.2 All action aimed at preventing excess weight gain, improving diet (and reducing energy intake) and increasing activity levels in children should involve parents and carers.
- 1.1.4.3 Nurseries and other childcare facilities should:
- minimise sedentary activities during play time, and provide regular opportunities for enjoyable active play and structured physical activity sessions
  - implement Department for Education and Skills, Food Standards Agency and Caroline Walker Trust<sup>[2]</sup> guidance on food procurement and healthy catering.
- 1.1.4.4 Staff should ensure that children eat regular, healthy meals in a pleasant, sociable environment free from other distractions (such as television). Children should be supervised at mealtimes and, if possible, staff should eat with children.

### **1.1.5 Schools**

During their school years, people often develop life-long patterns of behaviour that affect their ability to keep a healthy weight. Schools play an important role in this by providing opportunities for children to be active and develop healthy eating habits, and by providing role models. Improving children's diet and activity levels may also have wider benefits: regular physical activity is associated with higher academic achievement, better health in childhood and later life, higher motivation at school and reduced anxiety and depression.

There is no evidence that school-based interventions to prevent obesity, improve diet and increase activity levels foster eating disorders or extreme dieting or exercise behaviour.

These recommendations apply to:

- directors of children's services
- staff, including senior management, in schools
- school governors
- health professionals working in or with schools
- children and young people's strategic partnerships
- children's trusts.

Implementing these recommendations will contribute to meeting the target to halt the annual rise in obesity in children younger than 11 years by 2010 and implementing the Children's NSFs for England and Wales, the National Healthy Schools Programme (and the Welsh Network of Healthy Schools Schemes), and 'Every child matters' and similar initiatives in Wales.

Section 3 has links to tools to help with implementing the recommendations and meeting training needs.

Recommendations that refer to the planning of buildings, and stair use in particular, should be implemented in the context of existing building regulations and policies, particularly in relation to access for disabled people.

### ***Overarching recommendation***

1.1.5.1 All schools should ensure that improving the diet and activity levels of children and young people is a priority for action to help prevent excess weight gain. A whole-school approach should be used to develop life-long healthy eating and physical activity practices.

### ***Strategy: for head teachers and chairs of governors***

1.1.5.2 Head teachers and chairs of governors, in collaboration with parents and pupils, should assess the whole school environment and ensure that the ethos of all school policies helps children and young people to maintain a healthy weight, eat a healthy diet and be physically active, in line with existing standards and guidance. This includes policies relating to building layout and recreational spaces, catering (including vending machines) and the food and drink children bring into school<sup>[3]</sup>, the taught curriculum (including PE), school travel plans and

provision for cycling, and policies relating to the National Healthy Schools Programme and extended schools.

- 1.1.5.3 Head teachers and chairs of governors should ensure that teaching, support and catering staff receive training on the importance of healthy-school policies and how to support their implementation.
- 1.1.5.4 Schools should establish links with relevant organisations and professionals, including health professionals and those involved in local strategies and partnerships to promote sports for children and young people.
- 1.1.5.5 Interventions should be sustained, multicomponent and address the whole school, including after-school clubs and other activities. Short-term interventions and one-off events are insufficient on their own and should be part of a long-term integrated programme.

### ***Delivery: for teachers and other professionals***

- 1.1.5.6 Staff delivering physical education, sport and physical activity should promote activities that children and young people find enjoyable and can take part in outside school, through into adulthood. Children's confidence and understanding of why they need to continue physical activity throughout life (physical literacy) should be developed as early as possible.
- 1.1.5.7 Children and young people should eat meals (including packed lunches) in school in a pleasant, sociable environment. Younger children should be supervised at mealtimes and, if possible, staff should eat with children.
- 1.1.5.8 Staff planning interventions should consider the views of children and young people, any differences in preferences between boys and girls, and potential barriers (such as cost or the expectation that healthier foods do not taste as good).
- 1.1.5.9 Where possible, parents should be involved in school-based interventions through, for example, special events, newsletters and information about lunch menus and after-school activities.



### 1.1.6 Workplaces

The workplace may have an impact on a person's ability to maintain a healthy weight both directly, by providing healthy eating choices and opportunities for physical activity (such as the option to use stairs instead of lifts, staff gym, cycle parking and changing and shower facilities), and indirectly, through the overall culture of the organisation (for example, through policies and incentive schemes). Taking action may result in significant benefit for employers as well as employees.

These recommendations apply to:

- senior managers
- health and safety managers
- occupational health staff
- unions and staff representatives
- employers' organisations and chambers of commerce
- health professionals working with businesses.

The recommendations are divided into:

- those that all organisations may be able to achieve, with sufficient input and support from a range of staff, including senior management
- those that are resource intensive and may only be fully achieved by large organisations with on-site occupational health staff, such as the NHS, public bodies and larger private organisations.

The recommendations are likely to build on existing initiatives – such as catering awards, Investors in People and Investors in Health, and the Corporate Health Standard in Wales.

Section 3 has links to tools to help with implementing the recommendations and meeting training needs.

Recommendations that refer to the planning of buildings, and stair use in particular, should be implemented in the context of existing building regulations and policies, particularly in relation to access for disabled people.

### ***Overarching recommendation***

1.1.6.1 All workplaces, particularly large organisations such as the NHS and local authorities, should address the prevention and management of obesity, because of the considerable impact on the health of the workforce and associated costs to industry. Workplaces are encouraged to collaborate with local strategic partnerships and to ensure that action is in line with the local obesity strategy (in England).

### ***For all workplaces***

1.1.6.2 Workplaces should provide opportunities for staff to eat a healthy diet and be more physically active, through:

- active and continuous promotion of healthy choices in restaurants, hospitality, vending machines and shops for staff and clients, in line with existing Food Standards Agency guidance
- working practices and policies, such as active travel policies for staff and visitors
- a supportive physical environment, such as improvements to stairwells and providing showers and secure cycle parking
- recreational opportunities, such as supporting out-of-hours social activities, lunchtime walks and use of local leisure facilities.

1.1.6.3 Incentive schemes (such as policies on travel expenses, the price of food and drinks sold in the workplace and contributions to gym membership) that are used in a workplace should be sustained and part of a wider programme to support staff in managing weight, improving diet and increasing activity levels.

### ***For NHS, public organisations and large commercial organisations***

1.1.6.4 Workplaces providing health checks for staff should ensure that they address weight, diet and activity, and provide ongoing support.

1.1.6.5 Action to improve food and drink provision in the workplace, including restaurants, hospitality and vending machines, should be supported by tailored educational and promotional programmes, such as a behavioural intervention or environmental changes (for example, food labelling or changes to availability).

For this to be effective, commitment from senior management, enthusiastic catering management, a strong occupational health lead, links to other on-site health initiatives, supportive pricing policies and heavy promotion and advertisement at point of purchase are likely to be needed.

### 1.1.7 Self-help, commercial and community programmes

This section has been replaced by [Managing overweight and obesity in adults – lifestyle weight management services](#) (2014) NICE guideline PH53.

## 1.2 *Clinical recommendations*

This section has been replaced by [Obesity: Identification, assessment and management of overweight and obesity in children, young people and adults](#) (2014) NICE guideline CG189.

---

<sup>[2]</sup> [Caroline Walker Trust](#)

<sup>[3]</sup> See [School Food Trust](#).

## 2 Notes on the scope of the guidance

NICE guidelines are developed in accordance with a scope that defines what the guideline will and will not cover.

The scope specified that the guideline should cover adults and children aged 2 years or older and should include advice on the following aspects of overweight and obesity:

- identification and assessment in primary and secondary care
- clinical management in primary and secondary care
- clinical management of morbid obesity – in sufficient detail to inform and identify key aspects of care
- prevention in people who are currently a healthy weight, mainly outside the clinical setting, including
  - raising awareness
  - identifying children and adults who may benefit the most from participating in prevention programmes
  - maintaining energy balance
  - developing local strategies, with a focus on multifaceted interventions in:
    - ◇ the community – services and the wider environment
    - ◇ workplaces
    - ◇ schools
    - ◇ children aged 2–5
    - ◇ black and minority ethnic groups and vulnerable groups.

During the development of the guidance it was noted that the management of overweight and obesity in non-clinical settings had been omitted from the scope; this topic was also considered.

### How this guideline was developed

NICE commissioned the National Collaborating Centre for Primary Care and the Health Development Agency to develop this guidance. In April 2005, the Health Development Agency merged with NICE, and the work on this guidance was continued by NICE's Centre for Public Health Excellence. The National Collaborating Centre and the Centre for Public Health Excellence established two Guidance Development Groups (see appendix A), which reviewed the evidence and developed the recommendations. An independent Guideline Review Panel oversaw the development of the guidance (see appendix B).

There is more information about [how NICE clinical guidelines are developed](#) on the NICE website, including a booklet, 'How NICE clinical guidelines are developed: an overview for stakeholders, the public and the NHS'.

## 3 Implementation

### 3.1 *The NHS*

The Healthcare Commission assesses the performance of NHS organisations in meeting core and developmental standards set by the Department of Health in 'Standards for better health', issued in July 2004. Implementation of clinical guidelines forms part of the developmental standard D2. Core standard C5 says that national agreed guidance should be taken into account when NHS organisations are planning and delivering care.

NICE has developed [tools](#) to help organisations implement this guidance (listed below).

- Slides highlighting key messages for local discussion.
- Costing tools
  - Costing report to estimate the national savings and costs associated with implementation.
  - Costing template to estimate the local costs and savings involved.
- A signposting document on how to put the guidance into practice and national initiatives that support this locally.
- Audit criteria to monitor local practice.

### 3.2 *Other audiences and settings*

The guidance also makes recommendations for the following audiences and settings:

- public bodies – including local authorities; government, government agencies and arm's length bodies; schools, colleges and childcare in early years settings; forces, prisons and police service
- private and voluntary organisations
  - large employers (more than 250 employees)
  - small and medium employers (less than 50 and less than 250 employees, respectively)
- the general public including parents, and the media and others providing advice for different population groups.

## 4 Research recommendations

The Guidance Development Groups have made the following recommendations for research, based on their review of evidence, to improve NICE guidance and patient care in the future.

### 4.1 *What are the most effective interventions to prevent or manage obesity in children and adults in the UK?*

#### Why this is important

Many studies of interventions to prevent and manage obesity were of short duration, with little or no follow-up, were conducted outside the UK and were poorly reported. There is an urgent need for randomised controlled trials (or other appropriately designed studies, in line, for example, with the 'TREND statement'<sup>[4]</sup>), with at least 12 months' postintervention follow-up.

Studies should use validated methods to estimate body fatness (BMI), dietary intake and physical activity, and should assess the benefits of measures additional to BMI (such as waist circumference in children). Details of the intervention, provider, setting and follow-up times should be reported. The development of a 'CONSORT'-type<sup>[4]</sup> statement for public health research is strongly recommended. In research on managing obesity in clinical settings, the effects of different levels of intensity of non-pharmacological interventions and follow-up should be assessed. Further research is also needed on the effectiveness of pharmacological and surgical interventions in people with comorbidities such as type 2 diabetes or cardiovascular disease.

### 4.2 *How does the effectiveness of interventions to prevent or manage obesity vary by population group, setting and source of delivery?*

#### Why this is important

There is little UK-based evidence on the effectiveness of multicomponent interventions among key at-risk groups (for example, young children and families and black and minority ethnic groups), vulnerable groups (for example, looked-after children and young people, lower-income groups and people with disabilities) and people at vulnerable life stages (for example, women during and after pregnancy and people stopping smoking).

Interventions should be undertaken in 'real world' everyday clinical and non-clinical settings and should investigate how the setting, mode and source of delivery influence effectiveness. There is a need for research evaluating multicomponent interventions to manage obesity in primary care,

because factors such as the types of participant, the training of staff and the availability of resources may affect the results. Future research should:

- assess the feasibility of using in the UK interventions shown to be effective in other developed countries
- collect sufficient data to assess how the effectiveness of the intervention varies by age, gender, ethnic, religious and/or social group
- consider the value of corroborative evidence, such as associated qualitative studies on acceptability to participants
- consider the potential negative effects of an intervention as well as the intended positive effects (particularly for studies of children and young people).

### 4.3 *What is the cost effectiveness of interventions to prevent or manage obesity in children and adults in the UK?*

#### **Why this is important**

There is little evidence on the cost effectiveness of interventions, partly because of a lack of outcome measures that are amenable to health economic evaluations. Much of the evidence on the effectiveness of prevention strategies concerns crude measures such as average weight loss rather than response rates. Follow-up is usually short. In clinical research, more information from quality-of-life questionnaires throughout the intervention and follow-up period would help assess how valuable any clinical improvement is to the individual. This would allow greater comparison between types of intervention and improve assumptions made in cost-effectiveness analyses. It would be valuable to run cost-effectiveness studies in parallel with clinical trials, so that patient-level data can be collected.

### 4.4 *What elements make an intervention effective and sustainable, and what training do staff need?*

#### **Why this is important**

There are considerable barriers to the implementation of interventions, including organisational structures and personal views of both health professionals and patients. The enthusiasm and motivational skills of the health professional providing support and advice are likely to be key, and interventions may be more effective when tailored to the individual's needs. Further research is required to identify:



- what elements make an intervention effective and sustainable
- what staff training is needed.

## 4.5 *Evaluation and monitoring*

### 4.5.1 Population trends in overweight and obesity

Data on the prevalence of overweight and obesity at national and regional levels (with subgroup analysis by age, gender and social status) are published annually by the 'Health survey for England' (HSE) and the 'Welsh health survey'. The continued collection of such data at national and regional levels is strongly recommended. The 'Health survey for England' also provides detailed data on children and on black and minority ethnic groups about every 5 years. To allow full analysis of trends, more frequent collection of data among these and other vulnerable groups at national and local levels is encouraged.

### 4.5.2 Local and national action

Although considerable action is being undertaken at a local level that could directly or indirectly have an impact on the prevention or management of obesity, little evaluation is being undertaken. This observation is reflected in the 2005 Dr Foster survey<sup>[5]</sup> of obesity services, which found that only 15% of primary care organisations monitored interventions such as physical activity programmes and exercise on prescription. Many potentially important broader community policies are also not evaluated in terms of their health impact – examples include congestion charging, which is implemented to address traffic rather than health issues, and safer routes to schools.

It is therefore recommended that all local action – including action in childcare settings, schools and workplaces – be monitored and evaluated with the potential impact on health in mind. An audit of health impact should also be undertaken after each change has taken place. The need to evaluate projects should be taken into account when planning funding for those projects. It is recommended that the evaluation of local initiatives is carried out in partnership with local centres that have expertise in evaluation methods, such as health authorities, public health observatories and universities.

There is also limited high quality long-term evaluation of national schemes that are implemented locally and may have an impact on weight, diet or physical activity (such as interventions promoting a 'whole-school approach' to health, Sure Start initiatives and exercise referral schemes for children). It is therefore recommended that all current and future actions be rigorously monitored and evaluated with their potential health impact in mind. Evaluation of campaigns (including social

marketing campaigns) should go beyond the 'reach' of the campaigns and more fully explore their effectiveness in changing behaviour.

### 4.5.3 Clinical practice

In clinical practice there is a need to set up a registry on the use of orlistat in young people. There is also a need to undertake arrangements for prospective audits of bariatric surgery, so that the outcomes and complications of different procedures, their impact on quality of life and nutritional status, and the effect on comorbidities can be monitored in both the short and the long term.

---

<sup>[4]</sup> TREND – Transparent Reporting of Evaluations of Nonrandomized Designs.

<sup>[5]</sup> Dr Foster Survey of obesity services.

## 5 Other versions of this guideline

### 5.1 *Full guideline*

The full guideline, [Obesity: the prevention, identification, assessment and management of overweight and obesity in adults and children](#), contains details of the methods and evidence used to develop the guideline. It is published by the National Collaborating Centre for Primary Care and the Centre for Public Health Excellence at NICE.

## 6 Related NICE guidance

This guidance has updated, and replaces, the following NICE technology appraisals:

- Guidance on the use of surgery to aid weight reduction for people with morbid obesity (2002) NICE technology appraisal guidance 46
- Guidance on the use of orlistat for the treatment of obesity in adults (2001) NICE technology appraisal guidance 22

NICE has published the following related guidance:

- Four commonly used methods to increase physical activity (2006) NICE guideline PH2 [Replaced by [NICE guideline PH41](#) and [NICE guideline PH44](#)]
- [Eating disorders: core interventions in the treatment and management of anorexia nervosa, bulimia nervosa and related eating disorders](#) (2004) NICE guideline CG9
- Management of type 2 diabetes: management of blood pressure and blood lipids (2002) NICE guideline H [Replaced by [NICE guideline CG66](#)]
- [Nutrition support in adults: oral nutrition support, enteral tube feeding and parenteral nutrition](#) (2006) NICE guideline CG32
- [Improving the nutrition of pregnant and breastfeeding mothers and children in low income households](#) (2008). NICE guideline PH11
- [Physical activity and the environment](#) (2008) NICE guideline PH8
- [Promoting physical activity in children and young people](#) (2009) NICE guideline PH17

## 7 Updating the guideline

NICE clinical guidelines are updated as needed so that recommendations take into account important new information. We check for new evidence 2 and 4 years after publication, to decide whether all or part of the guideline should be updated. If important new evidence is published at other times, we may decide to do a more rapid update of some recommendations.

Sections 1.2 and 1.1.7 were updated in 2014 and section 1.1.1 was updated in 2015.

## Appendix A: The Guidance Development Groups

**Professor James McEwen (Chair)**

Emeritus Professor in Public Health and Honorary Senior Research Fellow, University of Glasgow

### *Public health*

**Mrs Mary Amos (from October 2005)**

Health and Social Policy Manager at Eastleigh Borough Council and South West Hampshire Primary Care Trusts Alliance

**Miss Elizabeth Biggs**

Hertfordshire Healthy Schools Coordinator and East of England Regional Coordinator

**Dr Mary Corcoran**

Director of Public Health, Gedling Primary Care Trust

**Dr Sara Kirk**

Principal Research Fellow, University of Leeds

**Mrs Esther Kurland**

Planning Advisor, Commission for Architecture and the Built Environment

**Dr Louis Levy**

Branch Head, Nutrition Policy and Advice, Food Standards Agency

**Ms Sue Mabley (to February 2005)**

Public Health Specialist, Welsh Local Government Association

**Dr Harshad Mistri**

Lay representative

**Mr Andy Ramwell**

Director of the Manchester Institute of Sport and Physical Activity, Manchester Metropolitan University

**Mrs Tracy Sortwell**

Lay representative

**Ms Helen Storer**

Dietetic Services Manager, Nottingham City Primary Care Trust

**Mr Malcolm Ward**

Principal Public Health Practitioner, National Public Health Service for Wales

*Clinical management*

**Mrs Mandakini Amin**

Health Visitor, Hinckley and Bosworth Primary Care Trust

**Ms Jude Cohen**

Lay representative

**Ms Helen Croker**

Clinical Research Dietitian, University College London

**Dr Penelope Gibson**

Consultant Community Paediatrician, Blackwater Valley and Hart Primary Care Trust

**Professor Paul Little**

General Practitioner and Professor of Primary Care Research, University of Southampton

**Mrs Suzanne Lucas**

Consumer representative

**Ms Mary O'Kane**

Clinical Specialist Dietitian, The General Infirmary at Leeds

**Mrs Sara Richards**

Practice Nurse, Slough Primary Care Trust

**Dr Ken Snider**

Public Health Physician, Director, County Durham and Tees Valley Public Health Network

**Professor John Wilding**

Professor of Medicine and Honorary Consultant, University Hospital Aintree, University of Liverpool

## *Guidance Development Group co-optees*

### **Public health**

#### **Professor Gerard Hasting**

Director, Institute for Social Marketing and Centre for Tobacco Control Research, University of Stirling and the Open University

#### **Dr Gill Hawksworth**

Community Pharmacist and Lecturer/Practitioner, University of Bradford

#### **Ms Wendy Hicks**

Specialist Nurse, Weight Management Service, Newcastle Primary Care Trust

#### **Mrs Anne Hollis**

Clinical Manager for School Nursing, Fareham and Gosport Primary Care Trust

#### **Dr Ira Madan, nominated by the Faculty of Occupational Medicine**

Consultant Occupational Physician, Guy's and St Thomas' NHS Trust

#### **Mr Lindley Owen**

Manager, Sustrans Cornwall

#### **Dr David Wilson**

Senior Lecturer in Paediatric Gastroenterology and Nutrition, Child Life and Health, University of Edinburgh

### **Clinical management**

#### **Dr John Buckley**

Exercise Physiologist, Keele University

#### **Laurel Edmunds**

Research Psychologist and Independent Consultant, Bristol Royal Children's Hospital

#### **Dr Nicholas Finer**

Senior Research Associate, University of Cambridge, and Honorary Consultant in Obesity Medicine, University of Cambridge Hospitals Trust



**Mr Tam Fry**

Honorary Chair, Child Growth Foundation

**Professor Philip James**

Chairman of the International Obesity Taskforce and Senior Vice President of the International Association for the Study of Obesity

**Mr David Kerrigan**

Consultant Surgeon, University Hospital Aintree

**Dr Krystyn Matyka**

Senior Lecturer in Paediatrics, University of Warwick

**Professor Mary Rudolf**

Consultant Paediatrician and Professor of Child Health, Leeds Primary Care Trust and the University of Leeds

**Russell Viner**

Consultant/ Honorary Senior Lecturer in Adolescent Medicine and Endocrinology, University College London Hospitals and Great Ormond St Hospital

**Professor Jane Wardle**

Professor of Clinical Psychology, University College London

*National Institute for Health and Clinical Excellence*

**Project Team, Centre for Public Health Excellence**

**Dr Hugo Crombie**

Analyst

**Dr Adrienne Cullum**

Analyst (Technical Lead)

**Mr Simon Ellis**

Associate Director (Methodology)

**Professor Mike Kelly**

Director

**Dr Caroline Mulvihill**

Analyst

**Ms Susan Murray**

Analyst

**Dr Bhash Naidoo**

Analyst

**Ms Karen Peplow**

Analyst

**Dr Nichole Taske**

Analyst

### *Obesity Collaborating Centre – Cardiff*

**Ms Sally Fry** (from October 2004 to September 2005)

Information Specialist

**Ms Hilary Kitcher** (from April 2005 to September 2005)

Information Specialist

**Dr Helen Morgan** (from January 2006)

Information Specialist

**Ms Lesley Sander** (from April 2005)

Information Specialist

**Dr Alison Weightman**

Associate Director and Head of Library Service Development; Director, Support Unit for Research Evidence

## ***Obesity Collaborating Centre – University of Teesside***

**Ms Tamara Brown**

Research Fellow

**Mr Phil Ray**

Research Assistant (from September 2004 to September 2005)

**Professor Carolyn Summerbell**

Professor of Human Nutrition and Assistant Dean for Research

## ***External Collaborators – University of York***

**Dr Catriona McDaid**, Research Fellow, Centre for Reviews and Dissemination

**Dr Amanda Sowden**, Associate Director, Centre for Reviews and Dissemination

**Ms Sarah Redmond**, Research Fellow, York Health Economics Consortium

**Mr Paul Trueman**, Director, York Health Economics Consortium

## ***National Collaborating Centre for Primary Care (NCC-PC)***

**Ms Janette Camosso-Stefinovic**, Information Librarian, NCC-PC, Department of Health Sciences, University of Leicester

**Ms Charmaine Larment**, Project Manager, NCC-PC

**Mr Richard Norman**, Health Economist, NCC-PC, and Queen Mary College, University of London

**Ms Vanessa Nunes**, Research Associate (Children's Lead); NCC-PC, Department of Health Sciences, University of Leicester

**Ms Elizabeth Shaw**, Research Fellow (Adult's Lead); NCC-PC, Department of Health Sciences, University of Leicester

**Dr Tim Stokes**, Project Lead; Clinical Director, NCC-PC, Senior Lecturer in General Practice, Department of Health Sciences, University of Leicester

**Dr Kathy DeMott**, Senior Health Research Fellow, NCC-PC

**Ms Katie Pike**, Statistician, Department of Health Sciences, University of Leicester

**Mrs Nancy Turnbull**, Chief Executive, NCC-PC

## Appendix B: The Guideline Review Panel

The Guideline Review Panel is an independent panel that oversees the development of the guideline and takes responsibility for monitoring adherence to NICE guideline development processes. In particular, the panel ensures that stakeholder comments have been adequately considered and responded to. The Panel includes members from the following perspectives: primary care, secondary care, lay, public health and industry.

### **Professor Mike Drummond (Chair)**

Professor of Health Economics, Centre for Health Economics, University of York

### **Dr Ann Hoskins**

Deputy Regional Director of Public Health, NHS North West

### **Dr Matt Kearney**

GP Public Health Practitioner, Knowsley PCT, and General Practitioner, Castlefields, Runcorn

### **Professor Ruth Hall**

Regional Director, Health Protection Agency South West

### **Dr John Harley**

Clinical Governance and Prescribing Lead, North Tees PCT

### **Mr Barry Stables**

Patient/Lay Representative

### **Dr Robert Walker**

General Practitioner, West Cumbria

## Appendix C: The algorithms

The recommendations from this guideline have been incorporated into [diet](#), [obesity](#) and [physical activity](#) NICE pathways. The [full guideline](#) also contains the algorithms.

## Appendix D: Existing guidance on diet, physical activity and preventing obesity

The recommendations from this guideline have been incorporated into [diet](#), [obesity](#) and [physical activity](#) NICE pathways. The [full guideline](#) also contains the information on existing guidance on diet, physical activity and preventing obesity.

## Changes after publication

March 2015: Section 1.1.1 was updated and replaced by [NICE guideline NG7](#).

January 2015: Updated Introduction to reflect recent alterations to guideline content.

November 2014: Section 1.2 was updated and replaced by [NICE guideline CG189](#).

May 2014: Section 1.1.7 was updated and replaced by [NICE guideline PH53](#).

December 2011: Minor maintenance

January 2010 CG43 Obesity replaces the following pieces of guidance:

- [TA22 Obesity – orlistat](#)
- [TA31 Obesity – sibutramine](#)
- [TA46 Obesity \(morbid\) – surgery](#)

**Sibutramine (Reductil): marketing authorisation suspended:**

On 21 January 2010, the MHRA announced the suspension of the marketing authorisation for the obesity drug sibutramine (Reductil). This follows a review by the European Medicines Agency which found that the cardiovascular risks of sibutramine outweigh its benefits. Emerging evidence suggests that there is an increased risk of non-fatal heart attacks and strokes with this medicine.

The MHRA advises that:

- Prescribers should not issue any new prescriptions for sibutramine (Reductil) and should review the treatment of patients taking the drug.
- Pharmacists should stop dispensing Reductil and should advise patients to make an appointment to see their doctor at the next convenient time.
- People who are currently taking Reductil should make a routine appointment with their doctor to discuss alternative measures to lose weight, including use of diet and exercise regimens. Patients may stop treatment before their appointment if they wish.



NICE clinical guideline 43 recommended sibutramine for the treatment of obesity in certain circumstances. These recommendations have now been withdrawn and healthcare professionals should follow the MHRA advice.

**December 2007 Amendment to Understanding NICE Guidance documents:**

The Understanding NICE Guidance (UNG) versions 'Preventing obesity and staying a healthy weight' and 'Treatment for people who are overweight or obese' previously referred to The Obesity Awareness and Solutions Trust (TOAST) as a source of additional information and support for people wanting to stay a healthy weight or to lose weight. Please note that this organisation closed in December 2007 and is no longer able to provide such information and support, and so reference to the organisation has been removed. The remaining organisations listed in the UNG documents are still able to offer information about preventing or treating obesity.

## About this guideline

NICE clinical guidelines are recommendations about the treatment and care of people with specific diseases and conditions in the NHS in England and Wales.

The guideline was developed by the National Collaborating Centre for Primary Care and the Centre for Public Health Excellence at NICE. They worked with a group of healthcare professionals (including consultants, GPs and nurses), patients and carers, and technical staff, who reviewed the evidence and drafted the recommendations. The recommendations were finalised after public consultation.

The methods and processes for developing NICE clinical guidelines are described in [the guidelines manual](#).

This guideline updates and replaces NICE technology appraisals 22, 31 and 46.

The recommendations from this guideline have been incorporated into [diet](#), [obesity](#) and [physical activity](#) NICE pathways. Tools to help you put the guideline into practice and information about the evidence it is based on are also [available](#).

Sections 1.2 and 1.1.7 were updated in 2014 and section 1.1.1 was updated in 2015.

### Your responsibility

This guidance represents the view of NICE, which was arrived at after careful consideration of the evidence available. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer, and informed by the summary of product characteristics of any drugs they are considering.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have regard to promoting equality of opportunity. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

### Copyright

© National Institute for Health and Clinical Excellence 2006. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

ISBN: 978-1-4731-0893-6