

2020 surveillance of acutely ill adults in hospital: recognising and responding to deterioration (2007) NICE guideline CG50 – summary of evidence

Overview

Studies identified in searches are summarised from the information presented in their abstracts.

We have noted where abstracts report studies as being undertaken in the United Kingdom. We have assumed that study populations are adults, unless otherwise reported in the abstract.

Evidence specific to sepsis is not summarised in this surveillance review as this is covered by the NICE related guideline on [sepsis: recognition, diagnosis and early management](#) (NG51).

Feedback from topic experts was considered alongside the evidence to reach a view on the need to update each section of the guideline.

Evidence from previous surveillance reviews in 2010 and 2016 and the exceptional surveillance review in 2019 for this topic was also considered.

Identification and evaluation of risk scoring tools: physiological observations in acute hospital settings

Surveillance proposal

The section of the guideline on identification and evaluation of risk scoring tools: physiological observations in acute hospital settings should not be updated.

Physiological observations in acute hospital settings

Previous surveillance

2010 surveillance summary

No relevant evidence was identified.

2016 surveillance summary

Three studies on staff competencies, education and training were suggested by topic experts. This new evidence highlighted the relevance of skills required for recording of physiological observations by staff and was considered consistent with current recommendations to ensure staff are appropriately trained.

One study on minimum physiological observations was highlighted by topic experts. This systematic review described signs and symptoms that trigger nurses to be concerned about a patient's condition. This new evidence was consistent with recommendation 1.8 that the response strategy for patients at risk of deterioration should be triggered by physiological track and trigger score or clinical concern.

2020 surveillance summary

A before and after study (1) assessed the impact of a short multidisciplinary training intervention for recognition of patient deterioration. Nursing, medical and allied nursing staff took part in a 1-hour long training session with real-life scenarios, simple tools and structured debriefing. Following training, staff were more likely to correctly calculate scores and perform observations at the correct frequency.

Intelligence gathering

A topic expert queried whether the age cut-off used in this guidance needed to be revised (i.e. 18 years and above or 16 years and above).

Impact statement

One study identified in the 2020 surveillance review, supported by the evidence from 3 studies on training identified in the 2016 surveillance review,

indicated the value of staff training in recognition of patient deterioration and is consistent with recommendation 1.1. that states that staff recording and acting upon physiological observations should have appropriate training to undertake these procedures and understand the clinical relevance.

A topic expert queried the age cut-off used in this guideline. However, the guideline took a pragmatic approach and did not define 'adult'.

New evidence is unlikely to change guideline recommendations.

Identification and evaluation of risk scoring tools: identifying patients whose clinical condition is deteriorating or is at risk of deterioration

Surveillance proposal

The section of the guideline on identification and evaluation of risk scoring tools: identifying patients whose clinical condition is deteriorating or is at risk of deterioration should not be updated.

Identifying patients whose clinical condition is deteriorating or is at risk of deterioration

Previous surveillance

2010 surveillance summary

No evidence was identified in the 2010 surveillance review.

2016 surveillance summary

Three studies on the frequency of recording of physiological observations were identified and highlighted the importance of monitoring patients and setting monitoring plans based on their risk level.

2020 surveillance summary

One randomised controlled trial (RCT) (2) explored the effect of timing of early warning score (EWS) measurements (score and country of setting not

reported) on patient outcomes in acutely admitted surgical and medical patients (n=1,346) at a university hospital. Patients (with an initial EWS of 0 or 1 [assumed low risk score]) were monitored at 8-hour or 12-hour intervals. No significant differences in clinical deterioration at 24 hours post-admission, cardiac arrests, intensive care unit (ICU) admissions, medical emergency team (MET) review, length of hospital stay, or elevated EWS at 48 hours were identified.

Intelligence gathering

A topic expert commented that, because of changes in policy and their view of possible new evidence, this section of the guideline needed to be reviewed. This topic expert also noted multiple local variations in practice and failure to integrate information. The topic expert stated there are examples of this within the 2019 [report on recognising and responding to critically unwell patients](#) by the Healthcare Safety Investigation Branch (HSIB). For example, in the reference patient event in this report, patient information was reported to be dispersed across a range of documentation and clinical staff, with the design and presentation of the information failing to support staff in assessment of the patient. Safety recommendations from this HSIB report advised that the Royal College of Physicians should continue to evaluate the implementation and use of NEWS2 and that NHS England/NHS Improvement expand the remit of the Cross-System Sepsis Programme Board to also include physical patient deterioration. A safety observation in the report noted that NEWS2 is not intended as a stand-alone tool but should be combined with other patient information and clinical observations.

Topic expert feedback also noted that a review should cover the need to integrate track and trigger with wider information and correct presentation and use of that information.

Impact statement

Recommendation 1.3 states that the frequency of monitoring of physiological observations should increase if abnormal physiology is detected and evidence from the 2016 surveillance review was consistent with this recommendation.

However, one RCT was identified in this 2020 surveillance review that

compared the effect of 8-hour or 12-hour EWS measurements on patient outcomes (assumed in low risk patients) and did not find any significant differences in negative clinical outcomes between groups. However, this was a single study and so overall findings support recommendation 1.3 that frequency of monitoring should be at least every 12 hours and with increasing frequency in patients with abnormal physiology. This surveillance review has identified an ongoing UK-based study funded by the National Institute for Health Research, which aims to produce an externally validated vital signs monitoring protocol (including information on how frequently observations should be recorded). The status of this study and any potential impact of publication will be monitored.

A topic expert highlighted the 2019 HSIB [report on recognising and responding to critically unwell patients](#), which includes safety recommendations and observations. However, the key issues were considered to relate primarily to implementation issues for systems for recognition and response to critically unwell patients and so were not considered to affect current guideline recommendations.

Identification and evaluation of risk scoring tools - choice of physiological track and trigger system

Surveillance proposal

The section of the guideline on identification and evaluation of risk scoring tools: choice of physiological track and trigger system should not be updated.

Choice of physiological track and trigger system

Previous surveillance

2010 surveillance summary

Twelve studies were included. It was noted that there was wide variation in the performance of different single and multiple-parameter track and trigger systems, with no evidence directly comparing the accuracy of different systems. A systematic review highlighted there was no evidence for clearly

identified cutoffs or weighting. A study describing the development of ViEWS was included (3). For hospital mortality, the Area Under Receiver Operating Characteristic curve (AUROC) of ViEWS was higher than the other aggregate weighted track and trigger tools tested. Two studies evaluating the use of the Modified Early Warning System (MEWS) were also identified. Overall, the evidence from the 2010 surveillance review reported variation in accuracy of the studied track and trigger systems, with no clearly defined cutoffs or weighting scores.

2016 surveillance summary

A systematic review indicated that EWS had good predictive values, but impact of implementation was uncertain. Six studies on accuracy of track and trigger systems were also identified by topic experts. National Early Warning Score (NEWS) was considered to perform better than other track and trigger systems in some clinical outcomes, but the 2016 surveillance review concluded that further research was required. This evidence was consistent with guideline recommendations that track and trigger systems should be used to monitor all adult patients in acute hospital settings.

Three systematic reviews on the impact of EWS on outcomes were identified. These were consistent with current recommendations on the use of multiple-parameter or aggregate weighted scoring systems.

2019 exceptional surveillance summary

This exceptional review was published in February 2019. Fifteen studies submitted by NHS England and the Royal College of Physicians were considered. The studies evaluated 10 different assessment tools. It was noted in the surveillance review that results were very mixed, and no single tool stood out as the most predictive or accurate. Recommendation 1.4 was revised to state that the EWS NEWS2 has been endorsed by NHS England. The exceptional surveillance review noted that this additional text is factually correct, does not change the intent of recommendations, and helps to provide a consistent message to healthcare professionals.

The exceptional surveillance review concluded that the guideline's research recommendations on the evaluation of early warning scores/track and trigger systems should be promoted with the National Institute for Health Research, with the aim of generating further evidence on EWS (including NEWS2).

2020 surveillance summary

Evidence was identified on the [National Early Warning Score \(NEWS\)](#), the [Modified Early Warning Score \(MEWS\)](#), [other early warning scores](#), and [mode of delivery](#).

National Early Warning Score (NEWS)

The 28 comparative studies identified on NEWS are presented in Table 1. No systematic reviews or RCTs were identified.

NEWS was compared with other scores or systems (listed alphabetically):

- Acute Physiology and Chronic Health Evaluation II (APACHE II) (4)
- Dyspnoea, eosinopenia, consolidation, acidaemia, atrial fibrillation (DECAF) (5)
- Early Deterioration Indicator (6)
- Goodacre physiological score (7)
- Groake physiological score (7)
- Laboratory Decision Tree risk index (8)
- Logistic EWS (9)
- MET criteria (44 sets) (10)
- Modified Early Warning Score (MEWS) (6), (11), (12), (4) (13)
- Modified Hypotension, Oxygen Saturation, Temperature, ECG, Loss of Independence (mHOTEL) score (13)
- NEWS-Lactate (14), (15), (16)
- NEWS-base excess (14)
- NEWS plus fractional inspired oxygen concentration (17)
- NEWS minus systolic blood pressure (BP) (18)
- NEWS minus temperature (18)

- NEWS plus white blood cell count, procalcitonin and midregional-proadrenomedullin or NEWS plus midregional-proadrenomedullin (19)
- NEWS plus age, sex and soluble urokinase plasminogen activator receptor (20)
- NEWS2 (21) (5)
- NEWS2 ALL COPD (5)
- NEWS at triage plus NEWS after ED management, coronary artery disease as comorbidity, use of vasoactive agent, initial serum bicarbonate level (22)
- Patient at Risk Score (23)
- Rapid Acute Physiology Score (RAPS) (7)
- Rapid Emergency Medicine Score (REMS) (7)
- SpO₂/FiO₂ (SF) ratio (11)
- Tachypnea, Oxygen saturation, Temperature, Alertness, Loss of Independence (TOTAL) score (13)
- VitalPAC EWS (11)
- Worthing physiological score (7)

The only scores that were compared with NEWS in more than one study (where specific scores were named in abstracts) were MEWS and NEWS-Lactate. In all 5 studies comparing NEWS with MEWS, NEWS showed better performance. Of the 3 studies comparing NEWS with NEWS-Lactate, NEWS-Lactate performed better than NEWS in 2 studies.

NEWS was compared with NEWS2 in two studies (21) (5). NEWS was reported to perform better than NEWS2 in prediction of in-hospital mortality (21). However, this study population was specific to patients at risk of type 2 respiratory failure or with documented type 2 respiratory failure (21). NEWS2 and NEWS2 ALL COPD performed better than NEWS in patients with exacerbation of chronic obstructive pulmonary disease.(5)

NEWS2 was also compared with the Hamilton EWS in patients from two Canadian hospitals (24) and NEWS2 was reported to have a lower AUROC value than the Hamilton score.

The performance of NEWS2 in the prediction of mortality within 48 hours was compared with other tools in the prehospital setting in two linked study reports from Spain.(25)(26) In the first report (25), NEWS2 had better predictive performance (AUC=0.891, 95% CI 0.84 to 0.94) than Shock Index (AUC=0.616, 95% CI 0.54 to 0.68), Glasgow Coma Scale-Age-Pressure Score (AUC=0.834, 95% CI 0.77 to 0.89) and Revised Trauma Score (AUC=0.800, 95% CI 0.73 to 0.86). In the second report (26), NEWS2 had better predictive performance (AUC=0.896, 95% CI 0.82 to 0.95) than the Scottish EWS (AUC=0.884, 95% CI 0.80 to 0.96), VitalPAC EWS (ViEWS) (AUC=0.894, 95% CI 0.82 to 0.96), Hamilton EWS (AUC=0.890, 95% CI 0.81 to 0.96), MEWS (AUC=0.848, 95% CI 0.76 to 0.93) and EWS (AUC=0.885, 95% CI 0.80 to 0.96).

Table 1. National Early Warning Score (NEWS)

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Abbott <i>et al.</i> (2015) (23) Prospective cohort study	All adult general medical patients admitted to single hospital over 20 days (sample size NR)	NEWS at admission	Composite of critical care admission or death within 2 days of admission	OR 1.54, p<0.001, NEWS ≥ 3 associated with outcome (OR=7.03, p=0.03)
		Patient at Risk Score (PARS) at admission		OR=1.42, p=0.056
Abbott <i>et al.</i> (2016) (14) Single centre prospective cohort study	Adult medical admissions (sample size NR)	NEWS (recorded at admission)	Composite of mortality or critical care escalation within 2 days of hospital admission	OR=1.46, p<0.01
		NEWS-Lactate		OR=1.18, p=0.01
		NEWS-base excess		OR=1.13, p=0.03
Abbott <i>et al.</i> (2018) (27) Prospective cohort study UK	Prehospital score calculated retrospectively for medical ward admissions at 1 UK teaching hospital (n=189)	NEWS (prehospital)	Composite of death or critical care escalation within 48 hours of hospital admission	OR=1.25, p=0.02
		NEWS (on admission)		OR=1.52, p<0.01
Bedoya <i>et al.</i> (2019) (28) Retrospective before and after cohort study	Hospitalised adults at a tertiary care academic facility and a community hospital (85,322 patients)	Implementation of NEWS in electronic health record and best practice alert	Rate of ICU transfer or death (academic hospital)	Adjusted hazard ratio = 0.94 (95% confidence interval, 95% CI 0.84 to 1.05)
			Rate of ICU transfer or death (community hospital)	Adjusted hazard ratio = 0.90 (95% CI 0.77 to 1.05)

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Brabrand <i>et al.</i> (2017) (7) Single centre observational study	Adults admitted to an acute medical unit at a teaching hospital (n=5,784)	NEWS RAPS REMS Goodacre physiological score Groake physiological score Worthing physiological score	All-cause 24-hour mortality and overall in-hospital mortality	Discriminatory power for 24-hour mortality > 0.8 for all scores (except Groake score = 0.587) and was highest for the Worthing score (0.847). Discriminatory power for prediction of in-hospital mortality was highest for Goodacre (0.810) and Worthing (0.800) scores and below 0.8 for other scores
Chiu <i>et al.</i> (2019) (9) Multicentre study UK	Patients discharged from ICU after cardiac surgery in 4 centres (all used VitalPAC for electronic collection of postoperative NEWS data) (13,631 patients)	Logistic EWS 24 h observation period	Composite: in-hospital death, cardiac arrest or unplanned ICU admission	AUROC=0.779 (95% CI 0.771 to 0.786)
		NEWS 24 h observation period		AUROC= 0.754 (95% CI 0.746 to 0.761), p<0.001
		Logistic EWS 6 h observation period		AUROC= 0.841 (95% CI 0.829 to 0.853)
		NEWS 6 h observation period		AUROC= 0.813 (95% CI 0.800 to 0.825), p<0.001
Dundar <i>et al.</i> (2019) (15) Retrospective observational study	Geriatric patients at emergency department (ED) (n=455)	NEWS	In-hospital mortality	AUC= 0.686 (95% CI 0.628 to 0.744)
		NEWS-Lactate		AUC= 0.714 (95% CI 0.658 to 0.770)
Echevarria <i>et al.</i> (2019) (5) Observational study UK	2645 patients with chronic obstructive pulmonary disease exacerbation at 6 UK hospitals. Scores calculated at admission.	NEWS	Inpatient mortality	AUROC=0.65
		NEWS2		AUROC=0.70, p=0.090
		NEWS2 ALL COPD		AUROC=0.72, p<0.001
		DECAF		AUROC=0.82
Eckart <i>et al.</i> (2019) (19) Multinational observational study (TRIAGE)	Adult medical patients needing ED care at 3 tertiary care centres in France, Switzerland and USA. NEWS calculated from admission data (1303 patients)	NEWS	All-cause 30-day mortality	AUROC=0.73
		NEWS plus white blood cell (WBC) count, procalcitonin (PCT) and midregional-proadrenomedullin (MR-proADM) or NEWS plus MR-proADM		AUROC=0.82 (p=0.002)
		NEWS	ICU admission	AUROC=0.65
	NEWS plus WBC count, PCT and MR-proADM		AUROC=0.70 (p=0.006)	

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Faisal <i>et al.</i> (2018) (29) Logistic regression modelling	Emergency admission patients with NEWS data over 24-month period (4.05% with hospital-acquired acute kidney injury) (33,608)	Models using index NEWS	Prediction of hospital-acquired acute kidney injury	Lower AUC (0.59 to 0.68) and lower sensitivity (19.84%) than models using maximum NEWS
		Models using maximum NEWS		Higher AUC (0.75 to 0.77) and higher sensitivity (67.6%) than models using index NEWS
Fernando <i>et al.</i> (2019) (24) Retrospective analysis of prospectively collected data	Consecutive rapid response team (RRT) patients from 2 hospitals (5,491 patients)	NEWS2 (low risk threshold)	Mortality	Sens 84.5% (95% CI 82.8% to 86.2%) Spec 49.0% (95% CI 47.4% to 50.7%) AUROC=0.72 (95% CI 0.71 to 0.74)
		Hamilton EWS (HEWS) ≥ 5 (low risk threshold)		Sens 75.9% (95% CI 73.9 to 77.9) Spec 67.6% (95% CI 66.1 to 69.1%) AUROC=0.76 (95% CI 0.75 to 0.77)
Foy <i>et al.</i> (2019) (13) Prospective observational study	Adult surgical patients (excluding obstetrics) admitted for > 24 hours to university teaching hospital (254 patients)	NEWS	30-day mortality	AUROC=0.805 (95% CI 0.688 to 0.92, p<0.001)
		MEWS		AUROC=0.76 (95% CI 0.63 to 0.88, p<0.001)
		mHOTEL		AUROC= 0.759 (95% CI 0.63 to 0.89, p<0.001)
		TOTAL		AUROC= 0.782 (95% CI 0.66 to 0.90, p<0.001)
Ghosh <i>et al.</i> (2018) (6) Model development and validation study	General ward patients (11,864 admissions for development, 2,418 general ward stays for validation)	NEWS	Deterioration	Validation AUROC=0.6569
		MEWS		Validation AUROC=0.6487
		Early Deterioration Indicator (electronic)		Validation AUROC=0.7655
Hydes <i>et al.</i> (2018) (30) Observational study England	1 acute care hospital in England (sample size NR), NEWS assessed in patients with primary liver disease, non-primary liver disease, no diagnosis of liver disease, alcohol-related liver disease	NEWS (primary liver disease)	Discrimination for 24-hour mortality, cardiac arrest or unanticipated admission to ICU	AUROC=0.873 (95% CI 0.860 to 0.886)
		NEWS (non-primary liver disease)		AUROC=0.898 (95% CI 0.891 to 0.905)
		NEWS (no diagnosis of liver disease)		AUROC=0.879 (95% CI 0.877 to 0.881)
		NEWS (alcohol-related liver disease)		AUROC=0.927 (95% CI 0.912 to 0.941)
Jo <i>et al.</i> (2016) (16) Retrospective cohort study	Adult patients at ED of academic tertiary care university hospital (n=4,624)	NEWS	2-day mortality	AUROC=0.94 (95% CI 0.91 to 0.96)
		NEWS-Lactate		AUROC= 0.96 (95% CI 0.94 to 0.98) (p=0.002)
Kwack <i>et al.</i> (2017) (11) Retrospective study	Adult patients admitted to respiratory ward (n=456)	NEWS	Unexpected ICU transfer	0.667
		MEWS		0.653
		VitalPAC EWS (ViEWS)		0.744

First author and year, study type	Population	System(s) evaluated	Outcome	Result
		SpO ₂ /FiO ₂ (SF) ratio		0.744 (p>0.05 vs. NEWS, MEWS, p=0.06 vs. ViEWS)
Kim <i>et al.</i> (2018) (31) Retrospective study	Patients with upper gastrointestinal bleeding (n=530)	NEWS-Lactate	Composite of in-hospital death, ICU admission, and need for ≥ 5 packs of red blood cell transfusion within 24 hours	AUROC=0.76 (95% CI 0.70 to 0.82) Sens by score: 3 (100%), 5 (98.3%), 5 (96.6%)
		Pre-endoscopic Rockall score		AUROC=0.66 (95% CI 0.59 to 0.73), p=0.004
		Glasgow-Blatchford score		AUROC=0.70 (95% CI 0.64 to 0.77), p=0.141
		AIMS65		AUROC=0.76 (95% CI 0.70 to 0.83), p=0.999 Sens by score: 0 (91.5%)
Lee <i>et al.</i> (2018) (32) Retrospective observational study	Number of patients NR	NEWS	In-hospital mortality	AUROC 0.765, 95% CI 0.659 to 0.846
		Combination model incorporating other factors, e.g. age, diagnosis		AUROC 0.861, p<0.005
Luis <i>et al.</i> (2018) (18) Observational study	Patients from 6 Portuguese hospital wards (sample size NR)	NEWS	Detection of patient clinical deterioration in preceding 24 hours	AUROC=0.944
		Model minus temperature		AUROC=0.965
		Model minus systolic BP		AUROC=0.903
Malycha <i>et al.</i> (2019) (17) Multicentre retrospective observational cohort study UK	Adult admissions with ≥ 1 complete set of vital sign observations recorded electronically at 5 hospitals from 2 UK NHS trusts (83,304 patients prescribed oxygen)	NEWS	In-hospital death or unplanned ICU admission within 24 hours of a complete set of vital sign observations	AUROC=0.811 (95% CI 0.809 to 0.814)
		NEWS plus fractional inspired oxygen concentration		AUROC=0.823 (95% CI 0.819 to 0.824)
Mitsunaga <i>et al.</i> (2019) (12) Retrospective single centre observational study	Prehospital (sample size NR)	Prehospital NEWS	In-hospital mortality	AUROC=0.678
		Prehospital MEWS		AUROC=0.652 (no significant difference between prehospital scores, p=0.081)
		ED NEWS		AUROC=0.789
		ED MEWS		AUROC=0.720 (p<0.001 between ED scores)
Pimentel <i>et al.</i> (2019) (21)	Adult admissions at 5 acute hospitals from 2 UK NHS trusts with type II respiratory failure (48,898 patients at risk of (by	NEWS2 in patients at risk of T2RF	In-hospital mortality within 24 hours	c-statistic=0.860 (95% CI 0.857 to 0.864)
		NEWS in patients at risk of T2RF		c-statistic=0.881 (95% CI 0.878 to 0.884)

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Multicentre retrospective observational study UK	diagnostic coding), 1,394 patients with documented type II respiratory failure (T2RF))	NEWS2 in patients with documented T2RF		c-statistic=0.841 (95% CI 0.827 to 0.855)
		NEWS in patients with documented T2RF		c-statistic=0.862 (95% CI 0.848 to 0.875)
Rasmussen <i>et al.</i> (2018) (20) Registry-based cohort study	Admitted acute medical patients (17,312 patients, admission NEWS available for 16,244)	NEWS	Prediction of 30-day mortality	AUC=0.86 (95% CI 0.85 to 0.87)
		NEWS plus age, sex and soluble urokinase plasminogen activator receptor		AUC=0.90 (95% CI 0.89 to 0.91), p<0.0001
Redfern <i>et al.</i> (2018) (8) Model development and validation study	Adult emergency medical admissions with vital signs and lab tests measured within their hospitalisation. Validation in 2 hospitals (Development n=97,933 admissions, validation n=21,028 + 16,383)	NEWS	Unanticipated ICU admission or in-hospital mortality within 24 hours	Validation c-statistic=0.877 (95% CI 0.873 to 0.882) to 0.898 (95% CI 0.898 to 0.904)
		Laboratory Decision Tree EWS: NEWS risk index		Validation c-statistic=0.901 (95% CI 0.898 to 0.905) to 0.916 (95% CI 0.911 to 0.921)
Smith <i>et al.</i> (2016) (10) Retrospective cohort study	Adults hospitalised at a large UK NHS district general hospital (2,245,778 vital signs sets, 103,998 admissions)	NEWS	Combined outcome: death, cardiac arrest, or unanticipated ICU admission	AUROC=0.88
		44 sets of MET criteria (no further details reported in abstract)		NEWS score of 7: Sens 44.5% NEWS score of 7: Spec 97.4% Sens 19.6 to 71.2% Spec 71.5 to 98.5% Position of NEWS ROC curve above and left of all MET criteria points (i.e. better discrimination). Positions of all MET criteria points above and to left of NEWS score efficiency curve (i.e. higher workload/trigger rate)
Srivilaithon <i>et al.</i> (2019) Case-control study	Cases were hospitalised patients experiencing in-hospital cardiac arrest (IHCA) (n=250), controls were adults, non-traumatic cases not experiencing IHCA within 24 hours of ED triage	NEWS	In-hospital cardiac arrest	AUROC=0.78 (95% CI 0.74 to 0.81)
		NEWS at triage plus NEWS after ED management, coronary artery disease as comorbidity, use of vasoactive agent, and initial serum bicarbonate level		AUROC=0.91 (95% CI 0.89 to 0.93)
Sutherasan <i>et al.</i> (2018) (33) Prospective cohort study	General wards (1,145 patients)	Hospital protocol in response to deterioration stratified using NEWS	In-hospital mortality and ICU transfer	No significant difference between intervention (protocol) and control (pre-protocol) groups

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Yuan <i>et al.</i> (2018) (4) Observational study	Patients in resuscitation room of a university hospital (621 cases)	NEWS	ICU admission	AUROC=0.760
		Modified EWS (MEWS)		AUROC=0.729
		Acute Physiology and Chronic Health Evaluation II (APACHE II)		AUROC=0.817
		NEWS	28-day mortality	AUROC=0.827
		MEWS		AUROC=0.723
		APACHE II		AUROC=0.883

Four additional, non-comparative studies (not summarised in Table 1) were suggested by topic experts and are summarised below.

The ability of NEWS to predict ICU admission within 48 hours or 30 day mortality in the prehospital setting (via clinical observations from 1,713 patients recorded by emergency ambulance crews) (AUROC NEWS total score = 0.740, 95% CI 0.685 to 0.795) was demonstrated in an observational study (34).

A further observational study (35) of the performance of NEWS in a prehospital setting (n=35,800 patients) reported AUROC values for death within 1 day of 0.840 (95% CI 0.832 to 0.858), with a higher AUROC for this outcome in the trauma subgroup (AUROC 0.901, 95% CI 0.859 to 0.942).

A mixed methods study (36) based at English hospitals participating in the National Cardiac Arrest Audit found that introduction of the NEWS score was associated with an additional 8.4% reduction in in-hospital cardiac arrests.

The application of the NEWS2 oxygen saturation parameters to a previously studied cohort (37) indicated that 44% (n=27/62) of patients who were scored ≥ 7 on the original NEWS and who then died would have been placed in a lower call-out threshold.

Modified Early Warning Score (MEWS)

Eight comparative studies on MEWS were included (and are tabulated below). Additional studies where MEWS was compared with NEWS are presented in Table 1.

Evidence was identified in which MEWS was compared with other scores or systems (listed alphabetically):

- Gastrointestinal EWS (38)
- Hypotension, Low Oxygen Saturation, Low Temperature, Abnormal Electrocardiogram (ECG), Loss of Independence (HOTEL) score (39)
- Linear predictors model (40)
- Logistic regression model with spline predictors (40)
- Novel metabolic score (41)
- Random forest model (40)
- RAPS (42)
- REMS (42)
- Simple Clinical Score (39)
- VitalPAC EWS (43)

All direct comparisons of MEWS with another score (other than the comparisons with NEWS described above) are from single studies.

Table 2. Modified Early Warning Score (MEWS)

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Churpek <i>et al.</i> (2016) (40) Observational cohort study (model development and validation)	Hospitalised ward patients at 5 hospitals (269,999 patients admitted)	MEWS	Combined: cardiac arrest, ICU transfer, or death	AUC=0.70
		Random forest model		AUC=0.80
		Logistic regression model with spline predictors		AUC=0.77
		Linear predictors model		AUC=0.74
				Machine learning methods more accurate in predicting clinical deterioration than logistic regression

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Dundar <i>et al.</i> (2016) (43) Prospective single-centred observational study	Geriatric (aged ≥ 65 years) ED patients at university hospital. Vital parameters measured on admission (n=671)	MEWS	Hospitalisation	AUC=0.727 (95% CI 0.689 to 0.765)
		VitalPac EWS		AUC=0.756 (95% CI 0.720 to 0.792)
		MEWS	In-hospital mortality	AUC=0.891 (95% CI 0.844 to 0.937)
		VitalPac EWS		AUC=0.900 (95% CI 0.860 to 0.941)
Jafar <i>et al.</i> (2016) (41) Prospective observational study	Patients presenting to resuscitation area of ED in teaching hospital (n=200)	MEWS	48-hour organ failure	OR=1.19, p>0.05
		Novel metabolic score (derived from a blood gas) (no further details)		OR=1.34, p>0.05
		MEWS	48-hour death	OR=1.32, p>0.05
		Novel metabolic score (derived from a blood gas) (no further details)		OR=1.56, p>0.05
		MEWS	48-hour escalation of care	Neither score was predictive (no further detailed)
		Novel metabolic score (derived from a blood gas) (no further details)		
		Metabolic score still significantly predictive of organ failure or death after controlling for MEWS parameters, p>0.05		
Kim <i>et al.</i> (2017) (38) Observational study	Consecutive events triggering MET calls in adults admitted to gastroenterology wards of medical centre (1,219 patients)	MEWS	ICU transfer	AUC=0.64
		Gastrointestinal EWS (EWS-GI)		AUC=0.76, p<0.001
Norman <i>et al.</i> (2018) (44) Before and after interventional study	Teaching hospital (sample size NR)	Emergency Room Safer Transfer of Patients (ER-STOP) system to avoid unexpected patient deterioration. Includes use of MEWS as component of a checklist	Critical care response team response within 24 hours of admission from ED to adult medical and surgical wards	Decreased
			ED wait times	Unchanged
			Cardiac care unit admissions	Unchanged
			ICU admissions	Unchanged
Weenk <i>et al.</i> (2019) (45)	Patients on surgical and internal medicine ward	MEWS via Visi Mobile continuous monitoring	Detection of high MEWS periods	71 (14 patients) high MEWS periods detected between non-nurse-observed periods. Time between high MEWS and

First author and year, study type	Population	System(s) evaluated	Outcome	Result
RCT	with minimum expected hospitalisation of 3 days (60 patients randomised)			next regular nurse measurement ranged 0 to 10 hours
		MEWS via HealthPatch continuous monitoring		32 (7 patients) high MEWS periods detected between non-nurse-observed periods. Time between high MEWS and next regular nurse measurement ranged 0 to 9 hours
Wei <i>et al.</i> (2019) (42) Retrospective observational study	Adult patients presenting to emergency department of university hospital (39,977 patients presented to ED, 4,857 admitted, 213 died)	REMS	Hospital admission	AUROC=0.76
		RAPS		AUROC=0.59
		MEWS		AUROC=0.55
		REMS	Hospital mortality	AUROC=0.88
		RAPS		AUROC=0.72
		MEWS		AUROC=0.73
		REMS	Length of hospital stay	AUROC=0.76
		RAPS		AUROC=0.67
		MEWS		AUROC=0.65
Wong <i>et al.</i> (2016) (39) Retrospective cohort study	All interfacility transfer cases by ambulance with nurse-led or physician-led escort (n=659)	MEWS at triage	Prediction of en-route complication during interfacility transfer in emergency department	AUROC=0.662
		MEWS at ambulance departure		AUROC=0.479
		Hypotension, Low Oxygen Saturation, Low Temperature, Abnormal ECG, Loss of Independence (HOTEL) score at triage		AUROC=0.613
		HOTEL at ambulance departure		AUROC=0.597
		Simple Clinical Score (SCS) at triage		AUROC=0.600
		SCS at ambulance departure		AUROC=0.568

Other early warning scores

Studies on other EWS are presented in Table 3. No direct comparison between other EWS or systems was made in more than one study.

Table 3. Other early warning scores (EWS)

First author and year, study type	Population	System(s) evaluated	Outcome	Result
Bellomo <i>et al.</i> (2018) (46) Pilot RCT	Surgical ward of a tertiary hospital (205 patients)	Laboratory alerts to trigger early ICU team review compared with usual care	RRT activation during first alert	Significantly more likely
			Receipt of allied health referral	Significantly less likely
			Mortality at 24 hours	No significant difference
			Overall mortality	No significant difference
Churpek <i>et al.</i> (2016) (47) Observational cohort study	Patients admitted to 5 hospitals (269,999 patient admissions)	Vital signs models using current and trend values	Combined: cardiac arrest, ICU transfer, and death	Use of trends improved accuracy vs. model using only current vital signs (AUC 0.78 vs. 0.74, p<0.001)
Douw <i>et al.</i> (2016) (48) Prospective cohort study	Adult surgical patients admitted to 3 surgical wards (trauma, vascular, and abdominal/oncological surgery) at a university teaching hospital (n=3,522)	Dutch-Early-Nurse-Worry-Indicator-Score (DENWIS)	Composite of unplanned ICU/high dependency unit (HDU) admission or unexpected mortality	AUC=0.85
		'Worry'		AUC=0.81
		EWS		AUC=0.86
		DENWIS plus 'worry'		AUC=0.87
		DENWIS plus EWS		AUC=0.91
Dziadzko <i>et al.</i> (2018) (49) Model development and validation	Patients admitted to 4 hospitals in 2013 or 5 th hospital in 2017 (68,775 admissions in 2013, 2,258 in 2017)	APPROVE model	Risk of death or respiratory failure needing ≥ 48 hours of mechanical ventilation	AUROC=0.87 (95% CI 0.85 to 0.88) in 2013 and 0.90 (95% CI 0.84 to 0.95) in 2017
		MEWS		AUROC NR Compared to APPROVE in 2013, at threshold to achieve comparable positive predictive value (PPV) (19% at MEWS > 4 and 22% at NEWS >6), MEWS had lower sensitivity (16%). In 2017, at comparable sensitivity threshold, (64% APPROVE >0.25, 67% MEWS and MEWS >4), more patients who triggered alert developed event with APPROVE (PPV 16%) than MEWS (PPV 7%).
		NEWS		AUROC NR Compared to APPROVE in 2013, at threshold to achieve comparable PPV (19% at MEWS > 4 and 22% at NEWS >6), NEWS had lower sensitivity (16%). In 2017, at comparable sensitivity threshold, (64% APPROVE >0.25, 67% MEWS and NEWS >4), more patients who triggered alert developed event with APPROVE (PPV 16%) than NEWS (PPV 4%).
Kwon <i>et al.</i> (2018) (50)	Data from Korean National Emergency	Deep-learning-based Triage		AUROC=0.935 Area Under Precision and Recall Curve (AUPRC)=0.264

First author and year, study type	Population	System(s) evaluated	Outcome	Result	
Retrospective observational cohort study	Department Information System on visits from 151 EDs and 1 general hospital for score development and validation (11,656,559 patients)	and Acuity Score	Primary outcome: in-hospital mortality		
		Korean Triage and Acuity Score		AUROC=0.785	AUPRC=0.192
		MEWS		AUROC=0.810	AUPRC=0.116
		Logistic regression		AUROC=0.903	AUPRC=0.209
		Random forest		AUROC=0.910	AUPRC=0.179
Linnen <i>et al.</i> (2019) (51) Systematic review	Adult patients on general hospital wards (6 studies)	EWSs using statistical modelling	ICU transfer or death	Mean AUC=0.80	
		EWSs using aggregate-weighting		Mean AUC=0.73 (with approximately 50% relative workload increase compared with statistical modelling EWSs)	
Merriel <i>et al.</i> (2017) (52) Before and after study	Women in hospital (Zimbabwe) (sample size NR)	Modified Obstetric EWS	Preoperative stabilisation	Increased	
			Action taken	Significantly improved	
O'Connell <i>et al.</i> (2016) (53) Before and after study	Adult tertiary hospital (sample size NR)	Observation and response chart and altered calling criteria	RRT call rate	Significantly increased	
			Ward admissions to ICU	Significantly increased	
			Deaths	No change (no further details)	
			Cardiac arrest	No change (no further details)	
Peek <i>et al.</i> (2017) (54) Review of clinical records	Consecutive post-cardiac surgery patients (n=400)	EWS	MET events, inpatient deaths and ICU admissions	73 events, no inpatient deaths and 12 ICU readmissions in study cohort	
		New Zealand EWS (NZEWS)		Rescoring with NZEWS resulted in 53 events. 8 of 12 ICU admissions preceded by a NZEWS event	
Romero-Brufau <i>et al.</i> , 2017 (55) Retrospective cohort study	All patients discharged (January to December 2011) from 2 tertiary care academic medical centre hospitals (sample size NR)	Published EWS (no further details reported)	Included unscheduled transfers to ICU, activation of rapid response system (RRS), and calls for cardiorespiratory resuscitation	All EWS had significantly higher PPV and sensitivity in medical population compared with surgical population. All EWS had PPV < 25%	

First author and year, study type	Population	System(s) evaluated	Outcome	Result	
Ryan <i>et al.</i> (2017) (56) Observational study	46 cases, 138 randomly selected controls. Pregnant or recently delivered women admitted at 2 maternity units (Canada) for > 24 hours	Modified Early Obstetric Warning System (MEOWS)	ICU admission > 24 hours	Sens 96%	Spec 54%
				Sens 96% (≥ 1 red trigger)	Spec 73% (≥ 1 red trigger)
Street <i>et al.</i> (2017) (57) Non-randomised pre-post intervention study	Adults who had elective surgery at 3 hospitals. Tool implemented in Post Anaesthesia Care Unit (PACU). Pre (n=723 patients), post (n=694 patients) phases	Post-Anaesthetic Care Tool (new observation, response and discharge chart)	Medical consultations initiated by PACU nurses	Significantly increased	
			Patients with MET activation criteria modified by anaesthetist while in PACU	Significantly increased	
			Analgesia administration	Significantly increased	
			Nurse assessment of pain and documentation of ongoing analgesia before discharge	Significantly increased	
			Adverse events recorded in PACU	Significantly increased	
			Adverse events after discharge from PACU	No change	
			Cardiac events after PACU discharge	Significantly decreased	
			Clinical deterioration after PACU discharge	Significantly decreased	
Williams <i>et al.</i> (2016) (58) Systematic review	Patients transported by ambulance in prehospital setting (8 studies)	EWS vs. standard practice using clinical judgement alone	Identification of critical illness	EWS pooled diagnostic OR=10.9 (95% CI 4.2 to 27.9)	EWS summary AUROC=0.78 (95% CI=0.74 to 0.82)

A further systematic review (59) identified 17 studies (n=157,878 participants) of the predictive performance of EWS for detection of patient deterioration in prehospital settings. AUROC values were reported to range from 0.50 (95% CI NR) to 0.89 (95% CI 0.82 to 0.96). The review concluded that 5 studies were at low or unclear risk of bias, with the remainder having high risk of bias.

Mode of delivery

A focused search was used to identify evidence comparing the use of electronic early warning systems with paper-based systems in recognition of patient deterioration. Seventeen studies were included (Table 4).

Table 4. Mode of delivery

First author and year, study type	Population	Intervention	Comparator	Outcome	Result
Bartkowiak <i>et al.</i> (2019) (60) Retrospective cohort study	Adults hospitalised on wards after surgical procedure at academic medical centre (32,537 admissions)	eCART	Scores were compared with each other	Composite: ICU transfer, ward cardiac arrest, or ward death	AUROC=0.79 (95% CI 0.78 to 0.81)
		NEWS			AUROC= 0.76 (95% CI 0.75 to 0.78)
		Modified EWS			AUROC= 0.75 (95% CI 0.73 to 0.76)
Fletcher <i>et al.</i> (2017) (61) Repeated treatment study	Patients in inpatient acute care wards of academic medical centre (6,736 eligible admissions)	Electronic medical record based alerting dashboard. Displayed all hospital patients in single view ranked by severity score and updated in real time	Dashboard display turned on and off each week for 10 2-week cycles over 20-week period	Overall rapid response team (RRT) activations	No significant difference
				First RRT activations	Significant increase
				Unexpected ICU transfers	No significant difference
				Cardiopulmonary arrests on general wards	No significant difference
				Deaths on general wards	No significant difference
Gagne <i>et al.</i> (2018) (62) Before and after study	NR	EWS embedded electronically into medical records and communication bundle with	Data 3 months before implementation	RRT calls	Non-significantly increased
				RRT calls for patients with EWS >4	Declined (significance NR)

First author and year, study type	Population	Intervention	Comparator	Outcome	Result
		notification and telephone collaboration between medical/surgical and ICU nurses (sample size NR)		ICU admissions after RRT calls	Significantly decreased
				ICU admissions for patients with EWS>4	Significantly decreased
				Documented reassessment response time	Significantly decreased
Green <i>et al.</i> (2018) (63) Multicentre retrospective study	Patients admitted to 5 hospitals	Electronic Cardiac Arrest Risk Triage (eCART) score	Scores were compared with each other	Cardiac arrest, ICU transfer or death	AUC= 0.801 (95% CI 0.799 to 0.802)
		NEWS			AUC= 0.718 (95% CI 0.716 to 0.720)
		Modified EWS			AUC=0.698 (95% CI 0.696 to 0.700)
		Between the Flags calling criteria			AUC= 0.663 (95% CI 0.661 to 0.664)
Heller <i>et al.</i> (2018) (64) Before and after study	Patients on 2 wards recovering from highly complex surgery (3,827)	Automated multiparameter EWS (MEWS based early warning system with paging functionality)	Before phase	Cardiac arrests	Significantly decreased
				Unplanned ICU admissions	Significantly decreased
Hogan <i>et al.</i> (2019) (36) Mixed methods study England	English hospitals participating in National Cardiac Arrest Audit	Electronic track and trigger systems	Paper-based track and trigger systems	In-hospital cardiac arrests	7.6% decrease (significance NR)
Hu <i>et al.</i> (2016) (65) Retrospective cohort study	Hospitalised adult patients with haematological malignancies at academic medical centre (565 admissions)	Routine vital signs and laboratory values from electronic medical record and use of machine learning algorithm	VitalPac EWS (VIEWS)	Clinical deterioration events (ICU transfer and cardiac arrest)	Neural network model higher PPV than VIEWS model (82% vs. 24%, p NR)
Imperato <i>et al.</i> (2017) (66) Retrospective pre and post intervention study	Patient in ED (sample size)	Clinical triggers programme measuring predetermined vital signs to trigger a rapid assessment by an emergency physician-led multidisciplinary team	Before phase	Primary outcome: median days admitted	No significant difference
				Secondary outcome: median days in special care unit	No significant difference
				Secondary outcome: in-hospital 30-day mortality	No significant difference

First author and year, study type	Population	Intervention	Comparator	Outcome	Result
				Secondary outcome: upgrade in level of care within 24 hours	No significant difference
Kang <i>et al.</i> (2016) (67) Validation study	Adult inpatients (3,889)	Electronic Cardiac Arrest Risk Triage score (electronic health record based EWS)	Standard of care RRT activation	Cardiac arrest	eCART AUROC=0.88 8 of 10 patients with cardiac arrest had high risk eCART scores (RRT activated on 2 patients, p<0.05)
				ICU transfer	eCART AUROC=0.80 eCART identified 52% of ICU transfers (vs. 34% by current system), p<0.001 Patients reached high risk eCART threshold median 30 hours before cardiac arrest/ICU transfer vs. 1.7 hours by standard RRT activation.
Keim-Malpass <i>et al.</i> (2019) (68) Model development and validation study	Patient admissions to cardiovascular medicine and surgery ward (8,111 patients for model training, 4,059 patients for validation, subsequent comparison of 91 patients with large abrupt spikes in risk with 59 control patients)	Spikes in ICU transfer risk from continuous predictive analytics monitoring (real time physiological data from ECG, vital signs, laboratory results and other clinical assessments)	Control patients matched for baseline risk including NEWS	ICU transfer	Event rate significantly higher in patients with risk spikes (PPV 24% vs. 7%, p=0.006). Risk spikes driven by respiratory changes had highest PPV (30 to 35%) and risk spikes from heart rate had lowest PPV (7%).
Kipnis <i>et al.</i> (2016) (69) Development and validation study	Adult hospitalisations (374,838 patients, 649,418 hospitalisations, analysis set: 48,723,248 hourly observations)	An automated hourly EWS (named Advanced Alert Monitor) based on electronic medical record data	Scores were compared with each other	Unplanned ICU transfer within next 12 hours	AUC=0.82
		eCART			AUC=0.79
		NEWS			AUC=0.76
Pappas <i>et al.</i> (2016) (70) Retrospective study	Patients outside ICU (580 mobile cart activations for critical care support, 577 completed)	Tele-ICU support for RRT for patients outside ICU using mobile platform (eMobile platform). Most common interventions:	RRT programme before testing of eMobile	Cost	Mean 66% increase in projected cost avoidance from unnecessary ICU transfers. For 2014, return on investment < \$1.66 per \$1 invested in IT support.

First author and year, study type	Population	Intervention	Comparator	Outcome	Result
		medication orders and laboratory studies. 33% calls managed without ICU upgrade.			
Subbe <i>et al.</i> (2017) (71) Prospective before and after study UK	All patients admitted to 2 clinical wards in UK district general hospital (2,139 patients (before phase), 2,263 patients (after intervention))	Electronic automated vital signs monitoring and notification system for patient deterioration. Abnormal vital signs alerted to RRT	Before phase	Number of RRT notifications	Significantly increased
				Number of patients with do not resuscitate (DNR) orders	Significantly increased
				Mortality	Significantly decreased
				Cardiac arrests	Significantly decreased
				Severity of illness in patients admitted to ICU	Decreased (significance NR)
				Mortality in patients admitted to ICU	Significantly decreased
Watkinson <i>et al.</i> (2018) (72) Model development and validation study	In-hospital patients (development data: 12,153 admissions, 301,644 vital sign observations, validation data: 53,395 admissions, 1,459,422 vital sign observations)	cCEWS (centile based EWS generated from continuously acquired data from bedside monitors)	Scores were compared with each other	Risk of cardiac arrest, unanticipated ICU admission, or death within 24 hours of a given vital sign observation	AUC=0.808 (95% CI 0.804 to 0.812)
		mCEWS (centile based EWS from manually recorded data)			AUC=0.868 (95% CI 0.864 to 0.872)
		NEWS			AUC=0.867 (95% CI 0.863 to 0.871)
Weller <i>et al.</i> (2018) (73) Prospective observational pilot study	Adult hospitalised neurological and neurosurgical patients in 26 bed unit (non-ICU) in academic medical centre (sample size NR)	Wireless, portable, body-worn, continuous, multiparameter vital signs monitor with automated nurse alarm notification via smartphones	NR	RRT call rate	Significantly reduced

First author and year, study type	Population	Intervention	Comparator	Outcome	Result
Wilson <i>et al.</i> (2015) (74) RCT	Hospitalised patients at a university hospital with \geq stage 1 acute kidney injury (1,201 patients in intervention group, 1,192 patients in control group)	Automated electronic alerts for acute kidney injury (text-based alert sent to covering provider and unit pharmacist to flag new acute kidney injury)	Usual care	ICU transfers	Non-significantly reduced
Wilson <i>et al.</i> (2016) (75) Single centre observational cohort study	ED (472 patient episodes)	Data-fusion patient status index for continuous vital sign monitoring (calculated retrospectively from continuous vital sign data)	Paper track and trigger system, Electronic track and trigger system	Unplanned patient deaths	Non-significantly reduced

Intelligence gathering

Choice and standardisation of systems

Topic expert feedback highlighted changes in the use of track and trigger systems, particularly noting the introduction of the NEWS2 system.

A topic expert suggested that the area of the guideline on choice of physiological track and trigger system needed to be reviewed, based on change in policy and their view of possible new evidence. This expert also noted that there are multiple local variations used in practice and failure to integrate information, citing the HSIB report as described previously. Another topic expert noted that this guideline does not recommend use of a specific score and that choice of track and trigger score should be standardised with all trusts using a recommended scoring system.

One topic expert reported that NEWS2 was, in their experience, the most widely adopted track and trigger system. Another topic expert commented that, as NEWS2 is being standardised across the NHS, the guideline should also promote NEWS2 in the interests of consistency. It was unclear whether the revised statement within recommendation 1.4 that NEWS2 is endorsed by

NHS England was considered adequate or whether additional specific recommendations were being requested. A further topic expert noted the formal endorsement of NEWS2 from NHS England and NHS Improvement and stated that the guideline should reflect safe use and clear documentation of NEWS2.

Components of systems

One topic expert commented that practice still varies in the components of EWSs and that outcome metrics are poor (no further details provided).

Topic expert feedback stated that a review was needed of the outcome measures of a successful early warning system and that the historical use of evidence supporting cardiac arrest as a suitable outcome measure was poor, citing a study by Hogan *et al.*, 2019 (no further details provided). A publication (76), identified during this surveillance review, described a Delphi consensus study to explore endpoints for validation of EWSs. Endpoints related to death, cardiac arrest and ICU admission were described as current compromises for validation of EWSs, with additional endpoints suggested by the study as potentially feasible in the future with large datasets and multiple measured parameters.

Settings and populations

A topic expert stated that clarity was needed on the use of track and trigger scoring systems in ED patients (citing previous concerns that systems are not validated in this population). It was also commented that recognition of subgroups was required where track and trigger systems may not be as reliable, such as in pregnancy or people with spinal cord injury.

A topic expert reported potential issues with monitoring in people with delirium, dementia, or learning disabilities and a possible need to consider how to score people in these subgroups who cannot have a score recorded for an observation. No information was identified in this surveillance review.

Electronic systems

One topic expert raised the use of electronic alert systems, indicating that a comparison with electronic versus paper only systems would be of value.

Three NICE medical technology innovation briefings ([VitalPAC for assessing vital signs of patients in hospital](#) MIB79, [EarlySense for heart and respiratory monitoring and predicting patient deterioration](#) MIB49, and [Visensia for early detection of deteriorating vital signs in adults in hospital](#) MIB36) and an in-development digital health technologies pilot of [Lifelight First for monitoring vital signs](#) were identified in this surveillance review that have relevance to electronic monitoring of patient physiological observations.

Impact statement

Choice and standardisation of systems

The 2016 surveillance review concluded that NEWS appeared to perform better than other scores but that more research was required.

Recommendation 1.4 states that ‘track and trigger systems (NEWS2 has been endorsed by NHS England) should use multiple-parameter or aggregate weighted scoring systems, which allow a graded response. These systems should define the parameters to be measured and the frequency of observations, [and] include a clear and explicit statement of the parameters, cut-off points or scores that should trigger a response.’ Recommendation 1.4 does not currently recommend any specific track and trigger system but now states that NEWS2 has been endorsed by NHS England.

In this surveillance review, evidence was identified where NEWS was compared with another EWS or system. However, the only named scores that were compared with NEWS in more than one study were MEWS and NEWS-Lactate. There were 4 studies identified in which NEWS2 was compared with another score. Two of these were performed in the UK but in specific study populations with respiratory conditions. The other two studies were performed in the hospital and prehospital settings (in Canada and Spain respectively). Therefore, in the 4 studies identified for NEWS2, populations, settings, and tested comparisons varied between studies. All direct comparisons of MEWS with another score (other than the comparisons with NEWS described above)

were from single studies. For other EWSs, no direct comparison between scores or systems was made in more than one study.

As most identified comparisons between track and trigger systems were only available in single studies, the evidence identified in this surveillance review does not allow a definitive conclusion on which track and trigger tool may have superior performance. Therefore, the identified evidence is not sufficient to impact on recommendation 1.4.

Components of systems

One topic expert reported that practice varies in EWS components and that outcome metrics are poor, advising that a review of the outcome measures of a successful EWS was required. One study was identified that proposed the most suitable endpoints for studies of EWSs but, as current recommendations do not refer to the use of endpoints in EWS research, this study does not impact on recommendations.

Settings and populations

A topic expert stated that clarity was needed on the use of track and trigger scoring systems in emergency department (ED) patients (citing previous concerns that systems are not validated in this population). Evidence of the use of track and trigger systems in EDs is now available and was included in this surveillance review. Recommendation 1.1 advises that physiological observations be recorded for adult patients in acute hospital settings, including patients in the ED for whom a clinical decision to admit has been made. The identified evidence on use of track and trigger systems in patients in EDs is consistent with this recommendation.

Topic expert feedback noted that recognition of subgroups was required where track and trigger systems may not be as reliable, such as in pregnancy or people with spinal cord injury. Two studies were included that showed that the modified obstetric EWS may have high sensitivity and positively affect clinical outcomes. However, as the monitoring of physiological observations are already covered in the NICE guidelines on [intrapartum care for healthy women and babies](#) (CG190) and [intrapartum care for women with existing](#)

[medical conditions or obstetric complications and their babies](#) (NG121) it is considered that this evidence fits within these guidelines and so these studies are not considered to impact on current recommendations in this guideline (CG50). NICE guideline NG41 provides recommendations for [spinal injury: assessment and initial management](#). A NICE guideline on [rehabilitation after traumatic injury](#) is also in development.

Electronic systems

Recommendation 1.4 does not specify the mode of delivery of track and trigger systems or whether electronic systems should be used. The evidence identified in this 2020 surveillance review indicates that the use of electronic monitoring systems show promise in improving patient outcomes. However, the available evidence does not allow a single electronic system to be identified as most effective and so this evidence is not considered to have potential impact on recommendation 1.4.

Identification and evaluation of risk scoring tools: physiological parameters to be used by track and trigger systems

Surveillance proposal

The section of the guideline on identification and evaluation of risk scoring tools: physiological parameters to be used by track and trigger systems should not be updated.

Physiological parameters to be used by track and trigger systems

Previous surveillance

2010 surveillance summary

A total of 7 studies were identified and serum lactate was noted to be a potential additional parameter. However, the available evidence on predictive

accuracy and generalisability was not conclusive and the review concluded that no new evidence contradicted the existing guideline recommendations.

2016 surveillance summary

No new evidence on physiological parameters to be used by track and trigger systems was identified.

The 2016 surveillance review identified examples of other pieces of NICE guidance that describe the management of diseases or conditions where specific parameters should be measured (e.g. [chest pain of recent onset: assessment and diagnosis](#) [CG95], [head injury: assessment and early management](#) [CG176]). Further guidance on physiological observations in specific circumstances is available in such related NICE guidelines.

2020 surveillance summary

Two studies were identified in the 2020 surveillance review.

Parameters to be measured in specific clinical circumstances

Continuous ECG monitoring was evaluated in a cohort study of 8,105 acute care patient admissions (77). Models with addition of continuous ECG measures improved prediction of deterioration leading to ICU transfer and unanticipated death within the next 24 hours compared with models using only laboratory results and vital signs.

In a prospective observational study (78) of 148 non-critical care inpatients, networked blood glucose meters with a visual alert for out of range blood glucose levels and a clinical glucose alert escalation pathway significantly increased nursing and medical action in response to episodes of adverse glycaemia. Patient-days with hyperglycaemia and patient-days with mean blood glucose > 15 mmol/l were significantly decreased.

Intelligence gathering

Parameters to be measured by scoring systems

Two topic experts commented that this guidance needed to be updated to include development of confusion and use of supplemental oxygen. New

evidence for the differing EWSs and track and trigger systems (including NEWS2) is summarised in [choice of physiological track and trigger system](#).

Recognition of acute kidney injury

One topic expert suggested that the guideline considered recognition of acute kidney injury. It is noted that guidance on recognition of acute kidney injury is already covered by some related NICE products. NICE has produced a [quality standard on acute kidney injury](#) (QS76) and a clinical guideline on [acute kidney injury: prevention, detection and management](#), both of which are included in the NICE Pathway on [acutely ill patients in hospital](#). Statement 3 (monitoring in hospital for people at risk) of the [quality standard on acute kidney injury](#) (QS76) uses CG50 recommendation 1.6 as source guidance. The acute kidney injury guideline directly refers to CG50. [Recommendation 1.2.1](#) from the acute kidney injury guideline NG148 directly refers to CG50: 1.2.1 Follow the recommendations in [CG50] on the use of track and trigger systems (EWSs) to identify adults who are at risk of acute kidney injury because their clinical condition is deteriorating or is at risk of deteriorating. [Recommendation 1.2.2](#) from the acute kidney injury guideline NG148 also refers to monitoring of urine output, which can be considered in line with CG50 recommendation 1.6 that advises additional monitoring (e.g. hourly urine output) in specific clinical circumstances.

Impact statement

Parameters to be measured in specific clinical circumstances

Two studies were included in the 2020 surveillance review that showed that monitoring of glucose and ECG monitoring may be of value to patients. This evidence is considered consistent with recommendation 1.6 that additional monitoring should be considered in specific clinical circumstances.

Recommendation 1.6 refers to lactate as a physiological parameter that may be used in additional monitoring in specific clinical circumstances. Three studies were identified that supported the performance of the NEWS-Lactate tool (see [choice of physiological track and trigger system](#)). This evidence

supports the potential value of lactate as a physiological parameter to be used by track and trigger systems and so is consistent with this recommendation.

Recognition of acute kidney injury

One topic expert flagged the need for this guidance to consider recognition of acute kidney injury.

Existing recommendations do not refer to recognition of acute kidney injury. Two relevant studies were identified in this surveillance review on the use of automated electronic alerts for acute kidney injury and modelling for prediction of acute kidney injury. The evidence identified was from single studies and was not considered to impact on current recommendations.

Response strategies for patients identified as having a deteriorating clinical condition: critical care outreach services for patients whose clinical condition is deteriorating

Surveillance proposal

The section of the guideline on response strategies for patients identified as having a deteriorating clinical condition: critical care outreach services for patients whose clinical condition is deteriorating should not be updated.

Critical care outreach services for patients whose clinical condition is deteriorating

Previous surveillance

2010 surveillance summary

A focused search for evidence on whether provision of critical care outreach services improve outcomes in patients at risk of deteriorating and/or deteriorating identified 26 studies, including evidence on the effectiveness of response strategies (e.g. critical care outreach services [CCOS] and METs).

The new evidence was inconclusive with no available evidence directly comparing different response strategies. A stakeholder commented at consultation that it would be useful if this guideline included recommendations on standardised methods for communicating patient deterioration. The 2010 surveillance review concluded that there was no impact on the existing guideline recommendations.

2016 surveillance summary

One systematic review assessed EWS or emergency response teams in hospital survival of adult patients. While the included evidence was described as of poor quality by the review authors, it supported a global approach of including track and trigger systems and teams having critical care competencies. A second systematic review found RRSs in acute care settings to be associated with reduced cardiorespiratory arrests outside of ICU and reduced mortality. An observational study also reported that RRS results in decreased admissions and mortality.

This guideline does not recommend a specific service configuration and, since the identified evidence did not allow a specific service configuration to be recommended, the evidence identified in the 2016 surveillance review was consistent with existing recommendations.

2020 surveillance summary

Education and training

One RCT (79) with 67 nurses examined a 3-hour interactive web-based educational intervention for early recognition and response to patient deterioration in a simulated environment; the control group did not receive the educational intervention. There was a significant increase in respiratory rate monitoring by the intervention group compared with control group nurses. The intervention group also had significantly better post-test scores for knowledge, performance in assessment and management of clinical deterioration, and reporting clinical deterioration.

An RCT (80) assessed whether a training intervention could improve escalation of care by junior doctors. Before undertaking simulated escalation of care scenarios, postgraduate surgeons (n=33) were randomised to either receive an educational intervention or act as a control group with no intervention. The intervention group demonstrated significantly improved patient assessment, communication, and non-technical skills and detected significantly more medical errors compared with the control group.

A simulated RRT (81) assessed the impact of a cognitive aid booklet intervention on performance in volunteer residents from a medical school. Resident performance was significantly better in the intervention group compared with an undefined control group.

Response triggers

A pilot retrospective cohort study (82) (n=620) assessed outcomes in patients with modifications to hospital RRT call triggers (n=393) compared with patients without modifications in (i.e. with standardised) RRT call triggers. Repeat RRT calls and in-hospital mortality were increased in patients with modifications. Modifications that were more conservative than standardised calling criteria were associated with mortality.

Intelligence gathering

No information was identified.

Impact statement

Education and training

Two RCTs and 1 intervention study were identified in the 2020 surveillance review. These studies assessed the provision of educational/training interventions in 2 RCTs and a cognitive aid booklet intervention in 1 intervention study. These interventions resulted in improved performance by professionals in simulated test settings, improved knowledge and other post-test scores and skills. This evidence is consistent with recommendation 1.7 that staff caring for patients in acute settings should have competencies in monitoring, measurement, interpretation and response appropriate to the level

of care they provide, and that education and training should be provided to ensure staff have the required competencies.

Response triggers

One cohort study identified in the 2020 surveillance review found that modifications to RRT call triggers resulted in increased repeat RRT calls and in-hospital mortality compared with standardised RRT call triggers. This study does not contradict recommendations 1.8 and 1.9 that response strategies should be triggered by track and trigger score or clinical concern and that thresholds should be set locally.

Response strategies for patients identified as having a deteriorating clinical condition: graded response strategy

Surveillance proposal

The section of the guideline on response strategies for patients identified as having a deteriorating clinical condition: graded response strategy should not be updated.

Graded response strategy

Previous surveillance

2010 surveillance summary

Evidence identified in the 2010 surveillance review is summarised in the above section on [critical care outreach services for patients whose clinical condition is deteriorating](#).

2016 surveillance summary

Evidence identified in the 2016 surveillance review is summarised in the above section on [critical care outreach services for patients whose clinical condition is deteriorating](#).

2020 surveillance summary

The evidence identified in the 2020 surveillance review is presented in categories reflecting the terminology used in individual studies to describe the response strategy.

Critical care response team / critical care outreach team

Three included studies described critical care response teams (CCRT) or outreach teams (CCOT).

A stepped wedge cluster RCT (83) of patients on general hospital wards in Iran (n=7,802 patients admitted before implementation, 10,990 patients admitted after implementation) assessed impact of a CCOT delivered by a team of intensive care nurses. The order of implementation was randomised. The intervention did not reduce mortality and focus groups highlighted lack of management and nurse endorsement of the intervention.

Addition of a critical care medicine-trained physician assistant to the CCOT was evaluated in a retrospective study (84) of 3,099 adults in ED and hospital requiring CCOT consultation. Compared with a control hospital with no staffing change, time to ICU transfer and hospital mortality were both significantly reduced.

A before and after study (85) showed a 5.2% increase in initiation and completion of DNR orders after critical care response team implementation (n=5,406 CCRT activations).

Intensive care consult service

One study (86) in adult patients (sample size not reported) found that an ICU consult service with a dedicated intensivist (a dedicated daytime/weekday service without formal training of ward teams) did not reduce total hospital mortality but significantly decreased both 30-day mortality of patients admitted to ICU and the 14-day ICU readmission rate.

Medical emergency teams

Four studies were identified that assessed the use of METs.

A data mining study (87) of 13,656 MET calls from 7,936 patients modelled the impact of redesign of the MET service to include a pre-emptive management algorithm for direct treatment of patients without waiting for MET team attendance. Decreases of 69% in projected MET calls for hypotension and 20% in projected total MET calls were reported.

MET implementation in a tertiary care hospital led to a significant decrease in crude hospital mortality rate in a retrospective cohort study of 511 MET activations (88).

A prospective study (89) (83 MET activations) found no significant difference in in-hospital mortality after MET implementation but a significant decrease in serious events (no further details).

In inpatients requiring MET activations for respiratory decompensation (n=50) (90), a prospective study showed that lung ultrasonography during MET activations had a higher diagnostic accuracy than MET diagnosis (84% versus 75%, p=0.29).

Patient at risk teams

One single centre before and after study (91) (number of patients not reported) of a patient at risk team (composed predominantly of experienced ward nurses) found significant decreases in ward cardiac arrests and hospital length of stay and decreases in direct ward admissions to ICU and number of MET calls compared with a historical control.

Rapid response teams / systems

Twenty-six studies were included that evaluated the use of rapid response teams (RRT) or systems (RRS). These are summarised below according to the study designs used.

Rapid response teams / systems – systematic reviews

Three systematic reviews were identified. One systematic review and meta-analysis (92) included 29 studies and reported that RRTs resulted in significant reductions in adult hospital mortality and cardiopulmonary arrests.

A second systematic review and meta-analysis (93) included 20 studies

published in English, Portuguese or Spanish on patients in adult hospital units and reported reductions in mortality and cardiac arrest. However, the review authors noted a low quality of evidence, high heterogeneity and risk of bias in the included studies. A further systematic review and meta-analysis (94) of 30 studies in adults hospitalised in non-ICU care found reductions in hospital mortality and non-ICU cardiac arrests.

Rapid response teams / systems – RCTs

Three RCTs were also included. A cluster RCT (95) of 3,135 emergency team calls for patients at 23 hospitals evaluated outcomes from implementation of RRTs compared with standard practice (with conventional cardiac arrest team responses to emergencies). The 1-year trial found that the proportion of delayed calls was significantly decreased with intervention. In all hospitals, delayed calls were significantly associated with increased risk of unplanned ICU admission and death. An RRS including a standardised observation and communication protocol with a pragmatic medical response strategy was described in a pragmatic stepped wedge cluster RCT of patients on 28 wards of 7 hospitals (intervention n=35,389, control n=34,267) (96). There were no significant differences between groups in unexpected death rates, cardiac arrest rates, or unplanned ICU admissions. However, the study authors noted a lower than expected baseline incidence of unexpected death and cardiac arrests that reduced the study statistical power. In an RCT (97) of 714 patients in emergency wards, a rapid response nursing team including an intensive care nurse and anaesthesia technician was compared with a control group receiving conventional emergency ward treatment. Transfer to special care units and the level of care in the first 24 hours after admission were both significantly decreased.

Rapid response teams / systems – non-RCTs

Of the remaining 20 non-RCT primary studies, details of the composition or delivery of the RRT/RRS were available at abstract level for 10 studies.

A multicomponent intervention was tested in adults hospitalised in medical-surgical wards (sample size not reported) (98). This included an intensivist-led RRT (available 24 hours per day, 7 days per week), with educational modules,

publicity, and bedside simulation-based training. A single activation criterion (heart rate < 40/minute or > 140/minute, systolic BP < 80 mmHg, cardiac arrest, respiratory rate <8/min or > 30/min, pulse oximetry < 90% with oxygen, respiratory distress in tracheostomy, respiratory arrest, coma, or sudden change in consciousness, seizure) allowed any caregiver to directly contact the RRT via a dedicated telephone number. Significant decreases in unexpected mortality, overall mortality, and sequential organ failure assessment score upon ICU admission were observed. ICU admission significantly increased.

A different multicomponent intervention was assessed in patients in a teaching hospital (sample size not reported) (99). This involved an RRS with introduction of a medical emergency team (MET) and redesign of a ward observation chart with colour coding of vital sign variables to indicate variation from normal; mandated minimum frequency of vital sign measurements; 3 formal levels of escalation according to physiological instability measured by MEWS; an education and e-learning package; and practice in escalation and communication. All-cause hospital mortality and the hospital-standardised mortality ratio both decreased (significance not reported).

Changes in RRS were implemented and assessed in adult ward patients (28,914 patients in control period, 39,802 patients in intervention period) (100). Expanded administrative oversight of an existing RRS resulted in restructure of ward nurse education for early recognition of deterioration; system changes to facilitate RRT mobilisation; development of RRT treatment protocols; changes to data collection and analysis for system compliance and performance improvement. RRT activations significantly increased. Cardiac arrests and hospital mortality significantly decreased.

Across 225 hospitals in an Australian region, a Between the Flags system for recognition and management of patient deterioration was evaluated (sample size not reported) (101). This included implementation of a Standard Adult General Observation Chart, staff awareness training, and two-level clinical emergency response systems. Implementation resulted in a significant

decrease in cardiac arrest rate and a significant increase in rapid response rate.

A 2 tier RRS (including ICU outreach nurse and RRT) resulted in a decrease in patients (sample size not reported) meeting MET call criteria without current treatment and non-significant decreases in unplanned ICU admissions and cardiac arrests compared with before implementation (102).

An RRS with after-hours Clinical Team Coordinator was assessed in 300 medical patients (103). Compared with before implementation, there was an increase in identification of adverse events and major adverse events (significance not reported).

Pharmacist involvement on an RRT was evaluated (before phase n=234, after phase n=157 rapid response events) (104). Median time to medication administration from central pharmacy was significantly decreased.

An RRT programme was implemented in a before and after study (sample size not reported) to decrease non-ICU cardiopulmonary arrests and overall hospital mortality (105). Charge nurses undertook training as unit specific RRT members. All inpatient staff undertook annual training in RRT including surveillance and recognition of deterioration. Overall hospital mortality was significantly decreased.

In a study (106) of 40,177 admissions and 795 RRT activations, RRT triggers were displayed on each floor and in-house staff were trained on RRT activations. No significant difference in mortality was observed.

EWS-based proactive RRT rounding was assessed in hospitalised adults (n=12,148) and compared with an RRT model based on staff nurse identification of abnormalities in vital signs (107). Unplanned ICU transfer was more likely during the RRT baseline (OR=1.392, 95% CI 1.017 to 1.905) compared with EWS-based proactive rounding.

In the remaining 10 non-RCT primary studies, no details were available at abstract level on the composition and delivery of the RRT/RRS and results

were mixed (108), (109), (110), (111), (112), (113), (114), (115), (116), (117). Sample sizes (where reported) ranged from 82 to 166,569 patients.

Statistically significant results from RRT/RRS implementation were identified in some of these non-RCT primary studies for which no details of composition or delivery was available at abstract level. These statistically significant results included:

- increase in frequency of RRS activation (114)
- decrease in time from patient deterioration to intervention (110)
- decrease in cardiopulmonary arrests (115)
- decrease in number of ICU admissions (110)
- decrease in in-hospital mortality (115)
- decrease in cardiopulmonary arrest, unplanned ICU admission or death (115)
- decrease in illness severity scores at ICU admission (114)
- significant decrease in waiting time for ICU beds (108)
- decrease in ICU length of stay (114)
- increase in recognition of patients requiring palliative care (108)
- increase in detection of adverse events (109)
- decrease in rate of incidents occurring as a result of staff leaving normal duties to attend to MET calls (117)

In a multicentre prospective cohort study (118) (Grieve *et al.*, 2019) of 49 UK NHS hospitals, 9,192 ward patients experiencing clinical deterioration were assessed for ICU transfer or continuing care on general wards. Effects of ICU transfer were estimated to be increased for patients with higher physiological severity (based on NEWS), particularly for older patients.

Intelligence gathering

A topic expert (under potential areas requiring update) suggested the specific actions for staff during waiting for responsive personnel and on who to contact out of hours be considered.

A further topic expert commented that any guideline update could also consider effective response (no further details provided).

Impact statement

Evidence included in the 2020 surveillance review evaluated the effects of a range of response strategies, including RRTs or systems, METs, patient at risk teams, intensive care consult services, and critical care outreach teams. The included evidence showed some promising patient benefits from response strategies, including reductions in cardiac arrests and mortality. However, the response strategies tested varied between studies and no evidence was available that directly compared one response strategy configuration with another.

The guideline states that ‘no specific service configuration can be recommended as a preferred response strategy for individuals identified as having a deteriorating clinical condition.’ The evidence identified in the 2020 surveillance review is consistent with this statement.

New evidence is unlikely to change guideline recommendations.

Transfer of patients from critical care areas: transfer of patients from critical care areas to general wards

Surveillance proposal

The section of the guideline on transfer of patients from critical care areas to general wards should not be updated.

Transfer of patients from critical care areas to general wards

Previous surveillance

2010 surveillance summary

No new evidence was identified that contradicted the existing guideline recommendations.

2016 surveillance summary

No relevant evidence was identified on timing of patient transfer from critical care to general wards.

2020 surveillance summary

Topic experts did not suggest this area as being of particular interest for this surveillance review and so focused evidence searches were not performed. Key relevant evidence identified in the course of the surveillance review was summarised.

Timing of transfer from critical care areas to general wards

A systematic review and meta-analysis (119) assessed whether time of discharge from ICU was associated with hospital mortality. Fourteen studies were included that assessed timing of discharge in 953,312 patients. Hospital mortality was significantly increased for patients discharged from ICU during nighttime compared with daytime.

In a further systematic review and meta-analysis (120), 18 cohort studies (n=1,191,178 patients) were included. Patients who were discharged from ICU during 'out of hours' had significantly higher in-hospital mortality and readmission rates compared with patients discharged 'in hours.'

Crude hospital mortality was significantly greater for patients discharged from ICU at nighttime compared with daytime discharges in a multicentre retrospective cohort study (n=19,622 patients) (121). This finding was confirmed by multivariable analysis. No difference in ICU readmission was observed for patients discharged at nighttime compared with daytime.

A single centre retrospective cohort study (122) in a medical-surgical ICU in a Brazilian tertiary care hospital examined the impact of timing of ICU discharge on patient outcomes. Patients (n=4,313) were classed as nighttime or daytime ICU discharge. There were no significant differences in in-hospital mortality, frequency of ICU readmission and length of hospital stay between groups. Length of ICU stay was significantly lower in nighttime compared with daytime ICU discharged patients.

Prediction and prevention of poor outcomes at transfer from critical care areas to general wards

In a retrospective observational study (123) in a cohort of gastrointestinal surgery patients (n=124 patients) following transfer from an ICU/HDU, NEWS at transfer from ICU predicted readmission (OR 1.32, 95% CI 1.01 to 1.72, p=0.04).

The ability of the Rothman Index (generated automatically from vital signs, laboratory data, cardiac rhythms, and nursing assessments) to predict mortality and readmissions in 1,445 patients transferred from the surgical ICU to the surgical ward was assessed in a single centre retrospective study (124). Seventy-nine patients were readmitted to ICU within 48 hours of transfer. Readmitted patients had significantly lower Rothman Index scores at 72, 48 and 24 hours before transfer and at 24 and 48 hours after transfer.

A retrospective observational study (125) of 194 adults discharged from the ICU found that higher Rothman Index (≥ 50) was associated with significantly lower odds of an adverse event within 72 hours (including readmission to a higher level of care) compared with a Rothman Index < 50 .

After implementation of a nationwide quality improvement project with detailed public display via the internet of ICU discharge proportions and outcomes (n=163,371 patients), the prevalence of nighttime discharges from ICU significantly decreased (126).

Intelligence gathering

No intelligence was identified.

Impact statement

Timing of transfer from critical care areas to general wards

Recommendation 1.14 states that, after it has been decided that a patient is to be transferred from critical care to the general ward, that this transfer should be made as early as possible during the day. It is also recommended that transfer between 22.00 and 07.00 should be avoided where possible and documented as an adverse event if it happens.

Two systematic reviews and 2 cohort studies were identified in the 2020 surveillance review that contributed data on the effect of timing of transfer from critical care areas to general wards.

Both systematic reviews and a multicentre cohort study found that poor outcomes (i.e. mortality and readmission rates) were increased in patients discharged from ICU at nighttime or 'out of hours' compared with daytime or 'in hours.' One retrospective cohort study found no significant differences in mortality, ICU readmission or length of stay between patients discharged from ICU at nighttime or daytime. However, since this was a single centre, non-UK study, this study was not considered enough to affect recommendation 1.14. The evidence identified on timing of discharge was considered on balance to be supportive of recommendation 1.14 to avoid transfer out of critical care to the general ward during nighttime hours.

Prediction and prevention of poor outcomes at transfer from critical care areas to general wards

An observational study noted that applying NEWS at transfer from ICU predicted readmission. Two observational studies also found the Rothman Index to be associated with readmission to ICU or a higher level of care. Current recommendations do not describe the use of scores in predicting outcomes at the point of patient transfer from critical care to the general ward. However, since these findings were based from a relatively small number of observational studies, this evidence was not sufficient to affect existing recommendations.

One quality improvement project was reported to reduce nighttime discharges from ICU. Current recommendations do not describe interventions to reduce nighttime ICU discharges, but evidence from this single study was not considered adequate to impact the recommendations.

Transfer of patients from critical care areas: care on the general ward following transfer

Surveillance proposal

The section of the guideline on care on the general ward following transfer should not be updated.

Care on the general ward following transfer

Previous surveillance

2010 surveillance summary

No new evidence was identified that contradicted the existing guideline recommendations.

2016 surveillance summary

One systematic review studied the use of risk stratification tools for identification of patients at high risk of adverse events following ICU discharge. Identified tools had AUROC values ranging from 0.66 to 0.92. The review authors concluded that more research was required in this field.

A systematic review found some evidence that provision of continuity, coordination and transition for patients with advanced illness improved patient caregiver and satisfaction.

A second systematic review assessed the effectiveness of interventions for improving safety and efficiency of patient discharge from ICU to the general ward. The authors reported that there was no difference in mortality reduction between the interventions and that there was a lack of good quality evidence in the area.

An additional systematic review assessed the impact of critical care transition programmes on patients discharged from an ICU to general wards. This evidence supported the recommendations in this guideline and that critical

care transition programmes may have a role in patient transfer from ICU to general wards, but more research is required.

The identified evidence was consistent with guideline recommendation 1.15 that the critical care transferring team and receiving ward team should share responsibility and assure patient care continuity, including support by a care plan with a formal handover.

A systematic review demonstrated that interventions for the provision of information to family members and patients reduced transfer anxiety compared with standard care.

Decision tools were found to support patient knowledge and awareness of treatment choices in a further systematic review.

This evidence was supportive of recommendation 1.1.6 that patients being transferred from critical care to a general ward should be offered information about their condition and encouraged to actively participate in decision relating to their recovery.

2020 surveillance summary

Topic experts did not suggest this area as being of particular interest for this surveillance review and so focused evidence searches were not performed. Key relevant evidence identified in the course of the surveillance review was summarised.

Tools for risk stratification following discharge from critical care areas to general wards

A prospective cohort study (127) was performed in adults who had been admitted to the ICU in a single tertiary hospital for more than 24 hours. The effectiveness of selected scores in predicting unplanned ICU readmission or unexpected death in the first 48 hours after ICU discharge was compared. Scores were calculated on the day of ICU discharge. The Stability and Workload Index for Transfer score (SWIFT), Sequential Organ Failure Assessment score (SOFA) and simplified Therapeutic Intervention Scoring

System (TISS-28) had similar predictive performance (AUC 0.66, 0.65 and 0.67 respectively, $p=0.58$).

A novel machine learning algorithm was developed to predict post-ICU outcomes in patients ($n=2,018$ patient episodes) at a single UK hospital (128). The developed model showed better discrimination for unplanned ICU readmission or death compared with the SWIFT score (AUROC=0.7095 vs. 0.6082, $p=0.014$).

A prospective observational study (129) demonstrated that NEWS recorded immediately before ICU discharge was a significant independent predictor of early clinical deterioration (acute respiratory failure or circulatory shock) within 24 hours of ICU discharge. The AUROC of NEWS was 0.92 (95% CI 0.89 to 0.94), and a score >7 gave a sensitivity of 93.6% and a specificity of 82.2% for detection of early clinical deterioration.

Patient characteristics and physiological variables were collected from 3,109 surgical patients within 48 hours before surgical ICU discharge (130). Among these patients there were 141 unplanned ICU readmissions within 72 hours. A clinical nomogram was developed and found to have moderate performance (AUC=0.71) for prediction of ICU readmission risk.

Interventions and discharge practices for patients transferred from critical care areas to general wards –pharmacist input and medicines reconciliation

A clinical pharmacist was integrated at the point of transfer of patients from ICU to wards in a multicentre RCT (131). The involvement of a clinical pharmacist resulted in a significant decrease in drug-related problems.

A multicentre retrospective study (132) of 985 patients transferred from 58 ICUs to non-ICU locations was performed. Daily patient care rounds in the ICU and orders discontinued and rewritten at point of transfer from the ICU were associated with decreased odds of medication error.

A medication reconciliation programme by pharmacists at the point of ICU admission and before ICU discharge was reported in a prospective study at 2

ICUs (n=478 patients) to reduce the proportion of patients with 1 or more medication transfer errors after discharge (133).

Interventions and discharge practices for patients transferred from critical care areas to general wards – multidisciplinary quality improvement intervention

A quality improvement intervention involving a multidisciplinary team was used to improve the transfer of patients from the ICU to the surgical ward (134). The intervention included verbal handovers, time-sensitive patient evaluations, and visual cues and was piloted over 1 year in high risk surgical patients discharged from the ICU. Patients received a baseline bedside examination from the primary team and respiratory therapists within 60 minutes of arrival on the ward. The rate of readmissions was reported to have decreased non-significantly, but the intervention was valued by stakeholders and had become standard of care.

Interventions and discharge practices for patients transferred from critical care areas to general wards –transition programmes

An interrupted time series analysis (135) of adult patients (n=32,234) discharged from medical-surgical ICUs in 8 hospitals assessed the effects of a critical care transition programme on risk of readmission to ICU within 72 hours and 14-day mortality. A multidisciplinary ICU provider team (including a physician, nurse and respiratory therapist) evaluated each patient after ICU discharge in 3 hospitals but not in the other 5 hospitals. Following programme implementation, there were no significant effects on readmission to ICU within 72 hours or mortality.

A MET follow up programme for patients discharged from ICU to general wards (including respiratory care, regular visiting and staff communication) (n=1,229 patients enrolled in an intervention study) (136) reported no significant difference in readmission to ICU within 72 hours but significant improvements in cardiac arrest, SOFA score, and readmission to ICU within 72 hours for patients discharged to a step-down unit only.

A new model for transition of care from the medical ICU (MICU) to medical ward was implemented in a before and after study (n=966 patients) at a single academic centre (137). In this model, MICU staff continued management of all patients transferred to the medical ward from the MICU for at least 24 hours. This model resulted in a significant decrease in hospital length of stay after MICU transfer but no significant differences in adjusted mortality or MICU readmission rates.

Interventions and discharge practices for patients transferred from critical care areas to general wards –nurse-led interventions

The ability of an acute post-ICU nurse-led ward based review tool (PIRT) to identify high risk patients at ICU discharge and improve ward based review and outcomes was tested in a prospective cohort of patients (n=1,028) discharged from 2 adult UK ICUs (138). The cohort received risk scoring at ICU discharge and inpatient review using PIRT. The use of ward based PIRT review did not show significant correlation with reduced poor outcomes overall or overall readmission. However, use of PIRT significantly reduced early readmission within 48 hours from 4.5% to 3.6% (p=0.039) while significantly increasing rate of late readmission (48 hours to 14 days) from 2.7% to 5.8% (p=0.046).

In a prospective intervention study with historical control (139) of a nurse-led multidisciplinary ICU follow up programme in patients with respiratory issues discharged from ICU (n=369), ICU readmission within 72 hours, all ICU readmission and hospital mortality were all significantly reduced, with no significant difference reported for 90-day mortality.

A study used analysis of registry data on 42,040 ICU admissions and a questionnaire to all Dutch ICUs to assess the association between post-ICU outcomes and ICU discharge practices (140). The ICU discharge practices of interest included ICU discharge criteria, bed managers, early discharge planning, step-down facilities, medication reconciliation, verbal and written handover, monitoring of post-ICU patients, and consulting ICU nurses. It was reported that no association between ICU discharges and rates of ICU readmission or post-ICU in-hospital mortality could be demonstrated.

Patient and caregiver information

In a prospective cohort study (141) of 451 adult patients at 10 hospitals transferred from a medical-surgical ICU to a hospital ward, patients more frequently reported satisfaction with their transfer when they received more information, when their questions were addressed, when they met the ward physician before transfer, and were assessed by a nurse within 1 hour of their arrival on the ward.

Intelligence gathering

No information was identified.

Impact statement

Tools for risk stratification following discharge from critical care areas to general wards

The systematic review of risk stratification tools for identification of patients at high risk of adverse events that was identified in the 2016 surveillance review found variable predictive performance between tools and concluded that additional research was required.

In this 2020 surveillance review, a further 4 studies were identified on the use of scores in identification of adverse outcomes following discharge from critical care to the general ward. Current recommendations do not describe the use of scores or tools to identify patients at high risk of adverse outcomes following transfer from critical care to general wards. The evidence identified in the 2016 and 2020 surveillance reviews indicates that some scores and algorithms may show promise in supporting the identification of patients at risk following ICU transfer, but no single tool emerges as the most predictive of negative outcomes following transfer to the general ward from critical care. This evidence is not considered to have impact on current recommendations.

Interventions and discharge practices for patients transferred from critical care areas to general wards

In the 2016 surveillance review, 3 systematic reviews were identified that were consistent with recommendation 1.15 that the team transferring from critical

care and the receiving ward team should share responsibility and ensure continuity of patient care, with support from a care plan with a formal handover.

A further 9 primary studies were identified in this 2020 surveillance review. The guideline does not currently recommend any specific interventions for care on the ward or discharge practices and the evidence identified in this surveillance review does not demonstrate which specific interventions or practices would be most effective and so this evidence is considered to support existing recommendations.

Patient and caregiver information

The 2 systematic reviews identified in the 2016 surveillance review were consistent with recommendation 1.16 that patients transferred from critical care to general wards should be offered information about their condition and actively encouraged to participate in shared decision making. The one cohort study found in this 2020 surveillance review is consistent with recommendation 1.16 on provision of information to patients and encouraged involvement in shared decision making.

Research recommendations

Identification and evaluation of risk scoring tools

- What is the clinical effectiveness and cost effectiveness of automated (electronic) monitoring systems compared with manual recording systems in identifying people at risk of clinical deterioration in general hospital ward settings?
- What are the sensitivities and specificities of track and trigger systems in different clinical settings?
- Can track and trigger systems that have higher sensitivities and specificities than existing scores be developed and validated?

Focused searches were performed to identify evidence on the use of track and trigger and EWSs and electronic monitoring systems for recognition of

patient deterioration. New evidence was identified in the 2020 surveillance review. This evidence and potential impact on guideline recommendations is summarised under the following sections: [physiological observations in acute hospital settings](#), [identifying patients whose clinical condition is deteriorating or is at risk of deterioration](#), [choice of physiological track and trigger system](#), and [physiological parameters to be used by track and trigger systems](#). Since it is considered that existing evidence is unlikely to have potential impact on recommendations, further research would be of value.

Response strategies for patients identified as having a deteriorating clinical condition

- What is the clinical and cost effectiveness of a structured educational programme to improve recognition of and response to acute illness compared with no structured programme in improving outcomes for people who clinically deteriorate in general hospital ward settings?

[Two RCTs and 1 intervention study](#) on provision of educational interventions were identified in the 2020 surveillance review. These studies assessed the provision of educational/training interventions in 2 RCTs and a cognitive aid booklet intervention in 1 intervention study. These interventions resulted in improved performance by professionals in simulated test settings, improved knowledge and other post-test scores and skills. Further research on this area would be of value to determine the optimal configuration of the educational intervention.

- What is the clinical and cost effectiveness of CCOS compared with usual care or educational outreach in improving health outcomes for patients who clinically deteriorate in general hospital ward settings? Such research should:
 - use a cluster RCT design conducted on multiple sites, with analysis of the cluster at hospital level rather than ward level
 - investigate a range of health outcomes, including mortality, morbidity, quality of life measures and patient satisfaction

- include a parallel qualitative process evaluation to help establish which components of outreach (a complex intervention) are likely to be most effective
- consider 24-hour critical care outreach as well as daytime outreach.

Three cluster RCTs were identified in the 2020 surveillance review and are summarised under [graded response strategy](#). Additional research would benefit this area.

Transfer of patients from critical care areas

- What is the clinical and cost effectiveness of providing structured educational advice (such as an information booklet) compared with usual care to patients who have been transferred from critical care areas back to general hospital ward settings?

Focused evidence searches were not performed in this area, but key relevant evidence identified in the course of the surveillance review was summarised. No evidence relating to this research recommendation was identified.

- What is the clinical and cost effectiveness of a transfer facilitator for patients transferred from critical care to a general ward environment? Such research could include outcome measures on:
 - patient satisfaction
 - time to discharge from acute hospital
 - destination when transferred.

Focused evidence searches were not performed in this area, but key relevant evidence identified in the course of the surveillance review was summarised. No evidence relating to this research recommendation was identified.

Additional research on these 2 research recommendations would be useful.

References

1. Merriel A, van der Nelson H, Merriel S, Bennett J, Donald F, Draycott T, et al. (2016) Identifying Deteriorating Patients Through Multidisciplinary

- Team Training. *American Journal of Medical Quality : the Official Journal of the American College of Medical Quality* 31(6):589–95
2. Petersen JA, Antonsen K, Rasmussen LS (2016) Frequency of early warning score assessment and clinical deterioration in hospitalized patients: A randomized trial. *Resuscitation* 101:91–6
 3. Prytherch DR, Smith GB, Schmidt PE, Featherstone PI (2010) ViEWS-- Towards a national early warning score for detecting adult inpatient deterioration. *Resuscitation* 81(8):932–7
 4. W.C. Yuan, C. Tao, Z.D. Dan et al. (2018) The significance of National Early Warning Score for predicting prognosis and evaluating conditions of patients in resuscitation room. *Hong Kong Journal of Emergency Medicine* 25(6):324–30
 5. Echevarria C, Steer J, Bourke SC (2019) Comparison of early warning scores in patients with COPD exacerbation: DECAF and NEWS score. *Thorax* 74(10):941–6
 6. Ghosh E, Eshelman L, Yang L, Carlson E, Lord B (2018) Early Deterioration Indicator: Data-driven approach to detecting deterioration in general ward. *Resuscitation* 122:99–105
 7. Brabrand M, Hallas P, Hansen SN, Jensen KM, Madsen JLB, Posth S (2017) Using scores to identify patients at risk of short term mortality at arrival to the acute medical unit: A validation study of six existing scores. *European Journal of Internal Medicine* 45:32–6
 8. Redfern OC, Pimentel MAF, Prytherch D, Meredith P, Clifton DA, Tarassenko L, et al. (2018) Predicting in-hospital mortality and unanticipated admissions to the intensive care unit using routinely collected blood tests and vital signs: Development and validation of a multivariable model. *Resuscitation* 133:75–81
 9. Chiu Y-D, Villar SS, Brand JW, Patteril M V, Morrice DJ, Clayton J, et al. (2019) Logistic early warning scores to predict death, cardiac arrest or

unplanned intensive care unit re-admission after cardiac surgery.

Anaesthesia

10. Smith GB, Prytherch DR, Jarvis S, Kovacs C, Meredith P, Schmidt PE, et al. (2016) A Comparison of the Ability of the Physiologic Components of Medical Emergency Team Criteria and the U.K. National Early Warning Score to Discriminate Patients at Risk of a Range of Adverse Clinical Outcomes. *Critical Care Medicine* 44(12):2171–81
11. Kwack WG, Lee DS, Min H, Choi YY, Yun M, Kim Y, et al. (2018) Evaluation of the SpO₂/FiO₂ ratio as a predictor of intensive care unit transfers in respiratory ward patients for whom the rapid response system has been activated. *PloS One* 13(7):e0201632
12. Mitsunaga T, Hasegawa I, Uzura M, Okuno K, Otani K, Ohtaki Y, et al. (2019) Comparison of the National Early Warning Score (NEWS) and the Modified Early Warning Score (MEWS) for predicting admission and in-hospital mortality in elderly patients in the pre-hospital setting and in the emergency department. *PeerJ* 7:e6947
13. Foy KE, Pearson J, Kettleby L, Lal N, Blackwood H, Bould MD (2019) Four early warning scores predict mortality in emergency surgical patients at University Teaching Hospital, Lusaka: a prospective observational study. *Canadian Journal of Anaesthesia = Journal Canadien d'anesthésie*
14. Abbott TEF, Torrance HDT, Cron N, Vaid N, Emmanuel J (2016) A single-centre cohort study of National Early Warning Score (NEWS) and near patient testing in acute medical admissions. *European Journal of Internal Medicine* 35:78–82
15. Dunder ZD, Kocak S, Girisgin AS (2019) Lactate and NEWS-L are fair predictors of mortality in critically ill geriatric emergency department patients. *The American Journal of Emergency Medicine*
16. Jo S, Yoon J, Lee JB, Jin Y, Jeong T, Park B (2016) Predictive value of

- the National Early Warning Score-Lactate for mortality and the need for critical care among general emergency department patients. *Journal of Critical Care* 36:60–8
17. Malycha J, Farajidavar N, Pimentel MAF, Redfern O, Clifton DA, Tarassenko L, et al. (2019) The effect of fractional inspired oxygen concentration on early warning score performance: A database analysis. *Resuscitation* 139:192–9
 18. Luís L, Nunes C (2018) Short National Early Warning Score - Developing a Modified Early Warning Score. *Australian Critical Care : Official Journal of the Confederation of Australian Critical Care Nurses* 31(6):376–81
 19. Eckart A, Hauser SI, Kutz A, Haubitz S, Hausfater P, Amin D, et al. (2019) Combination of the National Early Warning Score (NEWS) and inflammatory biomarkers for early risk stratification in emergency department patients: results of a multinational, observational study. *BMJ Open* 9(1):e024636
 20. Rasmussen LJH, Ladelund S, Haupt TH, Ellekilde GE, Eugen-Olsen J, Andersen O (2018) Combining National Early Warning Score With Soluble Urokinase Plasminogen Activator Receptor (suPAR) Improves Risk Prediction in Acute Medical Patients: A Registry-Based Cohort Study. *Critical Care Medicine* 46(12):1961–8
 21. Pimentel MAF, Redfern OC, Gerry S, Collins GS, Malycha J, Prytherch D, et al. (2019) A comparison of the ability of the National Early Warning Score and the National Early Warning Score 2 to identify patients at risk of in-hospital mortality: A multi-centre database study. *Resuscitation* 134:147–56
 22. Srivilaithon W, Amnuaypattanapon K, Limjindaporn C, Imsuwan I, Daorattanachai K, Dasanadeba I, et al. (2019) Predictors of in-hospital cardiac arrest within 24 h after emergency department triage: A case-control study in urban Thailand. *Emergency medicine Australasia : EMA*

31(5):843–50

23. Abbott TEF, Vaid N, Ip D, Cron N, Wells M, Torrance HDT, et al. (2015) A single-centre observational cohort study of admission National Early Warning Score (NEWS). *Resuscitation* 92:89–93
24. Fernando SM, Fox-Robichaud AE, Rochweg B, Cardinal P, Seely AJE, Perry JJ, et al. (2019) Prognostic accuracy of the Hamilton Early Warning Score (HEWS) and the National Early Warning Score 2 (NEWS2) among hospitalized patients assessed by a rapid response team. *Critical Care (London, England)* 23(1):60
25. Martín-Rodríguez F, López-Izquierdo R, Del Pozo Vegas C, Delgado Benito JF, Carbajosa Rodríguez V, Diego Rasilla MN, et al. (2019) Accuracy of National Early Warning Score 2 (NEWS2) in Prehospital Triage on In-Hospital Early Mortality: A Multi-Center Observational Prospective Cohort Study. *Prehospital and Disaster Medicine* 34(6):610–8
26. Martín-Rodríguez F, Castro-Villamor MÁ, Del Pozo Vegas C, Martín-Conty JL, Mayo-Iscar A, Delgado Benito JF, et al. (2019) Analysis of the early warning score to detect critical or high-risk patients in the prehospital setting. *Internal and Emergency Medicine* 14(4):581–9
27. Abbott TEF, Cron N, Vaid N, Ip D, Torrance HDT, Emmanuel J (2018) Pre-hospital National Early Warning Score (NEWS) is associated with in-hospital mortality and critical care unit admission: A cohort study. *Annals of Medicine and Surgery* (2012) 27:17–21
28. Bedoya AD, Clement ME, Phelan M, Steorts RC, O'Brien C, Goldstein BA (2019) Minimal Impact of Implemented Early Warning Score and Best Practice Alert for Patient Deterioration. *Critical Care Medicine* 47(1):49–55
29. Faisal M, Scally A, Elgaali MA, Richardson D, Beatson K, Mohammed MA (2018) The National Early Warning Score and its subcomponents

- recorded within +/-24 h of emergency medical admission are poor predictors of hospital-acquired acute kidney injury. *Clinical Medicine (London, England)* 18(1):47–53
30. Hydes TJ, Meredith P, Schmidt PE, Smith GB, Prytherch DR, Aspinall RJ (2018) National Early Warning Score Accurately Discriminates the Risk of Serious Adverse Events in Patients With Liver Disease. *Clinical Gastroenterology and Hepatology : the Official Clinical Practice Journal of the American Gastroenterological Association* 16(10):1657–1666.e10
 31. Kim D, Jo S, Lee JB, Jin Y, Jeong T, Yoon J, et al. (2018) Comparison of the National Early Warning Score+Lactate score with the pre-endoscopic Rockall, Glasgow-Blatchford, and AIMS65 scores in patients with upper gastrointestinal bleeding. *Clinical and Experimental Emergency Medicine* 5(4):219–29
 32. Lee YS, Choi JW, Park YH, Chung C, Park D II, Lee JE, et al. (2018) Evaluation of the efficacy of the National Early Warning Score in predicting in-hospital mortality via the risk stratification. *Journal of Critical Care* 47:222–6
 33. Sutherasan Y, Theerawit P, Suporn A, Nongnuch A, Phanachet P, Kositchaiwat C (2018) The impact of introducing the early warning scoring system and protocol on clinical outcomes in tertiary referral university hospital. *Therapeutics and Clinical Risk Management* 14:2089–95
 34. Silcock DJ, Corfield AR, Rooney KD, Staines H (2018) Superior performance of National Early Warning Score compared with quick Sepsis-related Organ Failure Assessment Score in predicting adverse outcomes: a retrospective observational study of patients in the prehospital setting. *European Journal of Emergency Medicine : official journal of the European Society for Emergency Medicine*
 35. Pirneskoski J, Kuisma M, Olkkola KT, Nurmi J (2019) Prehospital National Early Warning Score predicts early mortality. *Acta*

36. Hogan H, Hutchings A, Wulff J, Carver C, Holdsworth E, Welch J, et al. (2019) Interventions to reduce mortality from in-hospital cardiac arrest: a mixed-methods study. *Health Services and Delivery Research* 7(2)
37. Hodgson LE, Congleton J, Venn R, Forni LG, Roderick PJ (2018) NEWS 2 - too little evidence to implement? *Clinical Medicine (London, England)* 18(5):371–3
38. Kim W-Y, Lee J, Lee J-R, Jung YK, Kim HJ, Huh JW, et al. (2017) A risk scoring model based on vital signs and laboratory data predicting transfer to the intensive care unit of patients admitted to gastroenterology wards. *Journal of Critical Care* 40:213–7
39. Wong YK, Lui CT, Li KK, Wong CY, Lee MM, Tong WL, et al. (2016) Prediction of en-route complications during interfacility transport by outcome predictive scores in ED. *The American Journal of Emergency Medicine* 34(5):877–82
40. Churpek MM, Yuen TC, Winslow C, Meltzer DO, Kattan MW, Edelson DP (2016) Multicenter Comparison of Machine Learning Methods and Conventional Regression for Predicting Clinical Deterioration on the Wards. *Critical Care Medicine* 44(2):368–74
41. Jafar AJN, Junghans C, Kwok CS, Hymers C, Monk KJ, Gold E, et al. (2016) Do physiological scoring and a novel point of care metabolic screen predict 48-h outcome in admissions from the emergency department resuscitation area?. *European Journal of Emergency Medicine : official journal of the European Society for Emergency Medicine* 23(2):130–6
42. Wei X, Ma H, Liu R, Zhao Y (2019) Comparing the effectiveness of three scoring systems in predicting adult patient outcomes in the emergency department. *Medicine* 98(5):e14289
43. Dundar ZD, Ergin M, Karamercan MA, Ayranci K, Colak T, Tuncar A, et

- al. (2016) Modified Early Warning Score and VitalPac Early Warning Score in geriatric patients admitted to emergency department. *European Journal of Emergency Medicine : Official Journal of the European Society for Emergency Medicine* 23(6):406–12
44. Norman S, DeCicco F, Sampson J, Fraser IM (2018) Emergency Room Safer Transfer of Patients (ER-STOP): a quality improvement initiative at a community-based hospital to improve the safety of emergency room patient handovers. *BMJ Open* 8(12):e019553
45. Weenk M, Koeneman M, van de Belt TH, Engelen LJLPG, van Goor H, Bredie SJH (2019) Wireless and continuous monitoring of vital signs in patients at the general ward. *Resuscitation* 136:47–53
46. Bellomo R, Chan M, Guy C, Proimos H, Franceschi F, Crisman M, et al. (2018) Laboratory alerts to guide early intensive care team review in surgical patients: A feasibility, safety, and efficacy pilot randomized controlled trial. *Resuscitation* 133:167–72
47. Churpek MM, Adhikari R, Edelson DP (2016) The value of vital sign trends for detecting clinical deterioration on the wards. *Resuscitation* 102:1–5
48. Douw G, Huisman-de Waal G, van Zanten ARH, van der Hoeven JG, Schoonhoven L (2016) Nurses' "worry" as predictor of deteriorating surgical ward patients: A prospective cohort study of the Dutch-Early-Nurse-Worry-Indicator-Score. *International Journal of Nursing Studies* 59:134–40
49. Dziadzko MA, Novotny PJ, Sloan J, Gajic O, Hrasevich V, Mirhaji P, et al. (2018) Multicenter derivation and validation of an early warning score for acute respiratory failure or death in the hospital. *Critical Care (London, England)* 22(1):286
50. Kwon J-M, Lee Y, Lee Y, Lee S, Park H, Park J (2018) Validation of deep-learning-based triage and acuity score using a large national

dataset. PloS One 13(10):e0205836

51. Linnen DT, Escobar GJ, Hu X, Scruth E, Liu V, Stephens C (2019) Statistical Modeling and Aggregate-Weighted Scoring Systems in Prediction of Mortality and ICU Transfer: A Systematic Review. *Journal of Hospital Medicine* 14(3):161–9
52. Merriel A, Murove BT, Merriel SWD, Sibanda T, Moyo S, Crofts J (2017) Implementation of a modified obstetric early warning system to improve the quality of obstetric care in Zimbabwe. *International Journal of Gynaecology and Obstetrics: the Official Organ of the International Federation of Gynaecology and Obstetrics* 136(2):175–9
53. O’Connell A, Flabouris A, Kim SW, Horwood C, Hakendorf P, Thompson CH (2016) A newly designed observation and response chart’s effect upon adverse inpatient outcomes and rapid response team activity. *Internal Medicine Journal* 46(8):909–16
54. Peek KN, Gillham M (2017) Is the New Zealand Early Warning Score useful following cardiac surgery?. *The New Zealand Medical Journal* 130(1461):9–14
55. Romero-Brufau S, Morlan BW, Johnson M, Hickman J, Kirkland LL, Naessens JM, et al. (2017) Evaluating Automated Rules for Rapid Response System Alarm Triggers in Medical and Surgical Patients. *Journal of Hospital Medicine* 12(4):217–23
56. Ryan HM, Jones MA, Payne BA, Sharma S, Hutfield AM, Lee T, et al. (2017) Validating the Performance of the Modified Early Obstetric Warning System Multivariable Model to Predict Maternal Intensive Care Unit Admission. *Journal of Obstetrics and Gynaecology Canada : JOGC = Journal d’obstetrique et Gynecologie du Canada : JOGC* 39(9):728–733.e3
57. Street M, Phillips NM, Mohebbi M, Kent B (2017) Effect of a newly designed observation, response and discharge chart in the Post

- Anaesthesia Care Unit on patient outcomes: a quasi-experimental study in Australia. *BMJ Open* 7(12):e015149
58. Williams TA, Tohira H, Finn J, Perkins GD, Ho KM (2016) The ability of early warning scores (EWS) to detect critical illness in the prehospital setting: A systematic review. *Resuscitation* 102:35–43
 59. Patel R, Nugawela MD, Edwards HB, Richards A, Le Roux H, Pullyblank A, et al. (2018) Can early warning scores identify deteriorating patients in pre-hospital settings? A systematic review. *Resuscitation* 132:101–11
 60. Bartkowiak B, Snyder AM, Benjamin A, Schneider A, Twu NM, Churpek MM, et al. (2019) Validating the Electronic Cardiac Arrest Risk Triage (eCART) Score for Risk Stratification of Surgical Inpatients in the Postoperative Setting: Retrospective Cohort Study. *Annals of Surgery* 269(6):1059–63
 61. Fletcher GS, Aaronson BA, White AA, Julka R (2017) Effect of a Real-Time Electronic Dashboard on a Rapid Response System. *Journal of Medical Systems* 42(1):5
 62. Gagne C, Fetzer S (2018) Early Warning Score Communication Bundle: A Pilot Study. *American Journal of Critical Care : an official publication, American Association of Critical-Care Nurses* 27(3):238–42
 63. Green M, Lander H, Snyder A, Hudson P, Churpek M, Edelson D (2018) Comparison of the Between the Flags calling criteria to the MEWS, NEWS and the electronic Cardiac Arrest Risk Triage (eCART) score for the identification of deteriorating ward patients. *Resuscitation* 123:86–91
 64. Heller AR, Mees ST, Lauterwald B, Reeps C, Koch T, Weitz J (2018) Detection of Deteriorating Patients on Surgical Wards Outside the ICU by an Automated MEWS-Based Early Warning System With Paging Functionality. *Annals of Surgery*
 65. Hu SB, Wong DJL, Correa A, Li N, Deng JC (2016) Prediction of Clinical

- Deterioration in Hospitalized Adult Patients with Hematologic Malignancies Using a Neural Network Model. *PloS One* 11(8):e0161401
66. Imperato J, Mehegan T, Henning DJ, Patrick J, Bushey C, Setnik G, et al. (2017) Can an emergency department clinical “triggers” program based on abnormal vital signs improve patient outcomes?. *CJEM* 19(4):249–55
 67. Kang MA, Churpek MM, Zadavec FJ, Adhikari R, Twu NM, Edelson DP (2016) Real-Time Risk Prediction on the Wards: A Feasibility Study. *Critical Care Medicine* 44(8):1468–73
 68. Keim-Malpass J, Clark MT, Lake DE, Moorman JR (2019) Towards development of alert thresholds for clinical deterioration using continuous predictive analytics monitoring. *Journal of Clinical Monitoring and computing*
 69. Kipnis P, Turk BJ, Wulf DA, LaGuardia JC, Liu V, Churpek MM, et al. (2016) Development and validation of an electronic medical record-based alert score for detection of inpatient deterioration outside the ICU. *Journal of Biomedical Informatics* 64:10–9
 70. Pappas PA, Tirelli L, Shaffer J, Gettings S (2016) Projecting Critical Care Beyond the ICU: An Analysis of Tele-ICU Support for Rapid Response Teams. *Telemedicine Journal and e-health : the official journal of the American Telemedicine Association* 22(6):529–33
 71. Subbe CP, Duller B, Bellomo R (2017) Effect of an automated notification system for deteriorating ward patients on clinical outcomes. *Critical Care (London, England)* 21(1):52
 72. Watkinson PJ, Pimentel MAF, Clifton DA, Tarassenko L (2018) Manual centile-based early warning scores derived from statistical distributions of observational vital-sign data. *Resuscitation* 129:55–60
 73. Weller RS, Foard KL, Harwood TN (2018) Evaluation of a wireless, portable, wearable multi-parameter vital signs monitor in hospitalized

- neurological and neurosurgical patients. *Journal of Clinical Monitoring and computing* 32(5):945–51
74. Wilson FP, Shashaty M, Testani J, Aqeel I, Borovskiy Y, Ellenberg SS, et al. (2015) Automated, electronic alerts for acute kidney injury: a single-blind, parallel-group, randomised controlled trial. *Lancet (London, England)* 385(9981):1966–74
 75. Wilson SJ, Wong D, Pullinger RM, Way R, Clifton DA, Tarassenko L (2016) Analysis of a data-fusion system for continuous vital sign monitoring in an emergency department. *European Journal of Emergency Medicine : official journal of the European Society for Emergency Medicine* 23(1):28–32
 76. Pedersen NE, Oestergaard D, Lippert A (2016) End points for validating early warning scores in the context of rapid response systems: a Delphi consensus study. *Acta Anaesthesiologica Scandinavica* 60(5):616–22
 77. Moss TJ, Clark MT, Calland JF, Enfield KB, Voss JD, Lake DE, et al. (2017) Cardiorespiratory dynamics measured from continuous ECG monitoring improves detection of deterioration in acute care patients: A retrospective cohort study. *PloS One* 12(8):e0181448
 78. Kyi M, Wraight PR, Rowan LM, Marley KA, Colman PG, Furlanos S (2018) Glucose alert system improves health professional responses to adverse glycaemia and reduces the number of hyperglycaemic episodes in non-critical care inpatients. *Diabetic Medicine : a Journal of the British Diabetic Association* 35(6):816–23
 79. Liaw SY, Wong LF, Ang SBL, Ho JTY, Siau C, Ang ENK (2016) Strengthening the afferent limb of rapid response systems: an educational intervention using web-based learning for early recognition and responding to deteriorating patients. *BMJ Quality & Safety* 25(6):448–56
 80. Johnston MJ, Arora S, Pucher PH, McCartan N, Reissis Y, Chana P, et

- al. (2016) Improving Escalation of Care: A Double-blinded Randomized Controlled Trial. *Annals of Surgery* 263(3):421–6
81. Mitchell OJL, Lehr A, Lo M, Kam LM, Andriotis A, Felner K, et al. (2019) Development and Evaluation of a Cognitive Aid Booklet for Use in Rapid Response Scenarios. *Simulation in Healthcare : Journal of the Society for Simulation in Healthcare* 14(4):217–22
82. Ganju A, Kapitola K, Chalwin R (2019) Modifications to predefined rapid response team calling criteria: prevalence, characteristics and associated outcomes. *Critical Care and Resuscitation : journal of the Australasian Academy of Critical Care Medicine* 21(1):32–8
83. Jeddian A, Hemming K, Lindenmeyer A, Rashidian A, Sayadi L, Jafari N, et al. (2016) Evaluation of a critical care outreach service in a middle-income country: A stepped wedge cluster randomized trial and nested qualitative study. *Journal of Critical Care* 36:212–7
84. Gershengorn HB, Xu Y, Chan CW, Armony M, Gong MN (2016) The Impact of Adding a Physician Assistant to a Critical Care Outreach Team. *PloS One* 11(12):e0167959
85. Gouda AM, Alqahtani SM (2016) Do not resuscitate: An expanding role for critical care response team. *Indian Journal of Critical Care Medicine : peer-reviewed, official publication of Indian Society of Critical Care Medicine* 20(3):146–9
86. Al-Rajhi A, Mardini L, Jayaraman D (2016) The Impact of Implementation of an ICU Consult Service on Hospital-Wide Outcomes and ICU-Specific Outcomes. *Journal of Intensive Care Medicine* 31(7):478–84
87. Bergmeir C, Bilgrami I, Bain C, Webb GI, Orosz J, Pilcher D (2017) Designing a more efficient, effective and safe Medical Emergency Team (MET) service using data analysis. *PloS One* 12(12):e0188688
88. Silva R, Saraiva M, Cardoso T, Aragao IC (2016) Medical Emergency

- Team: How do we play when we stay? Characterization of MET actions at the scene. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 24:33
89. Kawaguchi R, Nakada T-A, Oshima T, Abe R, Matsumura Y, Oda S (2015) Reduction of unexpected serious adverse events after introducing medical emergency team. *Acute Medicine & Surgery* 2(4):244–9
 90. Sen S, Acash G, Sarwar A, Lei Y, Dargin JM (2017) Utility and diagnostic accuracy of bedside lung ultrasonography during medical emergency team (MET) activations for respiratory deterioration. *Journal of Critical Care* 40:58–62
 91. Pirret AM, Takerei SF, Kazula LM (2015) The effectiveness of a patient at risk team comprised of predominantly ward experienced nurses: A before and after study. *Intensive & Critical Care Nursing* 31(3):133–40
 92. Maharaj R, Raffaele I, Wendon J (2015) Rapid response systems: a systematic review and meta-analysis. *Critical Care (London, England)* 19:254
 93. Rocha HAL, Alcantara AC de C, Rocha SGM, Toscano CM (2018) Effectiveness of rapid response teams in reducing intrahospital cardiac arrests and deaths: a systematic review and meta-analysis. *Revista Brasileira de Terapia Intensiva* 30(3):366–75
 94. Solomon RS, Corwin GS, Barclay DC, Quddusi SF, Dannenberg MD (2016) Effectiveness of rapid response teams on rates of in-hospital cardiopulmonary arrest and mortality: A systematic review and meta-analysis. *Journal of Hospital Medicine* 11(6):438–45
 95. Chen J, Bellomo R, Flabouris A, Hillman K, Assareh H, Ou L (2015) Delayed Emergency Team Calls and Associated Hospital Mortality: A Multicenter Study. *Critical Care Medicine* 43(10):2059–65
 96. Haegdorens F, Van Bogaert P, Roelant E, De Meester K, Misselyn M,

- Wouters K, et al. (2018) The introduction of a rapid response system in acute hospitals: A pragmatic stepped wedge cluster randomised controlled trial. *Resuscitation* 129:127–34
97. YekeFallah L, Eskandari Z, Shahrokhi A, Javadi A (2018) Effect of rapid response nursing team on outcome of patient care. *Trauma Monthly* 23(4):e14488
98. Jung B, Daurat A, De Jong A, Chanques G, Mahul M, Monnin M, et al. (2016) Rapid response team and hospital mortality in hospitalized patients. *Intensive Care Medicine* 42(4):494–504
99. Mullany D V, Ziegenfuss M, Goleby MA, Ward HE (2016) Improved hospital mortality with a low MET dose: the importance of a modified early warning score and communication tool. *Anaesthesia and Intensive Care* 44(6):734–41
100. Sebat F, Vandegrift MA, Childers S, Lighthall GK (2018) A Novel Bedside-Focused Ward Surveillance and Response System. *Joint Commission journal on quality and patient safety* 44(2):94–100
101. Pain C, Green M, Duff C, Hyland D, Pantle A, Fitzpatrick K, et al. (2017) Between the flags: implementing a safety-net system at scale to recognise and manage deteriorating patients in the New South Wales Public Health System. *International Journal for Quality in Health Care : journal of the International Society for Quality in Health Care* 29(1):130–6
102. Aitken LM, Chaboyer W, Vaux A, Crouch S, Burmeister E, Daly M, et al. (2015) Effect of a 2-tier rapid response system on patient outcome and staff satisfaction. *Australian Critical Care : official journal of the Confederation of Australian Critical Care Nurses* 28(3):107–15
103. Massey D, Aitken LM, Chaboyer W (2015) The impact of a nurse led rapid response system on adverse, major adverse events and activation of the medical emergency team. *Intensive & Critical Care Nursing*

31(2):83–90

104. Feih J, Peppard WJ, Katz M (2017) Pharmacist involvement on a rapid response team. *American Journal of Health-System Pharmacy : AJHP : official journal of the American Society of Health-System Pharmacists* 74(5supplement1):10-s16
105. Davis DP, Aguilar SA, Graham PG, Lawrence B, Sell RE, Minokadeh A, et al. (2015) A novel configuration of a traditional rapid response team decreases non-intensive care unit arrests and overall hospital mortality. *Journal of Hospital Medicine* 10(6):352–7
106. Yousaf M, Bano S, Attaur-Rehman M, Nazar CMJ, Qadeer A, Khudaidad S, et al. (2018) Comparison of Hospital-Wide Code Rates and Mortality Before and After the Implementation of a Rapid Response Team. *Cureus* 10(1):e2043
107. Danesh V, Neff D, Jones TL, Aroian K, Unruh L, Andrews D, et al. (2019) Can proactive rapid response team rounding improve surveillance and reduce unplanned escalations in care? A controlled before and after study. *International Journal of Nursing Studies* 91:128–33
108. Almeida MC, Portela MC, Paiva EP, Guimaraes RR, Pereira Neto WC, Cardoso PR, et al. (2019) Implementation of a rapid response team in a large nonprofit Brazilian hospital: improving the quality of emergency care through Plan-Do-Study-Act. *Revista Brasileira de Terapia Intensiva* 31(2):217–26
109. Amaral ACK-B, McDonald A, Coburn NG, Xiong W, Shojania KG, Fowler RA, et al. (2015) Expanding the scope of Critical Care Rapid Response Teams: a feasible approach to identify adverse events. A prospective observational cohort. *BMJ Quality & Safety* 24(12):764–8
110. Blotsky A, Mardini L, Jayaraman D (2016) Impact of a Local Low-Cost Ward-Based Response System in a Canadian Tertiary Care Hospital.

111. Brunsveld-Reinders AH, Ludikhuizen J, Dijkgraaf MGW, Arbous MS, de Jonge E, group C study (2016) Unexpected versus all-cause mortality as the endpoint for investigating the effects of a Rapid Response System in hospitalized patients. *Critical Care (London, England)* 20(1):168
112. Considine J, Rawet J, Currey J (2015) The effect of a staged, emergency department specific rapid response system on reporting of clinical deterioration. *Australasian Emergency Nursing Journal : AENJ* 18(4):218–26
113. Hatlem T, Jones C, Woodard EK (2018) Reducing Mortality and Avoiding Preventable ICU Utilization: Analysis of a Successful Rapid Response Program Using APR DRGs. *Journal for Healthcare Quality : official publication of the National Association for Healthcare Quality*
114. Joshi K, Campbell V, Landy M, Anstey CM, Gooch R (2017) The effect of Rapid Response System revision on standard and specific intensive care unit outcomes in a regional hospital. *Anaesthesia and Intensive Care* 45(3):369–74
115. Ludikhuizen J, Brunsveld-Reinders AH, Dijkgraaf MGW, Smorenburg SM, de Rooij SEJA, Adams R, et al. (2015) Outcomes Associated With the Nationwide Introduction of Rapid Response Systems in The Netherlands. *Critical Care Medicine* 43(12):2544–51
116. Oh TK, Park YM, Do S-H, Hwang J-W, Song I-A (2017) ROSC rates and live discharge rates after cardiopulmonary resuscitation by different CPR teams - a retrospective cohort study. *BMC Anesthesiology* 17(1):166
117. Study Investigators TCMETM, Concord Medical Emergency Team (MET) 2 Study Investigators (2019) Outcomes following changing from a two-tiered to a three-tiered hospital rapid response system. *Australian*

Health Review : a publication of the Australian Hospital Association
43(2):178–87

118. Grieve R, O'Neill S, Basu A, Keele L, Rowan KM, Harris S (2019) Analysis of Benefit of Intensive Care Unit Transfer for Deteriorating Ward Patients: A Patient-Centered Approach to Clinical Evaluation. *JAMA Network Open* 2(2):e187704
119. Yang S, Wang Z, Liu Z, Wang J, Ma L (2016) Association between time of discharge from ICU and hospital mortality: a systematic review and meta-analysis. *Critical Care (London, England)* 20(1):390
120. Vollam S, Dutton S, Lamb S, Petrinic T, Young JD, Watkinson P (2018) Out-of-hours discharge from intensive care, in-hospital mortality and intensive care readmission rates: a systematic review and meta-analysis. *Intensive Care Medicine* 44(7):1115–29
121. Azevedo LCP, de Souza IA, Zygun DA, Stelfox HT, Bagshaw SM (2015) Association Between Nighttime Discharge from the Intensive Care Unit and Hospital Mortality: A Multi-Center Retrospective Cohort Study. *BMC Health Services Research* 15:378
122. Correa TD, Ponzoni CR, Filho RR, Neto AS, Chaves RC de F, Pardini A, et al. (2018) Nighttime intensive care unit discharge and outcomes: A propensity matched retrospective cohort study. *PloS One* 13(12):e0207268
123. Klepstad PK, Nordseth T, Sikora N, Klepstad P (2019) Use of National Early Warning Score for observation for increased risk for clinical deterioration during post-ICU care at a surgical ward. *Therapeutics and Clinical Risk Management* 15:315–22
124. Alarhayem AQ, Muir MT, Jenkins DJ, Pruitt BA, Eastridge BJ, Purohit MP, et al. (2019) Application of electronic medical record-derived analytics in critical care: Rothman Index predicts mortality and readmissions in surgical intensive care unit patients. *The Journal of*

125. Gotur DB, Masud F, Paraniyam J, Zimmerman JL (2018) Analysis of Rothman Index Data to Predict Postdischarge Adverse Events in a Medical Intensive Care Unit. *Journal of Intensive Care Medicine* :885066618770128
126. Parenmark F, Karlström G, Nolin T, Fredrikson M, Walther SM (2019) Reducing night-time discharge from intensive care. A nationwide improvement project with public display of ICU outcomes. *Journal of Critical Care* 49:7–13
127. Rosa RG, Roehrig C, Oliveira RP de, Maccari JG, Antonio ACP, Castro P de S, et al. (2015) Comparison of Unplanned Intensive Care Unit Readmission Scores: A Prospective Cohort Study. *PloS One* 10(11):e0143127
128. Desautels T, Das R, Calvert J, Trivedi M, Summers C, Wales DJ, et al. (2017) Prediction of early unplanned intensive care unit readmission in a UK tertiary care hospital: a cross-sectional machine learning approach. *BMJ Open* 7(9):e017199
129. Uppanisakorn S, Bhurayanontachai R, Boonyarat J, Kaewpradit J (2018) National Early Warning Score (NEWS) at ICU discharge can predict early clinical deterioration after ICU transfer. *Journal of Critical Care* 43:225–9
130. Martin LA, Kilpatrick JA, Al-Dulaimi R, Mone MC, Tonna JE, Barton RG, et al. (2019) Predicting ICU readmission among surgical ICU patients: Development and validation of a clinical nomogram. *Surgery* 165(2):373–80
131. Heselmans A, van Krieken J, Cootjans S, Nagels K, Filliers D, Dillen K, et al. (2015) Medication review by a clinical pharmacist at the transfer point from ICU to ward: a randomized controlled trial. *Journal of Clinical Pharmacy and Therapeutics* 40(5):578–83

132. Tully AP, Hammond DA, Li C, Jarrell AS, Kruer RM (2019) Evaluation of Medication Errors at the Transition of Care From an ICU to Non-ICU Location. *Critical Care Medicine* 47(4):543–9
133. Bosma LBE, Hunfeld NGM, Quax RAM, Meuwese E, Melief PHGJ, van Bommel J, et al. (2018) The effect of a medication reconciliation program in two intensive care units in the Netherlands: a prospective intervention study with a before and after design. *Annals of Intensive Care* 8(1):19
134. Hoffman RL, Saucier J, Dasani S, Collins T, Holena DN, Fitzpatrick M, et al. (2017) Development and implementation of a risk identification tool to facilitate critical care transitions for high-risk surgical patients. *International Journal for Quality in Health Care : journal of the International Society for Quality in Health Care* 29(3):412–9
135. Stelfox HT, Bastos J, Niven DJ, Bagshaw SM, Turin TC, Gao S (2016) Critical care transition programs and the risk of readmission or death after discharge from ICU. *Intensive Care Medicine* 42(3):401–10
136. Choi S, Lee J, Shin Y, Lee J, Jung J, Han M, et al. (2016) Effects of a medical emergency team follow-up programme on patients discharged from the medical intensive care unit to the general ward: a single-centre experience. *Journal of Evaluation in Clinical Practice* 22(3):356–62
137. Kheir F, Shawwa K, Nguyen D, Alraiyes AH, Simeone F, Nielsen ND (2016) A 24-Hour Postintensive Care Unit Transition-of-Care Model Shortens Hospital Stay. *Journal of Intensive Care Medicine* 31(9):597–602
138. Fabes J, Seligman W, Barrett C, McKechnie S, Griffiths J (2017) Does the implementation of a novel intensive care discharge risk score and nurse-led inpatient review tool improve outcome? A prospective cohort study in two intensive care units in the UK. *BMJ Open* 7(12):e018322
139. So HM, Yan WW, Chair SY (2018) A nurse-led critical care outreach

program to reduce readmission to the intensive care unit: A quasi-experimental study with a historical control group. *Australian Critical Care* : official journal of the Confederation of Australian Critical Care Nurses

140. van Sluisveld N, Bakhshi-Raiez F, de Keizer N, Holman R, Wester G, Wollersheim H, et al. (2017) Variation in rates of ICU readmissions and post-ICU in-hospital mortality and their association with ICU discharge practices. *BMC Health Services Research* 17(1):281
141. Stelfox HT, Leigh JP, Dodek PM, Turgeon AF, Forster AJ, Lamontagne F, et al. (2017) A multi-center prospective cohort study of patient transfers from the intensive care unit to the hospital ward. *Intensive Care Medicine* 43(10):1485–94

© NICE 2019. All rights reserved. Subject to [Notice of rights](#).