

National Institute for Health and Care Excellence

Early submission

Caesarean birth

Review questions

NICE guideline CG132 (update)

Review questions

August 2020

Early submission

Developed by the National Guideline Alliance which is a part of the Royal College of Obstetricians and Gynaecologists

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2020. All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

Review questions	5
-------------------------------	----------

Review questions

Review questions	
A	What are the benefits and risks (short and long-term) of planned caesarean birth compared with planned vaginal birth at term for women and neonates/infants/children?
B	What methods, apart from prophylactic antibiotics, should be used to reduce infectious morbidity in women having a caesarean birth?
C	What are the procedures to prevent and manage hypothermia and shivering in women having a caesarean birth in the pre-operative, peri-operative and post-operative periods?
D	What is the efficacy of single-layer closure of the uterus compared with double layer closure at caesarean birth?
E	What post-operative monitoring is required for women who have received intrathecal or epidural opioids at the time of caesarean birth, to identify or prevent potential complications (including the duration, frequency and features to be monitored)?
F	Are opioids safe and effective for pain management after caesarean birth?