

HTG246 – Distal iliotibial band lengthening for refractory greater trochanteric pain syndrome

SNOMED CT provides clinical terms for entry into the patient record to record clinical information relevant to that encounter; the mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure and device:

SNOMED CT preferred term (concept ID)

Division of iliotibial band (56142003)

OPCS-4 code(s):

T54.8 Other specified division of fascia

Z90.3 Upper leg NEC

Note: The NHS Classifications Service has advised NICE that currently these are the most suitable OPCS-4 codes to describe this procedure. The OPCS-4 classification is designed to categorise procedures for analysis and it is not always possible to identify a procedure uniquely.

Diagnosis or health condition:

SNOMED CT preferred term (concept ID)

Trochanteric bursitis (7674000)

Clinical coding recommendations for NICE guidance

For each published interventional procedure and medical technologies guidance, we work with NHS Digital to provide relevant clinical coding information.

SNOMED CT provides clinical terms for entry into the patient record to store clinical information relevant to that encounter.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

The UK Edition of SNOMED CT is managed by the Clinical Terminology Service of NHS Digital. For further information including licensing, see [UK Terminology Centre — NHS Digital](#).

The Clinical Classifications Service of NHS Digital is the central definitive source for clinical coding guidance and determines the coding standards associated with the classifications (OPCS-4 and ICD-10) to be used across the NHS. The Clinical Classifications Service and NICE work collaboratively to ensure the most appropriate classification codes are provided. [Clinical Classifications Service — NHS Digital](#).