

## **HTG29 - Complete cytoreduction for pseudomyxoma peritonei (Sugarbaker technique)**

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS-4 code(s):**

This procedure cannot be expressed in the OPCS-4 classification by a single code. The current guidance is to code each organ removed as per normal coding rules, and to combine this with the ICD-10 diagnosis code.

For the intraperitoneal chemotherapy, the following OPCS-4 code is assigned:

**T48.2 Introduction of cytotoxic substance into peritoneal cavity**

**Note:** The collection of codes classifying procurement is not required, therefore codes from categories **X70** and **X71** should not be assigned unless there is a local requirement to do so.

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**C78.6 Secondary malignant neoplasm of retroperitoneum and peritoneum**

---

## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).