

# Laparoscopic live donor simple nephrectomy

Understanding NICE guidance –  
information for people considering the  
procedure, and for the public

May 2004



Corporate member of  
Plain English Campaign.  
Committed to clearer communication.

**197**

## **Laparoscopic live donor simple nephrectomy**

### **Understanding NICE guidance – information for people considering the procedure, and for the public**

**Issue date:** May 2004

#### **To order copies**

Copies of this booklet can be ordered from the NHS Response Line; telephone 0870 1555 455 and quote reference number N0569. A version in Welsh and English is also available, reference number N0570. Mae fersiwn yn Gymraeg ac yn Saesneg ar gael hefyd, rhif cyfeirnod N0570. The NICE interventional procedures guidance on which this information is based is available from the NICE website ([www.nice.org.uk](http://www.nice.org.uk)). Copies can also be obtained from the NHS Response Line, reference number N0568.

---

## **National Institute for Clinical Excellence**

MidCity Place  
71 High Holborn  
London  
WC1V 6NA

Website: [www.nice.org.uk](http://www.nice.org.uk)

ISBN: 978-1-4731-8748-1

Published by the National Institute for Clinical Excellence

May 2004

Typeset by Icon Design, Eton

Print on Demand

© National Institute for Clinical Excellence, May 2004. All rights reserved. This material may be freely reproduced for educational and not-for-profit purposes within the NHS. No reproduction by or for commercial organisations is allowed without the express written permission of the National Institute for Clinical Excellence.

## Contents

About this information	4
About laparoscopic live donor simple nephrectomy	5
What has NICE decided?	8
What the decision means for you	9
Further information	10

## About this information

This information describes the guidance that the National Institute for Clinical Excellence (NICE) has issued to the NHS on a procedure called laparoscopic live donor simple nephrectomy. It is not a complete description of what is involved in the procedure – the healthcare team should describe it in detail.

NICE has looked at whether laparoscopic live donor simple nephrectomy is safe enough and works well enough for it to be used routinely when a person is donating a kidney.

To produce this guidance, NICE has:

- looked at the results of studies on the safety of laparoscopic live donor simple nephrectomy and how well it works
- asked experts for their opinions
- asked the views of the organisations that speak for the healthcare professionals and the people who will be affected by this guidance.

This guidance is part of NICE's work on 'interventional procedures' (see 'Further information' on page 10).

## About laparoscopic live donor simple nephrectomy

When a kidney is used in a kidney transplant, it's more likely that the transplant will be successful if the kidney has come from a living person (called a kidney donor) than if it has come from a person who has died.

The standard operation for taking the kidney out involves making an opening near to the kidney. The surgeon then works through this opening to remove the kidney. At the end of the operation, the opening is stitched closed. This is known as an 'open' procedure.

The operation that NICE has looked at is a way of removing the kidney from the kidney donor using keyhole surgery. This means that the surgeon uses special narrow equipment and does the operation through small cuts made in the kidney donor's abdomen.

## How well it works

### What the studies said

The studies NICE looked at showed that there were no differences in how well the transplanted kidney worked between patients whose donors had had the keyhole procedure and those whose donors had had the open operation. It was not possible to see whether this would be the case over a long period of time, though, because the studies didn't give any information on this.

One of the studies showed that around one-third of transplanted kidneys were rejected by the patients' bodies in the month after the transplant. This was the same for patients whose donors had had the keyhole procedure and for patients whose donors had had the open operation.

In general, the donors who had the keyhole surgery stayed in hospital for a shorter time than the donors who had the open operation, and they were also able to go back to work sooner.

### What the experts said

The experts did not have any concerns about how well the new procedure worked.

## Risks and possible problems

### What the studies said

The risks of the new procedure seem to be similar to those of the open operation for both the donors and the transplant patients (the patients who received a donated kidney). Up to 1 in 5 donors who had the new procedure had problems during or after the operation. Some of the studies of the open procedure did not give any information about what happened to donors. But the information that was given showed that up to around 1 in 3 donors had problems during or after the open operation.

For the transplant patients, there didn't seem to be any differences between the problems when the donor had had the new procedure and when the donor had had the open operation. But again, some of the studies didn't say what happened to all the transplant patients. Where information was given on problems affecting the tube joining the kidney to the bladder, the numbers for comparison were:

- 3 to 10 transplant patients in 100 had this type of problem where the donor had had the new procedure
- 3 to 6 transplant patients in 100 had this type of problem where the donor had had the open operation.

## What the experts said

The experts said that the main problems with laparoscopic live donor simple nephrectomy were likely to be bleeding and damage to other parts of the body near to the kidney. In some cases, a problem with this procedure may mean that part way through, the surgeon decides that it would be best to change to the open operation.

## What has NICE decided?

NICE has considered the evidence on laparoscopic live donor simple nephrectomy. It has recommended that when doctors use it for people who are donating a kidney, they should be sure that:

- the kidney donor understands what is involved and agrees (consents) to the procedure, and
- the results of the procedure are monitored.



## What the decision means for you

As a kidney donor, you may be offered laparoscopic live donor simple nephrectomy. NICE has considered this procedure because it is relatively new. NICE has decided that the procedure is safe enough and works well enough for use in the NHS. Nonetheless, you should understand the benefits and risks of laparoscopic live donor simple nephrectomy before you agree to it. These should be discussed with you. Some of these benefits and risks may be described above.

## Further information

You have the right to be fully informed and to share in decision-making about the procedure you have. You may want to discuss this guidance with the doctors and nurses looking after you.

You can visit the NICE website ([www.nice.org.uk](http://www.nice.org.uk)) for further information about the National Institute for Clinical Excellence and the Interventional Procedures Programme. A copy of the full guidance on laparoscopic live donor simple nephrectomy is on the NICE website ([www.nice.org.uk/IPG057guidance](http://www.nice.org.uk/IPG057guidance)), or you can order a copy from the website or by telephoning the NHS Response Line on 0870 1555 455 and quoting reference number N0568. The evidence that NICE considered in developing this guidance is also available from the NICE website.

If you want more information on kidney disease, a good starting point is NHS Direct, telephone 0845 4647, or NHS Direct Online ([www.nhsdirect.nhs.uk](http://www.nhsdirect.nhs.uk)).

Date: May 2004



**National Institute for  
Clinical Excellence**

MidCity Place  
71 High Holborn  
London  
WC1V 6NA

[www.nice.org.uk](http://www.nice.org.uk)