

## **HTG302 – Irreversible electroporation for treating primary lung cancer and metastases in the lung**

#The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS code(s):**

**E59.8 Other specified other operations on lung**

**Y12.3 Electrochemotherapy to lesion of organ NOC**

**Y53.- Approach to organ under image control**

Or

**Y68.- Other approach to organ under image control**

**Note:** Codes within category **Y53.-** and **Y68.-** are used as secondary codes to classify interventions that are percutaneous and require some form of image control: if the method of image control is unspecified, **Y53.9 Unspecified approach to organ under image control** is assigned.

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**C34.- Malignant neoplasm of bronchus and lung**

Or

**C78.0 Secondary malignant neoplasm of lung**

---

## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).