

## HTG338 - Faecal microbiota transplant for recurrent Clostridium difficile infection

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS-4 code(s):**

FMT using a colonoscope:

**H21.8 Other specified other therapeutic endoscopic operations on colon**

**Y37.8 Other specified introduction of other substance into organ NOC**

FMT using a nasoduodenal tube:

**G57.8 Other specified other operations on duodenum**

**Y37.8 Other specified introduction of other substance into organ NOC**

FMT using a nasogastric tube:

**G48.8 Other specified other operations on stomach**

**Y37.8 Other specified introduction of other substance into organ NOC**

FMT via rectal enema:

**H62.8 Other specified other operations on bowel**

**Y37.8 Other specified introduction of other substance into organ NOC**

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**A04.7 Enterocolitis due to Clostridium difficile**

---

## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).