

HTG410 - Balloon pulmonary angioplasty for chronic thromboembolic pulmonary hypertension

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS- 4 codes:

L13.5 Percutaneous transluminal balloon angioplasty of pulmonary artery NEC

Y53.- Approach to organ under image control

Or

Y68.- Other approach to organ under image control

Or

Y78 Arteriotomy approach to organ under image control

Note: A code from category **Y53.- Approach to organ under image control** or **Y68.- Other approach to organ under image control** may also be assigned to these procedures if image control has been used.

Note: Codes in categories **Y53.-** or **Y68.-** are used as secondary codes. When a procedure has been performed using image control and the code that classifies the procedure **does not** state the type of image control used, then a code from these categories **must be** assigned. If the specific method of image control is not stated, the fourth-character **.9** must be assigned

Diagnosis or health condition:

ICD-10 code(s):

I27.2 Other secondary pulmonary hypertension

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).