

HTG452 - Extracranial to intracranial bypass for intracranial atherosclerosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

L29.6 High-flow interposition extracranial to intracranial bypass from external carotid artery to middle cerebral artery

Y46.- Open approach to contents of cranium

or

L29.7 Bypass of carotid artery by anastomosis of superficial temporal artery to middle cerebral artery

Y46.- Open approach to contents of cranium

or

L29.8 Other specified reconstruction of carotid artery

Y46.- Open approach to contents of cranium

Note: **L29.8** is used to capture all other extracranial to intracranial bypass procedures not included at **L29.6** or **L29.7**. Additional codes are required if a graft is utilised.

Diagnosis or health condition:

ICD-10 code(s):

I67.2 Cerebral atherosclerosis

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).