

## HTG558 - Pressurised intraperitoneal aerosol chemotherapy for peritoneal carcinomatosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS-4 code(s):**

**T48.2 Introduction of cytotoxic substance into peritoneal cavity**

**Y75.2 Laparoscopic approach to abdominal cavity NEC**

**X72–X73 Delivery of chemotherapy for neoplasm**

**Note:** An OPCS-4 code from the range **X72–X73** from the National Tariff Chemotherapy Regimens List 2017-18 is assigned, as per the instructions in National Clinical Coding Standards OPCS-4 reference book **PCSX27: Delivery of chemotherapy for neoplasm (X72–X73)**.

**Note:** Only delivery of chemotherapy codes from categories **X72** and **X73** are required for national reporting purposes. The collection of codes classifying procurement is not required.

**Note:** It is not possible to classify the 'pressurised aerosolised' aspect of this treatment within OPCS-4.

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**C78.6 Secondary malignant neoplasm of retroperitoneum and peritoneum**

---

## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).