

HTG569 - Cytoreduction surgery with hyperthermic intraoperative peritoneal chemotherapy for peritoneal carcinomatosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure and device:

OPCS-4 code(s):

Note: Research indicates that 'Cytoreduction surgery' removes the maximum of visible tumour from the peritoneal cavity and may be performed along with the excision/resection of other organs where necessary.

'Cytoreduction' of the peritoneum is not specifically classifiable in OPCS-4, but would be assigned one of the following 'excision' of peritoneum codes, depending on the method of approach used.

T41.6 Excision of peritoneum

Or

T42.5 Endoscopic excision of peritoneum

Note: Heated chemotherapy is not specifically classifiable within OPCS-4. It is coded using the following code combination:

T48.2 Introduction of cytotoxic substance into peritoneal cavity

Plus

X72.- Delivery of chemotherapy for neoplasm

Note: If other organs are excised/resected in addition to the peritoneal cytoreduction, these procedures will require coding in addition.

Diagnosis or health condition:

ICD-10 code(s):

C78.6 Secondary malignant neoplasm of retroperitoneum and peritoneum

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).