

HTG662 Percutaneous transluminal renal sympathetic denervation for resistant hypertension

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

Renal sympathetic denervation using radiofrequency:

L43.6 Percutaneous transluminal radiofrequency denervation of renal artery

Y53.4 Approach to organ under fluoroscopic control

Z94.- Laterality of operation

Renal sympathetic denervation using ultrasound:

L43.8 Other specified transluminal operations on renal artery

Y11.5 Ultrasonic destruction of organ NOC

Y53.4 Approach to organ under fluoroscopic control

Z94.- Laterality of operation

Diagnosis or health condition:

ICD-10 code(s):

I10.X Essential (primary) hypertension

I15.- Secondary hypertension

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS Digital to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS Digital Information Representation Services work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS Digital is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).