

## **HTG737 Image-guided transperineal laser ablation (TPLA) for treating lower urinary tract symptoms caused by benign prostatic hyperplasia**

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS-4 code(s):**

**M70.8 Other specified other operations on outlet of male bladder**  
**Y08.3 Laser destruction of organ NOC**  
**Y53.2 Approach to organ under ultrasonic control**  
**Z42.2 Prostate**

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**N40.X Hyperplasia of prostate**

The appropriate code(s) to classify the lower urinary tract symptoms

---

## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).