

Laparoscopic repair of abdominal aortic aneurysm

Interventional procedures guidance

Published: 22 August 2007

[nice.org.uk/guidance/ipg229](https://www.nice.org.uk/guidance/ipg229)

Your responsibility

This guidance represents the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, healthcare professionals are expected to take this guidance fully into account. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Commissioners and/or providers have a responsibility to implement the guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity, and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

1 Guidance

- 1.1 There is adequate evidence of the safety and efficacy of laparoscopic repair of abdominal aortic aneurysm, but the technical demands are such that this procedure should not be used without special arrangements for consent and for audit or research.

- 1.2 Clinicians wishing to undertake laparoscopic repair of abdominal aortic aneurysm should take the following actions.
- Inform the clinical governance leads in their Trusts.
 - Ensure that patients understand the procedure and its place in the elective treatment of abdominal aortic aneurysm. They should provide patients with clear, written information. In addition, the patient should be informed of other available techniques, and told that conversion to open surgery may be necessary. Use of the Institute's [information for patients](#) ('Understanding NICE guidance') is recommended.
- 1.3 Clinicians undertaking this procedure should submit data on all patients to the National Vascular Database held by [The Vascular Society](#).
- 1.4 Selection of patients should be performed by a multidisciplinary team experienced in the management of aortic aneurysms and able to offer alternative treatment options.
- 1.5 This procedure should be performed by vascular surgeons who have had training in advanced laparoscopic surgery, and are mentored in these techniques.

2 The procedure

2.1 *Indications*

- 2.1.1 Dilatation of the aorta forming an aneurysm occurs in about 2% of men over the age of 65 (it is less common in women). Small aneurysms may present no problems, but some continue to grow, and larger aneurysms can leak or rupture. This carries a high risk of mortality, even when it is possible to offer emergency surgery. Preventive treatment is often advised for patients with aneurysms that represent an appreciable rupture risk.
- 2.1.2 The traditional treatment for abdominal aortic aneurysm is open surgical repair. The aneurysm is opened and a graft is then sewn in above and below the weakened area to allow normal blood flow. A less invasive approach is now commonly used, involving endovascular stent graft placement via the femoral arteries, but not all aneurysms are suitable for endovascular treatment.

2.2 *Outline of the procedure*

2.2.1 Laparoscopic repair of abdominal aortic aneurysm can be done by hand-assisted laparoscopic surgery (HALS) or by the technically more demanding totally laparoscopic surgery (TLS). In HALS, a midline minilaparotomy incision is made for insertion of one of the surgeon's hands to aid the procedure. In both techniques, small skin incisions are made for insertion of a laparoscope and instruments to guide and/or perform the repair. Lumbar arteries and the inferior mesenteric artery are dissected and clipped. Clamps are applied above and below the aneurysm, the sac is opened and thrombus removed. A prosthetic vascular graft is anastomosed to the proximal and distal ends of the aorta. The aneurysm wall and the posterior parietal peritoneum may be closed to cover the graft.

2.3 *Efficacy*

2.3.1 In three non-randomised controlled trials, mean hospital length of stay (LOS) was 5.9 (HALS), 6.2 and 6.3 (TLS) days, compared with 9.4, 10.0 and 10.2 days following open repair. One non-randomised controlled study reported that mean LOS was broadly similar following HALS (7.4 days) and endovascular stenting (6.4 days).

2.3.2 In one case series, mean LOS was 5 days for TLS (n = 131) and 7 days for HALS (n = 215). In a second case series, mean LOS for HALS was reported as 4.4 days. There was a statistically significant difference in mean LOS between the first 30 patients (5.3 days) and the last 92 patients (4.1 days) treated at one institution (p = 0.001). For more details, refer to the 'Sources of evidence' section.

2.3.3 All Specialist Advisers considered this procedure to be novel and of uncertain efficacy. They considered key efficacy outcomes for this procedure to be successful complete repair, open conversion rates, operative time, intensive care unit and overall hospital LOS, patient quality of life, renal function and need for return to theatre. Some Specialist Advisers suggested that there would be longer operating times, particularly early in the learning curve.

2.4 *Safety*

- 2.4.1 Postoperative death rates following laparoscopic aneurysm repair have been reported as 3% (1/29) and 4% (1/24) (HALS), and 5% (3/60) and 10% (2/20) (TLS).
- 2.4.2 One non-randomised controlled trial reported that the rate of renal insufficiency was 2% (1/60) following laparoscopic repair compared with 11% (11/100) following open repair. Other complications reported following laparoscopic aneurysm repair include bleeding from the hypogastric artery in <1% (1/122; HALS) and bleeding requiring reoperation in 2% (1/60; TLS).
- 2.4.3 In three non-randomised controlled trials that compared laparoscopic aneurysm repair with open surgery, the mean operative time was longer in the laparoscopic (181 [HALS], 462 and 468 minutes [TLS]) than in the open surgery groups (136, 300 and 301 minutes, respectively; significance not reported). A fourth non-randomised controlled study comparing HALS with endovascular stenting reported that mean operative time was longer in the HALS group (198 and 149 minutes, respectively; not statistically significant).
- 2.4.4 In one case series mean operative time was 257 minutes with HALS; in another it was 175 minutes with HALS and 265 minutes with TLS. For more details, refer to the 'Sources of evidence' section.
- 2.4.5 Safety outcomes highlighted by the Specialist Advisers were death within 30 days and late mortality, and major complications such as blood loss, infection, and multiple organ failure. They all agreed that the safety of this novel procedure is uncertain and that advanced training in laparoscopic surgical techniques is important.

2.5 *Other comments*

- 2.5.1 It was noted that the different laparoscopic techniques have differing technical demands and training requirements. The Committee considered that total cross-clamp time and long-term graft performance are important outcome measures.

3 Further information

- 3.1 The Institute has issued guidance on [stent-graft placement in abdominal aortic aneurysm](#) and [endovascular stent-graft placement in thoracic aortic aneurysms and dissections](#).

Andrew Dillon
Chief Executive
August 2007

Information for patients

NICE has produced [information describing its guidance on this procedure for patients and their carers](#) ('Understanding NICE guidance'). It explains the nature of the procedure and the decision made, and has been written with patient consent in mind.

Sources of evidence

The evidence considered by the Interventional Procedures Advisory Committee is described in the following document.

['Interventional procedure overview of laparoscopic repair of abdominal aortic aneurysm'](#), January 2007.

4 About this guidance

NICE interventional procedure guidance makes recommendations on the safety and efficacy of the procedure. It does not cover whether or not the NHS should fund a procedure. Funding decisions are taken by local NHS bodies after considering the clinical effectiveness of the procedure and whether it represents value for money for the NHS. It is for healthcare professionals and people using the NHS in England, Wales, Scotland and Northern Ireland, and is endorsed by Healthcare Improvement Scotland for implementation by NHSScotland.

This guidance was developed using the NICE [interventional procedure guidance](#) process.

We have produced a [summary of this guidance for patients and carers](#). Information about the evidence it is based on is also [available](#).

Changes since publication

14 January 2012: minor maintenance.

Your responsibility

This guidance represents the views of NICE and was arrived at after careful consideration of the available evidence. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of healthcare professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have regard to promoting equality of opportunity. Nothing in this guidance should be interpreted in a way which would be inconsistent with compliance with those duties.

Copyright

© National Institute for Health and Clinical Excellence 2007. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

Contact NICE

National Institute for Health and Clinical Excellence
Level 1A, City Tower, Piccadilly Plaza, Manchester M1 4BT

www.nice.org.uk

nice@nice.org.uk

0845 033 7780

Endorsing organisation

This guidance has been endorsed by [Healthcare Improvement Scotland](#).