

# Ex-vivo hepatic resection and reimplantation for liver cancer

Interventional procedures guidance

Published: 22 April 2009

[nice.org.uk/guidance/ipg298](https://www.nice.org.uk/guidance/ipg298)

## Your responsibility

This guidance represents the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, healthcare professionals are expected to take this guidance fully into account. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Commissioners and/or providers have a responsibility to implement the guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity, and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

## 1 Guidance

- 1.1 Current evidence on ex-vivo hepatic resection and reimplantation for liver cancer raises concerns about the safety and efficacy of the procedure. Therefore this procedure should only be used with special arrangements for clinical governance, consent and audit or research. It should only be used for

patients who would otherwise not survive and for whom other treatment options have failed or are inappropriate.

1.2 Clinicians wishing to undertake ex-vivo hepatic resection and reimplantation for liver cancer should take the following actions.

- Inform the clinical governance leads in their Trusts.
- Ensure that patients understand the uncertainty about the procedure's safety and efficacy; specifically the risks of death or serious morbidity, and the possible need for liver transplantation. Clear written information should be provided. In addition, the use of NICE's [information for patients](#) ('Understanding NICE guidance') is recommended.
- Audit and review clinical outcomes of all patients having ex-vivo hepatic resection and reimplantation for liver cancer (see section 3.1).

## 2 The procedure

### 2.1 *Indications and current treatments*

2.1.1 This procedure can be carried out in patients with primary or secondary (metastatic) liver cancer.

2.1.2 Treatment strategies for patients with liver cancer depend on tumour type, location, number and size. Most patients with liver cancer cannot benefit from surgical treatment and are treated with palliative intent. For some patients, liver resection surgery, either on its own or in combination with other treatments, may be beneficial.

### 2.2 *Outline of the procedure*

2.2.1 The procedure is carried out with the patient under general anaesthesia. The liver is removed through an abdominal incision and is perfused with a preservative solution. A bloodless resection of the diseased hepatic parenchyma is then performed, allowing complex reconstruction of the hepatic and portal vein structures, and the liver is reimplanted into the patient. The procedure can be performed with or without venovenous bypass.

Sections 2.3 and 2.4 describe efficacy and safety outcomes which were available in the published literature and which the Committee considered as part of the evidence about this procedure. For more detailed information on the evidence, see the [overview](#).

## 2.3 *Efficacy*

2.3.1 In a case series of 24 patients (22 with cancer) treated by ex-vivo hepatic resection and reimplantation, it was possible to resect and reimplant the liver in 91% (20/22) of the patients with cancer. In the case series of 24 patients (22 with cancer), 59% (13/22) survived the procedure and were discharged. Of the 13 patients who survived the procedure, 77% (10/13) died of tumour recurrence at between 12-month and 36-month follow-up. In a case series of 8 patients with liver metastases from colorectal cancer, 4 patients were treated by ex-vivo resection, of whom 2 were alive at 5-month follow-up (1 patient with tumour recurrence in the bone). The third patient died after 30 months and the fourth died 15 days after the operation. In a case series of 16 patients treated by liver resection with hepatic vein reconstruction, 2 patients were treated by ex-vivo resection. One of the 2 patients, with hepatocellular carcinoma, was alive and disease free at 52-month follow-up. The other patient, with colorectal metastases, was free of hepatic involvement at death following small bowel perforation after 4 months of follow-up. A case report of a single patient with hepatocellular carcinoma reported that the patient was alive with no recurrence 1 year after ex-vivo hepatic resection.

2.3.2 In the case series of 24 patients (22 with cancer), the mean operative time was 13.54 hours, and the mean anhepatic phase was 6.67 hours.

2.3.3 The Specialist Advisers stated that the key efficacy outcome of this procedure is survival.

## 2.4 *Safety*

2.4.1 Of the 22 patients with cancer in the case series of 24 patients treated by ex-vivo liver resection, 41% (9/22) died postoperatively during the same admission episode as the operation (exact timing of death not stated). In the same study, 32% of patients (7/22) required donor liver transplantation either immediately

(2 patients) or at a subsequent procedure (5 patients; exact timing of transplantation not stated).

- 2.4.2 In the case series of 8 patients, of the 4 patients undergoing ex-vivo resection, 1 patient died after 15 days from respiratory failure, renal failure and haemopneumothorax; and 1 patient developed inferior vena caval obstruction requiring stenting and pleural effusion requiring drainage.
- 2.4.3 The Specialist Advisers stated that adverse events (reported in the literature or anecdotally) include mortality, liver failure and bleeding/requirement for blood transfusion. The Specialist Advisers also commented that the procedure may increase the demand for donor livers.

### 3 Further information

- 3.1 This guidance requires that clinicians undertaking the procedure make special arrangements for audit. NICE has identified relevant audit criteria and has developed [audit support](#) (which is for use at local discretion).
- 3.2 NICE has published interventional procedures guidance on [radiofrequency ablation for the treatment of colorectal metastases in the liver](#), [selective internal radiation therapy for colorectal metastases in the liver](#), [microwave ablation for the treatment of metastases in the liver](#), [radiofrequency-assisted liver resection](#), [living-donor transplantation](#), [laparoscopic liver resection](#) and [radiofrequency ablation of hepatocellular carcinoma](#).

### *Information for patients*

NICE has produced [information on this procedure for patients and carers](#) ('Understanding NICE guidance'). It explains the nature of the procedure and the guidance issued by NICE, and has been written with patient consent in mind.

### 4 About this guidance

NICE interventional procedure guidance makes recommendations on the safety and efficacy of the procedure. It does not cover whether or not the NHS should fund a procedure. Funding decisions are taken by local NHS bodies after considering the clinical effectiveness of the procedure and whether it represents value for money for the NHS. It is for healthcare professionals and people

using the NHS in England, Wales, Scotland and Northern Ireland, and is endorsed by Healthcare Improvement Scotland for implementation by NHSScotland.

This guidance was developed using the NICE [interventional procedure guidance](#) process.

We have produced a [summary of this guidance for patients and carers](#). Tools to help you put the guidance into practice and information about the evidence it is based on are also [available](#).

## Changes since publication

7 January 2012: minor maintenance.

## Your responsibility

This guidance represents the views of NICE and was arrived at after careful consideration of the available evidence. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of healthcare professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the guidance, in their local context, in light of their duties to avoid unlawful discrimination and to have regard to promoting equality of opportunity. Nothing in this guidance should be interpreted in a way which would be inconsistent with compliance with those duties.

## Copyright

© National Institute for Health and Clinical Excellence 2009. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

## Contact NICE

National Institute for Health and Clinical Excellence  
Level 1A, City Tower, Piccadilly Plaza, Manchester M1 4BT

[www.nice.org.uk](http://www.nice.org.uk)

[nice@nice.org.uk](mailto:nice@nice.org.uk)

0845 033 7780

## *Endorsing organisation*

This guidance has been endorsed by [Healthcare Improvement Scotland](#).

## *Accreditation*

