

IPG418 – Percutaneous transluminal radiofrequency sympathetic denervation of the renal artery for resistant hypertension

SNOMED CT provides clinical terms for entry into the patient record to record clinical information relevant to that encounter; the mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure and device:

SNOMED CT preferred term (concept ID)

OPCS-4 code(s)

L43.6 Percutaneous transluminal radiofrequency denervation of renal artery

Y53.4 Approach to organ under fluoroscopic control

Diagnosis or health condition:

SNOMED CT preferred term (concept ID)

ICD-10 code(s):

Clinical coding recommendations for NICE guidance

For each published interventional procedure and medical technologies guidance, we work with NHS Digital to provide relevant clinical coding information.

SNOMED CT provides clinical terms for entry into the patient record to store clinical information relevant to that encounter.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS Digital Information Representation Services work collaboratively to ensure the most appropriate SNOMED CT, OPCS-4 and ICD codes are provided.

NHS Digital is the national release centre for the UK edition of SNOMED CT, ICD and OPCS-4. Further SNOMED CT information including licensing, see [here](#). Further Classifications information, see [here](#).