

Percutaneous interlaminar endoscopic lumbar discectomy for sciatica

Interventional procedures guidance

Published: 27 April 2016

[nice.org.uk/guidance/ipg555](https://www.nice.org.uk/guidance/ipg555)

Your responsibility

This guidance represents the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, healthcare professionals are expected to take this guidance fully into account. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

Commissioners and/or providers have a responsibility to implement the guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity, and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

1 Recommendations

- 1.1 Current evidence on the safety and efficacy of percutaneous interlaminar endoscopic lumbar discectomy for sciatica is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit.

- 1.2 Percutaneous interlaminar endoscopic lumbar discectomy for sciatica is a procedure that needs particular experience. Surgeons should acquire the necessary expertise through specific training and mentoring. It should only be done by surgeons who do the procedure regularly.
- 1.3 Details about all patients having percutaneous interlaminar endoscopic lumbar discectomy for sciatica should be entered onto the [British Spine Registry](#).

2 Indications and current treatments

- 2.1 Lumbar disc herniation occurs when the nucleus pulposus of an intervertebral disc protrudes through a tear in the surrounding annulus fibrosus. Symptoms include pain in the back or leg, and numbness or weakness in the leg. Serious neurological sequelae including painful foot drop, bladder dysfunction, and cauda equina syndrome, may sometimes occur.
- 2.2 Conservative treatments include analgesics, non-steroidal anti-inflammatory medication, manual therapy and acupuncture. Epidural corticosteroid injections can also be used to reduce nerve pain in the short term. Lumbar discectomy is considered if there is severe nerve compression or persistent symptoms that are unresponsive to conservative treatment. Surgical techniques include open discectomy, microdiscectomy or minimally invasive alternatives using percutaneous endoscopic approaches. The choice of operative technique may be influenced by several factors, including the presenting symptoms and signs, and the location and size of the disc prolapse.

3 The procedure

- 3.1 Percutaneous endoscopic lumbar discectomy aims to preserve bony structures and cause less damage to paravertebral muscles and ligaments than open discectomy, allowing a shorter hospital stay and faster recovery. An interlaminar approach provides an alternative to the transforaminal approach for treating central or centro-lateral disc extrusions, especially at the L5–S1 level where the transforaminal approach is difficult.
- 3.2 Percutaneous interlaminar endoscopic lumbar discectomy is usually done with the patient in the prone position using local or general anaesthesia. Under fluoroscopic guidance, a guidewire is inserted into the appropriate interlaminar

space. Dilators are used to expose the ligamentum flavum and the ruptured disc is accessed through this ligament. An endoscope and rongeurs are used to remove the herniated disc fragments. A laser may also be used to aid removal of the disc. The patient can usually mobilise within a few hours of the procedure.

4 Efficacy

This section describes efficacy outcomes from the published literature that the Committee considered as part of the evidence about this procedure. For more detailed information on the evidence, see the [interventional procedure overview](#).

- 4.1 A retrospective study of 60 patients comparing interlaminar endoscopic lumbar discectomy (n=30) against transforaminal endoscopic lumbar discectomy (n=30) reported a significant improvement in mean visual analogue scale (VAS) scores (ranging from 0 to 10 from best to worst), in both groups, for leg and back pain from before the procedure to a mean follow-up of 2.2 years. In the interlaminar group, back pain scores changed from 5.5 to 2.4 and leg pain scores changed from 7.6 to 1.7 (level of significance not reported). In the transforaminal group, back pain scores changed from 5.2 to 2.4 and leg pain scores changed from 7.4 to 1.6 (level of significance not reported). There was no significant difference between the interlaminar and transforaminal groups.
- 4.2 A case series of 400 patients treated by interlaminar endoscopic lumbar discectomy reported an improvement in mean VAS scores for back and leg pain from 7.9 before the procedure to 1.5 at 3 months after the procedure; it also reported that the VAS scores improved significantly in 90% of patients when compared against scores before the procedure.
- 4.3 The retrospective comparative study of 60 patients reported significant improvements in mean Oswestry Disability Index (ODI) scores (ranging from 0 to 100, from no disability to maximum disability) from before the procedure to a mean follow-up of 2.2 years; from 51% to 15% in the interlaminar group, and from 52% to 12% in the transforaminal group (no significant difference between groups). A case series of 372 patients treated by percutaneous interlaminar endoscopic discectomy reported improvement in mean ODI score from 79% before the procedure to 21% at 2 years after the procedure (level of significance not stated). The same study reported mean North American Spine Society

neurology scores (ranging from 1 to 6, from best to worst) of 3 before the procedure and 2 at 2 years (level of significance not stated).

- 4.4 A prospective comparative study of 200 patients with disc herniation treated by full-endoscopic discectomy (interlaminar approach, n=59; transforaminal approach, n=41) or microsurgical discectomy (n=100) reported recurrence rates at 2-year follow-up of 6% (3/53) in the interlaminar group, 8% (3/38) in the transforaminal group and 6% (5/87) in the microsurgical group (no significant difference between groups). All patients with recurrence were treated a second time by the same technique; in the transforaminal group, 2 patients had another recurrence.
- 4.5 A prospective comparative study of 100 patients with recurrent lumbar disc herniation treated by full-endoscopic discectomy (interlaminar approach, n=29; transforaminal approach, n=21) or microsurgical discectomy (n=50) reported re-recurrence rates at 2-year follow-up of 4% (1/24) in the interlaminar group, 10% (2/21) in the transforaminal group and 5% (2/42) in the microsurgical group (no significant difference between groups). All patients with re-recurrence were treated a second time by the same technique.
- 4.6 The retrospective comparative study of 60 patients reported recurrence in 7% (2/30) of patients treated by the interlaminar approach and in 3% (1/30) of patients treated by the transforaminal approach within a minimum of 2 years after the procedure (no significant difference between groups). The case series of 400 patients reported recurrence in 2 patients; they were treated again by surgery at 3 and 6 months after the first procedure.
- 4.7 The case series of 400 patients reported conversion to open surgery in 1 patient who had root protrusion after sustaining a dural tear during the procedure; the authors stated that this happened during the period when the surgeons were gaining experience in how to do the procedure. A case series of 163 patients (175 procedures) with lumbar disc herniations treated by interlaminar (n=104) or transforaminal (n=71) endoscopic lumbar discectomy reported no conversion to open surgery for either approach.
- 4.8 The retrospective comparative study of 60 patients reported complete removal of the disc fragment in 93% (28/30) of patients treated by the interlaminar

approach and in 97% (29/30) of patients treated by the transforaminal approach (no significant difference between groups).

- 4.9 The retrospective study of 60 patients reported that the mean time to return to work was 4.4 weeks for patients treated by the interlaminar approach and 4.9 weeks for patients treated by the transforaminal approach (no significant difference between groups). The case series of 372 patients reported that 98% (247/251) of patients who were not unemployed or retired returned to their occupation or sport activities; 2% (4/251) were not able to return to their occupation because of persistent paresis. Sick leave following hospitalisation ranged from 5 to 33 days (mean of 16 days).
- 4.10 The case series of 400 patients reported good-to-excellent results according to MacNab criteria in 91% (364/400) of patients; poor results were reported in 2% (8/400) of patients (no further details reported). The case series of 372 patients reported that 91% (301/331) of patients reported subjective satisfaction up to 2 years after the procedure and would have the procedure again; 9% (29/331) had a poor result (defined as no reduction in leg pain or having to be retreated by open surgery).
- 4.11 Specialists advisers listed key efficacy outcomes as resolution of leg pain, improvement in disability score, recurrence rate (reoccurrence of leg pain following an initial resolution of the leg pain), improvement in a generic quality of life measure (such as EQ-5D), return to activity, reduced operating time and hospital length of stay.

5 Safety

This section describes safety outcomes from the published literature that the Committee considered as part of the evidence about this procedure. For more detailed information on the evidence, see the [interventional procedure overview](#).

- 5.1 Single-facet injury during the procedure was reported in the first 3 patients in a case series of 400 patients with lumbar disc herniation treated by percutaneous interlaminar endoscopic lumbar discectomy (no further details provided).
- 5.2 Dural injury was reported in 1 patient who had recurrent lumbar disc herniation after conventional discectomy, treated by interlaminar endoscopic lumbar

discectomy in a prospective comparative study of 100 patients treated by full-endoscopic discectomy (interlaminar approach, n=29; transforaminal approach, n=21) or microsurgical discectomy (n=50); it was repaired with fibrin glue. Dural injury was reported in none of the patients in the transforaminal group and in 6% (3/50) of patients in the microsurgical group (no further details provided). Minor dural tear was reported in 2% (7/400) of patients in the case series of 400 patients (no further details provided). Dural tear was reported in 6% (6/104) of procedures using the interlaminar approach in a case series of 163 patients (175 procedures) with lumbar disc herniations treated by interlaminar or transforaminal endoscopic lumbar discectomy. In 5 procedures, patients were treated conservatively with 2 additional days of bed rest before mobilisation and discharge. In 1 procedure, an attempt was made to repair the dura by open surgery immediately after the procedure; this was complicated by an open cerebrospinal fluid fistula. The patient needed a second procedure to repair the dura and 5 days of bed rest and lumbar drainage.

- 5.3 Nerve root injury and persistent paraesthesia 2 years after the procedure were reported in 1 patient in the case series of 400 patients (no further details provided).
- 5.4 Motor deficit was reported in 3% (5/163) of patients (interlaminar approach, n=104 procedures; transforaminal approach, n=71 procedures) in the case series of 163 patients. In 2 of these 5 patients, 2-level discectomy was performed using an interlaminar approach for 1 level and a transforaminal approach for 1 level. In 4 patients these motor deficits were transient and they recovered completely, including the 2 patients who were treated by 2-level discectomies. In 1 patient there was a permanent motor deficit resulting in footdrop (no further details provided).
- 5.5 Transient dysaesthesia was reported in 3% (2/59) of patients with symptomatic lumbar disc herniation treated by interlaminar endoscopic lumbar discectomy in a prospective comparative study of 200 patients treated by full-endoscopic discectomy (interlaminar approach, n=59; transforaminal approach, n=41) or microsurgical discectomy (n=100). In the transforaminal group and in the microsurgical group, transient dysaesthesia was reported in 2% (1/41) and 5% (5/100) respectively (no further details provided). Transient dysaesthesia was reported in 6% (2/29) of patients who had recurrent lumbar disc herniation after conventional discectomy, treated by interlaminar endoscopic lumbar

discectomy, in the prospective comparative study of 100 patients treated by full-endoscopic discectomy or microsurgical discectomy; it was reported in none of the patients in the transforaminal group and in 10% (5/50) in the microsurgical group (no further details provided). Dysaesthesia was reported in 7% (2/30) of patients with symptomatic lumbar disc herniation treated by interlaminar endoscopic lumbar discectomy and in none of the 30 patients treated by the transforaminal approach in a retrospective comparative study of 60 patients (no further details provided). Transient dysaesthesia was reported in 3 patients treated by interlaminar endoscopic lumbar discectomy in a case series of 372 patients with symptomatic lumbar disc herniation (no further details provided).

- 5.6 Discitis was reported in 1% (2/400) of patients after the procedure in the case series of 400 patients; both patients were treated conservatively (no further details provided).
- 5.7 Pseudocysts were reported in 3% (9/298) of procedures in the group of patients treated by interlaminar endoscopic lumbar discectomy and in 1% (6/1205) of procedures in the group of patients treated by the transforaminal approach, in a case series of 1,406 patients with protruded or extruded disc materials compressing the lumbar root ($p=0.001$ for the comparison between groups). The interval between discectomy and pseudocyst detection on MRI was a mean of 53.7 (11–118) days. Five pseudocysts were treated surgically and 10 were treated conservatively.
- 5.8 In addition to safety outcomes reported in the literature, specialist advisers are asked about anecdotal adverse events (events which they have heard about) and about theoretical adverse events (events which they think might possibly occur, even if they have never done so). For this procedure, the specialist advisers did not report any anecdotal adverse events. They considered that the following were theoretical adverse events: bleeding, haematoma and scar tissue.

6 Further information

- 6.1 For related NICE guidance, see the [NICE website](#).

Information for patients

NICE has produced information on this procedure for patients and carers ([information for the public](#)). It explains the nature of the procedure and the guidance issued by NICE, and has been written with patient consent in mind.

ISBN: 978-1-4731-1812-6

Endorsing organisation

This guidance has been endorsed by [Healthcare Improvement Scotland](#).

Accreditation

