

# Artificial anal sphincter implantation

**Understanding NICE guidance –  
information for people considering the  
procedure, and for the public**

June 2004



Corporate member of  
Plain English Campaign.  
Committed to clearer communication.

**197**

## **Artificial anal sphincter implantation**

### **Understanding NICE guidance – information for people considering the procedure, and for the public**

**Issue date:** June 2004

#### **To order copies**

Copies of this booklet can be ordered from the NHS Response Line; telephone 0870 1555 455 and quote reference number N0598. A version in Welsh and English is also available, reference number N0599. Mae fersiwn yn Gymraeg ac yn Saesneg ar gael hefyd, rhif cyfeirnod N0599. The NICE interventional procedures guidance on which this information is based is available from the NICE website ([www.nice.org.uk](http://www.nice.org.uk)). Copies can also be obtained from the NHS Response Line, reference number N0597.

---

## **National Institute for Clinical Excellence**

MidCity Place  
71 High Holborn  
London  
WC1V 6NA

Website: [www.nice.org.uk](http://www.nice.org.uk)

ISBN: 1-84257-661-5

Published by the National Institute for Clinical Excellence  
June 2004

Typeset by Icon Design, Eton  
Print on Demand

© National Institute for Clinical Excellence, June 2004. All rights reserved. This material may be freely reproduced for educational and not-for-profit purposes within the NHS. No reproduction by or for commercial organisations is allowed without the express written permission of the National Institute for Clinical Excellence.

## Contents

<b>About this information</b>	<b>4</b>
<b>About artificial anal sphincter implantation</b>	<b>5</b>
<b>What has NICE decided?</b>	<b>8</b>
<b>What the decision means for you</b>	<b>9</b>
<b>Further information</b>	<b>10</b>

## About this information

This information describes the guidance that the National Institute for Clinical Excellence (NICE) has issued to the NHS on a procedure called artificial anal sphincter implantation. It is not a complete description of what is involved in the procedure – the patient’s healthcare team should describe it in detail.

NICE has looked at whether artificial anal sphincter implantation is safe enough and works well enough for it to be used routinely for the treatment of severe faecal incontinence.

To produce this guidance, NICE has:

- looked at the results of studies on the safety of artificial anal sphincter implantation and how well it works
- asked experts for their opinions
- asked the views of the organisations that speak for the healthcare professionals and the patients and carers who will be affected by this guidance.

This guidance is part of NICE’s work on ‘interventional procedures’ (see ‘Further information’ on page 10).

## About artificial anal sphincter implantation

Faecal incontinence is the medical term for what happens when a person passes faeces (stools) without the normal amount of control. It can happen as a result of problems with the anal sphincter, which is the ring of muscle that keeps the anus closed. Such problems may be due to sphincter damage, spinal injury or other neurological disorders.

Artificial anal sphincter implantation involves placing a circular cuff under the skin around the anus. The cuff is filled with fluid and sits tightly around the anus, keeping it closed. A tube runs under the skin from the cuff to a control pump. In a man, the pump is in his scrotum (the area of the testicles). In a woman, it's put into the area near to her vagina. A special balloon is placed into the patient's abdomen, and this is connected to the control pump by tubing that runs under the skin.

When the person is in the toilet, they can activate the pump, which takes the fluid out of the cuff and sends it up to the balloon. The cuff opens up and the person can pass faeces. Once finished, the fluid slowly returns to the cuff, which tightens up again.

## How well it works

### What the studies said

There were no good-quality studies that compared what happened in patients who had an artificial anal sphincter with what happened in similar patients who didn't have the artificial sphincter. In the studies that were found, some patients needed to have the artificial sphincter taken out. In one study, 10 out of 53 patients needed to have the sphincter removed (19% of patients). In another study, 7 out of 17 patients had to have it taken out (41% of patients). In the remaining patients, having the sphincter improved their incontinence, though the way this was measured was different in the different studies. In studies that looked at the pressure in the anus, the pressure was higher after the artificial sphincter had been implanted (which is what would be expected if the artificial sphincter was working).

### What the experts said

The experts thought that having to remove the sphincter was likely to be the main problem.

## Risks and possible problems

### What the studies said

The biggest study NICE found, which looked at what happened in 115 patients, showed that most patients had problems with the artificial sphincter (99 out of 115 patients had a problem). The most common ones are listed below, together with the numbers of patients who had the problem (out of the group of 115):

- infection (38 patients)
- pain (37 patients)
- erosion of the device, usually the cuff (24 patients)
- faeces becoming wedged behind the sphincter (21 patients)
- faecal incontinence (21 patients)
- constipation (20 patients)
- injury during the procedure (15 patients)
- problems with the openings made for the cuff, pump, balloon or tubing (11 patients)
- difficulty passing faeces (10 patients)
- the site of the cuff, pump, balloon or tubing opening up again (10 patients).

## What the experts said

The experts thought that the main problems would be infection, device erosion and problems passing faeces

## What has NICE decided?

NICE has decided that, if a doctor wants to carry out artificial anal sphincter implantation, he or she should make sure that the patient understands what is involved and that there are still uncertainties over the safety of the procedure and how well it works. There should be special arrangements in place so that the patient only agrees (consents) to the procedure after this discussion has taken place. NICE may look at the operation again if new information becomes available.

NICE has also recommended that the procedure should be carried out only in units that specialise in treating patients with faecal incontinence.

## Other comments from NICE

The procedure may be useful in the treatment of patients if sacral nerve stimulation is unsuitable (this is a way of treating incontinence by sending an electrical message to the nerves that control the muscles involved).



Also, as with all procedures, it's important that patients understand what is involved and what could go wrong. Problems are quite common following this procedure, and patients may need to have more operations as a result.

## **What the decision means for you**

Your doctor may have offered you artificial anal sphincter implantation at a specialist unit. NICE has considered this procedure because it is relatively new. NICE has decided that there are uncertainties about the benefits and risks of artificial anal sphincter implantation which you need to understand before you agree to it. Your doctor should discuss the benefits and risks with you. Some of these benefits and risks may be described above.

## Further information

You have the right to be fully informed and to share in decision-making about the treatment you receive. You may want to discuss this guidance with the doctors and nurses looking after you.

You can visit the NICE website ([www.nice.org.uk](http://www.nice.org.uk)) for further information about the National Institute for Clinical Excellence and the Interventional Procedures Programme. A copy of the full guidance on artificial anal sphincter implantation is on the NICE website ([www.nice.org.uk/IPG066guidance](http://www.nice.org.uk/IPG066guidance)), or you can order a copy from the website or by telephoning the NHS Response Line on 0870 1555 455 and quoting reference number N0597. The evidence that NICE considered in developing this guidance is also available from the NICE website.

If you want more information on incontinence, a good starting point would be NHS Direct (telephone 0845 4647) or NHS Direct Online ([www.nhsdirect.nhs.uk](http://www.nhsdirect.nhs.uk)).

Date: June 2004





*National Institute for  
Clinical Excellence*

**National Institute for  
Clinical Excellence**

MidCity Place  
71 High Holborn  
London  
WC1V 6NA

[www.nice.org.uk](http://www.nice.org.uk)