

Interventional procedure overview of VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Contents

The condition, current treatments, unmet need and procedure	2
Outcome measures.....	3
Cerebral performance category (CPC).....	3
Modified Rankin scale (mRS).....	3
Modified Glasgow Outcome Scale (MGOS)	3
Evidence summary	4
Population and studies description.....	4
Procedure technique	37
Efficacy.....	37
Safety	49
Validity and generalisability	52
Ongoing trials	54
Existing assessments of this procedure	54
Related NICE guidance	55
Interventional procedures	55
Professional societies	55
Company engagement.....	56
References.....	56
Appendix A: Methods and literature search strategy.....	58
Appendix B: Other relevant studies.....	83

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Table 1 Abbreviations

Abbreviation	Definition
ACS	Acute coronary syndrome
CA	Cardiac arrest
CCPR	Conventional cardiopulmonary resuscitation
CI	Confidence interval
CPC	Cerebral performance category
CPR	Cardiopulmonary resuscitation
CV	Cardiovascular
ECMO	Extracorporeal membrane oxygenation
ECPR	Extracorporeal cardiopulmonary resuscitation
IHCA	In-hospital cardiac arrest
ITT	Intention to treat
MGOS	Modified Glasgow Outcome Scale
mRS	Modified Rankin scale
OHCA	Out-of-hospital cardiac arrest
OR	Odds ratio
PEA	Pulseless electrical activity
PSM	Propensity-score matched
RCT	Randomised controlled trial
ROSC	Return of spontaneous circulation
RR	Relative risk
VA	Venoarterial
VF	Ventricular fibrillation
VT	Ventricular tachycardia

The condition, current treatments, unmet need and procedure

Information about the procedure, condition, current practice and unmet need is available in [NICE's interventional procedures guidance on VA ECMO for extracorporeal cardiopulmonary resuscitation \(ECPR\) in adults with refractory cardiac arrest](#).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Outcome measures

The main outcomes include survival and survival with favourable neurological outcome. The measures used are detailed in the following paragraphs.

Cerebral performance category (CPC)

The Glasgow-Pittsburgh cerebral performance category (CPC) assessment tool is a 5-category measure used to assess neurological outcome. Categories 1 (good cerebral performance: conscious, alert, capable of normal life) and 2 (moderate cerebral disability: conscious, alert, sufficient cerebral function for activities of daily life) are considered to show a good neurological outcome. Categories 3 (severe cerebral disability), 4 (coma/vegetative state) and 5 (certified brain death) are considered a poor neurological outcome.

Modified Rankin scale (mRS)

The modified Rankin scale (mRS) assessment tool grades functional recovery on a 7-category scale from 0 (no symptoms), to 6 (dead). A score of 0 to 3 is considered a good neurological outcome, and a score 4 or 5 as a poor neurological outcome.

Modified Glasgow Outcome Scale (MGOS)

The modified Glasgow Outcome Scale (MGOS) assessment tool grades functional recovery on a 5-category scale from 1 (dead), to 5 (good recovery). A score of 4 or 5 is considered a good neurological outcome, and a score 1 to 3 as a poor neurological outcome.

Evidence summary

Population and studies description

This interventional procedure overview is focused on VA ECMO use in ECPR for in hospital cardiac arrest (IHCA) and out of hospital cardiac arrest (OHCA). Two additional overviews have been developed focusing on VA ECMO in acute heart failure and postcardiotomy cardiogenic shock.

This overview is based on more than 300,000 people from 9 systematic reviews (Low 2023, Low 2024, Zhong 2024, Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024; Reddy 2025), 1 long-term RCT follow-up study (Rob 2024), 2 retrospective registry studies (Balucani 2024; Inoue 2022), and 1 single-centre retrospective PSM study (Shih 2024). This a rapid review of the literature, and a flow chart of the complete selection process is shown in [figure 1](#). This overview presents 13 studies as the key evidence in [table 2](#) and [table 3](#), and lists 60 other relevant studies in [appendix B, table 5](#).

All 9 systematic reviews (Low 2023, Low 2024, Zhong 2024, Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024; Reddy 2025), included the same 3 RCTs; ARREST (USA, 2020), Prague OHCA (Czech Republic, 2022), and INCEPTION (Netherlands, 2023). Of these systematic reviews, 3 included a fourth pilot RCT: EROCA (USA 2021) (Cheema 2023, Kiyohara 2023, Scquizzato 2023).

OHCA and IHCA studies

The selected key evidence includes 4 publications reporting on 3 systematic reviews and meta-analyses comparing ECPR and CCPR in people with IHCA and people with OHCA from studies done in Asia, Europe and North America (Low 2023, Low 2024, Zhong 2024, Reddy 2025). All other key evidence included people with OHCA only.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

The systematic review and meta-analyses by Low (2023), included 11 studies (3 RCTs and 8 PSMs) reporting on adults with OHCA or IHCA. This was updated in the publication by Low (2024) to include a further 2 PSM studies. Overall, both Low et al. publications report on 14,048 adults, of whom 6,336 had ECPR. Of the studies included in the review, 8 (3 RCTs and 5 PSMs) included people with OHCA, 4 PSM studies included people with IHCA, and 1 PSM included people with OHCA and IHCA. All RCTs were noted to be either at 'low risk' or 'some concerns' for bias, while all PSMs were noted to be of 'high quality'. Across the included studies, the mean age ranged from 55 to 73 years and 77% were male. The most common cause of cardiac arrest was acute coronary syndrome (33 to 77%), and of those who had ECPR, 57% had an initial shockable rhythm compared with 60% in the CCPR group.

The systematic review and meta-analyses by Zhong (2024) included 17 studies (3 RCTs, 5 prospective studies, and 9 retrospective studies) reporting on 167,728 people with OHCA or IHCA. Of the 17 included primary studies, 11 were also included in the Low systematic reviews, including the same 3 RCTs. Of the studies included by Zhong (2024), 10 included people with OHCA, 4 included people with IHCA, and 3 included people with OHCA and IHCA. Using the Cochrane Collaboration's risk assessment tool, no RCT study was considered 'high risk of bias' in each domain. Using the Newcastle Ottawa Scale, most of the 10 cohort studies were considered medium quality. Across the included studies, the mean age ranged from 50 to 75 years. People who had ECPR were considerably younger and more likely to have an initial shockable rhythm compared with those in the CCPR group.

The systematic review and meta-analyses by Reddy (2025) included 28 studies (3 RCTs and 25 observational studies) accounting for 304,360 people with refractory cardiac arrest. Using Cochrane Collaboration risk assessment tools for nonrandomised studies (ROBINS-I) and revised tool for assessing risk of bias in IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

randomised trials (ROB-2), in general, the risk of bias was high because of missing data as most of the studies were observational in nature and utilised retrospective chart review. Of the 28 included primary studies, 24 included people with OHCA and 4 included people with IHCA. The proportion of people with OHCA was 90% in the ECPR group and 92% in the CCPR group. The mean age was 59 years in people having ECPR and 66 years in people having CCPR. The proportion of males was 78% in the ECPR group and 67% in the CCPR group. People in the ECPR group had a higher incidence of shockable rhythm than those who had CCPR. Outcomes were reported up to 12 months.

All 3 systematic reviews reported on short-term (30 days from CA) and long-term (90 or more days from CA) outcomes, as well as survival up to 1 year.

OHCA studies

The systematic review and meta-analyses by Gomes (2023) included 3 RCTs comparing ECPR and CCPR in 418 people with OHCA. In the risk of bias assessment of the 3 RCTs, 2 were considered to have 'some concerns' due to crossover between the study groups. However, this potential bias was minimised by balanced deviations from intended interventions between study groups and by blinded outcomes assessment. The mean age across the ECPR groups ranged from 54 to 59 years compared with 57 to 58 years in the CCPR groups and the proportion of males was 82% to 93% and 73% to 89%, respectively. The primary cause of cardiac arrest was acute myocardial infarction in 50% of people in the Prague OHCA trial, and 77% in the INCEPTION trial. In 2 of the included RCTs (ARREST and INCEPTION) most people enrolled had an initial shockable rhythm (98 to 100%), whereas the Prague OHCA trial recruited a high proportion with non-shockable rhythm (mean 61% both arms), a subset known to have worse outcomes.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

The systematic reviews and meta-analyses by Cheema (2023), Kiyohara (2023) and Scquizzato (2023) all included the same 3 RCTs and an additional pilot trial (EROCA). The number of people included in each systematic review ranged from 433 to 435 depending on whether they included the intention to treat, per protocol or as-treated analysis set. The mean age across the ECPR groups ranged from 54 to 62 years, and 57 to 61 years in the CCPR group. The proportion of males ranged from 67 to 93% in the ECPR groups and 67 to 89% in the CCPR groups. Each systematic review reported slightly different meta-analysis results as they each used different statistical methods, pooled outcome endpoints differently, and did different subgroup analyses. All 3 systematic reviews conducted risk of bias assessments using the Cochrane Risk of Bias (RoB2) tool. Kiyohara (2023) considered all studies to be low risk of bias, Cheema (2023) considered 3 studies to have a low risk of bias and 1 (EROCA) to be high risk of bias. Scquizzato (2023) considered all studies to be intermediate risk of bias due to the absence of blinding of the treating team in all studies and unblinded assessors of neurological outcome in one study. All 3 systematic reviews reported on outcomes at hospital discharge or 30 days, and at 3 or 6 months.

The systematic review and meta-analyses by Pagura (2024) included 18 studies (3 RCTs and 15 observational studies [6 with PSM design]) reporting on 21,877 people with refractory OHCA. The mean age across the ECPR groups ranged from 46 to 66 years, and 54 to 77 years in the CCPR group. The proportion of males ranged from 70 to 93% in the ECPR groups and 62 to 89% in the CCPR groups. The mean proportion of people in the ECPR groups with initial shockable rhythm was 61% and 60% in the CCPR group. Outcomes were reported up to 6 months.

In the long-term follow-up study of the Prague OHCA RCT, 255 people who survived following the initial trial period were followed for a median of 5.3 years

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

(3.8 to 7.2) after initial cardiac arrest. The median age of people at recruitment was 58 years and 82% were male. More people in the CCPR arm had an initial shockable rhythm (VF) 64%, compared with the ECPR arm 58%. The authors noted that there was no formal power analysis for the long-term follow-up outcomes (Rob 2024).

The retrospective SAVE-J II multicentre registry study by Inoue (2022) included 1,644 adults with OHCA who had ECPR in Japan. The median age was 60 years and 85% of the population were male. Initial shockable rhythm was reported in 69% of people and 59% had a primary cause of CA of acute coronary syndrome. Outcomes were reported until hospital discharge.

The single centre retrospective PSM study by Shih (2024) reported on 1,193 people with OHCA resuscitated with ECPR or CCPR in a high volume emergency department in Taiwan. The PSM cohort included 231 people, of whom 77 had ECPR. The median age was 57 years, 81% were male and the proportion of people with an initial shockable rhythm was 86% and 88% in the ECPR and CCPR arms, respectively. Outcomes were reported until hospital discharge.

The retrospective analysis of data from the ELSO international registry by Balucani (2024) included 7,460 people who had ECPR for refractory cardiac arrest. The mean age was 53.4 years and 70% of the population were men. Pre-existing ischaemic heart disease was less frequent in women than in men (15% versus 19%; $p < 0.001$). Outcomes were reported until hospital discharge.

[Table 2](#) presents study details.


Figure 1 Flow chart of study selection

Table 2 Study details

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
1	Low, 2023 Low, 2024 Europe, Asia, North America	<p><u>Low, 2023</u> n=9,192 (4,595 ECPR) Mean age (years):</p> <ul style="list-style-type: none"> • ECPR: 59.0 • CCPR: 59.3 <p>Males:</p> <ul style="list-style-type: none"> • ECPR: 75.3% • CCPR: 78.3% <p><u>CA cause:</u></p> <ul style="list-style-type: none"> • Acute coronary syndrome (32.7% to 76.8%) <p><u>Shockable rhythm</u></p> <ul style="list-style-type: none"> • ECPR: 56.8% • CCPR: 59.5% <p><u>Low, 2024</u> n=14,048 (6,336 ECPR) Mean age ranged from 55 to 73 years Males: not reported</p>	<p><u>Low, 2023</u> Systematic review and meta-analysis of 11 studies: 3 RCTs and 8 PSMs Search date: April 2023</p> <p><u>Low, 2024</u> Systematic review and meta-analysis of 13 studies with 14 pairwise comparisons: 3 RCTs and 10 PSMs Search date: November 2023</p>	<p><u>Low, 2023</u> RCTs and PSMs comparing ECPR with CCPR in adults (aged 18 years and over) with OHCA and IHCA.</p> <p><u>CA location:</u></p> <ul style="list-style-type: none"> • OHCA: 6 studies (3 RCTs and 3 PSMs) • IHCA: (4 PSMs) • OHCA and IHCA: (1 PSM) <p><u>Low, 2024</u> <u>CA location:</u></p> <ul style="list-style-type: none"> • OHCA: 8 studies (3 	<ul style="list-style-type: none"> • Intervention: ECPR • Comparator: CCPR <p><u>Mean low flow time</u></p> <ul style="list-style-type: none"> • ECPR: 48 min • CCPR: 44 min 	<p><u>Low, 2024</u> In-hospital, 30-day post-discharge, 3mo, 6mo, 1-year</p> <p><u>Low, 2024</u> In-hospital 30-day post-discharge</p>

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
				RCTs and 5 PSMs) <ul style="list-style-type: none"> IHCA: (4 PSMs) OHCA and IHCA: (1 PSM) 		
2	Zhong, 2024 Czech Republic, Belgium, France, Germany, Japan, Korea, Netherlands, Taiwan, USA	n=167,728 (2,308 ECPR) Mean age (years, range): <ul style="list-style-type: none"> ECPR: 50 to 72 CCPR: 57 to 75 Males: not reported <u>Shockable rhythm</u> <ul style="list-style-type: none"> ECPR: 1.9% to 100% CCPR: 2.9% to 100% 	Systematic review and meta-analysis of 17 studies: 3 RCTs, 5 prospective studies, 9 retrospective studies. Search date: July 2023	Studies including people over 16 years with IHCA or OHCA comparing CCPR and ECPR. <u>CA location:</u> <ul style="list-style-type: none"> OHCA: 10 studies IHCA: 4 studies OHCA and IHCA: 3 studies 	<ul style="list-style-type: none"> Intervention: ECPR Comparator: CCPR 	
3	Gomes, 2023 Czech Republic, Netherlands, USA	n=418 (208 ECPR) <u>Mean age (years):</u>	Systematic review and meta-analysis of 3 RCTs (ARREST, Prague OHCA, INCEPTION)	RCTs which compared ECPR with standard CPR for OHCA.	<ul style="list-style-type: none"> Intervention: ECPR Comparator: standard CPR <u>Time from arrest to ECPR</u>	In-hospital, 6 months

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
		<ul style="list-style-type: none"> ECPR: Range 54 to 59 CPR: Range 57 to 58 <u>Male (%)</u> <ul style="list-style-type: none"> ECPR: Range 82 to 93 CPR: Range 73 to 89 <u>CA cause:</u> <ul style="list-style-type: none"> AMI (50% Prague OHCA, 77% INCEPTION) <u>Shockable rhythm</u> <ul style="list-style-type: none"> ECPR: 58 to 100% CCPR: 64 to 100% 	<p>Search date: March 2023</p> <p>All RCTs were open label.</p>		<ul style="list-style-type: none"> ARREST: 59 min Prague OHCA: 61 min INCEPTION: 74 min 	
4	Cheema, 2023 Czech Republic, Netherlands, USA	<p>n=434 (220 ECPR)</p> <p><u>Mean age:</u></p> <ul style="list-style-type: none"> Not reported <p><u>Male (%):</u></p> <ul style="list-style-type: none"> Not reported 	<p>Systematic review and meta-analysis of 4 RCTs (ARREST, Prague OHCA, INCEPTION, EROCA)</p> <p>Search date: March 2023</p>	RCTs comparing ECPR with conventional CPR in people with OHCA.	<ul style="list-style-type: none"> Intervention: ECPR Comparator: CCPR 	In-hospital or 30 days 3 or 6 months

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
			All RCTs were open label.			
5	Kiyohara, 2023 Czech Republic, Netherlands, USA	n=435 (221 ECPR) <u>Mean age (years):</u> <ul style="list-style-type: none"> ECPR: Range 54 to 62 CCPR: Range 57 to 61 <u>Male (%):</u> <ul style="list-style-type: none"> ECPR: Range 67 to 93 CPR: Range 67 to 89 	Systematic review and meta-analysis of 4 RCTs (ARREST, Prague OHCA, INCEPTION, EROCA) Search date: February 2023 All RCTs were open label.	RCTs comparing the clinical outcomes of ECPR and CCPR for people with OHCA	<ul style="list-style-type: none"> Intervention: ECPR Comparator: CCPR <u>Time from arrest to ECPR</u> <ul style="list-style-type: none"> ARREST: 59 min Prague OHCA: 61 min INCEPTION: 74 min EROCA: 66 min 	In-hospital or 30 days 6 months
6	Scquizzato, 2023 Czech Republic, Netherlands, USA	n=433 (220 ECPR) <u>Mean age:</u> <ul style="list-style-type: none"> Not reported <u>Male:</u> <ul style="list-style-type: none"> Not reported 	Systematic review and meta-analysis of 4 RCTs (ARREST, Prague OHCA, INCEPTION, EROCA) Search date: February 2023	RCTs enrolling adults with refractory OHCA randomised to have ECPR or CCPR alone	<ul style="list-style-type: none"> Intervention: ECPR Comparator: CCPR 	In-hospital or 30 days Longest follow-up available

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
			All RCTs were open label.			
7	Pagura, 2024 Asia, Europe, US	<p>n=21,877 (3,129 ECPR)</p> <p><u>Mean age (years):</u></p> <ul style="list-style-type: none"> • ECPR: Range 46 to 66 • CCPR: Range 54 to 77 <p><u>Male (%):</u></p> <ul style="list-style-type: none"> • ECPR: Range 70 to 93% • CCPR: Range 62 to 89% <p><u>CA cause ACS:</u></p> <ul style="list-style-type: none"> • ECPR: Range 21 to 85% • CCPR: Range 4 to 89% <p><u>Shockable rhythm</u></p> <ul style="list-style-type: none"> • ECPR: Range 0 to 100% (mean 61%) 	<p>Systematic review and meta-analysis of 18 studies: 3 RCTs (ARREST, Prague OHCA, INCEPTION) and 15 observational studies (6 with PSM design).</p> <p>Search date: April 2023</p>	Observational and RCTs, comparing the effect of ECPR and CCPR in refractory OHCA.	<ul style="list-style-type: none"> • Intervention: ECPR • Comparator: CCPR 	In-hospital or 30 days 6 months

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
		<ul style="list-style-type: none"> CCPR: Range 0 to 100% (mean 60%) 				
8	Rob, 2024 Czech Republic	<p>n=255 (123 ECPR) <u>Median age at randomisation (years):</u></p> <ul style="list-style-type: none"> ECPR: 59 (48 to 66) CCPR: 57 (47 to 65) <p><u>Male (%):</u></p> <ul style="list-style-type: none"> ECPR: 82% CCPR: 83% <p><u>Shockable rhythm (VF)</u></p> <ul style="list-style-type: none"> ECPR: 58% CCPR: 64% 	<p>Long-term follow-up of the Prague OHCA RCT.</p> <p>The original RCT was a single-centre, prospective, open label RCT.</p> <p>Randomisation was done on-scene (pre-hospital) by web-based system.</p> <p>Functional assessments during follow-up were done by evaluators who were blinded to group allocation.</p>	<p>Adults aged 18 to 65 years with witnessed OHCA of presumed cardiac aetiology, who had received a minimum of 5 minutes of advanced cardiac life support without ROSC.</p> <p>People with unwitnessed, non-cardiac cause CA, and had suspected or confirmed pregnancy, had ROSC within 5 minutes, had obvious life-limiting comorbidities, bleeding diathesis, DNR order or</p>	<ul style="list-style-type: none"> Intervention: ECPR <p>Comparator: CCPR</p> <p>20/256 people (7.8%) were crossed over (11 crossovers from the CCPR to the ECPR group and 9 from the ECPR group to the CCPR group)</p>	Median 5.3 years (IQR 3.8 to 7.2) from CA.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
				prearrest CPC ≥ 3 were excluded.		
9	Inoue, 2022 Japan	n=1,644 <u>Median age (years):</u> 60 (18 to 93) <u>Male (%):</u> 84.6% <u>Initial shockable rhythm:</u> 69.4% <u>CA cause ACS:</u> 59%	Retrospective SAVE-J II multicentre registry study Search date: 2013 to 2018	Adults with OHCA who had ECPR.	<ul style="list-style-type: none"> ECPR 	In-hospital
10	Shih, 2024 Taiwan	n=1,193 (85 ECPR) PSM cohort=231 (77 ECPR) <u>Median age (years):</u> <ul style="list-style-type: none"> ECPR: 57 (47 to 65) CCPR: 56 (44 to 66) <u>Male (%):</u> <ul style="list-style-type: none"> ECPR: 81.8% CCPR: 80.5% <u>Initial shockable rhythm:</u>	Single centre retrospective PSM study (Taiwan) Search date: 2016 to 2021	Adults over 20 years old with refractory OHCA resuscitated in the emergency department (ED). Excluded people with non-cardiac causes for arrest, and those who achieved sustained ROSC within 15 mins at the ED.	<ul style="list-style-type: none"> Intervention: ECPR Comparator: CCPR Estimated low flow time <100 min.	In-hospital or 30 days

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
		<ul style="list-style-type: none"> ECPR: 85.7% CCPR: 87.7% 				
11	Reddy S, 2025 Norway, France, Germany, Czech Republic, Mexico, Taiwan, Taipei, Korea, Japan, the Netherlands, Belgium, China, US.	<p>n=304,360 (9,656 ECPR)</p> <p>28 studies</p> <p>Mean age:</p> <ul style="list-style-type: none"> ECPR=59 years CCPR=66 years <p>% male:</p> <ul style="list-style-type: none"> ECPR=78% CCPR=67% <p>OHCA:</p> <ul style="list-style-type: none"> ECPR=90% CCPR=92% <p>People in the ECPR group had a higher incidence of shockable rhythm than those who had CCPR.</p>	<p>Systematic review and meta-analysis</p> <p>Search date: July 2024</p> <p>3 randomised controlled trials and 25 observational studies were included.</p>	<p>Randomised controlled trials and observational studies were eligible for inclusion. Studies that compared ECPR with CCPR after refractory cardiac arrest in adults (18 years or older) were eligible. Outcomes of interest included survival to hospital discharge, survival at 30 days, survival with favourable neurological outcomes at 30 days and 6 months, and major adverse events.</p>	<ul style="list-style-type: none"> ECPR=9,656 CCPR=294,704 Mean CPR duration=49 minutes 	Up to 12 months

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Study no.	First author, date country	Characteristics of people in the study (as reported by the study)	Study design	Inclusion criteria	Intervention	Follow up
12	Balucani C, 2024 International	<p>n=7,460</p> <p>Mean age=53.4 years (50.4 for women versus 54.7 for men, $p<0.001$)</p> <p>Men=69.5%</p> <p>Race or ethnicity:</p> <ul style="list-style-type: none"> • White=55.9% • Asian=23.7% • Black=8.9% • Hispanic=3.8% <p>Pre-existing ischaemic heart disease was less frequent in women than in men (14.6% versus 18.5%; $p<0.001$).</p>	<p>Retrospective analysis of data from the ELSO Registry.</p> <p>1992 to 2020</p>	<p>Inclusion criteria were adults aged over 18 years, having VA ECMO for refractory cardiac arrest, identified in the ELSO registry from 1992 to 2020.</p> <p>People who had multiple runs of ECPR were excluded to avoid complexity and bias in the analysis resulting from severe illness, as it was considered that the morbidity during the inter-ECMO periods could introduce significant confounders.</p>	<p>ECPR</p> <ul style="list-style-type: none"> • The mean duration of ECMO support was statistically significantly shorter in women than men (89.2 versus 101.3 hours, $p<0.001$). 	To hospital discharge

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Table 3 Study outcomes

First author, date	Efficacy outcomes	Safety outcomes
Low, 2023 Low, 2024	<p>The analysis in Low (2024) provides updated meta-analysis estimates for mortality, 30-day survival, and short term survival with neurologically favourable outcome.</p> <p>In-hospital mortality (Overall population) <u>Low, 2023</u> Meta-analysis 3 RCTs, 8 PSMs (n=9,192)</p> <ul style="list-style-type: none"> • ECPR: 75.2% (3,454 of 4,595) • CCPR: 80.7% (3,708 of 4,597) <p>OR 0.67, 95% CI: 0.51 to 0.87, $I^2=42\%$; $p=0.0034$</p> <p>Sensitivity analysis excluded one study that contributed substantial weight, due to its large sample size (n=7,652) OR 0.60, 95% CI 0.41 to 0.89; $p=0.010$</p> <p><u>Low, 2024</u> Meta-analysis 3 RCTs, 11 PSMs (n=14,048)</p> <ul style="list-style-type: none"> • ECPR: 76.7% (4,859 of 6,336) • CCPR: 83.1% (6,411 of 7,712) <p>OR 0.63, 95% CI: 0.50 to 0.79, $I^2=64\%$</p> <p>In-hospital mortality (OHCA) <u>Low, 2023</u> Meta-analysis 3 RCTs, 3 PSMs (n=8,662)</p> <p>OR 0.76, 95% CI: 0.54 to 1.07, $I^2=54\%$; $p=0.12$</p>	<p>Bleeding (Overall) <u>Low, 2023</u> Meta-analysis 4 studies OR 4.84, 95% CI: 1.91 to 12.24; $p=0.0009$</p>

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>Sensitivity analysis excluding one study that contributed substantial weight, due to its large sample size (n=7,652) did not significantly change the pooled estimates (OR 0.71, 95% CI: 0.39 to 1.27; p=0.24).</p> <p><u>Low, 2024</u></p> <p>Meta-analysis 3 RCTs, 6 PSMs (n=13,518) OR 0.67, 95% CI: 0.51 to 0.88, I²=74%</p> <p>In-hospital mortality (IHCA)</p> <p><u>Low, 2023</u></p> <p>Meta-analysis 4 PSMs (n=370) OR 0.42, 95% CI: 0.25 to 0.70, I²=0%; p=0.0009</p> <p>Short-term survival with favourable neurological outcome (CPC 1 or 2; 30 days after CA) (Overall population)</p> <p><u>Low, 2023</u></p> <p>Meta-analysis 7 studies OR 1.65, 95% CI: 1.02 to 2.68; p=0.042</p> <p><u>Low, 2024</u></p> <p>Meta-analysis 2 RCTs, 8 PSMs</p> <ul style="list-style-type: none"> • ECPR: 12.8% (306 of 2,391) • CCPR: 8.8% (331 of 3,767) <p>OR 1.57, 95% CI: 1.14 to 2.15, I²=56%</p> <p>Short-term survival with favourable neurological outcome (CPC 1 or 2; 30 days after CA) (OHCA)</p>	

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p><u>Low, 2023</u> Meta-analysis 3 studies OR 1.24, 95% CI: 0.65 to 2.36; p=0.51 Short-term survival with favourable neurological outcome (CPC 1 or 2; 30 days after CA) (IHCA)</p> <p><u>Low, 2023</u> Meta-analysis 4 studies OR 2.37, 95% CI: 1.34 to 4.19; p=0.0031 Long-term survival with favourable neurological outcome (CPC 1 or 2; 90 days or more after CA) (Overall population)</p> <p><u>Low, 2023</u> Meta-analysis 8 studies OR 2.04, 95% CI: 1.41 to 2.94; p=0.0001 Long-term survival with favourable neurological outcome (CPC 1 or 2; 90 days or more after CA) (OHCA)</p> <p><u>Low, 2023</u> Meta-analysis 4 studies OR 1.96, 95% CI: 1.02 to 3.79; p=0.045 Long-term survival with favourable neurological outcome (CPC 1 or 2; 90 days or more after CA) (IHCA)</p> <p><u>Low, 2023</u></p>	

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>Meta-analysis 3 studies OR 2.80, 95% CI: 1.31 to 6.00; p=0.008</p> <p>30-day survival <u>Low, 2023</u> Meta-analysis 7 studies OR 1.45, 95% CI: 1.08 to 1.96; p=0.015 <u>Low, 2024</u> Meta-analysis 1 RCTs, 8 PSMs</p> <ul style="list-style-type: none"> • ECPR: 24.2% (1,377 of 5,689) • CCPR: 18.1% (1,030 of 5,697) <p>OR 1.70, 95% CI: 1.29 to 2.26, I²=71%</p> <p>3-month survival <u>Low, 2023</u> Meta-analysis 3 studies OR 3.98, 95% CI: 1.12 to 14.16; p=0.033</p> <p>6-month survival <u>Low, 2023</u> Meta-analysis 6 studies OR 1.87, 95% CI: 1.36 to 2.57; p=0.0001</p> <p>1-year survival <u>Low, 2023</u></p>	

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	Meta-analysis 5 studies OR 1.72, 95% CI: 1.52 to 1.95; $p < 0.0001$	
Zhong 2024	<p>Short-term favourable neurological status (Overall population) Meta-analysis 11 studies (ECPR $n=1,442$, CCPR $n=51,221$) RR 2.88; 95% CI: 1.96 to 4.23, $I^2=76\%$; $p < 0.0001$ Subgroup analysis with matched data including 3 RCTs and 7 PMSs (RR 1.67, 95% CI: 1.16 to 2.40, $I^2=51\%$; $p=0.005$)</p> <p>Short-term favourable neurological status (OHCA) Meta-analysis 7 studies RR 1.50, 95% CI: 0.98 to 2.29, $I^2=55\%$</p> <p>Short-term favourable neurological status (IHCA) Meta-analysis 3 studies RR 2.18, 95% CI: 1.24 to 3.81, $I^2=9\%$</p> <p>Long-term favourable neurological status (Overall population) Meta-analysis 11 studies (ECPR $n=896$, CCPR $n=1,977$) RR 2.11, 95% CI 1.40 to 3.19, $I^2=69\%$; $p=0.0004$ Subgroup analysis with matched data including 3 RCTs and 6 PMSs (RR 1.83, 95% CI: 1.32 to 2.53, $I^2=14\%$; $p=0.0003$)</p>	No safety outcomes reported.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>Long-term favourable neurological status (OHCA) Meta-analysis 5 studies RR 1.95, 95% CI: 1.01 to 3.59, $I^2=11\%$</p> <p>Long-term favourable neurological status (IHCA) Meta-analysis 3 studies RR 2.17, 95% CI: 1.19 to 3.94, $I^2=0\%$</p> <p>Overall survival (Overall population) Meta-analysis matched data 3 RCTs, 9 PSMs RR 1.51, 95% CI: 1.20 to 1.89, $I^2=62\%$; $p=0.0004$</p> <p>Survival at discharge (OHCA and IHCA) Meta-analysis matched data RR 1.25, 95% CI 1.00 to 1.56, $I^2=57\%$; $p=0.05$,</p> <p>3- to 6- month survival (OHCA and IHCA) RR 2.73, 95% CI: 1.67 to 4.48, $I^2=0\%$; $p<0.0001$</p> <p>1 year survival (OHCA and IHCA) RR 1.92, 95% CI: 1.14 to 3.25, $I^2=0\%$; $p=0.01$</p> <p>Short-term survival (discharge or 1 month; OHCA) Meta-analysis 9 studies RR: 1.10, 95% CI: 0.91 to 1.34, $I^2=44\%$</p> <p>Short-term survival (discharge or 1 month; IHCA) Meta-analysis 3 studies RR: 2.03, 95% CI: 1.30 to 3.18, $I^2=0\%$</p> <p>Long-term survival (3- to 6- month or 1 year; OHCA)</p>	

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	Meta-analysis 3 studies RR: 3.16, 95% CI: 1.36 to 7.38, $I^2=0\%$ Long-term survival (3- to 6- month or 1 year; IHCA) Meta-analysis 3 studies RR: 1.92, 95% CI: 1.14 to 3.25, $I^2=0\%$	
Gomes, 2023	In-hospital mortality <ul style="list-style-type: none"> • ECPR: 71.1% • CPR: 78.6% RR 0.89, 95% CI: 0.74 to 1.07, $I^2=53\%$; $p=0.23$ Survival with favourable neurological status (shortest follow-up [in-hospital or 30 days]) <ul style="list-style-type: none"> • ECPR: 26.4% • CPR: 17.2% RR 1.47, 95% CI: 0.91 to 2.40, $I^2=23\%$; $p=0.12$ Subgroup analysis of people with shockable rhythms at presentation: RR 1.62, 95% CI 0.95 to 2.76; $p=0.07$ Survival with favourable neurological status (6 months) <ul style="list-style-type: none"> • ECPR: 28.3% • CPR: 18.6% RR 1.48, 95% CI: 0.88 to 2.49, $I^2=28\%$; $p=0.14$ <ul style="list-style-type: none"> • Subgroup analysis of people with shockable rhythms at presentation: RR 1.50, 95% CI 0.90 to 2.50; $p=0.12$ 	No safety outcomes reported.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<ul style="list-style-type: none"> Sensitivity analysis excluding ARREST trial (most significant result): RR 1.39, 95% CI 0.97 to 1.99; p=0.07). 	
Cheema, 2023	<p>Mid-term survival (in-hospital or 30 days) Meta-analysis 4 studies RR: 1.21, 95% CI: 0.64 to 2.28, I²=48%; p=0.55</p> <p>Long-term survival (3 or 6 months) Meta-analysis 3 studies RR 1.32, 95% CI: 0.18 to 9.5, I²=64%; p=0.79</p> <p>Mid-term favourable neurological outcome (in-hospital or 30 days) Meta-analysis 4 studies RR: 1.59, 95% CI: 1.09 to 2.33, I²=0%; p=0.02</p> <p>Long-term favourable neurological outcome (3 or 6 months) Meta-analysis 4 studies RR: 1.47, 95% CI: 0.89 to 2.43, I²=25%; p=0.13</p>	<p>Adverse events Meta-analysis 2 studies RR: 3.22, 95% CI: 1.18 to 8.80, I²=63%; p=0.02</p>
Kiyohara, 2023	<p>Short-term survival with favourable neurological outcome (in-hospital or 30 days) Meta-analysis 4 studies OR: 1.84, 95% CI: 1.14 to 2.99, I²=0%; p=0.01</p>	No safety outcomes reported.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>6 month survival Meta-analysis 4 studies OR: 1.50, 95% CI: 0.67 to 3.36, $I^2=50\%$; $p=0.33$</p> <p>6 month survival with favourable neurological outcome (Overall) Meta-analysis 4 studies OR: 1.74, 95% CI: 0.86 to 3.51, $I^2=35\%$; $p=0.12$</p> <p>6 month survival with favourable neurological outcome (shockable rhythm) Meta-analysis 4 studies OR: 1.91, 95% CI: 0.90 to 4.03, $I^2=36\%$; $p=0.09$</p> <p>6 month survival with favourable neurological outcome (non-shockable rhythm) Meta-analysis 4 studies (2 studies had no people with no shockable rhythm) OR: 3.92, 95% CI: 0.42 to 36.35, $I^2=NA$; $p=0.23$</p>	
Scquizzato, 2023	<p>Survival with good neurological outcome (3 or 6 months) Meta-analysis 4 studies</p>	<p>Survival with unfavourable neurological outcome</p> <ul style="list-style-type: none"> • ECPR: 0% (0/220) • CPR: 1.9% (4/214) <p>OR 0.24, 95% CI: 0.05 to 1.26, $I^2=0\%$; $p=0.780$</p>

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

	<ul style="list-style-type: none"> • ECPR: 27% (59/220) • CPR: 18% (39/213) <p>OR 1.72, 95% CI: 1.09 to 2.70, $I^2=26\%$; $p=0.02$</p> <p>Survival with good neurological outcome (3 or 6 months) (shockable rhythm)</p> <p>Meta-analysis 4 studies</p> <ul style="list-style-type: none"> • ECPR: 34% (55/164) • CPR: 23% (38/165) <p>OR 1.90, 95% CI: 1.16 to 3.13, $I^2=23\%$; $p=0.011$</p> <p>Survival with good neurological outcome (hospital discharge or 30 days)</p> <p>Meta-analysis 4 studies</p> <ul style="list-style-type: none"> • ECPR: 25% (55/220) • CPR: 16% (34/212) <p>OR 1.82, 95% CI: 1.13 to 2.92, $I^2=0\%$; $p=0.013$</p> <p>Survival with good neurological outcome (hospital discharge or 30 days) (shockable rhythm)</p> <p>Meta-analysis 4 studies</p> <ul style="list-style-type: none"> • ECPR: 31% (51/164) • CPR: 21% (34/164) <p>OR 1.93, 95% CI: 1.16 to 3.23, $I^2=0\%$; $p=0.012$</p> <p>Survival (longest follow-up available)</p> <p>Meta-analysis 4 studies</p>	
--	---	--

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<ul style="list-style-type: none"> • ECPR: 28% (61/220) • CPR: 22% (47/214) <p>OR 1.31, 95% CI: 0.49 to 3.49, $I^2=58\%$; $p=0.592$</p> <p>Survival (hospital discharge or 30 days) Meta-analysis 4 studies</p> <ul style="list-style-type: none"> • ECPR: 33% (72/220) • CPR: 27% (58/214) <p>OR 1.35, 95% CI: 0.55 to 3.29, $I^2=0\%$; $p=0.514$</p>	
Pagura, 2024	<p>Survival with favourable neurological outcome (in-hospital or 30 days) Meta-analysis 14 studies</p> <ul style="list-style-type: none"> • ECPR: 14% (385/2,842) • CPR: 7% (1,339/18,188) <p>OR: 2.35, 95% CI: 1.61 to 3.43, $I^2=80\%$; $p<0.0001$</p> <p>Subgroup analysis including only RCTs OR: 1.83, 95% CI: 1.13 to 2.96, $I^2=0\%$; $p=0.01$</p> <p>Subgroup analysis including in-hospital follow-up only (7 studies) OR: 1.77, 95% CI: 1.15 to 2.73, $I^2=73\%$; $p=0.009$</p> <p>Subgroup analysis including 30 day follow-up only (8 studies) OR: 3.04, 95% CI: 1.59 to 5.80, $I^2=83\%$; $p<0.001$</p>	No safety outcomes reported.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>Survival with favourable neurological outcome (6 months) Meta-analysis 6 studies</p> <ul style="list-style-type: none"> • ECPR: 16% (118/725) • CPR: 8% (61/793) <p>OR: 2.72, 95% CI: 1.47 to 5.04, $I^2=47\%$; $p=0.002$</p> <p>Survival (in-hospital or 30 days) Meta-analysis 13 studies</p> <ul style="list-style-type: none"> • ECPR: 20% (430/2,156) • CPR: 10% (1,621/16,149) <p>OR 1.71, 95% CI: 1.18 to 2.46, $I^2=81\%$; $p=0.004$ Subgroup analysis including in-hospital follow-up only OR: 1.38, 95% CI: 0.95 to 2.02, $I^2=69\%$; $p=0.094$ Subgroup analysis including 30 day follow-up only OR: 2.26, 95% CI: 1.09 to 4.68, $I^2=86\%$; $p=0.029$</p>	

Rob, 2024	<p>Long-term survival (ITT analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 27.6% (34/123) • CCPR: 19.7% (26/132) <p>Log rank p=0.01</p> <p>Long-term survival (per protocol analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 29.8% (34/114) • CCPR: 18.2% (22/121) <p>Log rank p=0.008</p> <p>Long-term survival (as-treated analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 30.4% (38/125) • CCPR: 16.9% (22/130) <p>Log rank p<0.001</p> <p>Long-term favourable neurological outcome (ITT analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 26.8% (33/123) • CCPR: 18.9% (25/132) <p>RR 0.90, 95% CI: 0.79 to 1.03, p=0.13</p> <p>Similar results were observed for CPC and mRS categories.</p> <p>Long-term survival (per protocol analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 28.9% (33/114) 	<p>Long-term poor neurological outcome (ITT analysis) Median follow-up 5.3 years</p> <ul style="list-style-type: none"> • ECPR: 2.9% (1/34) • CCPR: 3.8% (1/26) <p>Death after discharge during follow-up (ITT analysis)</p> <ul style="list-style-type: none"> • ECPR: 10.3% (4/39) • CCPR: 20% (6/30) <p>RR 0.51, 95% CI: 0.16 to 1.66, p=0.26</p> <p>Hospitalisation after discharge during follow-up (ITT analysis)</p> <ul style="list-style-type: none"> • ECPR: 76.9% (30/39) • CCPR: 60% (18/30) <p>RR 1.28, 95% CI: 0.91 to 1.8, p=0.15</p> <p>CV hospitalisation after discharge during follow-up (ITT analysis)</p> <ul style="list-style-type: none"> • ECPR: 64.1% (25/39) • CCPR: 50% (15/30) <p>RR 1.28, 95% CI: 0.84 to 1.97, p=0.26</p> <p>Myocardial infarction (ITT analysis)</p> <ul style="list-style-type: none"> • ECPR: 2.6% (1/39) • CCPR: 3.3% (1/30) <p>p=0.91</p> <p>Stroke (ITT analysis)</p>
-----------	---	--

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<ul style="list-style-type: none"> • CCPR: 17.4% (21/121) RR 0.86, 95% CI: 0.75 to 0.99, p=0.035 Similar results were observed for CPC and mRS categories. Long-term survival (as-treated analysis) Median follow-up 5.3 years <ul style="list-style-type: none"> • ECPR: 29.6% (37/125) • CCPR: 16.2% (21/130) RR 0.84, 95% CI: 0.73 to 0.96, p=0.007 Similar results were observed for CPC and mRS categories. Quality of life (Mean EQ-VAS value) <ul style="list-style-type: none"> • ECPR: 71.0 • CCPR: 76.3 P=0.30	<ul style="list-style-type: none"> • ECPR: 0% (0/39) • CCPR: 3.3% (1/30) p=0.34 Heart failure hospitalisation (ITT analysis) <ul style="list-style-type: none"> • ECPR: 5.1% (2/39) • CCPR: 10% (3/30) p=0.57 Ventricular arrhythmia hospitalisation (ITT analysis) <ul style="list-style-type: none"> • ECPR: 2.6% (1/39) • CCPR: 10% (3/30) p=0.22

First author, date	Efficacy outcomes	Safety outcomes
Inoue, 2022	<p>Favourable neurological outcome at hospital discharge</p> <ul style="list-style-type: none"> Overall: 14.1% (231 of 1,644) Shockable rhythm: 16.7% PEA rhythm: 9.2% Asystole: 3.9% <p>Survival to hospital discharge</p> <ul style="list-style-type: none"> Overall: 27.2% (447 of 1,644) Shockable rhythm: 32% PEA rhythm: 18.5% Asystole: 10.8% 	<p>Complications during ECPR: 32.7% (535 of 1,644)</p> <p>Procedure-related complications: 21.2% (346 of 1,644)</p> <ul style="list-style-type: none"> Cannula malposition: 4.9% (81 of 1,644) Unsuccessful cannulation: 0.7% (11 of 1,644) Cannulation-related bleeding: 16.4% (268 of 1,644) Other: 1.6% (26 of 1,644) <p>ECMO-related complications: 3.1% (50 of 1,644)</p> <p>Haemorrhage: 8.5% (139 of 1,644)</p> <p>Ischaemia: 1.6% (26 of 1,644)</p>
Shih, 2024	<p>Survival with favourable neurological outcome (follow-up unclear)</p> <ul style="list-style-type: none"> ECPR: 18.2% (14/77) CCPR: 5.2% (8/154) <p>PSM multivariate analysis: aOR 13.31, 95% CI: 1.61 to 109.9, p=0.016</p> <p>Survival (in-hospital or 30-day)</p> <ul style="list-style-type: none"> ECPR: 28.6% (22/77) CCPR: 7.8% (12/154) <p>PSM multivariate analysis: aOR 6.02, 95% CI: 2.19 to 16.52</p>	No safety outcomes reported.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
Reddy S, 2025	<p>Survival at hospital discharge</p> <ul style="list-style-type: none"> • ECPR=20.1% (1,682/8,362) • CCPR=3.33% (7,954/238,631) <p>OR=0.48, 95% CI 0.27 to 0.84, p=0.01; 19 studies, I²=97%</p> <p>Survival at 30 days</p> <ul style="list-style-type: none"> • ECPR=26.3% (162/617) • CCPR=9.50% (1,582/16,656) <p>OR=0.36, 95% CI 0.25 to 0.51, p<0.0001; 7 studies, I²=50%</p> <p>Favourable neurological outcome at hospital discharge</p> <ul style="list-style-type: none"> • ECPR=11.8% (865/7,308) • CCPR=1.9% (4,612/236,599) <p>OR=0.41, 95% CI 0.17 to 1.01, p=0.05; 9 studies, I²=97%</p> <p>Favourable neurological outcome at 30 days</p> <ul style="list-style-type: none"> • ECPR=13.4% (131/979) • CCPR=10.9% (1,635/15,054) <p>OR=0.94, 95% CI 0.36 to 2.44, p=0.90; 6 studies, I²=94%</p> <p>Favourable neurological outcome within 3 months</p> <ul style="list-style-type: none"> • ECPR=12.1% (1,028/8,472) • CCPR=2.5% (6,355/252,693) <p>OR=0.46, 95% CI 0.20 to 1.05, p=0.06; 18 studies,</p>	<p>Bleeding</p> <ul style="list-style-type: none"> • ECPR=35.4% (152/430) • CCPR=3.78% (34/899) <p>OR=0.08, 95% CI 0.03 to 0.24, p<0.0001</p>

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	<p>$I^2=98\%$</p> <p>Favourable neurological outcome at 3 months</p> <ul style="list-style-type: none"> • ECPR=17.4% (38/218) • CCPR=10.3% (117/1,140) <p>OR=0.22, 95% CI 0.04 to 1.16, $p=0.07$; 4 studies, $I^2=86\%$</p> <p>Favourable neurological outcome at 6 to 12 months</p> <ul style="list-style-type: none"> • ECPR=19.6% (127/647) • CCPR=9.7% (89/913) <p>OR=0.41, 95% CI 0.23 to 0.71, $p=0.002$; 7 studies, $I^2=64\%$</p> <p>In a subgroup analysis of the 3 randomised controlled trials, people in ECPR group had better neurological outcomes compared with the CCPR group (OR 0.58, 95% CI 0.28 to 1.19, $p=0.14$, $I^2=36\%$, $n=418$) and better survival within 30 days (OR 0.61, 95% CI 0.34 to 1.08, $p=0.09$, $I^2=20\%$, $n=419$).</p>	
Balucani C, 2024	<p>Survival at discharge</p> <ul style="list-style-type: none"> • Overall=28.5% (2,123/7460) • Men=27.8% • Women=30.0%, $p=0.138$ <p>In the adjusted multivariable analysis, older age was the</p>	<p>Any neurological complication</p> <ul style="list-style-type: none"> • Overall=17.2% (1,286/7,460) • Men=17.9% (928/5,184) • Women=15.7% (358/2,276), $p=0.022$ <p>Any haemorrhagic complication</p> <ul style="list-style-type: none"> • Overall=26.6% (1,988/7,460) • Men=25.9% (1,345/5,184)

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

First author, date	Efficacy outcomes	Safety outcomes
	most important contributor of in-hospital mortality.	<ul style="list-style-type: none"> • Women=28.2% (643/2,276), p=0.038 <p>Any infectious complication</p> <ul style="list-style-type: none"> • Overall=6.84% (510/7,460) • Men=7.18% (372/5,184) • Women=6.1% (138/2,276), p=0.08 <p>Any limb complication</p> <ul style="list-style-type: none"> • Overall=6.8% (504/7,460) • Men=6.4% (333/5,184) • Women=7.51% (171/2,276), p=0.084 <p>Any mechanical complication</p> <ul style="list-style-type: none"> • Overall=14.1% (1,051/7,460) • Men=13.9% (724/5,184) • Women=14.4% (327/2,276), p=0.646 <p>Any metabolic complication</p> <ul style="list-style-type: none"> • Overall=19.7% (1,473/7,460) • Men=20.2% (1,050/5,184) • Women=18.4% (423/2,276), p=0.095 <p>Any cardiovascular complication</p> <ul style="list-style-type: none"> • Overall=37.0% (2,764/7,460) • Men=38.0% (1,972/5,184) • Women=34.8% (792/2,276), p=0.008 <p>Female sex was associated with decreased odds of neurological, cardiovascular, and renal complications.</p>

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Procedure technique

Of the 13 studies, none detailed the ECMO device or combination of devices used. Only a few of the studies detailed ECMO procedures. ECPR was initiated at the hospital (either catheterisation laboratory or emergency department) in most studies reporting on the use of ECPR for OHCA (Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024, Inoue 2022, Shih 2024). Six OHCA studies included in the systematic reviews by Low et al. also reported ECPR initiation in hospital, but one did not specify the location of ECPR initiation. One systematic review detailed that targeted temperature management after cardiac arrest was used in 45% of people having ECPR and 15% of people having CCPR (Low 2023).

Efficacy

Short-term survival with favourable neurological outcomes

Short-term survival with favourable neurological outcome was reported in 10 out of 13 studies included in the key evidence. Most often favourable neurological outcome was defined as CPC 1 or 2 on the cerebral performance category. One systematic review also included studies using the MGOS (Zhong 2024), and 2 systematic reviews included studies using the mRS (Scquizzato 2023, Pagura 2024). Mostly, short-term was defined as hospital discharge or 30 days after initial CA.

OHCA and IHCA studies

Four publications reporting on 3 systematic reviews presented meta-analyses for short-term survival with favourable neurological outcomes in both OHCA and IHCA (Low 2023, Low 2024, Zhong 2024, Reddy 2025). Both systematic reviews defined short-term as to hospital discharge or 30 days after initial CA.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

In the systematic review of 11 studies reporting on adults with OHCA or IHCA, ECPR was associated with improved short-term survival with favourable neurological outcomes compared with CCPR in a meta-analysis of 7 studies (OR 1.65, 95% CI: 1.02 to 2.68; $p=0.042$) (Low 2023). The updated analysis of 10 studies also found ECPR was associated with favourable neurological outcomes at short-term follow-up (OR 1.57, 95% CI 1.14 to 2.15) (Low 2024). A meta-analysis of 11 studies in people with OHCA and IHCA (including 1,442 people who had ECPR and 51,221 who had CCPR), showed improved short-term survival with favourable neurological outcomes with ECPR compared with CCPR (RR 2.88, 95% CI: 1.96 to 4.23, $I^2=76\%$; $p<0.0001$; Zhong 2024). This result was consistent in a subgroup analysis using RCTs and PSM data only (RR 1.67, 95% CI: 1.16 to 2.40, $I^2=51\%$; $p=0.005$; Zhong 2024). In the systematic review of 28 studies in people with OHCA and IHCA (including 9,656 people who had ECPR and 294,704 people who had CCPR), favourable neurological outcome at hospital discharge was 12% for ECPR and 2% for CCPR (OR 0.41, 95% CI 0.17 to 1.01, $p=0.05$; 9 studies; Reddy 2025). Favourable neurological outcome at 30 days was 13% for ECPR and 11% for CCPR (OR 0.94, 95% CI 0.36 to 2.44, $p=0.90$; 6 studies; Reddy 2025).

After stratifying based on the location of cardiac arrest, subgroup meta-analysis of 4 studies, showed that statistically significantly more people having ECPR for IHCA had short-term favourable neurological outcomes compared with people having CCPR for IHCA (OR 2.37, 95% CI: 1.34 to 4.19; $p=0.0031$, Low 2023). Similar outcomes were also seen in a subgroup meta-analysis of 10 studies (RR 2.18, 95% CI: 1.24 to 3.81, $I^2=9\%$; Zhong 2024). In a subgroup meta-analysis of 3 studies, no significant differences in the rate of short-term favourable neurological outcome were observed in people with OHCA who had ECPR or CCPR (OR 1.24, 95% CI: 0.65 to 2.36; $p=0.51$). Similar outcomes were also

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

seen in a subgroup meta-analysis of 10 studies (RR 1.50, 95% CI: 0.98 to 2.29, $I^2=55\%$; Zhong 2024).

OHCA studies

Five systematic reviews (Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024), 1 retrospective registry study (Inoue 2022), and 1 single-centre retrospective PSM study (Shih 2024) presented short-term survival with favourable neurological outcomes in people with OHCA. Short-term was defined as hospital discharge or 30 days after initial CA (although this was labelled 'mid-term' in the Cheema (2023) systematic review).

In the systematic review of 3 RCTs including 418 people with OHCA, ECPR was associated with a non-statistically significant higher rate of survival with a favourable neurological outcome at the shortest follow-up (26%) compared with standard CPR (17%), RR 1.47 (95% CI: 0.91 to 2.40; $p=0.12$; Gomes 2023). A subgroup analysis of people with an initial shockable rhythm showed similar findings to the main analysis, with a non-statistically significant benefit of ECPR (RR 1.62, 95% CI: 0.95 to 2.76; $p=0.07$; Gomes 2023).

Across 3 systematic reviews and meta-analyses including the same 4 RCTs reporting on people with OHCA, the rate of short-term survival with favourable neurological outcome was statistically higher in the ECPR groups compared with CCPR (Cheema 2023, Kiyohara 2023, Scquizzato 2023). Pooled survival with favourable neurological outcome was 25% for those who had ECPR compared with 16% who had CCPR (Scquizzato 2023). Using a Mantel-Haenszel, random-effects meta-analysis, the odds ratios reported were 1.82, 95% CI: 1.13 to 2.92; $p=0.01$, $I^2=0\%$ (Scquizzato 2023) and 1.84, 95% CI: 1.14 to 2.99, $I^2=0\%$; $p=0.01$ (Kiyohara 2023). The risk ratio (RR) using an inverse-variance, random effects meta-analysis was 1.59, 95% CI: 1.09 to 2.33, $I^2=0\%$; $p=0.02$ (Cheema 2023). A subgroup analysis of people with an initial shockable rhythm showed higher rate

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

of short-term survival with favourable neurological outcome in the ECPR (31%) groups compared with CCPR (21%), OR 1.93, 95% CI: 1.16 to 3.23, $I^2=0\%$; $p=0.012$; Scquizzato 2023).

In a meta-analysis of 14 studies (3 RCTs and 11 observational studies) in people with OHCA, the rate of short-term survival with favourable neurological outcome was significantly improved with ECPR (14%) compared with CCPR (7%); OR 2.35, 95% CI: 1.61 to 3.43, $I^2=80\%$; $p<0.0001$. This was consistent in subgroup analysis including only RCTs OR: 1.83, 95% CI: 1.13 to 2.96, $I^2=0\%$; $p=0.01$ (Pagura 2024).

In the retrospective SAVE-J II multicentre registry study which included 1,644 adults with OHCA who had ECPR in Japan, the overall rate of survival with favourable neurological outcome at hospital discharge was 14% (Inoue 2022). This rate was higher in those with an initial shockable rhythm (17%), and lower in those with a non-shockable rhythm (9% PEA, 4% asystole) (Inoue 2022). Multivariable analysis showed that younger age, initial shockable rhythm at the scene, and location of cardiac arrest were significantly associated with both favourable outcome and survival to hospital discharge ($p<0.01$) (Inoue 2022).

In the single centre retrospective PSM study reporting on people with OHCA resuscitated with ECPR or CCPR in a high volume emergency department in Taiwan, the rate of survival with favourable neurological outcome was 18% among those who had ECPR, compared with 5% in those who had CCPR. The PSM multivariate analysis reported an adjusted odds ratio (aOR) of 13.31, 95% CI: 1.61 to 109.9, $p=0.016$ (Shih 2024). Higher rates of favourable neurological outcome were associated with younger age (48 versus 59 years, $p=0.001$), CPR duration (37 min versus 51 min, $p=0.006$) and collapse to ECMO flow initiation time (76 versus 98.0 min, $p=0.031$).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Long-term survival with favourable neurological outcomes

Long-term survival with favourable neurological outcome was reported in 9 out of 13 studies included in the key evidence. Most often favourable neurological outcome was defined as CPC 1 or 2 on the cerebral performance category. One systematic review also included studies using the MGOS (Zhong 2024), and 2 systematic reviews included studies using the mRS (Scquizzato 2023, Pagura 2024). Mostly, long-term was defined as 3 months or longer after initial CA.

OHCA and IHCA studies

Two systematic reviews presented meta-analyses for long-term survival with favourable neurological outcomes in both OHCA and IHCA (Low 2023, Zhong 2024). Both systematic reviews defined long-term as 3 months or more after initial CA.

In the systematic review of 11 studies reporting on adults with OHCA or IHCA, ECPR was associated with statistically significantly improved long-term survival with favourable neurological outcomes compared with CCPR in a meta-analysis of 8 studies (OR 2.04, 95% CI: 1.41 to 2.94; $p=0.0001$) (Low 2023). A meta-analysis of 11 studies in people with OHCA and IHCA (including 896 people who had ECPR and 1,977 who had CCPR), showed improved long-term survival with favourable neurological outcomes with ECPR compared with CCPR (RR 2.11, 95% CI 1.40 to 3.19, $I^2=69\%$; $p=0.0004$; Zhong 2024). This result was consistent with a subgroup analysis using RCTs and PSM data only (RR 1.83, 95% CI: 1.32 to 2.53, $I^2=14\%$; $p=0.0003$; Zhong, 2024). The systematic review and meta-analysis of 28 studies in people with OHCA or IHCA, favourable neurological outcome at 3 months was 12% in people who had ECPR and 10% in people who had CCPR (OR 0.22, 95% CI 0.04 to 1.016, $p=0.07$; 4 studies; Reddy 2025). Favourable neurological outcome at 6 to 12 months was 20% for ECPR and 10% for CCPR (OR 0.41, 95% CI 0.23 to 0.71, $p=0.002$; 7 studies; Reddy 2025).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

After stratifying based on the location of cardiac arrest, subgroup meta-analysis of 3 studies, showed a statistically significantly higher rate of long-term favourable neurological outcomes in ECPR groups compared with CCPR in people with IHCA (OR 2.80, 95% CI: 1.31 to 6.00; $p=0.008$), and in people with OHCA (4 studies: OR 1.96, 95% CI: 1.02 to 3.79; $p=0.045$; Low 2023). Similar outcomes were also seen in a subgroup meta-analysis of 3 studies in people with IHCA (RR 2.17, 95% CI: 1.19 to 3.94, $I^2=0\%$), and of 5 studies in people with OHCA (RR 1.95, 95% CI: 1.01 to 3.59, $I^2=11\%$; Zhong 2024).

OHCA studies

Five systematic reviews (Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024) and 1 long-term RCT follow-up study (Rob 2024), presented long-term survival with favourable neurological outcomes in people with OHCA. Long-term was defined as 3 or 6 months in 2 systematic reviews (Cheema 2023, Scquizzato 2023), or 6 months in 3 systematic reviews (Gomes 2023, Kiyohara 2023, Pagura 2024). Median follow-up was 5.3 years after initial CA in the RCT follow-up study (Rob 2024).

In the systematic review of 3 RCTs including 418 people with OHCA, ECPR was associated with a non-statistically significant higher rate of survival with a favourable neurological outcome at 6 months (28%) compared with standard CPR (19%), RR 1.48, 95% CI: 0.88 to 2.49, $I^2=28\%$; $p=0.14$ (Gomes 2023). A subgroup analysis of people with an initial shockable rhythm showed similar findings to the main analysis, with a non-significant benefit of ECPR (RR 1.50, 95% CI 0.90 to 2.50; $p=0.12$; Gomes 2023).

In 2 systematic reviews and meta-analyses including the same 4 RCTs reporting on people with OHCA, unlike short-term survival with favourable neurological outcomes, the rate of long-term survival with favourable neurological outcomes was not statistically significantly higher in the ECPR groups compared with

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

CCPR (Cheema 2023, Kiyohara 2023). In a systematic review using a Mantel-Haenszel, random-effects meta-analysis, the OR was 1.74, 95% CI: 0.86 to 3.51, $I^2=35\%$; $p=0.12$ (Kiyohara 2023) and in another, the risk ratio (RR) using an inverse-variance, random effects meta-analysis was 1.47, 95% CI: 0.89 to 2.43, $I^2=25\%$; $p=0.13$ (Cheema 2023). However, in a systematic review and meta-analysis of the same 4 RCTs, pooled survival with favourable neurological outcome was 27% for those who had ECPR compared with 18% on CCPR (OR 1.72, 95% CI: 1.09 to 2.70, $I^2=26\%$; $p=0.02$; Scquizzato 2023). Although trial sequential analysis confirmed the statistically significant beneficial effect of ECPR, the sample size included in the meta-analysis did not reach the required information size ($n=520$; Scquizzato 2023). The systematic review by Scquizzato (2023) also reported no difference in people surviving with poor neurological outcomes at the longest follow-up available (0% ECPR, 2% CCPR [OR 0.24, 95% CI: 0.05 to 1.26, $I^2=0\%$; $p=0.780$]).

Two systematic reviews considered long-term survival with favourable neurological outcomes in subgroup meta-analyses in people with an initial shockable rhythm. The rate of long-term survival with favourable neurological outcomes was 34% in the ECPR group compared with 23% in the CCPR group, OR 1.90, 95% CI: 1.16 to 3.13, $I^2=23\%$; $p=0.011$ (Scquizzato 2023), and OR 1.91, 95% CI: 0.90 to 4.03, $I^2=36\%$; $p=0.09$ (Kiyohara 2023).

One systematic review did a subgroup meta-analysis of studies by design (single-centre or multi-centre). A statistically significant difference in long-term survival with favourable neurological outcome with ECPR compared with CCPR was confirmed among single-centre studies (30% ECPR compared with 19% CCPR; OR 1.88, 95% CI: 1.11 to 3.19, $I^2=48\%$; $p=0.02$; Scquizzato 2023).

In a meta-analysis of 6 studies (including RCTs and observational studies) in people with OHCA, the rate of long-term survival with favourable neurological

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

outcome was statistically significantly improved with ECPR (16%) compared with CCPR (8%); 2.72, 95% CI: 1.47 to 5.04, $I^2=47\%$; $p=0.002$; Pagura 2024).

In the long-term follow-up study of the Prague OHCA RCT, 255 people who survived following the initial trial period were followed for a median of 5.3 years (3.8 to 7.2) after initial cardiac arrest. In the ITT analysis, the rate of survival with favourable neurological outcome was 27% in those with ECPR and 19% in those with CCPR (RR 0.90, 95% CI: 0.79 to 1.03, $p=0.13$). In the per protocol and as-treated analysis, RR were 0.86, 95% CI: 0.75 to 0.99, $p=0.035$ and RR 0.84, 95% CI: 0.73 to 0.96, $p=0.007$, respectively (Rob 2024). Only 1 person in each trial arm in the ITT analysis was reported as surviving with a poor neurological outcome (Rob 2024).

Short-term survival

OHCA and IHCA studies

Four publications reporting on 3 systematic reviews presented meta-analyses for short-term survival in both OHCA and IHCA (Low 2023, Low 2024, Zhong 2024; Reddy 2025). Short-term is defined as survival to discharge or 30 days after initial CA.

In the systematic review of 11 studies reporting on adults with OHCA or IHCA, ECPR was associated with improved 30-day survival compared with CCPR in a meta-analysis of 7 studies (OR 1.45, 95% CI: 1.08 to 1.96; $p=0.015$) (Low 2023). In the updated analysis of 9 studies (1 RCT and 8 PSM studies), 30-day survival rate was 24% in the ECPR group and 18% in the CCPR group (OR 1.70, 95% CI: 1.29 to 2.26, $I^2=71\%$; Low 2024). A meta-analysis of 3 RCTs and 9 PSM studies in people with OHCA and IHCA, showed improved short-term survival with ECPR compared with CCPR (RR 1.25, 95% CI 1.00 to 1.56, $I^2=57\%$; $p=0.05$; Zhong 2024). The systematic review and meta-analysis of 28 studies in people with

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

OHCA and IHCA demonstrated that survival at hospital discharge was 20% in the ECPR group compared with 3% in the CCPR group (OR 0.48, 95% CI 0.27 to 0.84, $p=0.01$; 19 studies) and survival at 30 days was 26% for ECPR and 10% for CCPR (OR 0.36, 95% CI 0.25 to 0.51, $p<0.0001$; 7 studies; Reddy 2025),

After stratifying based on the location of cardiac arrest, subgroup meta-analysis of 3 studies showed higher rates of survival at discharge or 1 month in ECPR groups compared with CCPR for people with IHCA (RR 2.03, 95% CI: 1.30 to 3.18, $I^2=0\%$), than for people with OHCA (RR 1.10, 95% CI: 0.91 to 1.34, $I^2=44\%$; Zhong 2024).

OHCA studies

Three systematic reviews (Cheema 2023, Scquizzato 2023, Pagura 2024), 1 retrospective registry study (Inoue 2022), 1 single-centre retrospective PSM study (Shih 2024), and a retrospective registry study (Balucani 2024) presented short-term survival outcomes in people with OHCA. Short-term was defined as hospital discharge or 30 days after initial CA.

In the 2 systematic reviews and meta-analyses including the same 4 RCTs reporting on people with OHCA, no statistically significant difference in short-term survival was observed in people who had ECPR compared with CCPR. In the inverse variance random effects meta-analysis of the 4 RCTs the RR was 1.21, 95% CI: 0.64 to 2.28, $I^2=48\%$; $p=0.55$ (Cheema 2023). In the Mantel-Haenszel random effects meta-analysis, short term survival was 33% in the ECPR group and 27% in the CCPR group (OR 1.35, 95% CI: 0.55 to 3.29, $I^2=0\%$; $p=0.514$; Scquizzato 2023).

In the systematic review of RCTs and observational studies, meta-analysis of 13 studies showed a higher rate of short-term survival with ECPR (20%) than

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

with CCPR (10%) for people with OHCA (OR 1.71, 95% CI: 1.18 to 2.46, $I^2=81\%$; $p=0.004$; Pagura 2024).

In the retrospective SAVE-J II multicentre registry study which included 1,644 adults with OHCA who had ECPR in Japan, the overall rate of survival to hospital discharge was 27% (Inoue 2022). This rate was higher in those with an initial shockable rhythm (32%), and lower in those with a non-shockable rhythm (19% PEA, 11% asystole). Multivariable analysis showed that shorter low flow time was significantly associated with survival to hospital discharge ($p<0.001$) (Inoue 2022).

In the single centre retrospective PSM study reporting on people with OHCA resuscitated with ECPR or CCPR in a high volume emergency department in Taiwan, the in-hospital or 30-day survival rate was 29% among those with ECPR, compared with 8% in those with CCPR. The PSM multivariate analysis reported an adjusted OR (aOR) of 6.02, 95% CI: 2.19 to 16.52 (Shih 2024).

In the retrospective analysis of data from the ELSO registry including 7,460 people with ECPR, the survival rate at discharge was 29% (30% women versus 28% men, $p=0.138$; Balucani 2024).

Long-term survival

OHCA and IHCA studies

Two systematic reviews presented meta-analyses for long-term survival in both OHCA and IHCA (Low 2023, Zhong 2024). Long-term is defined as 3- to 6-month survival after initial CA. Both studies also report on survival up to 1 year.

In the systematic review of 11 studies reporting on adults with OHCA or IHCA, ECPR was associated with improved survival compared with CCPR at 3 months (3 studies [OR 3.98, 95% CI: 1.12 to 14.16; $p=0.033$]), 6 months (6 studies [OR

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

1.87, 95% CI: 1.36 to 2.57; $p=0.0001$), and 1 year (5 studies [OR 1.72, 95% CI: 1.52 to 1.95; $p<0.0001$]) (Low 2023). The meta-analysis of 3 RCTs and 9 PSM studies in people with OHCA and IHCA, also showed improved long-term survival with ECPR compared with CCPR at 3 to 6 months (RR 2.73, 95% CI: 1.67 to 4.48, $I^2=0\%$; $p<0.0001$) and at 1 year (RR 1.92, 95% CI: 1.14 to 3.25, $I^2=0\%$; $p=0.01$; Zhong 2024).

After stratifying based on the location of cardiac arrest, subgroup meta-analysis of 3 studies, showed higher rates of survival to 3 to 6 months or 1 year in ECPR groups compared with CCPR for people with OHCA (RR 3.16, 95% CI: 1.36 to 7.38, $I^2=0\%$), than for people with IHCA (RR 1.92, 95% CI: 1.14 to 3.25, $I^2=0\%$; Zhong 2024).

OHCA studies

Three systematic reviews (Cheema 2023, Kiyohara 2023, Scquizzato 2023) and 1 long-term RCT follow-up study (Rob 2024), presented long-term survival in people with OHCA. Long-term was defined in 1 systematic review as 3 or 6 months (Cheema 2023), in 1 systematic review as 6 months (Kiyohara 2023), and in 1 systematic review as 30 days to 6 months. In Scquizzato (2023), survival was reported at the longest follow-up available. Median follow-up was 5.3 years after initial CA in the RCT follow-up study (Rob 2024).

In the 3 systematic reviews and meta-analyses including the same 4 RCTs reporting on people with OHCA, no statistically significant difference in long-term survival was observed in people who had ECPR compared with CCPR (Cheema 2023, Kiyohara 2023, Scquizzato 2023). In the inverse variance random effects meta-analysis of the 4 RCTs the RR for survival at 3 or 6 months was RR 1.32, 95% CI: 0.18 to 9.5, $I^2=64\%$; $p=0.79$ (Cheema 2023). In the Mantel-Haenszel random effects meta-analyses, survival at the longest follow-up was 28% in the ECPR group and 22% in the CCPR group (OR 1.31, 95% CI: 0.49 to 3.49,

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

$I^2=58\%$; $p=0.592$; Scquizzato 2023) and the OR for survival at 6 months was 1.50, 95% CI: 0.67 to 3.36, $I^2=50\%$; $p=0.33$ (Kiyohara 2023).

In the long-term follow-up study of the Prague OHCA RCT, 255 people who survived following the initial trial period were followed for a median of 5.3 years after initial cardiac arrest. In the ITT analysis, Kaplan-Meier estimates of survival were 28% in those with ECPR and 20% in those with CCPR (log rank $p=0.01$). In the per protocol and as-treated analysis, the difference between groups were log rank $p=0.008$ and $p<0.001$, respectively (Rob 2024).

Mortality

OHCA and IHCA studies

Two publications reporting on 1 systematic review presented meta-analyses for in-hospital mortality in both OHCA and IHCA (Low 2023, Low 2024).

In the meta-analyses of 3 RCTs and 8 PSMs of adults with OHCA or IHCA ($n=9,192$), ECPR was associated with significant reduction in mortality (OR 0.67, 95% CI: 0.51 to 0.87, $I^2=42\%$; $p=0.0034$ (Low 2023). In-hospital mortality in the updated analysis of 3 RCTs and 11 PSMs of adults with OHCA or IHCA ($n=14,048$) was 77% in ECPR group and 83% in the CCPR group (OR 0.63, 95% CI: 0.50 to 0.79, $I^2=64\%$; Low 2024).

After stratifying based on the location of cardiac arrest, subgroup meta-analysis of 4 studies, showed a significant reduction in mortality in people having ECPR compared with CCPR in those with IHCA (OR 0.42, 95% CI: 0.25 to 0.70, $I^2=0\%$; $p=0.0009$; Low 2023), and in people with OHCA (3 RCTs and 6 PSMs: OR 0.67, 95% CI: 0.51 to 0.88, $I^2=74\%$; Low 2024). Pooled HRs across studies showed that longer duration of CPR was associated with increased mortality (HR per min 1.01, 95% CI 1.00 to 1.01; $p=0.0001$), and an initial presentation with a shockable rhythm was associated with reduced mortality (HR 0.52, 95% CI 0.32

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

to 0.86; $p=0.011$). However, it did not show an association between age and mortality (HR per year 1.02, 95% CI 0.98 to 1.06; $p=0.41$; Low 2023).

OHCA

One systematic review reported in-hospital mortality in people with OHCA. In the systematic review of 3 RCTs including 418 people with OHCA, the mean absolute rate of in-hospital mortality was not significantly lower in the ECPR group (71%) compared with the CCPR group (79%; RR 0.89, 95% CI: 0.74 to 1.07, $I^2=53\%$; $p=0.23$) (Gomes 2023).

Safety

Of the 11 included studies in the key evidence, only 4 studies reported on any safety outcomes (2 systematic reviews [Low 2023, Cheema 2023], 1 registry study [Inoue 2022] and 1 long-term RCT follow-up study [Rob 2024]).

Bleeding

Two systematic reviews and 1 registry study reported bleeding events. In the systematic review of 11 studies including adults with OHCA or IHCA, people who had ECPR were more likely to have bleeding than those who had CCPR (meta-analysis 4 studies: OR 4.84, 95% CI: 1.91 to 12.24; $p=0.0009$) (Low 2023).

Similarly, in a meta-analysis of 6 studies in people with OHCA and IHCA, the bleeding rate was 35% in the ECPR group and 4% in the CCPR group (OR 0.08, 95% CI 0.03 to 0.24, $p<0.0001$; Reddy 2025).

In the retrospective SAVE-J II multicentre registry study which included 1,644 adults with OHCA who had ECPR in Japan, rates of cannulation-related bleeding were 16% (268 of 1,644), and rates of haemorrhage were 8.5% (139 of 1,644; Inoue 2022).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

In the retrospective ELSO registry study of 7,460 people with ECPR, the haemorrhagic complication rate was 27% (Balucani 2024).

Adverse events

One systematic review of 4 RCTs in people with OHCA reported the comparative rates of adverse events between ECPR and CCPR. Of the 4 RCTs included in the systematic review, only 2 studies reported the rate of adverse events. Meta-analysis indicated that ECPR was associated with higher rate of adverse events than CCPR (RR: 3.22, 95% CI: 1.18 to 8.80, $I^2=63\%$; $p=0.02$; Cheema 2023).

In the retrospective SAVE-J II multicentre registry study which included 1,644 adults with OHCA who had ECPR in Japan, the overall reported complication rate during ECPR was 33% (535 of 1,644). This included procedure-related complications (21%, 346 of 1,644) such as cannula malposition and unsuccessful cannulation, ECMO-related complications (3%, 50 of 1,644) and ischaemia (2%, 26 of 1,644) as well as the bleeding complications reported in the section above (Inoue 2022).

In the retrospective ELSO registry study of 7,460 people with ECPR, the overall in-hospital rate of neurological complications was 17% at hospital discharge (Balucani 2024). The overall in-hospital rate of infectious, limb, mechanical, metabolic and cardiovascular complications was 7%, 7%, 15%, 20% and 37%, respectively.

Long-term adverse events

In the long-term follow-up study of the Prague OHCA RCT, 255 people who survived following the initial trial period were followed for a median of 5.3 years after initial cardiac arrest. During the follow-up, 39 people (32%) in the ECPR group and 30 (23%) in the CCPR group were discharged from the hospital or long-term hospital facilities after the initial CA event (median time to discharge

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

19.5 days, IQR 12.5 to 32 days). Of these, 10% (4 of 39) people in the ECPR group and 20% (6 of 30) in the CCPR group died during the follow-up (RR 0.51, 95% CI: 0.16 to 1.66, $p=0.26$). At least one rehospitalisation after discharge occurred in 77% (30 of 39) of the ECPR group and 60% (18 of 30) of the CCPR group (RR 1.28, 95% CI: 0.91 to 1.8, $p=0.15$). At least one cardiovascular rehospitalisation occurred in 64% (25 of 39) of people in the ECPR group and 50% (15 of 30) of those in the CCPR group (RR 1.28, 95% CI: 0.84 to 1.97, $p=0.26$). One person in each group reported myocardial infarction after discharge, and 1 person who had CCPR had a stroke. Hospitalisation for heart failure occurred in 5% (2 of 39) of people in the ECPR group and 10% (3 of 30) in the CCPR group, and for ventricular arrhythmia in 3% (1 of 39) people in the ECPR group and 10% (3 of 30) in the CCPR group (Rob 2024).

Anecdotal and theoretical adverse events

Expert advice was sought from consultants who have been nominated or ratified by their professional society or royal college. They were asked if they knew of any other adverse events for this procedure that they had heard about (anecdotal), which were not reported in the literature. They were also asked if they thought there were other adverse events that might possibly occur, even if they had never happened (theoretical).

They listed the following anecdotal and theoretical adverse events:

- Left ventricle overloading
- Deep vein thrombosis
- Arteriovenous fistula
- Pseudoaneurysm
- Harlequin syndrome
- Haemolysis
- Intracerebral haemorrhage

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- Major pulmonary bleed
- Failure to cannulate during cardiac arrest
- Malposition of the cannula
- Device clotting
- Air entrainment/embolus
- Embolism
- Oxygenator failure
- Consumption coagulopathy
- Acquired Von Willebrand syndrome
- Systemic inflammatory response syndrome (SIRS)
- Multi-organ failure including kidney, liver, and pancreas.

Sixteen professional expert questionnaires were submitted. Find full details of what the professional experts said about the procedure in the [specialist advice questionnaires for this procedure](#).

Validity and generalisability

- Most of the key papers included are systematic reviews with meta-analyses. There was a significant amount of overlap identified across the systematic reviews included in the overview; much of the available evidence identified in this review is based on the same RCTs. Evidence was mainly for adults resuscitated from OHCA; 2 systematic reviews included studies done in IHCA.
- No RCTs were available for the IHCA population.
- Recent systematic reviews of RCTs in OHCA have varying conclusions of the benefit of ECPR compared with CCPR. This is reflected in the conflicting conclusions of the RCTs themselves.
 - All 3 RCTs and 1 pilot RCT were small and had a high rate of crossovers and deviations from the intended intervention.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- Authors of the systematic reviews noted that the RCTs may have insufficient power to observe significant differences between intervention arms.
- There was a high heterogeneity across RCTs, such as the proportion of people with non-shockable rhythms enrolled, the definition of refractory OHCA (5 minutes or 15 minutes without ROSC), the time between cardiac arrest and ECMO initiation, whether there was a standardised emergency medical services protocol in place, whether the RCT was a single or multicentre study, the centre's level of experience with ECPR, and outcome endpoints.
- In the 3 RCTs reported in 8 systematic reviews (Low 2023, Low 2024, Zhong 2024, Gomes 2023, Cheema 2023, Kiyohara 2023, Scquizzato 2023, Pagura 2024), ECPR was initiated in 64% (Prague OHCA), 66% (INCEPTION) and 86% (ARREST) of the people randomised to the intervention arm. In the additional pilot RCT included in 3 systematic reviews (Cheema 2023, Kiyohara 2023, Scquizzato 2023) 42% randomised to the intervention arm received ECPR (Scquizzato 2023). In the ARREST trial, none of the people randomised to the standard CPR group received ECPR, while the rate of crossover was 8% in Prague OHCA and 5% in INCEPTION (Scquizzato 2023).
- Some systematic reviews included observational studies, which are at higher risk of bias, particularly in ECPR, as the decision to treat is based on clinician's evaluation of comorbidities and prognostic factors which may have strong impact on the outcome. Authors of the Shih (2024) observational study noted that there may be selection bias for ECPR as there was no protocol for who got ECPR, therefore it was likely people with better prognosis were selected for ECPR than randomised studies.
- Only 1 follow-up study for 1 RCT reported any outcomes with a follow-up longer than 6 months.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Ongoing trials

- [ON-SCENE Initiation of Extracorporeal CardioPulmonary Resuscitation During Refractory Out-of-Hospital Cardiac Arrest \(ON-SCENE\)](#) (NCT04620070); RCT; Netherlands; n=390; completion date July 2026
- [Pre-hospital ECMO or Conventional Resuscitation for Refractory Cardiac Arrest \(PACER\)](#) (NCT06177730); RCT; Australia and New Zealand; n=10; completion date December 2024

Existing assessments of this procedure

[European Resuscitation Council \(ERC\): Guidelines for resuscitation \(2021\)](#) on ECPR recommends as follows:

- Consider ECPR as a rescue therapy for selected patients with cardiac arrest when conventional advanced life support (ALS) measures are failing or to facilitate specific interventions (e.g. coronary angiography and percutaneous coronary intervention (PCI), pulmonary thrombectomy for massive pulmonary embolism, rewarming after hypothermic cardiac arrest) in settings in which it can be implemented.

[International Liaison Committee on Resuscitation \(ILCOR\) International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations \(CoSTR\) \(2024\)](#)

- ECPR may be considered as a rescue therapy for selected adults with out-of-hospital cardiac arrest when conventional cardiopulmonary resuscitation is failing to restore spontaneous circulation in settings where this can be implemented (weak recommendation, low-certainty evidence).
- ECPR may be considered as a rescue therapy for selected adults with in-hospital cardiac arrest when conventional cardiopulmonary resuscitation is

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

failing to restore spontaneous circulation in settings where this can be implemented (weak recommendation, very low–certainty evidence).

[American Heart Association Focused Update on Advanced Cardiovascular Life Support](#) 2019

- There is insufficient evidence to recommend the routine use of ECPR for patients with cardiac arrest.
- ECPR may be considered for selected patients as rescue therapy when conventional CPR efforts are failing in settings in which it can be expeditiously implemented and supported by skilled providers.

Related NICE guidance

Interventional procedures

[Extracorporeal membrane oxygenation \(ECMO\) for acute heart failure in adults](#) (2014) NICE interventional procedures guidance [IPG 482]. (Recommendation: special arrangements).

Professional societies

- The Intensive Care Society
- Society for Cardiothoracic Surgery in Great Britain & Ireland
- Royal College of Anaesthetists
- Royal College of Surgeons
- Faculty of Intensive Care Medicine
- British Society for Heart Failure
- NHS Blood and Transplant
- British Cardiovascular Society
- European Extracorporeal Life Support Organisation

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Company engagement

NICE asked companies who manufacture a device potentially relevant to this procedure for information on it. NICE received 2 completed submissions. These were considered by the interventional procedures technical team and any relevant points have been taken into consideration when preparing this overview.

References

1. Low CJW, Ramanathan K, Ling RR et al. (2023) Extracorporeal cardiopulmonary resuscitation versus conventional cardiopulmonary resuscitation in adults with cardiac arrest: a comparative meta-analysis and trial sequential analysis. *The Lancet. Respiratory Medicine* 11(10): 883-893
2. Low CJW, Ling RR, Ramanathan K et al. (2024) Extracorporeal cardiopulmonary resuscitation versus conventional CPR in cardiac arrest: an updated meta-analysis and trial sequential analysis. *Critical Care (London, England)* 28(1): 57
3. Zhong H, Yin Z, Wang, Y et al. (2024) Comparison of prognosis between extracorporeal CPR and conventional CPR for patients in cardiac arrest: a systematic review and meta-analysis. *BMC Emergency Medicine* 24(1): 128
4. Gomes DA, Presume J, Ferreira J et al. (2023) Extracorporeal cardiopulmonary resuscitation for refractory out-of-hospital cardiac arrest: a systematic review and meta-analysis of randomized clinical trials. *Internal and Emergency Medicine* 18(7): 2113-2120
5. Cheema HA, Shafiee A, Jafarabady K et al. (2023) Extracorporeal cardiopulmonary resuscitation for out-of-hospital cardiac arrest: A meta-analysis of randomized controlled trials. *Pacing and Clinical Electrophysiology: PACE* 46(10): 1246-1250
6. Kiyohara Y, Kampaktsis PN, Briasoulis A et al. (2023) Extracorporeal membrane oxygenation-facilitated resuscitation in out-of-hospital cardiac arrest: a meta-analysis of randomized controlled trials. *Journal of Cardiovascular Medicine* 24(7): 414-419
7. Scquizzato T, Bonaccorso A, Swol J et al. (2023) Refractory out-of-hospital cardiac arrest and extracorporeal cardiopulmonary resuscitation: A meta-analysis of randomized trials. *Artificial Organs* 47(5): 806-816
8. Pagura L, Fabris E, Rakar S et al. (2024) Does extracorporeal cardiopulmonary resuscitation improve survival with favorable neurological

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

outcome in out-of-hospital cardiac arrest? A systematic review and meta-analysis. *Journal of Critical Care* 84: 154882

9. Rob D, Farkasovska K, Kreckova M et al. (2024) Effect of intra-arrest transport, extracorporeal cardiopulmonary resuscitation and immediate invasive assessment in refractory out-of-hospital cardiac arrest: a long-term follow-up of the Prague OHCA trial. *Critical Care (London, England)* 28(1): 125
10. Inoue A, Hifumi T, Sakamoto T et al. (2022) Extracorporeal cardiopulmonary resuscitation in adult patients with out-of-hospital cardiac arrest: a retrospective large cohort multicenter study in Japan. *Critical Care (London, England)* 26(1): 129
11. Shih H-M, Lin W-J, Lin Y-C et al. (2024) Extracorporeal cardiopulmonary resuscitation for patients with refractory out-of-hospital cardiac arrest: a propensity score matching, observational study. *Scientific Reports* 14(1): 9912
12. Reddy S, Garcia S, Hostetter LJ et al. (2025) Extracorporeal-CPR Versus Conventional-CPR for Adult Patients in Out of Hospital Cardiac Arrest-Systematic Review and Meta-Analysis. *Journal of Intensive Care Medicine* 40: 207–17
13. Balucani C, Canner JK, Tonna JE et al. (2024) Sex-Related Differences in Utilization and Outcomes of Extracorporeal Cardio-Pulmonary Resuscitation for Refractory Cardiac Arrest. *ASAIO Journal* 70 750-757

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Appendix A: Methods and literature search strategy

Methods and literature search strategy

NICE has identified studies and reviews relevant to venoarterial extracorporeal membrane oxygenation (VA ECMO) in the following indications from the medical literature:

- acute heart failure in adults
- extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest
- postcardiotomy cardiogenic shock in adults.

The search was initially developed for the acute heart failure indication only (Tables 4a and 4b) and then modified and updated to cover the additional 2 indications (Table 4c).

Search strategy design and peer review

This search report is informed by the [Preferred Reporting Items for Systematic reviews and Meta-Analyses literature search extension \(PRISMA-S\)](#).

A NICE information specialist ran the literature searches for acute heart failure in adults on 18th September 2024 and updated them on 12th May 2025. The search strategy was modified and updated on 19th June 2025 to incorporate the 2 additional interventions. See the [search strategy history](#) for the full search strategy for each database. Relevant published studies identified during consultation or resolution that are published after this date may also be considered for inclusion.

The principal search strategy was developed in MEDLINE ALL (Ovid interface). It was adapted for use in each of the databases listed in [table 4a](#), taking into account the database's size, search functionality and subject coverage. The MEDLINE ALL strategy was quality assured by a NICE senior information specialist. All translated search strategies were peer reviewed to ensure their accuracy. The quality assurance and peer review procedures were adapted from IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

the [Peer Review of Electronic Search Strategies \(PRESS\) 2015 evidence-based checklist](#).

Review management

The search results were managed in EPPI-Reviewer version 5 (EPPI-R5). Duplicates were removed in EPPI-R5 using a 2-step process. First, automated deduplication was done using a high-value algorithm. Second, manual deduplication was used to assess low-probability matches. All decisions about inclusion, exclusion and deduplication were recorded and stored.

Limits and restrictions

The CENTRAL database search removed trial registry records and conference material. The Embase search excluded conference material, letters and editorial. We excluded the following publication types in MEDLINE: letter, historical article, comment, editorial, news and case reports. English language limits were applied to the search when possible in the database.

The search was limited from March 2013 to the latest update. The date limit was included to update searches undertaken for an earlier version of this guidance.

The limit to remove animal studies in the searches is standard NICE practice, which has been adapted from [Dickersin K, Scherer R, Lefebvre C \(1994\) Systematic Reviews: Identifying relevant studies for systematic reviews. BMJ 309\(6964\): 1286](#).

Main search**Table 4a Main search results**

Database	Date searched	Database platform	Database segment or version	Number of results downloaded
Cochrane Central Register of Controlled Trials (CENTRAL)	18/09/24	Wiley	Issue 8 of 12, August 2024	410
Cochrane Database of Systematic Reviews (CDSR)	20/09/24	Wiley	Issue 9 of 12, September 2024	13
Embase	20/09/24	Ovid	1974 to 2024 September 17	2101
INAHTA International HTA Database	18/09/24	https://database.inahta.org /	-	24
MEDLINE ALL	18/09/24	Ovid	1946 to Sept 17, 2024	1454

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Update search

Table 4b Update search results 1

Database	Date searched	Database platform	Database segment or version	Number of results downloaded
Cochrane Central Register of Controlled Trials (CENTRAL)	12/05/2025	Wiley	Issue 4 of 12, April 2025	39
Cochrane Database of Systematic Reviews (CDSR)	12/05/2025	Wiley	Issue 5 of 12, May 2025	0
Embase	12/05/2025	Ovid	1974 to 2025 May 09	54
INAHTA International HTA Database	12/05/2025	https://database.inahta.org/		4
MEDLINE ALL	12/05/2025	Ovid	1946 to May 09, 2025	195

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Additional update search

Table 4c Update search results 2

This version of the search was modified to include 2 additional indications and searched from March 2013 to latest update.

Database	Date searched	Database platform	Database segment or version	Number of results downloaded
Cochrane Central Register of Controlled Trials (CENTRAL)	19/06/25	Wiley	Issue 6 of 12, 2025	295
Cochrane Database of Systematic Reviews (CDSR)	19/05/25	Wiley	Issue 6 of 12, 2025.	0
Embase	19/06/25	Ovid	1974 to 2025 June 17	4461
INAHTA International HTA Database	19/06/25	https://database.inahta.org/	-	29
MEDLINE ALL	19/06/25	Ovid	1946 to June 18, 2025	4707

Search strategy history – initial search strategy September 2024

MEDLINE ALL search strategy

1 , Heart Failure/th , 29,868

2 , Acute disease/th , 1,194

3 , 1 and 2 , 11

4 , *Cardiomyopathies/th , 1,150

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- 5 , *Shock cardiogenic/th , 2,135
- 6 , Myocardial Stunning/th [Therapy] , 155
- 7 , Myocarditis/th [Therapy] , 1,294
- 8 , *Myocardial infarction/ , 138,977
- 9 , Out-of-Hospital Cardiac Arrest/th [Therapy] , 5,734
- 10 , ((acute* or server*) adj (heart* or cardiac* or myocard* or cardio* or ventric*)
adj (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or
fault* or shock*)).ti,ab. , 9,513
- 11 , Myocardit*.ti,ab. , 21,440
- 12 , ((Postpartum* or post-partum* or peripartum* or peri-partum*) adj
cardiomyopath*).ti,ab. , 1,697
- 13 , PPCM.ti,ab. , 671
- 14 , (myocard* adj (stun* or hibernat* or infract*)).ti,ab. , 2,258
- 15 , Primary Graft Dysfunction/th [Therapy] , 99
- 16 , (primary* adj graft* adj dysfunct*).ti,ab. , 1,392
- 17 , or/3-16 , 182,062
- 18 , *Cardiopulmonary Resuscitation/mt [Methods] , 4,116
- 19 , *Extracorporeal Membrane Oxygenation/ , 13,895
- 20 , ECMO.ti. , 3,217
- 21 , *Extracorporeal Circulation/mt [Methods] , 1,090
- 22 , (extracorp* adj circulat*).ti,ab. , 8,596
- 23 , (extracorp* adj ((cardiopulmon* adj resuscitat*) or CPR)).ti,ab. , 1,229
- 24 , ECPR.ti. , 154
- 25 , (Biomedicus adj pump*).ti,ab. , 45
- 26 , (Maquet* adj rotaflow*).ti,ab. , 12
- 27 , (jostra adj (pump* or rotaflow*)).ti,ab. , 5

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

28 , (levitronix adj (centrimag* or pump* or system* or oxygen*)).ti,ab. , 54
 29 , (Medos adj (Hilite* or oxygen*)).ti,ab. , 22
 30 , left ventricle assist device.ti,ab. , 106
 31 , or/18-30 , 28,477
 32 , 17 and 31 , 2,725
 33 , animals/ not human/ , 5,225,551
 34 , 32 not 33 , 2,680
 35 , limit 34 to english language , 2,503
 36 , limit 35 to ed=20130331-20240930 , 2,028
 37 , limit 36 to (letter or historical article or comment or editorial or news or case reports) , 574
 38 , 36 not 37 , 1,454

[Embase] search strategy

1 , heart failure/th [Therapy] , 15,752
 2 , acute disease/th [Therapy] , 2,395
 3 , 1 and 2 , 10
 4 , *cardiomyopathy/th [Therapy] , 1,144
 5 , *cardiogenic shock/th [Therapy] , 2,129
 6 , stunned heart muscle/th [Therapy] , 53
 7 , myocarditis/th [Therapy] , 864
 8 , *heart infarction/ , 110,365
 9 , primary graft dysfunction/th [Therapy] , 94
 10 , "out of hospital cardiac arrest"/th [Therapy] , 3,862

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

11 , ((acute* or server*) adj (heart* or cardiac* or myocard* or cardio* or ventric*) adj (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock*)).ti,ab. , 17,537

12 , Myocardit*.ti,ab. , 31,093

13 , ((Postpartum* or post-partum* or peripartum* or peri-partum*) adj cardiomyopath*).ti,ab. , 2,835

14 , PPCM.tw. , 1,261

15 , (myocard* adj (stun* or hibernat* or infract*)).ti,ab. , 3,555

16 , (primary* adj graft* adj dysfunct*).tw. , 3,009

17 , or/3-16 , 173,201

18 , *resuscitation/ , 60,473

19 , *extracorporeal oxygenation/ , 16,545

20 , ECMO.ti. , 7,837

21 , *extracorporeal circulation/ , 9,094

22 , (extracorp* adj circulat*).ti,ab. , 9,683

23 , (extracorp* adj ((cardiopulmon* adj resuscitat*) or CPR)).ti,ab. , 1,851

24 , ECPR.ti. , 352

25 , (Biomedicus adj pump*).ti,ab. , 50

26 , (Maquet* adj rotaflow*).ti,ab. , 31

27 , (jostra adj (pump* or rotaflow*)).ti,ab. , 16

28 , (levitronix adj (centrimag* or pump* or system* or oxygen*)).ti,ab. , 150

29 , (Medos adj (Hilite* or oxygen*)).ti,ab. , 44

30 , left ventricle assist device.ti,ab. , 217

31 , or/18-30 , 96,434

32 , 17 and 31 , 5,350

33 , Nonhuman/ not Human/ , 5,532,522

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

34 , 32 not 33 , 5,275

35 , limit 34 to letter/ or (letter or editorial).pt. , 2,165,352

36 , 34 not 35 , 4,904

37 , limit 36 to dc=20130331-20240930 , 3,599

38 , limit 37 to english language , 3,481

39 , (conference abstract* or conference review or conference paper or
conference proceeding).db,pt,su. , 6,020,541

40 , 38 not 39 , 2,101

Cochrane Library (CDSR) search strategy

#1 MeSH descriptor: [Heart Failure] explode all trees and with qualifier(s):
[therapy - TH] 2591

#2 MeSH descriptor: [Acute Disease] explode all trees and with qualifier(s):
[therapy - TH] 118

#3 #1 and #2 0

#4 MeSH descriptor: [Cardiomyopathies] explode all trees and with
qualifier(s): [therapy - TH] 248

#5 MeSH descriptor: [Shock, Cardiogenic] explode all trees and with
qualifier(s): [therapy - TH] 177

#6 MeSH descriptor: [Myocardial Stunning] explode all trees and with
qualifier(s): [therapy - TH] 3

#7 MeSH descriptor: [Myocarditis] explode all trees and with qualifier(s):
[therapy - TH] 13

#8 MeSH descriptor: [Myocardial Infarction] explode all trees and with
qualifier(s): [therapy - TH] 3337

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- #9 MeSH descriptor: [Primary Graft Dysfunction] explode all trees and with qualifier(s): [therapy - TH] 3
- #10 MeSH descriptor: [Out-of-Hospital Cardiac Arrest] explode all trees and with qualifier(s): [therapy - TH] 539
- #11 ((acute* or server*) near/1 (heart* or cardiac* or myocard* or cardio* or ventric*) near/1 (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock*)) 2663
- #12 Myocardit* 1421
- #13 (Postpartum* or post-partum* or peripartum* or peri-partum*) near/1 cardiomyopath* 47
- #14 PPCM39
- #15 (myocard* near/1 (stun* or hibernat* or infract*)) 342
- #16 (primary* near/1 graft* near dysfunct*) 146
- #17 #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 8646
- #18 MeSH descriptor: [Cardiopulmonary Resuscitation] this term only 1688
- #19 MeSH descriptor: [Extracorporeal Membrane Oxygenation] this term only 361
- #20 ECMO 1101
- #21 MeSH descriptor: [Extracorporeal Circulation] this term only and with qualifier(s): [methods - MT] 120
- #22 (extracorp* near/1 circulat*) 1423
- #23 (extracorp* near/1 ((cardiopulmon* near resuscitat*) or CPR)) 71
- #24 ECPR 112
- #25 (Biomedicus near/1 pump*) 3
- #26 (Maquet* rotaflow*) 3

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- #27 jostra near/1 (pump* or rotaflow*) 1
- #28 (levitronix near/1 (centrimag* or pump* or system* or oxygen*)) 0
- #29 Medos near/1 (Hilite* or oxygen*) 0
- #30 left ventricle assist device 219
- #31 #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28
or #29 or #30 4577
- #32 #17 AND #31 494
- #33 "conference":pt or (clinicaltrials or trialsearch):so 777352
- #34 #32 NOT #33 with Cochrane Library publication date Between Mar 2013
and Sep 2024, in Cochrane Reviews 13

Cochrane Library CENTRAL search strategy

- #1 MeSH descriptor: [Heart Failure] explode all trees and with qualifier(s):
[therapy - TH] 2591
- #2 MeSH descriptor: [Acute Disease] explode all trees and with qualifier(s):
[therapy - TH] 118
- #3 #1 and #2 0
- #4 MeSH descriptor: [Cardiomyopathies] explode all trees and with
qualifier(s): [therapy - TH] 248
- #5 MeSH descriptor: [Shock, Cardiogenic] explode all trees and with
qualifier(s): [therapy - TH] 177
- #6 MeSH descriptor: [Myocardial Stunning] explode all trees and with
qualifier(s): [therapy - TH] 3
- #7 MeSH descriptor: [Myocarditis] explode all trees and with qualifier(s):
[therapy - TH] 13

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- #8 MeSH descriptor: [Myocardial Infarction] explode all trees and with
qualifier(s): [therapy - TH] 3337
- #9 MeSH descriptor: [Primary Graft Dysfunction] explode all trees and with
qualifier(s): [therapy - TH] 3
- #10 MeSH descriptor: [Out-of-Hospital Cardiac Arrest] explode all trees and
with qualifier(s): [therapy - TH] 539
- #11 ((acute* or server*) near/1 (heart* or cardiac* or myocard* or cardio* or
ventric*) near/1 (failur* or decompensation* or insufficient* or dysfunct* or stand*
or still* or fault* or shock*)) 2663
- #12 Myocardit* 1421
- #13 (Postpartum* or post-partum* or peripartum* or peri-partum*) near/1
cardiomyopath* 47
- #14 PPCM39
- #15 (myocard* near/1 (stun* or hibernat* or infract*)) 342
- #16 (primary* near/1 graft* near dysfunct*) 146
- #17 #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14
or #15 or #168646
- #18 MeSH descriptor: [Cardiopulmonary Resuscitation] this term only 1688
- #19 MeSH descriptor: [Extracorporeal Membrane Oxygenation] this term only
361
- #20 ECMO 1101
- #21 MeSH descriptor: [Extracorporeal Circulation] this term only and with
qualifier(s): [methods - MT] 120
- #22 (extracorp* near/1 circulat*) 1423
- #23 (extracorp* near/1 ((cardiopulmon* near resuscitat*) or CPR)) 71
- #24 ECPR 112

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

#25 (Biomedicus near/1 pump*) 3
 #26 (Maquet* rotaflow*) 3
 #27 jostra near/1 (pump* or rotaflow*) 1
 #28 (levitronix near/1 (centrimag* or pump* or system* or oxygen*)) 0
 #29 Medos near/1 (Hilite* or oxygen*) 0
 #30 left ventricle assist device 219
 #31 #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28
 or #29 or #30 4577
 #32 #17 AND #31 494
 #33 "conference":pt or (clinicaltrials or trialsearch):so 777352
 #34 #32 NOT #33 with Cochrane Library publication date Between Mar 2013
 and Sep 2024, in Trials 410

INAHTA HTA Database search strategy

1 , "Heart Failure"[mh] , 252
 2 , "Acute Disease"[mh] , 46
 3 , #2 AND #1 , 2
 4 , "Cardiomyopathies"[mh] , 21
 5 , "Shock, Cardiogenic"[mh] , 11
 6 , "Myocardial Stunning"[mh] , 1
 7 , "Myocarditis"[mh] , 1
 8 , "Myocardial Infarction"[mh] , 123
 9 , "Out-of-Hospital Cardiac Arrest"[mh] , 10
 10 , ((acute* or server*) and (heart* or cardiac* or myocard* or cardio* or ventric*)
 and (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or
 fault* or shock*)). , 149

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- 11 , Myocardit* , 5
- 12 , ((Postpartum* or post-partum* or peripartum* or peri-partum*) AND cardiomyopath*) , 1
- 13 , PPCM , 0
- 14 , (myocard* and (stun* or hibernat* or infract*)) , 2
- 15 , "Primary Graft Dysfunction"[mh] , 0
- 16 , (primary* AND graft* AND dysfunct*) , 3
- 17 , #16 OR #15 OR #14 OR #13 OR #12 OR #11 OR #10 OR #9 OR #8 OR #7 OR #6 OR #5 OR #4 OR #3 , 291
- 18 , "Cardiopulmonary Resuscitation"[mh] , 23
- 19 , "Extracorporeal Membrane Oxygenation"[mh] , 29
- 20 , ECMO , 31
- 21 , "Extracorporeal Circulation"[mh] , 9
- 22 , (extracorp* AND circulat*) , 13
- 23 , (extracorp* AND ((cardiopulmon* AND resuscitat*) or CPR)) , 8
- 24 , ECPR , 4
- 25 , (Biomedicus AND pump*) , 0
- 26 , Maquet* and rotaflow* , 0
- 27 , (jostra and (pump* or rotaflow*)) , 0
- 28 , (levitronix AND (centrimag* or pump* or system* or oxygen*)) , 0
- 29 , (Medos AND (Hilite* or oxygen*)) , 0
- 30 , left ventricle assist device , 3
- 31 , #30 OR #29 OR #28 OR #27 OR #26 OR #25 OR #24 OR #23 OR #22 OR #21 OR #20 OR #19 OR #18 , 74
- 32 , #31 AND #17 , 24

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Search strategy history – update search strategy June 2026

This version of the search was modified to include 2 additional indications and searched from March 2013 to latest update.

MEDLINE ALL search strategy

- 1, Heart Failure/th [Therapy], 31,048
- 2, Acute Disease/th [Therapy], 1,222
- 3, 1 and 2, 11
- 4, *Cardiomyopathies/th, 1,952
- 5, *Shock, Cardiogenic/th, 2,922
- 6, Myocardial Stunning/th [Therapy], 155
- 7, Myocarditis/th [Therapy], 1,333
- 8, *Myocardial infarction/th [Therapy], 19,394
- 9, Out-of-Hospital Cardiac Arrest/th [Therapy], 6,031
- 10, ((acute* or severe* or refract*) adj (heart* or cardiac* or myocard* or cardio* or ventric*) adj (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock* or arrest* or stunn*)).tw., 18,262
- 11, (cardiogen* adj shock).tw., 17,331
- 12, Myocardit*.tw., 22,410
- 13, ((Postpartum* or post-partum* or peripartum* or peri-partum*) adj cardiomyopath*).tw., 1,811
- 14, (postcardiotomy or Post-cardiotomy).tw., 1,230
- 15, PPCM.tw., 717
- 16, (myocard* adj (stun* or hibernat* or infarct*)).tw., 241,791
- 17, Primary Graft Dysfunction/th [Therapy], 109
- 18, (primary* adj graft* adj dysfunct*).tw., 1,543

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- 19, or/3-18, 301,507
- 20, *Cardiopulmonary Resuscitation/mt [Methods], 5,700
- 21, *Extracorporeal Membrane Oxygenation/, 14,654
- 22, ECMO.tw., 14,889
- 23, *Extracorporeal Circulation/mt, 1,259
- 24, (extracorp* adj circulat*).tw., 8,706
- 25, (extracorp* adj ((cardiopulmon* adj resuscitat*) or CPR)).tw., 1,377
- 26, ECPR.ti,ab., 1,006
- 27, (Biomedicus adj pump*).tw., 45
- 28, (Maquet* adj rotaflow*).tw., 12
- 29, (jostra adj (pump* or rotaflow*)).tw., 5
- 30, (levitronix adj (centrimag* or pump* or system* or oxygen*)).tw., 54
- 31, (Medos adj (Hilite* or oxygen*)).tw., 22
- 32, (left adj ventricle adj assist adj device).tw., 107
- 33, or/20-32, 35,941
- 34, 19 and 33, 6,390
- 35, animals/ not humans/, 5,314,500
- 36, 34 not 35, 6,296
- 37, (exp child/ or exp pediatrics/ or exp infant/ or exp adolescent/) not (exp adult/ or exp middle age/ or exp aged/), 2,281,857
- 38, 36 not 37, 5,768
- 39, limit 38 to english language, 5,398
- 40, limit 39 to ed=20130901-20250630, 3,883
- 41, limit 39 to dt=20130901-20250630, 4,610
- 42, 40 or 41, 4,707

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

EMBASE search strategy

- 1, heart failure/th [Therapy], 15,823
- 2, acute disease/th [Therapy], 2,430
- 3, 1 and 2, 10
- 4, *cardiomyopathy/th [Therapy], 1,155
- 5, *cardiogenic shock/th [Therapy], 2,198
- 6, stunned heart muscle/th [Therapy], 53
- 7, myocarditis/th [Therapy], 874
- 8, *heart infarction/th [Therapy], 9,266
- 9, "out of hospital cardiac arrest"/th [Therapy], 3,990
- 10, ((acute* or severe* or refract*) adj (heart* or cardiac* or myocard* or cardio* or ventric*) adj (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock* or arrest* or stunn*)).tw., 32,685
- 11, (cardiogen* adj shock).tw., 33,626
- 12, Myocardit*.tw., 33,287
- 13, ((Postpartum* or post-partum* or peripartum* or peri-partum*) adj cardiomyopath*).tw., 3,113
- 14, (postcardiotomy or Post-cardiotomy).tw., 2,079
- 15, PPCM.tw., 1,382
- 16, (myocard* adj (stun* or hibernat* or infarct*)).tw., 360,780
- 17, primary graft dysfunction/th [Therapy], 94
- 18, (primary* adj graft* adj dysfunct*).tw., 3,390
- 19, or/3-18, 448,325
- 20, *resuscitation/, 62,739
- 21, *extracorporeal oxygenation/, 18,275

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- 22, ECMO.tw., 32,585
- 23, *extracorporeal circulation/, 9,325
- 24, (extracorp* adj circulat*).tw., 10,199
- 25, (extracorp* adj ((cardiopulmon* adj resuscitat*) or CPR)).tw., 2,110
- 26, ECPR.tw., 2,050
- 27, (Biomedicus adj pump*).tw., 50
- 28, (Maquet* adj rotaflow*).tw., 33
- 29, (jostra adj (pump* or rotaflow*)).tw., 17
- 30, (levitronix adj (centrimag* or pump* or system* or oxygen*)).tw., 154
- 31, (Medos adj (Hilite* or oxygen*)).tw., 46
- 32, (left adj ventricle adj assist adj device).tw., 230
- 33, or/20-32, 115,373
- 34, 19 and 33, 15,247
- 35, Nonhuman/ not Human/, 5,720,207
- 36, 34 not 35, 14,988
- 37, (conference abstract* or conference review or conference paper or conference proceeding).db,pt,su., 6,294,375
- 38, 36 not 37, 6,986
- 39, (exp child/ or exp pediatrics/ or exp adolescent/) not exp adult/, 2,766,023
- 40, 38 not 39, 6,431
- 41, limit 40 to english language, 5,795
- 42, limit 41 to dd=20130901-20250630, 4,668
- 43, limit 41 to dc=20130901-20250630, 4,655
- 44, 42 or 43, 4,669
- 45, Clinical trial.pt., 533,511

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

46, 44 not 45, 4,461

Cochrane Library (CDSR) search strategy

#1 MeSH descriptor: [Heart Failure] this term only and with qualifier(s): [therapy - TH] 2567

#2 MeSH descriptor: [Acute Disease] this term only and with qualifier(s): [therapy - TH] 115

#3 #1 and #2 0

#4 MeSH descriptor: [Cardiomyopathies] this term only and with qualifier(s): [therapy - TH] 86

#5 MeSH descriptor: [Shock, Cardiogenic] this term only and with qualifier(s): [therapy - TH] 194

#6 MeSH descriptor: [Myocardial Stunning] this term only and with qualifier(s): [therapy - TH] 3

#7 MeSH descriptor: [Myocarditis] this term only and with qualifier(s): [therapy - TH] 13

#8 MeSH descriptor: [Myocardial Infarction] this term only and with qualifier(s): [therapy - TH] 2729

#9 MeSH descriptor: [Out-of-Hospital Cardiac Arrest] this term only and with qualifier(s): [therapy - TH] 568

#10 ((acute* or severe* or refract*) next (heart* or cardiac* or myocard* or cardio* or ventric*) next (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock* or arrest* or stunn*)) 2872

#11 (cardiogen* next shock) 1767

#12 Myocardit* 1473

#13 ((Postpartum* or post-partum* or peripartum* or peri-partum*) next cardiomyopath*) 50

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

#14 (postcardiotomy or Post-cardiotomy) 48

#15 PPCM 41

#16 (myocard* next (stun* or hibernat* or infarct*)) 39538

#17 MeSH descriptor: [Primary Graft Dysfunction] this term only and with
qualifier(s): [therapy - TH] 3

#18 (primary* next graft* next dysfunct*) 150

#19 #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or
#15 or #16 or #17 or #18 44787

#20 MeSH descriptor: [Cardiopulmonary Resuscitation] this term only and with
qualifier(s): [methods - MT] 761

#21 MeSH descriptor: [Extracorporeal Membrane Oxygenation] this term only
and with qualifier(s): [methods - MT] 103

#22 ECMO 1170

#23 MeSH descriptor: [Extracorporeal Circulation] this term only and with
qualifier(s): [methods - MT] 123

#24 (extracorp* next circulat*) 1423

#25 (extracorp* next ((cardiopulmon* next resuscitat*) or CPR)) 80

#26 ECPR 127

#27 (Biomedicus next pump*) 2

#28 (Maquet* next rotaflow*) 2

#29 (jostra next (pump* or rotaflow*)) 0

#30 (levitronix next (centrimag* or pump* or system* or oxygen*)) 0

#31 (Medos next (Hilite* or oxygen*)) 0

#32 (left next ventricle next assist next device) 1

#33 #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or
#31 or #32 3455

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with
refractory cardiac arrest

#34 #19 AND #33 with Cochrane Library publication date Between Sep 2013 and Jun 2025, in Cochrane Reviews 30

#35 conference:pt or (clinicaltrials or trialsearch or clinicaltrials.gov or

www.who.int) 861611

#36 #34 not #35 0

Cochrane Central (CDSR) search strategy

#1 MeSH descriptor: [Heart Failure] this term only and with qualifier(s): [therapy - TH] 2567

#2 MeSH descriptor: [Acute Disease] this term only and with qualifier(s): [therapy - TH] 115

#3 #1 and #2 0

#4 MeSH descriptor: [Cardiomyopathies] this term only and with qualifier(s): [therapy - TH] 86

#5 MeSH descriptor: [Shock, Cardiogenic] this term only and with qualifier(s): [therapy - TH] 194

#6 MeSH descriptor: [Myocardial Stunning] this term only and with qualifier(s): [therapy - TH] 3

#7 MeSH descriptor: [Myocarditis] this term only and with qualifier(s): [therapy - TH] 13

#8 MeSH descriptor: [Myocardial Infarction] this term only and with qualifier(s): [therapy - TH] 2729

#9 MeSH descriptor: [Out-of-Hospital Cardiac Arrest] this term only and with qualifier(s): [therapy - TH] 568

#10 ((acute* or severe or *refract*) next (heart* or cardiac* or myocard* or cardio* or ventric*) next (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock* or arrest* or stunn*)) 2872

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

#11 (cardiogen* next shock) 1767

#12 Myocardit* 1473

#13 ((Postpartum* or post-partum* or peripartum* or peri-partum*) next cardiomyopath*) 50

#14 (postcardiotomy or Post-cardiotomy) 48

#15 PPCM 41

#16 (myocard* next (stun* or hibernat* or infarct*)) 39538

#17 MeSH descriptor: [Primary Graft Dysfunction] this term only and with qualifier(s): [therapy - TH] 3

#18 (primary* next graft* next dysfunct*) 150

#19 #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 44787

#20 MeSH descriptor: [Cardiopulmonary Resuscitation] this term only and with qualifier(s): [methods - MT] 761

#21 MeSH descriptor: [Extracorporeal Membrane Oxygenation] this term only and with qualifier(s): [methods - MT] 103

#22 ECMO 1170 #23 MeSH descriptor: [Extracorporeal Circulation] this term only and with qualifier(s): [methods - MT] 123

#24 (extracorp* next circulat*) 1423

#25 (extracorp* next ((cardiopulmon* next resuscitat*) or CPR)) 80

#26 ECPR 127

#27 (Biomedicus next pump*) 2

#28 (Maquet* next rotaflow*) 2

#29 (jostra next (pump* or rotaflow*)) 0

#30 (levitronix next (centrimag* or pump* or system* or oxygen*)) 0

#31 (Medos next (Hilite* or oxygen*)) 0

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

#32 (left next ventricle next assist next device) 1

#33 #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 3455

#34 #19 AND #33 with Cochrane Library publication date Between Sep 2013 and Jun 2025, in Trials 499

#35 conference:pt or (clinicaltrials or trialsearch or clinicaltrials.gov or www.who.int) 861611 #36 #34 not #35 295

INHTA HTA Database search strategy

1 , "Heart Failure"[mh] , 271

2 , "Acute Disease"[mh] , 44

3 , #2 AND #1 , 2

4 , "Cardiomyopathies"[mh] , 25

5 , "Shock, Cardiogenic"[mh] , 11

6 , "Myocardial Stunning"[mh] , 1

7 , "Myocarditis"[mh] , 2

8 , "Myocardial Infarction"[mh] , 122

9 , "Out-of-Hospital Cardiac Arrest"[mh] , 11

10 , ((acute* or severe* or refract*) AND (heart* or cardiac* or myocard* or cardio* or ventric*) AND (failur* or decompensation* or insufficient* or dysfunct* or stand* or still* or fault* or shock* or arrest* or stunn*)) , 268

11 , (cardiogen* AND shock) , 19

12 , Myocardit* , 5

13 , ((Postpartum* or post-partum* or peripartum* or peri-partum*) AND cardiomyopath*) , 1

14 , (postcardiotomy or Post-cardiotomy) , 0

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

- 15 , PPCM , 0
- 16 , (myocard* AND (stun* or hibernat* or infarct*)) , 236
- 17 , "Primary Graft Dysfunction"[mh] , 0
- 18 , (primary* AND graft* AND dysfunct*) , 3
- 19 , #18 OR #17 OR #16 OR #15 OR #14 OR #13 OR #12 OR #11 OR #10 OR #9 OR #8 OR #7 OR #6 OR #5 OR #4 OR #3 , 523
- 20 , "Cardiopulmonary Resuscitation"[mh] , 23
- 21 , "Extracorporeal Membrane Oxygenation"[mh] , 28
- 22 , ECMO , 30
- 23 , "Extracorporeal Circulation"[mh] , 8
- 24 , (extracorp* AND circulat*) , 13
- 25 , (extracorp* AND ((cardiopulmon* AND resuscitat*) or CPR)): , 8
- 26 , ECPR , 4
- 27 , (Biomedicus AND pump*) , 0
- 28 , (Maquet* AND rotaflow*) , 0
- 29 , (jostra AND (pump* or rotaflow*)) , 0
- 30 , (levitronix AND (centrimag* or pump* or system* or oxygen*)) , 0
- 31 , (Medos AND (Hilite* or oxygen*)) , 0
- 32 , (Left AND ventricle AND assist AND device) , 3
- 33 , #32 OR #31 OR #30 OR #29 OR #28 OR #27 OR #26 OR #25 OR #24 OR #23 OR #22 OR #21 OR #20 , 72
- 34 , #33 AND #19 , 29

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Inclusion criteria

The following inclusion criteria were applied to the abstracts identified by the literature search.

Publication type: clinical studies were included with emphasis on identifying good quality studies. Abstracts were excluded if they did not report clinical outcomes. Reviews, editorials, and laboratory or animal studies, were also excluded and so were conference abstracts, because of the difficulty of appraising study methodology, unless they reported specific adverse events not available in the published literature.

Population: adults with refractory cardiac arrest.

Intervention or test: VA ECMO for ECPR.

Outcome: articles were retrieved if the abstract contained information relevant to the safety, efficacy, or both.

If selection criteria could not be determined from the abstracts the full paper was retrieved.

Potentially relevant studies not included in the main evidence summary are listed in [Appendix B: Other relevant studies](#).

Find out more about [how NICE selects the evidence for the committee](#).

Appendix B: Other relevant studies

Other potentially relevant studies that were not included in the main evidence summary ([tables 2 and 3](#)) are listed in table 5 below.

For the original search results identified from March 2013 to September 2024, case studies and observational studies with fewer than 100 people were excluded unless they included outcomes that were not frequently reported.

For the updated search results identified from September 2024 to June 2025, systematic reviews published before 2020 were excluded.

Table 5 additional studies identified

Study	Number of people and follow up	Direction of conclusions	Reason study was not included in main evidence summary
Alenazi A, Aljanoubi M, Yeung J et al. (2024) Variability in patient selection criteria across extracorporeal cardiopulmonary resuscitation (ECPR) systems: A systematic review. Resuscitation 204: 110403	Systematic review 90 studies (93 ECPR systems)	Across systems, ECPR eligibility criteria included age (n=75, 81%), arrest witnessed status (n=64, 69%), any initial rhythm (n=55, 59%), and bystander CPR (n=33, 36%). Within criteria, we observed marked variability. The age cut-off varied from 50 to 80 years, with the most common age cut-off being 75 years and 18 (19%) systems having no cut-off. Whilst most systems limited ECPR to shockable rhythms (n=28, 30%), some systems included	Review focuses on variability in patient selection criteria.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		shockable rhythms or pulseless electrical activity only (n=10, 11%).	
Alexy T, Kalra R, Kosmopoulos M et al. (2023) Initial hospital length of stay and long-term survival of patients successfully resuscitated using extracorporeal cardiopulmonary resuscitation for refractory out-of-hospital cardiac arrest. European Heart Journal. Acute Cardiovascular Care 12(3): 175-183	Single centre retrospective study, US n=160 Follow-up: 4 years (median follow-up: 3 years)	34% people survived the index admission. These survivors required a median 16 days of intensive care and 24 days total hospital stay. Of these, 80% and 72% were alive at 1 and 4 years, respectively. Most deaths within the first year occurred among the people requiring discharge to a long-term acute care facility.	Larger, more comprehensive systematic literature reviews and meta-analysis included. Longer-term data included from RCT.
Alhuneafat L, Ghanem F, Brankovic M et al. (2025) Predictors of extracorporeal membrane oxygenation utilization and survival during cardiopulmonary resuscitation in out and in-hospital cardiac arrest. Cardiovascular Revascularization Medicine https://doi.org/10.1016/j.carrev.2025.05.008	Retrospective cohort study n=1,585,960	ECPR rates were 1% for OHCA and 1.4% for IHCA, with inpatient mortality rates of 52% and 67%, respectively. In OHCA ECPR, Asian race (aOR 2.31), diabetes (aOR: 1.29), and liver disease (aOR: 1.77) predicted mortality, while shockable rhythms (aOR: 0.75), systolic heart failure (aOR: 0.67), and treatment in southern states (aOR: 0.72) predicted survival. In IHCA ECPR, acute myocardial infarction (aOR: 0.73) and private insurance (aOR: 0.63) were associated with improved survival, whereas liver disease (aOR: 1.59) predicted higher mortality.	Study reports limited outcomes for ECPR.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Belohlavek J, Smalcova J, Rob D et al. (2022) Effect of intra-arrest transport, extracorporeal cardiopulmonary resuscitation, and immediate invasive assessment and treatment on functional neurologic outcome in refractory out-of-hospital cardiac arrest: a randomized clinical trial. JAMA 327(8): 737-747	Prague OHCA randomised controlled trial n=256 (124 ECPR) Follow-up: 180 days (6 months)	In the main analysis, 32% of the ECPR group and 22% of the CCPR group survived to 180 days with good neurologic outcome (OR, 1.63, 95% CI 0.93 to 2.85; p=0.09). At 30 days, neurologic recovery had occurred in 31% in the ECPR group and in 18% in the CCPR group (OR, 1.99, 95% CI 1.11 to 3.57; p=0.02). Bleeding occurred more frequently in the ECPR versus CCPR group (31% vs 15%, respectively).	RCT included in all SLRs
Belohlavek J, Yannopoulos D, Smalcova J et al. (2023) Intraarrest transport, extracorporeal cardiopulmonary resuscitation, and early invasive management in refractory out-of-hospital cardiac arrest: an individual patient data pooled analysis of two randomised trials. EClinicalMedicine 59: 101988	Pooled individual patient data analysis of 2 randomised controlled trials (ARREST and PRAGUE-OHCA) n=286 Follow-up=30 days	In people with refractory OHCA, ECPR statistically significantly improved 30- and 180-day neurologically favourable survival.	Trials are already included in the systematic reviews in the key evidence.
Beyea MM, Tillmann BW, Iansavichene AE et al. (2018) Neurologic outcomes after extracorporeal membrane oxygenation assisted CPR for resuscitation of out-of-hospital cardiac arrest	Systematic review and meta-analysis n=NR, 75 studies (case series and cohort studies) Follow-up: hospital discharge	Among case series, 0 to 71% of people who had ECPR survived to discharge with a good neurological outcome. Subgroup analysis of cohort studies demonstrated survival-to-hospital discharge with good neurological	More recent systematic literature reviews with meta-analysis included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

patients: A systematic review. Resuscitation 130: 146-158		recovery in the ECPR group ranged from 8 to 42% compared with 2 to 9% in the CCPR group.	
Bougouin W, Dumas F, Lamhaut L et al. (2020) Extracorporeal cardiopulmonary resuscitation in out-of-hospital cardiac arrest: a registry study. European Heart Journal 41(21): 1961-1971	Retrospective registry study (France). n=13,191 Follow-up: hospital discharge	Survival was 8% in ECPR group and 9% in CCPR group (p=0.91). By adjusted multivariate analysis, ECPR was not associated with hospital survival (OR 1.3, 95% CI 0.8 to 2.1; p=0.24). PSM analysis found similar results (OR 0.8, 95% CI 0.5 to 1.3; p=0.41). In the ECPR group, factors associated with hospital survival were initial shockable rhythm (p=0.005), transient ROSC before ECMO (p=0.03), and prehospital ECMO implantation (p=0.002)	Included in SLRs (Pagura 2024)
Bourcier S, Desnos C, Clement M et al. (2022) Extracorporeal cardiopulmonary resuscitation for refractory in-hospital cardiac arrest: A retrospective cohort study. International Journal of Cardiology 350: 48-54	Retrospective cohort study n=137	1-year survival was 19% and most survivors had favourable neurological outcome. An algorithm based on age, initial rhythm (shockable or not) and low-flow duration identified those with the highest and lowest probability of survival.	Larger studies are included.
Bunya N, Ohnishi H, Kasai T et al. (2025) Signs of life as a favorable predictor for non-shockable cardiac arrest undergoing extracorporeal cardiopulmonary resuscitation post	Retrospective single-centre observational study n=227	Initial cardiac rhythms were shockable in 154 people (68%) and non-shockable in 73 (32%). Favourable neurological outcomes were observed in 4%, 48%, 0%, and 39% of people with shockable rhythm without signs of life, shockable rhythm with signs of life,	Larger studies are included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

non-shockable cardiac arrest. The American Journal of Emergency Medicine 87: 95–104		non-shockable rhythm without signs of life, and non-shockable rhythm with signs of life, respectively, with statistically significant differences. Multivariate logistic regression analysis showed an independent association of signs of life during resuscitation with favourable neurological outcomes. The adjusted odds ratios for people with shockable and non-shockable rhythm with signs of life were 34.33 and 96.51, respectively, compared with those without signs of life	
Carlson JM, Etchill E, Whitman G et al. (2022) Early withdrawal of life sustaining therapy in extracorporeal cardiopulmonary resuscitation (ECPR): Results from the Extracorporeal Life Support Organization registry. Resuscitation 179: 71-77	Registry (ELSO) n=411	More than half of people who had ECPR had early withdrawal of life sustaining therapy within 72 hours. The people with early withdrawal of life sustaining therapy had worse markers of severe critical illness at 24 hours and higher rates of complications.	Study focuses on early withdrawal of life sustaining therapy.
Chahine J, Kosmopoulos M, Raveendran G et al. (2023) Impact of age on survival for patients receiving ECPR for refractory out-of-hospital VT/VF cardiac arrest. Resuscitation 193: 109998	Single centre retrospective study, US n=391 Follow-up: hospital discharge	Age was independently associated with neurologically favourable survival to discharge, with a 30% decrease in survival with every 10-year increase in age (OR 0.7, 95% CI 0.57 to 0.87, p=0.001).	Larger, more comprehensive systematic literature reviews and meta-analysis included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Chen Z, Liu C, Huang J et al. (2019) Clinical efficacy of extracorporeal cardiopulmonary resuscitation for adults with cardiac arrest: meta-analysis with trial sequential analysis. BioMed Research International 2019: 6414673	Systematic review and meta-analysis n=NR, 13 observational studies Follow-up: 1 year	ECPR in OHCA and IHCA was associated with a significantly better 30-day survival (RR 1.60, 95% CI 1.25 to 2.06) and 30-day neurologic outcome (RR 2.69, 95% CI 1.63 to 4.46) than CCPR. Relative to CCPR, ECPR improved the survival and neurological outcome of people who had IHCA. Trial sequential analysis could not confirm better survival and neurological outcome of ECPR in people with OHCA, suggesting that further studies are needed.	More recent systematic literature reviews included.
Choi DS, Kim TR, Young S et al. (2016) Extracorporeal life support and survival after out-of-hospital cardiac arrest in a nationwide registry: A propensity score-matched analysis. Resuscitation 99: 26-32	Retrospective registry study (Korea). n=36,547 (320 ECPR) Follow-up: hospital discharge	There was no significant difference in neurologically favourable survival to discharge between the ECLS group and the non-ECLS group after adjusting for covariates (adjusted OR 0.65, 95% CI 0.41 to 1.04). In the PSM cohort, there was also no significant difference between the 2 groups (adjusted OR, 0.94, 95% CI 0.41 to 2.14).	Included in SLRs (Zhong 2024, Pagura 2024)
Choi Y, Park JH, Jeong J et al. (2023) Extracorporeal cardiopulmonary resuscitation for adult out-of-hospital cardiac arrest patients: time-dependent propensity score-sequential matching analysis from a nationwide	PSM retrospective registry study (Korea). n=2,290 (458 ECPR) Follow-up: hospital discharge	ECPR itself was not associated with good neurological recovery (10% in ECPR and 7% in no ECPR; RR 1.28, 95% CI 0.85 to 1.93), but early ECPR was positively associated with good neurological recovery.	Included in SLRs (Zhong 2024, Pagura 2024)

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

population-based registry. Critical Care 27(1): 87			
Crespo-Diaz R, Kosmopoulos M, Raveendran G et al. (2024) Effects of Perfusion, Coronary Artery Disease Burden, and Revascularization in Establishing Organized Cardiac Rhythm During Extracorporeal Cardiopulmonary Resuscitation for Shockable Refractory Out-of-Hospital Cardiac Arrest. Journal of the American Heart Association 13: e033907	Retrospective cohort study n=289	<p>Standard advanced cardiac life support before hospital arrival resulted in 148 of 289 (51%) people attaining an organised rhythm while 87 of 289 (30%) achieved an organised rhythm post ECMO cannulation but before percutaneous coronary intervention (PCI), and 37 of 289 (13%) achieved an organised rhythm following PCI.</p> <p>Most people who had ECPR for refractory OHCA due to shockable rhythms achieved an organised rhythm before PCI independent of coronary artery disease burden. Also, neurologically favourable survival was more prevalent in those attaining an organised rhythm before PCI.</p>	Larger studies are included.
Daou O, Winiszewski H, Besch G et al. (2020) Initial pH and shockable rhythm are associated with favorable neurological outcome in cardiac arrest patients resuscitated with extracorporeal cardiopulmonary resuscitation. Journal of Thoracic Disease 12: 849	Retrospective single-centre cohort study n=113	Overall low-flow time was 84 [55 to 122] minutes. Eighteen people (16%) survived with a favourable neurological outcome. By multivariate logistic regression analysis, initial shockable rhythm, and arterial blood pH at the time of ECPR implantation 7.0 or higher, were independent predictors of survival with favourable neurological	Larger studies are included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		outcome. All those presenting with both non-shockable rhythm and pH below 7.0 at the time of ECPR implantation died in the intensive care unit.	
Downing J, Al F, Reem CS et al. (2022) How effective is extracorporeal cardiopulmonary resuscitation (ECPR) for out-of-hospital cardiac arrest? A systematic review and meta-analysis. The American journal of emergency medicine 51: 127-138	Systematic review and meta-analysis n=3,097, 44 studies Follow-up: 90 days	ECPR for OHCA showed survival-to-discharge rate of 24%; 18% survived with favourable neurologic function. 30- and 90-days survival rates were both around 18%.	Larger, more recent and comparative systematic literature reviews and meta-analyses included.
Haas NL, Coute RA, Hsu CH et al. (2017) Descriptive analysis of extracorporeal cardiopulmonary resuscitation following out-of-hospital cardiac arrest-An ELSO registry study. Resuscitation 119: 56-62	Retrospective ELSO registry study n=217 Follow-up: hospital discharge	Reported complications included haemorrhage (31%), limb complications (11%), circuit complications (9%), infection (7%), and seizures (6%). Survival to hospital discharge was 28% (95% CI 22.1 to 34.0%), and male gender was independently associated with mortality (aOR 2.1 (95% CI 1.1 to 4.2, p<0.05). Survival did not differ by region, race, age, or year.	Larger, more recent and comparative systematic literature reviews and meta-analyses included.
Hashem A, Mohamed MS, Alabdullah K, et al. (2023). Predictors of mortality in patients with refractory cardiac arrest supported with VA-ECMO: a systematic review and a meta-analysis. Current Problems in	Systematic review and meta-analysis n=931, 10 studies Follow-up: 90 days	The overall mortality was 69%. The predictors for mortality were age over 65 (OR 4.61, 95% CI 1.63 to 13.03, p<0.01), history of chronic kidney disease (OR 2.42, 95% CI 1.37 to 4.28, p<0.01), cardiopulmonary resuscitation duration prior to ECMO more than	Larger, comparative systematic literature reviews and meta-analyses included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Cardiology, 48(6), 101658.		40 minutes (OR 6.62, 95% CI 1.39 to 9.02, $p<0.01$), having an initial non-shockable rhythm (OR 2.62, 95% CI 1.85 to 3.70, $p<0.01$) and sequential organ failure assessment score higher than 14 (OR 12.29, 95% CI 2.71 to 55.74, $p<0.01$).	
Havranek S, Fingrova Z, Rob D et al. (2022) Initial rhythm and survival in refractory out-of-hospital cardiac arrest. Post-hoc analysis of the Prague OHCA randomized trial. Resuscitation 181: 289-296	Post-hoc analysis of the Prague OHCA randomised controlled trial n=256 Follow-up: 180 days (6 months)	Favourable neurological survival at 180 days was achieved in 40% of people with a shockable rhythm and in 5% of people with a non-shockable rhythm ($p<0.001$). The difference between shockable and non-shockable initial rhythms remained statistically significant (35/72 [49%] versus 4/52 [8%] in the ECPR group and 28/84 [33%] versus 1/48 [2%] in the CCPR group; $p<0.001$).	Original RCT include in all key evidence systematic reviews.
Heuts S, Ubben JFH, Kawczynski MJ et al. (2024) Extracorporeal cardiopulmonary resuscitation versus standard treatment for refractory out-of-hospital cardiac arrest: a Bayesian meta-analysis. Critical Care 28(1): 217	Bayesian meta-analysis of ARREST, Prague OHCA and INCEPTION RCTs n=420 Follow-up: 6 months	The Bayesian meta-analysis found a 71% and 76% posterior probability of a clinically relevant ECPR-based treatment effect on 6-month neurologically favourable survival in people with all rhythms and shockable rhythms.	Original RCTs included in all key evidence systematic reviews.
Heuts S, van de Koolwijk AF, Gabrio A et al. (2024) Extracorporeal life support in cardiac arrest: a post hoc Bayesian re-analysis	Bayesian meta-analysis of the INCEPTION RCT n=134 Follow-up: 30 days	Bayesian re-analysis of the INCEPTION trial estimated a 42% probability of an MCID between ECPR and CCPR in refractory OHCA in terms of 30-day	Original RCT included in all key evidence systematic reviews.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

of the INCEPTION trial. European heart journal. Acute Cardiovascular Care 13(2): 191-200		survival with a favourable neurologic outcome.	
Holmberg MJ, Geri G, Wiberg, S et al. (2018) Extracorporeal cardiopulmonary resuscitation for cardiac arrest: A systematic review. Resuscitation 131: 91-100	Systematic review n=25 studies Follow-up: long-term (unclear)	There is inconclusive evidence to either support or refute the use of ECPR for OHCA and IHCA in adults and children.	Larger, more recent comparative systematic literature reviews and meta-analyses included.
Holmberg MJ, Granfeldt A, Guerguerian AM et al. (2023) Extracorporeal cardiopulmonary resuscitation for cardiac arrest: An updated systematic review. Resuscitation 182: 109665	Updated systematic review n=3 RCTs and 27 observational studies Follow-up: in-hospital	Results of individual studies were inconsistent, although many studies favoured ECPR. The risk of bias was intermediate for trials and critical for observational studies. The certainty of evidence was very low to low. Study heterogeneity precluded meta-analyses.	Larger, systematic literature reviews and meta-analyses were included.
Hongo T, Naito H, Nasu M et al. (2024) Prognostic performance of gray-white matter ratio in adult out-of-hospital cardiac arrest patients after receiving extracorporeal cardiopulmonary resuscitation. Resuscitation 203: 110351	Registry (SAVE-J II) n=1,146	Overall, people with lower average grey-to-white matter ratio (aGWR) more likely had poor neurological outcomes, aGWR 1.00 to 1.09 (95%), aGWR 1.10 to 1.19 (88%), aGWR 1.20 to 1.29 (79%), and aGWR 1.30 to 1.39 (70%). Multivariable logistic regression showed that lower aGWR was associated with poor neurological outcome at 30-day, aGWR 1.30 to 1.39: reference, aGWR 1.00 to 1.09: adjusted OR (aOR) 10.01 (95% CI 3.58 to 27.99), aGWR	Subanalysis of study that is already included in table 2 (Inoue, 2022).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		1.10 to 1.19: aOR 4.83 (95% CI 2.31 to 10.12), aGWR 1.20 to 1.29: aOR 2.16 (95% CI 1.02 to 4.55).	
Ijuin S, Inoue A, Hifumi T et al. (2025) Analysis of factors associated with favorable neurological outcomes in patients with initial PEA who underwent ECPR-A secondary analysis of the SAVE-J II study. Journal of Critical Care 85: 154917	Registry (SAVE-J II) n=473	Overall proportion of people with favourable neurological outcomes and survival rate at hospital discharge were 8% and 17%, respectively. Among the cause of cardiac arrest, people with acute coronary syndrome and pulmonary embolism had higher proportions of favourable neurological outcomes (10% and 19%), whereas no people with acute aortic disease or primary cerebral disease survived. The application of strict criteria for PEA using classification and regression tree analysis resulted in favourable neurological outcomes in 33% of the people.	Subanalysis of study that is already included in table 2 (Inoue, 2022).
Inoue F, Otani T, Abe T et al. (2025) Characteristics of Intracranial Hemorrhage and Acute Aortic Syndromes Resuscitated with Extracorporeal Cardiopulmonary Resuscitation (ECPR). The Journal of Emergency Medicine 69: 1–12	Secondary analysis of SAVE-J II study n=1,589	People in the intracranial haemorrhage (ICH) group were younger (ICH: 47 years [IQR 40 to 63], acute aortic syndromes [AAS]: 68 years [IQR 59 to 74], cardiac causes: 61 years [IQR 50 to 68]; $p<0.001$), and had fewer signs of life on hospital arrival (ICH: 2%, AAS: 12%, cardiac causes: 19%; $p=0.001$) compared with the cardiac causes group. People in the ICH and AAS groups were	Subanalysis of study that is already included in table 2 (Inoue, 2022).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		also less likely to present with an initial shockable rhythm (ICH, 19%; AAS, 27%; cardiac causes, 74%; $p<0.001$). Survival to hospital discharge was significantly lower in the ICH and AAS groups than in the cardiac cause group (ICH, 4%; AAS, 4%; cardiac causes, 29%; $p<0.001$).	
Kawahara N, Takayama W, Morishita K et al. (2025) Effects of prehospital advanced airway management on cardiac arrest patients who underwent extracorporeal cardiopulmonary resuscitation. Scientific Reports 15: 10670	Registry (SAVE-J II) n=1,789	Those who had prehospital advanced airway management had significantly lower favourable neurological outcomes at discharge, lower 30-day survival, longer low-flow time, shorter mechanical ventilation days, and shorter intensive care unit stay days than those who did not have prehospital advanced airway management. Propensity score-matched analysis confirmed these findings.	Subanalysis of study that is already included in table 2 (Inoue, 2022).
Kawauchi A, Okada Y, Aoki M et al. (2024) Sex differences in extracorporeal cardiopulmonary resuscitation for out-of-hospital cardiac arrest: nationwide multicenter retrospective study in Japan. Critical Care 28: 302	Registry (SAVE-J II) n=1,819	The proportion of favourable neurological outcomes was 12% in males and 16% in females ($p=0.10$). Multilevel logistic regression analysis showed that the female sex was significantly associated with a favourable neurological outcome at discharge (adjusted OR: 1.60 [95% CI: 1.05 to 2.43]; $p=0.03$). This advantage in women	Subanalysis of study that is already included in table 2 (Inoue, 2022).

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		was consistently observed in the sensitivity analyses.	
Khoury J, Soumagnac T, Vimpere D et al. (2025) Long-term heart function in refractory out-of-hospital cardiac arrest treated with prehospital extracorporeal cardiopulmonary resuscitation. Resuscitation 207: 110449	Retrospective single centre observational study n=114 Follow-up: median 6.8 years	24/114 (21%) survived at 1 year with good neurological outcomes. At the time of data collection, 21 people were still alive, with a median follow-up time of 6.8 years. Half of these people were actively working, with a median time of 10 months to regain the ability to work since the onset of OHCA.	Larger studies are included.
Kikutani K, Nishikimi M, Ohshimo S et al. (2025) Association between increasing institutional experience with ECPR and outcomes in patients with out-of-hospital cardiac arrest: A nationwide multicenter observational study in Japan (the JAAM-OHCA registry). Resuscitation 209: 110487	Registry (JAAM-OHCA) n=2,315	Increasing institutional experience of ECPR did not significantly improve 30-day survival after OHCA but was associated with a shorter interval between hospital arrival and initiation of ECMO. In people with non-shockable OHCA, increasing experience of ECPR improved 30-day survival.	Data from registry is included in systematic review by Low (2024). This study focuses on the association between increasing experience with ECPR and outcomes.
Kim SJ, Kim HJL, Hee Y et al. (2016) Comparing extracorporeal cardiopulmonary resuscitation with conventional cardiopulmonary resuscitation: A meta-analysis. Resuscitation 103: 106-116	Systematic review and meta-analysis n=10 studies Follow-up: 1 year	Survival and good neurological outcome tended to be superior in the ECPR group at 3 to 6 months after arrest. The effect of ECPR on survival to discharge in OHCA was not clearly shown.	Larger, more recent systematic literature reviews and meta-analyses included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Kosmopoulos M, Bartos JA, Kalra R et al. (2021) Patients treated with venoarterial extracorporeal membrane oxygenation have different baseline risk and outcomes dependent on indication and route of cannulation. Hellenic Journal of Cardiology 62: 38-45	Retrospective single centre cohort study n=317	Peripherally cannulated people with ECPR had significantly worse Acute Physiology and Chronic Health Evaluation 2 and Survival After Venoarterial ECMO scores compared with peripherally cannulated refractory cardiogenic shock or people with central ECMO, despite having similar mortality.	Larger studies are included.
Koukousaki D, Kosmopoulos M, Mallow J et al. (2024) Temporal trends in organ donation among cardiac arrest patients treated with extracorporeal cardiopulmonary resuscitation. Resuscitation 203: 110391	Retrospective, single centre observational study n=419	Out of 419 people who had ECPR presenting with refractory OHCA over the study period, 116 survived neurologically intact (28%). Among non-survivors (n=303), families of 41 (14%) consented to organ donation (median age 51 years, 76% male) and organs from 38 people were retrieved, leading to 74 organ transplants to 73 recipients.	Study focuses on organ donation after ECPR.
Kreling GAD, Mendes PV, Cardozo Junior LCM et al. (2025) Feasibility of implementing extracorporeal cardiopulmonary resuscitation in a middle-income country: systematic review and cardiac arrest case series. Critical Care Science 37: e20250320	Systematic review and meta-analysis 31 studies	The main and sensitivity analyses consistently demonstrated that ECPR was associated with favourable neurological outcomes (cerebral performance category 1 or 2, RR 1.45, 95% CI 1.19 to 1.77) and survival (RR 1.29, 95% CI 1.10 to 1.52). Age was inversely related to neurological outcome and survival.	There is a large amount of overlap with systematic reviews already included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Kruit N, Rattan N, Tian D et al. (2023) Prehospital extracorporeal cardiopulmonary resuscitation for out-of-hospital cardiac arrest: a systematic review and meta-analysis. Journal of Cardiothoracic and Vascular Anesthesia 37(5): 748-754	Systematic review and meta-analysis n=222 ECPR, 4 studies Follow-up: hospital discharge	Overall survival at discharge was 23% (95% CI 15.5 to 33.7; $I^2=62\%$). The quality of evidence was assessed to be low, and the overall risk of bias was assessed to be serious, with confounding being the primary source of bias.	Larger, more recent systematic literature reviews and meta-analyses included.
Lee SW, Han KS, Park JS et al. (2017) Prognostic indicators of survival and survival prediction model following extracorporeal cardiopulmonary resuscitation in patients with sudden refractory cardiac arrest. Annals of Intensive Care 7: 87	Retrospective analysis of prospective cohort n=111	Survival rate=19% Survivors showed younger age, shorter CPR duration ($p<0.05$) and had tendencies of higher rate of initial shockable rhythm ($p=0.055$) and higher rate of any return of spontaneous circulation event before ECPR ($p=0.066$) than non-survivors. 81% of survivors showed favourable neurological outcome at discharge. Based on multivariate logistic regression, predictors for survival in ECPR were: age 56 years or younger, no asystole as the initial arrest rhythm, CPR duration of 55 minutes or less, and any return of spontaneous circulation event before ECPR.	Larger studies are included.
Lunz D, Calabro L, Belliato M et al. (2020) Extracorporeal membrane oxygenation for refractory cardiac	Multicentre retrospective study, US n=423 Follow-up: 3 months	Eighty people (19%) had favourable neurological outcome. ICU survival was 24%. Favourable neurological outcome rate was lower (9%)	Larger, more recent systematic literature reviews and meta-

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

arrest: a retrospective multicenter study. Intensive care medicine 46(5): 973-982		versus 34%, $p<0.01$) in OHCA than IHCA and was significantly associated with shorter time from collapse to ECMO.	analyses included.
Magnet I, Behringer W, Eibensteiner F et al. (2025) Extracorporeal Cardiopulmonary Resuscitation: Outcomes Improve With Center Experience. Annals of Emergency Medicine 85: 421–27	Retrospective single centre observational study n=192	The proportion of people with favourable neurological outcomes was 25% (n=48), increasing each year: 15% (5/34) in 2020, 19% (8/42) in 2021, 23% (12/53) in 2022, and 37% (23/63) in 2023. This was particularly true for people with OHCA: 7% (2/29), 14% (4/29), 17% (7/41), and 32% (16/50), respectively. The observed learning curve illustrates that outcomes can improve with experience, a summation effect of training, patient selection, and process standardisation.	Larger studies are included.
Migdady I, Rice C, Deshpande A et al. (2020) Brain Injury and Neurologic Outcome in Patients Undergoing Extracorporeal Cardiopulmonary Resuscitation: A Systematic Review and Meta-Analysis. Critical Care Medicine 48: e611	Systematic review and meta-analysis n=50,049 (6,261 ECPR) 78 studies	The median ECPR duration was 3.2 days (interquartile range, 2.1 to 4.9). Overall, 27% (95% CI 0.17 to 0.39%) had at least 1 neurological complication, 23% (95% CI 0.14 to 0.32%) hypoxic-ischaemic brain injury, 6% (95% CI 0.02 to 0.11%) ischaemic stroke, 6% (95% CI, 0.01 to 0.16%) seizures, and 4% (95% CI 0.01 to 0.1%) intracerebral haemorrhage. 17% (95% CI 0.12 to 0.23%) developed brain death.	More recent systematic reviews are included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		The overall survival rate after ECPR was 29% (95% CI 0.26 to 0.33%) and good neurological outcome was achieved in 24% (95% CI 0.21 to 0.28%).	
Nanjayya VB, Fulcher B, Nehme E et al. (2025) Long-term health-related quality of life in survivors of extracorporeal cardiopulmonary resuscitation compared to conventional cardiopulmonary resuscitation- A cohort study using Australian and New Zealand extracorporeal membrane oxygenation registry and the Victorian Ambulance Cardiac Arrest Registry. Resuscitation 210: 110601	Retrospective cohort study n=9,112 (122 ECPR) Follow-up: 12 months	There were 33/122 (28%) ECPR and 1,074/8,990 (12%) CCPR OHCA survivors at 12 months. Of these, 24 (73%) ECPR and 754 (70%) CCPR survivors had HRQoL data. Despite statistically significant baseline differences between ECPR and CCPR, there were no differences in HRQoL at 12 months.	Data was only available for a small number of people who had ECPR.
Okada Y, Komukai S, Irisawa T et al. (2023) In-hospital extracorporeal cardiopulmonary resuscitation for patients with out-of-hospital cardiac arrest: an analysis by time-dependent propensity score matching using a nationwide database in Japan. Critical Care 27(1): 442	PSM retrospective JAAM-OHCA registry study (Japan) n=2,566 Follow-up: 30 days	The OR for 30-day survival in the ECPR group was 1.76 (95% CI 1.38 to 2.25) for shockable rhythm and 5.37 (95% CI 2.53 to 11.43) for non-shockable rhythm, compared with controls. For favourable neurological outcomes, the OR in the ECPR group was 1.11 (95% CI 0.82 to 1.49) for shockable rhythm and 4.25 (95% CI 1.43 to 12.63) for non-shockable	Included in SLRs (Low 2024)

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		rhythm, compared with controls.	
Patricio D, Peluso L, Brasseur A et al. (2019) Comparison of extracorporeal and conventional cardiopulmonary resuscitation: a retrospective propensity score matched study. Critical Care 23(1): 27	PSM retrospective study (Belgium) n=160 (80 ECPR) Follow-up: hospital discharge and 3 months	Survival to ICU discharge was 23% versus 18% in the ECPR and CCPR groups, respectively (p=0.42). At 3 months, 21% people with ECPR and 11% people with CCPR had a favourable outcome (p=0.11).	Included in SLRs (Low 2024, Low 2023, Zhong 2024)
Rob D, Farkasovska K, Kavalkova P et al. (2024) Heart rhythm at hospital admission: A factor for survival and neurological outcome among ECPR recipients? Resuscitation 204: 110412	Single-centre observational study n=196	The rhythm at hospital admission affects ECPR outcomes. People presenting with and maintaining ventricular fibrillation have a higher chance of favourable neurological survival, whereas those presenting with or converting to asystole have poor outcomes.	Larger studies are included.
Rob D, Komarek A, Smalcova J et al. (2024) Effect of intra-arrest transport, extracorporeal cardiopulmonary resuscitation, and invasive treatment: a post hoc bayesian reanalysis of a randomized clinical trial. Chest 165(2): 368-370	Bayesian meta-analysis of the Prague OHCA RCT n=256 Follow-up: 6 months	Bayesian reanalysis of the study primary outcome showed a benefit of the invasive approach compared with standard resuscitation under a broad set of scenarios.	Original RCT included in all key evidence systematic reviews.
Sakamoto T, Morimura N, Nagao K et al. (2014) Extracorporeal cardiopulmonary resuscitation versus conventional cardiopulmonary	Single centre prospective study (Japan) n=160 (80 ECPR) Follow-up: 6 months	In people with OHCA with VF/VT on the initial ECG, a treatment bundle including ECPR, therapeutic hypothermia and IABP was associated with improved neurological outcome at 1	Included in SLRs (Zhong 2024, Pagura 2024)

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

resuscitation in adults with out-of-hospital cardiac arrest: a prospective observational study. Resuscitation 85(6): 762-8		and 6 months after OHCA. CPC 1 or 2 were 12% in the ECPR group and 2% in the non-ECPR group at 1 month ($p<0.0001$), and 11% and 3% at 6 months ($p=0.001$), respectively.	
Smalcova J, Suen J, Huptych M et al. (2025) The significance of possible non-occlusive mesenteric ischemia in relation to neurological outcomes in patients with refractory cardiac arrest - Secondary analysis of the Prague OHCA study. Resuscitation 110642	Post-hoc analysis of Prague OHCA study n=256	The development of profuse diarrhoea, abdominal distension and other signs suggesting non-occlusive mesenteric ischaemia in people with refractory out-of-hospital cardiac arrest are observed more frequently in people with poor neurological outcome at day 180, especially in those who have ECPR.	Original trial is included in all key evidence systematic reviews.
Spangenberg T, Schewel J, Dreher A et al. (2018) Health related quality of life after extracorporeal cardiopulmonary resuscitation in refractory cardiac arrest. Resuscitation 127: 73-78	Single centre retrospective study (Germany) n=60 Follow-up: 1 year	12-month survival was 31%. HRQoL SF-36 scores of survivors ranged markedly below controls ($p<0.0001$)	Larger, more recent systematic literature reviews and meta-analyses included.
Springer A, Dreher A, Reimers J et al. (2023) Gender disparities in patients undergoing extracorporeal cardiopulmonary resuscitation. Frontiers in Cardiovascular Medicine 10: 1265978	Retrospective cohort study n=377	The results suggest a statistically significant survival benefit for females who have ECPR, possibly driven by a higher prevalence of witnessed collapse and bystander CPR.	Larger studies are included.
Sun P, Liu W, Li M et al. (2023) Extracorporeal	Retrospective cohort study n=133 (33 ECPR)	Among 22 propensity score matched pairs, 3 months after discharge,	Larger studies are included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

cardiopulmonary resuscitation versus conventional cardiopulmonary resuscitation for patients with refractory out-of-hospital cardiac arrest: A retrospective propensity matching analysis. Perfusion: 2676591231222365		the survival rate of those with good neurological function was 14% (3 out of 22) in the ECPR group and 4% (1 out of 22 cases) in the CCPR group, $p=0.61$, and the survival rate at discharge was 18% (4 out of 22) in the ECPR group and 4% (1 out of 22) in the CCPR group, $p=0.34$.	
Suverein MM, Delnoij TSR, Lorusso R et al. (2023) Early extracorporeal CPR for refractory out-of-hospital cardiac arrest. The New England Journal of Medicine 388(4): 299-309	INCEPTION randomised controlled trial $n=160$ (70 ECPR) Follow-up: 6 months	In people with refractory OHCA, ECPR and CCPR had similar effects on survival with a favourable neurological outcome. At 30 days, 14 people (20%) in the ECPR group were alive with a favourable neurological outcome, compared with 10 people (16%) in the CCPR group (OR 1.4; 95% CI 0.5 to 3.5; $p=0.52$). The number of serious adverse events per patient was similar in the 2 groups.	RCT included all SLRs
Tanimoto A, Sugiyama K, Tanabe M et al. (2020) Out-of-hospital cardiac arrest patients with an initial non-shockable rhythm could be candidates for extracorporeal cardiopulmonary resuscitation: a retrospective study. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine 28(1): 101	Single centre retrospective study (Japan) $n=186$ Follow-up: hospital discharge	The rate of good outcomes at hospital discharge was not significantly different between the shockable and non-shockable groups (19% versus 16%, $p=0.69$).	Larger, more recent systematic literature reviews and meta-analyses included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

Tonna J, Boonstra P, MacLaren G et al. (2024) Extracorporeal Life Support Organization Registry International Report 2022: 100,000 Survivors. ASAIO Journal 70: 131–43	Registry (ELSO) n=154,568 (8,050 ECPR procedures in adults)	Survival to hospital discharge was 30% (4,162/14,097) for adult ECPR. 67% of all adults with ECPR, and 74% of those with a reported Cerebral Performance Category value, had a score of 5 (brain death) or death.	Registry data includes all ages and all indications for ECMO.
Ubben JFH, Suverein MM, Delnoij TSR et al. (2024) Early extracorporeal CPR for refractory out-of-hospital cardiac arrest - A pre-planned per-protocol analysis of the INCEPTION-trial. Resuscitation 194: 110033	Per protocol analysis and Bayesian meta-analysis of the INCEPTION RCT n=81 Follow-up: 30 days	30-day survival with CPC1 or 2 was 15% in the ECPR group versus 9% in the CCPR group (adjusted OR 1.9, 95% CI 0.4 to 9.3; p=0.393). Bayesian analysis showed an 84% posterior probability of any ECPR benefit and a 61% posterior probability of a 5% absolute risk reduction for the primary outcome.	Original RCT included in all key evidence systematic reviews.
van de Koolwijk AF, Delnoij TSR, Suverein MM et al. (2024) Health-related quality of life one year after refractory cardiac arrest treated with conventional or extracorporeal CPR; a secondary analysis of the INCEPTION-trial. Resuscitation plus 19: 100669	Secondary analysis of INCEPTION trial n=134	Hospital survival was 20% (n=27). EQ-5D-5L data were available for 25 people (5 ECPR and 20 CCPR). One year after OHCA, the estimated mean health utility index was 0.73 (0.05) in all people, 0.84 (0.12) in ECPR survivors, and 0.71 (0.05) in CCPR survivors (p=0.31). Eight (32%) survivors had a poor HRQoL. HRQoL was good in 17 (68%) people, with 100% in ECPR survivors versus 60% in CCPR survivors (p=0.14).	Original trial is included in all key evidence systematic reviews.
Wang J-Y, Chen Y, Dong R et al. (2024) Extracorporeal vs.	Systematic review and meta-analysis	The pooled meta-analysis demonstrated that compared with CCRP,	Larger, more comprehensive

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

conventional CPR for out-of-hospital cardiac arrest: A systematic review and meta-analysis. The American journal of emergency medicine 80: 185-193	n=4,669 2 RCTs and 10 observational studies Follow-up: hospital discharge, 6 months	ECPR did not improve survival and neurological outcomes at 180 days following OHCA (RR 3.39, 95% CI 0.79 to 14.64; RR 2.35, 95% CI 0.97 to 5.67). While a beneficial effect of ECPR was obtained regarding 30-day survival and neurological outcomes.	systematic literature reviews and meta-analyses included.
Wongtanasarasin W, Krintratun S, Techasatian W et al. (2023) How effective is extracorporeal life support for patients with out-of-hospital cardiac arrest initiated at the emergency department? A systematic review and meta-analysis. PloS one 18(11): e0289054	Systematic review and meta-analysis n=51,173, 8 studies Follow-up: 30 days	ED-initiated ECPR may not be associated with a significant increase in favourable neurological outcomes (OR 1.43, 95% CI 0.30 to 6.70, $I^2=96\%$). However, this intervention may be linked to improved survival to hospital discharge (OR 3.34, 95% CI 2.23 to 5.01, $I^2=17\%$).	Larger, more comprehensive systematic literature reviews and meta-analyses included.
Yannopoulos D, Bartos J, Raveendran G et al. (2020) Advanced reperfusion strategies for patients with out-of-hospital cardiac arrest and refractory ventricular fibrillation (ARREST): a phase 2, single centre, open-label, randomised controlled trial. Lancet 396 (10265): 1807-1816	ARREST randomised controlled trial n=35 (15 ECPR) Follow-up: 6 months	Survival to hospital discharge was observed in one (7%) of 15 people in the standard ACLS treatment group versus 6 (43%) of 14 people in the early ECMO-facilitated resuscitation group (risk difference 36.2%, 3.7 to 59.2; posterior probability of ECMO superiority 0.9861). The study was terminated at the first preplanned interim analysis after enrolling 30 people because the posterior probability of ECMO superiority exceeded the prespecified monitoring boundary. Cumulative 6-month survival was	RCT included in all SLRs

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest

		significantly better in the early ECMO group than in the standard ACLS group.	
Yuan Z, Liu Y, Wei G et al. (2024) Clinical characteristics of extracorporeal cardiopulmonary resuscitation in China: a multicenter retrospective study. BMC Anesthesiology 24: 230	Retrospective cohort study n=648	The number of people who had ECPR increased from 33 in 2017 to 274 in 2020, and the survival rate increased from 24% to 34%.	Larger studies are included.

IP overview: VA ECMO for extracorporeal cardiopulmonary resuscitation (ECPR) in adults with refractory cardiac arrest