

## Post-traumatic stress disorder: management

### Review questions

*NICE guideline tbc*

*Review questions*

*April 2018*

*Draft for Consultation*

*Evidence reviews were developed by  
National Guideline Alliance hosted by the  
Royal College of Obstetricians and  
Gynaecologists*



## **Disclaimer**

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

## **Copyright**

© National Institute for Health and Care Excellence, 2016. All rights reserved. Subject to Notice of rights.

ISBN:

## **Contents**

<b>Review questions .....</b>	<b>5</b>
-------------------------------	----------

## Review questions

### Review questions

RQ. 1.1 For children and young people at risk of PTSD, what are the relative benefits and harms of psychological, psychosocial or other non-pharmacological interventions targeted at PTSD symptoms?

RQ. 1.2 For children and young people with clinically important post-traumatic stress symptoms, what are the relative benefits and harms of psychological, psychosocial or other non-pharmacological interventions targeted at PTSD symptoms?

RQ. 2.1 For adults at risk of PTSD, what are the relative benefits and harms of psychological, psychosocial or other non-pharmacological interventions targeted at PTSD symptoms?

RQ. 2.2 For adults with clinically important post-traumatic stress symptoms, what are the relative benefits and harms of psychological, psychosocial or other non-pharmacological interventions targeted at PTSD symptoms?

RQ. 3.1 For children and young people at risk of PTSD, what are the relative benefits and harms of specific pharmacological interventions?

RQ. 3.2 For children and young people with clinically important post-traumatic stress symptoms, what are the relative benefits and harms of specific pharmacological interventions?

RQ. 4.1 For adults at risk of PTSD, what are the relative benefits and harms of specific pharmacological interventions?

RQ. 4.2 For adults with clinically important post-traumatic stress symptoms, what are the relative benefits and harms of specific pharmacological interventions?

RQ. 5.1 For family members (including children and carers) of people at risk of PTSD, do specific psychological, psychosocial or other non-pharmacological interventions result in an improvement in their mental health and wellbeing, a reduction in burden and improved social and occupational outcomes?

RQ. 5.2 For family members (including children and carers) of people with clinically important post-traumatic stress symptoms, do specific psychological, psychosocial or other non-pharmacological interventions result in an improvement in their mental health and wellbeing, a reduction in burden and improved social and occupational outcomes?

RQ. 6.1 For adults, children and young people with clinically important post-traumatic stress symptoms, what factors should be taken into account in order to provide access to care, optimal care and coordination of care?

RQ. 7.1 Which service delivery models are effective at meeting the needs of adults, children and young people with clinically important post-traumatic stress symptoms?

RQ. 7.2 For adults, children and young people with clinically important post-traumatic stress symptoms, what are the aspects of a clinical care pathway that are associated with better outcomes?