



## **Evidence reviews**

Evidence review

Published: 25 September 2019

www.nice.org.uk

## **Evidence reviews 2019**

The following documents contain the evidence that was used to develop the recommendations:

- Accessibility and sustainability of abortion services
- Information needs of women undergoing an abortion
- Anti-D prophylaxis for women up to 13+6 weeks' gestation (replaced with <u>supporting</u> document Q for the recommendations on anti-D prophylaxis)
- Antibiotic prophylaxis for medical and surgical abortion
- VTE prophylaxis for women having abortion
- Abortion before ultrasound evidence
- Expulsion at home for early medical abortion
- <u>Simultaneous versus delayed mifepristone + misoprostol administration for medical</u> abortion up to 10+0 weeks' gestation
- Follow-up after medical abortion up to 10+0 weeks
- Misoprostol after mifepristone for inducing medical abortion between 10+1 and 24+0 weeks' gestation
- Medical versus surgical abortion between 13+0 and 24+0 weeks' gestation
- Medical abortion after 24 weeks' gestation
- Cervical priming before surgical abortion
- Anaesthesia or sedation for surgical abortion
- Support after abortion
- Contraception after abortion

## Other supporting evidence

- Methods
- Health economics
- Supporting document Q for the recommendations on anti-D prophylaxis