

Draft for consultation

Colorectal cancer (update)

[C8] Optimal duration of adjuvant chemotherapy for colorectal cancer

NICE guideline TBC

Evidence reviews

July 2019

Draft for Consultation

These evidence reviews were developed by the National Guideline Alliance hosted by the Royal College of Obstetricians and Gynaecologists

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2019. All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

Optimal duration of adjuvant chemotherapy for colorectal cancer	6
Review question	6
Introduction	6
Summary of the protocol	6
Methods and process	7
Clinical evidence	7
Summary of clinical studies included in the evidence review	7
Quality assessment of clinical studies included in the evidence review	9
Economic evidence	9
Economic model.....	9
Evidence statements	10
The committee's discussion of the evidence.....	14
References.....	18
Appendices.....	21
Appendix A – Review protocol.....	21
Review protocol for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer	21
Appendix B – Literature search strategies	25
Literature search strategies for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	25
Appendix C – Clinical evidence study selection	27
Clinical study selection for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	27
Appendix D – Clinical evidence tables	28
Clinical evidence tables for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	28
Appendix E – Forest plots.....	52
Forest plots for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	52
Appendix F – GRADE tables	60
GRADE tables for review question: What is the optimal duration of adjuvant chemotherapy for non-metastatic colorectal cancer?	60
Appendix G – Economic evidence study selection.....	68
Economic evidence study selection for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?.....	68
Appendix H – Economic evidence tables	69

Economic evidence tables for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	69
Appendix I – Economic evidence profiles	72
Economic evidence profiles for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	72
Appendix J – Economic analysis	74
Economic analysis for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	74
Appendix K – Excluded studies	75
Excluded clinical studies for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	75
Appendix L – Research recommendations	78
Research recommendations for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?	78

1 **Optimal duration of adjuvant 2 chemotherapy for colorectal cancer**

3 This evidence review supports recommendation 1.3.14.

4 **Review question**

5 What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

6 **Introduction**

7 Adjuvant chemotherapy for 24 weeks (6 months) has previously been established as
8 the standard of care for stage III colorectal cancer (Andre 2003). In recent years a
9 shorter duration of adjuvant chemotherapy have been suggested in order to minimise
10 the adverse long-term effects of chemotherapy, mainly neurotoxicity. This review
11 aims to find out what is the optimal duration of adjuvant chemotherapy for colorectal
12 cancer taking into consideration its effects on for example survival and cancer
13 recurrence, neurotoxicity and quality of life.

14 **Summary of the protocol**

15 Please see Table 1 for a summary of the population, intervention, comparison and
16 outcomes (PICO) characteristics of this review.

17 **Table 1: Summary of the protocol (PICO table)**

Population	Adults with non-metastatic colorectal cancer after receiving surgery with curative intent Subgroups to be considered separately: <ul style="list-style-type: none">• pT4• pT3/T4 N0 with vascular invasion• right versus left sided tumour• age over 70 years
Intervention	• Adjuvant chemotherapy for less than 24 weeks (6 months)
Comparison	• Adjuvant chemotherapy for 24 weeks (6 months)
Outcomes	Critical <ul style="list-style-type: none">• Disease-free survival• Overall survival• Neuropathy (lasting for 2 years considered permanent) Important <ul style="list-style-type: none">• Overall quality of life• Distant metastasis• Treatment-related mortality• Dose reduction

18 *N: nodal stage; p: pathological staging; T: tumour stage*

1 Methods and process

2 This evidence review was developed using the methods and process described in
3 [Developing NICE guidelines: the manual 2014](#). Methods specific to this review
4 question are described in the review protocol in appendix A.
5 Declarations of interest were recorded according to NICE's 2014 conflicts of interest
6 policy until 31 March 2018. From 1 April 2018, declarations of interest were recorded
7 according to NICE's 2018 [conflicts of interest policy](#). Those interests declared until
8 April 2018 were reclassified according to NICE's 2018 conflicts of interest policy (see
9 Register of Interests).

10 Clinical evidence

11 Included studies

12 Seven RCTs (reported in 7 publications) were included. Data from 6 RCTs (TOSCA,
13 SCOT, IDEA France, CALGB/SWOG, HORG, ACHIEVE) were reported in 1
14 collaborative paper (Grothey 2018) which pooled individual patient data from these
15 studies. Additional data from 3 of these 6 RCTs were also reported in separate
16 publications (IDEA France [Andre 2018]; SCOT [Iveson 2018]; TOSCA [Lonardi
17 2016]). One RCT (reported in 3 publications) was also included (Chau 2005a; Chau
18 2005b; Saini 2003).

19 The included studies are summarised in Table 2.

20 The collaborative paper (Grothey 2018), combined individual patient data from 6
21 RCTs (TOSCA, SCOT, IDEA France, CALGB/SWOG, HORG, ACHIEVE) to compare
22 3 months and 6 months of oxaliplatin-based adjuvant chemotherapy, either folinic
23 acid plus fluorouracil plus oxaliplatin (FOLFOX) or oxaliplatin plus capecitabine
24 (CAPOX), in people with colon cancer (IDEA Collaboration 2018; IDEA France
25 [Andre 2018]; SCOT [Iveson 2018]; TOSCA [Lonardi 2016]). An RCT within the IDEA
26 Collaboration also included people with rectal cancer (SCOT).

27 One RCT compared 3 months of protracted fluorouracil infusion to 6 months of bolus
28 fluorouracil plus folinic acid (leucovorin) (Chau 2005; Chau 2005a; Chau 2005b; Saini
29 2003).

30 See the literature search strategy in appendix B and study selection flow chart in
31 appendix C.

32 Excluded studies

33 Studies not included in this review with reasons for their exclusions are provided in
34 appendix K.

35 Summary of clinical studies included in the evidence review

36 Summaries of the studies that were included in this review are presented in Table 2.

37 Table 2: Summary of included clinical studies

Study	Population	Intervention/Comparison	Outcomes
Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy			

DRAFT FOR CONSULTATION
Optimal duration of adjuvant chemotherapy for colorectal cancer

Study	Population	Intervention/Comparison	Outcomes
IDEA collaboration (Grothey 2018) Individual patient data from 6 RCTs: <ul style="list-style-type: none">• TOSCA (Italy)• SCOT (UK, Denmark, Spain, Australia, Sweden, New Zealand)• IDEA France• CALGB/SWOG (US, Canada)• HORG (Greece)• ACHIEVE (Japan)	N=12,834 people with stage III colon cancer <ul style="list-style-type: none">• TOSCA n=2,402• SCOT n=3,983• IDEA France n=2,010• CALGB/SWOG n=2,440• HORG n=708• ACHIEVE n=1,291	3 months versus 6 months of adjuvant chemotherapy Type of chemotherapy: <ul style="list-style-type: none">• folinic acid, fluorouracil and oxaliplatin (FOLFOX), 60%• capecitabine and oxaliplatin (CAPOX), 40%	<ul style="list-style-type: none">• Disease-free survival• Grade 3 or 4 peripheral neurotoxicity• Percentage of chemotherapy dose delivered
IDEA France trial (Andre 2018) RCT France	N=2,022 people aged 18 years or older with stage III colon cancer	3 months versus 6 months of adjuvant chemotherapy Type of chemotherapy: <ul style="list-style-type: none">• folinic acid, fluorouracil and oxaliplatin (FOLFOX), 90%• capecitabine and oxaliplatin (CAPOX), 10%	<ul style="list-style-type: none">• Overall survival• Treatment-related mortality Disease-free survival, neurotoxicity and dose reduction from the IDEA France trial are reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.
SCOT trial (Iveson 2018) RCT UK, Denmark, Spain, Sweden, Australia, and New Zealand	N=6,088 people aged 18 years or older with stage III or high-risk stage II colon or rectal cancer	3 months versus 6 months of adjuvant chemotherapy Type of chemotherapy: <ul style="list-style-type: none">• folinic acid, fluorouracil and oxaliplatin (FOLFOX), 33%• capecitabine and oxaliplatin (CAPOX), 67%	<ul style="list-style-type: none">• Disease-free survival (for people with rectal cancer)• Overall survival• Quality of life• Treatment-related mortality Disease-free survival (for people with colon cancer), neurotoxicity, and dose reduction from the SCOT trial are reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.
TOSCA trial (Lonardi 2016) RCT	N=3,759 people aged 18 years or older with stage III or high-risk stage II colon cancer	3 months versus 6 months of adjuvant chemotherapy	<ul style="list-style-type: none">• Treatment-related mortality Disease-free survival and

Study	Population	Intervention/Comparison	Outcomes
Italy		Type of chemotherapy: • folinic acid, fluorouracil and oxaliplatin (FOLFOX), 64% • capecitabine and oxaliplatin (CAPOX), 36%	neurotoxicity are reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.
Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil/leucovorin			
Chau 2005 (Chau 2005a; Chau 2005b; Saini 2003)	N=826 people with curatively resected stage II and III adenocarcinoma of the colon or rectum	12 weeks of protracted venous infusion of fluorouracil versus 6 months of bolus fluorouracil/leucovorin	<ul style="list-style-type: none"> Disease-free survival Overall survival Quality of life Distant metastasis Chemotherapy-related mortality Percentage of chemotherapy dose delivered
RCT			
UK			

1 *N: number; RCT: randomised controlled trial*

2 See the full evidence tables in appendix D and the forest plots in appendix E.

3 Quality assessment of clinical studies included in the evidence review

4 See the clinical evidence profiles in appendix F.

5 Economic evidence

6 Included studies

7 One relevant study was identified in a literature review of published cost-effectiveness analyses on this topic (Robles-Zurita 2018; see appendix H and appendix I for summary and full evidence tables). The study compared a 3 month to a 6 month regimen of adjuvant chemotherapy in patients with fully resected high-risk stage II or stage III colorectal cancer. The study compared the length of adjuvant chemotherapy separately for both CAPOX and FOLFOX.

13 The economic analysis was a within study cost-utility analysis with all resource use and outcome data collected alongside the SCOT RCT considered in the clinical evidence review (Iveson 2018). The study took a NHS & PSS perspective.

16 Excluded studies

17 A global search of economic evidence was undertaken for all review questions in this guideline. See Supplement 2 for further information.

19 Economic model

20 No economic modelling was undertaken for this review because the committee agreed that other topics were higher priorities for economic evaluation.

1 **Evidence statements**

2 **Clinical evidence statements**

3 **Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant
4 chemotherapy for colorectal cancer**

5 **Critical outcomes**

6 **Disease-free survival**

- 7 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
8 (N=12,834; median follow-up 3.5 years) showed that there may be a clinically
9 important lower disease-free survival in people who received 3 months adjuvant
10 chemotherapy compared to 6 months of adjuvant chemotherapy for non-
11 metastatic colon cancer, but there is uncertainty around the estimate.
- 12 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
13 (N=5,071; median follow-up 3.5 years) showed no clinically important difference in
14 disease-free survival in the subpopulation of people who received 3 months or 6
15 months of adjuvant CAPOX chemotherapy for non-metastatic colon cancer.
- 16 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
17 (N=7,763; median follow-up 3.5 years) showed a clinically important lower
18 disease-free survival for subpopulation of people who received 3 months of
19 adjuvant FOLFOX chemotherapy compared to 6 months of adjuvant FOLFOX
20 chemotherapy for non-metastatic colon cancer.
- 21 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
22 (N=2,655; median follow-up 3.5 years) showed a clinically important lower
23 disease-free survival for subpopulation of people with stage T4 colon cancer who
24 received 3 months of adjuvant chemotherapy compared to 6 months of adjuvant
25 chemotherapy.
- 26 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
27 (N=7,471; median follow-up 3.5 years) showed no clinically important difference in
28 disease-free survival in the subpopulation of people with stage T1-3N1 colon
29 cancer who received 3 months or 6 months of adjuvant chemotherapy.
- 30 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
31 (N not reported; median follow-up 3.5 years) showed that there may be a clinically
32 important better disease-free survival in the subpopulation of people with stage
33 T1-3N1 colon cancer who received 3 months of adjuvant CAPOX chemotherapy
34 compared to 6 months of adjuvant CAPOX chemotherapy, but there is uncertainty
35 around the estimate.
- 36 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
37 (N not reported; median follow-up 3.5 years) showed no clinically important
38 difference in disease-free survival in the subpopulation of people with stage T1-
39 3N1 colon cancer who received 3 months or 6 months of adjuvant FOLFOX
40 chemotherapy.
- 41 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
42 (N=5,256; median follow-up 3.5 years) showed a clinically important lower
43 disease-free survival for the subpopulation of people with stage T4 and/or N2
44 colon cancer who received 3 months of adjuvant chemotherapy compared to 6
45 months of adjuvant chemotherapy.
- 46 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
47 (N not reported; median follow-up 3.5 years) showed no clinically important
48 difference in disease-free survival in the subpopulation of people with stage T4

- 1 and/or N2 colon cancer who received 3 months or 6 months of adjuvant CAPOX
2 chemotherapy.
- 3 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
4 (N not reported; median follow-up 3.5 years) showed a clinically important lower
5 disease-free survival for the subpopulation of people with stage T4 and/or N2
6 colon cancer who received 3 months of adjuvant FOLFOX chemotherapy
7 compared to 6 months of adjuvant FOLFOX chemotherapy.
 - 8 • Moderate quality evidence from 1 RCT (N=1,098; median follow-up 3.1 years)
9 showed no clinically important difference in disease-free survival in the
10 subpopulation of people with non-metastatic rectal cancer who received 3 months
11 or 6 months of adjuvant chemotherapy.

12 **Overall survival**

- 13 • High quality evidence from 2 RCTs (N=8,075; median follow-up 3.1 to 4.3 years)
14 showed no clinically important difference in overall survival between 3 months and
15 6 months of adjuvant chemotherapy for people with non-metastatic colorectal
16 cancer.
- 17 • Moderate quality evidence from 1 RCT (N=201; median follow-up 4.3 years)
18 showed no clinically important difference in overall survival in the subpopulation of
19 people who received 3 months or 6 months of adjuvant CAPOX chemotherapy for
20 non-metastatic colon cancer.
- 21 • Moderate quality evidence from 1 RCT (N=1,809; median follow-up 4.3 years)
22 showed no clinically important difference in overall survival in the subpopulation of
23 people who received 3 months or 6 months of adjuvant FOLFOX chemotherapy
24 for non-metastatic colon cancer.

25 **Neuropathy**

- 26 • Moderate quality evidence from meta-analysis of individual patient data from 6
27 RCTs (N=12,834) showed a clinically important lower risk of grade 3 or 4
28 neuropathy in people who received 3 months of adjuvant chemotherapy compared
29 to 6 months of adjuvant chemotherapy for non-metastatic colon cancer.
- 30 • Moderate quality evidence from meta-analysis of individual patient data from 6
31 RCTs (N=12,834) showed a clinically important lower risk of grade 3 or 4
32 neuropathy in the subpopulation of people who received 3 months of adjuvant
33 CAPOX chemotherapy compared to 6 months of adjuvant chemotherapy for non-
34 metastatic colon cancer.
- 35 • Moderate quality evidence from meta-analysis of individual patient data from 6
36 RCTs (N=12,834) showed a clinically important lower risk of grade 3 or 4
37 neuropathy in the subpopulation of people who received 3 months of adjuvant
38 FOLFOX chemotherapy compared to 6 months of adjuvant chemotherapy for non-
39 metastatic colon cancer.

40 **Important outcomes**

41 **Overall quality of life**

- 42 • Moderate quality evidence from 1 RCT (N=6,088) showed a clinically important
43 better quality of life (measure using QLQ-C30 global health status score and EQ-
44 5D VAS) at 6 months in people who received 3 months of adjuvant chemotherapy
45 compared to 6 months of adjuvant chemotherapy for non-metastatic colorectal
46 cancer. There was no difference in quality of life at 12 months (measured using
47 the same scales).

1 **Distant metastasis**

2 No evidence was identified to inform this outcome.

3 **Treatment-related mortality**

- 4 • Moderate quality evidence from 3 RCTs (N=11,729) showed no clinically important
5 difference in treatment-related mortality between 3 months and 6 months of
6 adjuvant FOLFOX chemotherapy for people with non-metastatic colorectal cancer.

7 **Dose reduction**

- 8 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
9 (N=5,071) showed a clinically important higher percentage of planned oxaliplatin
10 dose delivered in the subpopulation of people who received 3 months of adjuvant
11 CAPOX chemotherapy compared to 6 months of adjuvant CAPOX chemotherapy
12 for non-metastatic colon cancer.
- 13 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
14 (N=7,763) showed a clinically important higher percentage of planned oxaliplatin
15 dose delivered in the subpopulation of people who received 3 months of adjuvant
16 FOLFOX chemotherapy compared to 6 months of adjuvant FOLFOX
17 chemotherapy for non-metastatic colon cancer.
- 18 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
19 (N=7,763) showed a clinically important higher percentage of planned fluorouracil
20 dose delivered in the subpopulation of people who received 3 months of adjuvant
21 FOLFOX chemotherapy compared to 6 months of adjuvant FOLFOX
22 chemotherapy for non-metastatic colon cancer.
- 23 • High quality evidence from meta-analysis of individual patient data from 6 RCTs
24 (N=5,071) showed a clinically important higher percentage of planned
25 capecitabine dose delivered in the subpopulation of people who received 3
26 months of adjuvant CAPOX chemotherapy compared to 6 months of adjuvant
27 CAPOX chemotherapy for non-metastatic colon cancer.

28 **Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus
29 fluorouracil/leucovorin**

30 **Critical outcomes**

31 **Disease-free survival**

- 32 • Moderate quality evidence from 1 RCT (N=801; median follow-up 5.4 years)
33 showed that there may be a clinically important better disease-free survival in
34 people who received 3 months of fluorouracil infusion compared to 6 months of
35 bolus fluorouracil and leucovorin for non-metastatic colorectal cancer, but there is
36 uncertainty around the estimate.
- 37 • Moderate quality evidence from 1 RCT (N=323; median follow-up 4.6 years)
38 showed a clinically important better survival in the subpopulation of people with
39 non-metastatic rectal cancer who received 3 months of fluorouracil infusion
40 compared to 6 months of bolus fluorouracil and leucovorin.

41 **Overall survival**

- 42 • Moderate quality evidence from 1 RCT (N=801; median follow-up 5.4 years)
43 showed that there may be a clinically important better overall survival in people
44 who received 3 months of fluorouracil infusion compared to 6 months of bolus

1 fluorouracil and leucovorin for non-metastatic colorectal cancer, but there is
2 uncertainty around the estimate.
3 • Moderate quality evidence from 1 RCT (N=323; median follow-up 4.6 years)
4 showed that there may be a clinically important better overall survival in the
5 subpopulation of people with non-metastatic rectal cancer who received 3 months
6 of fluorouracil infusion compared to 6 months of bolus fluorouracil and leucovorin
7 for non-metastatic colorectal cancer, but there is uncertainty around the estimate.

8 **Neuropathy**

9 No evidence was identified to inform this outcome.

10 **Important outcomes**

11 **Overall quality of life**

12 • Very low quality evidence from 1 RCT (N=801) showed no clinically important
13 difference in quality of life at 24 weeks and at 2 years after randomisation
14 (measured using QLQ-C30 global health status score) in people who received 3
15 months of fluorouracil infusion compared to 6 months of bolus fluorouracil and
16 leucovorin for non-metastatic colorectal cancer.

17 **Distant metastasis**

18 • Moderate quality evidence from 1 RCT (N=801) showed no clinically important
19 difference in distant metastasis in people who received 3 months of fluorouracil
20 infusion or 6 months of bolus fluorouracil and leucovorin for non-metastatic
21 colorectal cancer.

22 **Treatment-related mortality**

23 • High quality evidence from 1 RCT (N=801) showed that there were no
24 chemotherapy-related deaths in people with non-metastatic colorectal cancer in
25 either the 3-month or 6-month chemotherapy arms.

26 **Dose reduction**

27 • High quality evidence from 1 RCT (N=692) showed a clinically important higher
28 percentage of dose delivered in people who received 3 months of fluorouracil
29 infusion compared to 6 months of bolus fluorouracil and leucovorin for non-
30 metastatic colorectal cancer.

31 **Economic evidence statements**

32 One cost utility analysis showed that for people receiving CAPOX chemotherapy a 3
33 month course is both cost saving and more effective compared to a 6 month course
34 with a less than 1% probability that the 6-month regimen provides a cost effective
35 use of NHS resources when QALYs are valued at £20,000 each. The same analysis
36 showed that whilst a 3 month course of FOLFOX chemotherapy would be less
37 effective it would also be cost saving. The saving per QALY forgone was greater than
38 £50,000 suggesting that the 3 month course would be cost effective if QALYs are
39 valued at £20,000 each. There was a less than 10% probability of a 6 month course
40 being cost effective when QALYs are valued at £20,000 each. The study took an
41 NHS perspective and was considered directly applicable to the decision problem with
42 only minor methodological limitations.

1 The committee's discussion of the evidence

2 Interpreting the evidence

3 *The outcomes that matter most*

4 The aim of this review was to compare the effectiveness and safety of a shorter
5 duration of adjuvant chemotherapy to the standard 6 months of adjuvant
6 chemotherapy for people with colorectal cancer. Disease-free survival and overall
7 survival were considered critical outcomes for decision-making because ultimately
8 the aim of cancer treatment is to improve survival. Neuropathy was also considered a
9 critical outcome as it is the main long-term adverse effect of chemotherapy which can
10 have a significant negative impact on people's lives.

11 Quality of life was considered an important outcome as it can capture many aspects
12 of the benefits and harms of treatment. Distant metastasis was also considered an
13 important outcome as it often related to survival and the need for further treatment.
14 Dose reduction was considered an important outcome as it reflects the short-term
15 toxicity of the treatment as well as its effectiveness.

16 *The quality of the evidence*

17 Evidence was available for the comparison of 3 months versus 6 months of
18 oxaliplatin-based adjuvant chemotherapy for colorectal cancer (comparison 1), and
19 for the comparison of 3 months of fluorouracil infusion versus six months of bolus
20 fluorouracil/leucovorin (comparison 2).

21 For comparison 1, evidence was available for all of the outcomes except distant
22 metastasis and for comparison 2, evidence was available for all of the outcomes
23 except neuropathy. The quality of the evidence was assessed using GRADE and
24 was mostly of high quality, varying from low to high.

25 The main reasons for downgrading the quality of the evidence were imprecision of
26 the effect estimate, and lack of blinding for outcomes that were considered to be
27 subjective (quality of life and neuropathy were both measured using patient self-
28 reported questionnaires).

29 *Benefits and harms*

30 Six months of adjuvant oxaliplatin-based chemotherapy has been the standard of
31 care after surgery for stage III colorectal cancer patients (apart from rectal cancer
32 patients who have received long-course chemoradiotherapy preoperatively).
33 However, the major long-term adverse effect of oxaliplatin-based chemotherapy is
34 peripheral neuropathy that can have significant effects on a person's quality of life.
35 The committee agreed that it is important to find a balance where the long-term
36 toxicity of chemotherapy could be minimised without compromising its beneficial
37 effect on survival and disease recurrence. In addition, the committee discussed that it
38 is important to consider patient characteristics, including age, performance status,
39 and comorbidities as well as the histopathology of the cancer when considering
40 difference options of treatment, particularly in relation to the side-effects of the
41 different treatment options.

42 The evidence showed that the rate of severe neuropathy was considerably lower with
43 3 months of chemotherapy compared to the standard 6 months of chemotherapy
44 (1.8% versus 10% of the patients having grade 3 or 4 peripheral neuropathy). There
45 was no difference in chemotherapy-related mortality. The percentage of total planned
46 dose of chemotherapy received was higher among people in the 3-month group

1 (around 90% for oxaliplatin) than those in the 6-month group (around 70% for
2 oxaliplatin). This finding suggests that 3 months of chemotherapy was better
3 tolerated and there was less need for dose reductions due to side-effects.

4 In terms of disease-free survival the findings were more complex. In the total
5 population, regardless of disease stage or chemotherapy regimen, the evidence
6 suggested that disease-free survival might be worse in the 3-month group (74.6%
7 [95% CI 73.5% to 75.7%] at 3 years) than in the 6-month group (75.5% at 3 years),
8 although there is uncertainty around the estimate and the differences are small.
9 However, when stratifying according to the treatment regimen (CAPOX or FOLFOX),
10 the findings showed that 3 months of CAPOX chemotherapy was as effective as 6
11 months of CAPOX (75.9% [95% CI 74.2% to 77.6%] versus 74.8% at 3 years)
12 whereas FOLFOX chemotherapy for 3 months showed worse disease-free survival
13 than FOLFOX for 6 months (73.6% [95% CI 72.2% to 75.1%] versus 76.0% at 3
14 years). Stratifying further according cancer stage, the evidence suggested that for
15 people who received CAPOX and had “low risk” cancer (stage T1-3 and N1), 3
16 months of CAPOX may actually be better than 6 months of CAPOX in terms of
17 disease-free survival (85.0% [95% CI 83.1% to 86.9%] versus 83.1% at 3 years). For
18 people who received FOLFOX and had “low risk” cancer there was no statistically
19 significant difference between the groups (81.9% [95% CI 80.2% to 83.6%] versus
20 83.5% at 3 years). For people with “high risk” cancer (stage T4 and/or N2) who
21 received CAPOX, there was no difference in disease-free survival between the
22 groups suggesting that 3 months is as effective as 6 months of CAPOX
23 chemotherapy (64.1% [95% CI 61.3% to 67.1%] versus 64.0% at 3 years). On the
24 other hand, those with “high risk” cancer who received FOLFOX, 3 months of
25 FOLFOX showed worse disease-free survival than 6 months of FOLFOX (61.5%
26 [95% CI 58.9% to 64.1%] versus 64.7% at 3 years).

27 Evidence on overall survival showed no difference between the different durations
28 although evidence for the total population was only available from 2 of the IDEA
29 collaboration trials and from only 1 trial stratified by treatment regimen and thus lacks
30 statistical power.

31 The committee were aware that the trials making up the IDEA collaboration were
32 designed to compare 3 months with 6 months of chemotherapy but were not
33 designed to compare the duration according to the chemotherapy regimen (FOLFOX
34 and CAPOX) or by cancer stage. The analyses of regimen and cancer stage were
35 thus exploratory in nature. The large size of the IDEA collaboration however provides
36 high quality evidence. In addition, it should be noted that the IDEA collaboration
37 based their interpretation of the results to a non-inferiority level of 1.12 (meaning that
38 non-inferiority of 3 months was proven only if the upper confidence interval of the
39 hazard ratio did not exceed 1.12, which corresponds to about 2.7 percentage point
40 difference in disease-free survival). As outlined in the review protocol, the committee
41 on the other hand considered statistical significance as the basis for judging if there
42 is a clinically important difference.

43 Weighing the benefits and harms of the different chemotherapy durations and
44 regimens, the committee agreed that people with colon cancer should be offered
45 CAPOX chemotherapy for 3 months as it has lower long-term toxicity but equal effect
46 on disease relapse and survival. Non-inferiority was seen for the entire treatment
47 group and in the exploratory analysis of “low risk” patients. In the “high risk” patient
48 population no difference in 3-year disease free survival was observed even though
49 the IDEA collaboration concluded that statistical non-inferiority was not confirmed
50 due to the upper confidence interval of the hazard ratio exceeding 1.12. Regardless,
51 the committee agreed that the risk of peripheral neuropathy significantly outweighs
52 any possible benefit of continuing CAPOX for 6 months in the “high risk” group of

1 patients. The committee therefore recommended 3 months of CAPOX chemotherapy
2 for all patients. Both capecitabine and oxaliplatin (drugs in the CAPOX regimen) are
3 generic drugs and used widely in current practice in the UK.

4 However, the committee recognised that CAPOX is not appropriate for all people
5 because of the side-effects of treatment. CAPOX is taken every 3 weeks with a large
6 dose of oxaliplatin and may cause side-effects such as fatigue, nausea, and
7 diarrhoea that people with comorbidities or who are frail might have difficulty
8 tolerating. Analysis from the SCOT trial showed higher rates of severe diarrhoea and
9 hand foot syndrome with CAPOX chemotherapy. Therefore, the committee agreed
10 that FOLFOX chemotherapy should be discussed as an alternative for such people.
11 The optimal duration of FOLFOX chemotherapy was difficult to determine as 6-month
12 course showed better disease-free survival but 3-month course showed less long-
13 term toxicity (neuropathy) and considerably less costs. Another alternative to
14 oxaliplatin containing chemotherapy which the committee recommended is single
15 agent capecitabine chemotherapy for 6 months. Single agent treatment is known to
16 be less effective than combination chemotherapy including capecitabine but is
17 associated with lower incidence of severe side-effects. This treatment is particularly
18 well tolerated in patients over 70 years of age who appear to gain little benefit from
19 the addition of oxaliplatin. Single agent capecitabine has been recommended by
20 previous NICE technology appraisal on [capecitabine and oxaliplatin in the adjuvant
treatment of stage III \(Dukes' C\) colon cancer \(TA100\)](#).

22 Not surprisingly, the 3-month group had a better overall quality of life at 6 months,
23 when the 6-month group is still on or about to finish chemotherapy. However, the
24 committee was surprised that no difference was observed in the quality of life at 1
25 year between the groups even though severe neuropathy was much more common
26 in the 6-month group.

27 People who received preoperative long-course radiotherapy or chemoradiotherapy
28 were not eligible for the IDEA collaboration trials. Therefore, rectal cancer patients
29 who had received preoperative long-course chemoradiotherapy are not covered by
30 these recommendations. Despite the lack of evidence among these people, a
31 research recommendation was not made because trials of adjuvant chemotherapy
32 after long-course radiotherapy or chemoradiotherapy have previously been
33 attempted but closed early due to poor recruitment (Glynne-Jones 2014). The focus
34 of national and international treatment has moved to intensifying pre-operative
35 treatment with studies investigating intensified chemo-radiotherapy regimens and
36 sequences of pre-operative systemic chemotherapy and pelvic radiotherapy
37 (ARITOTLE trial, www.isrctn.com/ISRCTN09351447, accessed 6 June 2019).

38 The SCOT trial, the largest trial within the IDEA collaboration, however, included
39 rectal cancer patients if they had had either short-course radiotherapy or no
40 preoperative therapy. Stratified analysis among these rectal cancer patients showed
41 that the main outcomes were similar for them as for the colon cancer patients,
42 therefore, the committee agreed that the recommendations for colon cancer patients
43 apply equally to those rectal cancer patients who have not received preoperative
44 long-course chemoradiotherapy.

45 The committee was also interested in the influence of age on the effectiveness of the
46 different chemotherapy durations. They were aware that previous studies comparing
47 fluorouracil therapy with or without oxaliplatin have shown reduced or no benefit of
48 oxaliplatin in patients over 70 years of age (McCleary 2013) and because of these
49 findings, many patients aged 70 years or older are considered for single agent
50 treatment alone (without oxaliplatin) in current practice. The available studies did not
51 provide results according to age and even if they would have the IDEA collaboration

1 trials included oxaliplatin in both intervention arms so the effect on age shown in the
2 previous studies was not applicable. Because there is no age-stratified evidence on
3 the effect of duration of chemotherapy, the committee was not able to make
4 recommendations according to the age of the patient.

5 The review also included an earlier UK trial comparing 3 months of protracted
6 fluorouracil infusion to 6 months of bolus fluorouracil and folinic acid (leucovorin). The
7 findings showed that 3-month chemotherapy may be better in terms of disease-free
8 survival and overall survival. However, the committee discussed that this trial did not
9 change practice when it was published because it was relatively small and under-
10 powered. In current UK practice, oral capecitabine is given as the treatment of
11 choice. Neither the bolus or infusional fluorouracil regimens investigated in the trial
12 are used routinely and therefore the interventions were not considered to be relevant
13 for current practice and the findings from this trial did not inform the
14 recommendations made.

15 Cost effectiveness and resource use

16 The literature search of previous economic evidence identified 1 economic evaluation
17 relevant to this topic. The study (Robles-Zurita 2018) compared a 3 month to a 6
18 month regimen of adjuvant chemotherapy in patients with fully resected high-risk
19 stage II or stage III colorectal cancer. The study compared the length of adjuvant
20 chemotherapy for 2 regimens-CAPOX and FOLFOX.

21 The study took a NHS & PSS perspective and was deemed to only have minor
22 methodological issues. The economic analysis was a within study cost-utility analysis
23 with all resource use and outcome data collected alongside the SCOT RCT
24 considered in the clinical evidence review (Iveson 2018). The RCT was conducted
25 across 2 Australasian and 4 European countries including the UK and the quality of
26 all clinical outcomes were rated either moderate or high using GRADE criteria.
27 Quality of life in the study was collected at baseline and all follow-up meetings in a
28 subsection of 1,832 patients equating to about 30% of the entire trial population. The
29 EQ-5D-3L questionnaire was used and scored using the UK general population
30 tariffs, the preferred method of NICE. All resource use was costed using publically
31 available UK costs.

32 The study found that a 3-month regimen of either CAPOX or FOLFOX led to cost
33 savings of £3,853 and £6,481 respectively during a maximum follow-up of 8 years.
34 CAPOX for 3 months led to an increase in both life expectancy (0.07 years) and
35 QALYs (0.19) dominating the 6-month regimen. The increase in QALYs is most likely
36 driven by reduced toxicity with a shorter duration of treatment and a reduction in
37 significant peripheral neuropathy. In the FOLFOX group moving from a 6- to a 3-
38 month regimen led to a reduction in both life expectancy (-0.22 years) and QALYs (-
39 0.12) although neither were statistically significant. This led to a saving of over
40 £50,000 for every QALY forgone. The 3-month FOLFOX regimen would be cost
41 effective under the conventional NICE threshold of £20,000 per QALY. Both these
42 results were robust during probabilistic sensitivity analysis with a >99% probability of
43 3-month CAPOX and >90% probability of 3-month FOLFOX being the preferred
44 option at a threshold of £20,000 per QALY. No deterministic sensitivity analyses were
45 presented.

46 The committee acknowledged that directly applicable, high quality economic
47 evidence was identified for this topic area and noted that it strongly favoured the 3-
48 month regimen for both CAPOX and FOLFOX. The committee were concerned that
49 recommending a 3-month treatment length for both CAPOX and FOLFOX in line with
50 the conclusions of the economic evidence may encourage an increased use of

1 CAPOX in patient groups where it may not be the most appropriate treatment.
2 Patients may, even after explaining the toxicity risks, opt for 3-month CAPOX given
3 the modest disease-free survival benefit from 3-month FOLFOX if an alternative of 6-
4 month FOLFOX is not available. This is a higher risk group of Grade 3/4/5 diarrhoea
5 with CAPOX chemotherapy. The committee noted during the recruitment of the
6 SCOT trial 8 patients died in the CAPOX arm from diarrhoea and vomiting. Due to
7 this the SCOT trial group specifically advised investigators on the management of
8 severe diarrhoea in patients receiving CAPOX. The committee considered whether
9 recommendations could be worded to prevent such crossover. The committee
10 considered that doing so could go against the NICE technology appraisal on
11 [capecitabine and oxaliplatin in the adjuvant treatment of stage III \(Dukes' C\) colon](#)
12 [cancer \(TA100\)](#). In particular the section on joint decision making when choosing the
13 most appropriate treatment. I would also likely go against further principles of patient
14 autonomy, consent and informed involvement in decision making.

15 Patients at high risk of severe chemotherapy-related diarrhoea may have worse
16 quality of life outcomes, higher treatment-related morbidity and mortality and costs
17 through treatment and hospitalisation from severe adverse events if they receive
18 CAPOX. It is difficult to estimate a proportion, given this is not currently a decision
19 faced in practise, but it would not be insignificant. This switch from FOLFOX to
20 CAPOX was not considered in the economic evaluation. It would not have been
21 picked up by 'intention to treat' as 6-month FOLFOX was available and clinical
22 evidence was limited at the time.

23 The committee also consider that whilst 3-8 years follow-up was a long time in the
24 context of an RCT, restricting the economic evaluation to this time horizon may
25 underestimate the true lifetime costs for the 6-month group. This is an area with
26 expensive downstream treatments and a patient group in which life expectancy is
27 increasing. The clinical evidence review showed that disease-free survival was worse
28 for 3 months FOLFOX chemotherapy than for 6 months FOLFOX (73.6% versus
29 76.0%, respectively, at 3 years), and that this result was statistically significant.
30 Prolonged courses of expensive palliative treatments downstream maybe forgone or
31 delayed in patients if they remain disease free longer. This again will decrease the
32 certainty around the cost effectiveness conclusions.

33 The committee concluded there was strong cost effectiveness evidence, based on
34 strong clinical evidence for recommending a 3-month course of CAPOX
35 chemotherapy. The committee acknowledge the strong economic evidence for a 3-
36 month FOLFOX regimen however given the clinical concerns, it was decided that
37 there should be an individualised consideration for the duration of FOLFOX for those
38 who are not suitable for 3-month CAPOX chemotherapy, taking into account the
39 benefits and short- and long-term harms of each option, and the person's
40 comorbidities and performance status and preference. The committee recognised
41 that this approach favours individual values over population values and that there is
42 potential for societal harm through inefficient allocation of resources.

43 References

44 Andre 2003

45 Andre T, Colin P, Louvet C, et al. (2003) Semimonthly versus monthly regimen of
46 fluorouracil and leucovorin administered for 24 or 36 weeks as adjuvant therapy in
47 stage II and III colon cancer: results of a randomized trial. Journal of Clinical
48 Oncology 21(15):2896-903

49 Chau 2005

- 1 Chau I, Norman A, Cunningham D, et al. (2005a) A randomised comparison between
2 6 months of bolus fluorouracil/leucovorin and 12 weeks of protracted venous infusion
3 fluorouracil as adjuvant treatment in colorectal cancer. Annals of Oncology 16(4):
4 549-57
- 5 Chau I, Norman A, Cunningham D, et al. (2005b) Longitudinal quality of life and
6 quality adjusted survival in a randomised controlled trial comparing six months of
7 bolus fluorouracil/leucovorin vs. twelve weeks of protracted venous infusion
8 fluorouracil as adjuvant chemotherapy for colorectal cancer. European Journal of
9 Cancer 41(11): 1551-9
- 10 Saini A, Norman A, Cunningham D, et al. (2005) Twelve weeks of protracted venous
11 infusion of fluorouracil (5-FU) is as effective as 6 months of bolus 5-FU and folinic
12 acid as adjuvant treatment in colorectal cancer. British Journal of Cancer (16):88
13 1859-65
- 14 **Glynne-Jones 2017**
- 15 Glynne-Jones R, Counsell N, Quirke P, et al. (2017) Chronicle: results of a
16 randomised phase III trial in locally advanced rectal cancer after neoadjuvant
17 chemoradiation randomising postoperative adjuvant capecitabine plus oxaliplatin
18 (XELOX) versus control. Annals of Oncology 25(7):1356-62
- 19 **IDEA collaboration 2018**
- 20 Grothey A, Sobrero A, Shields A, et al. (2018) Duration of adjuvant chemotherapy for
21 stage III colon cancer. New England Journal of Medicine 378: 1177-88
- 22 **IDEA France trial 2018**
- 23 Andre T, Vernerey D, Mineur L et al. (2018) Three versus 6 months of oxaliplatin-
24 based adjuvant chemotherapy for patients with stage III colon cancer: Disease-free
25 survival results from a randomized, open-label, International Duration Evaluation of
26 Adjuvant (IDEA) France, phase III trial. Journal of Clinical Oncology 36(15): 1469-77
- 27 **McCleary 2013**
- 28 McCleary N, Meyerhardt J, Green E, et al. (2013) Impact of age on the efficacy of
29 newer adjuvant therapies in patients with stage II/III colon cancer: Findings from the
30 ACCENT database. Journal of Clinical Oncology 31(20): 2600-6
- 31 **Robles-Zurita 2018**
- 32 Robles-Zurita J, Boyd K, Briggs A, et al. (2018) SCOT: a comparison of cost-
33 effectiveness from a large randomised phase III trial of two durations of adjuvant
34 Oxaliplatin combination chemotherapy for colorectal cancer. British Journal of Cancer
35 119(11):1332-1338
- 36 **SCOT trial 2018**
- 37 Iveson TJ, Kerr RS, Saunders M, et al. (2018) 3 versus 6 months of adjuvant
38 oxaliplatin-fluoropyrimidine combination therapy for colorectal cancer (SCOT): an
39 international, randomised, phase 3, non-inferiority trial. Lancet Oncology 19(4): 562-
40 78

1 **TOSCA trial 2016**

2 Lonardi S, Sobrero A, Rosati G, et al. (2016) Phase III trial comparing 3-6 months of
3 adjuvant FOLFOX4/XELOX in stage II-III colon cancer: safety and compliance in the
4 TOSCA trial. Annals of Oncology 27(11): 2074-81

1 Appendices

2 Appendix A – Review protocol

3 Review protocol for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer

Field (based on PRISMA-P)	Content
Review question	What is the optimal duration of adjuvant chemotherapy for colorectal cancer?
Type of review question	Intervention
Objective of the review	Previous clinical trials have established 24 weeks of adjuvant chemotherapy to be the standard of care for colorectal cancer. In recent years a shorter duration of adjuvant chemotherapy have been suggested in order to minimise the adverse long-term effects of chemotherapy, mainly neurotoxicity. This review aims to find out what is the optimal duration of adjuvant chemotherapy for colorectal cancer taking into consideration its effects on for example survival and cancer recurrence, neurotoxicity and quality of life.
Eligibility criteria – population/disease/condition/issue/domain	<p>Adults with non-metastatic colorectal cancer after receiving surgery with curative intent</p> <p>Non-metastatic cancer defined as:</p> <ul style="list-style-type: none">• pT_{any} pN1-2• pT3• pT4• M0 <p>Subgroups to be considered separately:</p> <ul style="list-style-type: none">• pT4• pT3/T4 N0 with vascular invasion• right versus left sided tumour• age over 70 years
Eligibility criteria – intervention(s)/exposure(s)/prognostic factor(s)	Adjuvant chemotherapy for less than 24 weeks (6 months), different durations analysed separately
Eligibility criteria – comparator(s)/control or reference (gold) standard	Adjuvant chemotherapy for 24 weeks (6 months) Studies that compare two durations shorter than 24 weeks to each other (for example 12 weeks versus 18 weeks) will also be considered and analysed separately.
Outcomes and prioritisation	<p>Critical outcomes:</p> <ul style="list-style-type: none">• Disease-free survival (minimally important difference [MID]: statistical significance)• Overall survival (MID: statistical significance)• Neuropathy (lasting for 2 years considered permanent) (MID: statistical significance) <p>Important outcomes:</p>

Field (based on PRISMA-P)	Content
	<ul style="list-style-type: none"> • Overall quality of life measured using validated scales (MID: published MIDs from literature) • Distant metastasis (MID: statistical significance) • Treatment-related mortality (MID: statistical significance) • Dose reduction (MID: statistical significance) <p>Quality of life MIDs from the literature:</p> <ul style="list-style-type: none"> • EORTC QLQ-C30: 5 points* • EORTC QLQ-CR29: 5 points* • EORTC QLQ-CR38: 5 points* • EQ-5D: 0.09 using FACT-G quintiles • FACT-C: 5 points* • FACT-G: 5 points* • SF-12: > 3.77 for the mental component summary (MCS) and > 3.29 for the physical component summary (PCS) of the Short Form SF-12 (SF-12) • SF-36: > 7.1 for the physical functioning scale, > 4.9 for the bodily pain scale, and > 7.2 for the physical component summary <p>*Confirmed with guideline committee.</p>
Eligibility criteria – study design	<p>Systematic reviews of randomised controlled trials (RCTs) RCTs</p> <p>Non-randomised studies will not be considered.</p>
Other inclusion/exclusion criteria	<p>Inclusion:</p> <ul style="list-style-type: none"> • English-language • Published full text papers • All settings will be considered that consider medications and treatments available in the UK • Studies published 2000 onwards <p>Studies published 2000 onwards will be considered for this review question because the guideline committee considered that evidence prior to 2000 would not be relevant any longer because the duration of adjuvant chemotherapy for colorectal cancer used to be longer than the current standard of 24 weeks.</p>
Proposed sensitivity/sub-group analysis, or meta-regression	<p>In case of high heterogeneity, the following factors will be considered:</p> <ul style="list-style-type: none"> • Type of chemotherapy
Selection process – duplicate screening/selection/analysis	<p>Sifting, data extraction, appraisal of methodological quality and GRADE assessment will be performed by the systematic reviewer. Resolution of any disputes will be with the senior systematic reviewer and the Topic Advisor. Quality control will be performed by the senior systematic reviewer.</p> <p>Dual sifting will be undertaken for this question for a random 10% sample of the titles and abstracts identified by the search.</p>

Field (based on PRISMA-P)	Content
Data management (software)	<p>Pairwise meta-analyses will be performed using Cochrane Review Manager (RevMan5).</p> <p>'GRADEpro' will be used to assess the quality of evidence for each outcome.</p> <p>NGA STAR software will be used for study sifting, data extraction, recording quality assessment using checklists and generating bibliographies/citations.</p>
Information sources – databases and dates	<p>Potential sources to be searched: Medline, Medline In-Process, CCTR, CDSR, DARE, HTA, Embase</p> <p>Limits (e.g. date, study design):</p> <p>Apply standard animal/non-English language exclusion</p> <p>Limit to RCTs and systematic reviews in first instance, but download all results</p> <p>Dates: from 2000</p>
Identify if an update	Not an update
Author contacts	<p>https://www.nice.org.uk/guidance/indevelopment/gid-ng10060</p> <p>Developer: NGA</p>
Highlight if amendment to previous protocol	For details please see section 4.5 of Developing NICE guidelines: the manual
Search strategy – for one database	For details please see appendix B.
Data collection process – forms/duplicate	For details please see appendix B.
Data items – define all variables to be collected	A standardised evidence table format will be used, and published as appendix D (clinical evidence tables) or H (economic evidence tables).
Methods for assessing bias at outcome/study level	<p>Standard study checklists were used to critically appraise individual studies. For details please see section 6.2 of Developing NICE guidelines: the manual</p> <p>Appraisal of methodological quality:</p> <p>The methodological quality of each study will be assessed using an appropriate checklist:</p> <ul style="list-style-type: none"> • ROBIS for systematic reviews • Cochrane risk of bias tool for RCTs <p>The quality of the evidence for an outcome (i.e. across studies) will be assessed using GRADE.</p> <p>The risk of bias across all available evidence was evaluated for each outcome using an adaptation of the 'Grading of Recommendations Assessment, Development and Evaluation (GRADE) toolbox' developed by the international GRADE working group http://www.gradeworkinggroup.org/</p>
Criteria for quantitative synthesis	For details please see section 6.4 of Developing NICE guidelines: the manual
Methods for quantitative analysis – combining studies and exploring (in)consistency	<p>Synthesis of data:</p> <p>Pairwise meta-analysis of randomised trials will be conducted where appropriate.</p>

Field (based on PRISMA-P)	Content
	<p>When meta-analysing continuous data, final and change scores will be pooled if baselines are comparable. If any studies report both, the method used in the majority of studies will be analysed.</p> <p>Minimally important differences: The guideline committee identified statistically significant differences as appropriate indicators for clinical significance for all outcomes except for quality of life for which published MIDs from literature will be used (see outcomes section for more information).</p>
Meta-bias assessment – publication bias, selective reporting bias	<p>For details please see section 6.2 of Developing NICE guidelines: the manual.</p> <p>If sufficient relevant RCT evidence is available, publication bias will be explored using RevMan software to examine funnel plots.</p>
Confidence in cumulative evidence	For details see sections 6.4 and 9.1 of Developing NICE guidelines: the manual .
Rationale/context – what is known	For details please see the introduction to the evidence review.
Describe contributions of authors and guarantor	<p>A multidisciplinary committee developed the guideline. The committee was convened by The National Guideline Alliance and chaired by Peter Hoskin in line with section 3 of Developing NICE guidelines: the manual.</p> <p>Staff from The National Guideline Alliance undertook systematic literature searches, appraised the evidence, conducted meta-analysis and cost-effectiveness analysis where appropriate, and drafted the guideline in collaboration with the committee. For details please see Supplement 1: methods.</p>
Sources of funding/support	The NGA is funded by NICE and hosted by the Royal College of Obstetricians and Gynaecologists
Name of sponsor	The NGA is funded by NICE and hosted by the Royal College of Obstetricians and Gynaecologists
Roles of sponsor	NICE funds the NGA to develop guidelines for those working in the NHS, public health, and social care in England
PROSPERO registration number	Not registered

1 CCTR: Cochrane Central Register of Controlled Trials; CDSR: Cochrane Database of Systematic
 2 Reviews; DARE: Database of Abstracts of Reviews of Effects; EQ-5D: EuroQol five dimensions
 3 questionnaire; EORTC QLQ-C30: European Organisation for Research and Treatment of Cancer
 4 Quality of Life Questionnaire Core 30 Items; EORTC QLQ-CR29: European Organisation for Research
 5 and Treatment of Cancer Quality of Life Questionnaire colorectal cancer module (29 items); EORTC
 6 QLQ-CR38: European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire
 7 colorectal cancer module (38 items); FACT-C: Functional Assessment of Cancer Therapy questionnaire
 8 (colorectal cancer); FACT-G: Functional Assessment of Cancer Therapy questionnaire (general);
 9 GRADE: Grading of Recommendations Assessment, Development and Evaluation; HTA: Health
 10 Technology Assessment; M0: distant metastasis stage; MCS: mental component summary; MID:
 11 minimal important difference; N: nodal stage; NGA: National Guideline Alliance; NHS: National Health
 12 Service; NICE: National Institute for Health and Care Excellence; PCS: physical component summary;
 13 RCT: randomised controlled trial; RevMan5: Review Manager version 5; ROBIS: a tool for assessing
 14 risk of bias in systematic reviews; SF-12: 12-Item Short Form Survey; SF-36: 36-Item Short Form
 15 Survey; T: tumour stage

1 Appendix B – Literature search strategies

2 Literature search strategies for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

4 Database: Embase/Medline

5 Last searched on: 15/02/2019

#	Searches
1	exp Colorectal Neoplasms/
2	1 use prmz
3	exp colorectal tumor/ or colorectal cancer/
4	3 use oemezd
5	((colorect* or colo rect*) adj3 (cancer* or neoplas* or oncolog* or malignan* or tumo?r* or carcinoma* or adenocarcinoma*).ti,ab.
6	bowel cancer.ti,ab.
7	or/2,4-6
8	exp Antineoplastic Agents/ or exp Antineoplastic Protocols/ or exp Antineoplastic Combined Chemotherapy Protocols/ or exp Organoplatinum Compounds/ or exp Fluorouracil/ or exp Capecitabine/ or exp Chemotherapy, Adjuvant/ or exp Treatment Failure/ or exp Treatment Outcome/
9	8 use prmz
10	exp antineoplastic agent/ or exp clinical protocol/ or exp adjuvant chemotherapy/ or exp platinum complex/ or exp fluorouracil/ or exp capecitabine/ or exp oxaliplatin/ or exp capecitabine plus oxaliplatin/ or exp drug combination/ or exp drug efficacy/ or exp treatment response/ or exp cancer adjuvant therapy/ or exp cancer combination chemotherapy/ or exp treatment failure/ or exp clinical effectiveness/
11	10 use oemezd
12	(adjuvant chemotherap* or 5?FU or fluorouracil or capecitabine or oxaliplatin or XELOX or FOLFOX or FOLFIRI or XELIRI).ti,ab.
13	or/9,11-12
14	7 and 13
15	exp Time Factors/ use prmz
16	exp time factor/ or exp statistical significance/ or exp treatment duration/
17	16 use oemezd
18	(admin* or dose* or dosing* or dosage* or duration* or time* or course* or day* or month* or week*).ti,ab.
19	or/15,17-18
20	14 and 19
21	clinical Trials as topic.sh. or (controlled clinical trial or pragmatic clinical trial or randomized controlled trial).pt. or (placebo or randomi#ed or randomly).ab. or trial.ti.
22	21 use prmz
23	crossover procedure/ or double blind procedure/ or randomized controlled trial/ or single blind procedure/ or (assign* or allocat* or crossover* or cross over* or ((doubt* or singl*) adj blind*) or factorial* or placebo* or random* or volunteer*).ti,ab.
24	23 use oemezd
25	or/22,24
26	20 and 25
27	(conference abstract or letter).pt. or letter/ or editorial.pt. or note.pt. or case report/ or case study/ use oemezd
28	Letter/ or editorial/ or news/ or historical article/ or anecdotes as topic/ or comment/ or case report/ use prmz
29	(letter or comment* or abstracts).ti.
30	or/27-29
31	randomized controlled trial/ use prmz
32	randomized controlled trial/ use oemezd
33	random*.ti,ab.
34	or/31-33
35	30 not 34
36	(animals/ not humans/) or exp animals, laboratory/ or exp animal experimentation/ or exp models, animal/ or exp rodentia/ use prmz
37	(animal/ not human/) or nonhuman/ or exp animal experiment/ or exp experimental animal/ or animal model/ or exp rodent/ use oemezd
38	(rat or rats or mouse or mice).ti.
39	35 or 36 or 37 or 38
40	26 not 39
41	limit 40 to english language
42	limit 41 to yr="2000 -Current"

1 Database: Cochrane Library

2 Last searched on: 15/02/2019

#	Search
1	MeSH descriptor: [Colorectal Neoplasms] explode all trees
2	colorect* or colo rect*
3	#1 or #2
4	MeSH descriptor: [Antineoplastic Agents] explode all trees
5	MeSH descriptor: [Antineoplastic Protocols] explode all trees
6	MeSH descriptor: [Antineoplastic Combined Chemotherapy Protocols] explode all trees
7	MeSH descriptor: [Organoplatinum Compounds] explode all trees
8	MeSH descriptor: [Fluorouracil] explode all trees
9	MeSH descriptor: [Capecitabine] explode all trees
10	MeSH descriptor: [Chemotherapy, Adjuvant] explode all trees
11	MeSH descriptor: [Treatment Failure] explode all trees
12	MeSH descriptor: [Treatment Outcome] explode all trees
13	adjuvant chemotherap* or 5?FU or fluorouracil or capecitabine or oxaliplatin or XELOX or FOLFOX or FOLFIRI or XELIRI
14	#4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or 13
15	MeSH descriptor: [Time Factors] explode all trees
16	admin* or dose* or dosing* or dosage* or duration* or time* or course* or day* or month* or week*
17	#15 or #16
18	#3 and #14 and #17 Publication Year from 2000 to 2018

3


4

1 Appendix C – Clinical evidence study selection

2 **Clinical study selection for review question: What is the optimal duration of
3 adjuvant chemotherapy for colorectal cancer?**

4 **Figure 1: Study selection flow chart**

5



6

1 Appendix D – Clinical evidence tables

2 Clinical evidence tables for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

3 Table 3: Clinical evidence tables

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Full citation Andre, T., Vernerey, D., Mineur, L., Bennouna, J., Desrame, J., Faroux, R., Fratte, S., Hug de Larauze, M., Paget-Bailly, S., Chibaudel, B., Bez, J., Dauba, J., Louvet, C., Lepere, C., Dupuis, O., Becouarn, Y., Mabro, M., Egretteau, J., Bouche, O., Deplanque, G., Ychou, M., Galais, M. P., Ghiringhelli, F., Dourthe, L. M., Bachet, J. B., Khalil, A., Bonnetaire, F., de Gramont, A., Taieb, J., for Prodigie investigators, Gercor Federation Francaise de Cancerologie Digestive, Unicancer, Three Versus 6 Months of Oxaliplatin-Based Adjuvant Chemotherapy for Patients With Stage	Sample size N=2,022 randomised; n=1,008 allocated to the 3-month therapy (n=900 received mFOLFOX6 and n=108 received CAPOX); n=1,014 assigned to the 6-month therapy (n=920 received mFOLFOX6 and n=94 received CAPOX) Characteristics Age in years, mean±SD 3 months 63.9±9.4 6 months 63.9±9.3 Male sex, n (%) 3 months 563 (56) 6 months 581 (58) Tumour stage, n (%) T1 3 months 45 (4) 6 months 33 (3)	Interventions 3 months of adjuvant chemotherapy versus 6 months of adjuvant chemotherapy Type of chemotherapy, n (%) Modified FOLFOX6 (infusional fluorouracil, leucovorin, and oxaliplatin for 6 or 12 cycles) 1,820 (90) CAPOX (capecitabine and oxaliplatin for four or eight cycles) 202 (10) The choice between modified FOLFOX6 and CAPOX was left to the patient and investigator decision.	Details Randomisation and allocation concealment Random allocation was done centrally via web, randomisation was stratified by centre, T stage, N stage, ECOG performance status, and age (<70 years or ≥70 years). Blinding No blinding. Follow-up/outcomes The primary endpoint was disease-free survival, defined as the time from random assignment to relapse or death, whichever occurred first (secondary colorectal cancers were regarded as events in the disease-free survival	Results <u>Outcome: Overall survival (median 4.3 years of follow-up)</u> Whole population 3 months 145 events, n=1,002 6 months 129 events, n=1,008 HR 1.15 95% CI 0.91 to 1.46 Subpopulation who received mFOLFOX6 3 months 131 events, n=895 6 months 118 events, n=914 HR 1.16 95% CI 0.90 to 1.48 Subpopulation who received CAPOX 3 months 14 events, n=107	Limitations Cochrane risk of bias tool Selection bias Random sequence generation: low risk Allocation concealment: unclear (Details not reported.) Performance bias Blinding of participants and personnel: unclear risk (No blinding, but unclear how much it would have an effect on performance.) Detection bias Blinding of outcome assessment: low/high (No blinding. For subjectively measured outcomes there could be a high risk of bias

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
III Colon Cancer: Disease-Free Survival Results From a Randomized, Open-Label, International Duration Evaluation of Adjuvant (IDEA) France, Phase III Trial, J Clin Oncol, 36, 1469-1477, 2018	T2 3 months 76 (8) 6 months 85 (8) T3 3 months 711 (71) 6 months 688 (68) T4 3 months 170 (17) 6 months 202 (20)		outcome, whereas non-colorectal cancers were disregarded in the analysis). Secondary endpoints were overall survival, treatment compliance and toxicities. Peripheral sensory neuropathy was assessed during the whole study period. Residual peripheral sensory neuropathy was defined as the last available measurement for neuropathy toxicity. Maximal neuropathy was defined as the maximum grade observed at any study or follow-up period.	6 months 11 events, n=94 HR 1.08 95% CI 0.49 to 2.37 <u>Outcome: Treatment-related mortality</u> 3 months 7/1,002 6 months 5/1,008	but low risk of objective outcomes.) Attrition bias Incomplete outcome data: low risk Reporting bias Selective reporting: low risk
Ref Id 861108	Node stage, n (%) N0 3 months 1 (0.1) 6 months 2 (0.2) N1 3 months 748 (75) 6 months 753 (75) N2 3 months 253 (25) 6 months 253 (25)			Disease-free survival, neuropathy and dose reduction from the IDEA France trial are reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.	Other bias Other sources of bias: -
Country/ies where the study was carried out France	Tumour and node stage, n (%) T1-2 and N1 3 months 633 (63) 6 months 612 (61) T4 and/or N2 3 months 368 (37) 6 months 396 (39)		Statistical analysis Analysis of the primary and secondary endpoints was performed on the basis of the modified intention-to-treat population (patients who did not receive any therapy whatsoever were excluded from the analysis, otherwise		Other information None

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Source of funding French National Institute of Cancer; the French Ministry of Health by Program Hospitalier de Recherche Clinique 2009; Groupe Coopérateur Multidisciplinaire en Oncologie	Location of colon tumour, n (%) Left 3 months 569 (60) 6 months 592 (61) Right 3 months 377 (39) 6 months 369 (38) Both 3 months 8 (1) 6 months 8 (1) Missing 3 months 48 6 months 39 Inclusion criteria Age \geq 18 years; stage III (according to TNM staging defined by the American Joint Cancer Committee); histologically confirmed colon cancer (tumour location greater than 12 cm from the anal verge by endoscopy and/or above the peritoneal reflection at surgery); had undergone curative intent surgery no more than 8 weeks before randomisation; without		patients were analysed according to original randomisation).		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>microscopic evidence of residual disease; had an ECOG performance status of 0 or 1; had postoperative CEA levels ≤ 10 ng/mL (2 x normal value); and had signed written informed consent.</p> <p>informed consent obtained before any study-specific procedures occurred.</p> <p>Exclusion criteria</p> <p>None reported.</p>				
Full citation Chau, I., Norman, A. R., Cunningham, D., Iveson, T., Hill, M., Hickish, T., Loftis, F., Jodrell, D., Webb, A., Tait, D., Ross, P. J., Shellito, P., Oates, J. R., Longitudinal quality of life and quality adjusted survival in a randomised controlled trial comparing six months of bolus fluorouracil/leucovorin vs. twelve weeks of protracted venous infusion fluorouracil as adjuvant chemotherapy	Sample size See Chau et al. (2005) A randomised comparison between 6 months of bolus fluorouracil/leucovorin and 12 weeks of protracted venous infusion fluorouracil as adjuvant treatment in colorectal cancer. Annals of oncology: official journal of the European society for medical oncology, 16, 549-557	Interventions	Details	Results	Limitations Other information None

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
for colorectal cancer, European Journal of Cancer, 41, 1551-1559, 2005 Ref Id 860893	Inclusion criteria Exclusion criteria				
Country/ies where the study was carried out					
Study type					
Aim of the study					
Study dates					
Source of funding					
Full citation Chau, I, Norman, Ar, Cunningham, D, Tait, D, Ross, Pj, Iveson, T, Hill, M, Hickish, T, Loftis, F, Jodrell, D, Webb, A, Oates, Jr, A randomised comparison between 6 months of bolus fluorouracil/leucovorin and 12 weeks of protracted venous infusion fluorouracil as adjuvant treatment in colorectal cancer,	Sample size N=826 randomised; n=411 allocated to 12 weeks 5-FU infusion; n=415 allocated to 6 months of bolus 5-FU/leucovorin Characteristics Age in years, median (range) 12 weeks 5-FU infusion 63 (27-82)	Interventions 12 weeks of protracted venous infusion of 5-FU versus bolus 5-FU/leucovorin for 6 months Protracted venous infusion of 5-FU was administered at a dose of 300 mg/m ² per day for 12 weeks. Bolus 5-FU 425 mg/m ² and leucovorin 20 mg/m ² were	Details Randomisation and allocation concealment Randomisation was done by an independent randomisation office on 1:1 ratio using random permuted blocks. Randomisation was stratified by treatment centre and in cases of rectal cancer, whether preoperative radiotherapy was given.	Results <u>Outcome: Relapse-free survival (median 65 months of follow-up; event is cancer recurrence or development of metachronous primary colorectal cancer)</u> Whole population 12 weeks 5-FU infusion 104 events, n=397 6 months bolus 5-FU/leucovorin 127 events, n=404	Limitations Cochrane risk of bias tool Selection bias Random sequence generation: low risk Allocation concealment: unclear risk (Details not reported.) Performance bias Blinding of participants and personnel: unclear risk (No blinding,

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Annals of oncology : official journal of the european society for medical oncology, 16, 549-557, 2005	6 months bolus 5-FU/leucovorin 62 (28-95) Male sex, n (%) 12 weeks 5-FU infusion 211 (53)	administered on days 1–5 every 4 weeks for six cycles. (Patients aged >70 years were treated with a reduced starting dose of 370 mg/m ² .)	Blinding No blinding.	HR 0.8 95% CI 0.62 to 1.04, p=0.1	unclear how much lack of blinding could affect performance of clinicians, probably low risk.)
Ref Id 836561	6 months bolus 5-FU/leucovorin 220 (55)	Adjuvant radiotherapy was given to patients with T4 tumour considered to be at high risk of locoregional failure, and was planned to start with the fourth cycle of bolus therapy or after completion of 12 weeks of protracted venous infusion 5-FU, which continued at a reduced dose of 200 mg/m ² until completion of radiotherapy.	Follow-up/outcomes Follow-up was done every 3 months for the first year, every 6 months for the second year and annually thereafter. Serum CEA was measured at baseline and at each clinic visit. Computer tomography scans of the thorax, abdomen and pelvis were performed at baseline, and at 12 and 24 months following initial start of chemotherapy. Colonoscopy was recommended at 12 months after the start of chemotherapy, subsequent colonoscopies was left to the surgeons' discretion.	Subpopulation with rectal cancer 12 weeks 5-FU infusion 39 events, n=156 6 months bolus 5-FU/leucovorin 59 events, n=167 HR 0.63 95% CI 0.43 to 0.94, p=0.0246	Detection bias Blinding of outcome assessment: high/low risk (No blinding). Potentially high risk of bias due to lack of blinding for subjective outcomes but low risk of objective outcomes.)
Country/ies where the study was carried out UK	Site of primary tumour Colon 12 weeks 5-FU infusion 241 (61) 6 months bolus 5-FU/leucovorin 237 (59)			<u>Outcome: Overall survival (median 65 months of follow-up; event is death from any cause)</u>	Attrition bias Incomplete outcome data: low risk
Study type Multicentre RCT	Rectum 12 weeks 5-FU infusion 156 (39) 6 months bolus 5-FU/leucovorin 167 (41)			Whole population 12 weeks 5-FU infusion 99 events, n=397 6 months bolus 5-FU/leucovorin 121 events, n=404 HR 0.79 95% CI 0.61 to 1.03, p=0.083*	Reporting bias Selective reporting: low risk
Aim of the study To compare the efficacy and toxicity of 12 weeks of protracted venous infusion 5-fluorouracil (5-FU) against the standard bolus monthly regimen of 5-FU/leucovorin given for 6 months as adjuvant treatment in colorectal cancer.	Duke's stage (colon cancer) B 12 weeks 5-FU infusion 105 (44) 6 months bolus 5-FU/leucovorin 106 (45)				Other bias Other sources of bias: -
Study dates 1993 to 2003	C 12 weeks 5-FU infusion 135 (56)			Note * calculated values used in the forest plot and corresponding GRADE table are different due to	Other information None
Source of funding					

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Not reported.	<p>6 months bolus 5-FU/leucovorin 131 (55)</p> <p>Duke's stage (rectal cancer) B 12 weeks 5-FU infusion 58 (37)</p> <p>6 months bolus 5-FU/leucovorin 95 (61)</p> <p>C 12 weeks 5-FU infusion 56 (34)</p> <p>6 months bolus 5-FU/leucovorin 109 (65)</p> <p>Radiotherapy received (rectal cancer) Preoperative 12 weeks 5-FU infusion 9 (6)</p> <p>6 months bolus 5-FU/leucovorin 16 (10)</p> <p>Postoperative 12 weeks 5-FU infusion 24 (15)</p> <p>6 months bolus 5-FU/leucovorin 10 (6)</p> <p>Inclusion criteria Curatively resected stage II and III adenocarcinoma of the</p>		<p>cause. Secondary end points were relapse-free survival (defined as the time from the date of randomisation to the date of either cancer recurrence or development of metachronous primary colorectal cancer), toxicity and quality of life. Quality of life was assessed using the European Organization for Research and Treatment of Cancer Quality of Life questionnaire (EORTC QLQ-C30) before randomisation, during adjuvant treatment and during follow-up.</p> <p>Statistical analysis Analysis was on intention-to-treat basis (although in total 25 participants were excluded from the analysis after randomisation due to ineligibility). Survival was analysed using the Kaplan-Meier method and were compared</p>	<p>rounding (0.78 [95% CI 0.60, 1.02])</p> <p>Subpopulation with rectal cancer 12 weeks 5-FU infusion, n=156 (number of events not reported)</p> <p>6 months bolus 5-FU/leucovorin, n=167 (number of events not reported)</p> <p>HR 0.66 95% CI 0.43 to 1.03, p=0.0697</p> <p><u>Outcome: Quality of life - Global health status (QLQ-C30; scale 0-100; higher score indicating better quality of life)</u> Change score from baseline at 24 weeks post-randomisation**</p> <p>12 weeks 5-FU infusion 5.6 (n not reported)</p> <p>6 months bolus 5-FU/leucovorin 2.2 (n not reported)</p> <p>p<0.001</p> <p>Change score from baseline at 2 years post-randomisation*</p> <p>12 weeks 5-FU infusion 9.3 (n not reported)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>colon or rectum; resection margins clear by at least 1 mm; adequate haematological, renal and liver function; no concurrent severe or life-threatening illness.</p> <p>Exclusion criteria Metastatic disease</p>		<p>between treatment groups using the log-rank test, stratified by treatment centre.</p>	<p>6 months bolus 5-FU/leucovorin 9.0 (n not reported) $p=0.999$</p> <p><u>Outcome: Distant metastasis</u> 12 weeks 5-FU infusion 84/397 6 months bolus 5-FU/leucovorin 97/404</p> <p><u>Outcome: Chemotherapy-related mortality</u> 12 weeks 5-FU infusion 0/397 6 months bolus 5-FU/leucovorin 0/404</p> <p><u>Outcome: Dose reduction - percentage of dose received</u> 12 weeks 5-FU infusion 90% 6 months bolus 5-FU/leucovorin 74% $p<0.001$</p> <p><u>Outcome: Dose reduction - percentage of patients with completed treatment</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>(no dose reductions, delayed or interruptions)</p> <p>12 weeks 5-FU infusion 45.7%</p> <p>6 months bolus 5-FU/leucovorin 13.3%</p> <p>*Data extracted from Chau, Norman et al. 2005.</p> <p>** Data extracted from Saini 2003.</p>	
Full citation Grothey, A., Sobrero, A. F., Shields, A. F., Yoshino, T., Paul, J., Taieb, J., Souglakos, J., Shi, Q., Kerr, R., Labianca, R., Meyerhardt, J. A., Verneroy, D., Yamanaka, T., Boukovinas, I., Meyers, J. P., Renfro, L. A., Niedzwiecki, D., Watanabe, T., Torri, V., Saunders, M., Sargent, D. J., Andre, T., Iveson, T., Duration of Adjuvant Chemotherapy for Stage III Colon Cancer, N Engl J Med, 378, 1177-1188, 2018	Sample size N=12,834 randomised; n=6,424 allocated to the 3-month group (n=3,870 received FOLFOX and n=2,554 received CAPOX); n=6,410 allocated to the 6-month group (n=3,893 received FOLFOX and n=2,517 received CAPOX)	Interventions 3 months of adjuvant chemotherapy versus 6 months of adjuvant chemotherapy Chemotherapy regimen, n (%): Capecitabine and oxaliplatin (CAPOX) 5,071 (39.5) Fluorouracil, leucovorin, and oxaliplatin (FOLFOX) 7,763 (60.5)	Details Randomisation and allocation concealment Randomisation method and allocation concealment for individual trials not reported. However, reports from the individual trials give more detail and appropriate methods were applied. Blinding No blinding. Follow-up/outcomes Primary outcome was disease-free survival at	Results <u>Outcome: Disease-free survival at 3 years (median 41.8 months of follow-up)</u> Whole population 3 months 74.6% (95% CI 73.5% to 75.7%) (n=6,424) 6 months 75.5% (95% CI 74.4% to 76.7%) (n=6,410) HR 1.07 95% CI 1.00 to 1.15 Subpopulation who received CAPOX	Limitations Cochrane risk of bias tool Selection bias Random sequence generation: unclear risk (The paper did not report methods of randomisation for individual trials, however, the reports and/or protocols from 3 of the 6 individual trials report sufficient detail to confirm appropriate methods were used.) Allocation concealment: unclear risk (The paper did not report methods of allocation concealment)

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Ref Id 860602 <p>Country/ies where the study was carried out Australia, Canada, Denmark, France, Greece, Italy, Japan, New Zealand, Spain, Sweden, UK, US</p> <p>Study type Pooled analysis from 6 collaborating RCTs (IDEA collaboration): TOSCA (Italy) N=2,402 SCOT (UK, Denmark, Spain, Australia, Sweden, New Zealand) N=3,983 IDEA France N=2,010 CALGB/SWOG (US, Canada) N=2,440 HORG (Greece) N=708 ACHIEVE (Japan) N=1,291</p> <p>Aim of the study To evaluate if 3 months of FOLFOX or CAPOX therapy would be non-inferior to 6 months of therapy in the rate of</p>	<p>Tumour stage, n (%): T1 493 (4) T2 1,197 (9) T3 8,400 (66) T4 2,655 (21) Missing 89 (1)</p> <p>Nodal stage, n (%): N1 9,168 (71) N2 3,567 (28) Missing 99 (1)</p> <p>Risk group, n (%): T1, T2, or T3N1 7,471 (59) T4, N2, or both 5,256 (41)</p> <p>Inclusion criteria Patients with stage III colon cancer (individual trials might have had slightly differing inclusion criteria)</p> <p>Exclusion criteria None reported.</p>		<p>3 years, defined as the time from randomisation to first relapse, the diagnosis of a secondary colorectal cancer after the initial diagnosis, or death from any cause, whichever occurred first.</p> <p>Statistical analysis Modified intention-to-treat analysis was performed (included all the patients who were randomised and had received at least one dose of a trial drug). Cox regression model stratified according to each trial was used to estimate hazard ratios and 95% CI for the comparison of 3 months versus 6 months of adjuvant chemotherapy.</p>	<p>3 months 75.9% (95% CI 74.2% to 77.6%) (n=2,554) 6 months 74.8% (95% CI 73.1% to 76.6%) (n=2,517) HR 0.95 95% CI 0.85 to 1.06</p> <p>Subpopulation who received FOLFOX 3 months 73.6% (95% CI 72.2% to 75.1%) (n=3,870) 6 months 76.0% (95% CI 74.6% to 77.5%) (n=3,893) HR 1.16 95% CI 1.06 to 1.26</p> <p>Subpopulation with T4 cancer 3 months 58.1% 6 months 61.4% HR 1.16 95% CI 1.03 to 1.31</p> <p>Subpopulation with T1-3 N1 cancer (low risk population) 3 months 83.1% (95% CI 81.8% to 84.4%)</p>	for individual trials, however, the reports and/or protocols from 3 of the 6 individual trials report sufficient detail to confirm appropriate methods were used.)

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>disease-free survival at 3 years.</p> <p>Study dates June 2007 to December 2015</p> <p>Source of funding National Cancer Institute; Institute National du Cancer; Programme Hospitalier de Recherche Clinique en Cancérologie; the National Institute for Health Research, Efficacy and Mechanism Evaluation; the National Institute for Health Research; Health Technology Assessment; Cancer Research United Kingdom; the Japanese Foundation for Multidisciplinary Cancer Treatment; L’Agenzia Italiana del Farmaco; the HORG Foundation.</p>				<p>6 months 83.3% (95% CI 82.1% to 84.6%) HR 1.01 95% CI 0.90 to 1.12*</p> <p>Note: * calculated values used in the forest plot and corresponding GRADE table differ due to rounding (1.01 [95% CI 0.9, 1.13])</p> <p>Subpopulation with T1-3 N1 cancer (low risk population) who received CAPOX 3 months 85.0 (95% CI 83.1% to 86.9%) 6 months 83.1% (95% CI 81.1% to 85.2%) HR 0.85 95% CI 0.71 to 1.01</p> <p>Subpopulation with T1-3 N1 cancer (low risk population) who received FOLFOX 3 months 81.9% (95% CI 80.2% to 83.6%) 6 months 83.5% (95% CI 81.9% to 85.1%) HR 1.10 95% CI 0.96 to 1.26</p>	<p>Reporting bias Selective reporting: low risk</p> <p>Other bias Other sources of bias: The inclusion/exclusion criteria as well as the details of the Interventions differ to an extent across the 6 different individual trials pooled in this collaborative paper. A BIT MORE TO BE ADDED ABOUT DIFFERENCES BETWEEN THE TRIALS. The 6 trials are also different in size accounting for unequal amount of weight in the analysis. The SCOT trial is the largest of the trials with almost 4,000 participants, TOSCA and CALGB/SWOG both had around 2,400 participants, IDEA France trial had around 2,000 participants, ACHIEVE trial has around 1,300 participants while</p>

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>Subpopulation with T4 and/or N2 cancer (high risk population)</p> <p>3 months 62.7% (95% CI 60.8% to 64.4%)</p> <p>6 months 64.4% (95% CI 62.6% to 66.4%)</p> <p>HR 1.12 95% CI 1.03 to 1.23</p> <p>Subpopulation with T4 and/or N2 cancer (high risk population) who received CAPOX</p> <p>3 months 64.1% (95% CI 61.3% to 67.1%)</p> <p>6 months 64.0% (95% CI 61.2% to 67.0%)</p> <p>HR 1.02 95% CI 0.89 to 1.17</p> <p>Subpopulation with T4 and/or N2 cancer (high risk population) who received FOLFOX</p> <p>3 months 61.5% (95% CI 58.9% to 64.1%)</p> <p>6 months 64.7% (95% CI 62.2% to 67.3%)</p> <p>HR 1.20 95% CI 1.07 to 1.35</p>	<p>HORG had around 700 participants.</p> <p>Other information None</p>

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><u>Outcome: Grade 3 or 4 peripheral sensory neurotoxicity</u></p> <p>Whole population 3 months 117/6,424 6 months 643/6,410</p> <p>Subpopulation who received CAPOX 3 months 37/2,554 6 months 124/2,517</p> <p>Subpopulation who received FOLFOX 3 months 80/3,870 6 months 519/3,893</p> <p><u>Outcome: Dose reduction - Percentage of dose delivered, mean ± SD</u></p> <p>Subpopulation who received FOLFOX Fluorouracil 3 months 92.4 ± 22.7 (n=3,870) 6 months 81.6 ± 26.6 (n=3,893) Oxaliplatin 3 months 91.4 ± 19.9 (n=3,870) 6</p>	

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>months 72.8 ± 25.6 (n=3,893)</p> <p>Subpopulation who received CAPOX Capecitabine 3 months 91.2 ± 23.5 (n=2,554)</p> <p>6 months 78.0 ± 29.4 (n=2,517)</p> <p>Oxaliplatin 3 months 89.8 ± 21.7 (n=2,554)</p> <p>6 months 69.3 ± 28.3 (n=2,517)</p>	
Full citation Iveson, T. J., Kerr, R. S., Saunders, M. P., Cassidy, J., Hollander, N. H., Tabernero, J., Haydon, A., Glimelius, B., Harkin, A., Allan, K., McQueen, J., Scudder, C., Boyd, K. A., Briggs, A., Waterston, A., Medley, L., Wilson, C., Ellis, R., Essapen, S., Dhadda, A. S., Harrison, M., Falk, S., Raouf, S., Rees, C., Olesen, R. K., Propper, D., Bridgewater, J., Azzabi, A., Farrugia, D., Webb, A., Cunningham,	<p>Sample size N=6,088 randomised; n=3,044 allocated to 3-month therapy; n=3,044 allocated to 6-month therapy</p> <p>Characteristics Age in years, median (IQR) 3 months 65 (58-70) 6 months 65 (58-70)</p> <p>Male sex, n (%) 3 months 1843 (61) 6 months 1844 (61)</p>	<p>Interventions 3 months of adjuvant chemotherapy versus 6 months of adjuvant chemotherapy</p> <p>Type of chemotherapy, n (%) FOLFOX (bolus and infused fluorouracil with oxaliplatin) 3 months 993 (33) 6 months 988 (32) CAPOX (capecitabine and oxaliplatin) 3 months 2,051 (67) 6 months 2,056 (68)</p>	<p>Details Randomisation and allocation concealment Randomisation was done centrally via computer system in 1:1 ratio, stratified by centre, choice of chemotherapy, sex, disease site, N stage and T stage, and if the patient was going to receive CAPOX the starting dose of capecitabine. Once enrolled, patients were randomly assigned via computer programme</p>	<p>Results <u>Outcome: Disease-free survival (median 37 months of follow-up)</u> Subpopulation with rectal cancer 3 months 107 events, n=551 6 months 114 events, n=547 HR 0.926 95% CI 0.711 to 1.205</p> <p><u>Outcome: Overall survival (median 37 months of follow-up)</u></p>	<p>Limitations Cochrane risk of bias tool Selection bias Random sequence generation: low risk Allocation concealment: low risk</p> <p>Performance bias Blinding of participants and personnel: unclear risk (No blinding, unclear how much lack of blinding could affect performance of clinicians, probably low risk.)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
D., Hickish, T., Weaver, A., Gollins, S., Wasan, H. S., Paul, J., 3 versus 6 months of adjuvant oxaliplatin-fluoropyrimidine combination therapy for colorectal cancer (SCOT): an international, randomised, phase 3, non-inferiority trial, Lancet Oncology Lancet Oncol, 19, 562-578, 2018	Disease site, n (%) Colon 3 months 2492 (82) 6 months 2144 (70) Rectum 3 months 552 (28) 6 months 549 (18)	The choice of the type of chemotherapy was not randomised but was chosen by the treating physician and patient. FOLFOX was given every 2 weeks (i.e. 6 or 12 cycles depending on the allocation). IV oxaliplatin 85 mg/m ² was given over 2 hours on the first day concurrently with L-folinic acid 175 mg or folinic acid (leucovorin) 350 mg. This was followed by an IV bolus injection of fluorouracil 400 mg/m ² over 5 minutes, then a continuous IV infusion of fluorouracil 2400 mg/m ² over 46 hours.	and allocated a unique identification number. Blinding No blinding. Follow-up/outcomes The primary outcome was disease-free survival, defined as the time from randomisation (or trial registration for those randomised after 3 months of therapy) to relapse, development of a new colorectal cancer, or death from any cause. Secondary endpoints were overall survival (defined as the time from randomisation (registration for those randomised at 3 months) to death from any cause), safety, quality of life and cost-effectiveness.	3 months 393 events, n=3,035 6 months 394 events, n=3,030 HR 0.994 95% CI 0.964 to 1.143 <u>Outcome: Quality of life - QLQ-C30 global health status (scale 0-100, higher score indicating better quality of life)</u> At 6 months Mean difference in score between groups (3 months minus 6 months) 11.58 (95% CI 9.62 to 13.54), p<0.001 At 12 months Mean difference in score between groups (3 months minus 6 months) 1.48 (95% CI -0.19 to 3.14), p>0.05 <u>Outcome: Quality of life - EQ-5D visual analogue scale</u> At 6 months Mean difference in score between groups (3 months minus 6	Detection bias Blinding of outcome assessment: high/low risk (No blinding). Potentially high risk of bias due to lack of blinding for subjective outcomes but low risk of objective outcomes.) Attrition bias Incomplete outcome data: low risk Reporting bias Selective reporting: low risk Other bias Other sources of bias: -
Ref Id 860452					
Country/ies where the study was carried out UK, Denmark, Spain, Sweden, Australia, and New Zealand					
Study type RCT (SCOT)	3 months 917 (30) 6 months 915 (30)	CAPOX was given every 3 weeks (i.e. 4 or 8 cycles depending on the allocation). IV oxaliplatin 130 mg/m ² was given on the first day over 2 hours. Oral capecitabine 1000 mg/m ² was taken twice	Quality of life was assessed with the European Organisation for Research and Treatment of Cancer (EORTC) QLQ-C30 and QLQ-CR29 and EQ-5D-		
Aim of the study To investigate whether 3 months of oxaliplatin-containing chemotherapy would be	N stage, n (%) N0 3 months 559 (18) 6 months 557 (18) N1 3 months 1731 (57)				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
non-inferior to the usual 6 months of treatment.	6 months 1732 (57) N2 3 months 754 (25) 6 months 755 (25)	per day for the first 14 days of each cycle. (Patients older than 70 years of age could be given 75% of the capecitabine full dose if deemed appropriate depending on the fitness of the patient.)	3L (visual analogue scale and health index). The quality of life questionnaires were administered at baseline and before each treatment cycle, each month in the 3 month group for the first 3 months after treatment, and at months 9 and 12 for the EORTC questionnaires and months 9, 12, 18, and 24, then annually for EQ-5D-3L.	months) 9.80 (SE 1.04), p<0.001 At 12 months Mean difference in score between groups (3 months minus 6 months) 1.45 (SE 0.88), p>0.05	
Study dates March 27 2008 to Nov 29 2013	High risk stage II*, n (%) 3 months 551 (18) 6 months 545 (81)				
Source of funding Medical Research Council; Swedish Cancer Society; Cancer Research UK Core Clinical Trials Unit Funding; NHS Greater Glasgow & Clyde; University of Glasgow	*defined as one or more of the following: T4 disease, tumour obstruction with or without perforation of the primary tumour preoperatively, fewer than ten lymph nodes harvested, poorly differentiated histology, perineural invasion, or extramural venous or lymphatic vascular invasion.		Neuropathy was assessed with the FACT/GOG-Ntx4 questionnaire.	<u>Outcome: Treatment-related mortality</u> 3 months 16/3,035 6 months 16/3,030	Disease-free survival, neuropathy, and dose reduction among people with colon cancer from the SCOT trial are reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.
	Inclusion criteria Aged ≥18 years; had undergone a curative resection for stage III or high-risk stage II* adenocarcinoma of the colon or rectum; WHO performance status 0 or 1; adequate organ		Statistical analysis Analysis was done on intention-to-treat population as much as possible. Kaplan-Meier techniques and Cox proportional hazard ratios were used to analyse disease-free survival and overall survival.		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>function; life expectancy of greater than 5 years with reference to non-cancer related diseases; normal CT scan of the chest, abdomen, and pelvis before study enrolment; carcinoembryonic antigen less than 1.2 times the local upper limit of normal within 1 week before randomisation; patients with rectal cancer had to have undergone total mesorectal excision surgery with negative resection margins (defined as >1 mm clearance).</p> <p>*defined as having one or more of the following: T4 disease, tumour obstruction with or without perforation of the primary tumour preoperatively, fewer than ten lymph nodes harvested, poorly differentiated histology, perineural invasion, or extramural venous or lymphatic vascular invasion</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Exclusion criteria</p> <p>Chemotherapy (except if administered with curative intent and completed >5 years ago and from which there were no residual complications); previous long-course chemoradiotherapy (preoperative short-course radiotherapy alone was allowed); moderate or severe renal impairment (glomerular filtration rate or creatinine clearance <30 mL/min, as calculated with the Cockcroft-Gault equation); haemoglobin less than 9 g/dL; absolute neutrophil count less than 1.5 x 10⁹/L; platelet count <100 x 10⁹/L; aspartate aminotransferase or alanine aminotransferase greater than 2.5 times the upper limit of normal; clinically significant cardiovascular disease; pregnancy or lactation</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>or being of child-bearing potential and not using, or willing to use, medically approved contraception (postmenopausal women must have been amenorrhoeic for at least 12 months to be considered of non-childbearing potential); previous malignancy other than adequately treated in-situ carcinoma of the uterine cervix or basal or squamous cell carcinoma of the skin (unless there had been a disease-free interval of at least 5 years); known or suspected dihydropyrimidine dehydrogenase deficiency.</p>				
Full citation Lonardi, S, Sobrero, A, Rosati, G, Bartolomeo, M, Ronzoni, M, Aprile, G, Massida, B, Scartozzi, M, Banzi, M, Zampino, Mg, Pasini, F, Marchetti, P, Cantore, M, Zaniboni, A, Rimassa, L, Ciuffreda,	Sample size N=3,759 randomised; n=1,870 allocated to 3 months adjuvant chemotherapy; n=1,889 allocated to 6 months adjuvant chemotherapy	Interventions 3-month adjuvant chemotherapy versus 6-month adjuvant chemotherapy Characteristics Type of chemotherapy FOLFOX4 (oxaliplatin and 5-FU) 3 months 64%	Details Randomisation and allocation concealment Randomisation was done centrally at Mario Negri Institute with the use of permuted blocks of variable size, randomisation was	Results <u>Outcome: Treatment-related mortality (within 30 days)</u> 3 months 3/1,820 6 months 7/1,834 Disease-free survival and neuropathy from the TOSCA trial are	Limitations Cochrane risk of bias tool Selection bias Random sequence generation: low risk Allocation concealment: low risk Performance bias

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
L, Ferrari, D, Barni, S, Zagonel, V, Maiello, E, Rulli, E, Labianca, R, Phase III trial comparing 3-6 months of adjuvant FOLFOX4/XELOX in stage II-III colon cancer: safety and compliance in the TOSCA trial, Annals of oncology : official journal of the European Society for Medical Oncology, 27, 2074-2081, 2016	Age in years, mean (range) 3 months 63.4 (21-83) 6 months 63.1 (21-84)	6 months 64% CAPOX (capecitabine and oxaliplatin) 36% 3 month 36% 6 months 36%	stratified by centre and cancer stage. Blinding No blinding.	reported within the pooled analysis of the IDEA collaboration trials in Grothey 2018.	Blinding of participants and personnel: unclear risk (No blinding, unclear how much lack of blinding could affect performance of clinicians, probably low risk.)
Ref Id 859553	Male sex, n (%) 3 months 1,035 (56) 6 months 1,027 (55)	FOLFOX4 therapy was administered as IV infusion of oxaliplatin 85 mg/m ² over 2 hours, concurrently with LV 100 mg/m ² , followed by 5-FU 400 mg/m ² as bolus injection and 5-FU 600 mg/m ² as IV infusion over 22 hours on the first day. On day 2, LV 100 mg/m ² , 5-FU 400mg/m ² bolus injection, and 5-FU 600 mg/m ² IV infusion over 22 hours were administered as previous day. Cycles were repeated every 2 weeks (i.e. either 6 or 12 cycles depending on the allocation).	Follow-up/outcomes Follow-up visits happened every 4 months during the first 3 years after completion of study treatment phase and annually after that. Laboratory assessment and abdominal ultrasound were carried out, and a yearly abdominal CT scan and chest X-ray were required as a minimum.		Detection bias Blinding of outcome assessment: high/low risk (No blinding). Potentially high risk of bias due to lack of blinding for subjective outcomes but low risk of objective outcomes.)
Country/ies where the study was carried out Italy	Stage, n (%) II 3 months 641 (35) 6 months 648 (35)				
Study type RCT (TOSCA)	T stage, n (%) Tx 3 months 7 (0.4) 6 months 5 (0.3)				Attrition bias Incomplete outcome data: low risk
Aim of the study To evaluate the efficacy and safety of a shorter course of treatment in radically resected stage II/III colon cancer patients.	T0 3 months 0 (0) 6 months 2 (0.1)				Reporting bias Selective reporting: low risk
Study dates	T1 3 months 33 (1.8) 6 months 44 (2.4)	CAPOX therapy was administered as IV infusion of oxaliplatin 130 mg/m ² over 2 hours on the first day, followed by oral	Primary outcome was relapse-free survival, defined as time from date of randomisation up to date of first relapse or death from		Other bias Other sources of bias: -
	T2a 3 months 74 (4)				
	T2b 6 months 98 (5)				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
June 2007 to March 2013 Source of funding Agenzia Italiana del Farmaco	3 months 43 (2.4) 6 months 35 (1.9) T3 3 months 1,391 (76) 6 months 1,358 (73) T4 3 months 285 (16) 6 months 313 (17) Unknown 3 months 15 6 months 12 N stage, n (%) Nx 3 months 4 (0.2) 6 months 9 (0.5) N0 3 months 623 (34) 6 months 362 (34) N1 3 months 869 (48) 6 months 895 (48) N2 3 months 4 (0.2) 6 months 3 (0.2) N3 3 months 4 (0.2) 6 months 3 (0.2) Unknown 3 months 21 6 months 15	capecitabine 1000 mg/m ² twice daily on day 1–14. Cycles were repeated every 3 weeks (i.e. 4 or 8 cycles depending on the allocation).	any cause. Secondary outcomes were overall survival and safety. Statistical analysis Analysis for safety outcomes was done on modified intention-to-treat population (defined as all randomised patients without major violations of eligibility criteria).		

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Inclusion criteria</p> <p>Histologically confirmed stage III or high-risk stage II (fulfilling at least 1 of the following criteria: T4 tumour, grade >3, onset with bowel obstruction/perforation, vascular or lymphatic/perineural invasion, <12 nodes examined) colon cancer; age >18 years; curative surgery carried out no less than 3 and no more than 10 weeks before randomisation; ECOG performance status ≤1; signed written informed consent.</p> <p>Exclusion criteria</p> <p>Macroscopic or microscopic evidence of residual tumour; prior cytotoxic chemotherapy/radiotherapy/immunotherapy for colon cancer; other malignancies within the last 5 years; lactating women;</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	history/presence of other dysfunction or clinical laboratory findings suggesting a disease or condition that contraindicates experimental therapy or high risk of treatment complications; chronic daily treatment with high-dose aspirin (>325 mg/day).				
Full citation	Sample size	Interventions	Details	Results	Limitations
Saini, A., Norman, A. R., Cunningham, D., Chau, I., Hill, M., Tait, D., Hickish, T., Iveson, T., Loftis, F., Jodrell, D., Ross, P. J., Oates, J., Twelve weeks of protracted venous infusion of fluorouracil (5-FU) is as effective as 6 months of bolus 5-FU and folinic acid as adjuvant treatment in colorectal cancer, British Journal of Cancer, 88, 1859-1865, 2003	See Chau et al. (2005) A randomised comparison between 6 months of bolus fluorouracil/leucovorin and 12 weeks of protracted venous infusion fluorouracil as adjuvant treatment in colorectal cancer. Annals of oncology: official journal of the european society for medical oncology, 16, 549-557				
Ref Id	Characteristics				Other information
859983	Inclusion criteria				
Country/ies where the study was carried out	Exclusion criteria				
Study type					

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Aim of the study					
Study dates					
Source of funding					


1 5-FU: fluorouracil; CAPOX: capecitabine and oxaliplatin; CEA: carcinoembryonic antigen; CI: confidence interval; CT: computer tomography; ECOG: Eastern Cooperative
 2 Oncology Group; EORTC: European Organisation for Research and Treatment of Cancer; EQ-5D: EuroQol five dimensions; FACT/GOG-Ntx4: Functional Assessment of Cancer
 3 Therapy/Gynecologic Oncology Group Neurotoxicity questionnaire; FOLFOX: folinic acid, fluorouracil and oxaliplatin; HR: hazard ratio; IQR: interquartile range; IV: intravenous; LV:
 4 leucovorin; N: nodal stage; NHS: National Health Service; QLQ-C30: Quality of Life Questionnaire Core 30 Items; QLQ-CR29: Quality of Life Questionnaire colorectal cancer
 5 module (29 items); RCT: randomised controlled trial; SD: standard deviation; SE: standard error; T: tumour stage; TNM: tumour, node and metastasis

6

1 Appendix E – Forest plots


2 Forest plots for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

3 **Figure 2: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 4 **Disease-free survival (follow-up median 3.5 years for all subgroups except for rectal cancer population (follow up median**
 5 **3.1years); event is relapse, diagnosis of secondary colorectal cancer or death from any cause**




1 CAPOX: capecitabine and oxaliplatin; CI: confidence interval; FOLFOX: folinic acid, fluorouracil and oxaliplatin; N: nodal stage; O-E: observed minus expected; T: tumour stage; V: variance

3 **Figure 3: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 4 **Overall survival (follow-up median 3.1 to 4.3 years); event is death from any cause**




5 CAPOX: capecitabine and oxaliplatin; CI: confidence interval; FOLFOX: folinic acid, fluorouracil and oxaliplatin; O-E: observed minus expected; V: variance

1 **Figure 4: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 2 **Grade 3 or 4 peripheral neuropathy (maximal level at any time point after randomisation)**




3 CAPOX: *capecitabine and oxaliplatin*; CI: *confidence interval*; FOLFOX: *folinic acid, fluorouracil and oxaliplatin*; M-H: *Mantel-Haenszel*

5 **Figure 5: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 6 **Quality of life – QLQ-C30 global health status (scale 0-100; better indicated by higher values)**



7 CI: *confidence interval*; IV: *inverse variance*; QLQ-C30: *Quality of Life Questionnaire Core 30 Items*; SE: *standard error*

1 **Figure 6: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 2 **Quality of life – EQ-5D VAS (scale 0-100; better indicated by higher values)**




3 *CI: confidence interval; EQ-5D VAS: EuroQol five dimensions visual analogue scale; IV: inverse variance; SE: standard error*

5 **Figure 7: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 6 **Treatment-related mortality**




7 *CI: confidence interval; M-H: Mantel-Haenszel*

9 **Figure 8: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
 10 **Percentage of oxaliplatin dose delivered (better indicated by higher values)**




1 CAPOX: *capecitabine and oxaliplatin*; *CI*: *confidence interval*; *FOLFOX*: *folinic acid, fluorouracil and oxaliplatin*; *IV*: *inverse variance*; *SD*: *standard deviation*

3 **Figure 9: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
4 **Percentage of fluorouracil dose delivered (FOLFOX subpopulation) (better indicated by higher values)**




5 *CI*: *confidence interval*; *FOLFOX*: *folinic acid, fluorouracil and oxaliplatin*; *IV*: *inverse variance*; *SD*: *standard deviation*

7 **Figure 10: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for non-metastatic colorectal cancer –**
8 **Percentage of capecitabine dose delivered (CAPOX subpopulation) (better indicated by higher values)**




9 CAPOX: *capecitabine and oxaliplatin*; *CI*: *confidence interval*; *IV*: *inverse variance*; *SD*: *standard deviation*

1 **Figure 11: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Disease-free survival**
 2 (follow-up median 5.4 years for whole population, median 4.6 years for rectal cancer population); event is cancer recurrence or
 3 development of metachronous primary colorectal cancer



4 *CI: confidence interval; O-E: observed minus expected; V: variance*

5 **Figure 12: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Overall survival**
 6 (follow-up median 5.4 years for whole population, median 4.6 years for rectal cancer population)); event is cancer recurrence
 7 or development of metachronous primary colorectal cancer



Footnotes


(1) Number of events not reported.

9

10


11 *CI: confidence interval; O-E: observed minus expected; V: variance*

1 **Figure 13: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Quality of life – QLQ-**
 2 **C30 global health status (scale 0-100; better indicated by higher values)**

Footnotes(1) SE calculated using p-value 0.0009 ($p < 0.001$ reported)


3 CI: confidence interval; IV: inverse variance; QLQ-C30: Quality of Life Questionnaire Core 30 items; SE: standard error
 4

5 **Figure 14: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Distant metastasis**
 6 **(follow-up median 5.4 years)**




7 CI: confidence interval; M-H: Mantel-Haenszel
 8

1 **Figure 15: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Chemotherapy-**
 2 **related mortality**



3 CI: confidence interval; M-H: Mantel-Haenszel
 4

5 **Figure 16: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and leucovorin – Percentage of dose**
 6 **delivered (Better indicated by higher values)**



Footnotes

(1) SE calculated using p-value 0.0009 ($p < 0.001$ reported)

7 CI: confidence interval; IV: inverse variance; SE: standard error
 8

1 Appendix F – GRADE tables

2 GRADE tables for review question: What is the optimal duration of adjuvant chemotherapy for non-metastatic colorectal cancer?

4 **Table 4: Clinical evidence profile: Comparison 1: 3 months versus 6 months of oxaliplatin-based adjuvant chemotherapy for colorectal cancer**

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Disease-free survival - Whole population (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	6424 (number of events not reported, 3263 across both arms)	6410 (number of events not reported, 3263 across both arms)	HR 1.07 (1.00 to 1.15)	At 3 years for 6-month arm 75.5% ¹ , for 3-month arm 74.6% (73.5% to 75.7%)	HIGH	CRITICAL
Disease-free survival - Subpopulation who received CAPOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	2554 (number of events not reported, 1299 across both arms)	2517 (number of events not reported, 1299 across both arms)	HR 0.95 (0.85 to 1.06)	At 3 years for 6-month arm 74.8% ¹ , for 3-month arm 75.9% (74.2% to 77.6%)	HIGH	CRITICAL
Disease-free survival - Subpopulation who received FOLFOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	3870 (number of events not reported, 1964 across both arms)	3893 (number of events not reported, 1964 across both arms)	HR 1.16 (1.06 to 1.27)	At 3 years for 6-month arm 76.0% ¹ , for 3-month arm 73.6% (72.2% to 75.1%)	HIGH	CRITICAL

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Disease-free survival - Subpopulation with stage T4 cancer (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported (2655 patients with 1075 events across both arms)	Not reported (2655 patients with 1075 events across both arms)	HR 1.16 (1.03 to 1.31)	At 3 years for 6-month arm 61.4%, for 3-month arm 56.8% (52.8% to 60.5%) ²	HIGH	CRITICAL
Disease-free survival - Subpopulation with stage T1-3 N1 (low risk) cancer (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported (7471 patients with 1313 events across both arms)	Not reported (7471 patients with 1313 events across both arms)	HR 1.01 (0.91 to 1.13)	At 3 years for 6-month arm 83.3% ¹ , for 3-month arm 83.1% (81.8% to 84.4%)	HIGH	CRITICAL
Disease-free survival - Subpopulation with stage T1-3 N1 cancer who received CAPOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	HR 0.85 (0.71 to 1.01)	At 3 years for 6-month arm 83.1% ¹ , for 3-month arm 85.0% (83.1% to 86.9%)	HIGH	CRITICAL
Disease-free survival - Subpopulation with stage T1-3 N1 (low risk) cancer who received FOLFOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	HR 1.10 (0.96 to 1.26)	At 3 years for 6-month arm 83.5% ¹ , for 3-month arm 81.9% (80.2% to 83.6%)	HIGH	CRITICAL

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Disease-free survival - Subpopulation with stage T4 and/or N2 (high risk) cancer (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported (5256 patients with 1935 events across both arms)	Not reported (5256 patients with 1935 events across both arms)	HR 1.12 (1.02 to 1.22)	At 3 years for 6-month arm 64.4% ¹ , for 3-month arm 62.7% (60.8% to 64.4%)	HIGH	CRITICAL
Disease-free survival - Subpopulation with stage T4 and/or N2 (high risk) cancer who received CAPOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	HR 1.02 (0.89 to 1.17)	At 3 years for 6-month arm 64.0% ¹ , for 3-month arm 64.1% (61.3% to 67.1%)	HIGH	CRITICAL
Disease-free survival - Subpopulation with stage T4 and/or N2 (high risk) cancer who received FOLFOX (follow-up median 3.5 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	HR 1.20 (1.07 to 1.35)	At 3 years for 6-month arm 64.7% ¹ , for 3-month arm 61.5% (58.9% to 64.1%)	HIGH	CRITICAL
Disease-free survival - Subpopulation with rectal cancer (follow-up median 3.1 years; event is relapse, diagnosis of secondary colorectal cancer or death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ³	none	107/551 (19.4%)	114/547 (20.8%)	HR 0.93 (0.71 to 1.21)	At 3 years for 6-month arm 77% ⁴ , for 3-month arm 78.5% (73% to 83%)	MODERATE	CRITICAL

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Overall survival - Whole population (follow-up median 3.1 to 4.3 years; event is death from any cause)												
2	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	538/4037 (13.3%)	523/4038 (13%)	HR 1.01 (0.93 to 1.10)	At 3 years for 6-month arm 90.0% ⁵ , for 3-month arm 89.9% (89.1% to 90.7%)	HIGH	CRITICAL
Overall survival - Subpopulation who received CAPOX (follow-up median 4.3 years; event is death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ³	none	14/107 (13.1%)	11/94 (11.7%)	HR 1.08 (0.49 to 2.38)	At 3 years for 6-month arm 89% ⁶ , for 3-month arm 92% (85% to 96%)	MODERATE	CRITICAL
Overall survival - Subpopulation who received FOLFOX (follow-up median 4.3 years; event is death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ³	none	131/895 (14.6%)	118/914 (12.9%)	HR 1.16 (0.90 to 1.49)	At 3 years for 6-month arm 93% ⁶ , for 3-month arm 91% (89% to 93%)	MODERATE	CRITICAL
Grade 3 or 4 peripheral neuropathy - Whole population (maximal level at any time point after randomisation)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	117/6424 (1.8%)	643/6410 (10%)	RR 0.18 (0.15 to 0.22)	82 fewer per 1000 (from 78 fewer to 85 fewer)	MODERATE	CRITICAL
Grade 3 or 4 peripheral neuropathy - Subpopulation who received CAPOX (maximal level at any time point after randomisation)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	37/2554 (1.4%)	124/2517 (4.9%)	RR 0.29 (0.20 to 0.42)	35 fewer per 1000 (from 29 fewer to 39 fewer)	MODERATE	CRITICAL

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Grade 3 or 4 peripheral neuropathy - Subpopulation who received FOLFOX (maximal level at any time point after randomisation)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	80/3870 (2.1%)	519/3893 (13.3%)	RR 0.16 (0.12 to 0.20)	112 fewer per 1000 (from 107 fewer to 117 fewer)	MODERATE	CRITICAL
Quality of life - global health status (QLQ-C30) – Change from baseline at 6 months (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	-	MD 11.58 higher (9.62 to 13.54 higher)	MODERATE	IMPORTANT
Quality of life - global health status (QLQ-C30) – Change from baseline at 12 months (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	-	MD 1.48 higher (0.19 lower to 3.15 higher)	MODERATE	IMPORTANT
Quality of life - EQ-5D VAS - Change from baseline at 6 months (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision ⁸	none	Not reported	Not reported	-	MD 9.80 higher (7.76 to 11.84 higher)	MODERATE	IMPORTANT
Quality of life - EQ-5D VAS - Change from baseline at 12 months (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁷	no serious inconsistency	no serious indirectness	no serious imprecision	none	Not reported	Not reported	-	MD 1.45 higher (0.27 lower to 3.17 higher)	MODERATE	IMPORTANT
Distant metastasis												
0	No evidence available	-	-	-	-	-	-	-	-	-	-	IMPORTANT
Treatment-related mortality												
3	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ³	none	26/5857 (0.44%)	28/5872 (0.48%)	RR 0.93 (0.55 to 1.58)	0 fewer per 1000 (from 2 fewer to 3 more)	MODERATE	IMPORTANT

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months	6 months	Relative (95% CI)	Absolute		
Percentage of oxaliplatin dose delivered - Subpopulation who received CAPOX (Better indicated by higher values)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	2554	2517	-	MD 20.60 higher (19.21 to 21.99 higher)	HIGH	IMPORTANT
Percentage of oxaliplatin dose delivered - Subpopulation who received FOLFOX (Better indicated by higher values)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	3870	3893	-	MD 18.60 higher (17.58 to 19.62 higher)	HIGH	IMPORTANT
Percentage of fluorouracil dose delivered (FOLFOX subpopulation) (Better indicated by higher values)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	3870	3893	-	MD 10.80 higher (9.70 to 11.90 higher)	HIGH	IMPORTANT
Percentage of capecitabine dose delivered (CAPOX subpopulation) (Better indicated by higher values)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	2554	2517	-	MD 13.20 higher (11.73 to 14.67 higher)	HIGH	IMPORTANT

1 CAPOX: capecitabine and oxaliplatin; CI: confidence interval; EQ-5D VAS: EuroQoL five dimensions questionnaire visual analogue scale; FOLFOX: folinic acid, fluorouracil and oxaliplatin; HR: hazard ratio; MD: mean difference; N: nodal stage; QLQ-C30: Quality of life questionnaire Core 30 items; RR: relative risk; T: tumour stage

2 From IDEA Collaboration (Grothey 2018).

3 Calculated from the control group survival and HR.

4 Quality of evidence downgraded by 1 because of imprecision of the effect estimate (less than 300 events).

5 Estimated from the whole population in the SCOT trial (Iveson 2018).

6 Estimated from SCOT trial (Iveson 2018) and IDEA France trial (Andre 2018).

7 From IDEA France trial (Andre 2018).

8 Quality of evidence downgraded by 1 because of no of blinding.

9 Imprecision estimated based on the MID for EQ-5D VAS scale 0-100 being 5.

1 **Table 5: Clinical evidence profile: Comparison 2: 3 months of fluorouracil infusion versus 6 months of bolus fluorouracil and
2 leucovorin**

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months fluorouracil infusion	6 months bolus fluorouracil/leucovorin	Relative (95% CI)	Absolute		
Disease-free survival - Whole population (follow-up median 5.4 years; event is cancer recurrence or development of metachronous primary colorectal cancer)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ¹	none	104/397 (26.2%)	127/404 (31.4%)	HR 0.80 (0.62 to 1.04)	At 5 years for 6-month arm 66.7% ² , for 3-month arm 73.3% (68.4% to 77.6%) ³	MODERATE	CRITICAL
Disease-free survival - Subpopulation with rectal cancer (follow-up median 4.6 years; event is cancer recurrence or development of metachronous primary colorectal cancer)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ¹	none	39/156 (25%)	59/167 (35.3%)	HR 0.63 (0.43 to 0.93)	At 5 years for 6-month arm 57.7% ² , for 3-month arm 74% (65.5% to 80.7%)	MODERATE	CRITICAL
Overall survival - Whole population (follow-up median 5.4 years; event is death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ¹	none	99/397 (24.9%)	121/404 (30%)	HR 0.78 (0.60 to 1.02)	At 5 years for 6-month arm 71.5% ² , for 3-month arm 75.7% (70.8% to 79.9%)	MODERATE	CRITICAL
Overall survival - Subpopulation with rectal cancer (follow-up median 4.6 years; event is death from any cause)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ¹	none	156	167	HR 0.66 (0.43 to 1.02)	At 5 years for 6-month arm 65.3% ² , for 3-month arm 78.8% (70.2% to 85.1%) ³	MODERATE	CRITICAL
Neuropathy												
0	No evidence available	-	-	-	-	-	-	-	-	-	-	CRITICAL

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	3 months fluorouracil infusion	6 months bolus fluorouracil/leucovorin	Relative (95% CI)	Absolute		
Quality of life - global health status (QLQ-C30) - Change from baseline at 24 weeks post-randomisation (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁴	no serious inconsistency	no serious indirectness	serious ⁵	none	Not reported	Not reported	-	MD 3.40 higher (1.40 to 5.40 higher)	LOW	IMPORTANT
Quality of life - global health status (QLQ-C30) - Change from baseline at 2 years post-randomisation (range of scores: 0-100; Better indicated by higher values)												
1	randomised trials	serious ⁴	no serious inconsistency	no serious indirectness	very serious ⁶	none	Not reported	Not reported	-	MD 0.30 higher (468.59 lower to 469.19 higher)	VERY LOW	IMPORTANT
Distant metastasis (follow-up median 5.4 years)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	serious ¹	none	84/397 (21.2%)	97/404 (24%)	RR 0.88 (0.68 to 1.14)	29 fewer per 1000 (from 77 fewer to 34 more)	MODERATE	IMPORTANT
Chemotherapy-related mortality												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	Not estimable ⁷	none	0/397 (0%)	0/404 (0%)	RD 0.00 (-0.00, 0.00)	- ⁷	HIGH	IMPORTANT
Percentage of dose delivered (Better indicated by higher values)												
1	randomised trials	no serious risk of bias	no serious inconsistency	no serious indirectness	no serious imprecision	none	342	350	-	MD 16.00 higher (6.60 to 25.40 higher)	HIGH	IMPORTANT

¹ CI: confidence interval; HR: hazard ratio; MD: mean difference; QLQ-C30: Quality of Life Questionnaire core 30 items; RD: risk difference; RR: relative risk

² Quality of evidence was downgraded by 1 because of imprecision of the effect estimate (less than 300 events).

³ From Chau 2005a.

⁴ Although there appears to be difference at 5 years, as reported in the paper, overall the HR was not significantly different so there was unlikely to be a clinically important difference.

⁵ Quality of evidence was downgraded by 1 because of no blinding.

⁶ Quality of evidence was downgraded by 1 because the 95% CI of the absolute effect crosses 1 MID.

⁷ Quality of evidence was downgraded by 2 because the 95% CI of the absolute effect crosses 2 MIDs.

⁹ Not estimable because of 0 events in both arms.

1 Appendix G – Economic evidence study selection

2 Economic evidence study selection for review question: What is the 3 optimal duration of adjuvant chemotherapy for colorectal cancer?

4 A global search of economic evidence was undertaken for all review questions in this
5 guideline. See Supplement 2 for further information.

1 Appendix H – Economic evidence tables

2 Economic evidence tables for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

4 Table 6: Economic evidence tables for the length of adjuvant treatment for colorectal cancer

Study details	Treatment strategies	Study population, design and data sources	Results	Comments
Author & year: Robles-Zurita 2018 Country: United Kingdom Type of economic analysis: With-in Trial Cost Utility Analysis (CUA) Source of funding: The study was funded by the Medical Research	Interventions in detail: Intervention: 3 months of either oxaliplatin-containing adjuvant chemotherapy (CAPOX or FOLFOX). Comparator: 6 months of either oxaliplatin-containing adjuvant chemotherapy (CAPOX or FOLFOX). The choice of either CAPOX or FOLFOX was decided by the doctor and patient prior to randomisation.	Population characteristics: Patients with fully resected high-risk stage II or stage III colorectal cancer suitable for adjuvant therapy. The mean age of the cohort was 64 years old and was 67% male. Modelling approach: With-in trial economic evaluation Source of base-line and effectiveness data: All effectiveness data was taken from the SCOT trial (Iveson 2018) reported in detail in the clinical evidence review. Source of cost data:	CAPOX Total cost 3 month regimen: £17,650 Total cost 6 month regimen £21,503 Total QALYs 3 month regimen: 5.34 Total QALYs 6 month regimen 5.16 Incremental Costs -£3,853 Incremental QALYs 0.19	Perspective: Third-party payer perspective – UK NHS and Personal Social Services. Currency: UK pound sterling (£). Cost year: 2016 Time horizon: 3-8 years as per the SCOT trial Discounting: 3.5% per year for both costs and QALYs

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study details	Treatment strategies	Study population, design and data sources	Results	Comments
Council and Cancer Research UK		<p>Resource use was collected at the patient level. All treatment, follow-up and other medical costs were collected during the follow-up period of the trial. (3-8 years)</p> <p>Chemotherapy and other drugs were costed using the British National Formulary. Direct and non-direct hospitalisation costs were obtained from Information Services Division (ISD) of the National Health Service Scotland.</p> <p>Source of QoL data:</p> <p>A subsample of 1832 patients were given the EQ-5D-3L at baseline and all study follow-up appointments. The responses were scored using the UK general population tariffs.</p>	<p>ICER 3 month regimen dominant</p> <p>FOLFOX</p> <p>Total cost 3 month regimen: £19,641</p> <p>Total cost 6 month regimen £26,483</p> <p>Total QALYs 3 month regimen: 5.21</p> <p>Total QALYs 6 month regimen 5.33</p> <p>Incremental Costs -£6,841</p> <p>Incremental QALYs -0.12</p> <p>ICER £57,008 per QALY (as both incremental costs and incremental QALYs are negative a higher QALY favours the intervention (3 month regimen) as it</p>	<p>Applicability: The study was deemed <i>directly applicable</i> to the decision problem.</p> <p>Limitations: The study was deemed to have only minor methodological limitations. The time horizon was potentially too short to capture all key cost and health differences between the groups and no attempt was made to model beyond the time horizon of the trial. There may particularly be large differences between the groups on the use of expensive palliative treatments.</p> <p>Other comments: The 'headline results reported in the study were for combined groups for FOLFOX and CAPOX. As the</p>

Study details	Treatment strategies	Study population, design and data sources	Results	Comments
			<p>represents a cost saving per QALY forgone)</p> <p>Probabilistic sensitivity analysis: Probabilistic sensitivity analysis was conducted. At the NICE threshold of £30,000 per QALY, the 3 month regimen had a probability of being cost effective of 99.9% for CALPOX and 77.2% for FOLFOX. For £20,000 per QALY threshold (read from the reported CEACs) the probabilities were greater than 99.9% and 90% respectively.</p>	committee found this combined analysis of limited use the disaggregated results reported in the supplementary material are presented here.

1 CAPOX: capecitabine and oxaliplatin
 2 CEAC: Cost Effectiveness Acceptability Curve; FOLFOX: folinic acid, fluorouracil and oxaliplatin ICER: incremental cost effectiveness ratio; QALY: quality adjusted life year; QoL: quality of life

1 Appendix I – Economic evidence profiles

2 Economic evidence profiles for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?

4 Table 7: Economic evidence profiles for the length of adjuvant treatment for colorectal cancer

Study	Population	Comparators	Costs	Effects	Incr costs	Incr effects	ICER	Uncertainty	Applicability and limitations
Robles-Zurita 2018	Patients with fully resected high-risk stage II or stage III colorectal cancer suitable for adjuvant therapy.	CAPOX							
	6 month	£21,503	5.16 QALYs	Reference					Probabilistic sensitivity analysis was conducted. At the NICE threshold of £30,000 per QALY, the 3 month regimen had a probability of being cost effective of 99.9% for CALPOX and 77.2% for FOLFOX. For £20,000 per QALY threshold (read from the reported CEACs) the probabilities were greater than 99.9% and 90% respectively.
	3 month	£17,650	5.34 QALYs	-£3,853	0.19 QALYs	Dominant			The study was deemed <i>directly applicable</i> to the decision problem.
	FOLFOX								
	6 month	£26,483	5.33	Reference					The study was deemed to have only minor methodological limitations. The time horizon was potentially too short to capture all key cost and health differences between the groups and no attempt was made to model beyond the time horizon of the trial. There may particularly be large differences between the groups on the use of expensive palliative treatments.
	3 month	£19,641	5.21 QALYs	-£6,841	-0.12 QALYs	£57,008 per QALY			

DRAFT FOR CONSULTATION

Optimal duration of adjuvant chemotherapy for colorectal cancer

Study	Population	Comparators	Costs	Effects	Incr costs	Incr effects	ICER	Uncertainty	Applicability and limitations
1 2									
Notes: The 'headline results reported in the study were for combined groups for FOLFOX and CAPOX. As the committee found this combined analysis of limited use the disaggregated results reported in the supplementary material are presented here.									
CAPOX: capecitabine and oxaliplatin CEAC: Cost Effectiveness Acceptability Curve; FOLFOX: folinic acid, fluorouracil and oxaliplatin ICER: incremental cost effectiveness ratio; Incr: Incremental QALY: quality adjusted life year; QoL: quality of life									

1 Appendix J – Economic analysis

2 Economic analysis for review question: What is the optimal duration of 3 adjuvant chemotherapy for colorectal cancer?

4 No economic analysis was conducted for this review question.

5

1 Appendix K – Excluded studies

2 Excluded clinical studies for review question: What is the optimal duration 3 of adjuvant chemotherapy for colorectal cancer?

4 Table 8: Excluded studies and reasons for their exclusion

Study	Reason for exclusion
Chau, I., Norman, A. R., Cunningham, D., Tait, D., Ross, P. J., Iveson, T., Hill, M., Hickish, T., Loftis, F., Jodrell, D., Webb, A., Oates, J. R. A randomised comparison between 6 months of bolus fluorouracil/leucovorin and 12 weeks of protracted venous infusion fluorouracil as adjuvant treatment in colorectal cancer. Ann Oncol, 549-557, 2005	Duplicate
Des Guetz, G., Uzzan, B., Morere, J. F., Perret, G., Nicolas, P., Duration of adjuvant chemotherapy for patients with non-metastatic colorectal cancer, Cochrane database of systematic reviews (Online), CD007046, 2010	Cochrane systematic review from 2010. None of the included studies are relevant for to this review because they compare longer durations
Haller, Dg, Catalano, Pj, Macdonald, Js, O'Rourke, Ma, Frontiera, Ms, Jackson, Dv, Mayer, Rj, Phase III study of fluorouracil, leucovorin, and levamisole in high-risk stage II and III colon cancer: final report of Intergroup 0089, Journal of clinical oncology, 23, 8671-8678, 2005	Intervention and comparison not according to review protocol
Hirata, K, Nakahara, S, Shimokobe, T, Imamura, T, Sakamoto, Y, Hirano, T, Abe, R, Kuroki, N, Konomi, K, Kato, H, Fujiwara, H, Fukuyama, N, Hotokezaka, M, Miyazaki, Y, Terasaka, R, Shiraishi, M, Miyazaki, R, Iwashita, A, Nakano, S, Ito, H, A randomized controlled trial of postoperative adjuvant chemotherapy for colorectal cancer-optimal duration of the treatment, Gan to kagaku ryoho. Cancer & chemotherapy, 36, 77-82, 2009	Full text in Japanese
Ito, K, Okushiba, S, Morikawa, T, Kondo, S, Katoh, H, Appropriate duration of postoperative oral adjuvant chemotherapy with HCFU for colorectal cancer, Gan to kagaku ryoho. Cancer & chemotherapy, 31, 55-59, 2004	Full text in Japanese
Ito, K., Kato, T., Koike, A., Miura, K., Yamaguchi, A., Sakou, T., Takagi, H., Optimum duration of oral adjuvant chemotherapy of HCFU for colorectal cancer; review of 5-year follow-up, Anticancer Research, 20, 4681-4686, 2000	This study compares adjuvant chemotherapy duration of 3 months to 18 months, not relevant according to the protocol
Iveson, T, Kerr, R, Saunders, Mp, Hollander, Nh, Tabernero, J, Haydon, Am, Glimelius, B, Harkin, A, Scudder, C, Boyd, K, Waterston, Am, Medley, Lc, Wilson, C, Ellis, R, Essapen, S, Dhadda, As, Harrison, M, Falk, S, Raouf, S, Paul, J, Final DFS results of the SCOT study: an International Phase III Randomised (1: 1) Noninferiority Trial comparing 3 versus 6 months of oxaliplatin based adjuvant chemotherapy for colorectal cancer, Journal of clinical oncology. Conference: 2017 annual meeting of the american society of clinical oncology, ASCO. United states, 35, 2017	Conference abstract

DRAFT FOR CONSULTATION
Optimal duration of adjuvant chemotherapy for colorectal cancer

Study	Reason for exclusion
Iveson, T., Kerr, R., Saunders, M., Hollander, N., Tabernero, J., Haydon, A., Glimelius, B., Harkin, A., Scudder, C., Boyd, K., Waterston, A., Medley, L., Wilson, C., Ellis, R., Essapen, S., Dhadda, A., Harrison, M., Falk, S., Abdel-Raouf, S., Paul, J., Updated results of the SCOT study: An international phase III randomised (1:1) non-inferiority trial comparing 3 versus 6 months of oxaliplatin based adjuvant chemotherapy for colorectal cancer, Annals of Oncology, 28 (Supplement 5), v613, 2017	Conference abstract
Jonker, D. J., Spithoff, K., Maroun, J., Gastrointestinal Cancer Disease Site Group of Cancer Care Ontario's Program in Evidence-based, Care, Adjuvant systemic chemotherapy for Stage II and III colon cancer after complete resection: an updated practice guideline, Clinical Oncology (Royal College of Radiologists)Clin Oncol (R Coll Radiol), 23, 314-22, 2011	A practice guideline and systematic review. No relevant comparisons
Nakamura, T., Ohno, M., Tabuchi, Y., Kamigaki, T., Fujii, H., Yamagishi, H., Kuroda, Y., Kansai Carmofur Study, Group, Optimal duration of oral adjuvant chemotherapy with Carmofur in the colorectal cancer patients: the Kansai Carmofur Study Group trial III, International Journal of Oncology, 19, 291-8, 2001	This study compares adjuvant chemotherapy duration of 6 months to 12 months, not relevant according to the protocol
Neugut, A. I., Matasar, M., Wang, X., McBride, R., Jacobson, J. S., Tsai, W. Y., Grann, V. R., Hershman, D. L., Duration of adjuvant chemotherapy for colon cancer and survival among the elderly, Journal of Clinical Oncology, 24, 2368-75, 2006	Observational study
Sadahiro, S., Tsuchiya, T., Sasaki, K., Kondo, K., Katsumata, K., Nishimura, G., Kakeji, Y., Baba, H., Sato, S., Koda, K., Yamaguchi, Y., Morita, T., Matsuoka, J., Usuki, H., Hamada, C., Kodaira, S., Randomized phase III trial of treatment duration for oral uracil and tegafur plus leucovorin as adjuvant chemotherapy for patients with stage IIB/III colon cancer: final results of JFMC33-0502, Annals of oncology : official journal of the european society for medical oncology, 26, 2274-2280, 2015	This study compares adjuvant chemotherapy duration of 6 months to 18 months, not relevant according to the protocol
Sobrero, A., Lonardi, S., Rosati, G., Di Bartolomeo, M., Ronzoni, M., Pella, N., Scartozzi, M., Banzi, M., Zampino, M. G., Pasini, F., Marchetti, P., Cantore, M., Zaniboni, A., Rimassa, L., Ciuffreda, L., Ferrari, D., Zagonel, V., Maiello, E., Barni, S., Rulli, E., Labianca, R., Tosca Investigators, FOLFOX or CAPOX in Stage II to III Colon Cancer: Efficacy Results of the Italian Three or Six Colon Adjuvant Trial, J Clin Oncol, 36, 1478-1485, 2018	This trial (TOSCA trial, part of the IDEA collaboration) has been included in the review but this publication does not report any additional outcomes (see Lonardi 2016 and Grothey 2018)
Suto, T., Ishiguro, M., Hamada, C., Kunieda, K., Masuko, H., Kondo, K., Ishida, H., Nishimura, G., Sasaki, K., Morita, T., Hazama, S., Maeda, K., Mishima, H., Ike, H., Sadahiro, S., Sugihara, K., Okajima, M., Saji, S., Sakamoto, J., Tomita, N., Preplanned safety analysis of the JFMC37-0801 trial: a randomized phase III study of six months versus twelve months of capecitabine as adjuvant chemotherapy for stage III colon cancer.[Erratum appears in Int J Clin Oncol. 2017]	This study compares adjuvant chemotherapy duration of 6 months to 12 months, not relevant according to the protocol

Study	Reason for exclusion
Aug;22(4):805-806; PMID: 28608229], International Journal of Clinical Oncology, 22, 494-504, 2017	
Tsuchiya, T., Sadahiro, S., Sasaki, K., Kondo, K., Katsumata, K., Nishimura, G., Kakeji, Y., Baba, H., Morita, T., Koda, K., Sato, S., Matsuoka, J., Yamaguchi, Y., Usuki, H., Hamada, C., Kodaira, S., Saji, S., Safety analysis of two different regimens of uracil-tegafur plus leucovorin as adjuvant chemotherapy for high-risk stage II and III colon cancer in a phase III trial comparing 6 with 18 months of treatment: JFMC33-0502 trial, Cancer Chemotherapy & PharmacologyCancer Chemother Pharmacol, 73, 1253-61, 2014	This study compares adjuvant chemotherapy duration of 6 months to 18 months, not relevant according to the protocol
You, K. Y., Huang, R., Yu, X., Liu, Y. M., Gao, Y. H., Is it possible to shorten the duration of adjuvant chemotherapy for locally advanced rectal cancer?, Medicine (United States), 95 (16) (no pagination), 2016	Wrong study design, this is a retrospective observational study

1

1 Appendix L – Research recommendations

- 2 Research recommendations for review question: What is the optimal duration of adjuvant chemotherapy for colorectal cancer?**
- 3**
- 4** No research recommendations were made for this review question.