

Neonatal Parenteral Nutrition

Review questions

NICE guideline tbc

Review questions

July 2019

Early submission

*Developed by National Guideline Alliance
which is part of the Royal College of
Obstetricians and Gynaecologists*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2019. All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

Review questions	5
-------------------------------	----------

Review questions

Number in the scope	Alphanumeric document number when published	Review questions
1.1	A1	What are the predictors for enteral feeding success?
1.2	A2	For those neonates where parenteral nutrition is required, what is the optimal timeframe for doing this?
1.3	D7	What are the most effective relative amounts of nitrogen and non-nitrogen energy (starting and target dose)?
1.4	D8	What are the most effective relative amounts of carbohydrate and lipids (starting and target dose)?
2.1	C	How many kcal/kg/day should be given to preterm and term babies receiving parenteral nutrition?
3.1	D2	What is the optimal target dosage for amino acids in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.2		What is the optimal way (starting dose and approach to increment, if employed) to achieve this target dosage for amino acids?
3.3	D1	What is the optimal target dosage for carbohydrates in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.4		What is the optimal way (starting dose and approach to increment, if employed) to achieve this target dosage for carbohydrates?
3.5	D3	What is the optimal target for lipid dosage in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.6		What is the optimal way (starting dose and approach to increment, if employed) to achieve that?
3.7	D4	What is the clinical effectiveness, efficacy and safety of lipid formulations from different sources (for example, soya, fish oil, or mixed sources)?
3.8	D9	What are the optimal target dosages for calcium and phosphate in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.9	D10	What is the optimal ratio of phosphate to amino acid in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.10	D5	What is the most effective and safe iron supplementation compared to no iron or different dosage/formulations in preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.11	D6	How much (if any) intravenous acetate should be provided to preterm and term babies who are receiving parenteral nutrition and neonatal care?
3.12	E	What is the effectiveness, efficacy and safety of standardised parenteral nutrition bags compared with individualised bags?
4.1	B	What overall osmolality, (concentration of calcium and glucose/dextrose) in parenteral nutrition can determine whether to administer centrally or peripherally?

Number in the scope	Alphanumeric document number when published	Review questions
5.1	F	In babies on parenteral nutrition, what is the optimal frequency of blood sampling and monitoring of glucose, calcium, phosphate, potassium, and serum triglycerides?
6.2	G	What amount of enteral feed (measured in terms of ml/kg/d, or kcal/kg/d, or ratios in relation to parenteral nutrition and enteral feeds) indicates that parenteral nutrition is no longer required?
7.1	H	Are nutrition care/support teams effective in providing parenteral nutrition in preterm and term babies?
8.1	I	What are the most effective methods of information provision about parenteral nutrition, and what information and support about this do parents or carers perceive as useful?
9.1	J	What are the general principles of neonatal parenteral nutrition for preterm and term babies?