

Epilepsies in children, young people and adults: diagnosis and management

[20] Evidence review: Transition from paediatric to adult epilepsy services

NICE guideline <number>

Evidence reviews underpinning recommendations 11.2.1 – 11.2.6 in the NICE guideline

November 2021

Draft for Consultation

*Developed by the National Guideline Centre,
hosted by the Royal College of Physicians*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to Notice of rights.

Content

1	Transition from paediatric to adult epilepsy services.....	5
1.1	Review question.....	5
1.1.1	Introduction.....	5
1.1.2	Summary of the protocol.....	5
1.1.3	Methods and process.....	5
1.1.4	Qualitative evidence.....	6
1.1.5	Summary of studies included in the qualitative evidence.....	6
1.1.6	Summary of the qualitative evidence.....	7
1.1.7	Qualitative evidence summary.....	11
1.1.8	Economic evidence.....	16
1.1.9	The committee’s discussion and interpretation of the evidence.....	17
1.1.10	Other factors the committee took into account.....	20
1.1.11	Recommendations supported by this evidence review.....	20
	References.....	21
	Appendices.....	27
Appendix A	Review protocols.....	27
Appendix B	Literature search strategies.....	34
Appendix C	Qualitative evidence study selection.....	39
Appendix D	Qualitative evidence.....	40
Appendix E	Excluded studies.....	43

1 Transition from paediatric to adult epilepsy services

1.1 Review question

How should the transition from children's and young people's services to adults' epilepsy services be managed?

1.1.1 Introduction

The NICE guideline “Transition from children’s to adults’ services for young people using health or social care services” sets out generic guidance covering the period before, during and after a young person moves from children’s to adults’ services. Young people with epilepsy face additional important challenges that may include uncertainty of prognosis; stigma; risk of injury or death owing, for example, to SUDEP; mental health co-morbidities; side effects of anti-seizure medications and in young women there may be additional risks of medications on the unborn child during pregnancy. Transition planning forms an important part of young people’s epilepsy service design and delivery of care when moving into adult services. This chapter examines the available evidence related to the specific needs of young people with epilepsy in planning and preparing for transition.

1.1.2 Summary of the protocol

For full details see the review protocol in Appendix A.

Table 1: PICO characteristics of review question

Objective	The aim of the review is to identify factors that will help prepare a young person to transition between child and adult epilepsy services. When changing, the young person will be expected to become more involved in decision making on the medical issues affecting them. This review will identify information that is needed to aid the transition.
Population and setting	Inclusion: children and young people with epilepsy, their families and carers 3 groups for access to service strata <ul style="list-style-type: none">• People with drug-resistant epilepsies• People with learning disabilities (access to service-strata)• People with otherwise well-controlled epilepsy
Context	Adolescence is a period of rapid change, both physical and psychosocial, for any young person. Transition can be a difficult time and should be a planned process of addressing the medical and associated comorbid conditions from paediatric to adult care. Significant apprehension may arise as a young person and their family move from a service where they have built therapeutic alliances to an unknown adult service. Both paediatric and adult epilepsy teams need to be motivated to provide a successful and safe transition for these patients.
Review strategy	Synthesis of qualitative research. Results will be presented in a narrative table format. The quality of the evidence will be assessed by a GRADE CerQual approach for each review finding.

1.1.3 Methods and process

This evidence review was developed using the methods and process described in [Developing NICE guidelines: the manual](#). Methods specific to this review question are described in the review protocol in appendix A and the methods document.

Declarations of interest were recorded according to [NICE’s conflicts of interest policy](#).

1 1.1.4 Qualitative evidence

2 1.1.4.1 Included studies

3 We searched for qualitative studies exploring young people’s, parents or carer’s perceptions
4 of their experiences of transitioning from paediatric to adult epilepsy services.

5 Two qualitative studies were included in the review; ^{34,62} one study focussed on people with
6 epilepsy and the other on people with epilepsy and cognitive impairments. The findings were
7 extracted as narratively synthesised information into broad overarching themes, summarised
8 in Table 2 below. Key findings from these studies are summarised in the clinical evidence
9 summary below (Table 3). See also the study selection flow chart in Figure 1, study evidence
10 tables in Appendix D, and excluded studies listed in Appendix E.

11 1.1.4.2 Excluded studies

12 Excluded studies can be found in Table 6 in Appendix E.

13 1.1.5 Summary of studies included in the qualitative evidence

14 **Table 2: Summary of studies included in the evidence review**

Study	Design	Population	Research aim	Comments
Lewis 2013 ³⁴	Semi – structured Interviews or focus groups with two-stage analytical approach: thematic analysis and pattern matching UK n=58 (n=30 young people and n=28 parents of young people who participated)	Young people aged between 13–19 years with epilepsy. Parents of young people with epilepsy who met the inclusion criteria were also invited to participate.	To explore the views of young people with epilepsy (and their parents) about their experience of communication, information and knowledge exchange in two epilepsy services.	
Schultz 2013 ⁶²	Semi-structured Interviews using grounded theory methods, using coding and constant comparison analysis USA n=7	Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	To explicate processes that parents of adolescents with epilepsy and cognitive impairments undergo as they help their adolescent transition from paediatric to adult health care	

15 See Appendix D for full qualitative evidence tables.

1 **1.1.6 Summary of the qualitative evidence**

2 **Table 3: Review findings**

Main findings	Population	Statement of finding
Clarity	Young people aged between 13–19 years with epilepsy and their parents.	Young people perceived that when healthcare professionals used medical jargon, they were keeping information about their epilepsy from them. Most girls (aged 13-<18 years) had yet to receive information, or could not remember receiving information about contraception and pregnancy.
Communication barriers	Young people aged between 13–19 years with epilepsy and their parents.	Young people not knowing what questions to ask healthcare professionals about their epilepsy, what they needed to know about their own epilepsy, and parents being unable to seek information with their child present.
Information	- Young people aged between 13–19 years with epilepsy and their parents. - Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	Continuity of information encompassed the importance of young people remembering facts through repeating information at staged clinical intervals and the negative impact that lack of continuity of information had on the behaviours of parents and lives of young people. Parents expressed difficulty in locating information and understanding the ‘ins and outs’ of it.
Disengaging from healthcare	Young people aged between 13–19 years with epilepsy and their parents.	Disengaging with healthcare professionals and services was linked with young people and parents’ first experience with healthcare professionals in the hand-over clinic.
Ongoing engagement with healthcare professionals	Young people aged between 13–19 years with epilepsy and their parents.	On-going and beneficial engagement was helped by befriending and continuity of care.
Negative stigma	Young people aged between 13–19 years with epilepsy and their parents.	Stigma originating from peers had a significant impact on the behaviour of young women and men, which may reinforce negative feelings about continued engagement with services.
Turmoil	Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	Experienced by all parents. They experienced feelings of fear, rejection or uncertainty.
Parents as advocates	Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	Experienced by all parents. They assumed the role of protector and information gatherer.
Waiting	Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	Discontinuity between agencies resulted in waiting for answers before being able to move forward.
Transition sparked by crisis	Parents or guardians of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments	Transitioning is not planned and is usually a result of a health issue.

3 See Appendix D for full details of qualitative evidence.

1 **1.1.6.1 Narrative summary of review findings**

2 **Review finding 1: Clarity**

3 Young people perceived that when healthcare professionals used medical jargon, they were
4 keeping information about their epilepsy from them, with one stating, "The Doctors were just,
5 "I don't know, I don't know". They just didn't want to say too much sort of thing." When
6 imparting a diagnosis of epilepsy, the issue of lack of clarity tended to cause confusion,
7 stress and anxiety. Another stated they 'didn't know what was going on for ages.' Girls aged
8 16–17 years who had not yet received reproductive health information thought that the ideal
9 age to start discussing gender-specific issues was 14 years of age. The majority of girls
10 (aged >18-19 years) had received some information at various clinic appointments in
11 children's and adult services but could not be specific about what they were told. A number
12 of the girls remembered being told by the nurse not to become pregnant whilst taking their
13 medication due to the risk of congenital malformations. All the girls worried about starving
14 their baby of oxygen during a seizure and about harming their baby should they fall. They did
15 not know if they could give birth 'naturally'.

16 Explanation of quality assessment: the study had no or very minor concerns of
17 methodological limitations in the contributing studies; no or very minor concerns regarding
18 coherence, relevance or adequacy with nothing to lower our confidence. There was a
19 judgement of high confidence in this finding due to no concerns regarding the applicability of
20 this finding to the overarching question.

21 **Review finding 2: Communication barriers**

22 Young people stated not knowing what questions to ask healthcare professionals about their
23 epilepsy, what they needed to know about their own epilepsy, and parents felt unable to seek
24 information with their child present. Some young people were afraid of receiving some
25 epilepsy information, developed an inability to adjust, and were in fear of being alone in case
26 something bad happened. One stated, "I didn't want to speak to anyone about it at all
27 [epilepsy] and I was just really upset all of the time". Parents found it difficult to ask questions
28 and were frequently unable to seek information with their child present. For example, one
29 parent could not ask questions as she was afraid that the information might be frightening to
30 both of them. One common concern was the risk of fatal accidents and sudden death due to
31 seizures. Some parents were too afraid to ask about the risks associated with seizures;
32 however, young people appeared less worried about this in comparison to parents. Shutting
33 down communication with their parents occurred when young people took ownership of their
34 epilepsy in an overly self-protecting way and only talked about epilepsy with their parents
35 and peers when they had to.

36 Explanation of quality assessment: the study had no or very minor concerns of
37 methodological limitations in the contributing studies; no or very minor concerns regarding
38 coherence, relevance or adequacy with nothing to lower our confidence. There was a
39 judgement of high confidence in this finding due to no concerns regarding the applicability of
40 this finding to the overarching question.

41 **Review finding 3: Information**

42 The continuity of information encompassed the importance of young people remembering
43 facts through repeating information at staged clinical intervals and the negative impact that
44 lack of continuity of information had on the behaviours of parents and lives of young people.
45 One stated, "I need something told to me three times before I take it in...perhaps be told
46 something and then read it, go through it, you know". Children and young people wanted
47 verbal information given by healthcare professionals to be backed up with written materials.
48 They were more interested in the format, presentation and relevance of information rather
49 than the type of information, such as information in leaflets and books versus DVDs and web
50 resources. Repeating information at times annoyed some young people, however, the

1 majority benefited from this approach of repetition of information as it strengthened recall and
2 understanding, enabling ability to self-care.

3 Explanation of quality assessment: the study had no or very minor concerns of
4 methodological limitations in the contributing studies; no or very minor concerns regarding
5 coherence, relevance or adequacy with nothing to lower our confidence. There was a
6 judgement of high confidence in this finding due to no concerns regarding the applicability of
7 this finding to the overarching question.

8 **Review finding 4: Disengaging from healthcare**

9 Disengaging with healthcare professionals and services was linked with young people and
10 parents' first experience with healthcare professionals in the hand-over clinic. Once in adult
11 services young people felt forgotten by adult healthcare professionals due to infrequent clinic
12 visits, which were fewer than when following the children's epilepsy care pathway. Parents
13 felt that healthcare professionals in the handover clinic lacked facilitative skills when
14 communicating with their children. The outcome was that some young people disengaged
15 with healthcare professionals and subsequently the adult clinic, as they refused to attend.

16 Explanation of quality assessment: the study had no or very minor concerns of
17 methodological limitations in the contributing studies; no or very minor concerns regarding
18 coherence, relevance or adequacy with nothing to lower our confidence. There was a
19 judgement of high confidence in this finding due to no concerns regarding the applicability of
20 this finding to the overarching question.

21 **Review finding 5: Ongoing engagement with healthcare professionals**

22 On-going and beneficial engagement was helped by befriending and continuity of care.
23 Young people who entered the teenager clinic at an early age (14–15 years) and were
24 befriended/supported by a multi-disciplinary team that included the children's epilepsy nurse
25 appeared more confident and able to communicate. This confidence continued when
26 engaging with adult services, with one stating, "I felt a bit shy at first moving ... because I
27 have been with that doctor for years and I was used to him, moving on was a bit scary".
28 Young people who saw the same healthcare professional, who was responsive to their
29 needs, at frequently staged intervals, developed a positive rapport. Even when older, many
30 young people said being able to bring parents into the clinic room helped. They felt that if
31 they forgot or did not understand what had been said, then at least they could turn to their
32 parent for the information. Including parents during consultations seemed to improve family
33 functioning, and young people could see that this alleviated their parents' stress and worry.

34 Explanation of quality assessment: the study had no or very minor concerns of
35 methodological limitations in the contributing studies; no or very minor concerns regarding
36 coherence, relevance or adequacy with nothing to lower our confidence. There was a
37 judgement of high confidence in this finding due to no concerns regarding the applicability of
38 this finding to the overarching question.

39 **Review finding 6: Negative stigma**

40 The stigma of having epilepsy originating from peers had a significant impact on the
41 behaviour of girls and boys, which may reinforce negative feelings about continued
42 engagement with services. An inability to cope with stigma, especially for girls, originated
43 from being stigmatised by peers at school and work. One participant mentioned, "Someone
44 came up to me the other day and they said, 'do you remember that time when you had a fit',
45 and I said 'no' 'well we call it a funky chicken' and I said 'why?' (she mimicked flapping her
46 arms to demonstrate what the other person did)...I didn't like going to school if I had one so I
47 would stay at home for about a week afterwards...Because I was scared what people would

1 say". Young people were more inclined to stay engaged with epilepsy services if they could
2 experience some tangible benefits from continued attendance.

3 Explanation of quality assessment: the study had no or very minor concerns of
4 methodological limitations in the contributing studies; no or very minor concerns regarding
5 coherence, relevance or adequacy with nothing to lower our confidence. There was a
6 judgement of high confidence in this finding due to no concerns regarding the applicability of
7 this finding to the overarching question.

8 1.1.6.2 Narrative summary of review findings from those with learning disabilities

9 Review finding 1: Turmoil

10 Turmoil was experienced by all parents. They experienced feelings of fear, rejection or
11 uncertainty. One participant stated that "changes are scary. And especially when it has to do
12 with health - living or dying. To me, it's very frightening." Parents also felt uncertainty
13 regarding where they could receive healthcare or the source of insurance coverage. Another
14 participant said, "there are a lot of unknowns... there's too many of them – just to leave them
15 hanging".

16 Explanation of quality assessment: the study had moderate concerns of methodological
17 limitations in the contributing studies; no or very minor concerns regarding coherence,
18 relevance or adequacy with nothing to lower our confidence. There was a judgement of
19 moderate confidence in this finding due to concerns regarding the ethical considerations and
20 the relationship between the researcher and participants not being adequately considered.

21 Review finding 2: Parents as advocates

22 This was experienced by all parents. They assumed the role of protector and information
23 gatherer, with one participant mentioning, "you have to be very diligent in the way that you
24 cover yourself in terms of if you really don't want that service". Parents took charge of
25 locating and sustaining essential health care services. As an information gatherer they
26 needed to identify resources whilst understanding the differences in eligibility requirements
27 and delivery of services for various agencies.

28 Explanation of quality assessment: the study had moderate concerns of methodological
29 limitations in the contributing studies; no or very minor concerns regarding coherence,
30 relevance or adequacy with nothing to lower our confidence. There was a judgement of
31 moderate confidence in this finding due to concerns regarding the ethical considerations and
32 the relationship between the researcher and participants' not being adequately considered.

33 Review finding 3: Information

34
35 Parents expressed difficulty in locating information and understanding the 'ins and outs' of it.
36 One parent mentioned the difficulty in locating information and understanding it. They stated
37 it is a "...huge web of trying to figure out how it all works", and parents found support groups
38 the most helpful source. Along with support groups 'just networking with parents' was also
39 claimed to be a valuable source.

40 Explanation of quality assessment: the study had moderate concerns of methodological
41 limitations in the contributing studies; no or very minor concerns regarding coherence,
42 relevance or adequacy with nothing to lower our confidence. There was a judgement of
43 moderate confidence in this finding due to concerns regarding the ethical considerations and
44 the relationship between the researcher and participants not being adequately considered.

1 **Review finding 4: Waiting**

2 Because of participants experiencing discontinuity between agencies, they were confined to
3 a state of ‘captive waiting’ for answers before being able to move forward. This captivity
4 generated feelings of uncertainty and a fear of loss of services. Waiting on lists to acquire
5 new services was not new to many of the families. Furthermore, parents often did not have
6 the choice of coverage; they were at the mercy of the ‘system’ to determine when and where
7 they were going to receive services.

8 Explanation of quality assessment: the study had moderate concerns of methodological
9 limitations in the contributing studies; no or very minor concerns regarding coherence,
10 relevance or adequacy with nothing to lower our confidence. There was a judgement of
11 moderate confidence in this finding due to concerns regarding the ethical considerations and
12 the relationship between the researcher and participants’ not being adequately considered.

13 **Review finding 5: Transition sparked by crisis**

14 Participants explained how transition is not planned and is usually a result of a health issue.
15 It was mentioned how there is no preparation, and they are just transferred to the adult
16 system. One participant stated, “...nothing was planned or nothing. It was like this is an
17 emergency, so now we’ve got to fix it”.

18 Explanation of quality assessment: the study had moderate concerns of methodological
19 limitations in the contributing studies; no or very minor concerns regarding coherence,
20 relevance or adequacy with nothing to lower our confidence. There was a judgement of
21 moderate confidence in this finding due to concerns regarding the ethical considerations and
22 the relationship between the researcher and participants’ not being adequately considered.

23 **1.1.7 Qualitative evidence summary**

24 Table 4: Summary of evidence

Study design and sample size		Finding	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Clarity					
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	Young people perceived that when healthcare professionals used medical jargon, they were keeping information about their epilepsy from them.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

1

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Communication barriers					
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	Young people not knowing what questions to ask healthcare professionals about their epilepsy, what they needed to know about their own epilepsy, and parents being unable to seek information with their child present.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

2

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Information					
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	Continuity of information encompassed the importance of young people remembering facts through repeating information at staged clinical intervals and the negative impact that lack of continuity of information had on the behaviours of parents and lives of young people.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

3

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Disengaging from healthcare					
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	Disengaging with healthcare professionals and services was linked with young people and parents' first experience with healthcare professionals in the hand-over clinic.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

1

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Ongoing engagement with healthcare professionals					
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	On-going and beneficial engagement was helped by befriending and continuity of care.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

2

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Negative stigma					

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
1 study Lewis 2013 ³⁴	A combination of interviews and focus groups	Stigma originating from peers had a significant impact on the behaviour of girls and boys, which may reinforce negative feelings about continued engagement with services.	Limitations	No or very minor limitations	HIGH
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	No concerns	

1
2

Those with learning disabilities

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Turmoil					
1 study Schultz 2013 ⁶²	Individual interviews	Experienced by all parents. They experienced feelings of fear, rejection or uncertainty.	Limitations	Moderate limitations ¹	LOW
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	Moderate concerns ²	

¹ downgraded for lack of information on; participant/researcher relationship and ethical considerations
² downgraded for small sample size, n=7

3
4
5

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Parents as advocates					
1 study Schultz 2013 ⁶²	Individual interviews	Experienced by all parents. They assumed the role of protector and information gatherer.	Limitations	Moderate limitations ¹	LOW
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	Moderate concerns ²	

¹ downgraded for lack of information on; participant/researcher relationship and ethical considerations

² downgraded for small sample size, n=7

1
2
3

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Information					
1 study Schultz 2013 ⁶²	A combination of interviews and focus groups	Parents expressed difficulty in locating information and understanding the 'ins and outs' of it.	Limitations	Moderate limitations ¹	LOW
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	Moderate concerns ²	

¹ downgraded for lack of information on; participant/researcher relationship and ethical considerations

² downgraded for small sample size, n=7

4
5
6
7

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Waiting					
1 study Schultz 2013 ⁶²	Individual interviews	Discontinuity between agencies resulted in waiting for answers before being able to move forward.	Limitations	Moderate limitations ¹	LOW
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	Moderate concerns ²	

¹ downgraded for lack of information on; participant/researcher relationship and ethical considerations

² downgraded for small sample size, n=7

1
2
3
4

Study design and sample size		Findings	Quality assessment		
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence
Transition sparked by crisis					
1 study Schultz 2013 ⁶²	Individual interviews	Transitioning is not planned and is usually a result of health issue or change in insurance.	Limitations	Moderate limitations ¹	LOW
			Coherence	No or very minor concerns	
			Relevance	No or very minor concerns	
			Adequacy	Moderate concerns ²	

¹ downgraded for lack of information on; participant/researcher relationship and ethical considerations

² downgraded for small sample size, n=7

5
6
7

8 1.1.8 Economic evidence

9 The committee agreed that health economic studies would not be relevant to this review
10 question, and so were not sought.

1 **1.1.9 The committee's discussion and interpretation of the evidence**

2 **1.1.9.1 The quality of the evidence**

3 High-quality evidence was found for the themes of clarity, communication barriers,
4 disengaging from healthcare, ongoing engagement with healthcare professionals and
5 negative stigma.

6 Low-quality evidence was found for the themes of information giving, turmoil, parents as
7 advocates, waiting, and transition sparked by a crisis.

8 The committee noted one small study was from the US and that configuration of services
9 would be quite different to UK practice. However, the committee agreed the description of the
10 concerns reported by parents of young people with learning disabilities were common and
11 still applicable.

12 **1.1.9.2 Findings identified in the evidence synthesis**

13 Clarity, communication, and lack of information were identified as important aspects of the
14 evidence that concerned information giving during transition from children to adult epilepsy
15 services. Young people reported a lack of clarity caused by the use of medical jargon, and
16 this led to a perception that health professions were withholding information. Young women
17 expressed concern that they were not provided information on contraception or pregnancy
18 and the associated risks of epilepsy in pregnancy. There is a risk that unclear information
19 provided to young women might result in their abruptly stopping medication without
20 understanding the consequences if they were to become pregnant. The importance of
21 providing clear and balanced information to young people about their medication and any
22 associated risks was acknowledged by the committee.

23 Barriers to communication expressed by young people included not knowing what to ask or
24 being afraid to ask about risks associated with epilepsy such as SUDEP. Young people
25 considered information about their plan for transition could be facilitated by health
26 professionals repeating information at staged intervals to ensure their management plan and
27 the support that is to be provided is understood and agreed. Parents of young people with
28 epilepsy reported difficulty discussing epilepsy in front of their children. The committee
29 acknowledged that both parents and young people could feel inhibited in discussing sensitive
30 issues in front of each other and agreed health professionals should provide the opportunity
31 for joint and individual meetings with parents and the young person to facilitate open
32 discussion.

33 The committee noted the evidence highlighted the importance of providing information
34 without jargon and in an accessible format which should be repeated at different time points
35 over the transition period. The evidence reported children and young people valued having
36 verbal information given by healthcare professionals backed up with written materials. The
37 committee decided to make a recommendation that information should be provided on the
38 individual's management plan for transition in a format that meets the young person's needs
39 using language they understand. The committee also recommended that this may require
40 regularly revisiting the individualised management plan over a period of time to ensure the
41 young person fully understands their care plan and the support to be provided.

42 The themes of engagement and disengagement with healthcare professionals and services
43 were also identified from the evidence. The committee acknowledged that transition could be
44 a distressing time for both children and adults. The evidence found that young people who
45 engaged with a multidisciplinary team (MDT) felt more confident and able to communicate
46 their views. Health care professionals befriending the young person also facilitated
47 engagement with services and continuity of care. The evidence also described how young
48 people disengaged with services and healthcare professionals due to negative experiences
49 when attending 'hand-over' appointments and at subsequent clinic visits. Less frequent or

1 delayed clinic appointments in adult services than had occurred in children's services
2 resulted in young people feeling forgotten about or were interpreted as being unsupportive.
3 The committee agreed that encouraging discussion of the expectations of service provision
4 between the young person and their new adult epilepsy team can be helpful to establish the
5 person's needs and what the service is able to provide. Making sure young people are aware
6 of the importance of attending appointments was discussed, and the committee agreed adult
7 services should not automatically discharge young people who miss appointments during the
8 transition, although they were aware this does happen occasionally.

9 The committee discussed the value of having a management plan developed by the epilepsy
10 multidisciplinary teams who can then facilitate the transition rather than transition occurring in
11 crisis (for example, by a sixteen-year-old attending the accident and emergency department
12 and meeting the adult care team for the first time in this context). It was considered essential
13 that planning should occur with both paediatric services and the adult MDT working together.
14 Other external agencies may also need to be included as part of the MDT such as social
15 care services and school nurses. Descriptions of barriers to a smooth transition were noted
16 by the committee, who acknowledged the importance of clear communication channels and
17 sharing of information between multi-professional epilepsy teams, young people and their
18 parents or carers. The committee also noted that a young person's transition is bespoke to
19 that person and should be recognised as such by the paediatric and adult MDT. The
20 committee discussed how the severity of a young person's epilepsy may change over time,
21 and planning for transition should include a review of the diagnosis and medical
22 management of the person with epilepsy. The committee discussed that there could be a
23 lack of certainty around the diagnosis of epilepsy, for example, that dissociative seizures may
24 look similar to epileptic seizures indicating the necessity of the clinician to review the original
25 diagnosis. The committee agreed this opportunity for review should not be missed and
26 recommended this should involve both paediatric and adult MDT.

27 The evidence showed stigma of having epilepsy originating from peers had a significant
28 impact on the behaviour of young people, which may reinforce negative feelings about
29 continued engagement with services. Stigma can have a significant impact on young
30 people's everyday activities and school attainment. Participation in social activities can also
31 be adversely curtailed because of seizure activity or the young person avoiding social events
32 due to embarrassment or negative self-image. The committee acknowledged the importance
33 of encouraging young people to discuss their experiences and providing support during
34 consultations on transition.

35 The themes arising from the evidence on young people with learning disabilities were turmoil,
36 parents as advocates, information, waiting, and transition as a result of a crisis. Parents of
37 young people with learning disabilities expressed feelings of fear, rejection or uncertainty
38 around their child's transition from paediatric to adult services. Parents assumed the role of
39 protector and information gatherer and described difficulty in finding information and
40 understanding the implications of the changes in service provision for their child. The
41 committee noted that parents and carers should be empowered to have the confidence to
42 speak up and advocate for the young person, and health professionals need to be mindful of
43 how intimidating the system can be. Parents observed discontinuity between agencies
44 involved with transition of care, who often acted as a barrier to successful transfer due to lack
45 of coordination between teams, leading to delays in moving plans forward. It was noted that
46 transitioning was often not planned and would occur as a consequence of a crisis. The
47 committee acknowledged the significant anxiety that some parents or carers can experience
48 as the young person prepares to transfer to the adult service, and how important it is to make
49 time to include parents and carers in the planning stage and explain the process fully.

50 The committee considered the themes described by parents and carers of young people with
51 learning disabilities and observed that these often overlapped with the themes described in
52 the other study of young people who didn't have a learning disability. The committee agreed
53 this demonstrated the generalisability of the findings across the studies and were applicable

1 to all young people, their parents and carers during transition, although difficulties can be
2 particularly marked in young people with complex needs.

3 The committee agreed that people with learning disabilities may have very complex needs,
4 and transition to adult services could require more planning, need longer appointments and
5 involve many other specialities, including professionals from a learning disabilities MDT,
6 CAMHS and special needs schools. The committee noted that transition of young people
7 with learning disabilities could occur considerably later than other young people with
8 epilepsy, even up to the age of 24 years, and this was often because of the multiple agencies
9 required to organise an optimal package of care and the different age cut-offs used in
10 services for young people with learning disabilities. The committee acknowledged some
11 young people with complex health and social care needs, such as treatment-resistant
12 epilepsies may require additional management support or input from multiple agencies to
13 provide appropriate packages of care. The committee decided to make a consensus
14 recommendation based on their experience that planning transition for these young people
15 should start earlier, although agreed it would be inappropriate to specify a timeframe as this
16 would be dependent on individual circumstances and needs.

17 The committee made recommendations based on the themes found in the evidence review,
18 but they also identified issues from their experience that they considered important to include
19 in discussions as part of transition. These were topics often raised by the young person
20 during preparations for transfer. The committee decided to make a consensus
21 recommendation that highlighted these key concerns, or questions, that are commonly
22 raised. These include adherence to medication, the effects epilepsy and medication may
23 have on cognition and behaviour, and other work and education-related issues such as
24 advice on how to access benefits or special arrangements that may be required for taking
25 exams. Young people often described disturbance of sleep patterns, and the committee
26 noted the importance of discussing attending social activities safely, such as parties or
27 staying overnight at other people's houses, and ensuring the young person is aware of the
28 need to adhere to a healthy sleep regime. Other common concerns raised include being able
29 to participate in activities such as sports or driving and being aware of triggers such as
30 flashing lights that can induce a seizure. Many young people want to discuss preparing for
31 independent living with someone outside of their family and turn to the MDT for advice and
32 guidance. The committee agreed it was important to discuss potential co-morbidities of
33 epilepsy, especially impaired memory and low mood. The committee also acknowledged
34 social aspects of epilepsy such as decisions on when or if to disclose epilepsy status and the
35 emotional health and psychological wellbeing of the young person as issues of stigma and
36 cognitive consequences of the person's epilepsy can have a negative impact on mental
37 health.

38 1.1.9.3 Cost-effectiveness and resource use

39 Cost-effectiveness evidence was not sought as this was a qualitative review. The
40 recommendations generally provide guidance regarding the content of information and
41 support specific to people transitioning from child to adult epilepsy services in line with the
42 general principles of provision of information already established in the existing NICE
43 Transition from Children's to Adults' Services Guideline and so the recommendations
44 concerning the information and support provided to young people with epilepsy transitioning
45 were not considered to have a substantial resource impact over and above this.

46 The committee noted that transition for children to adult epilepsy services should be
47 individually tailored to the young person and acknowledged it is current best practice to plan
48 earlier transition for people who have complex or additional social care needs but for a small
49 proportion of people, this does not happen. It was discussed that although planning transition
50 earlier will result in additional costs for the NHS at the pre-transition phase, these will likely
51 be offset by cost savings observed in the form of better health care received in adult
52 services.

1 Reviewing the diagnosis and management plan of a young person with epilepsy during
2 transition to adult services is generally current practice at epilepsy centres, but the level of
3 review and involvement of both paediatric and adult multidisciplinary teams can vary from
4 centre to centre, especially for people with more complex epilepsy. The committee stressed
5 the importance of an 'in-depth' review of a person's epilepsy as this may be an ideal time and
6 opportunity for a full holistic review to be conducted. The committee noted that taking the
7 time to review the young person's epilepsy has the potential to lead to substantial health
8 gains if it means people subsequently receive better-tailored health care plans. A review of a
9 young person's epilepsy will generally consist of; communication between both the children's
10 and adult epilepsy teams, an initial meeting with the adult epilepsy services team, and any
11 tests a person may receive as a result of these. The need for tests will be individualised to a
12 person's epilepsy and will only be conducted if the clinician believes young people may
13 benefit from long-term health gains. The committee noted that around 20%-30% of young
14 people might undergo additional testing when transitioning from children to adults' services,
15 with most of these tests being outpatient appointments. It is important to have early
16 discussions about the potential need for inpatient tests at the time of transition or in the future
17 to reduce anxiety and allow preparatory hospital/ward visits. As reviewing a person's
18 epilepsy is generally current practice at epilepsy centres, this is unlikely to have a substantial
19 resource impact. At epilepsy centres where the level of reviewing is less than stated in the
20 recommendation, there may be a small resource impact, but these additional costs have the
21 potential to be offset by health gains achieved by optimal diagnosis and individualised
22 management.

23 **1.1.10 Other factors the committee took into account**

24 The committee acknowledged the Ready Steady Go programme for transition to adult
25 services developed by Southampton Children's Hospital, now widely adopted within epilepsy
26 healthcare services. The committee considered the recommendations in the NICE guideline
27 Transition from children to adult services to be applicable and provided guidance on the
28 general principles for planning and organising the transfer of young people to an adult
29 epilepsy service. A cross-reference to the guideline was made.

30 **1.1.11 Recommendations supported by this evidence review**

31 This evidence review supports recommendations 11.2.1 – 11.2.6 in the NICE guideline.
32

References

1. Abubakar A, Kariuki SM, Tumaini JD, Gona J, Katana K, Owen JA et al. Community perceptions of developmental and behavioral problems experienced by children living with epilepsy on the Kenyan coast: A qualitative study. *Epilepsy & Behavior*. 2015; 45:74-78
2. Alkhamra H, Tannous A, Hadidi M, Alkhateeb J. Knowledge and attitudes toward epilepsy among school teachers and counselors in Jordan. *Epilepsy & Behavior*. 2012; 24(4):430-434
3. Amjad RN, Navab E, Nasrabadi AN. Parents of children with epilepsy captured by epilepsy: A qualitative study. *Acta Medica Mediterranea*. 2016; 32(Special Issue 4):1303-1309
4. Andrade DM, Bassett AS, Bercovici E, Borlot F, Bui E, Camfield P et al. Epilepsy: Transition from pediatric to adult care. Recommendations of the Ontario epilepsy implementation task force. *Epilepsia*. 2017; 58(9):1502-1517
5. Aytch LS, Hammond R, White C. Seizures in infants and young children: an exploratory study of family experiences and needs for information and support. *Journal of Neuroscience Nursing*. 2001; 33(5):278-285
6. Baca CM, Barry F, Berg AT. The epilepsy transition care gap in young adults with childhood-onset epilepsy. *Epilepsy & Behavior*. 2018; 87:146-151
7. Baird AD, Wilson SJ, Bladin PF, Saling MM, Reutens DC. Sexual outcome after epilepsy surgery. *Epilepsy & Behavior*. 2003; 4(3):268-278
8. Benson A, Lambert V, Gallagher P, Shahwan A, Austin JK. "I don't want them to look at me and think of my illness, I just want them to look at me and see me": Child perspectives on the challenges associated with disclosing an epilepsy diagnosis to others. *Epilepsy & Behavior*. 2015; 53:83-91
9. Benson A, Lambert V, Gallagher P, Shahwan A, Austin JK. Parent perspectives of the challenging aspects of disclosing a child's epilepsy diagnosis to others: Why don't they tell? *Chronic Illness*. 2017; 13(1):28-48
10. Benson A, O'Toole S, Lambert V, Gallagher P, Shahwan A, Austin JK. The stigma experiences and perceptions of families living with epilepsy: Implications for epilepsy-related communication within and external to the family unit. *Patient Education and Counseling*. 2016; 99(9):1473-1481
11. Blank L, Baird W, Reuber M. Patient perceptions of the referral of older adults to an epilepsy clinic: do patients and professionals agree who should be referred to a specialist? *Epilepsy & Behavior*. 2014; 34:120-123
12. Burke L, Kirkham J, Arnott J, Gray V, Peak M, Beresford MW. The transition of adolescents with juvenile idiopathic arthritis or epilepsy from paediatric health-care services to adult health-care services: A scoping review of the literature and a synthesis of the evidence. *Journal of Child Health Care*. 2018; 22(3):332-358
13. Butau T, Piachaud J. Knowledge and beliefs about epilepsy in mothers of children with epilepsy: a view from a developing country. *Central African Journal of Medicine*. 1993; 39(9):183-188
14. Camfield P, Camfield C, Busiah K, Cohen D, Pack A, Nabbout R. The transition from pediatric to adult care for youth with epilepsy: Basic biological, sociological, and psychological issues. *Epilepsy & Behavior*. 2017; 69:170-176

- 1 15. Canvin K, Jacoby A. Duty, desire or indifference? A qualitative study of patient
2 decisions about recruitment to an epilepsy treatment trial. *Trials*. 2006; 7:32
- 3 16. Chen HJ, Chen YC, Yang HC, Chi CS. Lived experience of epilepsy from the
4 perspective of children in Taiwan. *Journal of Clinical Nursing*. 2010; 19(9-10):1415-
5 1423
- 6 17. Chew J, Carpenter J, Haase AM. Young people's experiences of living with epilepsy:
7 The significance of family resilience. *Social Work in Health Care*. 2018; 57(5):332-354
- 8 18. Chiu YP, Lee TY, Lin KL, Laadt VL. Adjusting to a seizure-free "new normal" life
9 following discontinuation of antiepileptic drugs during adolescence. *Epilepsy &
10 Behavior*. 2014; 33:54-58
- 11 19. Chong L, Jamieson NJ, Gill D, Singh-Grewal D, Craig JC, Ju A et al. Children's
12 experiences of epilepsy: A systematic review of qualitative studies. *Pediatrics*. 2016;
13 138(3):e20160658
- 14 20. Collard SS, Regmi P. Qualitative insights into the feelings, knowledge, and impact of
15 SUDEP: A narrative synthesis. *Epilepsy & Behavior*. 2019; 94:20-28
- 16 21. Eklund PG, Sivberg B. Adolescents' lived experience of epilepsy. *Journal of
17 Neuroscience Nursing*. 2003; 35(1):40-49
- 18 22. Elliott IM, Lach L, Smith ML. I just want to be normal: a qualitative study exploring
19 how children and adolescents view the impact of intractable epilepsy on their quality
20 of life. *Epilepsy & Behavior*. 2005; 7(4):664-678
- 21 23. Gauffin H, Flensner G, Landtblom AM. Being parents with epilepsy: thoughts on its
22 consequences and difficulties affecting their children. *Neuropsychiatric Disease and
23 Treatment*. 2015; 11:1291-1298
- 24 24. Geerlings RP, Aldenkamp AP, de With PH, Zinger S, Gottmer-Welschen LM, de
25 Louw AJ. Transition to adult medical care for adolescents with epilepsy. *Epilepsy &
26 Behavior*. 2015; 44:127-135
- 27 25. Hames A, Appleton R. Living with a brother or sister with epilepsy: siblings'
28 experiences. *Seizure*. 2009; 18(10):699-701
- 29 26. Harden J, Black R, Chin RFM. Families' experiences of living with pediatric epilepsy:
30 A qualitative systematic review. *Epilepsy & Behavior*. 2016; 60:225-237
- 31 27. Harden J, Tonberg A, Chin RF, McLellan A, Duncan S. 'If you're gonna die, you're
32 gonna die': Young adults' perceptions of sudden unexpected death in epilepsy.
33 *Chronic Illness*. 2015; 11(3):230-241
- 34 28. Heath G, Abdin S, Begum R, Kearney S. Putting children forward for epilepsy
35 surgery: A qualitative study of UK parents' and health professionals' decision-making
36 experiences. *Epilepsy & Behavior*. 2016; 61:185-191
- 37 29. Hightower S, Carmon M, Minick P. A qualitative descriptive study of the lived
38 experiences of school-aged children with epilepsy. *Journal of Pediatric Health Care*.
39 2002; 16(3):131-137
- 40 30. Joung WJ. Pregnancy and childbirth experiences of women with epilepsy: A
41 phenomenological approach. *Asian Nursing Research*. 2019; 13(2):122-129
- 42 31. Kaddumukasa M, Nalubwama H, Kaddumukasa MN, Lhatoo S, Sewankambo N,
43 Katabira E et al. Barriers to epilepsy care in Central Uganda, a qualitative interview
44 and focus group study involving PLWE and their caregivers. *BMC Neurology*. 2019;
45 19(1):161

- 1 32. Kampra M, Tzerakis N, Lund Holm Thomsen L, Katsarou E, Voudris K, S DM et al.
2 The challenges that parents of children with epilepsy face: A qualitative study.
3 *Epilepsy & Behavior*. 2017; 71(Pt A):94-103
- 4 33. Kuchenbuch M, Chemaly N, Chiron C, Dulac O, Nabbout R. Transition and transfer
5 from pediatric to adult health care in epilepsy: a families' survey on Dravet syndrome.
6 *Epilepsy & Behavior*. 2013; 29(1):161-165
- 7 34. Lewis SA, Noyes J. Effective process or dangerous precipice: qualitative comparative
8 embedded case study with young people with epilepsy and their parents during
9 transition from children's to adult services. *BMC Pediatrics*. 2013; 13:169
- 10 35. Louik J, Doumlele K, Hussain F, Crandall L, Buchhalter J, Hesdorffer D et al.
11 Experiences with premorbid SUDEP discussion among participants in the North
12 American SUDEP Registry (NASR). *Epilepsy & Behavior*. 2017; 70(Pt A):131-134
- 13 36. MacLeod JS. The everyday lives of adolescent girls with epilepsy: A qualitative
14 description. Indianapolis, IN. Indiana University-Purdue University Indianapolis. 2009
- 15 37. Mc Govern EM, Maillart E, Bourgninaud M, Manzato E, Guillonnet C, Mochel F et al.
16 Making a 'JUMP' from paediatric to adult healthcare: A transitional program for young
17 adults with chronic neurological disease. *Journal of the Neurological Sciences*. 2018;
18 395:77-83
- 19 38. McAuley JW, Patankar C, Lang C, Prasad M. Evaluating the concerns of pregnant
20 women with epilepsy: a focus group approach. *Epilepsy & Behavior*. 2012; 24(2):246-
21 248
- 22 39. McEwan MJ, Espie CA, Metcalfe J, Brodie MJ, Wilson MT. Quality of life and
23 psychosocial development in adolescents with epilepsy: a qualitative investigation
24 using focus group methods. *Seizure*. 2004; 13(1):15-31
- 25 40. McNelis AM, Buelow J, Myers J, Johnson EA. Concerns and needs of children with
26 epilepsy and their parents. *Clinical Nurse Specialist*. 2007; 21(4):195-202
- 27 41. Mengoni SE, Gates B, Parkes G, Wellsted D, Barton G, Ring H et al. "Sometimes, it
28 just stops me from doing anything": A qualitative exploration of epilepsy management
29 in people with intellectual disabilities and their carers. *Epilepsy & Behavior*. 2016;
30 64(Pt A):133-139
- 31 42. Miller WR, Young N, Friedman D, Buelow JM, Devinsky O. Discussing sudden
32 unexpected death in epilepsy (SUDEP) with patients: practices of health-care
33 providers. *Epilepsy & Behavior*. 2014; 32:38-41
- 34 43. Moffat C, Dorris L, Connor L, Espie CA. The impact of childhood epilepsy on quality
35 of life: A qualitative investigation using focus group methods to obtain children's
36 perspectives on living with epilepsy. *Epilepsy & Behavior*. 2009; 14(1):179-189
- 37 44. Mu PF. Transition experience of parents caring of children with epilepsy: a
38 phenomenological study. *International Journal of Nursing Studies*. 2008; 45(4):543-
39 551
- 40 45. Nashef L, Garner S, Sander JW, Fish DR, Shorvon SD. Circumstances of death in
41 sudden death in epilepsy: interviews of bereaved relatives. *Journal of Neurology,*
42 *Neurosurgery and Psychiatry*. 1998; 64(3):349-352
- 43 46. National Institute for Health and Care Excellence. Developing NICE guidelines: the
44 manual [updated October 2020]. London. National Institute for Health and Care
45 Excellence, 2014. Available from:
46 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20overview>

- 1 47. Ninnoni JPK. A qualitative study of the communication and information needs of
2 people with learning disabilities and epilepsy with physicians, nurses and carers.
3 BMC Neurology. 2019; 19(1):12
- 4 48. Nisbet T, Turbull S, Mulhern S, Razvi S. 'Breaking Good News': Neurologists'
5 experiences of discussing SUDEP with patients in Scotland. *Epilepsy & Behavior*.
6 2017; 70(Pt A):72-79
- 7 49. O'Toole S, Lambert V, Gallagher P, Shahwan A, Austin JK. Talking about epilepsy:
8 Challenges parents face when communicating with their child about epilepsy and
9 epilepsy-related issues. *Epilepsy & Behavior*. 2016; 57(Pt A):9-15
- 10 50. Pashley S, O'Donoghue MF. The safety of anti-epileptic drug regimens: a qualitative
11 study of factors determining the success of counselling women before conception.
12 *Journal of Family Planning and Reproductive Health Care*. 2009; 35(3):153-156
- 13 51. Prinjha S, Chapple A, Herxheimer A, McPherson A. Many people with epilepsy want
14 to know more: a qualitative study. *Family Practice*. 2005; 22(4):435-441
- 15 52. RamachandranNair R, Jack SM. SUDEP: What do adult patients want to know?
16 *Epilepsy & Behavior*. 2016; 64(Pt A):195-199
- 17 53. RamachandranNair R, Jack SM, Meaney BF, Ronen GM. SUDEP: what do parents
18 want to know? *Epilepsy & Behavior*. 2013; 29(3):560-564
- 19 54. RamachandranNair R, Jack SM, Strohm S. SUDEP: To discuss or not?
20 Recommendations from bereaved relatives. *Epilepsy & Behavior*. 2016; 56:20-25
- 21 55. Ridsdale L, Viridi C, Noble A, Morgan M. Explanations given by people with epilepsy
22 for using emergency medical services: A qualitative study. *Epilepsy & Behavior*.
23 2012; 25(4):529-533
- 24 56. Roberts J, Whiting C. Caregivers of school children with epilepsy: findings of a
25 phenomenological study. *British Journal of Special Education*. 2011; 38(4):169-177
- 26 57. Ronen GM, Rosenbaum P, Law M, Streiner DL. Health-related quality of life in
27 childhood epilepsy: the results of children's participation in identifying the
28 components. *Developmental Medicine and Child Neurology*. 1999; 41(8):554-559
- 29 58. Rushe H. Living with a sibling with refractory epilepsy: Children's experience of
30 having a sibling with epilepsy. University of Essex. 2011
- 31 59. Saburi GL, Mapanga KG, Mapanga MB. Perceived family reactions and quality of life
32 of adults with epilepsy. *Journal of Neuroscience Nursing*. 2006; 38(3):156-165
- 33 60. Sample PL, Ferguson PL, Wagner JL, Elisabeth Pickelsimer E, Selassie AW.
34 Experiences of persons with epilepsy and their families as they look for medical and
35 community care: A focus group study from South Carolina. *Epilepsy & Behavior*.
36 2006; 9(4):649-662
- 37 61. Scheffer IE, Dravet C. Transition to adult life in the monogenic epilepsies. *Epilepsia*.
38 2014; 55(Suppl 3):12-15
- 39 62. Schultz RJ. Parental experiences transitioning their adolescent with epilepsy and
40 cognitive impairments to adult health care. *Journal of Pediatric Health Care*. 2013;
41 27(5):359-366
- 42 63. Shih P, Nikpour A, Bleasel A, Herkes G, Mitchell R, Seah R et al. Leading up to
43 saying "yes": A qualitative study on the experience of patients with refractory epilepsy
44 regarding presurgical investigation for resective surgery. *Epilepsy & Behavior*. 2018;
45 83:36-43

- 1 64. Smith G, Wagner J, Andrews J, Austin J, Mueller M, Carter E et al. Caregiving in
2 pediatric epilepsy: results of focus groups and implications for research and practice.
3 *Epilepsy & Behavior*. 2014; 34:34-41
- 4 65. Snape D, Wang W, Wu J, Jacoby A, Baker GA. Knowledge gaps and uncertainties
5 about epilepsy: Findings from an ethnographic study in China. *Epilepsy & Behavior*.
6 2009; 14(1):172-178
- 7 66. Sonecha S, Noble AJ, Morgan M, Ridsdale L. Perceptions and experiences of
8 epilepsy among patients from black ethnic groups in South London. *Primary Health
9 Care Research & Development (Cambridge University Press/UK)*. 2015; 16(5):450-
10 460
- 11 67. Swarztrauber K, Dewar S, Engel Jr J. Patient attitudes about treatments for
12 intractable epilepsy. *Epilepsy & Behavior*. 2003; 4(1):19-25
- 13 68. Tanaka C, Tuliao MTR, Tanaka E, Yamashita T, Matsuo H. A qualitative study on the
14 stigma experienced by people with mental health problems and epilepsy in the
15 Philippines. *BMC Psychiatry*. 2018; 18(1):325
- 16 69. Thompson D, Thomas H, Solomon J, Nashef L, Kendall S. Chronic illness,
17 reproductive health and moral work: women's experiences of epilepsy. *Chronic
18 Illness*. 2008; 4(1):54-64
- 19 70. Thompson R, Linehan C, Glynn M, Kerr MP. A qualitative study of carers' and
20 professionals' views on the management of people with intellectual disability and
21 epilepsy: A neglected population. *Epilepsy & Behavior*. 2013; 28(3):379-385
- 22 71. Tonberg A, Harden J, McLellan A, Chin RF, Duncan S. A qualitative study of the
23 reactions of young adults with epilepsy to SUDEP disclosure, perceptions of risks,
24 views on the timing of disclosure, and behavioural change. *Epilepsy & Behavior*.
25 2015; 42:98-106
- 26 72. Van Naarden Braun K, Yeargin-Allsopp M, Lollar D. A multi-dimensional approach to
27 the transition of children with developmental disabilities into young adulthood: The
28 acquisition of adult social roles. *Disability and Rehabilitation*. 2006; 28(15):915-928
- 29 73. Varley J, Delanty N, Normand C, Fitzsimons M. The health care journeys
30 experienced by people with epilepsy in Ireland: what are the implications for future
31 service reform and development? *Epilepsy & Behavior*. 2011; 20(2):299-307
- 32 74. Wagner JL, Sample PL, Ferguson PL, Pickelsimer EE, Smith GM, Selassie AW.
33 Impact of pediatric epilepsy: voices from a focus group and implications for public
34 policy change. *Epilepsy & Behavior*. 2009; 16(1):161-165
- 35 75. Wallace HK, Solomon JK. Quality of epilepsy treatment and services: the views of
36 women with epilepsy. *Seizure*. 1999; 8(2):81-87
- 37 76. Watkins J, Espie CA, Curtice L, Mantala K, Corp A, Foley J. Development of a
38 measure to assess the impact of epilepsy on people with an intellectual disability: the
39 Glasgow Epilepsy Outcome Scale - Client version (GEOS-C). *Journal of Intellectual
40 Disability Research*. 2006; 50(Pt 3):161-171
- 41 77. Webster M, Gabe J. Diet and identity: being a good parent in the face of
42 contradictions presented by the ketogenic diet. *Sociology of Health and Illness*. 2016;
43 38(1):123-136
- 44 78. Weckesser A, Denny E. Women living with epilepsy, experiences of pregnancy and
45 reproductive health: a review of the literature. *Seizure*. 2013; 22(2):91-98

- 1 79. Westin E, Sund Levander M. Parent's experiences of their children suffering febrile
2 seizures. *Journal of Pediatric Nursing*. 2018; 38:68-73
- 3 80. Widnes SF, Schjott J, Granas AG. Risk perception and medicines information needs
4 in pregnant women with epilepsy--a qualitative study. *Seizure*. 2012; 21(8):597-602
- 5 81. Wilde M, Haslam C. Living with epilepsy: a qualitative study investigating the
6 experiences of young people attending outpatients clinics in Leicester. *Seizure*. 1996;
7 5(1):63-72
- 8 82. Wo SW, Ong LC, Low WY, Lai PSM. Exploring the needs and challenges of parents
9 and their children in childhood epilepsy care: A qualitative study. *Epilepsy & Behavior*.
10 2018; 88:268-276
- 11

Appendices

Appendix A Review protocols

A.1 Review protocol for transition to care from children's and young people's services to adults' services

ID	Field	Content
1.	Review title	How should the transition from children's and young people's services to adults' epilepsy services be managed?
2.	Review question	How should the transition from children's and young people's services to adults' epilepsy services be managed?
3.	Objective	The aim of the review is to identify factors that will help prepare a young person to transition between services. When changing the young person will be expected to become more involved in decision making on the medical issues affecting them. The review will identify information that is needed to aid in the transition.
4.	Searches	The following databases (from inception) will be searched: <ul style="list-style-type: none">• Embase• MEDLINE• PsycINFO• CINAHL Searches will be restricted by: <ul style="list-style-type: none">• English language studies• Human studies

		<p>Other searches:</p> <ul style="list-style-type: none"> • Inclusion lists of systematic reviews <p>The searches may be re-run 6 weeks before the final committee meeting and further studies retrieved for inclusion if relevant.</p> <p>The full search strategies will be published in the final review.</p>
5.	Condition or domain being studied	Children and young people with epilepsy transitioning from children's and young people's services to adult services
6.	Population	<p>Inclusion: children and young people with epilepsy, their families and carers</p> <p>3 groups for access to service strata</p> <ul style="list-style-type: none"> • People with drug resistant epilepsies • People with learning disability (access to service- strata) • People with otherwise well-controlled-epilepsy <p>Exclusion: New-born babies (under 28 days) with acute symptomatic seizures</p>
7.	Intervention/Exposure/Test	Views, opinions and experiences in relation to information, education or support that will help prepare a young person to transition between services.
8.	Comparator/Reference standard/Confounding factors	Not applicable
9.	Types of study to be included	Qualitative studies using any appropriate methodology (e.g., semi-structured interviews or focus groups with ethnography or grounded theory-based analysis) and systematic reviews of qualitative studies will be considered for inclusion. Qualitative studies for transition of care will provide the personal experiences of people with epilepsy and their carers and this will feed into the recommendations.

10.	Other exclusion criteria	<ul style="list-style-type: none"> • Non-English language studies. • Conference abstracts will be excluded because these do not typically provide sufficient information to fully assess risk of bias • Studies conducted in non-OECD countries
11.	Context	Adolescence is a period of rapid change, both physical and psychosocial for any young person. Transition can be a difficult time and should be a planned process of addressing the medical and associated comorbid conditions from paediatric to adult care. Significant apprehension may arise as a young person and their family move from a service where they have built therapeutic alliances to an unknown adult service. Both the paediatric and adult epilepsy teams need to be motivated to provide a successful and safe transition for these patients.
12.	Primary outcomes (critical outcomes)	Themes will be derived from the evidence identified for this review and may include driving, teratogenicity of certain anti-epileptic medications, interaction of anti-epileptic medications with contraception, effect of alcohol/recreational drugs on seizures, psychosocial aspects of epilepsy as people move out of the parental home e.g., to work, to University, independence. Memory, stigma, mental health
13.	Secondary outcomes (important outcomes)	Not applicable.
14.	Data extraction (selection and coding)	<p>EndNote will be used for reference management, sifting, citations and bibliographies. All references identified by the searches and from other sources will be screened for inclusion. 10% of the abstracts will be reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer. The full text of potentially eligible studies will be retrieved and will be assessed in line with the criteria outlined above.</p> <p>A standardised form will be used to extract data from studies (see Developing NICE guidelines: the manual section 6.4) https://www.nice.org.uk/process/pmg20/chapter/reviewing-research-evidence#reviewing-research-evidence</p> <p>Once saturation is considered to have been reached (all the themes are already covered in the data extraction) data from other included papers will not be extracted or critically appraised, but the paper will still be read to check for any additional themes and will be noted in the included studies. The point at which data extraction is reached will be noted within the review.</p>

15.	Risk of bias (quality) assessment	<p>The methodological quality of each study will be assessed using the CASP qualitative checklist and the GRADE CERQual approach used to provide overall risk of bias at outcome level</p> <p>10% of all evidence reviews are quality assured by a senior research fellow. This includes checking:</p> <ul style="list-style-type: none"> • papers were included /excluded appropriately • a sample of the data extractions • correct methods are used to synthesise data • a sample of the risk of bias assessments <p>Disagreements between the review authors over the risk of bias in particular studies will be resolved by discussion, with involvement</p>	
16.	Strategy for data synthesis	<ul style="list-style-type: none"> • CERQual will be used to synthesise data from qualitative studies. <p>The synthesis of qualitative data will follow a thematic analysis approach. Information will be synthesised into main review findings. Results will be presented in a detailed narrative and in table format with summary statements of main review findings.</p> <p>If suggested by the evidence, themes may be reported separately for patients, families and carers.</p>	
17.	Analysis of sub-groups	None.	
18.	Type and method of review	<input type="checkbox"/>	Intervention
		<input type="checkbox"/>	Diagnostic
		<input type="checkbox"/>	Prognostic
		<input checked="" type="checkbox"/>	Qualitative
		<input type="checkbox"/>	Epidemiologic
		<input type="checkbox"/>	Service Delivery
		<input type="checkbox"/>	Other (please specify)

19.	Language	English		
20.	Country	England		
23.	Stage of review at time of this submission	Review stage	Started	Completed
		Preliminary searches	<input type="checkbox"/>	<input checked="" type="checkbox"/>
		Piloting of the study selection process	<input type="checkbox"/>	<input checked="" type="checkbox"/>
		Formal screening of search results against eligibility criteria	<input type="checkbox"/>	<input checked="" type="checkbox"/>
		Data extraction	<input type="checkbox"/>	<input checked="" type="checkbox"/>
		Risk of bias (quality) assessment	<input type="checkbox"/>	<input checked="" type="checkbox"/>
		Data analysis	<input type="checkbox"/>	<input checked="" type="checkbox"/>
24.	Named contact	<p>5a. Named contact Angela Cooper National Guideline Centre Angela.cooper@rcplondon.ac.uk</p> <p>5b Named contact e-mail epilepsies@nice.org.uk</p> <p>5e Organisational affiliation of the review National Institute for Health and Care Excellence (NICE) and the National Guideline Centre</p>		

25.	Review team members	<p>From the National Guideline Centre:</p> <p>Gill Ritchie, Guideline Lead</p> <p>Angela Cooper, Senior Research Fellow</p> <p>Rafina Yarde, Systematic reviewer</p> <p>Margaret Constanti, Senior Health economist</p> <p>Joseph Runicles, Information specialist</p>
26.	Funding sources/sponsor	This systematic review is being completed by the National Guideline Centre which receives funding from NICE.
27.	Conflicts of interest	All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline.
28.	Collaborators	Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of Developing NICE guidelines: the manual . Members of the guideline committee are available on the NICE website: https://www.nice.org.uk/guidance/indevelopment/gid-ng10112/documents
29.	Other registration details	
30.	Reference/URL for published protocol	
31.	Dissemination plans	<p>NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as:</p> <ul style="list-style-type: none"> • notifying registered stakeholders of publication • publicising the guideline through NICE's newsletter and alerts

		<ul style="list-style-type: none"> • issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE. 	
32.	Keywords		
33.	Details of existing review of same topic by same authors		
34.	Current review status	<input checked="" type="checkbox"/>	Ongoing
		<input checked="" type="checkbox"/>	Completed but not published
		<input type="checkbox"/>	Completed and published
		<input type="checkbox"/>	Completed, published and being updated
		<input type="checkbox"/>	Discontinued
35.	Additional information		
36.	Details of final publication	www.nice.org.uk	

Appendix B Literature search strategies

This literature search strategy was used for the following reviews:

- What information and support is needed by people, parents or carers in relation to epilepsy, and when should this be provided?
- How should the transition from children's and young people's services to adults' epilepsy services be managed?

The literature searches for this review are detailed below and complied with the methodology outlined in Developing NICE guidelines: the manual.⁴⁶

For more information, please see the Methodology review published as part of the accompanying documents for this guideline.

B.1 Clinical search literature search strategy

Searches for patient views were run in Medline (OVID), Embase (OVID), CINAHL, Current Nursing and Allied Health Literature (EBSCO) and PsycINFO (ProQuest). Search filters were applied to the search where appropriate.

Table 5: Database date parameters and filters used

Database	Dates searched	Search filter used
Medline (OVID)	1946 – 30 December 2019	Qualitative studies Exclusions
Embase (OVID)	1974 – 30 December 2019	Qualitative studies Exclusions
CINAHL, Current Nursing and Allied Health Literature (EBSCO)	Inception – 30 December 2019	
PsycINFO (ProQuest)	Inception – 30 December 2019	

Medline (Ovid) search terms

1.	exp epilepsy/
2.	seizures/
3.	exp status epilepticus/
4.	seizures, febrile/
5.	(dravet syndrome or epilep* or continuous spike wave or slow sleep or landau kleffner syndrome or lennox gastaut syndrome or infant* spasm* or seizure* or west syndrome).ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.

15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16
18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23
25.	6 not 24
26.	limit 25 to English language
27.	"patient acceptance of health care"/ or exp patient satisfaction/ or consumer health information/ or needs assessment/
28.	Patient Education as Topic/ or exp patients/ or exp family/ or caregivers/ or patient preference/
29.	((information* or advice or advising or advised or support*) adj3 (patient* or need* or requirement* or assess* or seek* or access* or disseminat*)).ti,ab.
30.	(information* adj2 support*).ti,ab.
31.	((client* or patient* or user* or carer* or consumer* or customer* or parent* or famil* or spouse*) adj2 (attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact* or inform* or experience or experiences or opinion* or preference* or focus group*)).ti,ab.
32.	or/27-30
33.	"Continuity of Patient Care"/
34.	Patient Transfer/
35.	Patient Care Planning/
36.	"Delivery of Health Care, Integrated"/
37.	shared care.ti,ab.
38.	shared service*.ti,ab.
39.	((healthcare or care or service*) adj3 integrat*).ti,ab.
40.	transition*.ti.
41.	(transfer? or transferred or transferral or transferring).ti.
42.	(transition* adj10 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department? or patient?)).ab.
43.	((transfer? or transferred or transferral or transferring) adj10 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department? or patient?)).ab.
44.	(transition* and (adult? adj3 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department?))).ab.
45.	((transfer? or transferred or transferral or transferring) and (adult? adj3 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department?))).ab.
46.	(transfer? or transferred or transferal or transferring or transition).ab. and (adult?.ti. or (adult focused or adult oriented).ti,ab.)
47.	(continuity adj3 (care or health care or healthcare or treatment? or therapy or therapies or patient? or "doctor-patient" or nurse patient)).ti,ab.
48.	or/33-47
49.	Qualitative research/ or Narration/ or exp Interviews as Topic/ or exp "Surveys and Questionnaires"/ or Health care surveys/
50.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*).ti,ab.

51.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* adj3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*).ti,ab.
52.	or/49-51
53.	26 and 52 and (32 or 48)

1

Embase (Ovid) search terms

1.	exp epilepsy/
2.	seizure/
3.	epileptic state/
4.	febrile convulsion/
5.	(dravet syndrome or epilep* or continuous spike wave or slow sleep or landau kleffner syndrome or lennox gastaut syndrome or infant* spasm* or seizure* or west syndrome).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	case report/ or case study/
11.	(letter or comment*).ti.
12.	or/7-11
13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental Animal/
19.	animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	limit 23 to English language
25.	patient attitude/ or patient preference/ or patient satisfaction/ or consumer attitude/ or needs assessment/
26.	patient information/ or consumer health information/ or patients/ or family/ or caregivers/
27.	((information* or advice or advising or advised or support*) adj3 (patient* or need* or requirement* or assess* or seek* or access* or disseminat*)).ti,ab.
28.	(information* adj2 support*).ti,ab.
29.	((client* or patient* or user* or carer* or consumer* or customer* or parent* or famil* or spouse*) adj2 (attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact* or inform* or experience or experiences or opinion* or preference* or focus group*)).ti,ab.
30.	patient education/
31.	or/25-30
32.	*patient care/
33.	*patient transport/

34.	patient care planning/
35.	integrated health care system/
36.	shared care.ti,ab.
37.	shared service*.ti,ab.
38.	((healthcare or care or service*) adj3 integrat*).ti,ab.
39.	transition*.ti.
40.	(transfer? or transferred or transferral or transferring).ti.
41.	(transition* adj10 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department? or patient?)).ab.
42.	((transfer? or transferred or transferral or transferring) adj10 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department? or patient?)).ab.
43.	(transition* and (adult? adj3 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department?))).ab.
44.	((transfer? or transferred or transferral or transferring) and (adult? adj3 (care or service? or center? or centre? or clinic? or facility or facilities or unit? or department?))).ab.
45.	(transfer? or transferred or transferal or transferring or transition).ab. and (adult?.ti. or (adult focused or adult oriented).ti,ab.)
46.	(continuity adj3 (care or health care or healthcare or treatment? or therapy or therapies or patient? or "doctor-patient" or nurse patient)).ti,ab.
47.	or/32-46
48.	health survey/ or exp questionnaire/ or exp interview/ or qualitative research/ or narrative/
49.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*).ti,ab.
50.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* adj3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*).ti,ab.
51.	or/48-50
52.	24 and 51 and (31 or 47)

1

CINAHL (EBSCO) search terms

S1.	(MM "Seizures")
S2.	(MH "Status Epilepticus+")
S3.	(MH "Epilepsy+")
S4.	(MH "Convulsions, Febrile")
S5.	(dravet syndrome or epilep* or continuous spike wave or slow sleep or landau kleffner syndrome or lennox gastaut syndrome or infant* spasm* or seizure* or west syndrome)
S6.	S1 OR S2 OR S3 OR S4 OR S5
S7.	(MH "Consumer Satisfaction+") OR (MH "Patient Education") OR (MH "Health Education")
S8.	((information* or advice or advising or advised or support*) n3 (patient* or need* or requirement* or assess* or seek* or access* or disseminat*))
S9.	(information* n2 support*)
S10.	((client* or patient* or user* or carer* or consumer* or customer*) n2 (attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact* or inform* or experience or experiences or opinion*))
S11.	S7 OR S8 OR S9 OR S10
S12.	(MH "Qualitative Studies+")

S13.	(MH "Qualitative Validity+")
S14.	(MH "Interviews+") OR (MH "Focus Groups") OR (MH "Surveys") OR (MH "Questionnaires+")
S15.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*)
S16.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* n3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*)
S17.	S12 OR S13 OR S14 OR S15 OR S16
S18.	S6 AND S11 AND S17

1

PsycINFO (ProQuest) search terms


1.	(MJMAINSUBJECT.EXACT.EXPLODE("Epilepsy") OR MAINSUBJECT.EXACT("Seizures") OR MAINSUBJECT.EXACT("Status Epilepticus") OR TI,AB(dravet syndrome OR epilep* OR continuous spike wave or slow sleep OR landau kleffner syndrome OR lennox gastaut syndrome OR infant* spasm* OR seizure* OR west syndrome)) AND (SU.EXACT("Client Education") OR SU.EXACT.EXPLODE("Client Attitudes") OR TI,AB((information* OR advice OR advising OR advised OR support*) NEAR/3 (patient* OR need* OR requirement* OR assess* OR seek* OR access* OR disseminat*)) OR TI,AB(information* NEAR/2 support*) OR TI,AB((client* OR patient* OR user* OR carer* OR consumer* OR customer*) NEAR/2 (attitud* OR priorit* OR perception* OR preferen* OR expectation* OR choice* OR perspective* OR view* OR satisfact* OR inform* OR experience OR experiences OR opinion*))) AND ((su.exact.explode("qualitative methods") OR su.exact("narratives") OR su.exact.explode("questionnaires") OR su.exact.explode("interviews") OR su.exact.explode("health care services") OR ti,ab(qualitative OR interview* OR focus group* OR theme* OR questionnaire* OR survey*) OR ti,ab(metasynthes* OR meta-synthes* OR metasummar* OR meta-summar* OR metastud* OR meta-stud* OR metathem* OR meta-them* OR ethno* OR emic OR etic OR phenomenolog* OR grounded theory OR constant compar* OR (thematic* NEAR/3 analys*) OR theoretical-sampl* OR purposive-sampl* OR hermeneutic* OR heidegger* OR husserl* OR colaizzi* OR van kaam* OR van manen* OR giorgi* OR glaser* OR strauss* OR ricoeur* OR spiegelberg* OR merleau*)))Limits applied
----	---

2

3

1 Appendix C Qualitative evidence study selection

2 **Figure 1: Flow chart of qualitative study selection for the review of transition to care**
3 **from children's and young people's services to adult services**



4

Appendix D Qualitative evidence

Study	Lewis 2013 ³⁴
Aim	To explore the views of young people with epilepsy (and their parents) about their experience of communication, information and knowledge exchange in two epilepsy services, during transition from children's to adult epilepsy services.
Population	<p>All young people aged between 13–19 years with epilepsy attending the children's, transitional teenager, handover and adult epilepsy clinics were identified in each Case. The qualitative unit of analysis was children's and adult epilepsy services in two District General Hospitals (Case 1 and Case 2) and two embedded units of analysis were young people and their parents within each Case.</p> <p>Young people were included if they had a diagnosis of epilepsy and who had reported experience of a seizure in the past 12 months. Young people with severe learning disabilities (and their parents) were not included in this study as they were considered a different group with specific information and communication needs. Parents of young people with epilepsy who met the inclusion criteria were also invited to participate, n=58.</p>
Setting	Children's and adult epilepsy services in two District General Hospitals. UK.
Study design	A qualitative comparative embedded case study.
Methods and analysis	Semi-structured focus groups and interviews with a two-stage analytical approach was guided by the theoretical framework and propositions. First, thematic analysis using the Framework approach was used to explore experiences of young people with epilepsy and their parents attending epilepsy services. Second, pattern matching using theory-based evaluation was applied to test the theoretical framework and propositions and better understand which models of service delivery worked during transition to adult services, for whom, and in what contexts.
Findings	Clarity: Young people perceived that when healthcare professionals used medical jargon, they were keeping information about their epilepsy from them. Most girls (aged 13-<18 years) had yet to receive information, or could not remember receiving information, about contraception and pregnancy.

Study	Lewis 2013 ³⁴
	<p>Communication barriers: Young people not knowing what questions to ask healthcare professionals about their epilepsy, what they needed to know about their own epilepsy, and parents being unable to seek information with their child present.</p> <p>Information: Continuity of information encompassed the importance of young people remembering facts through repeating information at staged clinical intervals, and the negative impact that lack of continuity of information had on the behaviours of parents and lives of young people.</p> <p>Disengaging from healthcare: Disengaging with healthcare professionals and services was linked with young people and parents' first experience with healthcare professionals in the hand-over clinic.</p> <p>Ongoing engagement with healthcare professionals: On-going and beneficial engagement was helped by befriending and continuity of care.</p> <p>Negative stigma: Stigma originating from peers had a significant impact on the behaviour of girls and boys, which may reinforce negative feelings about continued engagement with services.</p>
Limitations and applicability of evidence	The study had minor methodological limitations providing valuable research and findings. There was a judgement of high confidence in this finding due to no concerns regarding the applicability of this finding to the overarching question.

Study	Schultz 2013 ⁶²
Aim	To explicate processes that parents of adolescents with epilepsy and cognitive impairments undergo as they help their adolescents to transition from paediatric to adult health care. Because parents are the healthcare agents for their adolescents with cognitive impairments and epilepsy, an improved understanding of the processes that these parents experience as they help their adolescents transition from paediatric to adult healthcare is needed to develop effective, individualised transition plans.
Population	<p>Participants were parents or guardians (mothers, fathers or both) of adolescents 18 years or older with epilepsy and severe to profound cognitive impairments who had transitioned to an adult provider more than 1 year ago but less than 5 years ago and spoke English.</p> <p>n=7</p>

Study	Schultz 2013 ⁶²
Setting	Participant's home, USA
Study design	Qualitative interview study
Methods and analysis	<p>Semi-structured individual interviews with grounded theory methodology. Individual interviews were conducted from June through September 2009 using open-ended semi-structured questions. All interviews were conducted in the participant's home, audio-taped, transcribed verbatim, and reviewed to ensure accuracy.</p> <p>Data collection and analysis occurred concurrently according to the tenets of grounded theory methodology using coding and constant comparison analysis. Open coding was done to identify conceptual labels that described participants' experiences.</p>
Themes with findings	<p>Turmoil: Experienced by all parents. They experienced feelings of fear, rejection or uncertainty.</p> <p>Parents as advocates: Experienced by all parents. They assumed the role of protector and information gatherer.</p> <p>Information: Continuity of information encompassed the importance of young people remembering facts through repeating information at staged clinical intervals and the negative impact that lack of continuity of information had on the behaviours of parents and lives of young people.</p> <p>Waiting: Discontinuity between agencies resulted in waiting for answers before being able to move forward.</p> <p>Transition sparked by crisis: Transitioning is not planned and is usually a result of a health issue or change in insurance.</p>
Limitations and applicability of evidence	<p>The study had moderate methodological limitations providing valuable research and findings. There was a judgement of moderate confidence in this finding due to concerns regarding the ethical considerations and the relationship between the researcher and participants' being adequately considered. There were no concerns regarding the applicability of the evidence.</p>

1 Appendix E Excluded studies

2 E.1 Clinical studies

3 **Table 6: Studies excluded from the qualitative review**

Reference	Reason for exclusion
Abubakar 2015 ¹	Incorrect population, included those without epilepsy.
Alkhamra 2012 ²	Incorrect study design, questionnaire.
Amjad 2016 ³	Included in information review, no information on transition.
Andrade 2017 ⁴	Incorrect study design, recommendations.
Aytch 2001 ⁵	Incorrect population, included those without epilepsy.
Baca 2018 ⁶	Incorrect study design, not qualitative.
Baird 2003 ⁷	Quantitative study with some themes reported, doesn't mention quantitative analysis.
Benson 2015 ⁸	Included in information review, no information on transition.
Benson 2016 ¹⁰	Included in information review, no information on transition.
Benson 2017 ⁹	Included in information review, no information on transition.
Blank 2014 ¹¹	Incorrect study objectives, addressing why elderly patients don't regularly attend clinic.
Burke 2018 ¹²	Systematic review, references individually assessed, not all looked at epilepsy, those that did have already been ordered.
Butau 1993 ¹³	Incorrect study design: close ended questions, no clear themes identified.
Camfield 2017 ¹⁴	Incorrect study design, conference proceedings.
Canvin 2006 ¹⁵	Incorrect objective; asking opinions on joining an epilepsy trial.
Chen 2010 ¹⁶	Included in information review, no information on transition.
Chew 2018 ¹⁷	Included in information review, no information on transition.
Chiu 2014 ¹⁸	Included in information review, no information on transition.
Chong 2016 ¹⁹	Systematic review: not all qualitative studies, already have the ones that are.
Collard 2019 ²⁰	Systematic review, not all studies qualitative, ordered relevant ones.
Eklund, 2003 ²¹	Included in information review; no information on transition.
Elliot, 2005 ²²	Included in information review, no information on transition.
Gauffin 2015 ²³	Included in information review, no information on transition.

Reference	Reason for exclusion
Geerlings 2015 ²⁴	Systematic review: not all qualitative studies, already have the ones that are.
Hames, 2009 ²⁵	Incorrect study design, written accounts.
Harden 2015 ²⁷	Included in information review, no information on transition.
Harden 2016 ²⁶	Systematic review, not all studies qualitative, ordered relevant ones.
Heath 2016 ²⁸	Incorrect study objective, about surgery.
Hightower, 2002 ²⁹	Included in information review, no information on transition.
Joung 2019 ³⁰	Included in information review; no information on transition.
Kaddumukasa 2019 ³¹	Included in information review, no information on transition.
Kampra 2017 ³²	Included in information review, no information on transition.
Kuchenbuch 2013 ³³	Incorrect study design, questionnaire.
Louik, 2017 ³⁵	Incorrect study design, questionnaire.
MacLeod, 2009 ³⁶	Included in information review, no information on transition.
Mc Govern 2018 ³⁷	Incorrect study design, actual transition programme.
McAuley, 2012 ³⁸	Incorrect study objective.
McEwan, 2004 ³⁹	Included in information review, no information on transition.
McNelis 2007 ⁴⁰	Included in information review, no information on transition.
Mengoni 2016 ⁴¹	Included in information review, no information on transition.
Miller, 2014 ⁴²	Incorrect population, providers e.g., neurologists and doctors.
Moffat, 2009 ⁴³	Included in information review, no information on transition.
Mu 2008 ⁴⁴	Included in information review, no information on transition.
Nashef, 1998 ⁴⁵	Incorrect population and study design, not all SUDEP, not qualitative data analysis.
Ninnoni 2019 ⁴⁷	Included in information review, no information on transition.
Nisbet, 2017 ⁴⁸	Incorrect population, neurologists and doctors.
O'Toole 2016 ⁴⁹	Included in information review, no information on transition.
Pashley, 2009 ⁵⁰	Included in information review, no information on transition.
Prinjha 2005 ⁵¹	Included in information review, no information on transition.
Prinjha, 2005 ⁵¹	Duplicate.
Ramachandran Nair, 2016 ⁵⁴	Incorrect study comparisons.

Reference	Reason for exclusion
Ramachandrannair, 2013 ⁵³	Included in information review, no information on transition.
RamachandranNair, 2016 ⁵²	Included in information review, no information on transition.
Ridsdale, 2012 ⁵⁵	Incorrect study objective.
Roberts, 2011 ⁵⁶	Included in information review, no information on transition.
Ronen, 1999 ⁵⁷	Included in information review, no information on transition.
Rushe, 2011 ⁵⁸	Unavailable.
Saburi 2006 ⁵⁹	Incorrect study design, survey
Sample 2006 ⁶⁰	Included in information review, no information on transition.
Scheffer 2014 ⁶¹	Incorrect study design, not qualitative study.
Shih 2018 ⁶³	Incorrect study objective, discussing people considering surgery.
Smith 2014 ⁶⁴	Included in information review, no information on transition.
Snape 2009 ⁶⁵	Included in information review, no information on transition.
Sonecha 2015 ⁶⁶	Incorrect study objective.
Swarztrauber 2003 ⁶⁷	Incorrect study objective, more about surgery.
Tanaka 2018 ⁶⁸	Incorrect study population, people with mental health issues.
Thompson 2013 ⁷⁰	Incorrect study design, online survey.
Thompson, 2008 ⁶⁹	Included in information review, no information on transition.
Tonberg 2015 ⁷¹	Included in information review, no information on transition.
Van Naarden Braun 2006 ⁷²	Incorrect study design, cross sectional study.
Varley, 2011 ⁷³	Included in information review, no information on transition.
Wagner, 2009 ⁷⁴	Included in information review, no information on transition.
Wallace 1999 ⁷⁵	Included in information review, no information on transition.
Watkins 2006 ⁷⁶	Incorrect study design, not qualitative.
Webster 2016 ⁷⁷	Included in information review, no information on transition.
Weckesser 2013 ⁷⁸	Systematic review, not all papers looking at epilepsy, relevant ones ordered/we have.
Westin 2018 ⁷⁹	Included in information review, no information on transition.
Widnes 2012 ⁸⁰	Included in information review; no information on transition
Wilde 1996 ⁸¹	Included in information review, no information on transition.
Wo 2018 ⁸²	Included in information review, no information on transition.

