

Draft for consultation

Depression in adults: treatment and management

Chapter 16 References

NICE Guideline <...>

References

March 2018

Draft for consultation

*Developed by the National Guideline
Alliance, hosted by the Royal College of
Obstetricians and Gynaecologists*

Disclaimer

Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and/or their guardian or carer.

Copyright

National Institute for Health and Care Excellence [2018]. All rights reserved. Subject to Notice of Rights.

Contents

16 References	5
----------------------------	----------

Update 2018

16 References

- 2 Abas, M. (1996). Depression and anxiety among older Caribbean people in the UK:
3 screening, unmet need and the provision of appropriate services. International Journal of
4 Geriatric Psychiatry, 11(4), 377-382.
- 5 Abas, M. A., Phillips, C., Carter, J., Walter, J., Banerjee, S., & Levy, R. (1998). Culturally
6 sensitive validation of screening questionnaires for depression in older African-Caribbean
7 people living in south London. Br J Psychiatry, 173, 249-254.
- 8 Abbass, A., Town, J., & Driessen, E. (2011). The efficacy of short-term psychodynamic
9 psychotherapy for depressive disorders with comorbid personality disorder. Psychiatry,
10 74(1), 58-71.
- 11 Addis, M. E. (2008). Gender and depression in men. Clinical Psychology: Science and
12 Practice, 15(3), 153-168.
- 13 Adler, D., Bungay, K., Wilson, I., Pei, Y., Supran, S., Peckham, E., Cynn, D.J., & Rogers,
14 W.H. (2004). The impact of a pharmacist intervention on 6-month outcomes in depressed
15 primary care patients. General Hospital Psychiatry, 26, 199-209.
- 16 Agosti V, Ocepek-Welikson K. The efficacy of imipramine and psychotherapy in early-onset
17 chronic depression: a reanalysis of the National Institute of Mental Health Treatment of
18 Depression Collaborative Research Program. Journal of Affective Disorders. 1997 May
19 1;43(3):181-6.
- 20 AGREE-Collaboration. (2003). Development and validation of an international appraisal
21 instrument for assessing the quality of clinical practice guidelines: the AGREE project.
22 Quality and Safety in Health Care, 12(1), 18-23.
- 23 Akiskal, H. S. (1986). A developmental perspective on recurrent mood disorders: a review of
24 studies in man. Psychopharmacol Bull, 22(3), 579-586.
- 25 Alexopoulos, G. S., Meyers, B. S., Young, R. C., Kalayam, B., Kakuma, T., Gabrielle,
26 M....Hull, J. (2000). Executive dysfunction and long-term outcomes of geriatric depression.
27 Archives of General Psychiatry, 57(3), 285-290.
- 28 Alexopoulos GS, Raue PJ, McCulloch C, Kanellopoulos D, Seirup JK, Sirey JA, Banerjee S,
29 Kirosses DN, Areán PA. Clinical case management versus case management with problem-
30 solving therapy in low-income, disabled elders with major depression: a randomized clinical
31 trial. The American Journal of Geriatric Psychiatry. 2016 Jan 31; 24(1):50-9.
- 32 Aljumah K, Hassali MA. Impact of pharmacist intervention on adherence and measurable
33 patient outcomes among depressed patients: a randomised controlled study. BMC
34 psychiatry. 2015 Sep 16; 15(1):219.
- 35 Allart-Van Dam, E., Hosman, C. M. H., Hoogduin, C. A. L., & Schaap, C. P. D. R. (2003) The
36 Coping with Depression course: Short-term outcomes and mediating effects of a randomized
37 controlled trial in the treatment of subclinical depression. Behavior Therapy, 34, 381-396.
- 38 Altamura AC, Dell'Osso B, Buoli M, Zanoni S, Mundo E. Intravenous augmentative
39 citalopram versus clomipramine in partial/nonresponder depressed patients: a short-term,
40 low dose, randomized, placebo-controlled study. Journal of Clinical Psychopharmacology.
41 2008 Aug 1; 28(4):406-10.
- 42 Altamura AC, Montgomery SA, Wernicke JF. The evidence for 20mg a day of fluoxetine as
43 the optimal dose in the treatment of depression. British Journal of Psychiatry -
44 Supplementum 1988 ;(3):109-112.

- 1 Altman, D. G., & Bland, J. M. (1994). Diagnostic tests 2: Predictive values. *BMJ*, 309(6947), 102.
- 3 Altman, D. G., & Bland, J. M. (1994). Diagnostic tests. 1: Sensitivity and specificity. *BMJ*, 308(6943), 1552.
- 5 American Psychiatric Association. (1994). Diagnostic and statistical manual of mental disorders: DSM-IV (4th ed.). Washington, DC: American Psychiatric Association.
- 7 American Psychiatric Association. (2000). Diagnostic and statistical manual of mental disorders: DSM-IV-TR (4th ed.). Washington, DC: American Psychiatric Association.
- 9 American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders: DSM-5 (5th ed.). Washington, D.C.: American Psychiatric Association.
- 11 Amini H, Aghayan S, Jalili SA, Akhondzadeh S, Yahyazadeh O, Pakravan-Nejad M.
12 Comparison of mirtazapine and fluoxetine in the treatment of major depressive disorder: a
13 double-blind, randomized trial. *Journal of Clinical Pharmacy and Therapeutics* 2005;
14 30(2):133-8.
- 15 Amore M, Jori MC. Faster response on amisulpride 50 mg versus sertraline 50-100 mg in
16 patients with dysthymia or double depression: a randomized, double-blind, parallel group
17 study. *International clinical psychopharmacology*. 2001 Nov 1; 16(6):317-24.
- 18 Amsterdam, J., Case, W., Csanalosi, E., Singer, M., & Rickels, K. (1986). A double-blind
19 comparative trial of zimelidine, amitriptyline, and placebo in patients with mixed anxiety and
20 depression. *Pharmacopsychiatry*, 19(03), 115-119.
- 21 Anand, A. S., & Cochrane, R. (2005). The mental health status of South Asian women in
22 Britain. *Psychology and Developing Societies*, 17(2), 195-214.
- 23 Anderson, H. D., Pace, W. D., Libby, A. M., West, D. R., & Valuck, R. J. (2012). Rates of 5
24 common antidepressant side effects among new adult and adolescent cases of depression:
25 a retrospective US claims study. *Clin Ther*, 34(1), 113-123.
- 26 Anderson, I. M., & Fergusson, G. (2013). Mechanism of action of ECT. In E. A. Waite J (Ed.),
27 The ECT Handbook. 3rd edn. (3rd ed., pp. 1-7): Royal College of Psychiatrists.
- 28 Anderson, I. M., Ferrier, I. N., Baldwin, R. C., Cowen, P. J., Howard, L., Lewis, G....Tylee, A.
29 (2008). Evidence-based guidelines for treating depressive disorders with antidepressants: a
30 revision of the 2000 British Association for Psychopharmacology guidelines. *J
31 Psychopharmacol*, 22(4), 343-396.
- 32 Andersson, G., Bergstrom, J., Hollandare, F., Carlbring, P., Kaldo, V., & Ekselius, L. (2005)
33 Internet-based self-help for depression: randomised controlled trial. *British Journal of
34 Psychiatry*, 187, 456-461.
- 35 Andrade, L., Caraveo-Anduaga, J. J., Berglund, P., Bijl, R. V., De Graaf, R., Vollebergh, W.,
36 Wittchen, H. U. (2003). The epidemiology of major depressive episodes: results from the
37 International Consortium of Psychiatric Epidemiology (ICPE) Surveys. *Int J Methods
38 Psychiatr Res*, 12(1), 3-21.
- 39 Andreeescu, C., Glick, R. M., Emeremni, C. A., Houck, P. R., & Mulsant, B. H. (2011).
40 Acupuncture for the treatment of major depressive disorder: a randomized controlled trial.
41 *Journal of Clinical Psychiatry*, 72(8), 1129-1135.
- 42 Andreeescu, C., Mulsant, B. H., Peasley-Miklus, C., Rothschild, A. J., Flint, A. J., Heo, M., . . .
43 Group, S.-P. S. (2007). Persisting low use of antipsychotics in the treatment of major
44 depressive disorder with psychotic features. *J Clin Psychiatry*, 68(2), 194-200.

- 1 Andreoli V, Caillard V, Deo RS, Rybakowski JK, Versiani M. Reboxetine, a new noradrenaline selective antidepressant, is at least as effective as fluoxetine in the treatment of depression. *Journal of Clinical Psychopharmacology* 2002; 22(4):393-399.
- 4 Andrews, G., Anderson, T. M., Slade, T., & Sunderland, M. (2008). Classification of anxiety and depressive disorders: problems and solutions. *Depression and Anxiety*, 25(4), 274-281.
- 6 Andrews, G., & Jenkins, R. (Eds.). (1999). *Management of Mental Disorders* (UK edn, vol. 1). Sydney: WHO Collaborating Centre for Mental Health and Substance Misuse.
- 8 Anglin, R., Yuan, Y., Moayyedi, P., Tse, F., Armstrong, D., & Leontiadis, G. I. (2014). Risk of upper gastrointestinal bleeding with selective serotonin reuptake inhibitors with or without concurrent nonsteroidal anti-inflammatory use: a systematic review and meta-analysis. *Am J Gastroenterol*, 109(6), 811-819.
- 12 Anisman, H., Ravindran, A. V., Griffiths, J., & Merali, Z. (1999). Interleukin-1 β production in dysthymia before and after pharmacotherapy. *Biological Psychiatry*, 46(12), 1649-1655.
- 14 Anton, R. F., Jr., & Burch, E. A., Jr. (1990). Amoxapine versus amitriptyline combined with perphenazine in the treatment of psychotic depression. *Am J Psychiatry*, 147(9), 1203-1208.
- 16 Apil, S. R., Hoencamp, E., Judith Haffmans, P. M., & Spinhoven, P. (2012). A stepped care relapse prevention program for depression in older people: a randomized controlled trial. *International Journal of Geriatric Psychiatry*, 27(6), 583-591.
- 19 Appelberg, B. G., Syvälähti, E., Koskinen, T. E., Mehtonen, O.-P., Muuronen, T. T., & Naukkarinen, H. H. (2001). Patients with severe depression may benefit from buspirone augmentation of selective serotonin reuptake inhibitors: results from a placebo-controlled, randomized, double-blind, placebo wash-in study. *The Journal of Clinical Psychiatry*, 62(6), 448-452.
- 24 Aragones E, Lopez-Cortacans G, Sanchez-Iriso E, Pinol JL, Caballero A, Salvador-Carulla L, Cabasés J. Cost-effectiveness analysis of a collaborative care programme for depression in primary care. *Journal of Affective Disorders* 2014; 159:85-93.
- 27 Aragonès, E., Piñol, J. L., Caballero, A., López-Cortacans, G., Casaus, P., Hernández, J. M., Folch, S. (2012). Effectiveness of a multi-component programme for managing depression in primary care: a cluster randomized trial. The INDI project. *Journal of Affective Disorders*, 142(1-3), 297-305.
- 31 Aragones, E., Pinol, J. L., & Labad, A. (2006). The overdiagnosis of depression in non-depressed patients in primary care. *Fam Pract*, 23(3), 363-368.
- 33 Araya, R., Rojas, G., Fritsch, R., Gaete, J., Rojas, M., Simon, G., & Peters, T.J. (2003). Treating depression in primary care in low-income women in Santiago, Chile: a randomised controlled trial. *Lancet*, 361, 995-1000.
- 36 Areán PA, Ayalon L, Jin C, McCulloch CE, Linkins K, Chen H, McDonnell-Herr B, Levkoff S, Estes C. Integrated specialty mental health care among older minorities improves access but not outcomes: results of the PRISMe study. *International journal of geriatric psychiatry*. 2008 Oct 1; 23(10):1086-92.
- 40 Arnow, B. A., & Constantino, M. J. (2003). Effectiveness of psychotherapy and combination treatment for chronic depression. *J Clin Psychol*, 59(8), 893-905.
- 42 Arvidsdotter T, Marklund B, Taft C. Effects of an integrative treatment, therapeutic acupuncture and conventional treatment in alleviating psychological distress in primary care patients-a pragmatic randomized controlled trial. *BMC complementary and alternative medicine*. 2013 Nov 7; 13(1):308.

- 1 Ashouri A, Atef-Vahid M-K, Gharaee B, Rasoulian M. Effectiveness of meta-cognitive and cognitive-behavioral therapy in patients with major depressive disorder. *Iranian journal of psychiatry and behavioral sciences*. 2013; 7:24-34.
- 4 Avery, D., Kizer, D., Bolte, M., & Hellekson, C. (2001). Bright light therapy of subsyndromal seasonal affective disorder in the workplace: morning vs. afternoon exposure. *Acta Psychiatrica Scandinavica*, 103(4), 267-274.
- 7 Avery, D. H., Bolte, M. A., Dager, S. R., Wilson, L. G., Weyer, M., Cox, G. B., & Dunner, D. L. (1993). Dawn simulation treatment of winter depression: a controlled study. *American Journal of Psychiatry*, 150, 113-113.
- 10 Bagby, R.M., Quily, L.C., Segal, Z.V., McBride, C.C., Kennedy, S.H., & Costa, P.T. (2008) Personality and differential treatment response in major depression: A randomized controlled trial comparing cognitive-behavioural therapy and pharmacotherapy. *The Canadian Journal of Psychiatry*, 53, 361-370.
- 14 Baker AL, Kavanagh DJ, Kay-Lambkin FJ, Hunt SA, Lewin TJ, Carr VJ, Connolly J. Randomized controlled trial of cognitive-behavioural therapy for coexisting depression and alcohol problems: short-term outcome. *Addiction*. 2010 Jan 1; 105(1):87-99.
- 17 Bakish, D., Bradwejn, J., Nair, N., McClure, J., Remick, R., & Bulger, L. (1992). A comparison of moclobemide, amitriptyline and placebo in depression: a Canadian multicentre study. *Psychopharmacology*, 106, S98-S101.
- 20 Bakish, D., Lapierre, Y., Weinstein, R., Klein, J., Wiens, A., Jones, B.... Blanchard, A. (1993). Ritanserin, imipramine, and placebo in the treatment of dysthymic disorder. *Journal of Clinical Psychopharmacology*, 13(6), 409-414.
- 23 Baldwin, R. (2012). Mood disorders in the elderly New Oxford Textbook of Psychiatry (pp. 1551-1558): Oxford University Press (OUP).
- 25 Balestrieri A, Benassi P, Cassano GB, et al. Clinical comparative evaluation of maprotiline, a new antidepressant drug. *International Pharmacopsychiatry* 1972; 6:236-248.
- 27 Barber, J. P., et al. (2012) Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. *The Journal of clinical psychiatry* 73, 66-73.
- 30 Ballard, C., Bannister, C., Solis, M., Oyebode, F., & Wilcock, G. (1996). The prevalence, associations and symptoms of depression amongst dementia sufferers. *J Affect Disord*, 36(3-4), 135-144.
- 33 Ballesteros, J., & Callado, L. F. (2004). Effectiveness of pindolol plus serotonin uptake inhibitors in depression: a meta-analysis of early and late outcomes from randomised controlled trials. *Journal of Affective Disorders*, 79(1), 137-147.
- 36 Barbee, J. G., Thompson, T. R., Jamhour, N. J., Stewart, J. W., Conrad, E. J., Reimherr, F. W., ... Shelton, R. C. (2011). A double-blind placebo-controlled trial of lamotrigine as an antidepressant augmentation agent in treatment-refractory unipolar depression. *The Journal of Clinical Psychiatry*, 72(10), 1405-1412.
- 40 Barge-Schaapveld, D. Q. & Nicolson, N. A. (2002). Effects of antidepressant treatment on the quality of daily life: an experience sampling study. *Journal of Clinical Psychiatry*, 63, 477-485.
- 43 Barnhofer, T., Crane, C., Hargus, E., Amarasinghe, M., Winder, R., & Williams, J. M. (2009). Mindfulness-based cognitive therapy as a treatment for chronic depression: A preliminary study. *Behaviour research and therapy*, 47(5), 366-373.

- 1 Barr, B., Taylor-Robinson, D., Scott-Samuel, A., McKee, M., & Stuckler, D. (2012). Suicides associated with the 2008-10 economic recession in England: time trend analysis. *BMJ*, 345(aug13 2), e5142-e5142.
- 4 Bartels, S. J., Coakley, E. H., Zubritsky, C., Ware, J. H., Miles, K. M., Arean, P. A.,...
5 Investigators, P.-E. (2004). Improving access to geriatric mental health services: a
6 randomized trial comparing treatment engagement with integrated versus enhanced referral
7 care for depression, anxiety, and at-risk alcohol use. *Am J Psychiatry*, 161(8), 1455-1462.
- 8 Bastos AG, Guimaraes LS, Trentini CM. The efficacy of long-term psychodynamic
9 psychotherapy, fluoxetine and their combination in the outpatient treatment of depression.
10 *Psychotherapy Research*. 2015 Sep 3; 25(5):612-24.
- 11 Bauer, M., Adli, M., Bschor, T., Pilhatsch, M., Pfennig, A., Sasse, J., ... Lewitzka, U. (2010).
12 Lithium's emerging role in the treatment of refractory major depressive episodes:
13 augmentation of antidepressants. *Neuropsychobiology*, 62(1), 36-42.
- 14 Bauer, M., Bschor, T., Kunz, D., Berghöfer, A., Ströhle, A., & Müller-Oerlinghausen, B.
15 (2000). Double-blind, placebo-controlled trial of the use of lithium to augment antidepressant
16 medication in continuation treatment of unipolar major depression. *American Journal of
17 Psychiatry*, 157(9), 1429-1435.
- 18 Bauer, M., Pfennig, A., Linden, M., Smolka, M. N., Neu, P., & Adli, M. (2009). Efficacy of an
19 algorithm-guided treatment compared with treatment as usual: a randomized, controlled
20 study of inpatients with depression. *Journal of Clinical Psychopharmacology*, 29(4), 327-333.
- 21 Bauer, M., Pfennig, A., Severus, E., Whybrow, P. C., Angst, J., Moller, H. J., ... Yamada, K.
22 (2013). World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for
23 Biological Treatment of Unipolar Depressive Disorders, Part 1: Update 2013 on the acute
24 and continuation treatment of unipolar depressive disorders. *World Journal of Biological
25 Psychiatry*, 14(5), 334-385
- 26 Bauer, M., Pretorius, H. W., Constant, E. L., Earley, W. R., Szamosi, J., & Brecher, M.
27 (2009). Extended-release quetiapine as adjunct to an antidepressant in patients with major
28 depressive disorder: results of a randomized, placebo-controlled, double-blind study. *Journal
29 of Clinical Psychiatry*, 70(4), 540-549.
- 30 Bauer, M. S., & Dunner, D. L. (1993). Validity of seasonal pattern as a modifier for recurrent
31 mood disorders for DSM-IV. *Comprehensive Psychiatry*, 34(3), 159-170
- 32 Baumann, P., Nil, R., Souche, A., Montaldi, S., Baettig, D., Lambert, S., ... Jonzier-Perey, M.
33 (1996). A double-blind, placebo-controlled study of citalopram with and without lithium in the
34 treatment of therapy-resistant depressive patients: a clinical, pharmacokinetic, and
35 pharmacogenetic investigation. *Journal of Clinical Psychopharmacology*, 16(4), 307-314.
- 36 Be, D., Whisman, M. A., & Uebelacker, L. A. (2013). Prospective associations between
37 marital adjustment and life satisfaction. *Personal Relationships*, 20(4), 728-739.
- 38 Beach, M. C., Gary, T. L., Price, E. G., Robinson, K., Gozu, A., Palacio, A., ... Cooper, L. A.
39 (2006). Improving health care quality for racial/ethnic minorities: a systematic review of the
40 best evidence regarding provider and organization interventions. *BMC Public Health*, 6, 104.
- 41 Beach, S. R., Kostis, W. J., Celano, C. M., Januzzi, J. L., Ruskin, J. N., Noseworthy, P. A., &
42 Huffman, J. C. (2014). Meta-analysis of selective serotonin reuptake inhibitor-associated QTc
43 prolongation. *J Clin Psychiatry*, 75(5), e441-449.
- 44 Beach, S. R., & O'Leary, K. D. (1992). Treating depression in the context of marital discord:
45 Outcome and predictors of response of marital therapy versus cognitive therapy. *Behavior
46 therapy*, 23(4), 507-528.

- 1 Beach, S. R. H., Katz, J., Kim, S., & Brody, G. H. (2003). Prospective effects of marital satisfaction on depressive symptoms in established marriages: a dyadic model. *Journal of Social and Personal Relationships*, 20(3), 355-371.
- 4 Beach, S. R. H., Kim, S., Cercone-Keeney, J., Gupta, M., Arias, I., & Brody, G. H. (2004). Physical aggression and depressive symptoms: gender asymmetry in effects? *Journal of Social and Personal Relationships*, 21(3), 341-360.
- 7 Beasley CM, Jr. & Potvin JH. Fluoxetine: activating and sedating effects. [Review] [13 refs].
8 International Clinical Psychopharmacology 1993; 8(4):271-275.
- 9 Beasley CM, Jr., Bosomworth JC, Wernicke JF. Fluoxetine: relationships among dose,
10 response, adverse events, and plasma concentrations in the treatment of depression.
11 Psychopharmacology Bulletin 1990; 26(1):18-24.
- 12 Beasley CM, Nilsson ME, Koke SC, Gonzales JS. Efficacy, adverse events, and treatment
13 discontinuations in fluoxetine clinical studies of major depression: A meta-analysis of the 20-
14 mg/day dose. *Journal of Clinical Psychiatry* 2000; 61(10).
- 15 Beasley CM Jr, Sayler ME, Potvin JH. Fluoxetine versus amitriptyline in the treatment of
16 major depression: a multicenter trial. *International Clinical Psychopharmacology* 1993 Fall;
17 8(3):143-9.
- 18 Beasley, C. M. J., Sayler, M. E., Bosomworth, J. C., & Wernicke, J. F. (1991). High-dose
19 fluoxetine: efficacy and activating-sedating effects in agitated and retarded depression.
20 *Journal of Clinical Psychopharmacology*, 11, 166-174.
- 21 Beasley CM, Jr., Sayler ME, Weiss AM, Potvin JH. Fluoxetine: activating and sedating
22 effects at multiple fixed doses. *Journal of Clinical Psychopharmacology* 1992; 12(5):328-333.
- 23 Beck, A. T. (1964). Thinking and depression. *Archives of General Psychiatry*, 10(6), 561.
- 24 Beck, A. T. (2008). The Evolution of the Cognitive Model of Depression and Its
25 Neurobiological Correlates. *American Journal of Psychiatry*, 165(8), 969-977.
- 26 Beck, A. T., A., S. R., & Brown, G. K. (1996). Manual for the Beck Depression Inventory-II.
27 San Antonio. Texas: Psychological Corporation.
- 28 Beck, A. T., Guth, D., Steer, R. A., & Ball, R. (1997). Screening for major depression
29 disorders in medical inpatients with the Beck Depression Inventory for Primary Care. *Behav
Res Ther*, 35(8), 785-791.
- 31 Beck AT, Hollon SD, Young JE, Bedrosian RC, Budenz D. Treatment of depression with
32 cognitive therapy and amitriptyline. *Archives of general psychiatry*. 1985 Feb 1; 42(2):142-8.
- 33 Beck, A. T., Rial, W. Y., & Rickels, K. (1974). Short form of depression inventory: cross-
34 validation. *Psychological reports*.
- 35 Beck, A. T., Rush, A. J., & Shaw, B. F. (1979). Cognitive therapy of depression. New York:
36 Guilford Press.
- 37 Beck, A. T., Steer, R., & Brown, G. (1996). Beck Depression Inventory Manual. The
38 Psychological Corporation. San Antonio, TX.
- 39 Beck, A. T., Steer, R. A., & Brown, G. K. (2000). Beck Depression Inventory FastScree for
40 Medical Patients.
- 41 Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for
42 measuring depression. *Arch Gen Psychiatry*, 4, 561-571.

- 1 Bedi N, Chilvers C, Churchill R, Dewey M, Duggan C, Fielding K, Gretton V, Miller P,
- 2 Harrison G, Lee A, Williams I. Assessing effectiveness of treatment of depression in primary care. Partially randomised preference trial. *British Journal of Psychiatry* 2000; 177:312-318.
- 3 Beeber, L. S., Holditch-Davis, D., Perreira, K., A Schwartz, T., Lewis, V., Blanchard, H., ... Goldman, B. D. (2010). Short-term in-home intervention reduces depressive symptoms in Early Head Start Latina mothers of infants and toddlers. *Research in nursing & health*, 33(1), 60-76.
- 4 Behnke, K., Jensen, G. S., Graubaum, H.-J., & Gruenwald, J. (2002). Hypericum perforatum versus fluoxetine in the treatment of mild to moderate depression. *Advances in therapy*, 19(1), 43-52.
- 5 Bell, D. N. F., & Blanchflower, D. G. (2011). Young people and the Great Recession. *Oxford Review of Economic Policy*, 27(2), 241-267.
- 6 Bellino S, Barzega G, Bogetto F, Maina G, Venturello S, Ravizza L. An open-label, randomized, prospective comparison of sertraline and amisulpride in the treatment of dysthymia in the elderly. *Current therapeutic research*. 1997 Oct 31; 58(10):798-808.
- 7 Benedict, A., Arellano, J., De Cock, E., & Baird, J. (2010). Economic evaluation of duloxetine versus serotonin selective reuptake inhibitors and venlafaxine XR in treating major depressive disorder in Scotland. *Journal of Affective Disorders*, 120, 94-104.
- 8 Bentall, D. P., Richard. (1999). The medicalisation of misery: A critical realist analysis of the concept of depression. *Journal of Mental Health*, 8(3), 261-274.
- 9 Berger, T., et al. (2011) Internet-based treatment of depression: a randomized controlled trial comparing guided with unguided self-help. *Cognitive behaviour therapy* 40, 251-266.
- 10 Berghöfer, A., Hartwich, A., Bauer, M., Unützer, J., Willich, S. N., & Pfennig, A. (2012). Efficacy of a systematic depression management program in high utilizers of primary care: a randomized trial. *BMC Health Services Research*, 12, 298.
- 11 Bergmann, R., Nussner, J., & Demling, J. (1993). Therapy of minor and moderate depressions. *Ts Neurologie Psychiatrie*, 7(4), 235-240.
- 12 Berlin, J. A. (1997). Does blinding of readers affect the results of meta-analyses? University of Pennsylvania Meta-analysis Blinding Study Group. *Lancet*, 350(9072), 185-186.
- 13 Berman, R. M., Fava, M., Thase, M. E., Trivedi, M. H., Swanink, R., McQuade, R. D., ... Marcus, R. N. (2009). Aripiprazole augmentation in major depressive disorder: a double-blind, placebo-controlled study in patients with inadequate response to antidepressants. *CNS Spectrums*, 14(4), 197-206.
- 14 Berman, R. M., Marcus, R. N., Swanink, R., McQuade, R. D., Carson, W. H., Corey-Lisle, P. K., & Khan, A. (2007). The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: a multicenter, randomized, double-blind, placebo-controlled study. *The Journal of Clinical Psychiatry*, 68(6), 843-853.
- 15 Bersani G, Rapisarda V, Ciani N, Bertolino A, Sorge G. A double-blind comparative study of sertraline and amitriptyline in outpatients with major depressive episodes. *HUM PSYCHOPHARMACOL* 1994; 9(1):63-68.
- 16 Beutler LE, Engle D, Mohr D, Daldrup RJ, Bergan J, Meredith K, Merry W. Predictors of differential response to cognitive experiential and self-directed psychotherapeutic procedures. *Journal of Consulting and Clinical Psychology* 1991; 59:333-340.

- 1 Bhargava, J. and Z. Y. Khan (2012) Comparative evaluation of the efficacy and side effects of imipramine, sertraline and an ayurvedic formulation in patients of depression. *Journal of clinical and diagnostic research* 6, 220-225.
- 4 Bhugra, D., & Cochrane, R. (2000). Psychiatry in a multi-ethnic context. In: Bhugra D, Cochrane R (eds). *Psychiatry in multicultural Britain* (pp. viii, 367 p.). London, Gaskell.
- 6 Bhui, K., Bhugra, D., & Goldberg, D. (2000). Cross-cultural validity of the Amritsar Depression Inventory and the General Health Questionnaire amongst English and Punjabi primary care attenders. *Soc Psychiatry Psychiatr Epidemiol*, 35(6), 248-254.
- 9 Bhui, K., Bhugra, D., Goldberg, D., Dunn, G., & Desai, M. (2001). Cultural influences on the prevalence of common mental disorder, general practitioners' assessments and help-seeking among Punjabi and English people visiting their general practitioner. *Psychol Med*, 31(5), 815-825.
- 13 Bhui, K., Stansfeld, S., Hull, S., Priebe, S., Mole, F., & Feder, G. (2003). Ethnic variations in pathways to and use of specialist mental health services in the UK. Systematic review. *Br J Psychiatry*, 182, 105-116.
- 16 Bi, X., Breland, J. Y., Moos, R. H., & Cronkite, R. C. (2015). Longitudinal changes in somatic symptoms and family disagreements among depression and community groups: a 23-year study. *BMC Psychiatry*, 15(1).
- 19 Biddle, L., Donovan, J. L., Gunnell, D., & Sharp, D. (2006). Young adults' perceptions of GPs as a help source for mental distress: a qualitative study. *Br J Gen Pract*, 56(533), 924-931.
- 21 Biddle, S., Fox, K., & Edmunds, L. (1994). Physical activity promotion in primary health care in England: Health Education Authority.
- 23 Bieling, P. J., Hawley, L. L., Bloch, R. T., Corcoran, K. M., Levitan, R. D., Young, L. T., . . . Segal, Z. V. (2012). Treatment-specific changes in decentering following mindfulness-based cognitive therapy versus antidepressant medication or placebo for prevention of depressive relapse. *Journal of Consulting and Clinical Psychology*, 80(3), 365.
- 27 Biesheuvel-Leliefeld, K. E., Kok, G. D., Bockting, C. L., Cuijpers, P., Hollon, S. D., van Marwijk, H. W., & Smit, F. (2015). Effectiveness of psychological interventions in preventing recurrence of depressive disorder: Meta-analysis and meta-regression. *Journal of Affective Disorders*, 174, 400-410.
- 31 Blackburn IM, Bishop S, Glen AI, Whalley LJ, Christie JE. The efficacy of cognitive therapy in depression: a treatment trial using cognitive therapy and pharmacotherapy, each alone and in combination. *The British Journal of Psychiatry*. 1981 Sep 1; 139(3):181-9.
- 34 Blackburn IM, Moore RM. Controlled acute and follow-up trial of cognitive therapy and pharmacotherapy in out-patients with recurrent depression. *British Journal of Psychiatry* 1997; 171:328-334.
- 37 Blackwell SE, Browning M, Mathews A, Pictet A, Welch J, Davies J, Watson P, Geddes JR, Holmes EA. Positive imagery-based cognitive bias modification as a web-based treatment tool for depressed adults: a randomized controlled trial. *Clinical Psychological Science*. 2015 Jan; 3(1):91-111.
- 41 Blanchard, M., Waterreus, A., & Mann, A. (1995). The effect of primary care nurse intervention upon older people screened as depressed. *International Journal of Geriatric Psychiatry*, 10, 289-298.
- 44 Blanco, C., Okuda, M., Markowitz, J. C., Liu, S.-M., Grant, B. F., & Hasin, D. S. (2010). The epidemiology of chronic major depressive disorder and dysthymic disorder and dysthymic

- 1 disorder: results from the national epidemiologic survey on alcohol and related conditions.
- 2 The Journal of Clinical Psychiatry, 71(12), 1645-1656.
- 3 Blom MB, Jonker K, Dusseldorp E, Spinhoven P, Hoencamp E, Haffmans J, van Dyck R.
- 4 Combination treatment for acute depression is superior only when psychotherapy is added to medication. Psychotherapy and Psychosomatics. 2007 Aug 16; 76(5):289-97.
- 5 Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, et al.
- 6 Exercise and pharmacotherapy in the treatment of major depressive disorder. Psychosomatic Medicine 2007; 69(7):587-96.
- 7 Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, Waugh R,
- 8 Napolitano MA, Forman LM, Appelbaum M, Doraiswamy PM. Effects of exercise training on older patients with major depression. Archives of internal medicine. 1999 Oct 25;
- 9 159(19):2349-56.
- 10 13 BMA, & Employers, N. (2006). Revisions to the GMS Contract 2006/07. Delivering investment in general practice. London.
- 11 15 Bockting, C. L., Schene, A. H., Spinhoven, P., Koeter, M. W., Wouters, L. F., Huyser, J., &
- 12 Kamphuis, J. H. (2005). Preventing relapse/recurrence in recurrent depression with cognitive therapy: a randomized controlled trial. Journal of Consulting and Clinical Psychology, 73(4), 647.
- 13 19 Bockting, C. L., Spinhoven, P., Koeter, M. W., Wouters, L. F., Schene, A. H., & Depression Evaluation Longitudinal Therapy Assessment Study, G. (2006). Prediction of recurrence in recurrent depression and the influence of consecutive episodes on vulnerability for depression: a 2-year prospective study. J Clin Psychiatry, 67(5), 747-755.
- 14 23 Bodenmann, G., Plancherel, B., Beach, S. R., Widmer, K., Gabriel, B., Meuwly, N., ...
- 15 24 Schramm, E. (2008). Effects of coping-oriented couples therapy on depression: A randomized clinical trial. Journal of Consulting and Clinical Psychology, 76(6), 944.
- 16 26 Bohlmeijer, E. T., et al. (2011) Efficacy of an early intervention based on acceptance and commitment therapy for adults with depressive symptomatology: Evaluation in a randomized controlled trial. Behaviour research and therapy 49, 62-67.
- 17 29 Bolier, L., et al. (2013) An Internet-based intervention to promote mental fitness for mildly depressed adults: Randomized controlled trial. Journal of Medical Internet Research 15.
- 18 31 Bondolfi, G., Jermann, F., Linden, M. V., Gex-Fabry, M., Bizzini, L., Rouget, B. W., ...
- 19 32 Bertschy, G. (2010). Depression relapse prophylaxis with Mindfulness-Based Cognitive Therapy: replication and extension in the Swiss health care system. Journal of Affective Disorders, 122(3), 224-231.
- 20 35 Booth, A. (2004). Chapter 6: Formulating research questions. In A. Booth & A. Brice (Eds.), Evidence-based practice for information professionals: a handbook (pp. xvi, 304 p.). London: Facet Pub.
- 21 38 Bosanquet, K., Bailey, D., Gilbody, S., Harden, M., Manea, L., Nutbrown, S., & McMillan, D. (2015). Diagnostic accuracy of the Whooley questions for the identification of depression: a diagnostic meta-analysis. BMJ Open, 5(12), e008913.
- 22 41 Bosch P, van den Noort M, Yeo S, Lim S, Coenen A, van Luijtelaar G. The effect of acupuncture on mood and working memory in patients with depression and schizophrenia. Journal of integrative medicine. 2015 Nov 1; 13(6):380-90.
- 23 44 Bose, A., Korotzer, A., Gommoll, C., & Li, D. (2008). Randomized placebo-controlled trial of escitalopram and venlafaxine XR in the treatment of generalized anxiety disorder. Depression and Anxiety, 25(10), 854-861.

- 1 Bose, A., Li, D. & Gandhi, C. (2008) Escitalopram in the acute treatment of depressed patients aged 60 years or older. *American Journal of Geriatric Psychiatry*, 16, 14-20.
- 3 Bose, A., Tsai, J., & Li, D. (2012). Early non-response in patients with severe depression: escitalopram up-titration versus switch to duloxetine. *Clinical Drug Investigation*, 32(6), 373-385.
- 6 Bosmans, J. E., Brook, O. H., Van Hout, H. P. J., De Bruijne, M. C., Nieuwenhuyse, H., Bouter, L. M., . . . Van Tulder, M. W. (2007). Cost effectiveness of a pharmacy-based coaching programme to improve adherence to antidepressants. *PharmacoEconomics*, 25, 25-37
- 10 Bostwick, J. M., & Pankratz, V. S. (2000). Affective disorders and suicide risk: A reexamination. *American Journal of Psychiatry*, 157(12), 1925-1932.
- 12 Bower, P., & Gilbody, S. (2005). Stepped care in psychological therapies: access, effectiveness and efficiency. *Narrative literature review*. *Br J Psychiatry*, 186, 11-17.
- 14 Bower, P., Rowland, N., & Hardy, R. (2003). The clinical effectiveness of counselling in primary care: a systematic review and meta-analysis. *Psychol Med*, 33(2), 203-215.
- 16 Bower, P., & Sibbald, B. (2000). On-site mental health workers in primary care: effects on professional practice. *Cochrane Database Syst Rev* (3), CD000532.
- 18 Boyer, P., & Lecriubier, Y. (1996). Atypical antipsychotic drugs in dysthymia: placebo controlled studies of amisulpride versus imipramine, versus amineptine. *European Psychiatry*, 11, 135s-140s.
- 21 Brakemeier EL, Merkl A, Wilbertz G, Quante A, Regen F, Bührsche N, van Hall F, Kischkel E, Danker-Hopfe H, Anghelescu I, Heuser I. Cognitive-behavioral therapy as continuation treatment to sustain response after electroconvulsive therapy in depression: a randomized controlled trial. *Biological psychiatry*. 2014 Aug 1; 76(3):194-202.
- 25 Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3(2), 77-101.
- 27 Bremner, J. D. (1995). A double-blind comparison of Org 3770, amitriptyline, and placebo in major depression. *Journal of Clinical Psychiatry*, 56, 519-525.
- 29 Brenes GA, Williamson JD, Messier SP, Rejeski WJ, Pahor M, Ip E, et al. Treatment of minor depression in older adults: a pilot study comparing sertraline and exercise. *Aging and Mental Health* 2007; 11(1):61–8.
- 32 Brenner, R., Azbel, V., Madhusoodanan, S., & Pawlowska, M. (2000). Comparison of an extract of hypericum (LI 160) and sertraline in the treatment of depression: a double-blind, randomized pilot study. *Clinical Therapeutics*, 22(4), 411-419.
- 35 Bridges, K. W., & Goldberg, D. P. (1987). Somatic presentations of depressive illness in primary care. In P. Freeling, L. J. Downey, & J. C. Malkin (Eds.), *The presentation of depression: Current approaches*. London: Royal College of General Practitioners.
- 38 Briggs, A., Sculpher, M., & Claxton, K. (2006). *Decision modelling for health economic evaluation*. New York, NY: Oxford University Press.
- 40 British-Association-for-Behavioural-&-Cognitive-Psychotherapies. (2016). Criteria and guidelines for re-accreditation as a behavioural and/or cognitive psychotherapist. .
- 42 Brodaty, H., Luscombe, G., Parker, G., Wilhelm, K., Hickie, I., Austin, M. P., & Mitchell, P. (1997). Increased rate of psychosis and psychomotor change in depression with age. *Psychol Med*, 27(5), 1205-1213.

- 1 Brogden, R. N., Heel, R. C., Speight, T. M., & Avery, G. S. (1981). Trazodone: a review of its pharmacological properties and therapeutic use in depression and anxiety. *Drugs*, 21(6), 401-429.
- 4 Brook, O. H., van Hout, H. P., Stalman, M., Nieuwenhuyse, H., Bakker, B., Heerdink, E., de Haan, M.. (2005). A pharmacy-based coaching program to improve adherence to antidepressant treatment among primary care patients. *Psychiatric Services*, 56 (4), 487-489.
- 7 Brown, G. W., & Harris, T. (1978). Social origins of depression: a study of psychiatric disorder in women. London: Routledge, 2001.
- 9 Brown, J. S., Casey, S. J., Bishop, A. J., Prytys, M., Whittinger, N., & Weinman, J. (2011). How black African and white British women perceive depression and help-seeking: a pilot vignette study. *Int J Soc Psychiatry*, 57(4), 362-374.
- 12 Brown, J. S., Elliott, S. A., Boardman, J., Ferns, J., & Morrison, J. (2004) Meeting the unmet need for depression services with psycho-educational self-confidence workshops: preliminary report. *British Journal of Psychiatry*, 185, 511-515.
- 15 Brown RA, Lewinsohn PM. A psychoeducational approach to the treatment of depression: Comparison of group, individual, and minimal contact procedures. *Journal of Consulting & Clinical Psychology* 1984; 52(5):774-783.
- 18 Brown, T. A., Campbell, L. A., Lehman, C. L., Grisham, J. R., & Mancill, R. B. (2001). Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *Journal of Abnormal Psychology*, 110(4), 585-599.
- 21 Browne, G., Steiner, M., Roberts, J., Gafni, A., Byrne, C., Dunn, E., ... Wallik, D. (2002). Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs. *Journal of Affective Disorders*, 68(2), 317-330.
- 25 Browne, M., Lapierre, Y., Hrdina, P., & Horn, E. (1990). Lithium as an adjunct in the treatment of major depression. *International Clinical Psychopharmacology*, 5(2), 103-110.
- 27 Bruce, M., Ten Have, T., Reynolds III, C., Katz, I., Schulberg, H., Mulsant, B., Brown, G.K., McAvay, G.J., Pearson, J.L., Alexopoulos, G.S. (2004). Reducing suicidal ideation and depressive symptoms in depressed older primary care patients: a randomized controlled trial. *Journal of the American Medical Association*, 291, 1081-1091.
- 31 Bruce, M. L., & Hoff, R. A. (1994). Social and physical health risk factors for first-onset major depressive disorder in a community sample. *Soc Psychiatry Psychiatr Epidemiol*, 29(4), 165-171.
- 34 Brunner E, Tohen M, Osuntokun O, Landry J, Thase ME. Efficacy and safety of olanzapine/fluoxetine combination vs fluoxetine monotherapy following successful combination therapy of treatment-resistant major depressive disorder. *Neuropsychopharmacology*. 2014 Oct; 39(11):2549.
- 38 Brunoni, A. R., Teng, C. T., Correa, C., Imamura, M., Brasil-Neto, J. P., Boechat, R., ... Fregni, F. (2010). Neuromodulation approaches for the treatment of major depression: challenges and recommendations from a working group meeting. *Arquivos de Neuro-Psiquiatria*, 68(3), 433-451.
- 42 Buckley, N. A., & McManus, P. R. (2002). Fatal toxicity of serotonergic and other antidepressant drugs: analysis of United Kingdom mortality data. *BMJ*, 325(7376), 1332-1333.

- 1 Bukh, J. D., Andersen, P. K., & Kessing, L. V. (2016). Rates and predictors of remission, recurrence and conversion to bipolar disorder after the first lifetime episode of depression--a prospective 5-year follow-up study. *Psychol Med*, 46(6), 1151-1161.
- 4 Bull, S. A., Hunkeler, E. M., Lee, J. Y., Rowland, C. R., Williamson, T. E., Schwab, J. R., & Hurt, S. W. (2002). Discontinuing or switching selective serotonin-reuptake inhibitors. *Ann Pharmacother*, 36(4), 578-584.
- 7 Buntrock, C., et al. (2015). "Effectiveness of a Web-Based Cognitive Behavioural Intervention for Subthreshold Depression: Pragmatic Randomised Controlled Trial." *Psychotherapy and psychosomatics* 84(6): 348-358.
- 10 Burke WJ, Gergel I, Bose A. Fixed-dose trial of the single isomer SSRI escitalopram in depressed outpatients. *Journal of Clinical Psychiatry* 2002; 63(4):331-336.
- 12 Burnand Y, Andreoli A, Kolatte E, Venturini A, Rosset N. Psychodynamic psychotherapy and clomipramine in the treatment of major depression. *Psychiatric Services* 2002; 53(5):585-590.
- 15 Burroughs, H. (2006). 'Justifiable depression': how primary care professionals and patients view late-life depression? a qualitative study. *Family Practice*, 23(3), 369-377.
- 17 Burrows, G. D., Norman, T. R., & Judd, F. K. (1994). Definition and differential diagnosis of treatment-resistant depression. *Int Clin Psychopharmacol*, 9 Suppl 2, 5-10.
- 19 Burton, C., Anderson, N., Wilde, K., & Simpson, C. R. (2012). Factors associated with duration of new antidepressant treatment: analysis of a large primary care database. *Br J Gen Pract*, 62(595), e104-112.
- 22 Buszewicz, M., Griffin, M., McMahon, E. M., Beecham, J., & King, M. (2010). Evaluation of a system of structured, pro-active care for chronic depression in primary care: a randomised controlled trial. *BMC Psychiatry*, 10, 61.
- 25 Buszewicz, M., Griffin, M., McMahon, E. M., Walters, K., & King, M. (2016). Practice nurse-led proactive care for chronic depression in primary care: a randomised controlled trial. *Br J Psychiatry*, 208(4), 374-380.
- 28 Button, K. S., Turner, N., Campbell, J., Kessler, D., Kuyken, W., Lewis, G., ... Wiles, N. (2015). Moderators of response to cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care. *Journal of Affective Disorders*, 174, 272-280.
- 32 Byerley WF, Reimherr FW, Wood DR, Grosser BI. Fluoxetine, a selective serotonin uptake inhibitor, for the treatment of outpatients with major depression. *J Clin Psychopharmacol* 1988; 8:112-5.
- 35 Byford, S., Barrett, B., Despiégel, N., & Wade, A. (2011). Impact of treatment success on health service use and cost in depression. *PharmacoEconomics*, 29(2), 157-170.
- 37 Caldwell, D.M. (2014). An overview of conducting systematic reviews with network meta-analysis. *Systematic Reviews*, 3: 109.
- 39 Caldwell, D. M., Ades, A. E., & Higgins, J. P. (2005). Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ*, 331(7521), 897-900.
- 41 Callahan, C. M., Hendrie, H. C., Dittus, R. S., Brater, D. C., Hui, S. L., & Tierney, W. M. (1994). Improving treatment of late life depression in primary care: a randomized clinical trial. *J Am Geriatr Soc*, 42(8), 839-846.
- 44 Callaghan P, Khalil E, Morres I, Carter T. Pragmatic randomised controlled trial of preferred intensity exercise in women living with depression. *BMC Public Health*. 2011 Jun 12; 11(1):1.

- 1 Cameron, D. E. (1947). The day hospital; an experimental form of hospitalization for psychiatric patients. *Modern Hospital*, 69(3), 60-62.
- 3 Cameron, I. M., Crawford, J. R., Lawton, K., & Reid, I. C. (2008). Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *Br J Gen Pract*, 58(546), 32-36.
- 6 Cano, A., & O'Leary, K. D. (2000). Infidelity and separations precipitate major depressive episodes and symptoms of nonspecific depression and anxiety. *Journal of Consulting and Clinical Psychology*, 68(5), 774-781.
- 9 Capoccia, K., Boudreau, D., Blough, D., Ellsworth, A., Clark, D., Stevens, N., Katon, W.J., Sullivan ,S.D. (2004). Randomized trial of pharmacist interventions to improve depression care and outcomes in primary care. *American Journal of Health-System Pharmacy*, 61, 364-372.
- 13 Carlbring P, Hägglund M, Luthström A, Dahlin M, Kadowaki Å, Vernmark K, Andersson G. Internet-based behavioral activation and acceptance-based treatment for depression: a randomized controlled trial. *Journal of Affective Disorders*. 2013 Jun 30; 148(2):331-7.
- 16 Carpenter, L. L., Yasmin, S., & Price, L. H. (2002). A double-blind, placebo-controlled study of antidepressant augmentation with mirtazapine. *Biological Psychiatry*, 51(2), 183-188.
- 18 Carter JD, Luty SE, McKenzie JM, Mulder RT, Frampton CM, Joyce PR. Patient predictors of response to cognitive behaviour therapy and interpersonal psychotherapy in a randomised clinical trial for depression. *Journal of affective disorders*. 2011 Feb 28; 128(3):252-61.
- 21 Carter, J. D., et al. (2013) Psychotherapy for depression: a randomized clinical trial comparing schema therapy and cognitive behavior therapy. *Journal of Affective Disorders* 151, 500-50.
- 24 Cartwright, N., & Munro, E. (2010). The limitations of randomized controlled trials in predicting effectiveness. *Journal of Evaluation in Clinical Practice*, 16(2), 260-266.
- 26 Casañas, R., et al. (2012) Effectiveness of a psycho-educational group program for major depression in primary care: a randomized controlled trial. *BMC psychiatry* 12, 230
- 28 Caspi, A., Sugden, K., Moffitt, T. E., Taylor, A., Craig, I. W., Harrington, H., ... Poulton, R. (2003). Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*, 301(5631), 386-389.
- 31 Cassano, P., & Fava, M. (2002). Depression and public health. *Journal of Psychosomatic Research*, 53(4), 849-857.
- 33 Cassano, G. B., Heinze, G., Loo, H., Mendlewicz, J., & Paes, D. (1996). A double-blind comparison of tianeptine, imipramine and placebo in the treatment of major depressive episodes. *European Psychiatry: the Journal of the Association of European Psychiatrists*, 11, 254-259.
- 37 Cassidy, S., & Henry, J. (1987). Fatal toxicity of antidepressant drugs in overdose. *Br Med J (Clin Res Ed)*, 295(6605), 1021-1024.
- 39 Castonguay, L. G., & Beutler, L. E. (2006). Principles of therapeutic change: a task force on participants, relationships, and techniques factors. *J Clin Psychol*, 62(6), 631-638.
- 41 Chaimani, A., Vasiliadis, H. S., Pandis, N., Schmid, C. H., Welton, N. J., & Salanti, G. (2013). Effects of study precision and risk of bias in networks of interventions: a network meta-epidemiological study. *International journal of epidemiology*, 42(4), 1120-1131.
- 44 Chalder, M., Wiles, N., Campbell, J., Hollinghurst, S., Searle, A., Haase, A., ... Davis, M. (2012). A pragmatic randomised controlled trial to evaluate the cost-effectiveness of a

- 1 physical activity intervention as a treatment for depression: the treating depression with physical activity (TREAD) trial. *HEALTH TECHNOLOGY ASSESSMENT*, 16 (10) 1-+.
- 3 Chan, A. S., et al. (2012) A Chinese Chan-based mind-body intervention for patients with depression. *Journal of Affective Disorders* 142, 283-289.
- 5 Chan, Y.-Y., Lo, W.-Y., Yang, S.-N., Chen, Y.-H., & Lin, J.-G. (2015). The benefit of combined acupuncture and antidepressant medication for depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 176, 106-117.
- 8 Chaput, Y., Magnan, A., & Gendron, A. (2008). The co-administration of quetiapine or placebo to cognitive-behavior therapy in treatment refractory depression: a preliminary trial. *BMC Psychiatry*, 8(1), 73.
- 11 Chaudhry, H. R., Najam, N., & Naqvi, A. (1998). The value of amineptine in depressed patients treated with cognitive behavioural psychotherapy. *Human Psychopharmacology*, 13, 419-424.
- 14 Chen, S., Conwell, Y., He, J., Lu, N., & Wu, J. (2015). Corrections. Depression care management for adults older than 60 years in primary care clinics in urban China: a cluster-randomised trial. *Lancet Psychiatry*, 2(5), 378.
- 17 Chen, S., Conwell, Y., He, J., Lu, N., & Wu, J. (2015). Depression care management for adults older than 60 years in primary care clinics in urban China: A cluster-randomised trial. *The Lancet Psychiatry*, 2(4), 332-339.
- 20 Chen, Y. L., Pan, A. W., Hsiung, P. C., Chung, L., Lai, J. S., Shur-Fen Gau, S., & Chen, T. J. (2015). Life Adaptation Skills Training (LAST) for persons with depression: A randomized controlled study. *Journal of Affective Disorders*, 185(103-2314-B-002-179-MY3 (MOST) *Ministry of Science and Technology*), 108-114.
- 24 Chew-Graham, C., Bashir, C., Chantler, K., Burman, E., & Batsleer, J. (2002). South Asian women, psychological distress and self-harm: lessons for primary care trusts. *Health Soc Care Community*, 10(5), 339-347.
- 27 Chew-Graham, C., Chamberlain, E., Turner, K., Folkes, L., Caulfield, L., & Sharp, D. (2008). GPs' and health visitors' views on the diagnosis and management of postnatal depression: a qualitative study. *Br J Gen Pract*, 58(548), 169-176.
- 30 Chew-Graham, C., Lovell, K., Roberts, C., Baldwin, R., Morley, M., Burns, A., Richards, D., Burroughs, H. (2007). A randomised controlled trial to test the feasibility of a collaborative care model for the management of depression in older people. *The British Journal of General Practice*, 57, 364-370.
- 34 Chew-Graham, C. A., May, C. R., Cole, H., & Hedley, S. (2000). The burden of depression in primary care: a qualitative investigation of general practitioners' constructs of depressed people in the inner city. *International Journal of Psychiatry in Clinical Practice*, 6(4), 137-141.
- 37 Chaudhry, H. R., Najam, N., & Naqvi, A. (1998). The value of amineptine in depressed patients treated with cognitive behavioural psychotherapy. *Human Psychopharmacology*, 13, 419-424.
- 40 Chiesa, A., Castagner, V., Andrisano, C., Serretti, A., Mandelli, L., Porcelli, S., & Giommi, F. (2015). Mindfulness-based cognitive therapy vs. psycho-education for patients with major depression who did not achieve remission following antidepressant treatment. *Psychiatry research*, 226(2-3), 474-483.
- 44 Chisholm D, Sweeny K, Sheehan P, Rasmussen B, Smit F, Cuijpers P, Saxena S. Scaling-up treatment of depression and anxiety: a global return on investment analysis. *Lancet Psychiatry*. 2016 May; 3(5):415-24.

- 1 Choi, H., & Marks, N. F. (2008). Marital conflict, depressive symptoms, and functional impairment. *Journal of Marriage and Family*, 70(2), 377-390.
- 3 Choi, I., Zou, J., Titov, N., Dear, B. F., Li, S., Johnston, L., ... Hunt, C. (2012). Culturally attuned Internet treatment for depression amongst Chinese Australians: a randomised controlled trial. *Journal of Affective Disorders*, 136(3), 459-468.
- 6 Chong, J., & Moreno, F. (2012). Feasibility and acceptability of clinic-based telepsychiatry for low-income Hispanic primary care patients. *Telemedicine Journal and e-Health*, 18(4), 297-304.
- 9 Christian, J. L., O'Leary, K. D., & Vivian, D. (1994). Depressive symptomatology in maritally discordant women and men: The role of individual and relationship variables. *Journal of Family Psychology*, 8, 32-42.
- 12 Christensen H, Griffiths KM, Jorm AF. Delivering interventions for depression by using the internet: randomised controlled trial. *Bmj*. 2004 Jan 29; 328(7434):265.
- 14 Chu IH, Buckworth J, Kirby TE, Emery CF. Effect of exercise intensity on depressive symptoms in women. *Mental Health and Physical Activity* 2009; 2(1):37–43.
- 16 Ciechanowski, P., Wagner, E., Schmaling, K., Schwartz, S., Williams, B., Diehr, P., Kulzer, J., Gray, S., Collier, C., LoGerfo, J. (2004). Community-integrated home-based depression treatment in older adults: a randomized controlled trial. *Journal of the American Medical Association*, 291, 1569-1577.
- 20 Cipriani, A., Furukawa, T. A., Salanti, G., Geddes, J. R., Higgins, J. P., Churchill, R., ... Barbui, C. (2009). Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *Lancet*, 373(9665), 746-758.
- 23 Claghorn, J., Gershon, S., Goldstein, B. J., Behrnetz, S., Bush, D. F., & Huitfeldt, B. (1983). A double-blind evaluation of zimelidine in comparison to placebo and amitriptyline in patients with major depressive disorder. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 7, 367-382.
- 27 Claghorn JL, Johnstone EE, Studebaker SL, Ajeman SA. The effectiveness of 6-azamianserin (Org 3770) in depressed outpatients. *Psychopharmacology Bulletin* 1987; 23(1):160-161.
- 30 Claghorn, J.L.; Lesem, M.D. A double-blind placebo-controlled study of Org 3770 in depressed outpatients. *Journal of Affective Disorders* 1995; 34(3):165-171.
- 32 Clarke G, Eubanks D, Reid E, Kelleher C, O'connor E, DeBar LL, Lynch F, Nunley S, Gullion C. Overcoming Depression on the Internet (ODIN)(2): a randomized trial of a self-help depression skills program with reminders. *Journal of medical Internet research*. 2005; 7(2):e16.
- 36 Clarke, G., Reid, E., Eubanks, D., O'Connor, E., DeBar, L. L., Kelleher, C. et al. (2002). Overcoming Depression on the Internet (ODIN): A randomized controlled trial of an internet depression skills intervention program. *Journal of Medical Internet Research*, 4, e14.
- 39 Clarke, K., Mayo-Wilson, E., Kenny, J., & Pilling, S. (2015). Can non-pharmacological interventions prevent relapse in adults who have recovered from depression? A systematic review and meta-analysis of randomised controlled trials. *Clinical Psychology Review*, 39, 58-70.
- 43 Cohn, J. B., Varga, L., & Lyford, A. (1984). A two-center double-blind study of nomifensine, imipramine, and placebo in depressed geriatric outpatients. *Journal of Clinical Psychiatry*, 45, 68-72.

- 1 Cole, M., McCusker, J., Elie, M., Dendukuri, N., Latimer, E., & Belzile, E. (2006). Systematic detection and multidisciplinary care of depression in older medical inpatients: a randomized trial. *Canadian Medical Association Journal*, 174, 38-44.
- 4 Coleman CC, King BR, Bolden-Watson C, Book MJ, Segraves RT, Richard N, Ascher J, Batey S, Jamerson B, Metz A. A placebo-controlled comparison of the effects on sexual functioning of bupropion sustained release and fluoxetine. *Clinical Therapeutics* 2001; 23(7):1040-1058.
- 8 Collaboration, C. (2014). Review Manager (RevMan) [Computer program]. Version 5.3. .
- 9 Colonna, L., Andersen, H.F. & Reines, E.H. (2005) A randomised, double-blind, 24 -week study of escitalopram (10 mg/day), versus citalopram (20 mg/day) in primary care patients with major depressive disorder. *Current Medical Research and Opinion*, 21, 1659-1668.
- 12 Commander, M. J., Dharan, S. P., Odell, S. M., & Surtees, P. G. (1997). Access to mental health care in an inner-city health district. I: Pathways into and within specialist psychiatric services. *Br J Psychiatry*, 170, 312-316.
- 15 Committee-on-Safety-of-Medicines. (2000). Reminder: St John's wort (*hypericum perforatum*) interactions. *Current Problems in Pharmacovigilance*, 26, 6-7.
- 17 Compton JS. Adapting marital therapy in older adults with depression [NCT00612807].
18 Available from: <https://clinicaltrials.gov/ct2/show/study/NCT00612807> [accessed 22.09.2017]
- 19 Constantino, M.J., Marnell, M.E., Haile, A.J., et al. (2008) Integrative cognitive therapy for depression: A randomized pilot comparison. *Psychotherapy: Theory, Research, Practice, Training*, 45 (2), 122-134.
- 22 Cook IA, Leuchter AF, Witte E, Abrams M, Uijtdehaage SH, Stubbeman W, Rosenberg-Thompson S, Anderson-Hanley C, Dunkin JJ. Neurophysiologic predictors of treatment response to fluoxetine in major depression. [erratum appears in Psychiatry Res 2000 Jul 24;95(1):87.]. *Psychiatry Research* 22-3-1999;85(3):263-273.
- 26 Coope, C., Gunnell, D., Hollingworth, W., Hawton, K., Kapur, N., Fearn, V., ... Metcalfe, C. (2014). Suicide and the 2008 economic recession: Who is most at risk? Trends in suicide rates in England and Wales 2001–2011. *Social Science & Medicine*, 117, 76-85.
- 29 Cooper, L. A., Ghods Dinoso, B. K., Ford, D. E., Roter, D. L., Primm, A. B., Larson, S. M., ... Wang, N. Y. (2013). Comparative effectiveness of standard versus patient-centered collaborative care interventions for depression among African Americans in primary care settings: the BRIDGE Study. *Health Services Research*, 48(1), 150-174.
- 33 Cooper-Kazaz, R., Apter, J. T., Cohen, R., Karagichev, L., Muhammed-Moussa, S., Grupper, D., ... Glaser, B. (2007). Combined treatment with sertraline and liothyronine in major depression: a randomized, double-blind, placebo-controlled trial. *Archives of General Psychiatry*, 64(6), 679-688.
- 37 Cooper-Patrick, L., Powe, N. R., Jenckes, M. W., Gonzales, J. J., Levine, D. M., & Ford, D. E. (1997). Identification of patient attitudes and preferences regarding treatment of depression. *J Gen Intern Med*, 12(7), 431-438.
- 40 Coplan, J. D., Gopinath, S., Abdallah, C. G., & Berry, B. R. (2014). A neurobiological hypothesis of treatment-resistant depression - mechanisms for selective serotonin reuptake inhibitor non-efficacy. *Front Behav Neurosci*, 8, 189.
- 43 Coppen, A., Ghose, K., Montgomery, S., Rao, V. R., Bailey, J., & Jorgensen, A. (1978). Continuation therapy with amitriptyline in depression. *The British Journal of Psychiatry*, 133(1), 28-33.

- 1 Cornelius JR, Bukstein OG, Douaihy AB, Clark DB, Chung TA, Daley DC, Wood DS, Brown SJ. Double-blind fluoxetine trial in comorbid MDD–CUD youth and young adults. *Drug and alcohol dependence*. 2010 Nov 1; 112(1):39-45.
- 4 Corrigan MH, Denahan AQ, Wright CE, Ragual RJ, Evans DL. Comparison of pramipexole, fluoxetine, and placebo in patients with major depression. *Depression & Anxiety* 2000; 11(2):58-65.
- 7 Corya, S. A., Williamson, D., Sanger, T. M., Briggs, S. D., Case, M., & Tollefson, G. (2006). A randomized, double-blind comparison of olanzapine/fluoxetine combination, olanzapine, fluoxetine, and venlafaxine in treatment-resistant depression. *Depression and Anxiety*, 23(6), 364-372.
- 11 Coryell, W., Akiskal, H. S., Leon, A. C., Winokur, G., Maser, J. D., Mueller, T. I., & Keller, M. B. (1994). The time course of nonchronic major depressive disorder. Uniformity across episodes and samples. National Institute of Mental Health Collaborative Program on the Psychobiology of Depression--Clinical Studies. *Arch Gen Psychiatry*, 51(5), 405-410.
- 15 Coryell, W., Endicott, J., & Keller, M. B. (1991). Predictors of relapse into major depressive disorder in a nonclinical population. *Am J Psychiatry*, 148(10), 1353-1358.
- 17 Coryell, W., Leon, A., Winokur, G., Endicott, J., Keller, M., Akiskal, H., & Solomon, D. (1996). Importance of psychotic features to long-term course in major depressive disorder. *Am J Psychiatry*, 153(4), 483-489.
- 20 Coupland, C., Dhiman, P., Morriss, R., Arthur, A., Barton, G., & Hippisley-Cox, J. (2011). Antidepressant use and risk of adverse outcomes in older people: population based cohort study. *BMJ*, 343, d4551.
- 23 Coventry, P., Hudson, J., Kontopantelis, E., Archer, J., Richards, D., Gilbody, S., ... Bower, P. (2014). Characteristics of Effective Collaborative Care for Treatment of Depression: A Systematic Review and Meta-Regression of 74 Randomised Controlled Trials. *PLoS ONE*, 9, e108114.
- 27 Covi L, Lipman RS. Cognitive behavioral group psychotherapy combined with imipramine in major depression. *Psychopharmacol Bull* 1987; 23(1):173-176.
- 29 Cowen, P. J. (2012). Phototherapy. In J. Lopez-Ibor & J. R. Geddes (Eds.), *New Oxford Textbook of Psychiatry* (2nd ed., pp. 1261-1263): Oxford University Press (OUP).
- 31 Coyne, J. C., Pepper, C. M., & Flynn, H. (1999). Significance of prior episodes of depression in two patient populations. *J Consult Clin Psychol*, 67(1), 76-81.
- 33 Crabb, R. and Hunsley, J. (2006). Utilization of mental health care services among older adults with depression. *J. Clin. Psychology*, 62(3), 299–312.
- 35 Craig, T. K. J., & Bhugra, D. (2012). Mental health services for ethnic minorities *New Oxford Textbook of Psychiatry* (pp. 1503-1504): Oxford University Press (OUP).
- 37 Cramer, H., Anheyer, D., Lauche, R., & Dobos, G. (2017). A systematic review of yoga for major depressive disorder. *J Affect Disord*, 213, 70-77.
- 39 Cramer, H., Salisbury, C., Conrad, J., Eldred, J., & Araya, R. (2011). Group cognitive behavioural therapy for women with depression: pilot and feasibility study for a randomised controlled trial using mixed methods. *BMC Psychiatry*, 11, 82.
- 42 Crawford, M. J., Prince, M., Menezes, P., & Mann, A. H. (1998). The recognition and treatment of depression in older people in primary care. *Int J Geriatr Psychiatry*, 13(3), 172-176.

- 1 Creed, F., Black, D., Anthony, P., Osborn, M., Thomas, P., & Tomenson, B. (1990). Randomised controlled trial of day patient versus inpatient psychiatric treatment. *BMJ*, 300(6731), 1033-1037.
- 4 Creed, F., & Marks, B. (1989). Liaison psychiatry in general practice: a comparison of the liaison-attachment scheme and shifted outpatient clinic models. *J R Coll Gen Pract*, 39(329), 514-517.
- 7 Creed, F., Mbaya, P., Lancashire, S., Tomenson, B., Williams, B., & Holme, S. (1997). Cost effectiveness of day and inpatient psychiatric treatment: results of a randomised controlled trial. *BMJ*, 314(7091), 1381.
- 10 Croft H. A placebo-controlled comparison of the antidepressant efficacy and effects on sexual functioning of sustained-release bupropion and sertraline. *Clinical Therapeutics* 21(4):643-58, 1999.
- 13 Crowther, R. E., Marshall, M., Bond, G. R., & Huxley, P. (2001). Helping people with severe mental illness to obtain work: systematic review. *BMJ*, 322(7280), 204-208.
- 15 Cuijpers, P., & Gentili, C. (2017). Psychological treatments are as effective as pharmacotherapies in the treatment of adult depression: a summary from Randomized Clinical Trials and neuroscience evidence. *Research in Psychotherapy: Psychopathology, Process and Outcome*, 20(2).
- 19 Cuijpers, P., Sijbrandij, M., Koole, S. L., Andersson, G., Beekman, A. T., & Reynolds, C. F., 3rd. (2014). Adding psychotherapy to antidepressant medication in depression and anxiety disorders: a meta-analysis. *World Psychiatry*, 13(1), 56-67.
- 22 Cuijpers, P., Smit, F., & van Straten, A. (2007). Psychological treatments of subthreshold depression: a meta-analytic review. *Acta Psychiatrica Scandinavica*, 115(6), 434-441.
- 24 Curtis, L. (2014). *Unit Costs of Health & Social Care 2014*. Canterbury.
- 25 Curtis, L., & Burns, A. (2016). *Unit Costs of Health & Social Care 2016*. Canterbury.
- 26 Dalgard, O. S., Dowrick, C., Lehtinen, V., Vazquez-Barquero, J. L., Casey, P., Wilkinson, G., ODIN Group. (2006). Negative life events, social support and gender difference in depression. *Social psychiatry and psychiatric epidemiology*, 41(6), 444-451.
- 29 Danielsson L, Papoulias I, Petersson EL, Carlsson J, Waern M. Exercise or basic body awareness therapy as add-on treatment for major depression: a controlled study. *Journal of Affective Disorders*. 2014 Oct 15; 168:98-106.
- 32 Darling, C., & Tyrer, P. (1990). Brief encounters in general practice: liaison in general practice psychiatry clinics. *Psychiatric Bulletin*, 14(10), 592.
- 34 Datto, C., Thompson, R., Horowitz, D., Disbot, M., & Oslin, D. (2003). The pilot study of a telephone disease management program for depression. *General Hospital Psychiatry*, 25, 169-177.
- 37 David, D., Szentagotai, A., Lupu, V., & Cosman, D. (2008) Rational emotive behaviour therapy, cognitive therapy, and medication in the treatment of major depressive disorder: A randomised clinical trial, posttreatment outcomes, and six-month follow-up. *Journal of Clinical Psychology*, 64, 728-746.
- 41 Davidson, J., Gadde, K., Fairbank, J., Rama Krishnan, K. R., Calif, R., Binanay, C., ... Weisler, R. (2002). Effect of Hypericum perforatum (St John's wort) in major depressive disorder: a randomized controlled trial. *JAMA*, 287(14), 1807-1814.

- 1 Davila, J., Karney, B. R., Hall, T. W., & Bradbury, T. N. (2003). Depressive symptoms and marital satisfaction: Within-subject associations and the moderating effects of gender and neuroticism. *Journal of Family Psychology*, 17(4), 557-570.
- 4 Davison, G. C. (2000). Stepped care: doing more with less? *J Consult Clin Psychol*, 68(4), 580-585.
- 6 de Graaf, L. E., et al. (2009) Clinical effectiveness of online computerised cognitive-behavioural therapy without support for depression in primary care: randomised trial. *British journal of psychiatry* 195, 73-80.
- 9 de Jonghe FE, Kool S, Van Aalst G, Dekker J, Peen J. Combining psychotherapy and antidepressants in the treatment of depression. *Journal of Affective Disorders*. 2001 May 31; 64(2):217-29.
- 12 de Jonghe F, HendrickSEN M, van Aalst G, Kool S, Peen V, Van R, van den Eijnden E, Dekker J. Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. *The British Journal of Psychiatry*. 2004 Jul 1; 185(1):37-45.
- 15 de la Cerda P, Cervelló E, Cocca A, Viciana J. Effect of an aerobic training program as complementary therapy in patients with moderate depression. *Perceptual and motor skills*. 2011 Jun 1; 112(3):761-9.
- 18 de Mello, M. F., Myczcowisk, L. M., & Menezes, P. R. (2001). A randomized controlled trial comparing moclobemide and moclobemide plus interpersonal psychotherapy in the treatment of dysthymic disorder. *The Journal of psychotherapy practice and research*, 10(2), 117.
- 22 De Picker, L., Van Den Eede, F., Dumont, G., Moorkens, G., & Sabbe, B. G. (2014). Antidepressants and the risk of hyponatremia: a class-by-class review of literature. *Psychosomatics*, 55(6), 536-547.
- 25 De Ronchi D, Rucci P, Lodi M, Ravaglia G, Forti P, Volterra V. Fluoxetine and amitriptyline in elderly depressed patients. A 10-week, double-blind study on course of neurocognitive adverse events and depressive symptoms. *Archives Gerontology Geriatrics* 1998; 6 suppl: 125-140.
- 29 Dekker, J.J.M., Koelen, J.A., Van, H.L., et al. (2008) Speed of action: The relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. *Journal of Affective Disorders*, 109, 183-188.
- 32 Dekker J, Molenaar PJ, Kool S, Van Aalst G, Peen J, De Jonghe F. Dose–effect relations in time-limited combined psycho-pharmacological treatment for depression. *Psychological medicine*. 2005; 35(01):47-58.
- 35 del Carmen Macías-Cortés E, Llanes-González L, Aguilar-Faisal L, Asbun-Bojalil J. Individualized homeopathic treatment and fluoxetine for moderate to severe depression in peri-and postmenopausal women (HOMDEP-MENOP study): a randomized, double-dummy, double-blind, placebo-controlled trial. *PloS one*. 2015 Mar 13; 10(3):e0118440.
- 39 Del Piccolo, L., Saltini, A., & Zimmermann, C. (1998). Which patients talk about stressful life events and social problems to the general practitioner? *Psychological Medicine*, 28(6), 1289-1299.
- 42 Delgadillo J, Gore S, Ali S, Ekers D, Gilbody S, Gilchrist G, McMillan D, Hughes E. Feasibility randomized controlled trial of cognitive and behavioral interventions for depression symptoms in patients accessing drug and alcohol treatment. *Journal of Substance Abuse Treatment*. 2015 Aug 31; 55:6-14.

- 1 Demyttenaere K, Van Ganse E, Gregoire J, Gaens E, Mesters P. Compliance in depressed patients treated with fluoxetine or amitriptyline. International Clinical Psychopharmacology 1998; 13:11-17.
- 4 Dennehy, E. B., Robinson, R. L., Stephenson, J. J., Faries, D., Grabner, M., Palli, S. R., ...
5 Marangell, L. B. (2015). Impact of non-remission of depression on costs and resource
6 utilization: from the COnorbidities and symptoms of DEpression (CODE) study. Current
7 Medical Research and Opinion, 31(6), 1165-1177.
- 8 Denton, W. H., Carmody, T. J., Rush, A. J., Thase, M. E., Trivedi, M. H., Arnow, B. A., ...
9 Keller, M. B. (2009). Dyadic discord at baseline is associated with lack of remission in the
10 acute treatment of chronic depression. Psychological Medicine, 40(03), 415.
- 11 Department of Health. (1999). National Service Framework for mental health: modern
12 standards and service models; executive summary: Great Britain, Department of Health.
- 13 Department of Health. (2001). Exercise referral systems: a national quality assurance
14 framework: Great Britain, Department of Health.
- 15 Department of Health. (2003). Fast-forwarding primary care mental health: graduate primary
16 care mental healthworkers - best practice guidance. London.
- 17 Department of Health. (2005). Delivering race equality in mental health care: an action plan
18 for reform inside and outside services: and the Government's response to the independent
19 inquiry into the death of David Bennett. London: Dept. of Health.
- 20 Department of Health. (2007). Improving access to psychological therapies: specification for
21 the commissioner-led pathfinder programme. .
- 22 Department of Health. (2008). Report on self reported experience of patients from black and
23 minority ethnic groups. .
- 24 Department of Health. (2011). Talking therapies: a four-year plan of action. .
- 25 Department of Health, & Social Care Institute for Excellence. (2009). Think child, think
26 parent, think family: A guide to parental mental health and child welfare. .
- 27 Department of Health, & Social Care Institute for Excellence & Care Services Improvement
28 Partnership. (2008). Care programme approach (CPA) briefing: Parents with mental health
29 problems and their children
- 30 DeRubeis, R. J., Cohen, Z. D., Forand, N. R., Fournier, J. C., Gelfand, L. A., & Lorenzo-
31 Luaces, L. (2014). The Personalized Advantage Index: translating research on prediction into
32 individualized treatment recommendations. A demonstration. PLoS ONE, 9(1), e83875.
- 33 Desan, P. H., Weinstein, A. J., Michalak, E. E., Tam, E. M., Meesters, Y., Ruiter, M. J., ...
34 Boivin, D. B. (2007). A controlled trial of the Litebook light-emitting diode (LED) light therapy
35 device for treatment of Seasonal Affective Disorder (SAD). BMC Psychiatry, 7(1), 38.
- 36 Devanand DP, Nobler MS, Cheng J, Turret N, Pelton GH, Roose SP, Sackeim HA.
37 Randomized, double-blind, placebo-controlled trial of fluoxetine treatment for elderly patients
38 with dysthymic disorder. The American journal of geriatric psychiatry. 2005 Jan 31;13(1):59-
39 68.
- 40 Dias, S., Welton, N. J., Sutton, A. J., & Ades, A. E. (2011a [last updated 2013]). NICE DSU
41 Technical support document 2: a generalised linear modelling framework for pairwise and
42 network meta-analysis of randomised controlled trials. Retrieved from
43 <http://www.nicedsu.org.uk>

- 1 Dias, S., Welton, N. J., Sutton, A. J., Caldwell, D., Lu, G., & Ades, A. E. (2011b). NICE DSU
- 2 Technical support document 4: Inconsistency in networks of evidence based on randomised
- 3 controlled trials.
- 4 Dick, P., Cameron, L., Cohen, D., Barlow, M., & Ince, A. (1985). Day and full time psychiatric
- 5 treatment: a controlled comparison. *The British Journal of Psychiatry*, 147(3), 246-249.
- 6 Dick, P. H., Sweeney, M. L., & Crombie, I. K. (1991). Controlled comparison of day-patient
- 7 and out-patient treatment for persistent anxiety and depression. *The British Journal of*
- 8 *Psychiatry*, 158(1), 24-27.
- 9 Dietrich, A., Oxman, T., Williams, J., Schulberg, H., Bruce, M., Lee, P., Barry, S., Raue, .P.J.
- 10 Lefever, J.J., Heo, M., Rost, K., Kroenke, K., Gerrity, M., Nutting, P.A. (2004). Re-
- 11 engineering systems for the treatment of depression in primary care: cluster randomised
- 12 controlled trial. *British Medical Journal*, 329, 602.
- 13 Dimidjian, S., Hollon, S.D., Dobson, K.S., Schmaling, K.B., et al. (2006) Randomized trial of
- 14 behavioural activation, cognitive therapy, and antidepressant medication in the acute
- 15 treatment of adults with major depression. *Journal of Consulting and Clinical Psychology*, 74
- 16 (4), 658-670.
- 17 Dimidjian, S., Martell, C. R., Addis, M. E., Herman-Dunn, R., & Barlow, D. (2008). Behavioral
- 18 activation for depression. *Clinical handbook of psychological disorders: A step-by-step*
- 19 treatment manual, 4, 328-364.
- 20 Dinger, U., Klipsch, O., Köhling, J., Ehrenthal, J. C., Nikendei, C., Herzog, W., &
- 21 Schauenburg, H. (2014). Day-clinic and inpatient psychotherapy for depression (DIP-D): a
- 22 randomized controlled pilot study in routine clinical care. *Psychotherapy and*
- 23 *psychosomatics*, 83(3), 194-195.
- 24 Dixon-Woods, M., Kirk, D., Agarwal, S., et al. (2005). Vulnerable groups and access to health
- 25 care: A critical interpretive review report for the National Coordinating Centre for NHS
- 26 Service Delivery and Organisation R&D (NCCSDO). London: NCCSDO.
- 27 Dobscha, S. K., Corson, K., Hickam, D. H., Perrin, N. A., Kraemer, D. F., & Gerrity, M. S.
- 28 (2006). Depression decision support in primary care: a cluster randomized trial. *Ann Intern*
- 29 *Med*, 145(7), 477-487.
- 30 Dobson, K. S., Hollon, S. D., Dimidjian, S., Schmaling, K. B., Kohlenberg, R. J., Gallop, R. J.,
- 31 ... Jacobson, N. S. (2008). Randomized trial of behavioral activation, cognitive therapy, and
- 32 antidepressant medication in the prevention of relapse and recurrence in major depression.
- 33 *Journal of Consulting and Clinical Psychology*, 76(3), 468.
- 34 Donoghue, J., Tylee, A., & Wildgust, H. (1996). Cross sectional database analysis of
- 35 antidepressant prescribing in general practice in the United Kingdom, 1993-5. *BMJ*,
- 36 313(7061), 861-862.
- 37 Doogan, D., & Caillard, V. (1992). Sertraline in the prevention of depression. *The British*
- 38 *Journal of Psychiatry*, 160(2), 217-222.
- 39 Doogan DP, Langdon CJ. A double-blind, placebo-controlled comparison of sertraline and
- 40 dothiepin in the treatment of major depression in general practice. *International Clinical*
- 41 *Psychopharmacology* 1994 summer; 9(2):95-100.
- 42 Doongaji DR, Bal S, Rajkumar S et al. Multicentre double-blind comparison of lofepramine
- 43 and amitriptyline in the treatment of major depressive disorders in Indian patients. *British*
- 44 *Journal Clinical Research* 1993; 4:45-53.

- 1 Doose M, Ziegenbein M, Hoos O, Reim D, Stengert W, Hoffer N, Vogel C, Ziert Y, Sieberer M. Self-selected intensity exercise in the treatment of major depression: A pragmatic RCT. International journal of psychiatry in clinical practice. 2015 Oct 2; 19(4):266-75.
- 4 Dorée, J.-P., Rosiers, J. D., Lew, V., Gendron, A., Elie, R., Stip, E., & Valérie Tourjman, S. (2007). Quetiapine augmentation of treatment-resistant depression: a comparison with lithium. Current Medical Research and Opinion, 23(2), 333-341.
- 7 Dornseif BE, Dunlop SR, Potvin JH, Wernicke JF. Effect of dose escalation after low-dose fluoxetine therapy. Psychopharmacology bulletin. 1989; 25(1):71.
- 9 Dorstyn, D. S., Saniotis, A., & Sobhanian, F. (2013). A systematic review of telecounselling and its effectiveness in managing depression amongst minority ethnic communities. Journal of Telemedicine and Telecare, 19(6), 338-346.
- 12 Dougherty, D. D., Rezai, A. R., Carpenter, L. L., Howland, R. H., Bhati, M. T., O'Reardon, J. P., . . . Malone, D. A., Jr. (2015). A randomized sham-controlled trial of deep brain stimulation of the ventral capsule/ventral striatum for chronic treatment-resistant depression. Biol Psychiatry, 78(4), 240-248.
- 16 Dowrick, C. (2004). Beyond depression: a new approach to understanding and management. Oxford; New York: Oxford University Press.
- 18 Dozois, D. J., et al. (2009) Changes in self-schema structure in cognitive therapy for major depressive disorder: a randomized clinical trial. Journal of Consulting and Clinical Psychology 77, 1078-1088.
- 21 Drevets, W. C., Price, J. L., & Furey, M. L. (2008). Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. Brain Structure and Function, 213(1-2), 93-118.
- 24 Driessens, E., Cuijpers, P., de Maat, S. C., Abbass, A. A., de Jonghe, F., & Dekker, J. J. (2010). The efficacy of short-term psychodynamic psychotherapy for depression: a meta-analysis. Clin Psychol Rev, 30(1), 25-36.
- 27 Driessens, E., Hollon, S. D., Bockting, C. L., Cuijpers, P., & Turner, E. H. (2015). Does publication bias inflate the apparent efficacy of psychological treatment for major depressive disorder? A systematic review and meta-analysis of US National Institutes of Health-funded trials. PLoS ONE, 10(9), e0137864.
- 31 Duan, D. M., Tu, Y., Chen, L. P., & Wu, Z. J. (2009). Efficacy evaluation for depression with somatic symptoms treated by electroacupuncture combined with Fluoxetine. Journal of Traditional Chinese Medicine / Chung i tsa chih ying wen pan, 29(3), 167-173
- 34 Duarte A, Mikkelsen H, Delini-Stula A. Moclobemide versus fluoxetine for double depression: a randomized double-blind study. Journal of psychiatric research. 1996 Dec 31; 30(6):453-8.
- 36 Dubini A. Do noradrenaline and serotonin differentially affect social motivation and behaviour? European Neuropsychopharmacology Vol 7 Suppl 1, pp S49-55; discussion S71-3, 1997.
- 39 Dunlop SR, Dornseif BE, Wernicke JF, Potvin JH. Pattern analysis shows beneficial effect of fluoxetine treatment in mild depression. Psychopharmacology Bulletin 1990; 26(2):173-180.
- 41 Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. The DOSE study: a clinical trial to examine efficacy and dose response of exercise as a treatment for depression. Controlled Clinical Trials 2002; 23:584-603.

- 1 Dunner, D. (1996). Efficacy of selective serotonin re-uptake inhibitors in the treatment of dysthymia. Selective Serotonin Re-uptake Inhibitors: Advances in Basic Research and Clinical Practice. Chichester: John Wiley & Sons, 223-232.
- 4 Dunner, D. L., Amsterdam, J. D., Shelton, R. C., Loebel, A., & Romano, S. J. (2007). Efficacy and tolerability of adjunctive ziprasidone in treatment-resistant depression: a randomized, open-label, pilot study. *The Journal of Clinical Psychiatry*, 68(7), 1071-1077.
- 7 Dwamena, B. (2007). MIDAS: Stata module for meta-analytical integration of diagnostic test accuracy studies. Boston, Massachusetts: Boston College Department of Economics.
- 9 Dwight-Johnson, M., Aisenberg, E., Golinelli, D., Hong, S., O'Brien, M., & Ludman, E. (2011). Telephone-based cognitive-behavioral therapy for Latino patients living in rural areas: a randomized pilot study. *Psychiatric Services*, 62(8), 936-942.
- 12 Dwight-Johnson M, Lagomasino I, Hay J, Zhang L, Tang L, Green J, et al. Effectiveness of collaborative care in addressing depression treatment preferences among low-income Latinos. *Psychiatric Services* 2010; 61(11):1112-8.
- 15 Eagles, J. M., Wileman, S. M., Cameron, I. M., Howie, F. L., Lawton, K., Gray, D. A., ... Naji, S. A. (1999). Seasonal affective disorder among primary care attenders and a community sample in Aberdeen. *The British Journal of Psychiatry*, 175(5), 472-475.
- 18 Eastman, C. I., Young, M. A., Fogg, L. F., Liu, L., & Meaden, P. M. (1998). Bright light treatment of winter depression: a placebo-controlled trial. *Archives of General Psychiatry*, 55(10), 883-889.
- 21 Eaton, W. W., Kalaydjian, A., Scharfstein, D. O., Mezuk, B., & Ding, Y. (2007). Prevalence and incidence of depressive disorder: the Baltimore ECA follow-up, 1981-2004. *Acta Psychiatrica Scandinavica*, 116(3), 182-188.
- 24 Eaton, W. W., Shao, H., Nestadt, G., Lee, H. B., Bienvenu, O. J., & Zandi, P. (2008). Population-based study of first onset and chronicity in major depressive disorder. *Arch Gen Psychiatry*, 65(5), 513-520.
- 27 Eccles, M., Freemantle, N., & Mason, J. (1998). North of England evidence based guidelines development project: methods of developing guidelines for efficient drug use in primary care. *BMJ*, 316(7139), 1232-1235.
- 30 Edwards, S., Tinning, L., Brown, J. S., Boardman, J., & Weinman, J. (2007). Reluctance to seek help and the perception of anxiety and depression in the United Kingdom: a pilot vignette study. *J Nerv Ment Dis*, 195(3), 258-261.
- 33 Edwards, S. J. (2013). Lithium or an atypical antipsychotic drug in the management of treatment-resistant depression: A systematic review and economic evaluation. *Health Technology Assessment*, 17(54), 1-34.
- 36 Egan, G. (1990). The skilled helper: A systematic approach to effective helping. Pacific Grove, California: Brooks/Cole.
- 38 Eisendrath, S. J., Gillung, E., Delucchi, K. L., Segal, Z. V., Nelson, J. C., McInnes, L. A., ... Feldman, M. D. (2016). A Randomized Controlled Trial of Mindfulness-Based Cognitive Therapy for Treatment-Resistant Depression. *Psychotherapy and psychosomatics*, 85(2), 99-110.
- 42 Ekers, D., Richards, D., McMillan, D., Bland, J. M., & Gilbody, S. (2011). Behavioural activation delivered by the non-specialist: phase II randomised controlled trial. *British Journal of Psychiatry*, 198(1), 66-72.

- 1 Ekkers, W., Korrelboom, K., Huijbrechts, I., Smits, N., Cuijpers, P., & Gaag, M. (2011).
- 2 Competitive Memory Training for treating depression and rumination in depressed older adults: a randomized controlled trial. *Behaviour research and therapy*, 49(10), 588-596.
- 3 Elgie, R. (2006). A patient and primary care perspective: a patient's perspective on the treatment of depression. *J Clin Psychiatry*, 67 Suppl 6, 38-40.
- 4 El-Khalili, N., Joyce, M., Atkinson, S., Buynak, R. J., Datto, C., Lindgren, P., & Eriksson, H. (2010). Extended-release quetiapine fumarate (quetiapine XR) as adjunctive therapy in major depressive disorder (MDD) in patients with an inadequate response to ongoing antidepressant treatment: a multicentre, randomized, double-blind, placebo-controlled study. *International Journal of Neuropsychopharmacology*, 13(7), 917-932.
- 5 Elkin, I., Shea, M., Watkins, J. T. & et al. (1989). National institute of mental health treatment of depression collaborative research program: General effectiveness of treatments. *Archives of General Psychiatry* 46, 971-982.
- 6 Ell, K., Unützer, J., Aranda, M., Gibbs, N., Lee, P.-J., & Xie, B. (2007). Managing depression in home health care: a randomized clinical trial. *Home Health Care Services Quarterly*, 26, 81-104.
- 7 Ellingrod, V. L., & Perry, P. J. (1994). Venlafaxine: a heterocyclic antidepressant. *Am J Hosp Pharm*, 51(24), 3033-3046.
- 8 Ellis, A. (1955). New approaches to psychotherapy techniques. *Journal of Clinical Psychology*, 11(3), 207-260.
- 9 Ellis, C. G. (1996). Chronic unhappiness. Investigating the phenomenon in family practice. *Can Fam Physician*, 42, 645-651.
- 10 Emanuels-Zuurveen, L., & Emmelkamp, P. (1996). Individual behavioural-cognitive therapy v. marital therapy for depression in maritally distressed couples. *The British Journal of Psychiatry*, 169(2), 181-188.
- 11 Everett, B. (2015). Sexual orientation identity change and depressive symptoms. *Journal of Health and Social Behavior*, 56(1), 37-58.
- 12 Executive, H. a. S. (2008). Self-reported work-related illness and workplace injuries in 2006/07: Results from the labour force survey. .
- 13 Fabre, L. F. (1992). A 6-week, double-blind trial of paroxetine, imipramine, and placebo in depressed outpatients. *Journal of Clinical Psychiatry*, 53 (Suppl.), 40-43
- 14 Fabre, L. & Crismon, L. Efficacy of fluoxetine in outpatients with major depression. *Current therapeutic research* 1985; 37(1):115-123.
- 15 Fabre, L. & McLendon, D. (1980). A double-blind study comparing the efficacy and safety of alprazolam with imipramine and placebo in primary depression, *Current Therapeutic Research*, 27, 474-482.
- 16 Fang, Y., Yuan, C., Xu, Y., Chen, J., Wu, Z., Cao, L., ... Gao, K. (2011). A pilot study of the efficacy and safety of paroxetine augmented with risperidone, valproate, buspirone, trazodone, or thyroid hormone in adult Chinese patients with treatment-resistant major depression. *Journal of Clinical Psychopharmacology*, 31(5), 638-642.
- 17 Fang, Y., Yuan, C., Xu, Y., Chen, J., Wu, Z., Cao, L., ... Nierenberg, A. A. (2010). Comparisons of the efficacy and tolerability of extended-release venlafaxine, mirtazapine, and paroxetine in treatment-resistant depression: a double-blind, randomized pilot study in a Chinese population. *Journal of Clinical Psychopharmacology*, 30(4), 357-364.

- 1 Farah, W. H., Alsawas, M., Mainou, M., Alahdab, F., Farah, M. H., Ahmed, A. T., ... LeBlanc, A. (2016). Non-pharmacological treatment of depression: a systematic review and evidence map. *Evid Based Med*, 21(6), 214-221.
- 4 Farrer L, Christensen H, Griffiths KM, Mackinnon A. Internet-based CBT for depression with and without telephone tracking in a national helpline: randomised controlled trial. *PloS one*. 2011 Nov 30; 6(11):e28099.
- 7 Fava, G. A., Rafanelli, C., Grandi, S., Conti, S., & Belluardo, P. (1998). Prevention of recurrent depression with cognitive behavioral therapy: preliminary findings. *Arch Gen Psychiatry*, 55(9), 816-820.
- 10 Fava, M. (2003). Diagnosis and definition of treatment-resistant depression. *Biol Psychiatry*, 53(8), 649-659.
- 12 Fava, M., & Kendler, K. S. (2000). Major depressive disorder. *Neuron*, 28(2), 335-341.
- 13 Fava M, Alpert J, Nierenberg AA, Mischoulon D, Otto MW, Zajecka J, Murck H, Rosenbaum JF. A double-blind, randomized trial of St John's wort, fluoxetine, and placebo in major depressive disorder. *Journal of clinical psychopharmacology*. 2005 Oct 1; 25(5):441-7.
- 16 Fava, G. A., Grandi, S., Zielezny, M., & Canestrari, R. (1994). Cognitive behavioral treatment of residual symptoms in primary major depressive disorder. *The American Journal of Psychiatry*, 9, 1295–1299.
- 19 Fava, M., Mischoulon, D., Iosifescu, D., Witte, J., Pencina, M., Flynn, M., ... Pollack, M. (2012). A double-blind, placebo-controlled study of aripiprazole adjunctive to antidepressant therapy among depressed outpatients with inadequate response to prior antidepressant therapy (ADAPT-A Study). *Psychotherapy and psychosomatics*, 81(2), 87-97.
- 23 Fava, M., & Rankin, M. (2002). Sexual functioning and SSRIs. *J Clin Psychiatry*, 63 Suppl 5, 13-16; discussion 23-15.
- 25 Fava, M., Rosenbaum, J. F., McGrath, P. J., Stewart, J. W., Amsterdam, J. D., & Quitkin, F. M. (1994). Lithium and tricyclic augmentation of fluoxetine treatment for resistant major depression: a double-blind, controlled study. *American Journal of Psychiatry*, 151(9), 1372-1374.
- 29 Feiger, A. D. (1996). A double-blind comparison of gepirone extended release, imipramine, and placebo in the treatment of outpatient major depression. *Psychopharmacology Bulletin*, 32, 659-665.
- 32 Feiger, A. D., Bielski, R. J., Bremner, J., Heiser, J. F., Trivedi, M., Wilcox, C. S., ... Kaplita, S. B. (1999). Double-blind, placebo-substitution study of nefazodone in the prevention of relapse during continuation treatment of outpatients with major depression. *International Clinical Psychopharmacology*, 14(1), 19-28.
- 36 Feighner, J. P. & Boyer, W. F. (1989). Paroxetine in the treatment of depression: a comparison with imipramine and placebo. *Acta Psychiatrica Scandinavica Supplementum*, 350, 125-129.
- 39 Feighner JP & Overo K. Multicenter, placebo-controlled, fixed-dose study of citalopram in moderate-to-severe depression. *Journal of Clinical Psychiatry* 1999; 60(12):824-830.
- 41 Feighner, J. P., Brauzer, B., Gelenberg, A. J., Gomez, E., Kiev, A., Kurland, M. L. et al. (1979). A placebo-controlled multicenter trial of Limbitrol versus its components (amitriptyline and chlordiazepoxide) in the symptomatic treatment of depressive illness. *Psychopharmacology*, 61, 217-225.

- 1 Feighner, J. P., Cohn, J. B., Fabre, L. F. J., Fieve, R. R., Mendels, J., Shrivastava, R. K. et al. (1993). A study comparing paroxetine placebo and imipramine in depressed patients. *Journal of Affective Disorders*, 28, 71-79.
- 4 Feighner, J. P., Meridith, C. H., Dutt, J. E., & Hendrickson, G. G. (1982). A double blind comparison of lofepramine, imipramine and placebo in patients with depression. *Acta Psychiatrica Scandinavica*, 66, 100-108.
- 7 Feighner, J. P., Meredith, C. H., Frost, N. R., Chammas, S., & Hendrickson, G. (1983). A double-blind comparison of alprazolam vs. imipramine and placebo in the treatment of major depressive disorder. *Acta Psychiatrica Scandinavica*, 68, 223-233.
- 10 Fekadu, A., Wooderson, S., Donaldson, C., Markopoulos, K., Masterson, B., Poon, L., & Cleare, A. J. (2009). A multidimensional tool to quantify treatment resistance in depression: the Maudsley staging method. *J Clin Psychiatry*, 70(2), 177-184.
- 13 Fernandez-Pujals, A. M., Adams, M. J., Thomson, P., McKechnie, A. G., Blackwood, D. H., Smith, B. H., ... McIntosh, A. M. (2015). Epidemiology and Heritability of Major Depressive Disorder, Stratified by Age of Onset, Sex, and Illness Course in Generation Scotland: Scottish Family Health Study (GS:SFHS). *PLoS ONE*, 10(11), e0142197.
- 17 Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J. L., . . . Whiteford, H. A. (2013). Burden of depressive disorders by country, sex, age, and year: Findings from the global burden of disease study 2010. *PLoS Medicine*, 10(11), e1001547.
- 20 Ferreri, M., Lavergne, F., Berlin, I., Payan, C., & Puech, A. (2001). Benefits from mianserin augmentation of fluoxetine in patients with major depression non-responders to fluoxetine alone. *Acta Psychiatrica Scandinavica*, 103(1), 66-72.
- 23 Ferster, C. B. (1973). A functional analysis of depression. *Am Psychol*, 28(10), 857-870.
- 24 Fink, M. (2012). Electroconvulsive therapy. In M. Gelder, N. Andreasen, J. Lopez-Ibor, & J. R. Geddes (Eds.), *New Oxford Textbook of Psychiatry* (2nd ed., pp. 1252-1260): Oxford University Press (OUP).
- 27 Finley, P., Rens, H., Pont, J., Gess, S., Louie, C., Bull, S., Lee, J.Y., Bero, L.A. (2003). Impact of a collaborative care model on depression in a primary care setting: a randomized controlled trial. *Pharmacotherapy*, 23, 1175-1185.
- 30 Fischer, J. E., Bachmann, L. M., & Jaeschke, R. (2003). A readers' guide to the interpretation of diagnostic test properties: clinical example of sepsis. *Intensive Care Med*, 29(7), 1043-1051.
- 33 Fledderus, M., et al. (2012) Acceptance and commitment therapy as guided self-help for psychological distress and positive mental health: a randomized controlled trial. *Psychological Medicine* 42, 485-49.
- 36 Floyd, M., Scogin, F., McKendree-Smith, N. L., Floyd, D. L. & Rokke, P. D. (2004) Cognitive Therapy for depression: A comparison of individual psychotherapy and bibliotherapy for depressed older adults. *Behavior Modification*, 28, 297-318.
- 39 Foley LS, Papavassiliou H, Osuch EA, De Pace JA, Murphy BA, Podolinsky NJ. An examination of potential mechanisms for exercise as a treatment for depression: a pilot study. *Mental Health and Physical Activity*. 2008 Dec 31; 1(2):69-73.
- 42 Folke, F., et al. (2012) Acceptance and commitment therapy for depression: a preliminary randomized controlled trial for unemployed on long-term sick leave. *Cognitive and Behavioral Practice*, 19, 583-594.

- 1 Fonagy, P., Rost, F., Carlyle, J. a., McPherson, S., Thomas, R., Pasco Fearon, R., ... Taylor, D. (2015). Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression: the Tavistock Adult Depression Study (TADS). *World Psychiatry*, 14(3), 312-321.
- 5 Fontaine, R., Ontiveros, A., Elie, R., Kensler, T. T., Roberts, D. L., Kaplita, S. et al. (1994). A double-blind comparison of nefazodone, imipramine, and placebo in major depression. *Journal of Clinical Psychiatry*, 55, 234-241.
- 8 Foran, H. M., Whisman, M. A., & Beach, S. R. H. (2015). Intimate partner relationship distress in the DSM-5. *Family Process*, 54(1), 48-63.
- 10 Forest Research Institute. Double-blind comparison of the effects of lu 26-054 (escitalopram) and fluoxetine on sleep in depressed patients (SCT-MD-09). Report date: July 27, 2004.
- 12 Forest Research Institute. Flexible dose comparison of the safety and efficacy of escitalopram and fluoxetine in the treatment of major depressive disorder (SCT-MD-16). Report date: July 26, 2004.
- 15 Forest Research Institute. Two-week double-blind placebo controlled study of escitalopram in the treatment of severe major depression. Report date: November 26, 2003
- 17 Forest Laboratories Inc. Flexible dose comparison of the saftey and efficacy of lu 26-054, citalopram, and placebo in the treatment of major depressive disorder (SCT-MD-02). Report date: December 5, 2000.
- 20 Fortney, J., Pyne, J., Edlund, M., Williams, D., Robinson, D., Mittal, D., Henderson, K.L. (2007). A randomized trial of telemedicine-based collaborative care for depression. *Journal of General Internal Medicine*, 22, 1086-1093.
- 23 Fortney, J. C., Pyne, J. M., Mouden, S. B., Mittal, D., Hudson, T. J., Schroeder, G. W., ... Rost, K. M. (2013). Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *The American Journal of Psychiatry*, 170(4), 414-425.
- 27 Fountoulakis, K. N., & Möller, H.-J. (2010). Efficacy of antidepressants: a re-analysis and re-interpretation of the Kirsch data. *The International Journal of Neuropsychopharmacology*, 14(03), 405-412.
- 30 Fournier, J. C., DeRubeis, R. J., Hollon, S. D., Dimidjian, S., Amsterdam, J. D., Shelton, R. C., & Fawcett, J. (2010). Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA*, 303(1), 47.
- 33 Fournier J-P, Lane RM, Chouinard G, Watson DB, Amin M, Remick RA, Thorpe LU. A double-blind comparison of sertraline and imipramine in outpatients with major depression: Acute (8 weeks) and continuation (16 weeks) treatment. *Human Psychopharmacology* 1997; 12(3):203- 215.
- 37 Franchini, L., Gasperini, M., Perez, J., Smeraldi, E., & Zanardi, R. (1998). Dose-response efficacy of paroxetine in preventing depressive recurrences: a randomized, double-blind study. *The Journal of Clinical Psychiatry*, 59(5), 229-232.
- 40 Frank E, Cassano GB, Rucci P, Thompson WK, Kraemer HC, Fagioli A, Maggi L, Kupfer DJ, Shear MK, Houck PR, Calugi S. Predictors and moderators of time to remission of major depression with interpersonal psychotherapy and SSRI pharmacotherapy. *Psychological medicine*. 2011; 41(01):151-62.
- 44 Frank, E., Kupfer, D. J., Perel, J. M., Cornes, C., Jarrett, D. B., Mallinger, A. G., ... Grochocinski, V. J. (1990). Three-year outcomes for maintenance therapies in recurrent depression. *Arch Gen Psychiatry*, 47(12), 1093-1099.

- 1 Freeman, C. (2013). Cognitive adverse effects of ECT. In J. Waite & A. Easton (Eds.), *The ECT handbook* (Third edition ed.): RCPsych Publications.
- 3 Freire, E., et al. (2015). "Counselling versus low-intensity cognitive behavioural therapy for persistent sub-threshold and mild depression (CLICD): A pilot/feasibility randomised controlled trial." *BMC psychiatry* 15(1).
- 6 Fremont J, Craighead LW. Aerobic exercise and cognitive therapy in the treatment of dysphoric moods. *Cognitive therapy and Research*. 1987 Apr 1; 11(2):241-51.
- 8 Freud, S. (1917 (reprinted)). Mourning and melancholia. In J. Strachey (Ed.), *The standard edition of the complete psychological works of Sigmund Freud* (Vol. 14). London: Hogarth Press.
- 11 Fu, C. H., Williams, S. C., Brammer, M. J., Suckling, J., Kim, J., Cleare, A. J., . . . Pich, E. M. (2007). Neural responses to happy facial expressions in major depression following antidepressant treatment. *American Journal of Psychiatry*, 164(4), 599-607.
- 14 Furukawa, T. A., Barbui, C., Cipriani, A., Brambilla, P., & Watanabe, N. (2006). Imputing missing standard deviations in meta-analyses can provide accurate results. *J Clin Epidemiol*, 59(1), 7-10.
- 17 Furukawa, T. A., Kitamura, T., & Takahashi, K. (2000). Time to recovery of an inception cohort with hitherto untreated unipolar major depressive episodes. *Br J Psychiatry*, 177, 331-335.
- 20 Furukawa, T. A., et al. (2012) Telephone cognitive-behavioral therapy for subthreshold depression and presenteeism in workplace: a randomized controlled trial. *PloS One* 7.
- 22 Furukawa, T. A., Noma, H., Caldwell, D. M., Honyashiki, M., Shinohara, K., Imai, H., . . . Churchill, R. (2014). Waiting list may be a nocebo condition in psychotherapy trials: a contribution from network meta-analysis. *Acta Psychiatr Scand*, 130(3), 181-192.
- 25 Gabbard, G. O. (2017). Long-term psychodynamic psychotherapy: A basic text: American Psychiatric Pub.
- 27 Galante, J., Iribarren, S. J., & Pearce, P. F. (2013). Effects of mindfulness-based cognitive therapy on mental disorders: A systematic review and meta-analysis of randomised controlled trials. *Journal of Research in Nursing*, 18(2), 133-155.
- 30 Gallagher DE, Thompson LW. Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. *Psychother Theory Res Pract* 1982; 19(4):482-490.
- 32 Gallagher-Thompson DE, Steffen AM. Comparative effects of cognitive behavioral therapy and brief psychodynamic psychotherapies for depressed family caregivers. *J Consult Clin Psychol* 1994; 62(3):543-549.
- 35 Gansler, D. A., Suvak, M., Arean, P., & Alexopoulos, G. S. (2015). Role of executive dysfunction and dysexecutive behavior in late-life depression and disability. *The American Journal of Geriatric Psychiatry*, 23(10), 1038-1045.
- 38 Gask, L., Dowrick, C., Dixon, C., Sutton, C., Perry, R., Torgerson, D., & Usherwood, T. (2004). A pragmatic cluster randomized controlled trial of an educational intervention for GPs in the assessment and management of depression. *Psychological Medicine*, 34(1), 63-72.
- 41 Gask, L., Goldberg, D., Lesser, A. L., & Millar, T. (1988). Improving the psychiatric skills of the general practice trainee: an evaluation of a group training course. *Medical Education*, 22(2), 132-138.

- 1 Gask, L., Rogers, A., Oliver, D., May, C., & Roland, M. (2003). Qualitative study of patients' perceptions of the quality of care for depression in general practice. *Br J Gen Pract*, 53(489), 278-283.
- 4 Gask, L., Sibbald, B., & Creed, F. (1997). Evaluating models of working at the interface between mental health services and primary care. *Br J Psychiatry*, 170, 6-11.
- 6 Gastpar M, Singer A, Zeller K. Comparative efficacy and safety of a once-daily dosage of hypericum extract STW3-VI and citalopram in patients with moderate depression: A double-blind, randomised, multicentre, placebo-controlled study. *Pharmacopsychiatry* 2006; 39(2):66-75.
- 10 Gater R, Waheed W, Husain N, Tomenson B, Aseem S, Creed F. Social intervention for British Pakistani women with depression: randomised controlled trial. *The British Journal of Psychiatry*. 2010 Sep 1; 197(3):227-33.
- 13 Gaudiano, B. A., & Miller, I. W. (2013). The evidence-based practice of psychotherapy: facing the challenges that lie ahead. *Clin Psychol Rev*, 33(7), 813-824.
- 15 Gawrysiak, M., Nicholas, C., & Hopko, D. R. (2009). Behavioral activation for moderately depressed university students: Randomized controlled trial. *Journal of Counseling Psychology*, 56(3), 468-475.
- 18 Geddes, J. R., Carney, S. M., Davies, C., Furukawa, T. A., Kupfer, D. J., Frank, E., & Goodwin, G. M. (2003). Relapse prevention with antidepressant drug treatment in depressive disorders: a systematic review. *Lancet*, 361(9358), 653-661.
- 21 Gelenberg, A. J., Wojcik, J. D., Falk, W. E., Baldessarini, R. J., Zeisel, S. H., Schoenfeld, D. et al. (1990). Tyrosine for depression: a double-blind trial. *Journal of Affective Disorders*, 19, 125-132.
- 24 Gelenberg, A. J., Wojcik, J. D., Falk, W. E., Spring, B., Brotman, A. W., & Galvin-Nadeau, M. (1990). Clovoxamine in the treatment of depressed outpatients: a double-blind, parallel-group comparison against amitriptyline and placebo. *Comprehensive Psychiatry*, 31(4), 307-314.
- 27 Gensichen, J., Beyer, M., Muth, C., Gerlach, F. M., Von Korff, M., & Ormel, J. (2006). Case management to improve major depression in primary health care: a systematic review. *Psychol Med*, 36(1), 7-14.
- 30 Gensichen, J., vonKorff, M., Peitz, M., Muth, C., Beyer, M., Guthlin, C., Torge, M., Petersen, J.J., Rosemann, T., König, J., Gerlach, F.M.; PRoMPT (PRimary care Monitoring for depressive Patients Trial). (2009). Case management for depression by health care assistants in small primary care practices: a cluster randomized trial. *Annals of Internal Medicine*, 151(6), 369-378.
- 35 Georgotas, A., McCue, R., & Cooper, T. (1989). A placebo-controlled comparison of nortriptyline and phenelzine in maintenance therapy of elderly depressed patients. *Archives of General Psychiatry*, 46(9), 783-786.
- 38 Georgotas, A., McCue, R. E., Friedman, E., Kim, O. M., Welkowitz, J., Chang, I., & Cooper, T. B. (1986). Comparative efficacy and safety of MAOIs versus TCAs in treating depression in the elderly. *Biological Psychiatry*, 21(12), 1155-1166.
- 41 Georgotas, A., Krakowski, M., & Gershon, S. (1982). Controlled trial of zimelidine, a 5-HT reuptake inhibitor, for treatment of depression. *American Journal of Psychiatry*, 139, 1057-1058
- 44 Geraedts Anna, S., Kleiboer Annet, M., Twisk, J., Wiezer Noortje, M., Mechelen, W., & Cuijpers, P. (2014). Long-term results of a Web-based guided self-help intervention for

- 1 employees with depressive symptoms: Randomized controlled trial. Journal of Medical Internet Research, 16(7), 14-28.
- 3 Gerber, P. D., Barrett, J. E., Barrett, J. A., Oxman, T. E., Manheimer, E., Smith, R., & Whiting, R. D. (1992). The relationship of presenting physical complaints to depressive symptoms in primary care patients. Journal of General Internal Medicine, 7(2), 170-173.
- 6 Gertsik, L., Poland, R. E., Bressee, C., & Rapaport, M. H. (2012). Omega-3 fatty acid augmentation of citalopram treatment for patients with major depressive disorder. Journal of Clinical Psychopharmacology, 32(1), 61-64.
- 9 Gibbons, M. B., et al. (2012) Supportive-expressive dynamic psychotherapy in the community mental health system: a pilot effectiveness trial for the treatment of depression. Psychotherapy (Chicago, Ill.) 49, 303-31.
- 12 Gilaberte, I., Montejo, A. L., de la Gandara, J., Perez-Sola, V., Bernardo, M., Massana, J., . . . Casais, L. (2001). Fluoxetine in the prevention of depressive recurrences: a double-blind study. Journal of Clinical Psychopharmacology, 21(4), 417-424.
- 15 Gilbody, S. (2006). Should we screen for depression? BMJ, 332(7548), 1027-1030.
- 16 Gilbody, S., Bower, P., Fletcher, J., Richards, D., & Sutton, A. J. (2006). Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. Arch Intern Med, 166(21), 2314-2321.
- 19 Gilbody, S., Littlewood, E., Hewitt, C., Brierley, G., Tharmanathan, P., Araya, R., ... White, D. (2015). Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REACT trial): Large scale pragmatic randomised controlled trial. BMJ 2015; 351:h5627.
- 23 Gilbody, S., Richards, D., & Barkham, M. (2007). Diagnosing depression in primary care using self-completed instruments: UK validation of PHQ-9 and CORE-OM. Br J Gen Pract, 57(541), 650-652.
- 26 Gilbody, S., Richards, D., Brealey, S., & Hewitt, C. (2007). Screening for depression in medical settings with the Patient Health Questionnaire (PHQ): a diagnostic meta-analysis. J Gen Intern Med, 22(11), 1596-1602.
- 29 Gilbody, S., Sheldon, T., & House, A. (2008). Screening and case-finding instruments for depression: a meta-analysis. Canadian Medical Association Journal, 178(8), 997-1003.
- 31 Gilbody, S., Whitty, P., Grimshaw, J., & Thomas, R. (2003). Educational and organizational interventions to improve the management of depression in primary care. JAMA, 289(23), 3145.
- 34 Giles, D. E., Jarrett, R. B., Biggs, M. M., Guzick, D. S., & Rush, A. J. (1989). Clinical predictors of recurrence in depression. Am J Psychiatry, 146(6), 764-767.
- 36 Gili, M., Vicens, C., Roca, M., Andersen, P., & McMillan, D. (2015). Interventions for preventing relapse or recurrence of depression in primary health care settings: A systematic review. Preventive Medicine, 76(S), S16-S21. Gillen, R., Tennen, H., McKee, T. E., Gernert-Dott, P., & Affleck, G. (2001). Depressive symptoms and history of depression predict rehabilitation efficiency in stroke patients. Arch Phys Med Rehabil, 82(12), 1645-1649.
- 41 Ginty, A. T., & Conklin, S. M. (2015). Short-term supplementation of acute long-chain omega-3 polyunsaturated fatty acids may alter depression status and decrease symptomatology among young adults with depression: A preliminary randomized and placebo controlled trial. Psychiatry Research, 229(1-2), 485-489.

- 1 Ginzburg, K., Ein-Dor, T., & Solomon, Z. (2010). Comorbidity of posttraumatic stress disorder, anxiety and depression: a 20-year longitudinal study of war veterans. *J Affect Disord*, 123(1-3), 249-257.
- 4 Girlanda F, Cipriani A, Agrimi E, Appino MG, Barichello A, Beneduce R, Bighelli I, Bisoffi G, Bisogno A, Bortolaso P, Bosco M. Effectiveness of lithium in subjects with treatment-resistant depression and suicide risk: results and lessons of an underpowered randomised clinical trial. *BMC Research Notes*. 2014 Oct 17; 7(1):731.
- 8 Glasgow, R. E., & Rosen, G. M. (1978). Behavioral bibliotherapy: a review of self-help behavior therapy manuals. *Psychol Bull*, 85(1), 1-23.
- 10 Glassman, A. H., & Bigger, J. T., Jr. (2001). Antipsychotic drugs: prolonged QTc interval, torsade de pointes, and sudden death. *Am J Psychiatry*, 158(11), 1774-1782.
- 12 Glen AI, Johnson AL, Shepherd M. Continuation therapy with lithium and amitriptyline in unipolar depressive illness: a randomized, double-blind, controlled trial. *Psychological Medicine* 1984; 14(1):37-50.
- 15 Glick, I. D., Fleming, L., DeChillo, N., Meyerkopf, N., Jackson, C., Muscara, D., & Good-Ellis, M. (1986). A controlled study of transitional day care for non-chronically ill patients. *American Journal of Psychiatry*, 143(12), 1551-1556.
- 18 Godfrin, K. A., & Heeringen, C. (2010). The effects of mindfulness-based cognitive therapy on recurrence of depressive episodes, mental health and quality of life: A randomized controlled study. *Behaviour Research & Therapy*, 48(8), 738-746.
- 21 Goethe, J. W., Woolley, S. B., Cardoni, A. A., Woznicki, B. A., & Piez, D. A. (2007). Selective serotonin reuptake inhibitor discontinuation: side effects and other factors that influence medication adherence. *J Clin Psychopharmacol*, 27(5), 451-458.
- 24 Goldberg, D., Privett, M., Ustun, B., Simon, G., & Linden, M. (1998). The effects of detection and treatment on the outcome of major depression in primary care: a naturalistic study in 15 cities. *Br J Gen Pract*, 48(437), 1840-1844.
- 27 Goldberg, D., & Williams, P. (1991). A user's guide to the general health questionnaire: NFER-Nelson.
- 29 Goldberg, D. P., & Bridges, K. (1988). Somatic presentations of psychiatric illness in primary care setting. *Journal of Psychosomatic Research*, 32(2), 137-144.
- 31 Goldberg, D. P., & Huxley, P. (1980). Mental illness in the community: the pathway to psychiatric care. London; New York: Tavistock Publications.
- 33 Goldberg, D. P., & Huxley, P. (1992). Common mental disorders: a bio-social model. London; New York: Tavistock/Routledge.
- 35 Goldberg, D. P., Jenkins, L., Millar, T., & Faragher, E. B. (1993). The ability of trainee general practitioners to identify psychological distress among their patients. *Psychological Medicine*, 23(01), 185.
- 38 Goldman, R.N., Greenberg, L.S. & Angus, L. (2006) The effects of adding emotion-focused interventions to the client-centered relationship conditions in the treatment of depression. *Psychotherapy Research*, 16, 537-549.
- 41 Gonzales, L. R., Lewinsohn, P. M., & Clarke, G. N. (1985). Longitudinal follow-up of unipolar depressives: an investigation of predictors of relapse. *J Consult Clin Psychol*, 53(4), 461-469.
- 43 Goodwin, G. (2000). Neurobiological aetiology of mood disorders. In M. G. Gelder, J. J. Lopez-Ibor, & N. C. Andreasen (Eds.), *New Oxford textbook of psychiatry*. Oxford: Oxford University Press.

- 1 Goodwin, G. M., Emsley, R., Rembry, S., & Rouillon, F. (2009). Agomelatine prevents relapse in patients with major depressive disorder without evidence of a discontinuation syndrome: a 24-week randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry*, 70(8), 1128-1137.
- 5 Goorden, M., Huijbregts, K. M. L., Van Marwijk, H. W. J., Beekman, A. T. F., Feltz-Cornelis, C. M., & Hakkaart-van Roijen, L. (2015). Cost-utility of collaborative care for major depressive disorder in primary care in the Netherlands. *Journal of Psychosomatic Research*, 79, 316-323.
- 9 Goorden, M., Vlasveld, M. C., Anema, J. R., Van Mechelen, W., Beekman, A. T., Hoedeman, R., ... Hakkaart-van Roijen, L. (2014). Cost-utility analysis of a collaborative care intervention for major depressive disorder in an occupational healthcare setting. *Journal of occupational rehabilitation*, 24, 555-562.
- 13 Gorwood, P., Weiller, E., Lemming, O., & Katona, C. (2007). Escitalopram prevents relapse in older patients with major depressive disorder. *The American Journal of Geriatric Psychiatry*, 15(7), 581-593.
- 16 Gottschalk, M. G., Wesseling, H., Guest, P. C., & Bahn, S. (2014). Proteomic enrichment analysis of psychotic and affective disorders reveals common signatures in presynaptic glutamatergic signaling and energy metabolism. *Int J Neuropsychopharmacol*, 18(2) pii: pyu019.
- 20 Grant, B. F., Chou, S. P., Goldstein, R. B., Huang, B., Stinson, F. S., Saha, T. D., . . . Ruan, W. J. (2008). Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*, 69(4), 533-545.
- 24 Green, C., Richards, D. A., Hill, J. J., Gask, L., Lovell, K., Chew-Graham, C., . . . Barkham, M. (2014). Cost-effectiveness of collaborative care for depression in UK primary care: Economic evaluation of a randomised controlled trial (CADET). *PLoS ONE*, 9(8), e104225.
- 27 Greenberg, L.S. (1998). Experiential therapy of depression: differential effects of client-centred relationship conditions and process experiential interventions. *Psychotherapy Research*, 8, 210-224.
- 30 Greenberg, J. R., & Mitchell, S. A. (1983). Object relations in psychoanalytic theory. Cambridge, Mass.: Harvard University Press.
- 32 Greenberg, P. E., Fournier, A.-A., Sisitsky, T., Pike, C. T., & Kessler, R. C. (2015). The economic burden of adults with major depressive disorder in the United States (2005 and 2010). *The Journal of Clinical Psychiatry*, 155-162.
- 35 Greene, R., Pugh, R., & Roberts, D. (2008). Black and minority ethnic parents with mental health problems and their children.
- 37 Greenhalgh, J., Knight, C., Hind, D., Beverley, C., & Walters, S. (2005). Clinical and cost-effectiveness of electroconvulsive therapy for depressive illness, schizophrenia, catatonia and mania: Systematic reviews and economic modelling studies. *Health Technology Assessment*, 9(9).
- 41 Griffiths, K.M., Crisp, D. A., Barney, L. and Reid, R. (2011). Seeking help for depression from family and friends: A qualitative analysis of perceived advantages and disadvantages. *BMC Psychiatry* 11:196.
- 44 Griffiths, K. M., Mackinnon, A. J., Crisp, D. A., Christensen, H., Bennett, K., & Farrer, L. (2012). The effectiveness of an online support group for members of the community with depression: a randomised controlled trial. *PLoS ONE*, 7(12), e53244.

- 1 Grilo, C. M., White, M. A., & Masheb, R. M. (2009). DSM-IV psychiatric disorder comorbidity and its correlates in binge eating disorder. *Int J Eat Disord*, 42(3), 228-234.
- 3 Group, M. R. (2005). Do patients want to disclose psychological problems to GPs? *Family Practice*, 22, 631-637.
- 5 Group, M. R. (2005). General practitioners' perceptions of barriers to their provision of mental healthcare: a report on Mental Health and General Practice Investigation (MaGPIe). *The New Zealand Medical Journal*, 118(1222), U1654.
- 8 Group, U. E. R. (2003). Efficacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and meta-analysis. *Lancet*, 361(9360), 799-808.
- 10 Grunhaus, L., Hirschman, S., Dolberg, O. T., Schreiber, S., & Dannon, P. N. (2001). Coadministration of melatonin and fluoxetine does not improve the 3-month outcome following ECT. *The journal of ECT*, 17(2), 124-128.
- 13 Guasp, A. (2010). *Lesbian, gay & bisexual people in later life*. London: Stonewall.
- 14 Gulliford, M., Figueiroa-Munoz, J., Morgan, M., Hughes, D., Gibson, B., Beech, R., & Hudson, M. (2002). What does 'access to health care' mean? *J Health Serv Res Policy*, 7(3), 186-188.
- 16 Gulrez, G., Badyal, D. K., Deswal, R. S., & Sharma, A. (2012). Bupropion as an augmenting agent in patients of depression with partial response. *Basic & Clinical Pharmacology & Toxicology*, 110(3), 227-230.
- 19 Gunn, J., Diggens, J., Hegarty, K., & Blashki, G. (2006). A systematic review of complex system interventions designed to increase recovery from depression in primary care. *BMC Health Serv Res*, 6, 88.
- 22 Guo T, Xiang YT, Xiao L, Hu CQ, Chiu HF, Ungvari GS, Correll CU, Lai KY, Feng L, Geng Y, Feng Y. Measurement-based care versus standard care for major depression: a randomized controlled trial with blind raters. *American Journal of Psychiatry*. 2015 Aug 28; 172(10):1004-13.
- 26 Guy, W. (1976). Early clinical drug evaluation programme assessment manual for psychopharmacology - Revised. Rockville, MD: US Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, NIMH.
- 30 Guyatt, G., Oxman, A. D., Akl, E. A., Kunz, R., Vist, G., Brozek, J., ... Schunemann, H. J. (2011). GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*, 64(4), 383-394.
- 33 Haboush A, Floyd M, Caron J, LaSota M, Alvarez K. Ballroom dance lessons for geriatric depression: An exploratory study. *The Arts in Psychotherapy*. 2006 Dec 31; 33(2):89-97.
- 35 Haghghi M, Salehi I, Erfani P, Jahangard L, Bajoghli H, Holsboer-Trachsler E, Brand S. Additional ECT increases BDNF-levels in patients suffering from major depressive disorders compared to patients treated with citalopram only. *Journal of Psychiatric Research*. 2013 Jul 31; 47(7):908-15.
- 39 Halikas JA. Org 3770 (mirtazapine) versus trazodone: a placebo controlled trial in depressed elderly patients. *Human Psychopharmacology: Clinical and Experimental*. 1995 Jul 1; 10(S2):S125-33.
- 42 Hallgren, M., Kraepelien, M., Ojehagen, A., Lindefors, N., Zeebari, Z., Kaldo, V., & Forsell, Y. (2015). Physical exercise and internet-based cognitive-behavioural therapy in the treatment of depression: Randomised controlled trial. *British Journal of Psychiatry*, 207(3 // *Brain Foundation*), 227-234.

- 1 Hamidian S, Omidi A, Mousavinasab SM, Naziri G. Comparison of the effect of mindfulness-based cognitive therapy accompanied by pharmacotherapy with pharmacotherapy alone in treating dysthymic patients. *Iranian Red Crescent Medical Journal*. 2013 Mar; 15(3):239.
- 4 Hamamci Z. Integrating psychodrama and cognitive behavioral therapy to treat moderate depression. *The Arts in Psychotherapy*. 2006 Dec 31; 33(3):199-207.
- 6 Hamdan-Mansour AM, Puskar K, Bandak AG. Effectiveness of cognitive-behavioral therapy on depressive symptomatology, stress and coping strategies among Jordanian university students. *Issues in Mental Health Nursing*. 2009; 30(3):188-96.
- 9 Hamidian S, Omidi A, Mousavinasab SM, Naziri G. Comparison of the effect of mindfulness-based cognitive therapy accompanied by pharmacotherapy with pharmacotherapy alone in treating dysthymic patients. *Iranian Red Crescent Medical Journal*. 2013 Mar; 15(3):239.
- 12 Hammen, C., Henry, R., & Daley, S. E. (2000). Depression and sensitization to stressors among young women as a function of childhood adversity. *Journal of Consulting and Clinical Psychology*, 68(5), 782-787.
- 15 Hansen, D. G., Vach, W., Roshholm, J. U., Sondergaard, J., Gram, L. F., & Kragstrup, J. (2004). Early discontinuation of antidepressants in general practice: association with patient and prescriber characteristics. *Fam Pract*, 21(6), 623-629.
- 18 Hansson, M., Bodlund, O., Chotai, J. (2008) Patient education and group counselling to improve the treatment of depression in primary care: a randomized controlled trial. *Journal of Affective Disorders*, 105, 235-240.
- 21 Hansgen, K. D., & Vesper, J. (1996). Antidepressive effectiveness of a highly dosed hypericum extract. *Munchener Medizinische Wochenschrift*, 138(3), 35-39.
- 23 Hardeveld, F., Spijker, J., De Graaf, R., Nolen, W. A., & Beekman, A. T. (2010). Prevalence and predictors of recurrence of major depressive disorder in the adult population. *Acta Psychiatr Scand*, 122(3), 184-191.
- 26 Hardeveld, F., Spijker, J., De Graaf, R., Nolen, W. A., & Beekman, A. T. (2013). Recurrence of major depressive disorder and its predictors in the general population: results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Psychol Med*, 43(1), 39-48.
- 30 Hardy, G. E., Barkham, M., Shapiro, D. A., Stiles, W. B., Rees, A., & Reynolds, S. (1995). Impact of Cluster C personality disorders on outcomes of contrasting brief psychotherapies for depression. *J Consult Clin Psychol*, 63(6), 997-1004.
- 33 Haringsma, R., Engels, G. I., Beekman, A. T., & Spinhoven, P. (2004). The criterion validity of the Center for Epidemiological Studies Depression Scale (CES-D) in a sample of self-referred elders with depressive symptomatology. *Int J Geriatr Psychiatry*, 19(6), 558-563.
- 36 Haringsma, R., Engels, G. I., Cuijpers, P., & Spinhoven, P. (2006) Effectiveness of the Coping With Depression (CWD) course for older adults provided by the community-based mental health caresystem in the Netherlands: a randomized controlled field trial. *International Psychogeriatrics*, 18, 307-325.
- 40 Harris, T. O. (2000). Introduction to the work of George Brown. In T. O. Harris (Ed.), *Where inner and outer worlds meet: psychosocial research in the tradition of George W. Brown* (pp. 1-52). London, New York: Routledge.
- 43 Hasin, D. S., Goodwin, R. D., Stinson, F. S., & Grant, B. F. (2005). Epidemiology of major depressive disorder. *Archives of General Psychiatry*, 62(10), 1097.

- 1 Hayhurst, H., Cooper, Z., Paykel, E. S., Vearnals, S., & Ramana, R. (1997). Expressed emotion and depression. A longitudinal study. *Br J Psychiatry*, 171, 439-443.
- 3 Hedrick, S. C., Chaney, E. F., Felker, B., Liu, C. F., Hasenberg, N., Heagerty, P., ... Katon, W. (2003). Effectiveness of collaborative care depression treatment in Veterans' Affairs primary care. *J Gen Intern Med*, 18(1), 9-16.
- 6 Hegerl U, Hautzinger M, Mergl R, Kohnen R, Schutze M, Scheunemann W, et al. Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: A randomized, controlled trial including a patients' choice arm. *International Journal of Neuropsychopharmacology*. 2010; 13(1):31-44.
- 10 Hellerstein, D. J., Batchelder, S. T., Hyler, S., Arnaout, B., Toba, C., Benga, I., & Gangure, D. (2010). Escitalopram versus placebo in the treatment of dysthymic disorder. *International Clinical Psychopharmacology*, 25(3), 143-148.
- 13 Hellerstein, D. J., Little, S. A., Samstag, L. W., Batchelder, S., Muran, J. C., Fedak, M., ... Winston, A. (2001). Adding group psychotherapy to medication treatment in dysthymia: a randomized prospective pilot study. *The Journal of Psychotherapy Practice and Research*, 10(2), 93.
- 17 Hellerstein, D. J., Rosenthal, R. N., Pinsker, H., Samstag, L. W., Muran, J. C., & Winston, A. (1998). A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. *J Psychother Pract Res*, 7(4), 261-271.
- 20 Hellerstein, D. J., Stewart, J. W., McGrath, P. J., Deliyannides, D. A., Batchelder, S. T., Black, S. R., ... Chen, Y. (2012). A randomized controlled trial of duloxetine versus placebo in the treatment of nonmajor chronic depression. *Journal of Clinical Psychiatry*, 73(7), 984-991.
- 23 Hellerstein, D. J., Yanowitch, P., Rosenthal, J., Samstag, L. W., Maurer, M., Kasch, K., ... Winston, A. (1993). A randomized double-blind study of fluoxetine versus placebo in the treatment of dysthymia. *American Journal of Psychiatry*, 150, 1169-1169.
- 26 Hemat-Far A, Shahsavari A, Roholla Mousavi S. Effects of selected aerobic exercises on the depression and concentrations of plasma serotonin in the depressed female students aged 18 to 25. *Journal of Applied Research in Clinical and Experimental Therapeutics*. 2012; 12(1):47.
- 30 Hemeryck, A., & Belpaire, F. M. (2002). Selective serotonin reuptake inhibitors and cytochrome P-450 mediated drug-drug interactions: an update. *Curr Drug Metab*, 3(1), 13-37.
- 33 Henry, J. A., Alexander, C. A., & Sener, E. K. (1995). Relative mortality from overdose of antidepressants. *BMJ*, 310(6974), 221-224.
- 35 Herman S, Blumenthal JA, Babyak M, Khatri P, Craighead WE, Krishnan KR, Doraiswamy PM. Exercise therapy for depression in middle-aged and older adults: Predictors of early dropout and treatment failure. *Health Psychology*. 2002 Nov; 21(6):553.
- 38 Hersen M, Himmelhoch JM, Thase ME, Bellack AS. Effects of social skill training, amitriptyline, and psychotherapy in unipolar depressed women. *Behavior Therapy*. 1984 Jan 31; 15(1):21-40.
- 41 Hicks, F., Robins, E., & Murphy, G. E. (1988). Comparison of adinazolam, amitriptyline, and placebo in the treatment of melancholic depression. *Psychiatry Research*, 23(2), 221-227.
- 43 Higgins, J., & Green, S. (Eds.). (2011). *Cochrane handbook for systematic reviews of interventions Version 5.1.0 (updated March 2011)*: The Cochrane Collaboration, Available from www.cochrane-handbook.org.

- 1 Hirschfeld, R. M. (1999). Personality disorders and depression: comorbidity. *Depress & Anxiety*, 10(4), 142-146.
- 3 Ho, C. W., Chan, S., Wong, J., Cheung, W., Chung, D. W., & Lau, T. F. (2014). Effect of aerobic exercise training on Chinese population with mild to moderate depression in Hong Kong. *Rehabilitation research and practice*, 2014.
- 6 Ho, F. Y., Yeung, W. F., Ng, T. H., & Chan, C. S. (2016). The efficacy and cost-effectiveness of stepped care prevention and treatment for depressive and/or anxiety disorders: A systematic review and meta-analysis. *Sci Rep*, 6, 29281.
- 9 Hochstrasser, B., Isaksen, P., Koponen, H., Lauritzen, L., Mahnert, F., Rouillon, F., ... De Swart, J. (2001). Prophylactic effect of citalopram in unipolar, recurrent depression. *The British Journal of Psychiatry*, 178(4), 304-310.
- 12 Hodge, S., & Buley, N. (2014). ECTAS Fifth National Report. .
- 13 Hoffman BM, Blumenthal JA, Babyak MA, Smith PJ, Rogers SD, Doraiswamy PM, Sherwood A. Exercise fails to improve neurocognition in depressed middle-aged and older adults. Medicine and science in sports and exercise. 2008 Jul; 40(7):1344.
- 16 Hoifodt, R. S., Lillevoll, K. R., Griffiths, K. M., Wilsgaard, T., Eisemann, M., Waterloo, K., & Kolstrup, N. (2013). The clinical effectiveness of web-based cognitive behavioral therapy with face-to-face therapist support for depressed primary care patients: randomized controlled trial. *Journal of Medical Internet Research*, 15(8), e153.
- 20 Hollandare, F., Anthony, S. A., Randestad, M., Tillfors, M., Carlbring, P., Andersson, G., & Engstrom, I. (2013). Two-year outcome of internet-based relapse prevention for partially remitted depression. *Behaviour Research & Therapy*, 51(11), 719-722. Holländare F, Johnsson S, Randestad M, Tillfors M, Carlbring P, Andersson G, et al. Randomized trial of Internet-based relapse prevention for partially remitted depression. *Acta Psychiatrica Scandinavica* [Internet]. 2011; 124(4): [285-94 pp.].
- 26 Hollinghurst, S., Carroll, F. E., Abel, A., Campbell, J., Garland, A., Jerrom, B., ... Wiles, N. (2014). Cost-effectiveness of cognitive-behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: Economic evaluation of the CoBalT Trial. *British Journal of Psychiatry*, 204, 69-76.
- 30 Hollinghurst, S., Peters, T. J., Kaur, S., Wiles, N., Lewis, G., & Kessler, D. (2010). Cost-effectiveness of therapist-delivered online cognitive-behavioural therapy for depression: Randomised controlled trial. *British Journal of Psychiatry*, 197, 297-304.
- 33 Hollon, S. D., DeRubeis, R. J., Shelton, R. C., Amsterdam, J. D., Salomon, R. M., O'Reardon, J. P., . . . Freeman, B. B. (2005). Prevention of relapse following cognitive therapy vs medications in moderate to severe depression. *Archives of General Psychiatry*, 62(4), 417-422.
- 37 Hollon, S. D., DeRubeis, R. J., Evans, M. D., Wiemer, M. J., Garvey, M. J., Grove, W. M. & Tuason, V. B. (1992). Cognitive therapy and pharmacotherapy for depression. Singly and in combination. *Arch Gen Psychiatry* 49, 774-81.
- 40 Hollyman, J. A., Freeling, P., Paykel, E., Bhat, A., & Sedgwick, P. (1988). Double-blind placebo-controlled trial of amitriptyline among depressed patients in general practice. *JR Coll Gen Pract*, 38(314), 393-397.
- 43 Holma, K. M., Holma, I. A., Melartin, T. K., Rytsala, H. J., & Isometsa, E. T. (2008). Long-term outcome of major depressive disorder in psychiatric patients is variable. *J Clin Psychiatry*, 69(2), 196-205.

- 1 Holman, A. J., Serfaty, M. A., Leurent, B. E., & King, M. B. (2011). Cost-effectiveness of cognitive behaviour therapy versus talking and usual care for depressed older people in primary care. *BMC Health Services Research*, 11, 33.
- 4 Holmes, J. (2001). The search for the secure base: attachment theory and psychotherapy. Hove, East Sussex; Philadelphia, PA: Brunner-Routledge; Taylor & Francis.
- 6 Hölzel L, Härtel M, Reese C, Kriston L. (2011). Risk factors for chronic depression--a systematic review. *J Affect Disord*. 129(1-3):1-13.
- 8 Hooley, J. M., & Teasdale, J. D. (1989). Predictors of relapse in unipolar depressives: expressed emotion, marital distress, and perceived criticism. *J Abnorm Psychol*, 98(3), 229-235.
- 11 Hopko, D.R., Lejuez, C.W., LePage, J.P., Hopko, S.D., & McNeil, D.W. (2003) A brief behavioural activation treatment for depression. *Behaviour Modification*, 27 (4), 458-469.
- 13 Hopko, D. R., Lejuez, C. W., Ruggiero, K. J., & Eifert, G. H. (2003). Contemporary behavioral activation treatments for depression: procedures, principles, and progress. *Clin Psychol Rev*, 23(5), 699-717.
- 16 Hormazabal, L., Omer, L. M., & Ismail, S. (1985). Cianopramine and amitriptyline in the treatment of depressed patients--a placebo-controlled study. *Psychopharmacology*, 86, 205-208.
- 19 Horrell, L., Goldsmith, K. A., Tylee, A. T., Schmidt, U. H., Murphy, C. L., Bonin, E. M., ...
20 Brown, J. S. L. (2014). One-day cognitive-behavioural therapy self-confidence workshops for people with depression: Randomised controlled trial. *British Journal of Psychiatry*, 204, 222-233.
- 23 House, J., Marasali, P., Lister, M., & Brown, J. (2017). Male views on help-seeking for depression: a Q methodology study. *Psychol Psychother Theory Res Pract* 2044-8341.
- 25 Howe, D. (1995). Attachment theory for social work in practice. Basingstoke: Palgrave.
- 26 Hoyberg OJ, Maragakis B, Mullin J, Norum D, Stordall E, Ekdahl P, Ose E, Moksnes KM, Sennef C. A double-blind multicentre comparison of mirtazapine and amitriptyline in elderly depressed patients. *Acta Psychiatrica Scandinavica* 1996; 93(3):184-190.
- 29 HSCIC. (2015). Annual report on the use of IAPT services 2014-15. London.
- 30 HSCIC. (2015). Improving access to psychological therapies (IAPT). London.
- 31 HSCIC. (2016). Annual report on the use of IAPT services 2015-16. London.
- 32 Hsu J-W, Su T-P, Huang C-Y, Chen Y-S, Chou Y-H. Faster onset of antidepressant effects of citalopram compared with sertraline in drug-naïve first-episode major depressive disorder in a Chinese population: A 6-week double-blind, randomized comparative study. *Journal of Clinical Psychopharmacology* 2011; 31(5):577-81.
- 36 Huffman, J. C., Celano, C. M., Beach, S. R., Motiwala, S. R., & Januzzi, J. L. (2013). Depression and cardiac disease: epidemiology, mechanisms, and diagnosis. *Cardiovasc Psychiatry Neurol*, 2013, 695925.
- 39 Huijbers, M. J., Spinhoven, P., Spijker, J., Ruhe, H. G., Schaik, D. J. F., Oppen, P., ...
40 Speckens, A. E. M. (2015). Adding mindfulness-based cognitive therapy to maintenance antidepressant medication for prevention of relapse/recurrence in major depressive disorder: Randomised controlled trial. *Journal of Affective Disorders*, 187, 54-61.
- 43 Huijbers, M. J., Spinhoven, P., Spijker, J., Ruhé, H. G., van Schaik, D. J., van Oppen, P., ...
44 van der Wilt, G. J. (2016). Discontinuation of antidepressant medication after mindfulness-

- 1 based cognitive therapy for recurrent depression: randomised controlled non-inferiority trial.
- 2 The British Journal of Psychiatry, bjp. bp. 115.168971.
- 3 Huijbregts, K. M., Jong, F. J., Marwijk, H. W., Beekman, A. T., Adèr, H. J., Hakkaart-van
- 4 Roijen, L., . . . Feltz-Cornelis, C. M. (2013). A target-driven collaborative care model for Major
- 5 Depressive Disorder is effective in primary care in the Netherlands. A randomized clinical
- 6 trial from the depression initiative. *Journal of Affective Disorders*, 146(3), 328-337.
- 7 Hunter SB, Watkins KE, Hepner KA, Paddock SM, Ewing BA, Osilla KC, Perry S. Treating
- 8 depression and substance use: A randomized controlled trial. *Journal of substance abuse*
- 9 treatment. 2012 Sep 30; 43(2):137-51.
- 10 Husain N, Chaudhry N, Fatima B, Husain M, Amin R, Chaudhry IB, Rahman RU, Tomenson
- 11 B, Jafri F, Naeem F, Creed F. Antidepressant and group psychosocial treatment for
- 12 depression: a rater blind exploratory RCT from a low income country. *Behavioural and*
- 13 cognitive psychotherapy. 2014 Nov; 42(6):693-705.
- 14 Husain, N., Waheed, W., Tomenson, B., & Creed, F. (2007). The validation of personal
- 15 health questionnaire amongst people of Pakistani family origin living in the United Kingdom. *J*
- 16 *Affect Disord*, 97(1-3), 261-264.
- 17 Ince Burcin, U., et al. (2013) Internet-based, culturally sensitive, problem-solving therapy for
- 18 Turkish migrants with depression: Randomized controlled trial. *Journal of Medical Internet*
- 19 *Research* 15, 248-262.
- 20 Ingram, A., Saling, M. M., & Schweitzer, I. (2008). Cognitive side effects of brief pulse
- 21 electroconvulsive therapy: a review. *J ECT*, 24(1), 3-9.
- 22 Institute of Medicine (US). (2001). Neurological, psychiatric, and developmental disorders:
- 23 Meeting the challenge in the developing world. Washington (DC): Institute of Medicine (US),
- 24 Board on Global Health & Committee on Nervous System Disorders in Developing Countries.
- 25 Interian A, Lewis-Fernández R, Gara MA, Escobar JI. A randomized-controlled trial of an
- 26 intervention to improve antidepressant adherence among Latinos with depression.
- 27 *Depression and anxiety*. 2013 Jul 1; 30(7):688-96.
- 28 Jacobson, N. S., Dobson, K., Fruzzetti, A. E., Schmaling, K. B., & Salusky, S. (1991). Marital
- 29 therapy as a treatment for depression. *Journal of Consulting and Clinical Psychology*, 59(4),
- 30 547.
- 31 Jacobson, N.S., Dobson, K.S., Truax, P.A., et al. (1996) A component analysis of cognitive-
- 32 behavioural treatment for depression. *Journal of Consulting and Clinical Psychology*, 64 (2),
- 33 295-304.
- 34 Jacobson, N. S., Fruzzetti, A. E., Dobson, K., Whisman, M., & Hops, H. (1993). Couple
- 35 therapy as a treatment for depression: II. The effects of relationship quality and therapy on
- 36 depressive relapse. *J Consult Clin Psychol*, 61(3), 516-519.
- 37 Jacobson, N. S., Martell, C. R., & Dimidjian, S. (2001). Behavioral activation treatment for
- 38 depression: Returning to contextual roots. *Clinical Psychology: Science and Practice*, 8(3),
- 39 255-270.
- 40 Jadad, A. R., Moore, R. A., Carroll, D., Jenkinson, C., Reynolds, D. J., Gavaghan, D. J., &
- 41 McQuay, H. J. (1996). Assessing the quality of reports of randomized clinical trials: is
- 42 blinding necessary? *Control Clin Trials*, 17(1), 1-12.
- 43 Jakobsen JC, Gluud C, Kongerslev M, Larsen KA, Sorensen P, Winkel P, et al. Third-wave
- 44 cognitive therapy versus mentalisation-based treatment for major depressive disorder: a
- 45 randomised clinical trial. *BMJ Open*. 2014; 4(8):e004903.

- 1 Jakobsen, J. C., Katakam, K. K., Schou, A., Hellmuth, S. G., Stallknecht, S. E., Leth-Moller, K., ... Gluud, C. (2017). Selective serotonin reuptake inhibitors versus placebo in patients with major depressive disorder. A systematic review with meta-analysis and Trial Sequential Analysis. *BMC Psychiatry*, 17(1), 58.
- 5 Jamison, C., & Scogin, F. (1995). The outcome of cognitive bibliotherapy with depressed adults. *Journal of Consulting and Clinical Psychology*, 63(4), 644.
- 7 Jansen, J., & Naci, H. (2013). Is network meta-analysis as valid as standard pairwise meta-analysis? It all depends on the distribution of effect modifiers. *BMC Medicine*, 11:159.
- 9 Jarrett, R. B., Kraft, D., Doyle, J., Foster, B. M., Eaves, G. G., & Silver, P. C. (2001). Preventing recurrent depression using cognitive therapy with and without a continuation phase: a randomized clinical trial. *Arch Gen Psychiatry*, 58(4), 381-388.
- 12 Jarrett, R. B., Minhajuddin, A., Gershenson, H., Friedman, E. S., & Thase, M. E. (2013). Preventing depressive relapse and recurrence in higher-risk cognitive therapy responders: A randomized trial of continuation phase cognitive therapy, fluoxetine, or matched pill placebo. *JAMA Psychiatry*, 70(11), 1152-1160.
- 16 Jazayeri, S., Tehrani-Doost, M., Keshavarz, S. A., Hosseini, M., Djazayery, A., Amini, H., ... Peet, M. (2008). Comparison of therapeutic effects of omega-3 fatty acid eicosapentaenoic acid and fluoxetine, separately and in combination, in major depressive disorder. *Australian & New Zealand Journal of Psychiatry*, 42(3), 192-198.
- 20 Jeong H, Yim HW, Jo SJ, Nam B, Kwon SM, Choi JY, Yang SK. The effects of care management on depression treatment in a psychiatric clinic: a randomized controlled trial. *International journal of geriatric psychiatry*. 2013 Oct 1; 28(10):1023-30.
- 23 Jeong, J. H., et al. (2015). "Efficacy and safety of generic escitalopram (Lexacure) in patients with major depressive disorder: A 6-week multicenter, randomized, rater-blinded, escitalopram-comparative, non-inferiority study." *Neuropsychiatric disease and treatment* 11: 2557-2564.
- 27 Joffe, R. T., Moul, D. E., Lam, R. W., Levitt, A. J., Teicher, M. H., Lebegue, B., . . . Murray, M. G. (1993). Light visor treatment for seasonal affective disorder: A multicenter study. *Psychiatry research*, 46(1), 29-39.
- 30 Joffe, R. T., Sokolov, S. T., & Levitt, A. J. (2006). Lithium and triiodothyronine augmentation of antidepressants. *The Canadian Journal of Psychiatry*, 51(12), 791-793.
- 32 Johansson, R., et al. (2012). Psychodynamic guided self-help for adult depression through the internet: a randomised controlled trial. *PloS One* 7,
- 34 Johansson, R., et al. (2012). Tailored vs. standardized internet-based cognitive behavior therapy for depression and comorbid symptoms: a randomized controlled trial. *PloS One* 7
- 36 Johnson, J., Horwath, E., & Weissman, M. M. (1991). The validity of major depression with psychotic features based on a community study. *Arch Gen Psychiatry*, 48(12), 1075-1081.
- 38 Johnson, S., Nolan, F., Pilling, S., Sandor, A., Hoult, J., McKenzie, N., ... Bebbington, P. (2005). Randomised controlled trial of acute mental health care by a crisis resolution team: the north Islington crisis study. *BMJ*, 331(7517), 599.
- 41 Johnston, O., Kumar, S., Kendall, K., Peveler, R., Gabbay, J., & Kendrick, T. (2007). Qualitative study of depression management in primary care: GP and patient goals, and the value of listening. *Br J Gen Pract*, 57(544), 872-879.
- 44 Joling, K. J., Hout, H. P., van't Veer-Tazelaar, P. J., Horst, H. E., Cuijpers, P., Ven, P. M., & Marwijk, H. W. (2011). How effective is bibliotherapy for very old adults with subthreshold

- 1 depression? A randomized controlled trial. American Journal of Geriatric Psychiatry, 19(3),
2 256-265.
- 3 Jones, E., & Asen, E. (1999). Systemic couple therapy and depression. London: Karnac.
- 4 Jordan J, Carter JD, McIntosh VV, Fernando K, Frampton CM, Porter RJ, et al.
- 5 Metacognitive therapy versus cognitive behavioural therapy for depression: a randomized
6 pilot study. The Australian and New Zealand journal of psychiatry. 2014; 48(10):932-43.
- 7 Joy, C., Adams, C., & Rice, K. (2003). Crisis intervention for people with severe mental
8 illness (Cochrane Review). The Cochrane Library, 1.
- 9 Judd, L. L., Akiskal, H. S., Maser, J. D., Zeller, P. J., Endicott, J., Coryell, W., ... Keller, M. B.
10 (1998). Major depressive disorder: a prospective study of residual subthreshold depressive
11 symptoms as predictor of rapid relapse. J Affect Disord, 50(2-3), 97-108.
- 12 Judd, L. L., Akiskal, H. S., Maser, J. D., Zeller, P. J., Endicott, J., Coryell, W., ... Keller, M. B.
13 (1998). A prospective 12-year study of subsyndromal and syndromal depressive symptoms
14 in unipolar major depressive disorders. Arch Gen Psychiatry, 55(8), 694-700.
- 15 Judd FK, Moore K, Norman TR, Burrows GD, Gupta RK, Parker G. A multicentre double
16 blind trial of fluoxetine versus amitriptyline in the treatment of depressive illness. Australian &
17 New Zealand Journal of Psychiatry 1993 Mar; 27(1):49-55.
- 18 Kabat-Zinn, J., & University of Massachusetts Medical Center/Worcester. Stress Reduction
19 Clinic. (1990). Full catastrophe living: using the wisdom of your body and mind to face stress,
20 pain, and illness. New York, N.Y.: Delacorte Press.
- 21 Kalb, R., Trautmann-Sponsel, R. D., & Kieser, M. (2001). Efficacy and tolerability of
22 hypericum extract WS 5572 versus placebo in mildly to moderately depressed patients. A
23 randomized double-blind multicenter clinical trial. Pharmacopsychiatry, 34(3), 96-103.
- 24 Kallert, T. W., Priebe, S., McCabe, R., Kiejna, A., Rymaszewska, J., Nawka, P., ... Koch, R.
25 (2007). Are Day Hospitals Effective for Acutely Ill Psychiatric Patients? A European
26 Multicenter Randomized Controlled Trial. Journal of Clinical Psychiatry, 68(2), 278-287.
- 27 Kalenthaler, E., Brazier, J., De Nigris, E., Tumur, I., Ferriter, M., Beverley, C., ... Sutcliffe, P.
28 (2006). Computerised cognitive behaviour therapy for depression and anxiety update: a
29 systematic review and economic evaluation. Health Technology Assessment, 10(33), 1-186.
- 30 Kamijima K, Higuchi T, Ishigooka J, Ohmori T, Ozaki N, Kanba S, Kinoshita T, Koyama T,
31 ADMIRE Study Group. Aripiprazole augmentation to antidepressant therapy in Japanese
32 patients with major depressive disorder: a randomized, double-blind, placebo-controlled
33 study (ADMIRE study). Journal of affective disorders. 2013 Dec 31; 151(3):899-905.
- 34 Kanai, T., Takeuchi, H., Furukawa, T. A., Yoshimura, R., Imaizumi, T., Kitamura, T., &
35 Takahashi, K. (2003). Time to recurrence after recovery from major depressive episodes and
36 its predictors. Psychol Med, 33(5), 839-845.
- 37 Kanter JW, Santiago-Rivera AL, Santos MM, Nagy G, López M, Hurtado GD, West P. A
38 randomized hybrid efficacy and effectiveness trial of behavioral activation for Latinos with
39 depression. Behavior therapy. 2015 Mar 31; 46(2):177-92.
- 40 Kantor, D., McNevin, S., Leichner, P., Harper, D., & Krenn, M. (1986). The benefit of lithium
41 carbonate adjunct in refractory depression—fact or fiction? The Canadian Journal of
42 Psychiatry, 31(5), 416-418.
- 43 Kasper, S., de Swart, H., & Andersen, H. F. (2005). Escitalopram in the treatment of
44 depressed elderly patients. The American Journal of Geriatric Psychiatry, 13(10), 884-891.

- 1 Kasper, S., Wehr, T. A., Bartko, J. J., Gaist, P. A., & Rosenthal, N. E. (1989).
- 2 Epidemiological findings of seasonal changes in mood and behavior. A telephone survey of 3 Montgomery County, Maryland. *Arch Gen Psychiatry*, 46(9), 823-833.
- 4 Katon, W., Robinson, P., Von Korff, M., Lin, E., Bush, T., Ludman, E., Simon, G., Walker, E. 5 (1996). A multifaceted intervention to improve treatment of depression in primary care. 6 Archives of General Psychiatry, 53, 924.
- 7 Katon, W., von Korff, M., Lin, E., Bush, T., & Ormel, J. (1992). Adequacy and duration of 8 antidepressant treatment in primary care. *Med Care*, 30(1), 67-76.
- 9 Katon, W., Von Korff, M., Lin, E., & Simon, G. (2001). Rethinking practitioner roles in chronic 10 illness: the specialist, primary care physician, and the practice nurse. *Gen Hosp Psychiatry*, 11 23(3), 138-144.
- 12 Katon, W., Von Korff, M., Lin, E., Simon, G., Walker, E., Unutzer, J., ... Ludman, E. (1999). 13 Stepped collaborative care for primary care patients with persistent symptoms of depression: 14 a randomized trial. *Arch Gen Psychiatry*, 56(12), 1109-1115.
- 15 Katon, W., Von Korff, M., Lin, E., Walker, E., Simon, G., Bush, T., Robinson, P. & Russo J. 16 (1995). Collaborative management to achieve treatment guidelines: impact on depression in 17 primary care. *Journal of the American Medical Association*, 273, 1026-1031.
- 18 Katona, C., Abou-Saleh, M. T., Harrison, D. A., Nairac, B. A., Edwards, D., Lock, T., ... 19 Robertson, M. M. (1995). Placebo-controlled trial of lithium augmentation of fluoxetine and 20 lofepramine. *The British Journal of Psychiatry*, 166(1), 80-86.
- 21 Katz, I. R., Simpson, G. M., Curlik, S. M., & Parmelee, P. A. (1990). Pharmacologic treatment 22 of major depression for elderly patients in residential care settings. *Journal of Clinical 23 Psychiatry*.
- 24 Katzelnick, D., Simon, G., Pearson, S., Manning, W., Helstad, C., Henk, H., Cole, S.M., Lin 25 E.H., Taylor, L.H., Kobak, K.A. (2000). Randomized trial of a depression management 26 program in high utilizers of medical care. *Archives of Family Medicine*, 9, 345–351.
- 27 Kay-Lambkin FJ, Baker AL, Lewin TJ, Carr VJ. Computer-based psychological treatment for 28 comorbid depression and problematic alcohol and/or cannabis use: a randomized controlled 29 trial of clinical efficacy. *Addiction*. 2009 Mar 1; 104(3):378-88.
- 30 Keegan D, Bowen RC, Blackshaw S, Saleh S, Dayal N, Remillard F, Shrikhande S, Cebrian 31 Perez S, Boulton A. A comparison of fluoxetine and amitriptyline in the treatment of major 32 depression. *International Clinical Psychopharmacology* 1991 summer; 6(2):117-24.
- 33 Kehle, S. M., Greer, N., Rutks, I., & Wilt, T. (2011). Interventions to improve veterans' access 34 to care: a systematic review of the literature. *Journal of General Internal Medicine*, 26 Suppl 35 2, 689-696. Keitner, G. I., Garlow, S. J., Ryan, C. E., Ninan, P. T., Solomon, D. A., 36 Nemeroff, C. B., & Keller, M. B. (2009). A randomized, placebo-controlled trial of risperidone 37 augmentation for patients with difficult-to-treat unipolar, non-psychotic major depression. 38 *Journal of psychiatric research*, 43(3), 205-214.
- 39 Keller, M. (1985). Chronic and recurrent affective disorders: incidence, course and 40 influencing factors. In D. Kemali & G. Racagni (Eds.), *Chronic treatments in neuropsychiatry* 41 (pp. xv, 219 p.). New York: Raven Press.
- 42 Keller, M. B., & Boland, R. J. (1998). Implications of failing to achieve successful long-term 43 maintenance treatment of recurrent unipolar major depression. *Biol Psychiatry*, 44(5), 348- 44 360.
- 45 Keller, M. B., Harrison, W., Fawcett, J. A., Gelenberg, A., Hirschfeld, R. M., Klein, D. et al. 46 (1995). Treatment of chronic depression with sertraline or imipramine: preliminary blinded

- 1 response rates and high rates of undertreatment in the community. *Psychopharmacol Bull*,
2 31(2), 205-212.
- 3 Keller, M. B., Klerman, G. L., Lavori, P. W., Coryell, W., Endicott, J., & Taylor, J. (1984).
- 4 Long-term outcome of episodes of major depression. Clinical and public health significance.
5 *JAMA*, 252(6), 788-792.
- 6 Keller, M. B., Lavori, P. W., Mueller, T. I., Endicott, J., Coryell, W., Hirschfeld, R. M., & Shea,
7 T. (1992). Time to recovery, chronicity, and levels of psychopathology in major depression. A
8 5-year prospective follow-up of 431 subjects. *Arch Gen Psychiatry*, 49(10), 809-816.
- 9 Keller, M. B., Lavori, P. W., Rice, J., Coryell, W., & Hirschfeld, R. M. (1986). The persistent
10 risk of chronicity in recurrent episodes of nonbipolar major depressive disorder: a prospective
11 follow-up. *Am J Psychiatry*, 143(1), 24-28.
- 12 Keller, M. B., McCullough, J. P., Klein, D. N., Arnow, B., Dunner, D. L., Gelenberg, A. J., ...
13 Zajecka, J. (2000). A comparison of nefazodone, the cognitive behavioral-analysis system of
14 psychotherapy, and their combination for the treatment of chronic depression. *N Engl J Med*,
15 342(20), 1462-1470.
- 16 Keller, M. B., & Shapiro, R. W. (1981). Major depressive disorder. Initial results from a one-
17 year prospective naturalistic follow-up study. *J Nerv Ment Dis*, 169(12), 761-768.
- 18 Keller, M. B., Trivedi, M. H., Thase, M. E., Shelton, R. C., Kornstein, S. G., Nemeroff, C. B.,
19 Dunner, D. L. (2007). The Prevention of Recurrent Episodes of Depression with Venlafaxine
20 for Two Years (PREVENT) Study: Outcomes from the 2-year and combined maintenance
21 phases. *The Journal of Clinical Psychiatry*, 68(8), 1246-1256.
- 22 Kellner, C. H., Knapp, R. G., Petrides, G., Rummans, T. A., Husain, M. M., Rasmussen, K.,
23 Smith, G. (2006). Continuation electroconvulsive therapy vs pharmacotherapy for relapse
24 prevention in major depression: a multisite study from the Consortium for Research in
25 Electroconvulsive Therapy (CORE). *Archives of General Psychiatry*, 63(12), 1337-1344.
- 26 Kendler, K. S., Gardner, C. O., Neale, M. C., & Prescott, C. A. (2001). Genetic risk factors for
27 major depression in men and women: similar or different heritabilities and same or partly
28 distinct genes? *Psychological Medicine*, 31(04).
- 29 Kendler, K. S., & Prescott, C. A. (1999). A population-based twin study of lifetime major
30 depression in men and women. *Archives of General Psychiatry*, 56(1), 39.
- 31 Kendrick, T., Chatwin, J., Dowrick, C., Tylee, A., Morriss, R., Peveler, R., ... Moore, M.
32 (2009). Randomised controlled trial to determine the clinical effectiveness and cost-
33 effectiveness of selective serotonin reuptake inhibitors plus supportive care, versus
34 supportive care alone, for mild to moderate depression with somatic symptoms in primary
35 care: the THREAD (THREshold for AntiDepressant response) study. *Health Technology
36 Assessment*, 13(22), i-182.
- 37 Kendrick, T., Dowrick, C., McBride, A., Howe, A., Clarke, P., Maisey, S., ... Smith, P. W.
38 (2009). Management of depression in UK general practice in relation to scores on depression
39 severity questionnaires: analysis of medical record data. *BMJ*, 338(mar19 1), b750-b750.
- 40 Kendrick, T., King, F., Albertella, L., & Smith, P. W. (2005). GP treatment decisions for
41 patients with depression: an observational study. *Br J Gen Pract*, 55(513), 280-286.
- 42 Kendrick, T., Peveler, R., Longworth, L., Baldwin, D., Moore, M., Chatwin, J., ... Smith, H.
43 (2006). Cost-effectiveness and cost-utility of tricyclic antidepressants, selective serotonin
44 reuptake inhibitors and lofepramine. *The British Journal of Psychiatry*, 188(4), 337-345.
- 45 Kendrick, T., Simons, L., Mynors-Wallis, L., Gray, A., Lathlean, J., Pickering, R., ...
46 Thompson, C. (2005). A trial of problem-solving by community mental health nurses for

- 1 anxiety, depression and life difficulties among general practice patients. The CPN-GP study.
- 2 Health Technology Assessment, 9(37), 1-104.
- 3 Kendrick, T., Simons, L., Mynors-Wallis, L., Gray, A., Lathlean, J., Pickering, R., ...
- 4 Thompson, C. (2006). Cost-effectiveness of referral for generic care or problem-solving
- 5 treatment from community mental health nurses, compared with usual general practitioner
- 6 care for common mental disorders. The British Journal of Psychiatry, 189(1), 50-59.
- 7 Kendrick, T., Stevens, L., Bryant, A., Goddard, J., Stevens, A., Raftery, J., & Thompson, C.
- 8 (2001). Hampshire depression project: changes in the process of care and cost
- 9 consequences. Br J Gen Pract, 51(472), 911-913.
- 10 Kendrick, T., Stuart, B., Newell, C., Geraghty, A. W. A., & Moore, M. (2015). Changes in
- 11 rates of recorded depression in English primary care 2003–2013: Time trend analyses of
- 12 effects of the economic recession, and the GP contract quality outcomes framework (QOF).
- 13 Journal of Affective Disorders, 180, 68-78.
- 14 Kennedy, N., & Foy, K. (2005). The impact of residual symptoms on outcome of major
- 15 depression. Current Psychiatry Reports, 7(6), 441-446.
- 16 Kennedy, N., Abbott, R., & Paykel, E. S. (2003). Remission and recurrence of depression in
- 17 the maintenance era: long-term outcome in a Cambridge cohort. Psychol Med, 33(5), 827-
- 18 838.
- 19 Kennedy, S. H., Konarski, J. Z., Segal, Z. V., Lau, M. A., Bieling, P. J., McIntyre, R. S. &
- 20 Mayberg, H. S. (2007). Differences in brain glucose metabolism between responders to CBT
- 21 and venlafaxine in a 16-week randomized controlled trial. Am J Psychiatry 164, 778-88.
- 22 Kerling A, Tegtbjørn U, Gützlaff E, Kück M, Borchert L, Ates Z, von Bohlen A, Frieling H, Hüper
- 23 K, Hartung D, Schweiger U. Effects of adjunctive exercise on physiological and psychological
- 24 parameters in depression: a randomized pilot trial. Journal of affective disorders. 2015 May
- 25 15; 177:1-6.
- 26 Kessing, L. V. (2007). Epidemiology of subtypes of depression. Acta Psychiatrica
- 27 Scandinavica, 115(s433), 85-89.
- 28 Kessing, L. V., & Andersen, P. K. (1999). The effect of episodes on recurrence in affective
- 29 disorder: a case register study. J Affect Disord, 53(3), 225-231.
- 30 Kessler, D., Bennewith, O., Lewis, G., & Sharp, D. (2002). Detection of depression and
- 31 anxiety in primary care: follow up study. BMJ, 325(7371), 1016-1017.
- 32 Kessler, D., Lewis, G., Kaur, S., Wiles, N., King, M., Weich, S., ... Peters, T. J. (2009).
- 33 Therapist-delivered Internet psychotherapy for depression in primary care: a randomised
- 34 controlled trial. Lancet, 374(9690), 628-634.
- 35 Kessler, R. C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., ... Wang, P.
- 36 S. (2003). The epidemiology of major depressive disorder: results from the National
- 37 Comorbidity Survey Replication (NCS-R). JAMA, 289(23), 3095.
- 38 Kessler, R. C., & Bromet, E. J. (2013). The epidemiology of depression across cultures.
- 39 Annu Rev Public Health, 34, 119-138.
- 40 Kessler, R. C., Brown, R. L., & Broman, C. L. (1981). Sex differences in psychiatric help-
- 41 seeking: evidence from four large-scale surveys. J Health Soc Behav, 22(1), 49-64.
- 42 Kessler, R. C., Demler, O., Frank, R. G., Olfson, M., Pincus, H. A., Walters, E. E., ...
- 43 Zaslavsky, A. M. (2005). Prevalence and treatment of mental disorders, 1990 to 2003. New
- 44 England Journal of Medicine, 352(24), 2515-2523.

- 1 Kessler, R. C., Foster, C. L., Saunders, W. B., & Stang, P. E. (1995). Social consequences of psychiatric disorders, I: Educational attainment. *Am J Psychiatry*, 152(7), 1026-1032.
- 3 Kessler, R. C., Gruber, M., Hettema, J. M., Hwang, I., Sampson, N., & Yonkers, K. A. (2008). Co-morbid major depression and generalized anxiety disorders in the National Comorbidity Survey follow-up. *Psychol Med*, 38(3), 365-374.
- 6 Kessler, R. C., Petukhova, M., Sampson, N. A., Zaslavsky, A. M., & Wittchen, H. U. (2012). Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the United States. *Int J Methods Psychiatr Res*, 21(3), 169-184.
- 9 Khan, N., Bower, P., & Rogers, A. (2007). Guided self-help in primary care mental health: meta-synthesis of qualitative studies of patient experience. *Br J Psychiatry*, 191, 206-211.
- 11 Khoo, A. L., Zhou, H. J., Teng, M., Lin, L., Zhao, Y. J., Soh, L. B., ... Gwee, K. P. (2015). Network meta-analysis and cost-effectiveness analysis of new generation antidepressants. *CNS Drugs*, 29(8), 695-712.
- 14 Kiloh, L. G., Andrews, G., & Neilson, M. (1988). The long-term outcome of depressive illness. *Br J Psychiatry*, 153, 752-757.
- 16 Kinser PA, Bourguignon C, Whaley D, Hauenstein E, Taylor AG. Feasibility, acceptability, and effects of gentle hatha yoga for women with major depression: findings from a randomized controlled mixed-methods study. *Archives of psychiatric nursing*. 2013 Jun 30; 27(3):137-47.
- 20 Kirmayer, L. J., Robbins, J. M., Dworkind, M., & Yaffe, M. J. (1993). Somatization and the recognition of depression and anxiety in primary care. *Am J Psychiatry*, 150(5), 734-741.
- 22 Kirsch, I., Deacon, B. J., Huedo-Medina, T. B., Scoboria, A., Moore, T. J., & Johnson, B. T. (2008). Initial severity and antidepressant benefits: A meta-analysis of data submitted to the Food and Drug Administration. *PLoS Medicine*, 5(2), e45.
- 25 Kisely, S., Gater, R., & Goldberg, D. P. (1995). Results from the Manchester Centre. In T. B. Üstün & N. Sartorius (Eds.), *Mental illness in general health care: an international study* (pp. xii, 398 p.). Chichester; New York: Wiley.
- 28 Kishimoto, A., Mizukawa, R., Matsuzaki, F., Hazama, H., Kamase, H., Tanaka, K., & Kunimoto, N. (1994). Prophylactic effect of mianserin on recurrent depression. *Acta Psychiatrica Scandinavica*, 89(1), 46-51.
- 31 Kivi M, Eriksson MC, Hange D, Petersson EL, Vernmark K, Johansson B, Björkelund C. Internet-based therapy for mild to moderate depression in Swedish primary care: short term results from the PRIM-NET randomized controlled trial. *Cognitive behaviour therapy*. 2014 Oct 2; 43(4):289-98.
- 35 Klein, D. N., Leon, A. C., Li, C., D'Zurilla, T. J., Black, S. R., Vivian, D., ... Kocsis, J. H. (2011). Social problem solving and depressive symptoms over time: a randomized clinical trial of cognitive-behavioral analysis system of psychotherapy, brief supportive psychotherapy, and pharmacotherapy. *Journal of Consulting and Clinical Psychology*, 79(3), 342-352.
- 40 Klerman, G. L. (1984). *Interpersonal psychotherapy of depression*. New York: Basic Books.
- 41 Klug G, Hermann G, Fuchs-Nieder B, Panzer M, Haider-Stipacek A, Zapotoczky HG, Priebe S. Effectiveness of home treatment for elderly people with depression: Randomised controlled trial. *British Journal of Psychiatry* 2010; 197:463-467.
- 44 Klysner, R., Bent-hansen, J., Hansen, H. L., Lunde, M., Pleidrup, E., Poulsen, D. L., ... Petersen, H. E. H. (2002). Efficacy of citalopram in the prevention of recurrent depression in

- 1 elderly patients: placebo-controlled study of maintenance therapy. *The British Journal of Psychiatry*, 181(1), 29-35.
- 3 Kocsis, J. H., Frances, A. J., Voss, C., Mann, J. J., Mason, B. J., & Sweeney, J. (1988).
- 4 Imipramine treatment for chronic depression. *Archives of General Psychiatry*, 45(3), 253-257.
- 5 Kocsis, J. H., Leon, A. C., Markowitz, J. C., Manber, R., Arnow, B., Klein, D. N., & Thase, M.
- 6 E. (2009). Patient preference as a moderator of outcome for chronic forms of major
- 7 depressive disorder treated with nefazodone, cognitive behavioral analysis system of
- 8 psychotherapy, or their combination. *The Journal of Clinical Psychiatry*, 70(3), 354-361.
- 9 Kocsis JH, Thase ME, Trivedi MH, Shelton RC, Kornstein SG, Nemeroff CB, Friedman ES,
- 10 Gelenberg AJ, Dunner DL, Hirschfeld RM, Rothschild AJ. Prevention of recurrent episodes of
- 11 depression with venlafaxine ER in a 1-year maintenance phase from the PREVENT Study.
- 12 *The Journal of clinical psychiatry*. 2007 Jul; 68(7):1014-23.
- 13 Koeser, L., Donisi, V., Goldberg, D. P., & McCrone, P. (2015). Modelling the cost-effectiveness of pharmacotherapy compared with cognitive-behavioural therapy and combination therapy for the treatment of moderate to severe depression in the UK.
- 14 Psychological Medicine, 45, 3019-3031.
- 17 Kohtala, A., Lappalainen, R., Savonen, L., Timo, E., & Tolvanen, A. (2015). A four-session acceptance and commitment therapy based intervention for depressive symptoms delivered by masters degree level psychology students: a preliminary study. *Behavioural and cognitive psychotherapy*, 43(3), 360-373.
- 21 Kool, S., Dekker, J., Duijzens, I. J., de Jonghe, F., & Puite, B. (2003). Efficacy of combined therapy and pharmacotherapy for depressed patients with or without personality disorders.
- 23 *Harv Rev Psychiatry*, 11(3), 133-141.
- 24 Kornstein SG, Dunner DL, Meyers AL, Whitmyer VG, Mallinckrodt CH, Wohlreich MM, Detke MJ, Hollandbeck MS, Greist JH. A randomized, double-blind study of increasing or maintaining duloxetine dose in patients without remission of major depressive disorder after initial duloxetine therapy. *The Journal of clinical psychiatry*. 2008 Sep; 69(9):1383-92.
- 28
- 29 Kornstein, S., Bose, A., Li, D., Saikali, K., Gandhi, C. (2006) Escitalopram Maintenance Treatment for Prevention of Recurrent Depression: A Randomized, Placebo-Controlled Trial, *Journal of Clinical Psychiatry*, 67, 1767-1775.
- 32 Korkeila, J., Vahtera, J., Nabi, H., Kivimaki, M., Korkeila, K., Sumanen, M., ... Koskenvuo, M.
- 33 (2010). Childhood adversities, adulthood life events and depression. *J Affect Disord*, 127(1-3), 130-138.
- 35 Kornstein, S. G., Bose, A., Li, D., Saikali, K. G., & Gandhi, C. (2006). Escitalopram maintenance treatment for prevention of recurrent depression: a randomized, placebo-controlled trial. *The Journal of Clinical Psychiatry*, 67(11), 1767-1775.
- 38 Kornstein SG, Dunner DL, Meyers AL, Whitmyer VG, Mallinckrodt CH, Wohlreich MM, Detke MJ, Hollandbeck MS, Greist JH. A randomized, double-blind study of increasing or maintaining duloxetine dose in patients without remission of major depressive disorder after initial duloxetine therapy. *The Journal of clinical psychiatry*. 2008 Sep; 69(9):1383-92.
- 42 Kouros, C. D., Papp, L. M., & Cummings, E. M. (2008). Interrelations and moderators of longitudinal links between marital satisfaction and depressive symptoms among couples in established relationships. *Journal of Family Psychology*, 22(5), 667-677.
- 45 Kovacs, M. (1996). Presentation and course of major depressive disorder during childhood and later years of the life span. *Journal of the American Academy of Child & Adolescent Psychiatry*, 35(6), 705-715.

- 1 Krahm, D., Bartels, S., Coakley, E., Oslin, D., Chen, H., McIntyre, J., Chung, H., Maxwell, J., Ware, J., Levkoff, S.E. (2006). PRISM-E: comparison of integrated care and enhanced specialty referral models in depression outcomes. *Psychiatric Services*, 57, 946-953.
- 4 Kroenke, K. (2003). Patients presenting with somatic complaints: epidemiology, psychiatric co-morbidity and management. *International Journal of Methods in Psychiatric Research* 12 (1) 34 -43.
- 7 Kroenke, K., Spitzer, R. L., & Williams, J. B. (2003). The Patient Health Questionnaire-2: validity of a two-item depression screener. *Medical care*, 41(11), 1284-1292.
- 9 Krogh J, Saltin B, Nordentoft M. The DEMO trial: a randomized, parallel-group, observer-blinded clinical trial of strength versus aerobic versus relaxation training for patients with mild to moderate depression. *The Journal of clinical psychiatry*. 2009 Jun 15; 70(6):790-800.
- 12 Krogh J. DEMO II: A randomized, parallel-group, observer-blinded clinical trial of aerobic exercise versus stretching exercise for patients with light to moderate depression [NCT00695552]. Available from: <https://clinicaltrials.gov/ct2/show/NCT00695552> [accessed 19.09.2017]
- 16 Kuhn, R. (1958). The treatment of depressive states with g 22355 (imipramine hydrochloride). *American Journal of Psychiatry*, 115(5), 459-464.
- 18 Kunzel, H. E., Ackl, N., Hatzinger, M., Held, K., Holsboer-Trachsler, E., Ising, M., ... Steiger, A. (2009). Outcome in delusional depression comparing trimipramine monotherapy with a combination of amitriptyline and haloperidol--a double-blind multicenter trial. *J Psychiatr Res*, 43(7), 702-710.
- 22 Kupfer, D. J. (1991). Long-term treatment of depression. *J Clin Psychiatry*, 52 Suppl, 28-34.
- 23 Kusalic M, Engelsmann F, Bradwejn J. Thyroid functioning during treatment for depression. *Journal of Psychiatry and Neuroscience*. 1993 Nov; 18(5):260.
- 25 Kuyken, W., Byford, S., Taylor, R. S., Watkins, E., Holden, E., White, K., ... Teasdale, J. D. (2008). Mindfulness-based cognitive therapy to prevent relapse in recurrent depression. *J Consult Clin Psychol*, 76(6), 966-978.
- 28 Kuyken, W., Hayes, R., Barrett, B., Byng, R., Dalgleish, T., Kessler, D., ... Byford, S. (2015). The effectiveness and cost-effectiveness of mindfulness-based cognitive therapy compared with maintenance antidepressant treatment in the prevention of depressive relapse/recurrence: results of a randomised controlled trial (the PREVENT study). *Health Technol Assess*, 19(73), 1-124.
- 33 Kwon SH. Wheel of Wellness Counseling in Community Dwelling, Korean Elders: A Randomized, Controlled Trial. *Journal of Korean Academy of Nursing*. 2015 Jun 1; 45(3):459-68.
- 36 Kyle CJ, Petersen HEH, Overo KF. Comparison of the tolerability and efficacy of citalopram and amitriptyline in elderly depressed patients treated in general practice. *Depression & Anxiety* 1998; 8:147-153.
- 39 Laakman, G., Faltermaier-Temizel, M., Bossert-Zaudig, S., Baghai, T., & Lorkowski, G. (1995). Treatment of depressive outpatients with lorazepam, alprazolam, amitriptyline and placebo. *Psychopharmacology (Berl)*, 120(1), 109-115.
- 42 Laakmann, G., Schule, C., Baghai, T., & Kieser, M. (1998). St. John's wort in mild to moderate depression: the relevance of hyperforin for the clinical efficacy. *Pharmacopsychiatry*, 31 Suppl 1, 54-59.

- 1 Labbate, L. A., Lafer, B., Thibault, A., & Sachs, G. S. (1994). Side effects induced by bright light treatment for seasonal affective disorder. *Journal of Clinical Psychiatry*. 5: 189–191.
- 2 Laidlaw, K. (2008). Post-stroke depression and CBT with older people. *Handbook of behavioral and cognitive therapies with older adults*, 233-248.
- 3 Laidlaw, K., Davidson, K., Toner, H., et al (2008) A randomised controlled trial of cognitive behaviour therapy vs treatment as usual in the treatment of mild to moderate late life depression. *International Journal of Geriatric Psychiatry*, 23, 843-850.
- 4 Lam, R. W., Levitt, A. J., Levitan, R. D., Enns, M. W., Morehouse, R., Michalak, E. E., & Tam, E. M. (2006). The Can-SAD study: a randomized controlled trial of the effectiveness of light therapy and fluoxetine in patients with winter seasonal affective disorder. *American Journal of Psychiatry*, 163(5), 805-812.
- 5 Lam RW, Parikh SV, Ramasubbu R, Michalak EE, Tam EM, Axler A, Yatham LN, Kennedy SH, Manjunath CV. Effects of combined pharmacotherapy and psychotherapy for improving work functioning in major depressive disorder. *The British Journal of Psychiatry*. 2013 Nov 1; 203(5):358-65.
- 6 Lamers, S. M., Bohlmeijer, E. T., Korte, J., & Westerhof, G. J. (2015). The efficacy of life-review as online-guided self-help for adults: a randomized trial. *The journals of gerontology. Series B, Psychological sciences and social sciences*, 70(1), 24-34.
- 7 Landis, S., Gaynes, B., Morrissey, J., Vinson, N., Ellis, A., & Domino, M. (2007). Generalist care managers for the treatment of depressed medicaid patients in North Carolina: A pilot study. *BMC Family Practice*, 8, 7.
- 8 Landreville P. Effects of cognitive bibliotherapy for depressed older adults with a disability. *Clinical Gerontologist* 17(4):35-55, 1997.
- 9 Lappalainen P, Granlund A, Siltanen S, Ahonen S, Vitikainen M, Tolvanen A, Lappalainen R. ACT Internet-based vs face-to-face? A randomized controlled trial of two ways to deliver Acceptance and Commitment Therapy for depressive symptoms: An 18-month follow-up. *Behaviour research and therapy*. 2014 Oct 31; 61:43-54.
- 10 Lau, M. A., Segal, Z. V., & Williams, J. M. (2004). Teasdale's differential activation hypothesis: implications for mechanisms of depressive relapse and suicidal behaviour. *Behav Res Ther*, 42(9), 1001-1017.
- 11 Lauritzen, L., Odgaard, K., Clemmesen, L., Lunde, M., Öhrström, J., Black, C., & Bech, P. (1996). Relapse prevention by means of paroxetine in ECT-treated patients with major depression: a comparison with imipramine and placebo in medium-term continuation therapy. *Acta Psychiatrica Scandinavica*, 94(4), 241-251.
- 12 Lavretsky, H., Alstein, L. L., Olmstead, R. E., Ercoli, L. M., Riparetti-Brown, M., Cyr, N. S., & Irwin, M. R. (2011). Complementary use of tai chi chih augments escitalopram treatment of geriatric depression: a randomized controlled trial. *The American Journal of Geriatric Psychiatry*, 19(10), 839-850.
- 13 Law, D. D., & Crane, D. R. (2000). The influence of marital and family therapy on health care utilization in a health-maintenance organization. *Journal of Marital and Family Therapy*, 26(3), 281-291.
- 14 Law, D. D., Crane, D. R., & Berge, J. M. (2003). The influence of individual, marital, and family therapy on high utilizers of health care. *Journal of Marital and Family Therapy*, 29(3), 353-363.

- 1 Lawrence, V., Banerjee, S., Bhugra, D., Sangha, K., Turner, S., & Murray, J. (2006). Coping with depression in later life: a qualitative study of help-seeking in three ethnic groups. *Psychol Med*, 36(10), 1375-1383.
- 4 Layard, R. (2006). The case for psychological treatment centres. *BMJ*, 332(7548), 1030-1032.
- 6 Leadholm, A. K., Rothschild, A. J., Nolen, W. A., Bech, P., Munk-Jorgensen, P., & Ostergaard, S. D. (2013). The treatment of psychotic depression: is there consensus among guidelines and psychiatrists? *J Affect Disord*, 145(2), 214-220.
- 9 Lecrubier, Y., Boyer, P., Turjanski, S., Rein, W., & Group, A. S. (1997). Amisulpride versus imipramine and placebo in dysthymia and major depression. *Journal of Affective Disorders*, 43(2), 95-103.
- 12 Lecrubier, Y., Clerc, G., Didi, R., & Kieser, M. (2002). Efficacy of St. John's wort extract WS(R) 5570 in major depression - a double-blind, placebo-controlled trial. *European Psychiatry: the Journal of the Association of European Psychiatrists*, 17(Suppl1), 67.
- 15 Lee, A. S., & Murray, R. M. (1988). The long-term outcome of Maudsley depressives. *Br J Psychiatry*, 153, 741-751.
- 17 Lee, M. S., Choi, J., Posadzki, P., & Ernst, E. (2012). Aromatherapy for health care: An overview of systematic reviews. *Maturitas*, 71(3), 257-260.
- 19 Leff, J., Vearnals, S., Wolff, G., Alexander, B., Chisholm, D., Everitt, B., ... Dayson, D. (2000). The London Depression Intervention Trial - randomised controlled trial of antidepressants v. Couple therapy in the treatment and maintenance of people with depression living with a partner: Clinical outcome and costs. *The British Journal of Psychiatry*, 177(2), 95-100.
- 24 Legrand FD. Effects of exercise on physical self-concept, global self-esteem, and depression in women of low socioeconomic status with elevated depressive symptoms. *J Sport Exerc Psychol*. 2014 Aug 1; 36(4):357-65.
- 27 Lehmann, H. E., Fenton, F. R., Deutsch, M., Feldman, S., & Engelsmann, F. (1988). An 11-year follow-up study of 110 depressed patients. *Acta Psychiatr Scand*, 78(1), 57-65.
- 29 Leith, L. M. (1994). Foundations of exercise and mental health. Morgantown, WV: Fitness Information Technology.
- 31 Leinonen, E. & Niemi, H. (2007). The influence of educational information on depressed outpatients treated with escitalopram: a semi-naturalistic study. *Nordic Journal of Psychiatry*, 61, 109-114.
- 34 Leinonen E, Skarstein J, Behnke K, et al., Efficacy and tolerability of mirtazapine versus citalopram: A double-blind, randomized study in patients with major depressive disorder. *International Clinical Psychopharmacology* 1999; 14(6):329-337.
- 37 Lemmens, L., Arntz, A., Peeters, F., Hollon, S., Roefs, A., & Huibers, M. (2015). Clinical effectiveness of cognitive therapy v. interpersonal psychotherapy for depression: results of a randomized controlled trial. *Psychological Medicine*, 45(10), 2095-2110.
- 40 Lemmens, L. H., DeRubeis, R. J., Arntz, A., Peeters, F. P., & Huibers, M. J. (2016). Sudden gains in Cognitive Therapy and Interpersonal Psychotherapy for adult depression. *Behaviour research and therapy*, 77, 170-176.
- 43 Lenox-Smith, A., Greenstreet, L., Burslem, K., & Knight, C. (2009). Cost effectiveness of venlafaxine compared with generic fluoxetine or generic amitriptyline in major depressive disorder in the UK. *Clinical Drug Investigation*, 29, 173-184.

- 1 Lenox-Smith, A. J., & Jiang, Q. (2008). Venlafaxine extended release versus citalopram in patients with depression unresponsive to a selective serotonin reuptake inhibitor. *International Clinical Psychopharmacology*, 23(3), 113-119.
- 4 Lenze EJ, Mulsant BH, Blumberger DM, Karp JF, Newcomer JW, Anderson SJ, Dew MA, Butters MA, Stack JA, Begley AE, Reynolds CF. Efficacy, safety, and tolerability of augmentation pharmacotherapy with aripiprazole for treatment-resistant depression in late life: a randomised, double-blind, placebo-controlled trial. *The Lancet*. 2015 Dec 18; 386(10011):2404-12.
- 9 Lepine, J. P., Gastpar, M., Mendlewicz, J., & Tylee, A. (1997). Depression in the community: the first pan-European study DEPRES (Depression Research in European Society). *Int Clin Psychopharmacol*, 12(1), 19-29.
- 12 Lépine, J.-P., Caillard, V., Bisserbe, J.-C., Troy, S., Hotton, J.-M., & Boyer, P. (2004). A randomized, placebo-controlled trial of sertraline for prophylactic treatment of highly recurrent major depressive disorder. *American Journal of Psychiatry*, 161(5), 836-842.
- 15 Lepola, U.M., Loft, H. & Reines, H. (2003) Escitalopram (10-20 mg/day) is effective and well tolerated in a placebo-controlled study in depression in primary care. *International Clinical Psychopharmacology*, 18, 211-217.
- 18 Lepore, S. J. (1997). Expressive writing moderates the relation between intrusive thoughts and depressive symptoms. *J Pers Soc Psychol*, 73(5), 1030-1037.
- 20 Levesque DA, Van Marter DF, Schneider RJ, Bauer MR, Goldberg DN, Prochaska JO, Prochaska JM. Randomized trial of a computer-tailored intervention for patients with depression. *American Journal of Health Promotion*. 2011 Nov; 26(2):77-89.
- 23 Levin, W., Campbell, D. R., McGovern, K. B., Gau, J. M., Kosty, D. B., Seeley, J. R., & Lewinsohn, P. M. (2011). A computer-assisted depression intervention in primary care. *Psychological Medicine*, 41(7), 1373-1383.
- 26 Levine S, Deo R, Mahadevan K. A comparative trial of a new antidepressant, fluoxetine. *International Clinical Psychopharmacology* 1989 Jan; 4 Suppl 1:41-5.
- 28 Levitt, A. J., Wesson, V. A., Joffe, R. T., Mauder, R. G., & King, E. F. (1996). A controlled comparison of light box and head-mounted units in the treatment of seasonal depression. *The Journal of Clinical Psychiatry*, 57(3), 105-110.
- 31 Lewinsohn, P. M. (1975). The behavioural study and treatment of depression. In M. Hersen, R. M. Eisler, & P. M. Miller (Eds.), *Progress in behavior modification* (pp. 19-64). New York; Newbury Park, CA: Academic Press; Sage Publications.
- 34 Lewinsohn, P. M. (1984). The Coping with depression course: a psychoeducational intervention for unipolar depression. Eugene, OR: Castalia Pub. Co.
- 36 Lewinsohn, P. M., & Arconad, M. (1981). Behavioral treatment of depression: A social learning approach. *Depression: Behavioral and directive intervention strategies*, 33-67.
- 38 Lewinsohn, P. M., Solomon, A., Seeley, J. R., & Zeiss, A. (2000). Clinical implications of "subthreshold" depressive symptoms. *Journal of Abnormal Psychology*, 109(2), 345-351.
- 40 Lewis, E., Marcus, S. C., Olfson, M., Druss, B. G., & Pincus, H. A. (2004). Patients' early discontinuation of antidepressant prescriptions. *Psychiatr Serv*, 55(5), 494.
- 42 Lewis-Fernandez R, Vermes D. Motivational Antidepressant Therapy for Hispanics [NCT00564278]. 2007. Available from: <https://clinicaltrials.gov/ct2/show/study/NCT00564278> [accessed 11.08.2017].

- 1 Lewy, A., Sack, R., Miller, L., & Hoban, T. (1987). Antidepressant and circadian phase-shifting effects of light. *Science*, 235(4786), 352-354.
- 3 Lexchin, J. (2003). Pharmaceutical industry sponsorship and research outcome and quality: systematic review. *BMJ*, 326(7400), 1167-1170.
- 5 Li X, Xing B, Yu E, Chen W, Wu H. The combined treatment of venlafaxine and quetiapine for treatment-resistant depression: a clinical study. *The Journal of neuropsychiatry and clinical neurosciences*. 2013 Apr; 25(2):157-60.
- 8 Liberman, R. P., & Eckman, T. (1981). Behavior therapy vs insight-oriented therapy for repeated suicide attempts. *Arch Gen Psychiatry*, 38(10), 1126-1130.
- 10 Licht, R. W., & Qvitzau, S. (2002). Without Title. *Psychopharmacology*, 161(2), 143-151.
- 11 Liebowitz, M., Lam, R. W., Lepola, U., Datto, C., Sweitzer, D., & Eriksson, H. (2010). Efficacy and tolerability of extended release quetiapine fumarate monotherapy as maintenance treatment of major depressive disorder: a randomized, placebo-controlled trial. *Depression and Anxiety*, 27(10), 964-976.
- 15 Lieverse R, Van Someren EJ, Nielen MM, Uitdehaag BM, Smit JH, Hoogendoijk WJ. Bright light treatment in elderly patients with nonseasonal major depressive disorder: a randomized placebo-controlled trial. *Archives of General Psychiatry*. 2011 Jan 3; 68(1):61-70.
- 18 Linde, K., & Mulrow, C. D. (2004). St John's wort for major depression (Cochrane Review). *Cochrane Database Syst Rev*. 2005(2):Cd000448..
- 20 Linde, K., Rücker, G., Schneider, A., & Kriston, L. (2016). Questionable assumptions hampered interpretation of a network meta-analysis of primary care depression treatments. *Journal of Clinical Epidemiology* 71, 86-96.
- 23 Lindert, J., Schouler-Ocak, M., Heinz, A., & Priebe, S. (2008). Mental health, health care utilisation of migrants in Europe. *European Psychiatry*, 23, 14-20.
- 25 Lingjærde, O., Reichborn-Kjennerud, T., Haggag, A., Gärtner, I., Narud, K., & Berg, E. (1993). Treatment of winter depression in Norway. *Acta Psychiatrica Scandinavica*, 88(5), 372-380.
- 28 Littlewood, E., Duarte, A., Hewitt, C., Knowles, S., Palmer, S., Walker, S., ... Bower, P. (2015). A randomised controlled trial of computerised cognitive behaviour therapy for the treatment of depression in primary care: the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REACT) trial. *Health technology assessment* (Winchester, England), 19(101), 1.
- 33 Liu, E.-H., Chen, W. L., Li, Y. H., Wang, C. H., Mok, T. J., & Huang, H. S. (2009). Exploring the efficacy of cognitive bibliotherapy and a potential mechanism of change in the treatment of depressive symptoms among the Chinese: A randomized controlled trial. *Cognitive Therapy and Research*, 33(5), 449-461.
- 37 Lobello, K., Reddy, S., Musgnung, J., Pedersen, R., & Ninan, P. T. (2010). Patient outcomes with education, drug therapy, and support: a study of venlafaxine ER-treated outpatients with major depressive disorder. *Psychopharmacology Bulletin*, 43(2), 28-44.
- 40 Losada, A., Márquez-González, M., Romero-Moreno, R., Mausbach, B. T., López, J., Fernández-Fernández, V., & Nogales-González, C. (2015). Cognitive-behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for dementia family caregivers with significant depressive symptoms: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology*, 83(4), 760.

- 1 Lovell, K., Bower, P., Richards, D., Barkham, M., Sibbald, B., Roberts, C., ... Hennessy, S. (2008). Developing guided self-help for depression using the Medical Research Council complex interventions framework: a description of the modelling phase and results of an exploratory randomised controlled trial. *BMC Psychiatry*, 8(1), 91.
- 5 Lu, G., & Ades, A. E. (2004). Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med*, 23(20), 3105-3124.
- 7 Lucas M, Asselin G, Mérette C, Poulin MJ, Dodin S. Ethyl-eicosapentaenoic acid for the treatment of psychological distress and depressive symptoms in middle-aged women: a double-blind, placebo-controlled, randomized clinical trial. *The American journal of clinical nutrition*. 2009 Feb 1; 89(2):641-51.
- 11 Ludman, E., Simon, G., Grothaus, L., Luce, C., Markley, D., & Schaefer, J. (2007). A pilot study of telephone care management and structured disease self-management groups for chronic depression. *Psychiatric Services*, 58, 1065-1072.
- 14 Lunn, D. J., Thomas, A., Best, N., & Spiegelhalter, D. (2000). WinBUGS-a Bayesian modelling framework: concepts, structure, and extensibility. *Statistics and computing*, 10(4), 325-337.
- 17 Luo H, Shen Y, Zhou D, Jia Y. A comparative study of the treatment of depression by electroacupuncture and amitriptyline. *Acupuncture*. 1990; 1(2):20-6.
- 19 Luty, S.E., Carter, J.D., McKenzie, J.M., et al. (2007). Randomised controlled trial of interpersonal psychotherapy and cognitive-behavioural therapy for depression. *British Journal of Psychiatry*, 190, 496-502.
- 22 Ly, K. H., et al. (2015). "Smartphone-supported versus full behavioural activation for depression: A randomised controlled trial." *PloS one* 10(5) (no pagination).
- 24 Ly KH, Trüschel A, Jarl L, Magnusson S, Windahl T, Johansson R, Carlbring P, Andersson G. Behavioural activation versus mindfulness-based guided self-help treatment administered through a smartphone application: a randomised controlled trial. *BMJ open*. 2014 Jan 1; 4(1):e003440.
- 28 Lyall, D. M., Inskip, H. M., Mackay, D., Deary, I. J., McIntosh, A. M., Hotopf, M., ... Smith, D. J. (2016). Low birth weight and features of neuroticism and mood disorder in 83 545 participants of the UK Biobank cohort. *British Journal of Psychiatry Open*, 2(1), 38-44.
- 31 Lydiard RB, Laird LK, Morton WA Jr, Steele TE, Kellner C, Laraia MT, Ballenger JC. Fluvoxamine, imipramine, and placebo in the treatment of depressed outpatients: effects on depression. *Psychopharmacology Bulletin* 1989; 25(1):68-70.
- 34 Lydiard, R. B., Stahl, S. M., Hertzman, M., & Harrison, W. M. (1997). A double-blind, placebo-controlled study comparing the effects of sertraline versus amitriptyline in the treatment of major depression. *The Journal of Clinical Psychiatry*.
- 37 Lynch DJ, Tamburrino M, Nagel R. Telephone counseling for patients with minor depression: preliminary findings in a family practice setting. *Journal of family practice*. 1997 Mar 1; 44(3):293-9.
- 40 Ma, L., & Li, Y. (2014). Investigation of the influence of antidepressant therapy on blood pressure and quality of life in elderly patients both with hypertension and depression. *Cardiology (Switzerland)*, 129, 103-104.
- 43 Ma, S.H., & Teasdale, J.D. (2004). Mindfulness-based cognitive therapy for depression: Replication and exploration of differential relapse prevention effects. *Journal of Consulting and Clinical Psychology*, 1, 31–40.

- 1 Macaskill, N. D., & Macaskill, A. (1996). Rational-emotive therapy plus pharmacotherapy versus pharmacotherapy alone in the treatment of high cognitive dysfunction depression. *Cognitive Therapy and Research*, 20(6), 575-592.
- 2
- 3
- 4 MacMillan, H. L., Patterson, C. J., Wathen, C. N., Feightner, J. W., Bessette, P., Elford, R. W., . . . Canadian Task Force on Preventive Health, C. (2005). Screening for depression in primary care: recommendation statement from the Canadian Task Force on Preventive Health Care. *CMAJ*, 172(1), 33-35.
- 5
- 6
- 7
- 8 MacPherson, H., Richmond, S., Bland, M., Brealey, S., Gabe, R., Hopton, A., ... Sculpher, M. (2013). Acupuncture and counselling for depression in primary care: a randomised controlled trial. *PLoS Medicine*, 10(9), e1001518.
- 9
- 10
- 11 Maina G, Forner F, Bogetto F. Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. *Psychotherapy and psychosomatics*. 2004 Dec 23; 74 (1):43-50.
- 12
- 13
- 14 Maina G, Rosso G, Crespi C, Bogetto F. Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. *Psychotherapy and psychosomatics*. 2007 Aug 16; 76 (5):298-305.
- 15
- 16
- 17 Maina G, Rosso G, Rigardetto S, Chiadò Piat S, Bogetto F. No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. *Psychotherapy and psychosomatics*. 2010 Jul 9; 79 (5):295-302.
- 18
- 19
- 20
- 21 Maginn, S., Boardman, A. P., Craig, T. K., Haddad, M., Heath, G., & Stott, J. (2004). The detection of psychological problems by General Practitioners--influence of ethnicity and other demographic variables. *Soc Psychiatry Psychiatr Epidemiol*, 39(6), 464-471.
- 22
- 23
- 24 Magnusson, A., & Partonen, T. (2005). The diagnosis, symptomatology, and epidemiology of seasonal affective disorder. *CNS Spectr*, 10(8), 625-634; quiz 621-614.
- 25
- 26 Mahmoud, R. A., Pandina, G. J., Turkoz, I., Kosik-Gonzalez, C., Canuso, C. M., Kujawa, M. J., & Gharabawi-Garibaldi, G. M. (2007). Risperidone for Treatment-Refractory Major Depressive DisorderA Randomized TrialRisperidone for Major Depressive Disorder. *Annals of Internal Medicine*, 147(9), 593-602.
- 27
- 28
- 29
- 30 Maj, M., Veltro, F., Pirozzi, R., Lobrace, S., & Magliano, L. (1992). Pattern of recurrence of illness after recovery from an episode of major depression: a prospective study. *Am J Psychiatry*, 149(6), 795-800.
- 31
- 32
- 33 Malhi, G. S., Parker, G. B., & Greenwood, J. (2005). Structural and functional models of depression: from sub-types to substrates. *Acta Psychiatrica Scandinavica*, 111(2), 94-105.
- 34
- 35 Malone, D., Marriott, S., Newton-Howes, G., Simmonds, S., & Tyrer, P. (2007). Community mental health teams (CMHTs) for people with severe mental illnesses and disordered personality. *Cochrane Database of Systematic Reviews* 2007, Issue 3.
- 36
- 37
- 38 Malouff JM, Lanyon RI, Schutte NS. Effectiveness of a brief group RET treatment for divorce-related dysphoria. *Journal of Rational-Emotive & Cognitive-Behavior Therapy*. 1988 Sep 1; 6(3):162-71.
- 39
- 40
- 41 Mancuso, C. A., Rincon, M., McCulloch, C. E., & Charlson, M. E. (2001). Self-efficacy, depressive symptoms, and patients' expectations predict outcomes in asthma. *Med Care*, 39(12), 1326-1338.
- 42
- 43
- 44 Manicavasgar, V., et al. (2011) Mindfulness-based cognitive therapy vs cognitive behaviour therapy as a treatment for non-melancholic depression. *Journal of Affective Disorders* 130, 138-144.
- 45
- 46

- 1 Mann, A., Blizzard, R., Murray, J., Smith, J., Botega, N., MacDonald, E., Wilkinson, G. (1998).
- 2 An evaluation of practice nurses working with general practitioners to treat people with depression. *British Journal of General Practice*, 48, 875.
- 3
- 4 Mann, R., Gilbody, S., & Richards, D. (2009). Putting the 'Q' in depression QALYs: A comparison of utility measurement using EQ-5D and SF-6D health related quality of life measures. *Social psychiatry and psychiatric epidemiology*, 44(7), 569-578.
- 5
- 6
- 7 Mann, T. (1996). Clinical Guidelines: Using clinical guidelines to improve patient care within the NHS. .
- 8
- 9 Marchesi C, Ceccherinelli A, Rossi A, Maggini C. Is anxious-agitated major depression responsive to fluoxetine? A double-blind comparison with amitriptyline. *Pharmacopsychiatry* 1998; 31:216-221.
- 10
- 11
- 12 Marcus, S. M., Kerber, K. B., Rush, A. J., Wisniewski, S. R., Nierenberg, A., Balasubramani, G., ... Trivedi, M. H. (2008). Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study. *Comprehensive Psychiatry*, 49(3), 238-246.
- 13
- 14
- 15
- 16 Markowitz, J. C., Kocsis, J. H., Bleiberg, K. L., Christos, P. J., & Sacks, M. (2005). A comparative trial of psychotherapy and pharmacotherapy for "pure" dysthymic patients. *Journal of Affective Disorders*, 89(1), 167-175.
- 17
- 18
- 19 Marshall, M., Crowther, R., Almaraz-Serrano, A., Creed, F., Sledge, W., Kluiter, H., ... Bond, G. (2001). Systematic reviews of the effectiveness of day care for people with severe mental disorders: (1) acute day hospital versus admission; (2) vocational rehabilitation; (3) day hospital versus outpatient care. *Health technology assessment (Winchester, England)*, 5(21), 1-75.
- 20
- 21
- 22
- 23
- 24 Marshall, M., Crowther, R., Almaraz-Serrano, A. M., & Tyrer, P. (2001). Day hospital versus out-patient care for psychiatric disorders. *Cochrane Database of Systematic Reviews*, 2.
- 25
- 26 Marshall, M., Crowther, R., Sledge, W. H., Rathbone, J., & Soares-Weiser, K. (2011). Day hospital versus admission for acute psychiatric disorders. *Cochrane Database of Systematic Reviews*, 12.
- 27
- 28
- 29 Marshall, M.B., Zuroff, D.C., McBride, C., & Bagby, R.M. (2008) Self-criticism predicts differential response to treatment for major depression. *Journal of Clinical Psychology*, 64 (3), 231-244.
- 30
- 31
- 32 Marteau, T. M. (1989). Psychological costs of screening. *BMJ*, 299(6698), 527.
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44

- 1 Mather AS, Rodriguez C, Guthrie MF, McHARG AM, Reid IC, McMURDO ME. Effects of exercise on depressive symptoms in older adults with poorly responsive depressive disorder. *The British Journal of Psychiatry*. 2002 May 1; 180 (5):411-5.
- 4 Mattisson, C., Bogren, M., Horstmann, V., Munk-Jorgensen, P., & Nettelbladt, P. (2007). The long-term course of depressive disorders in the Lundby Study. *Psychol Med*, 37(6), 883-891.
- 6 Mavridis, D., Giannatsi, M., Cipriani, A., & Salanti, G. (2015). A primer on network meta-analysis with emphasis on mental health. *Evid Based Ment Health*, 18(2), 40-46.
- 8 May, C., Allison, G., Chapple, A., Chew-Graham, C., Dixon, C., Gask, L., ... Roland, M. (2004). Framing the doctor-patient relationship in chronic illness: a comparative study of general practitioners' accounts. *Sociol Health Illn*, 26(2), 135-158.
- 11 McCann IL, Holmes DS. Influence of aerobic exercise on depression. *Journal of personality and social psychology*. 1984 May; 46(5):1142.
- 13 McCallum, P., & Meares, R. (1975). A controlled trial of maprotiline (Ludiomil) in depressed outpatients. *The Medical journal of Australia*, 2(10), 392-393.
- 15 McClay, C. A., et al. (2015). A community-based pilot randomised controlled study of life skills classes for individuals with low mood and depression. *BMC Psychiatry* 15: 17.
- 17 McClelland HA, Kerr TA., Stephens DA, Howell RW. The comparative antidepressant value of lofepramine and amitriptyline. Results of a controlled trial with comments on the scales used. *Acta Psychiatrica Scandinavica* 1979; 60:190-198.
- 20 McClure, D. J., Low, G. L., & Gent, M. (1973). Clomipramine HCL--a double-blind study of a new antidepressant drug. *Can Psychiatr Assoc J*, 18(5), 403-408.
- 22 McCrone, P., Dhanasiri, S., Patel, A., & Knapp, M. (2008). Paying the price: the cost of mental health care in England to 2026. .
- 24 McCrone, P., Knapp, M., Proudfoot, J., Ryden, C., Cavanagh, K., Shapiro, D. A., ... Mann, A. (2004). Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: randomised controlled trial. *The British Journal of Psychiatry*, 185(1), 55-62.
- 28 McCullough, J. P. (2000). Treatment for chronic depression: cognitive behavioral analysis system of psychotherapy (CBASP). New York: Guilford Press.
- 30 McCusker, J., Cole, M., Yaffe, M., Cappeliez, P., Dawes, M., Sewitch, M., Ciampi, A., Latimer, E. (2008). Project DIRECT: Pilot study of a collaborative intervention for depressed seniors. *Canadian Journal of Community Mental Health*, 27(2), 201-218.
- 33 McGrath, C. L., Kelley, M. E., Holtzheimer, P. E., Dunlop, B. W., Craighead, W. E., Franco, A. R., . . . Mayberg, H. S. (2013). Toward a neuroimaging treatment selection biomarker for major depressive disorder. *JAMA Psychiatry*, 70(8), 821-829.
- 36 McGrath, P. J., Stewart, J. W., Quitkin, F. M., Chen, Y., Alpert, J. E., Nierenberg, A. A., et al. (2006). Predictors of relapse in a prospective study of fluoxetine treatment of major depression. *American Journal of Psychiatry*, 163, 1542-1548.
- 39 McGrath, P. J., Stewart, J. W., Quitkin, F. M., Chen, Y., Alpert, J. E., Nierenberg, A. A., . . . Petkova, E. (2006). Predictors of relapse in a prospective study of fluoxetine treatment of major depression. *American Journal of Psychiatry*, 163(9), 1542-1548.
- 42 McHugh, R. K., Whitton, S. W., Peckham, A. D., Welge, J. A., & Otto, M. W. (2013). Patient preference for psychological vs pharmacologic treatment of psychiatric disorders: a meta-analytic review. *J Clin Psychiatry*, 74(6), 595-602.

- 1 McIndoo, C. C., File, A. A., Preddy, T., Clark, C. G., & Hopko, D. R. (2016). Mindfulness-based therapy and behavioral activation: A randomized controlled trial with depressed college students. *Behaviour research and therapy*, 77, 118-128.
- 4 McIntyre, A., Gendron, A., & McIntyre, A. (2007). Quetiapine adjunct to selective serotonin reuptake inhibitors or venlafaxine in patients with major depression, comorbid anxiety, and residual depressive symptoms: a randomized, placebo-controlled pilot study. *Depression and Anxiety*, 24(7), 487-494.
- 8 McIntyre, R. S., Soczynska, J. K., Liauw, S. S., Woldeyohannes, H. O., Brietzke, E., Nathanson, J., . . . Kennedy, S. H. (2012). The association between childhood adversity and components of metabolic syndrome in adults with mood disorders: results from the international mood disorders collaborative project. *Int J Psychiatry Med*, 43(2), 165-177.
- 12 McKnight, D. L., Nelson-Gray, R. O. & Barnhill, J. (1992). Dexamethasone suppression test and response to cognitive therapy and antidepressant medication. *Behavior Therapy* 23, 99-111.
- 15 McMahon, L., Foran, K., Forrest, S., Taylor, M., Ingram, G., Rajwal, M., Cornwall, P.L., McAllister-Williams, R.H.. (2007). Graduate mental health worker case management of depression in UK primary care: a pilot study. *British Journal of General Practice*, 57, 880-885.
- 19 McManus, S., & Bebbington, P. (2009). Adult psychiatric morbidity in England, 2007: results of a household survey. London: National Centre for Social Research.
- 21 McManus, S., Bebbington, P., Jenkins, R., & Brugha, T. (2016). Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. .
- 23 McNamara K, Horan JJ. Experimental construct validity in the evaluation of cognitive and behavioral treatments for depression. *Journal of Counseling Psychology*. 1986 Jan; 33(1):23.
- 25 McNeil JK, LeBlanc EM, Joyner M. The effect of exercise on depressive symptoms in the moderately depressed elderly. *Psychology and Aging*. 1991 Sep; 6(3):487.
- 27 McRae K, Rekshan W, Williams LM, Cooper N, Gross JJ. Effects of antidepressant medication on emotion regulation in depressed patients: An iSPOT-D report. *Journal of Affective Disorders*. 2014 Apr 20; 159:127-32.
- 30 Mead, G. E., Morley, W., Campbell, P., Greig, C. A., McMurdo, M., & Lawlor, D. A. (2008). Exercise for depression. *Cochrane Database Syst Rev* (4), CD004366.
- 32 Meadows, G. N., Shawyer, F., Enticott, J. C., Graham, A. L., Judd, F., Martin, P. R., ... Segal, Z. (2014). Mindfulness-based cognitive therapy for recurrent depression: A translational research study with 2-year follow-up. *Australian and New Zealand Journal of Psychiatry*, 48(8), 743-755.
- 36 Medicine, A. C. o. S. (1980). Guidelines for graded exercise testing and exercise prescription: Lea & Febiger.
- 38 Meesters, Y., Beersma, D. G., Bouhuys, A. L., & van den Hoofdakker, R. H. (1999). Prophylactic treatment of seasonal affective disorder (SAD) by using light visors: bright white or infrared light? *Biological Psychiatry*, 46(2), 239-246.
- 41 Melander, H., Ahlvist-Rastad, J., Meijer, G., & Beermann, B. (2003). Evidence based medicine--selective reporting from studies sponsored by pharmaceutical industry: review of studies in new drug applications. *BMJ*, 326(7400), 1171-1173.

- 1 Melartin, T. K., Rytsala, H. J., Leskela, U. S., Lestela-Mielonen, P. S., Sokero, T. P., & Isometsa, E. T. (2004). Severity and comorbidity predict episode duration and recurrence of DSM-IV major depressive disorder. *J Clin Psychiatry*, 65(6), 810-819.
- 4 Meltzer, H., Bebbington, P., Brugha, T., Farrell, M., Jenkins, R., & Lewis, G. (2000). The reluctance to seek treatment for neurotic disorders. *Journal of Mental Health*, 9(3), 319-327.
- 6 Melville, J. L., Reed, S. D., Russo, J., Croicu, C. A., Ludman, E., LaRocco-Cockburn, A., & Katon, W. (2014). Improving care for depression in obstetrics and gynecology: a randomized controlled trial. *Obstetrics and Gynecology*, 123, 1237-1246.
- 9 Memon, A., Taylor, K., Mohebati, L. M., Sundin, J., Cooper, M., Scanlon, T., & de Visser, R. (2016). Perceived barriers to accessing mental health services among black and minority ethnic (BME) communities: a qualitative study in Southeast England. *BMJ Open*, 6(11), e012337.
- 13 Menchetti M, Rucci P, Bortolotti B, Bombi A, Scocco P, Kraemer HC, Berardi D. Moderators of remission with interpersonal counselling or drug treatment in primary care patients with depression: randomised controlled trial. *The British Journal of Psychiatry*. 2013.
- 16 Menchetti, M., Sighinolfi, C., Michele, V., Peloso, P., Nespeca, C., Bandieri, P. V., ... Berardi, D. (2013). Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. *General Hospital Psychiatry*, 35(6), 579-586.
- 19 Mendels J. Double-blind comparison of citalopram and placebo in depressed outpatients with melancholia. *Depression & Anxiety* 9(2):54-60, 1999.
- 21 Mendels J, Fabre L, Kiev A. A double-blind parallel comparison of citalopram and placebo in out-patients with major depression or bipolar disorder, depressed. American Psychiatric Association Meeting (new clinical drug evaluation unit [ncdeu]), 1990.
- 24 Mendels J. Citalopram in the Treatment of Moderate to Severe Depression Conference Abstract. 150th Annual Meeting of the American Psychiatric Association San Diego, California, USA 17-22 May, 1997.
- 27 Mental-Health-Taskforce. (2015). The five year forward view Mental Health Taskforce: public engagement findings. London.
- 29 Merideth, C. H. & Feighner, J. P. (1983). A double-blind, controlled evaluation of zimeldine, imipramine and placebo in patients with primary affective disorders. *Acta Psychiatrica Scandinavica Supplementum*, 308, 70-79.
- 32 Merson, S., Tyrer, P., Lack, S., Birkett, P., Lynch, S., Onyett, S., & Johnson, T. (1992). Early intervention in psychiatric emergencies: a controlled clinical trial. *The Lancet*, 339(8805), 1311-1314.
- 35 Meyer B, Bierbrodt J, Schröder J, Berger T, Beevers CG, Weiss M, Jacob G, Späth C, Andersson G, Lutz W, Hautzinger M. Effects of an Internet intervention (Deprexis) on severe depression symptoms: randomized controlled trial. *Internet Interventions*. 2015 Mar 31; 2 (1):48-59.
- 39 Meyers, B. S., Flint, A. J., Rothschild, A. J., Mulsant, B. H., Whyte, E. M., Peasley-Miklus, C., . . . Heo, M. (2009). A double-blind randomized controlled trial of olanzapine plus sertraline vs olanzapine plus placebo for psychotic depression: the study of pharmacotherapy of psychotic depression (STOP-PD). *Archives of General Psychiatry*, 66(8), 838-847.
- 43 Meyers, B. S., Klimstra, S. A., Gabriele, M., Hamilton, M., Kakuma, T., Tirumalasetti, F., & Alexopoulos, G. S. (2001). Continuation treatment of delusional depression in older adults. *Am J Geriatr Psychiatry*, 9(4), 415-422.

- 1 MHRA. (2011). Citalopram and escitalopram: QT interval prolongation. Retrieved from
2 <https://www.gov.uk/drug-safety-update/citalopram-and-escitalopram-qt-interval-prolongation>
- 3 Michalak, J., Schultze, M., Heidenreich, T., & Schramm, E. (2015). A randomized controlled
4 trial on the efficacy of mindfulness-based cognitive therapy and a group version of cognitive
5 behavioral analysis system of psychotherapy for chronically depressed patients. *Journal of
6 Consulting and Clinical Psychology*, 83(5), 951-963.
- 7
- 8 Miller L & Weissman M. Interpersonal psychotherapy delivered over the telephone to
9 recurrent depressives. A pilot study. *Depression and Anxiety*. 2002 Jan 1; 16(3):114-7.
- 10 Miller, P., Chilvers, C., Dewey, M., Fielding, K., Gretton, V., Palmer, B., ... Harrison, G.
11 (2003). Counseling versus antidepressant therapy for the treatment of mild to moderate
12 depression in primary care economic analysis. *International Journal of Technology
13 Assessment in Health Care*, 19, 80-90.
- 14 Miller IW, Norman W, Keitner G, Bishop SB, Dow MG. Cognitive-behavioral treatment of
15 depressed inpatients. *Behav Ther* 1989; 20:25-47.
- 16 Mills, E. J., Thorlund, K., & Ioannidis, J. P. (2013). Demystifying trial networks and network
17 meta-analysis. *BMJ*, 346, f2914.
- 18 MIND. (2013). We still need to talk - A report on access to talking therapies Retrieved from
19 London: http://www.mind.org.uk/media/494424/we-still-need-to-talk_report.pdf
- 20 Mindham, R. H. (1982). Tricyclic antidepressants Drugs in Psychiatric Practice (pp. 177-218):
21 Elsevier BV.
- 22 Miranda, J., Chung, J. Y., Green, B. L., Krupnick, J., Siddique, J., Revicki, D. A., & Belin, T.
23 (2003). Treating depression in predominantly low-income young minority women: a
24 randomized controlled trial. *JAMA*, 290(1), 57-65.
- 25 Mischoulon, D., Alpert, J. E., Arning, E., Bottiglieri, T., Fava, M., & Papakostas, G. I. (2012).
26 Bioavailability of S-Adenosyl Methionine and Impact on Response in a Randomized
27 Controlled Trial in Major Depressive Disorder. *The Journal of Clinical Psychiatry*, 73(6), 843.
- 28 Mischoulon, D., Nierenberg, A. A., Schettler, P. J., Kinhead, B. L., Fehling, K., Martinson, M.
29 A., & Rapaport, M. H. (2015). A double-blind, randomized controlled clinical trial comparing
30 eicosapentaenoic acid versus docosahexaenoic acid. *Journal of Clinical Psychiatry*, 76(1),
31 54-61.
- 32 Mischoulon, D., et al. (2014) A double-blind, randomized, placebo-controlled clinical trial of
33 S-ade syl-L-methionine (SAMe) versus escitalopram in major depressive disorder. *The
34 Journal of clinical psychiatry* 75, 370-376.
- 35 Mitchell, A. J., & Subramaniam, H. (2005). Prognosis of depression in old age compared to
36 middle age: A systematic review of comparative studies. *American Journal of Psychiatry*,
37 162(9), 1588-1601.
- 38 Mitchell, A. J., Vaze, A., & Rao, S. (2009). Clinical diagnosis of depression in primary care: a
39 meta-analysis. *Lancet*, 374(9690), 609-619.
- 40 Modell, J. G., Rosenthal, N. E., Harriett, A. E., Krishen, A., Asgharian, A., Foster, V. J., . . .
41 Wightman, D. S. (2005). Seasonal affective disorder and its prevention by anticipatory
42 treatment with bupropion XL. *Biological Psychiatry*, 58(8), 658-667.
- 43 Moffitt, T. E., Caspi, A., Taylor, A., Kokaua, J., Milne, B. J., Polanczyk, G., & Poulton, R.
44 (2010). How common are common mental disorders? Evidence that lifetime prevalence rates
45 are doubled by prospective versus retrospective ascertainment. *Psychol Med*, 40(6), 899-
46 909.

- 1 Mohr, D. C., et al. (2012) Effect of telephone-administered vs face-to-face cognitive behavioral therapy on adherence to therapy and depression outcomes among primary care patients: a randomized trial. *JAMA* 307, 2278-2285.
- 4 Mohr, D. C., Carmody, T., Erickson, L., Jin, L., & Leader, J. (2011). Telephone-administered cognitive behavioral therapy for veterans served by community-based outpatient clinics. *Journal of Consulting and Clinical Psychology*, 79(2), 261.
- 7 Moldovan, R., Cobeanu, O., & David, D. (2013). Cognitive bibliotherapy for mild depressive symptomatology: randomized clinical trial of efficacy and mechanisms of change. *Clinical Psychology & Psychotherapy*, 20(6), 482-493.
- 10 Molenaar PJ, Boom Y, Peen J, Schoevers RA, Van R, Dekker JJ. Is there a dose–effect relationship between the number of psychotherapy sessions and improvement of social functioning? *British Journal of Clinical Psychology*. 2011 Sep 1; 50 (3):268-82.
- 13 Moller HJ, Gallinat J, Hegerl U, et al. Double-blind, multicenter comparative study of sertraline and amitriptyline in hospitalized patients with major depression. *Pharmacopsychiatry* 1998; 31:170-177.
- 16 Moller HJ, Glaser K, Leverkus F et al. Double-blind, multicenter comparative study of sertraline versus amitriptyline in outpatients with major depression. *Pharmacopsychiatry* 2000; 33:206-212.
- 19 Moller-Leimkuhler, A. M. (2002). Barriers to help-seeking by men: a review of sociocultural and clinical literature with particular reference to depression. *J Affect Disord*, 71(1-3), 1-9.
- 21 Monroe, S. M., & Harkness, K. L. (2011). Recurrence in major depression: a conceptual analysis. *Psychol Rev*, 118(4), 655-674.
- 23 Montgomery, S. A., Dufour, H., Brion, S., & Galledreau, J. (1988). The prophylactic efficacy of fluoxetine in unipolar depression. *The British Journal of Psychiatry*, (3):69-76.
- 25 Montgomery, S. A., & Dunbar, G. (1993). Paroxetine is better than placebo in relapse prevention and the prophylaxis of recurrent depression. *International Clinical Psychopharmacology*, 8(3), 189-196.
- 28 Montgomery, S. A., Entsuah, R., Hackett, D., Kunz, N. R., & Rudolph, R. L. (2004). Venlafaxine versus placebo in the preventive treatment of recurrent major depression. *The Journal of Clinical Psychiatry*, 65(3), 328-336.
- 31 Montgomery SA, Rasmussen JG, Tanghoj P. A 24-week study of 20 mg citalopram, 40 mg citalopram, and placebo in the prevention of relapse of major depression. *International Clinical Psychopharmacology* 1993; 8(3):181-188.
- 34 Moore, N., Verdoux, H. & Fantino, B. (2005) Prospective, multicentre, randomized, double-blind study of the efficacy of escitalopram versus citalopram in outpatient treatment of major depressive disorder. *International Clinical Psychopharmacology*, 20, 131-137.
- 37 Moradveisi L, Huibers MJ, Renner F, Arasteh M, Arntz A. Behavioural activation v. antidepressant medication for treating depression in Iran: randomised trial. *The British Journal of Psychiatry*. 2013 Mar 1; 202(3):204-11.
- 40 Moreno, S. G., Sutton, A. J., Ades, A., Cooper, N. J., & Abrams, K. R. (2011). Adjusting for publication biases across similar interventions performed well when compared with gold standard data. *Journal of clinical epidemiology*, 64(11), 1230-1241.
- 43 Moreno, S. G., Sutton, A. J., Ades, A., Stanley, T. D., Abrams, K. R., Peters, J. L., & Cooper, N. J. (2009). Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study. *BMC Medical Research Methodology*, 9(1), 2.

- 1 Morgan, O., Griffiths, C., Baker, A., & Majeed, A. (2004). Fatal toxicity of antidepressants in England and Wales, 1993-2002. *Health Stat Q*(23), 18-24.
- 2 Moritz, S., et al. (2012) A randomized controlled trial of internet-based therapy in depression. *Behaviour Research and Therapy* 50, 513-521.
- 3 Morris RR, Schueller SM, Picard RW. Efficacy of a web-based, crowdsourced peer-to-peer cognitive reappraisal platform for depression: Randomized controlled trial. *Journal of medical Internet research*. 2015 Mar; 17 (3).
- 4 Moscovitch, A., Blashko, C. A., Eagles, J. M., Darcourt, G., Thompson, C., Kasper, S., & Lane, R. M. (2004). A placebo-controlled study of sertraline in the treatment of outpatients with seasonal affective disorder. *Psychopharmacology*, 171(4), 390-397.
- 5 Moss, K., Scogin, F., Napoli, E., & Presnell, A. (2012). A self-help behavioral activation treatment for geriatric depressive symptoms. *Aging & Mental Health*, 16(5), 625-635.
- 6 Mota-Pereira, J., Silverio, J., Carvalho, S., Ribeiro, J. C., Fonte, D., & Ramos, J. (2011). Moderate exercise improves depression parameters in treatment-resistant patients with major depressive disorder. *Journal of psychiatric research*, 45(8), 1005-1011.
- 7 Moussavi, S., Chatterji, S., Verdes, E., Tandon, A., Patel, V., & Ustun, B. (2007). Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *The Lancet*, 370(9590), 851-858.
- 8 Mozaffari-Khosravi H, Yassini-Ardakani M, Karamati M, Shariati-Bafghi SE. Eicosapentaenoic acid versus docosahexaenoic acid in mild-to-moderate depression: a randomized, double-blind, placebo-controlled trial. *European Neuropsychopharmacology*. 2013 Jul 31; 23(7):636-44.
- 9 Mrazek, D. A., Hornberger, J. C., Altar, C. A., & Degtiar, I. (2014). A review of the clinical, economic, and societal burden of treatment-resistant depression: 1996-2013. *Psychiatr Serv*, 65(8), 977-987.
- 10 Mueller, T. I., Keller, M. B., Leon, A. C., Solomon, D. A., Shea, M. T., Coryell, W., & Endicott, J. (1996). Recovery after 5 years of unremitting major depressive disorder. *Arch Gen Psychiatry*, 53(9), 794-799.
- 11 Mueller, T. I., Leon, A. C., Keller, M. B., Solomon, D. A., Endicott, J., Coryell, W., ... Maser, J. D. (1999). Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. *Am J Psychiatry*, 156(7), 1000-1006.
- 12 Muijen M, Roy D, Silverstone T, Mehmet A, Christie M. A comparative clinical trial of fluoxetine, mianserin and placebo in depressed outpatients. *Acta Psychiatrica Scandinavica* 1988 Sep; 78 (3):384-90.
- 13 Mukuria, C., Brazier, J., Barkham, M., Connell, J., Hardy, G., Hutten, R., ... Parry, G. (2013). Cost-effectiveness of an improving access to psychological therapies service. *British Journal of Psychiatry*, 202, 220-227.
- 14 Mullin J. A multicentre, double-blind, amitriptyline-controlled study of mirtazapine in patients with major depression. *Journal of Psychopharmacology* 1996; 10 (3):235-240.
- 15 Mullin, A., et al. (2015). The UniWellbeing course: A randomised controlled trial of a transdiagnostic internet-delivered cognitive behavioural therapy (CBT) programme for university students with symptoms of anxiety and depression. *Internet Interventions* 2(2 // (NHMRC) *National Health and Medical Research Council*): 128-136.
- 16 Mulsant, B. H., Sweet, R. A., Rosen, J., Pollock, B. G., Zubenko, G. S., Flynn, T., ... Reynolds, C. F., 3rd. (2001). A double-blind randomized comparison of nortriptyline plus

- 1 perphenazine versus nortriptyline plus placebo in the treatment of psychotic depression in
2 late life. *J Clin Psychiatry*, 62(8), 597-604.
- 3 Murphy GE, Carney RM, Knesevitch MA, Wetzel RD, Whitworth P. Cognitive behavior
4 therapy, relaxation training, and tricyclic antidepressant medication in the treatment of
5 depression. *Psychol Rep* 1995; 77(2):403-420.
- 6 Murphy, S. M., Irving, C. B., Adams, C. E., & Waqar, M. (2015). Crisis intervention for people
7 with severe mental illnesses. *Cochrane Database Syst Rev*. 16(5):CD001087.
- 8 Murray, C. J., Lopez, A. D., & Jamison, D. T. (1994). The global burden of disease in 1990:
9 summary results, sensitivity analysis and future directions. *Bull World Health Organ*, 72(3),
10 495-509.
- 11 Murphy GE, Simons AD, Wetzel RD, Lustman PJ. Cognitive therapy and pharmacotherapy
12 singly and together in the treatment of depression. *Arch Gen Psychiatry* 1984; 41:33-41.
- 13 Murray, C. J., Vos, T., Lozano, R., Naghavi, M., Flaxman, A. D., Michaud, C., ... Memish, Z.
14 A. (2012). Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions,
15 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*,
16 380(9859), 2197-2223.
- 17 Murray, G., Michalak, E. E., Axler, A., Yaxley, D., Hayashi, B., Westrin, A., ... Lam, R. W.
18 (2010). Relief of chronic or resistant depression (Re-ChORD): a pragmatic, randomized,
19 open-treatment trial of an integrative program intervention for chronic depression. *Journal of
Affective Disorders*, 123(1-3), 243-248.
- 21 Murray G, Michalak EE, Axler A, et al. Relief of Chronic or Resistant Depression (Re-
22 ChORD): A pragmatic, randomized, open-treatment trial of an integrative program
23 intervention for chronic depression. *Journal of affective disorders*. 2010 Jun 30; 123(1):243-
24 8.
- 25 Murray, J., Banerjee, S., Byng, R., Tylee, A., Bhugra, D., & Macdonald, A. (2006). Primary
26 care professionals' perceptions of depression in older people: a qualitative study. *Soc Sci
Med*, 63(5), 1363-1373.
- 28 Murri MB, Amore M, Menchetti M, Toni G, Neviani F, Cerri M, Rocchi MB, Zocchi D, Bagnoli
29 L, Tam E, Buffa A. Physical exercise for late-life major depression. *The British Journal of
Psychiatry*. 2015 Sep 1; 207(3):235-42.
- 31 Mustafa, R. A., Santesso, N., Brozek, J., Akl, E. A., Walter, S. D., Norman, G., ...
32 Schunemann, H. J. (2013). The GRADE approach is reproducible in assessing the quality of
33 evidence of quantitative evidence syntheses. *J Clin Epidemiol*, 66(7), 736-742; quiz 742
34 e731-735.
- 35 Mynors-Wallis, L. M., Gath, D. H., Day, A., & Baker, F. (2000). Randomised controlled trial of
36 problem solving treatment, antidepressant medication, and combined treatment for major
37 depression in primary care. *BMJ*, 320(7226), 26-30.
- 38 Mynors-Wallis, L. M., Gath, D. H., Lloyd-Thomas, A. R., & Tomlinson, D. (1995).
39 Randomised controlled trial comparing problem solving treatment with amitriptyline and
40 placebo for major depression in primary care. *BMJ*, 310(6977), 441-445.
- 41 Naeem, F., Gul, M., Irfan, M., Munshi, T., Asif, A., Rashid, S., ... Aslam, M. (2015). Brief
42 culturally adapted CBT (CaCBT) for depression: a randomized controlled trial from Pakistan.
43 *Journal of Affective Disorders*, 177, 101-107.
- 44 Naeem, F., Sarhandi, I., Gul, M., Khalid, M., Aslam, M., Anbrin, A., ... Minhas, F. (2014). A
45 multicentre randomised controlled trial of a carer supervised culturally adapted CBT (CaCBT)
46 based self-help for depression in Pakistan. *Journal of Affective Disorders*, 156, 224-227.

- 1 Naeem, F., et al. (2013) A multicentre randomised controlled trial of a carer supervised culturally adapted CBT (CACBT) based self-help for depression in Pakistan. *Journal of Affective Disorders* 156, 224-227.
- 4 Naeem F, Waheed W, Gobbi M, Ayub M, Kingdon D. Preliminary evaluation of culturally sensitive CBT for depression in Pakistan: findings from Developing Culturally-sensitive CBT Project (DCCP). *Behavioural and Cognitive Psychotherapy*. 2011 Mar 1; 39 (02):165-73.
- 7 Nair, N., Amin, M., Holm, P., Katona, C., Klitgaard, N., Kin, N. N. Y., ... Stage, K. (1995). Moclobemide and nortriptyline in elderly depressed patients. A randomized, multicentre trial against placebo. *Journal of Affective Disorders*, 33(1), 1-9.
- 10 Nandi, D. N., Ajmany, S., Ganguli, H., Banerjee, G., Boral, G. C., Ghosh, A. et al. (1976). A clinical evaluation of depressives found in a rural survey in India. *British Journal of Psychiatry*, 128, 523- 527.
- 13 Nanni, V., Uher, R., & Danese, A. (2012). Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: a meta-analysis. *Am J Psychiatry*, 169(2), 141-151.
- 16 Napadow, V., Makris, N., Liu, J., Kettner, N. W., Kwong, K. K., & Hui, K. K. (2005). Effects of electroacupuncture versus manual acupuncture on the human brain as measured by fMRI. *Hum Brain Mapp*, 24(3), 193-205.
- 19 NAPT. (2013). National audit of psychological therapy services for depression and anxiety
- 20 National College for Teaching and Leadership. (2016). Review of clinical and educational psychology training arrangements.
- 22 Navarro, V., Gasto, C., Torres, X., Masana, G., Penades, R., Guarch, J., ... Catalan, R. (2008). Continuation/maintenance treatment with nortriptyline versus combined nortriptyline and ECT in late-life psychotic depression: a two-year randomized study. *Am J Geriatr Psychiatry*, 16(6), 498-505.
- 26 Naylor, E. V., Antonuccio, D. O., Litt, M., Johnson, G. E., Spogen, D. R., Williams, R., ... Higgins, D. L. (2010). Bibliotherapy as a treatment for depression in primary care. *Journal of Clinical Psychology in Medical Settings*, 17(3), 258-271.
- 29 Nazroo, J. Y., Edwards, A. C., & Brown, G. W. (1997). Gender differences in the onset of depression following a shared life event: a study of couples. *Psychol Med*, 27(1), 9-19.
- 31 Newby, J. M., et al. (2013) Internet cognitive behavioural therapy for mixed anxiety and depression: a randomized controlled trial and evidence of effectiveness in primary care [Wellbeing 6]. *Psychological Medicine*, 43, 2635-2648.
- 34 Newton-Howes, G., Tyrer, P., & Johnson, T. (2006). Personality disorder and the outcome of depression: meta-analysis of published studies. *Br J Psychiatry*, 188, 13-20.
- 36 Nezu, A. M. (1987). A problem-solving formulation of depression: A literature review and proposal of a pluralistic model. *Clinical Psychology Review*, 7(2), 121-144.
- 38 Nezu, A. M., Nezu, C. M., & Perri, M. G. (1989). Problem-solving therapy for depression: theory, research, and clinical guidelines. New York: Wiley.
- 40 NHS. (2017). National Drug Tariff. London.
- 41 NHS England (Producer). (2016, 18 September 2016). More people than ever receiving psychological therapies and recovering.
- 43 NHS The Information Centre. (2007). Prescription Cost Analysis: England 2006.

- 1 NICE. (2003). Guidance on the use of electroconvulsive therapy. TA59. London. NICE.
- 2 NICE. (2004). Depression: management of depression in primary and secondary care. CG23. London. NICE.
- 3 NICE. (2005). Depression in children and young people: Identification and management in primary, community and secondary care. CG28. London. NICE.
- 4 NICE. (2009). Borderline personality disorder: recognition and management. CG78 London. NICE.
- 5 NICE. (2009). Depression in adults with a chronic physical health problem: treatment and management. CG91. London. NICE.
- 6 NICE. (2009). Depression in adults: recognition and management. CG90. London. NICE.
- 7 NICE. (2009). Vagus nerve stimulation for treatment-resistant depression. IPG330. London. NICE.
- 8 NICE. (2011). Common mental health disorders - identification and pathways to care. CG123. London. NICE.
- 9 NICE. (2011). Service user experience in adult mental health: improving the experience of care for people using adult NHS mental health services. CG136. London. NICE.
- 10 NICE. (2013). Guide to the methods of technology appraisal. London. NICE.
- 11 NICE. (2014). Antenatal and postnatal mental health: clinical management and service guidance. CG192. London. NICE.
- 12 NICE. (2014). Bipolar disorder: assessment and management. CG185. London. NICE.
- 13 NICE. (2014). Developing NICE guidelines: the manual. London. NICE.
- 14 NICE. (2015). Repetitive direct current stimulation for depression. IPG542. London. NICE.
- 15 NICE. (2015). Transcranial direct current stimulation for depression. IPG530. London. NICE.
- 16 NICE. (2015). Vortioxetine for treating major depressive episodes. TA367. London. NICE.
- 17 Nicholson, A., Kuper, H., & Hemingway, H. (2006). Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J*, 27(23), 2763-2774.
- 18 Nierenberg, A. A., Fava, M., Trivedi, M. H., Wisniewski, S. R., Thase, M. E., McGrath, P. J., ... Niederehe, G. (2006). A comparison of lithium and T 3 augmentation following two failed medication treatments for depression: a STAR* D report. *American Journal of Psychiatry*, 163(9), 1519-1530.
- 19 Nierenberg, A. A., Greist, J. H., Mallinckrodt, C. H., Prakash, A., Sambunaris, A., Tollefson, G. D. et al. (2007). Duloxetine versus escitalopram and placebo in the treatment of patients with major depressive disorder: onset of antidepressant action, a non-inferiority study. *Current Medical Research & Opinion*, 23, 401-416.
- 20 Nierenberg, A. A., Papakostas, G. I., Petersen, T., Kelly, K. E., Iacoviello, B. M., Worthington, J. J., ... Fava, M. (2003). Nortriptyline for treatment-resistant depression. *The Journal of Clinical Psychiatry*. 64(1):35-9.
- 21 Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspect Psychol Sci*, 3(5), 400-424.

- 1 Norbury, R., Taylor, M. J., Selvaraj, S., Murphy, S. E., Harmer, C. J., & Cowen, P. J. (2009).
- 2 Short-term antidepressant treatment modulates amygdala response to happy faces.
- 3 Psychopharmacology, 206(2), 197-204.
- 4 Nordenskjöld, A., Knorring, L., Ljung, T., Carlborg, A., Brus, O., & Engström, I. (2013).
- 5 Continuation electroconvulsive therapy with pharmacotherapy versus pharmacotherapy
- 6 alone for prevention of relapse of depression: a randomized controlled trial. Journal of ECT,
- 7 29(2), 86-92.
- 8 Nordstrom, G., Despiegel, N., Marteau, F., Danchenko, N., & Maman, K. (2010). Cost
- 9 effectiveness of escitalopram versus SNRIs in second-step treatment of major depressive
- 10 disorder in Sweden. Journal of Medical Economics, 13, 516-526.
- 11 Nutt, D. J. (2010). Rationale for, barriers to, and appropriate medication for the long-term
- 12 treatment of depression. The Journal of Clinical Psychiatry, 71(suppl E1).
- 13 Nuttbrock, L., Bockting, W., Rosenblum, A., Mason, M., Macri, M., & Becker, J. (2012).
- 14 Gender identity conflict/affirmation and major depression across the life course of
- 15 transgender women. International Journal of Transgenderism, 13(3), 91-103.
- 16 Nyth AL, Gottfries CG, Lyby K, Smedegaard-Andersen L, Gylding-Sabroe J, Kristensen M,
- 17 Refsum HE, Ofsti E, Eriksson S, Syversen S. A controlled multicenter clinical study of
- 18 citalopram and placebo in elderly depressed patients with and without concomitant dementia.
- 19 Acta Psychiatrica Scandinavica 1992; 86 (2):138-145.
- 20 Odell, S. M., Surtees, P. G., Wainwright, N. W., Commander, M. J., & Sashidharan, S. P.
- 21 (1997). Determinants of general practitioner recognition of psychological problems in a multi-
- 22 ethnic inner-city health district. Br J Psychiatry, 171, 537-541.
- 23 Oka, Y., Okamoto, K., Kawashita, N., Shirakuni, Y., & Takagi, T. (2014). Meta-analysis of the
- 24 risk of upper gastrointestinal hemorrhage with combination therapy of selective serotonin
- 25 reuptake inhibitors and non-steroidal anti-inflammatory drugs. Biol Pharm Bull, 37(6), 947-
- 26 953.
- 27 Okumura, Y., & Higuchi, T. (2011). Cost of depression among adults in Japan. The Primary
- 28 Care Companion For CNS Disorders, 13(3). pii:
- 29 Oladeji, B. D., Kola, L., Abiona, T., Montgomery, A. A., Araya, R., & Gureje, O. (2015). A pilot
- 30 randomized controlled trial of a stepped care intervention package for depression in primary
- 31 care in Nigeria. BMC Psychiatry, 15(1).
- 32 O'Leary, K. D., Riso, L. P., & Beach, S. R. (1990). Attributions about the marital
- 33 discord/depression link and therapy outcome. Behavior therapy, 21(4), 413-422.
- 34 Old Age Depression Interest Group. How long should the elderly take antidepressants? A
- 35 double-blind placebo-controlled study of continuation/prophylaxis therapy with dothiepin. Old
- 36 Age Depression Interest Group.[comment]. British Journal of Psychiatry 1993; 162:175-182.
- 37 Olfson, M., Marcus, S. C., Tedeschi, M., & Wan, G. J. (2006). Continuity of antidepressant
- 38 treatment for adults with depression in the United States. Am J Psychiatry, 163(1), 101-108.
- 39 Olgiati, P., Bajo, E., Bigelli, M., Montgomery, S., & Serretti, A. (2013). Challenging sequential
- 40 approach to treatment resistant depression: Cost-utility analysis based on the Sequenced
- 41 Treatment Alternatives to Relieve Depression (STAR*D) trial. European
- 42 Neuropsychopharmacology, 23(12), 1739-1746.
- 43 Olie JP, Gunn KP, Katz E. A double-blind placebo-controlled multicentre study of sertraline in
- 44 the acute and continuation treatment of major depression. European Psychiatry 1997;
- 45 12(1):34-41.

- 1 ONS. (2016). Suicides in the UK: 2015 registrations.
- 2 Ormel, J., Brink, W. V. D., Koeter, M. W. J., Giel, R., Meer, K. V. D., Willige, G. V. D., & Wilmink, F. W. (1990). Recognition, management and outcome of psychological disorders in primary care: a naturalistic follow-up study*. *Psychological Medicine*, 20(04), 909.
- 5 Ormel, J., & Costa de Silva, J. A. (1995). The impact of psychopathology on disability and health perceptions. In T. B. Üstün & N. Sartorius (Eds.), *Mental illness in general health care: an international study*. Chichester: Wiley.
- 8 Ormel, J., Oldehinkel, T., Brilman, E., & vanden Brink, W. (1993). Outcome of depression and anxiety in primary care. A three-wave 3 1/2-year study of psychopathology and disability. *Arch Gen Psychiatry*, 50(10), 759-766.
- 11 Ormel, J., Rijssdijk, F. V., Sullivan, M., van Sonderen, E., & Kempen, G. I. (2002). Temporal and reciprocal relationship between IADL/ADL disability and depressive symptoms in late life. *J Gerontol B Psychol Sci Soc Sci*, 57(4), P338-347.
- 14 Ormel, J., Vonkorff, M., Oldehinkel, A. J., Simon, G., Tiemens, B. G., & ÜStÜN, T. B. (1999). Onset of disability in depressed and non-depressed primary care patients. *Psychological Medicine*, 29(4), 847-853.
- 17 Oslin, D. W., Sayers, S., Ross, J., Kane, V., Ten Have, T., Conigliaro, J., & Cornelius, J. (2003). Disease management for depression and at-risk drinking via telephone in an older population of veterans. *Psychosom Med*, 65(6), 931-937.
- 20 Ostergaard, S. D., Bille, J., Soltoft-Jensen, H., Lauge, N., & Bech, P. (2012). The validity of the severity-psychosis hypothesis in depression. *J Affect Disord*, 140(1), 48-56.
- 22 Ostergaard, S. D., Rothschild, A. J., Flint, A. J., Mulsant, B. H., Whyte, E. M., Vermeulen, T., ... Meyers, B. S. (2016). Establishing the cut-off score for remission and severity-ranges on the Psychotic Depression Assessment Scale (PDAS). *J Affect Disord*, 190, 111-114.
- 25 Ostler, K. (2001). Influence of socio-economic deprivation on the prevalence and outcome of depression in primary care: The Hampshire Depression Project. *The British Journal of Psychiatry*, 178(1), 12-17.
- 28 Ou JJ, Xun GL, Wu RR, Li LH, Fang MS, Zhang HG, et al. Efficacy and safety of escitalopram versus citalopram in major depressive disorder: a 6-week, multicenter, randomized, double-blind, flexible-dose study. *Psychopharmacology (Berl)* 2011; 213:639-46.
- 32 Oxman TE, Hegel MT, Hull JG, Dietrich AJ. Problem-solving treatment and coping styles in primary care for minor depression. *Journal of consulting and clinical psychology*. 2008 Dec; 76(6):933.
- 35 Pace TM, Dixon DN. Changes in depressive self-schemata and depressive symptoms following cognitive therapy. *Journal of Counseling Psychology*. 1993 Jul; 40(3):288.
- 37 Pachankis, J. E., Cochran, S. D., & Mays, V. M. (2015). The mental health of sexual minority adults in and out of the closet: A population-based study. *Journal of Consulting and Clinical Psychology*, 83(5), 890-901.
- 40 Palmer, S. C., & Coyne, J. C. (2003). Screening for depression in medical care: pitfalls, alternatives, and revised priorities. *J Psychosom Res*, 54(4), 279-287.
- 42 Papakostas GI, Fava M, Baer L, Swee MB, Jaeger A, Bobo WV, Shelton RC. Ziprasidone augmentation of escitalopram for major depressive disorder: efficacy results from a randomized, double-blind, placebo-controlled study. *American Journal of Psychiatry*. 2015 Jun 18; 172(12):1251-8.

- 1 Papakostas, G. I., Fava, M., & Thase, M. E. (2008). Treatment of SSRI-resistant depression: a meta-analysis comparing within-versus across-class switches. *Biological Psychiatry*, 63(7), 699-704.
- 4 Park, Y., et al. (2015). "Supplementation of n-3 polyunsaturated fatty acids for major depressive disorder: A randomized, double-blind, 12-week, placebo-controlled trial in Korea." *Annals of Nutrition & Metabolism* 66(2-3): 141-148.
- 7 Park, S. C., Choi, J., Kim, J. M., Jun, T. Y., Lee, M. S., Kim, J. B., ... Park, Y. C. (2014). Is the Psychotic Depression Assessment Scale a useful diagnostic tool? The CRESCEND study. *J Affect Disord*, 166, 79-85.
- 10 Park, Y., Park, Y. S., Kim, S. H., Oh, D. H., & Park, Y. C. (2015). Supplementation of n-3 polyunsaturated fatty acids for major depressive disorder: A randomized, double-blind, 12-week, placebo-controlled trial in Korea. *Annals of Nutrition & Metabolism*, 66(2-3), 141-148.
- 13 Parker, G. (2012). Diagnosis, classification, and differential diagnosis of the mood disorders. In M. G. Gelder, N. Andreasen, J. Lopez-Ibor, & J. Geddes (Eds.), *New Oxford textbook of psychiatry* (2nd ed. ed.). Oxford: Oxford University Press.
- 16 Partonen, T., & Lönnqvist, J. (1996). Moclobemide and fluoxetine in treatment of seasonal affective disorder. *Journal of Affective Disorders*, 41(2), 93-99.
- 18 Patel, V., Weiss, H. A., Chowdhary, N., Naik, S., Pednekar, S., Chatterjee, S., Kirkwood, B. R. (2010). Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *Lancet*, 376(9758), 2086-2095.
- 22 Patkar, A. A., Peindl, K., Mago, R., Mannelli, P., & Masand, P. S. (2006). An open-label, rater-blinded, augmentation study of aripiprazole in treatment-resistant depression. *Primary care companion to the Journal of Clinical Psychiatry*, 8(2), 82.
- 25 Patten, S. B. (1991). Are the Brown and Harris "vulnerability factors" risk factors for depression? *J Psychiatry Neurosci*, 16(5), 267-271.
- 27 Patten, S. B. (2001). Long-term medical conditions and major depression in a Canadian population study at waves 1 and 2. *J Affect Disord*, 63(1-3), 35-41.
- 29 Paykel, E. S., Ramana, R., Cooper, Z., Hayhurst, H., Kerr, J., & Barocka, A. (1995). Residual symptoms after partial remission: an important outcome in depression. *Psychol Med*, 25(6), 1171-1180.
- 32 Peckham, C. S., & Dezateux, C. (1998). Issues underlying the evaluation of screening programmes. *Br Med Bull*, 54(4), 767-778.
- 34 Pedersen, R. D., Pallay, A. G., & Rudolph, R. L. (2002). Can improvement in well-being and functioning be distinguished from depression improvement in antidepressant clinical trials? *Quality of Life Research*, 11, 9-17.
- 37 Peet, M., & Horrobin, D. F. (2002). A dose-ranging study of the effects of ethyl-eicosapentaenoate in patients with ongoing depression despite apparently adequate treatment with standard drugs. *Archives of General Psychiatry*, 59(10), 913-919.
- 40 Pentecost, C., et al. (2015). Combining behavioural activation with physical activity promotion for adults with depression: Findings of a parallel-group pilot randomised controlled trial (BACPAc). *Trials* 16(1).
- 43 Perahia, D., Wang, F., Mallinckrodt, C., Walker, D., & Detke, M. (2006). Duloxetine in the treatment of major depressive disorder: a placebo-and paroxetine-controlled trial. *European Psychiatry*, 21(6), 367-378.

- 1 Perahia, D. G., Maina, G., Thase, M. E., Spann, M. E., Wang, F., Walker, D. J., & Detke, M.
- 2 J. (2009). Duloxetine in the prevention of depressive recurrences: a randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry*, 70(5), 706-716.
- 3
- 4 Perahia, D. G., Quail, D., Gandhi, P., Walker, D. J., & Peveler, R. C. (2008). A randomized, controlled trial of duloxetine alone vs. duloxetine plus a telephone intervention in the treatment of depression. *Journal of Affective Disorders*, 108, 33-41.
- 5
- 6
- 7 Perini, S., Titov, N., & Andrews, G. (2009). Clinician-assisted Internet-based treatment is effective for depression: randomized controlled trial. *Australian and New Zealand Journal of Psychiatry*, 43(6), 571-578.
- 8
- 9
- 10 Perlis, R. H. (2014). Pharmacogenomic testing and personalized treatment of depression. *Clin Chem*, 60(1), 53-59.
- 11
- 12 Perlis, R. H., Nierenberg, A. A., Alpert, J. E., Pava, J., Matthews, J. D., Buchin, J., ... Fava, M. (2002). Effects of adding cognitive therapy to fluoxetine dose increase on risk of relapse and residual depressive symptoms in continuation treatment of major depressive disorder. *Journal of Clinical Psychopharmacology*, 22(5), 474-480.
- 13
- 14
- 15
- 16 Peselow, E. D., Stanley, M., Filippi, A. M., Barouche, F., Goodnick, P., & Fieve, R. R. (1989).
- 17 The predictive value of the dexamethasone suppression test. A placebo-controlled study. *British Journal of Psychiatry*, 155, 667-672.
- 18
- 19 Petersen, T. J., Pava, J. A., Buchin, J., Matthews, J. D., Papakostas, G. I., Nierenberg, A. A., Fava, M. (2010). The role of cognitive-behavioral therapy and fluoxetine in prevention of recurrence of major depressive disorder. *Cognitive Therapy and Research*, 34(1), 13-23.
- 20
- 21
- 22 Peveler, R., George, C., Kinmonth, A. L., Campbell, M., & Thompson, C. (1999). Effect of antidepressant drug counselling and information leaflets on adherence to drug treatment in primary care: randomised controlled trial. *BMJ*, 319(7210), 612-615.
- 23
- 24
- 25 Peveler, R., Kendrick, A., Buxton, M., Longworth, L., Baldwin, D., Moore, M., ... Smith, H. (2005). A randomised controlled trial to compare the cost-effectiveness of tricyclic antidepressants, selective serotonin reuptake inhibitors and lofepramine. *Health Technol Assess*, 9(16):1-134, iii.
- 26
- 27
- 28
- 29 Pfaff JJ, Alfonso H, Newton RU, Sim M, Flicker L, Almeida OP. ACTIVEDEP: a randomised, controlled trial of a home-based exercise intervention to alleviate depression in middle-aged and older adults. *British journal of sports medicine*. 2014 Feb; 48(3):226-32. Epub 2013 Jul 6.
- 30
- 31
- 32
- 33 Philipp, M., Kohnen, R., & Hiller, K. O. (1999). Hypericum extract versus imipramine or placebo in patients with moderate depression: randomised multicentre study of treatment for eight weeks. *BMJ*, 319(7224), 1534-1538.
- 34
- 35
- 36 Phillips, R., Schneider, J., Molosankwe, I., Leese, M., Foroushani, P. S., Grime, P., ... Thornicroft, G. (2014). Randomized controlled trial of computerized cognitive behavioural therapy for depressive symptoms: effectiveness and costs of a workplace intervention. *Psychological Medicine*, 44, 741-752.
- 37
- 38
- 39
- 40 Pignone, M. P., Gaynes, B. N., Rushton, J. L., Burchell, C. M., Orleans, C. T., Mulrow, C. D., & Lohr, K. N. (2002). Screening for depression in adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med*, 136(10), 765-776.
- 41
- 42
- 43 Poirier, M., & Boyer, P. (1999). Venlafaxine and paroxetine in treatment-resistant depression. Double-blind, randomised comparison. *The British Journal of Psychiatry*, 175(1), 12-16.
- 44
- 45 Pollock, K., & Grime, J. (2002). Patients' perceptions of entitlement to time in general practice consultations for depression: qualitative study. *BMJ*, 325(7366), 687.
- 46

- 1 Popay, J., Kowarzik, U., Mallinson, S., Mackian, S., & Barker, J. (2007). Social problems, primary care and pathways to help and support: addressing health inequalities at the individual level. Part I: the GP perspective. *J Epidemiol Community Health*, 61(11), 966-971.
- 4 Porter, R. J., Douglas, K., & Knight, R. G. (2008). Monitoring of cognitive effects during a course of electroconvulsive therapy: recommendations for clinical practice. *J ECT*, 24(1), 25-34.
- 7 Posternak, M. A., & Miller, I. (2001). Untreated short-term course of major depression: a meta-analysis of outcomes from studies using wait-list control groups. *Journal of Affective Disorders*, 66(2-3), 139-146.
- 10 Pots WT, Meulenbeek PA, Veehof MM, Klungers J, Bohlmeijer ET. The efficacy of mindfulness-based cognitive therapy as a public mental health intervention for adults with mild to moderate depressive symptomatology: A randomized controlled trial. *PloS one*. 2014 Oct 15; 9(10):e109789.
- 14 Pots WT, Meulenbeek PA, Veehof MM, Klungers J, Bohlmeijer ET. The efficacy of mindfulness-based cognitive therapy as a public mental health intervention for adults with mild to moderate depressive symptomatology: A randomized controlled trial. *PloS one*. 2014 Oct 15;9(10):e109789.
- 18 Pouget, R., Yersin, B., Wietlisbach, V., Burnand, B., Bula, C.J. (2000). Depressed mood in a cohort of elderly medical inpatients: prevalence, clinical correlates and recognition rate. *Aging (Milano)*, 12: 301-307.
- 21 Pradeep J, Isaacs A, Shanbag D, Selvan S, Srinivasan K. Enhanced care by community health workers in improving treatment adherence to antidepressant medication in rural women with major depression. *The Indian journal of medical research*. 2014 Feb; 139(2):236.
- 25 Prescribing & Medicines Team & HSCIC. (2016). Prescription Cost Analysis: England 2015. Retrieved from
- 27 Preskorn SH, Silkey B, Beber J, Dorey C. Antidepressant response and plasma concentrations of fluoxetine. *Annals of Clinical Psychiatry* 1991; 3:147-151.
- 29 Prien, R. F., Kupfer, D. J., Mansky, P. A., Small, J. G., Tuason, V. B., Voss, C. B., & Johnson, W. E. (1984). Drug therapy in the prevention of recurrences in unipolar and bipolar affective disorders: report of the NIMH Collaborative Study Group comparing lithium carbonate, imipramine, and a lithium carbonate-imipramine combination. *Archives of General Psychiatry*, 41(11), 1096-1104.
- 34 Priest, R. G., Vize, C., Roberts, A., Roberts, M., & Tylee, A. (1996). Lay people's attitudes to treatment of depression: results of opinion poll for Defeat Depression Campaign just before its launch. *BMJ*, 313(7061), 858-859.
- 37 Pringle, A., & Harmer, C. J. (2015). The effects of drugs on human models of emotional processing: an account of antidepressant drug treatment. *Dialogues in clinical neuroscience*, 17(4), 477.
- 40 Prins, M. A., Verhaak, P. F., Bensing, J. M., & van der Meer, K. (2008). Health beliefs and perceived need for mental health care of anxiety and depression--the patients' perspective explored. *Clin Psychol Rev*, 28(6), 1038-1058.
- 43 Proudfoot, J., Ryden, C., Everitt, B., Shapiro, D. A., Goldberg, D., Mann, A., . . . Gray, J. A. (2004). Clinical efficacy of computerised cognitive-behavioural therapy for anxiety and depression in primary care: randomised controlled trial. *Br J Psychiatry*, 185, 46-54.

- 1 Pyne, J. M., Fortney, J. C., Mouden, S., Lu, L., Hudson, T. J., & Mittal, D. (2015). Cost-effectiveness of on-site versus off-site collaborative care for depression in rural FQHCs. *Psychiatric Services, 66*, 491-499.
- 4 Qirjazi, E., McArthur, E., Nash, D. M., Dixon, S. N., Weir, M. A., Vasudev, A., . . . Garg, A. X. (2016). Risk of ventricular arrhythmia with citalopram and escitalopram: A population-based study. *PLoS ONE, 11*(8), e0160768.
- 7 Qu, S. S., Huang, Y., Zhang, Z. J., Chen, J. Q., Lin, R. Y., Wang, C. Q., . . . Zhang, Y. C. (2013). A 6-week randomized controlled trial with 4-week follow-up of acupuncture combined with paroxetine in patients with major depressive disorder. *Journal of psychiatric research, 47*(6), 726-732.
- 11 Quah-Smith, I., Smith, C., Crawford, J. D., & Russell, J. (2013). Laser acupuncture for depression: A randomised double blind controlled trial using low intensity laser intervention. *Journal of Affective Disorders, 148*(2-3), 179-187.
- 14 Quinn, G. R., Singer, D. E., Chang, Y., Go, A. S., Borowsky, L. H., Udaltssova, N., & Fang, M. C. (2014). Effect of selective serotonin reuptake inhibitors on bleeding risk in patients with atrial fibrillation taking warfarin. *Am J Cardiol, 114*(4), 583-586.
- 17 Quilty LC, Dozois DJ, Lobo DS, Ravindran LN, Bagby RM. Cognitive structure and processing during cognitive behavioral therapy vs. pharmacotherapy for depression. *International Journal of Cognitive Therapy. 2014 Sep; 7*(3):235-50.
- 20 Quitkin, F. M. (1991). Response to phenelzine and imipramine in placebo nonresponders with atypical depression. *Archives of General Psychiatry, 48*(4), 319.
- 22 Quitkin, F. M., McGrath, P. J., Stewart, J. W., Harrison, W., Wager, S. G., Nunes, E. et al. (1989). Phenelzine and imipramine in mood reactive depressives. Further delineation of the syndrome of atypical depression. *Archives of General Psychiatry, 46*, 787-793.
- 25 Rabins, P. (1996). Barriers to diagnosis and treatment of depression in elderly patients. *American Journal of Geriatric Psychiatry, 4*, S79-S83.
- 27 Radhakrishnan, M., Hammond, G., Jones, P. B., Watson, A., McMillan-Shields, F., & Lafourture, L. (2013). Cost of improving Access to Psychological Therapies (IAPT) programme: an analysis of cost of session, treatment and recovery in selected Primary Care Trusts in the East of England region. *Behav Res Ther, 51*(1), 37-45.
- 31 Radloff, L. S. (1977). The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement, 1*, 385-401.
- 33 Raine, R., Lewis, L., Sensky, T., Hutchings, A., Hirsch, S., & Black, N. (2000). Patient determinants of mental health interventions in primary care. *Br J Gen Pract, 50*(457), 620-625.
- 36 Rait, G., Burns, A., Baldwin, R., Morley, M., Chew-Graham, C., St Leger, A. S., & Abas, M. (1999). Screening for depression in African-Caribbean elders. *Fam Pract, 16*(6), 591-595.
- 38 Ramachandani, P. Stein, A. (2003) The impact of parental psychiatric disorder on children. *BMJ, 327*(7409), 242-243.
- 40 Rapaport, M. H., Bose, A., & Zheng, H. (2004). Escitalopram continuation treatment prevents relapse of depressive episodes. *Journal of Clinical Psychiatry, 65*(1), 44-49.
- 42 Rapaport, M. H., Gharabawi, G. M., Canuso, C. M., Mahmoud, R. A., Keller, M. B., Bossie, C. A., . . . Bouhours, P. (2006). Effects of risperidone augmentation in patients with treatment-resistant depression: results of open-label treatment followed by double-blind continuation. *Neuropsychopharmacology, 31*(11), 2505.

- 1 Rastad, C., Ulfberg, J., & Lindberg, P. (2008). Light room therapy effective in mild forms of seasonal affective disorder—a randomised controlled study. *Journal of Affective Disorders*, 108(3), 291-296.
- 4 Ravindran, A. V., Cameron, C., Bhatla, R., Ravindran, L. N., & Silva, T. L. (2013). Paroxetine in the treatment of dysthymic disorder without co-morbidities: A double-blind, placebo-controlled, flexible-dose study. *Asian Journal of Psychiatry*, 6(2), 157-161.
- 7 Ravindran, L. N., Eisfeld, B. S., & Kennedy, S. H. (2008). Combining mirtazapine and duloxetine in treatment-resistant depression improves outcomes and sexual function. *Journal of Clinical Psychopharmacology*, 28(1), 107-108.
- 10 Ravindran AV, Guelfi JD, Lane RM, Cassano GB. Treatment of dysthymia with sertraline: a double-blind, placebo-controlled trial in dysthymic patients without major depression. *The Journal of clinical psychiatry*. 2000 Nov; 61 (11):821-7.
- 13 Ravindram AV, Teehan MD, Bakish D, Yatham L, O'Reilly R, Fernando ML, Manchanda R, Charbonneau Y, Buttars J. The impact of sertraline, desipramine, and placebo on psychomotor functioning in depression. *Hum Psychopharmacol* 1995; 10(4):273-281.
- 16 Ravizza L. Amisulpride in medium-term treatment of dysthymia: a six-month, double-blind safety study versus amitriptyline. *Journal of Psychopharmacology*. 1999 May; 13 (3):248-54.
- 18 Reeves, G. M., Postolache, T. T., & Snitker, S. (2008). Childhood obesity and depression: connection between these growing problems in growing children. *International journal of child health and human development: IJCHD*, 1(2), 103.
- 21 Reichstadt, J., Sengupta, G., Depp, C. A., Palinkas, L. A., & Jeste, D. V. (2010). Older adults' perspectives on successful aging: Qualitative interviews. *The American Journal of Geriatric Psychiatry*, 18(7), 567-575.
- 24 Reimherr, F., Byerley, W., Ward, M., Lebegue, B., & Wender, P. (1988). Sertraline, a selective inhibitor of serotonin uptake, for the treatment of outpatients with major depressive disorder. *Psychopharmacology Bulletin*, 24(1), 200-205.
- 27 Reimherr FW, Amsterdam JD, Quitkin FM, Rosenbaum JF, Fava M, Zajecka J, Beasley CM, Jr., Michelson D, Roback P, Sundell K. Optimal length of continuation therapy in depression: a prospective assessment during long-term fluoxetine treatment.[comment]. *American Journal of Psychiatry* 1998; 155(9):1247-1253.
- 31 Reimherr FW, Chouinard G, Cohn CK, Cole JO, Itil TM, LaPierre YD, Masco HL - Mendels J. Antidepressant efficacy of sertraline: a double-blind, placebo- and amitriptyline-controlled, multicenter comparison study in outpatients with major depression [see comments]. *Journal of Clinical Psychiatry* 1990 Dec; 51 Suppl B:18-27.
- 35 Reynolds III, C. F., Dew, M. A., Pollock, B. G., Mulsant, B. H., Frank, E., Miller, M. D., . . . Stack, J. A. (2006). Maintenance treatment of major depression in old age. *New England Journal of Medicine*, 354(11), 1130-1138.
- 38 Reynolds, C., Dew, M., Martire, L., Miller, M., Curanowski, J., Lenze, E., Whyte, E., Mulsant, B. et al (2010) Treating depression to remission in older adults: a controlled evaluation of combined escitalopram with interpersonal psychotherapy verus escitalopram with depression care management, *International Journal of Geriatric Psychiatry*, 25, 1134-1141
- 42 Reynolds III, C. F., Frank, E., Perel, J. M., Imber, S. D., Cornes, C., Miller, M. D., . . . Stack, J. A. (1999). Nortriptyline and interpersonal psychotherapy as maintenance therapies for recurrent major depression: a randomized controlled trial in patients older than 59 years. *JAMA*, 281(1), 39-45.

- 1 Richards, D. A., Bower, P., Chew-Graham, C., Gask, L., Lovell, K., Cape, J., ... Russell, A. (2016). Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): A cluster randomised controlled trial. *Health Technology Assessment*, 20, 1-192.
- 5 Richards, D., et al. (2015). A randomized controlled trial of an internet-delivered treatment: Its potential as a low-intensity community intervention for adults with symptoms of depression. *Behaviour research and therapy* 75: 20-31.
- 8 Richards, D. A., Hill, J. J., Gask, L., Lovell, K., Chew-Graham, C., Bower, P., ... Barkham, M. (2013). Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. *BMJ*, 347, f4913.
- 11 Richards DA, Lovell K, Gilbody S, Gask L, Torgerson, D, Barkham M, et al. Collaborative care for depression in UK primary care: a randomized controlled trial. *Psychological Medicine* 2008; 38(2):279–87.
- 14 Rickels, K., Feighner, J. P., & Smith, W. T. (1985). Alprazolam, amitriptyline, doxepin, and placebo in the treatment of depression. *Archives of General Psychiatry*, 42(2), 134-141.
- 16 Rickels, K., Montgomery, S. A., Tourian, K. A., Guelfi, J. D., Pitrosky, B., Padmanabhan, S. K., . . . Brisard, C. (2010). Desvenlafaxine for the prevention of relapse in major depressive disorder: results of a randomized trial. *Journal of Clinical Psychopharmacology*, 30(1), 18-24.
- 19 Rickels, K., Case, W. G., Werblowsky, J., Csanalosi, I., Schless, A., & Weise, C. C. (1981). Amoxapine and imipramine in the treatment of depressed outpatients: a controlled study. *American Journal of Psychiatry*, 138, 20-24.
- 22 Rickels K, Schweizer E, Clary C, Fox I, Weise C. Nefazodone and imipramine in major depression: a placebo-controlled trial. *British Journal of Psychiatry* 1994 Jun; 164(6):802-5.
- 24 Ricken, R., Wiethoff, K., Reinhold, T., Schietsch, K., Stamm, T., Kiermeir, J., ... Adli, M. (2011). Algorithm-guided treatment of depression reduces treatment costs--results from the randomized controlled German Algorithm Project (GAPII). *Journal of Affective Disorders*, 134(1-3), 249-256.
- 28 Rickles, N. M., Svarstad, B. L., Statz-Paynter, J. L., Taylor, L. V., & Kobak, K. A. (2005). Pharmacist telemonitoring of antidepressant use: effects on pharmacist-patient collaboration. *Journal of the American Pharmacists Association*, 45, 344-353.
- 31 Ridge, D., & Ziebland, S. (2006). "The old me could never have done that": how people give meaning to recovery following depression. *Qual Health Res*, 16(8), 1038-1053.
- 33 Rieu, J., et al. (2011) Efficacy of ultrabrief cognitive and behavioural therapy performed by psychiatric residents on depressed inpatients. *Psychotherapy and Psychosomatics* 80, 374-376.
- 36 Riihimaki, K. A., Vuorilehto, M. S., Melartin, T. K., & Isometsa, E. T. (2014). Five-year outcome of major depressive disorder in primary health care. *Psychol Med*, 44(7), 1369-1379.
- 39 Rinaldi, M., Perkins, R., Glynn, E., Montibeller, T., Clenaghan, M., & Rutherford, J. (2007). Individual placement and support: from research to practice. *Advances in Psychiatric Treatment*, 14(1), 50.
- 42 Robert, P., Montgomery, S., Darcourt, G., Robert, G., Houillon, P., Bouchard, J., ... Richou, H. (1995). Citalopram in doses of 20-60 mg is effective in depression relapse prevention: a placebo-controlled 6 month study. *International Clinical Psychopharmacology*, 10, 29-36.

- 1 Roberts, D., Bernard, M., Misca, G., & Head, E. (2008). Experiences of children and young people caring for a parent with a mental health problem. London: Social Care Institute for Excellence, (Research Briefings).
- 4 Robertson MM, Abou Saleh MT, Harrison DA, Nairac BL, Edwards DRL, Lock T, Burns RA, Katona CLE. A double-blind controlled comparison of fluoxetine and lofepramine in major depressive illness. *J Psychopharmacol*, 1994;8(2):98-103.
- 7 Robinson, D. (1991). Antidepressant efficacy of 5-HT 1A partial agonist drugs. Sandler M, Coppen A, Harnett S (eds), 158-165.
- 9 Rocca P, Fonzo V, Ravizza L, Rocca G, Scotta M, Zanalda E, Bogetto F. A comparison of paroxetine and amisulpride in the treatment of dysthymic disorder. *Journal of affective disorders*. 2002 Aug 31; 70(3):313-7.
- 12 Rocca P, Marchiaro L, Rasetti R, Rivoira E, Bogetto F. A comparison of paroxetine versus paroxetine plus amisulpride in the treatment of dysthymic disorder: efficacy and psychosocial outcomes. *Psychiatry Res*. 2002; 112(2):145–152
- 15 Rodin, I., & Thompson, C. (1997). Seasonal affective disorder. *Advances in Psychiatric Treatment*, 3(6), 352-359.
- 17 Roffman, M., Gould, E. E., Brewer, S. J., et al. (1982). A double-blind comparative study of oxaprotiline with amitriptyline and placebo in moderate depression. *Current Therapeutic Research, Clinical & Experimental*, 32, 247-256.
- 20 Rogers, C. M. D. O. A. (2001). Experiencing depression, experiencing the depressed: The separate worlds of patients and doctors. *Journal of Mental Health*, 10(3), 317-333.
- 22 Rogers, C. R. (1957). The necessary and sufficient conditions of therapeutic personality change. *J Consult Psychol*, 21(2), 95-103.
- 24 Rohan, K. J., Lindsey, K. T., Roecklein, K. A., & Lacy, T. J. (2004). Cognitive-behavioral therapy, light therapy, and their combination in treating seasonal affective disorder. *Journal of Affective Disorders*, 80(2), 273-283.
- 27 Rohan, K. J., Roecklein, K. A., Tierney Lindsey, K., Johnson, L. G., Lippy, R. D., Lacy, T. J., & Barton, F. B. (2007). A randomized controlled trial of cognitive-behavioral therapy, light therapy, and their combination for seasonal affective disorder. *Journal of Consulting and Clinical Psychology*, 75(3), 489.
- 31 Röhricht F, Papadopoulos N, Priebe S. An exploratory randomized controlled trial of body psychotherapy for patients with chronic depression. *Journal of affective disorders*. 2013 Oct 31; 151(1):85-91.
- 34 Roose SP. Antidepressant Treatment of Melancholia in Late life [NCT00000378]. Available from: <https://clinicaltrials.gov/ct2/show/NCT00000378> [accessed 26.09.2017]
- 36 Rose, D., Fleischmann, P., Wykes, T., Leese, M., & Bindman, J. (2003). Patients' perspectives on electroconvulsive therapy: systematic review. *BMJ*, 326(7403), 1363.
- 38 Rosenberg C, Damsbo N, Fuglum E, Jacobsen LV, Horsgard S. Citalopram and imipramine in the treatment of depressive patients in general practice. A Nordic multicentre clinical study. *International Clinical Psychopharmacology* 1994 Mar;9 Suppl 1:41-8.
- 41 Rosenthal, J. Z., Boyer, P., Vialet, C., Hwang, E., & Tourian, K. A. (2013). Efficacy and safety of desvenlafaxine 50 mg/d for prevention of relapse in major depressive disorder: a randomized controlled trial. *The Journal of Clinical Psychiatry*, 74(2), 158-166.
- 44 Rosenthal, N. E. (1984). Seasonal Affective Disorder. *Archives of General Psychiatry*, 41(1), 72.

- 1 Rosenthal, N. E., Moul, D. E., Hellekson, C. J., Oren, D. A., Frank, A., Brainard, G. C., . . .
- 2 Wehr, T. A. (1993). A multicenter study of the light visor for seasonal affective disorder: no difference in efficacy found between two different intensities. *Neuropsychopharmacology*, 8(2), 151-160.
- 5 Rosenthal, N. E., Sack, D. A., Gillin, J. C., Lewy, A. J., Goodwin, F. K., Davenport, Y., . . .
- 6 Wehr, T. A. (1984). Seasonal affective disorder. A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry*, 41(1), 72-80.
- 8 Ross, J. T., TenHave, T., Eakin, A. C., Difilippo, S., & Oslin, D. W. (2008). A randomized controlled trial of a close monitoring program for minor depression and distress. *J Gen Intern Med*, 23(9), 1379-1385.
- 11 Rosso G, Martini B, Maina G. Brief dynamic therapy and depression severity: A single-blind, randomized study. *Journal of affective disorders*. 2013 May 31; 147(1):101-6.
- 13 Rost, K., Humphrey, J., & Kelleher, K. (1994). Physician management preferences and barriers to care for rural patients with depression. *Arch Fam Med*, 3(5), 409-414.
- 15 Rost, K., Nutting, P., Smith, J., Elliott, C., & Dickinson, M. (2002). Managing depression as a chronic disease: a randomised trial of ongoing treatment in primary care. *British Medical Journal*, 325, 934.
- 18 Rost, K., Nutting, P., Smith, J., Werner, J., & Duan, N. (2001). Improving depression outcomes in community primary care practices: a randomized trial of the QuEST intervention. *Journal of General Internal Medicine*, 16, 143-149.
- 21 Rost, K., Williams, C., Wherry, J., & Jr, G. R. S. (1995). The process and outcomes of care for major depression in rural family practice settings. *The Journal of Rural Health*, 11(2), 114-121.
- 24 Rotella, F., & Mannucci, E. (2013). Depression as a risk factor for diabetes: a meta-analysis of longitudinal studies. *J Clin Psychiatry*, 74(1), 31-37.
- 26 Roter, D. L., Hall, J. A., Kern, D. E., Barker, L. R., Cole, K. A., & Roca, R. P. (1995). Improving physicians' interviewing skills and reducing patients' emotional distress. A randomized clinical trial. *Arch Intern Med*, 155(17), 1877-1884.
- 29 Roth, A., & Fonagy, P. (2005). What works for whom?: a critical review of psychotherapy research (2nd ed.). New York: Guilford Press.
- 31 Rothschild, A. J. (2003). Challenges in the treatment of depression with psychotic features. *Biol Psychiatry*, 53(8), 680-690.
- 33 Rothschild, A. J., Williamson, D. J., Tohen, M. F., Schatzberg, A., Andersen, S. W., Van Campen, L. E., ... Tollefson, G. D. (2004). A double-blind, randomized study of olanzapine and olanzapine/fluoxetine combination for major depression with psychotic features. *J Clin Psychopharmacol*, 24(4), 365-373.
- 37 Rothschild, A. J., Winer, J., Flint, A. J., Mulsant, B. H., Whyte, E. M., Heo, M., Fratoni, S., Gabriele, M., Kasapinovic, S., Meyers, B.S.; Study of Pharmacotherapy of Psychotic Depression (STOP-PD) Collaborative Study Group. (2008). Missed diagnosis of psychotic depression at 4 academic medical centers. *J Clin Psychiatry*, 69(8), 1293-1296.
- 41 Rouillon, F., Serrurier, D., Miller, H. D., & Gerard, M.-J. (1991). Prophylactic efficacy of maprotiline on unipolar depression relapse. *The Journal of Clinical Psychiatry*, 52(10):423-31.
- 44 Rowe, S. K., & Rapaport, M. H. (2006). Classification and treatment of sub-threshold depression. *Current Opinion in Psychiatry*, 19(1), 9-13.

- 1 Royal College of Psychiatrists. (2008). Mental health and work.
- 2 Rubenstein, L., Parker, L., Meredith, L., Altschuler, A., dePillis, E., Hernandez, J., Gordon, N.
- 3 P. (2002). Understanding team-based quality improvement for depression in primary care.
- 4 Health Services Research, 37, 1009–1029.
- 5 Rubio-Valera, M., Bosmans, J., Fernandez, A., Penarrubia-Maria, M., March, M., Trave,
- 6 P.,... Serrano-Blanco, A. (2013). Cost-Effectiveness of a Community Pharmacist Intervention
- 7 in Patients with Depression: A Randomized Controlled Trial (PRODEFAR Study). PLoS
- 8 ONE, 8(8), e70588.
- 9 Rudolph RL, Feiger AD. A double-blind, randomized, placebo-controlled trial of once-daily
- 10 venlafaxine extended release (XR) and fluoxetine for the treatment of depression. Journal of
- 11 Affective Disorders, 1999; 56(2-3):171-181.
- 12 Ruhé, H. G., Booij, J., Weert, H. C., Reitsma, J. B., Fransen, E. J., Michel, M. C., & Schene,
- 13 A. H. (2009). Evidence why paroxetine dose escalation is not effective in major depressive
- 14 disorder: a randomized controlled trial with assessment of serotonin transporter occupancy.
- 15 Neuropsychopharmacology, 34(4), 999.
- 16 Ruhe, H. G., van Rooijen, G., Spijker, J., Peeters, F. P., & Schene, A. H. (2012). Staging
- 17 methods for treatment resistant depression. A systematic review. J Affect Disord, 137(1-3),
- 18 35-45.
- 19 Ruscio, A. M., Stein, D. J., Chiu, W. T., & Kessler, R. C. (2010). The epidemiology of
- 20 obsessive-compulsive disorder in the National Comorbidity Survey Replication. Mol
- 21 Psychiatry, 15(1), 53-63.
- 22 Rush, A., Beck, A., Kovacs, M. & Hollon, S. (1977). Comparative efficacy of cognitive therapy
- 23 and pharmacotherapy in the treatment of depressed outpatients. Cognitive Therapy and
- 24 Research 1, 17-37.
- 25 Rush, A. J., Trivedi, M. H., Wisniewski, S. R., Nierenberg, A. A., Stewart, J. W., Warden, D.,
- 26 ... Fava, M. (2006). Acute and longer-term outcomes in depressed outpatients requiring one
- 27 or several treatment steps: a STAR*D report. Am J Psychiatry, 163(11), 1905-1917.
- 28 Rush, A. J., Trivedi, M. H., Wisniewski, S. R., Stewart, J. W., Nierenberg, A. A., Thase, M. E.,
- 29 ... Luther, J. F. (2006). Bupropion-SR, sertraline, or venlafaxine-XR after failure of SSRIs for
- 30 depression. New England Journal of Medicine, 354(12), 1231-1242.
- 31 Ruwaard, J., Schrieken, B., Schrijver, M., Broeksteeg, J., Dekker, J., Vermeulen, H., &
- 32 Lange, A. (2009). Standardized web-based cognitive behavioural therapy of mild to moderate
- 33 depression: a randomized controlled trial with a long-term follow-up. Cognitive Behaviour
- 34 Therapy, 38(4), 206-221.
- 35 Sackeim, H. A., Haskett, R. F., Mulsant, B. H., Thase, M. E., Mann, J. J., Pettinati, H. M., . . .
- 36 Prudic, J. (2001). Continuation pharmacotherapy in the prevention of relapse following
- 37 electroconvulsive therapy: a randomized controlled trial. JAMA, 285(10), 1299-1307.
- 38 Sainsbury Centre for Mental Health. (2007). Mental health at work: Developing the business
- 39 case.
- 40 Salkovskis, P., Rimes, K., Stephenson, D., Sacks, G., & Scott, J. (2006). A randomized
- 41 controlled trial of the use of self-help materials in addition to standard general practice
- 42 treatment of depression compared to standard treatment alone. Psychological Medicine,
- 43 36(3), 325-333.
- 44 Salminen, J.K., Karlsson, H., Hietala, J., et al. (2008) Short-term psychodynamic
- 45 psychotherapy and fluoxetine in major depressive disorder: A randomized comparative
- 46 study. Psychotherapy and Psychosomatics, 77, 351-357.

- 1 Sanderson, K., Andrews, G., Corry, J., & Lapsley, H. (2003). Reducing the burden of affective disorders: is evidence-based health care affordable? *Journal of Affective Disorders*, 77(2), 109-125.
- 4 Santos, M. A., Rocha, F. L., & Hara, C. (2008). Efficacy and safety of antidepressant augmentation with lamotrigine in patients with treatment-resistant depression: a randomized, placebo-controlled, double-blind study. *Primary care companion to the Journal of clinical psychiatry*, 10(3), 187.
- 8 Sapin, C., Fantino, B., Nowicki, M. L., & Kind, P. (2004). Usefulness of EQ-5D in assessing health status in primary care patients with major depressive disorder. *Health and Quality of Life Outcomes*, 2(20).
- 11 Sartorius, N. (2001). The economic and social burden of depression. *J Clin Psychiatry*, 62 Suppl 15, 8-11.
- 13 Saver, B. G., Van-Nguyen, V., Keppel, G., & Doescher, M. P. (2007). A qualitative study of depression in primary care: missed opportunities for diagnosis and education. *J Am Board Fam Med*, 20(1), 28-35.
- 16 Schatzberg, A. F., Blier, P., Delgado, P. L., Fava, M., Haddad, P. M., & Shelton, R. C. (2006). Antidepressant discontinuation syndrome: consensus panel recommendations for clinical management and additional research. *J Clin Psychiatry*, 67 Suppl 4, 27-30.
- 19 Schene, A., Wijngaarden, B. v., Poelijoe, N., & Gersons, B. (1993). The Utrecht comparative study on psychiatric day treatment and inpatient treatment. *Acta Psychiatrica Scandinavica*, 87(6), 427-436.
- 22 Schindler, F., & Anhelescu, I. G. (2007). Lithium versus lamotrigine augmentation in treatment resistant unipolar depression: a randomized, open-label study. *International Clinical Psychopharmacology*, 22(3), 179-182.
- 25 Schlager, D. S. (1994). Early-morning administration of short-acting B blockers for treatment of winter depression. *American Journal of Psychiatry*, 151(9), 1383-1385.
- 27 Schlogelhofer, M., Willinger, U., Wiesegger, G., Eder, H., Priesch, M., & Itzlinger, U. (2014). Clinical study results from a randomized controlled trial of cognitive behavioural guided self-help in patients with partially remitted depressive disorder. *Psychology and Psychotherapy: Theory, Research and Practice*, 87(2), 178-190.
- 31 Schmidt MM, & Miller WR. Amount of therapist contact and outcome in a multidimensional depression treatment program. *Acta Psychiatrica Scandinavica* 1983; 67(5):319-332.
- 33 Schmidt, M. E., Fava, M., Robinson, J. M., & Judge, R. (2000). The efficacy and safety of a new enteric-coated formulation of fluoxetine given once weekly during the continuation treatment of major depressive disorder. *The Journal of Clinical Psychiatry*, 61(11), 851-857.
- 36 Schneider LS, Nelson JC, Clary CM, Newhouse P, Krishnan KRR, Shiovitz T, Weihs K. An 8-week multicenter, parallel-group, double-blind, placebo-controlled study of sertraline in elderly outpatients with major depression. *American Journal of Psychiatry*. Vol 160(7) 1277-1285 2003.
- 40 Schrader, E. (2000). Equivalence of St John's wort extract (Ze 117) and fluoxetine: a randomized, controlled study in mild/moderate depression. *International Clinical Psychopharmacology*, 15(2), 61-68.
- 43 Schrader, E., Meier, B., & Brattstrom, A. (1998). Hypericum treatment of mild-moderate depression in a placebo-controlled study. A prospective, double-blind, randomized, placebo-controlled, multicentre study. *Human Psychopharmacology-Clinical and Experimental*, 13(3), 163-169.

- 1 E, Dietrich van Calker MD, Dykierk P, Lieb K, Kech S, Zobel I, Leonhart R, Berger M. An intensive treatment program of interpersonal psychotherapy plus pharmacotherapy for depressed inpatients: acute and long-term results. *American Journal of Psychiatry*. 2007 May 1.
- 5 Schramm, E., Schneider, D., Zobel, I., van Calker, D., Dykierk, P., Kech, S., ... Berger, M. (2008). Efficacy of interpersonal psychotherapy plus pharmacotherapy in chronically depressed inpatients. *Journal of Affective Disorders*, 109(1), 65-73.
- 8 Schramm, E., Zobel, I., Dykierk, P., Kech, S., Brakemeier, E. L., Külz, A., & Berger, M. (2011). Cognitive behavioral analysis system of psychotherapy versus interpersonal psychotherapy for early-onset chronic depression: a randomized pilot study. *Journal of Affective Disorders*, 129(1-3), 109-116.
- 12 Schramm, E., Zobel, I., Schoepf, D., Fangmeier, T., Schnell, K., Walter, H., ... Normann, C. (2015). Cognitive behavioral analysis system of psychotherapy versus escitalopram in chronic major depression. *Psychotherapy and psychosomatics*, 84(4), 227-240.
- 15 Schuch, F. B., Vasconcelos-Moreno, M. P., Borowsky, C., Zimmermann, A. B., Rocha, N. S., & Fleck, M. P. (2015). Exercise and severe major depression: effect on symptom severity and quality of life at discharge in an inpatient cohort. *Journal of psychiatric research*, 61, 25-32.
- 19 Schulberg, H. C., Block, M. R., Madonia, M. J., Scott, C. P., Rodriguez, E., Imber, S. D., ... Coulehan, J. L. (1996). Treating major depression in primary care practice: eight-month clinical outcomes. *Archives of General Psychiatry*, 53(10), 913-919.
- 22 Schulte, S., & Holland, M. (2008). Dual diagnosis in Manchester, UK: practitioners' estimates of prevalence rates in mental health and substance misuse services. *Mental Health and Substance Use*, 1(2), 118-124.
- 25 Schünemann, H. J., Best, D., Vist, G., Oxman, A., & Group, G. W. (2003). Letters, numbers, symbols and words: how to communicate grades of evidence and recommendations. . Canadian Medical Association Journal, 169, 677-680.
- 28 Schuver KJ & Lewis BA. Mindfulness-based yoga intervention for women with depression. Complementary therapies in medicine. 2016 Jun 30; 26:85-91.
- 30 Schweizer, E., Feighner, J., Mandos, L. A., & Rickels, K. (1994). Comparison of venlafaxine and imipramine in the acute treatment of major depression in outpatients. *Journal of Clinical Psychiatry*, 55, 104-108.
- 33 Schweizer E, Rickels K, Amsterdam JD, Fox I, Puzzioli G, Weise C (1990) What constitutes an adequate antidepressant trial for fluoxetine? *J Clin Psychiatry* 51:8-11
- 35 Schweizer, E., Rickels, K., Hassman, H., & Garcia-Espana, F. (1998). Buspirone and imipramine for the treatment of major depression in the elderly. *Journal of Clinical Psychiatry*, 59, 175-183.
- 38 Schweizer, E., Rynn, M., Mandos, L., Demartinis, N., Garcia-Espana, F., & Rickels, K. (2001). The antidepressant effect of sertraline is not enhanced by dose titration: results from an outpatient clinical trial. *International Clinical Psychopharmacology*, 16(3), 137-143.
- 41 Scogin, F. (1987). The concurrent validity of the Geriatric Depression Scale with depressed older adults. *Clinical Gerontologist*, 7(1), 23-32.
- 43 Scogin F, Jamison C, Gochneaur K. Comparative efficacy of cognitive and behavioral bibliotherapy for mildly and moderately depressed older adults. *Journal of consulting and clinical psychology*. 1989 Jun 1; 57(3):403-7.

- 1 Scott, A., & Freeman, C. (1992). Edinburgh primary care depression study: treatment outcome, patient satisfaction, and cost after 16 weeks. *BMJ*, 304(6831), 883-887.
- 3 Scott, J., Palmer, S., Paykel, E., Teasdale, J., & Hayhurst, H. (2003). Use of cognitive therapy for relapse prevention in chronic depression: Cost-effectiveness study. *British Journal of Psychiatry*, 182, 221-227.
- 6 Scott C, Tacchi MJ, Jones R, Scott J. Acute and one-year outcome of a randomised controlled trial of brief cognitive therapy for major depressive disorder in primary care. *British Journal of Psychiatry* 1997; 171:131-134.
- 9 Scott, J., Teasdale, J. D., Paykel, E. S., Johnson, A. L., Abbott, R., Hayhurst, H., ... Garland, A. (2000). Effects of cognitive therapy on psychological symptoms and social functioning in residual depression. *The British Journal of Psychiatry*, 177(5), 440-446.
- 12 Segal, Z. V., Bieling, P., Young, T., MacQueen, G., Cooke, R., Martin, L., ... Levitan, R. D. (2010). Antidepressant monotherapy vs sequential pharmacotherapy and mindfulness-based cognitive therapy, or placebo, for relapse prophylaxis in recurrent depression. *Archives of General Psychiatry*, 67(12), 1256-1264.
- 16 Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). Mindfulness-based cognitive therapy for depression: a new approach to preventing relapse. New York: Guilford Press.
- 18 Segrin, C. (2000). Social skills deficits associated with depression. *Clinical Psychology Review*, 20(3), 379-403.
- 20 Seidler, Z. E., Dawes, A. J., Rice, S. M., Oliffe, J. L., & Dhillon, H. M. (2016). The role of masculinity in men's help-seeking for depression: A systematic review. *Clin Psychol Rev*, 49, 106-118.
- 23 Selmi, P. M., Klein, M. H., Greist, J. H., Sorrell, S. P., & Erdman, H. P. (1990). Computer-administered cognitive-behavioral therapy for depression. *Am J Psychiatry*, 147(1), 51-56.
- 25 Serby, M., & Yu, M. (2003). Overview: depression in the elderly. *Mt Sinai J Med*, 70(1), 38-44.
- 27 Serretti A, Fabbri C, Pellegrini S, Porcelli S, Politi P, Bellino S, Menchetti M, Mariotti V, Demi C, Martinelli V, Cappucciati M. No effect of serotonergic gene variants on response to interpersonal counseling and antidepressants in major depression. *Psychiatry investigation*. 2013 Jun 1; 10(2):180-9.
- 31 Serfaty, M. A., Haworth, D., Blanchard, M., Buszewicz, M., Murad, S., & King, M. (2009). Clinical effectiveness of individual cognitive behavioral therapy for depressed older people in primary care: a randomized controlled trial. *Archives of General Psychiatry*, 66(12), 1332-1340.
- 35 Shallcross, A., Visvanathan, P., Ford, B., Shirk, S., Holm-Denoma, J., Goode, K., Cox, E., Gross, J., Kumar, N., Palfrey, A., Dimidjian, S., Chaplin, W., Mauss, I. (2015) Relapse prevention in major depressive disorder: mindfulness-based cognitive therapy versus an active control condition, *Journal of Consulting and Clinical Psychology*, 83, 964-975
- 39 Shamsaei F, Rahimi A, Zarabian MK et al. Efficacy of pharmacotherapy and cognitive therapy, alone and in combination in major depressive disorder. *Hong Kong J Psychiatry* 2008; 18:76-80.
- 42 Shapira, B., Zislin, J., Gelfin, Y., Osher, Y., Gorfine, M., Souery, D., ... Lerer, B. (1999). Social adjustment and self-esteem in remitted patients with unipolar and bipolar affective disorder: A case-control study. *Comprehensive Psychiatry*, 40(1), 24-30.

- 1 Shaw, C. M., Creed, F., Tomenson, B., Riste, L., & Cruickshank, J. K. (1999). Prevalence of anxiety and depressive illness and help seeking behaviour in African Caribbeans and white Europeans: two phase general population survey. *BMJ*, 318(7179), 302-305.
- 4 Sheeber LB, Seeley JR, Feil EG, Davis B, Sorensen E, Kosty DB, Lewinsohn PM.
- 5 Development and pilot evaluation of an Internet-facilitated cognitive-behavioral intervention
- 6 for maternal depression. *Journal of consulting and clinical psychology*. 2012 Oct; 80(5):739.
- 7 Sheehan, D. V., et al. (2009) Placebo-controlled inpatient comparison of venlafaxine and fluoxetine for the treatment of major depression with melancholic features. *International clinical psychopharmacology* 24, 61-86.
- 10 Shelton, C., Entsuah, R., Padmanabhan, S. K., & Vinall, P. E. (2005). Venlafaxine XR demonstrates higher rates of sustained remission compared to fluoxetine, paroxetine or placebo. *International Clinical Psychopharmacology*, 20(4), 233-238.
- 13 Shelton, R. C., Keller, M. B., Gelenberg, A., Dunner, D. L., Hirschfeld, R., Thase, M. E., ... Halbreich, U. (2001). Effectiveness of St John's wort in major depression: a randomized controlled trial. *JAMA*, 285(15), 1978-1986.
- 16 Shepard, M., Crow, T., Lader, M., & Johnson, A. (1981). Continuation therapy with lithium and amitriptyline in unipolar depressive illness: A controlled clinical trial. *Psychological Medicine*.
- 19 Shiels, C., Gabbay, M., Dowrick, C., & Hulbert, C. (2004). Depression in men attending a rural general practice: factors associated with prevalence of depressive symptoms and diagnosis. *Br J Psychiatry*, 185, 239-244.
- 22 Shorter, E., & Healy, D. (2007). Shock therapy: a history of electroconvulsive treatment in mental illness. New Brunswick, N.J.: Rutgers University Press.
- 24 Silverstone PH, Ravindran A. Once-daily venlafaxine extended release (XR) compared with fluoxetine in outpatients with depression and anxiety. *Journal of Clinical Psychiatry* 1999; 60(1):22-28.
- 27 Silverstone, T., bou-Saleh, M. T., Pathak, R., Lock, T., Bennie, E., Mohammed, Y. et al. (1994). A multicentre comparative trial of moclobemide, imipramine and placebo in major depressive disorder. UK moclobemide study group. *International Clinical Psychopharmacology*, 9, 109-113.
- 31 Simon, G., Ludman, E., Tutty, S., Operksalski, B., & Von Korff, M. (2004). Telephone psychotherapy and telephone care management for primary care patients starting antidepressant treatment: a randomized controlled trial. *Journal of the American Medical Association*, 292, 935-942.
- 35 Simon, G., VonKorff, M., Rutter, C., & Wagner, E. (2000). Randomised trial of monitoring, feedback, and management of care by telephone to improve treatment of depression in primary care. *British Medical Journal*, 320, 550–554.
- 38 Simon, G. E., Goldberg, D., Tiemens, B. G., & Ustun, T. B. (1999). Outcomes of recognized and unrecognized depression in an international primary care study. *Gen Hosp Psychiatry*, 21(2), 97-105.
- 41 Simon, G. E., Goldberg, D. P., Von Korff, M., & ÜStÜN, T. B. (2002). Understanding cross-national differences in depression prevalence. *Psychological Medicine*, 32(04).
- 43 Simon, G. E., Ralston, J. D., Savarino, J., Pabiniak, C., Wentzel, C., & Operksalski, B. H. (2011). Randomized trial of depression follow-up care by online messaging. *Journal of General Internal Medicine*, 26(7), 698-704.

- 1 Simon, G. E., Von Korff, M., Ludman, E. J., Katon, W. J., Rutter, C., Unutzer, J., ... Walker, E. (2002). Cost-effectiveness of a program to prevent depression relapse in primary care. *Medical care*, 40, 941-950.
- 4 Simon, J., Pilling, S., Burbeck, R., & Goldberg, D. (2006). Treatment options in moderate and severe depression: Decision analysis supporting a clinical guideline. *British Journal of Psychiatry*, 189, 494-501.
- 7 Simon, J. S., Aguiar, L. M., Kunz, N. R., & Lei, D. (2004). Extended-release venlafaxine in relapse prevention for patients with major depressive disorder. *Journal of psychiatric research*, 38(3), 249-257.
- 10 Simpson, S., Corney, R., & Beecham, J. (2003). A randomized controlled trial to evaluate the effectiveness and cost-effectiveness of psychodynamic counselling for general practice patients with chronic depression. *Psychological Medicine*, 33, 229-239.
- 13 Sims J, Hill K, Davidson S, Gunn J, Huang N. Exploring the feasibility of a community-based strength training program for older people with depressive symptoms and its impact on depressive symptoms. *BMC geriatrics*. 2006 Nov 30; 6(1):1.
- 16 Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 1997 Jan 1; 52(1):M27-35.
- 19 Singh NA, Stavrinos TM, Scarbek Y, Galambos G, Liber C, Singh MA. A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2005 Jun 1; 60(6):768-76.
- 23 Singh, G., Verma, H., Verma, R., & Kaur, H. (1974). A new depressive inventory. *Indian Journal of Psychiatry*, 161, 83-188.
- 25 Singleton, N., Bumpstead, R., & O'Brien, M. (2001). Psychiatric morbidity among adults living in private households, 2000: The Stationery Office.
- 27 Singleton, N., Meltzer, H., & Jenkins, R. (2003). Building a picture of psychiatric morbidity in a nation: a decade of epidemiological surveys in Great Britain. *International Review of Psychiatry*, 15(1-2), 19-28.
- 30 Skodol, A. E., Grilo, C. M., Keyes, K. M., Geier, T., Grant, B. F., & Hasin, D. S. (2011). Relationship of personality disorders to the course of major depressive disorder in a nationally representative sample. *Am J Psychiatry*, 168(3), 257-264.
- 33 Skodol, A. E., Oldham, J. M., & Gallaher, P. E. (1999). Axis II comorbidity of substance use disorders among patients referred for treatment of personality disorders. *Am J Psychiatry*, 156(5), 733-738.
- 36 Smeraldi E. Amisulpride versus fluoxetine in patients with dysthymia or major depression in partial remission: a double-blind, comparative study. *Journal of Affective Disorders*. 1998 Feb 1; 48(1):47-56.
- 39 Smith, W. T., Glaudin, V., Panagides, J., & Gilvary, E. (1990). Mirtazapine vs. amitriptyline vs. placebo in the treatment of major depressive disorder. *Psychopharmacology Bulletin*, 26, 191-196.
- 42 Smith, C. A., Hay, P. P., & Macpherson, H. (2010). Acupuncture for depression. *Cochrane Database Syst Rev* (1), CD004046.

- 1 Smith, D., Dempster, C., Glanville, J., Freemantle, N., & Anderson, I. (2002). Efficacy and tolerability of venlafaxine compared with selective serotonin reuptake inhibitors and other antidepressants: a meta-analysis. *Br J Psychiatry*, 180, 396-404.
- 4 Sneed JR, Reinlieb ME, Rutherford BR, Miyazaki M, Fitzsimons L, Turret N, Pelton GH, Devanand DP, Sackeim HA, Roose SP. Antidepressant treatment of melancholia in older adults. *The American Journal of Geriatric Psychiatry*. 2014 Jan 31; 22(1):46-52.
- 7 Sabin, C., Sackeim, H. A., Prudic, J., Devanand, D. P., Moody, B. J., & McElhiney, M. C. (1995). Predictors of retrograde amnesia following ECT. *Am J Psychiatry*, 152(7), 995-1001.
- 9 Sobocki, P., Ekman, M., Agren, H., Krakau, I., Runeson, B., Martensson, B., & Jonsson, B. (2007). Health-related quality of life measured with EQ-5D in patients treated for depression in primary care. *Value in Health*, 10(2), 153-160.
- 12 Sobocki, P., Ekman, M., Agren, H., Runeson, B., & Jonsson, B. (2006). The mission is remission: Health economic consequences of achieving full remission with antidepressant treatment for depression. *International Journal of Clinical Practice*, 60(7), 791-798.
- 15 Solomon, D. A., Keller, M. B., Leon, A. C., Mueller, T. I., Lavori, P. W., Shea, M. T., ... Endicott, J. (2000). Multiple recurrences of major depressive disorder. *Am J Psychiatry*, 157(2), 229-233.
- 18 Solomon, D. A., Keller, M. B., Leon, A. C., Mueller, T. I., Lavori, P. W., Shea, M. T., ... Endicott, J. (2000). Multiple recurrences of major depressive disorder. *Am J Psychiatry*, 157(2), 229-233.
- 21 Sotsky, S. M., Glass, D. R., Shea, M. T., Pilkonis, P. A., Collins, J. F., Elkin, I., ... Moyer, J. (1991). Patient predictors of response to psychotherapy and pharmacotherapy: findings in the NIMH Treatment of Depression Collaborative Research Program. *Am J Psychiatry*, 148(8), 997-1008.
- 25 Souery, D., Amsterdam, J., de Montigny, C., Lecribier, Y., Montgomery, S., Lipp, O., ... Mendlewicz, J. (1999). Treatment resistant depression: methodological overview and operational criteria. *Eur Neuropsychopharmacol*, 9(1-2), 83-91
- 28 Souery, D., Papakostas, G. I., & Trivedi, M. H. (2006). Treatment-resistant depression. *J Clin Psychiatry*, 67 Suppl 6, 16-22.
- 30 Souery, D., Serretti, A., Calati, R., Oswald, P., Massat, I., Konstantinidis, A., ... Mendlewicz, J. (2011). Citalopram versus desipramine in treatment resistant depression: effect of continuation or switching strategies: a randomized open study. *World Journal of Biological Psychiatry*, 12(5), 364-375.
- 34 Souza, L. H., Salum, G. A., Mosqueiro, B. P., Caldieraro, M. A., Guerra, T. A., & Fleck, M. P. (2016). Interpersonal psychotherapy as add-on for treatment-resistant depression: A pragmatic randomized controlled trial. *Journal of Affective Disorders*, 193, 373-380.
- 37 Spackman, E., Richmond, S., Sculpher, M., Bland, M., Brealey, S., Gabe, R., ... MacPherson, H. (2014). Cost-effectiveness analysis of acupuncture, counselling and usual care in treating patients with depression: The results of the ACUDep trial. *PLoS ONE*, 9.
- 40 Spek, V., Cuijpers, P., Nyklicek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: a meta-analysis. *Psychol Med*, 37(3), 319-328.
- 43 Spek, V., Cuijpers, P., Nyklicek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: a meta-analysis. *Psychol Med*, 37(3), 319-328.

- 1 Spiegelhalter, D., Thomas, A., Best, N., & Lunn, D. (2003). WinBUGS user manual: version 2 1.4.
- 3 Spiegelhalter, D. J. (2001). Bayesian methods for cluster randomized trials with continuous 4 responses. *Statistics in medicine*, 20(3), 435-452.
- 5 Spiers, N., Brugha, T. S., Bebbington, P., McManus, S., Jenkins, R., & Meltzer, H. (2012). 6 Age and birth cohort differences in depression in repeated cross-sectional surveys in 7 England: the National Psychiatric Morbidity Surveys, 1993 to 2007. *Psychological Medicine*, 8 42(10), 2047-2055.
- 9 Spijker, J., de Graaf, R., Bijl, R. V., Beekman, A. T., Ormel, J., & Nolen, W. A. (2002). 10 Duration of major depressive episodes in the general population: results from The 11 Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Br J Psychiatry*, 181, 12 208-213.
- 13 Spiker, D. G., & Kupfer, D. J. (1988). Placebo response rates in psychotic and nonpsychotic 14 depression. *J Affect Disord*, 14(1), 21-23.
- 15 Spiker, D. G., Weiss, J. C., Dealy, R. S., Griffin, S. J., Hanin, I., Neil, J. F., ... Soloff, P. H. 16 (1985). The pharmacological treatment of delusional depression. *Am J Psychiatry*, 142(4), 17 430-436.
- 18 Spitzer, R. L., Williams, J. B., Kroenke, K., Linzer, M., Verloin deGruy, F., Hahn, S. R., ... 19 Johnson, J. G. (1994). Utility of a new procedure for diagnosing mental disorders in primary 20 care: the PRIME-MD 1000 study. *JAMA*, 272(22), 1749-1756.
- 21 Spring, B., Gelenberg, A. J., Garvin, R., & Thompson, S. (1992). Amitriptyline, clovoxamine 22 and cognitive function: a placebo-controlled comparison in depressed outpatients. 23 *Psychopharmacology*, 108(3), 327-332.
- 24 Sramek JJ, Kashkin K, Jasinsky O, Kardatzke D, Kennedy S, Cutler NR. Placebo-controlled 25 study of ABT-200 versus fluoxetine in the treatment of major depressive disorder. 26 *Depression* 1995;3(4):199-203.
- 27 Stansfeld, S., Clark, C., Bebbington, P., King, M., Jenkins, R., & Hinchliffe, S. (2016). 28 Common mental disorders. In S. McManus, P. Bebbington, R. Jenkins, & T. Brugha (Eds.), 29 *Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014*. Leeds: 30 NHS Digital.
- 31 Stassen, H. H., Delini-Stula, A., & Angst, J. (1993). Time course of improvement under 32 antidepressant treatment: a survival-analytical approach. *European 33 Neuropsychopharmacology*, 3, 127-135.
- 34 Stegenga, B. T., Kamphuis, M. H., King, M., Nazareth, I., & Geerlings, M. I. (2012). The 35 natural course and outcome of major depressive disorder in primary care: the PREDICT-NL 36 study. *Soc Psychiatry Psychiatr Epidemiol*, 47(1), 87-95.
- 37 Stein, G., & Bernadt, M. (1993). Lithium augmentation therapy in tricyclic-resistant 38 depression. A controlled trial using lithium in low and normal doses. *The British Journal of 39 Psychiatry*, 162(5), 634-640.
- 40 Stein MK, Rickels K, Weise CC. Maintenance therapy with amitriptyline: a controlled trial. 41 *American Journal of Psychiatry* 1980; 137(3):370-371.
- 42 Steinert, C., Hofmann, M., Kruse, J., & Leichsenring, F. (2014). Relapse rates after 43 psychotherapy for depression - Stable long-term effects? A meta-analysis. *Journal of 44 Affective Disorders*, 168, 107-118.

- 1 Stewart, J. W., McGrath, P. J., Quitkin, F. M., Harrison, W., Markowitz, J., Wager, S., & Leibowitz, M. R. (1989). Relevance of DMS-III Depressive Subtype and Chronicity of Antidepressant Efficacy in Atypical Depression: Differential Response to Phenelzine, Imipramine, and Placebo. *Archives of General Psychiatry*, 46(12), 1080-1087.
- 5 Stewart, J. W., McGrath, P. J., Quitkin, F. M., Rabkin, J. G., Harrison, W., Wager, S., . . . Tricamo, E. (1993). Chronic depression: response to placebo, imipramine, and phenelzine. *Journal of Clinical Psychopharmacology*, 13(6), 391-396.
- 8 Stewart, J. W., Tricamo, E., McGrath, P. J., & Quitkin, F. M. (1997). Prophylactic efficacy of phenelzine and imipramine in chronic atypical depression: likelihood of recurrence on discontinuation after 6 months' remission. *American Journal of Psychiatry*, 154(1), 31-36.
- 11 Stice, E., Burton, E., Bearman, S. K., & Rohde, P. (2007). Randomized trial of a brief depression prevention program: An elusive search for a psychosocial placebo control condition. *Behaviour research and therapy*, 45(5), 863-876.
- 14 Strauss, C., Hayward, M., & Chadwick, P. (2012). Group person-based cognitive therapy for chronic depression: a pilot randomized controlled trial. *British Journal of Clinical Psychology*, 51(3), 345-350.
- 17 Ström M, Uckelstam CJ, Andersson G, Hassmén P, Umefjord G, Carlbring P. Internet-delivered therapist-guided physical activity for mild to moderate depression: a randomized controlled trial. *PeerJ*. 2013 Oct 3; 1:e178.
- 20 Strong, R. E., Marchant, B. K., Reimherr, F. W., Williams, E., Soni, P., & Mestas, R. (2009). Narrow-band blue-light treatment of seasonal affective disorder in adults and the influence of additional nonseasonal symptoms. *Depression and Anxiety*, 26(3), 273-278.
- 23 Sullivan, P. W., Valuck, R., Saseen, J., & MacFall, H. M. (2004). A comparison of the direct costs and cost effectiveness of serotonin reuptake inhibitors and associated adverse drug reactions. *CNS Drugs*, 18, 911-932.
- 26 Sun, H., Zhao, H., Ma, C., Bao, F., Zhang, J., Wang, D. H., ... He, W. (2013). Effects of electroacupuncture on depression and the production of glial cell line-derived neurotrophic factor compared with fluoxetine: a randomized controlled pilot study. *Journal of Alternative and Complementary Medicine*, 19(9), 733-739.
- 30 Suominen, K., Haukka, J., Valtonen, H. M., & Lonnqvist, J. (2009). Outcome of patients with major depressive disorder after serious suicide attempt. *J Clin Psychiatry*, 70(10), 1372-1378.
- 33 Suresh, K., & Bhui, K. (2006). Ethnic minority patients' access to mental health services. *Psychiatry*, 5(11), 413-416.
- 35 Swartz, H. A., Frank, E., Zuckoff, A., Cyranowski, J. M., Houck, P. R., Cheng, Y., ... Shear, M. K. (2008). Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment. *American Journal of Psychiatry*, 165(9), 1155-1162.
- 38 Swift, J. K., & Callahan, J. L. (2009). The impact of client treatment preferences on outcome: a meta-analysis. *Journal of Clinical Psychology*, 65(4), 368-381.
- 40 Swindle, R., Rao, J., Helmy, A., Plue, L., Zhou, X., Eckert, G., Weinberger, M. (2003). Integrating clinical nurse specialists into the treatment of primary care patients with depression. *International Journal of Psychiatry in Medicine*, 33, 17-37.
- 43 Tan RS, Barlow RJ, Abel C, Reddy S, Palmer AJ, Fletcher AE, Nicholl CG, Pitt BM, Bulpitt CJ. The effect of low dose lofepramine in depressed elderly patients in general medical wards. *British Journal of Clinical Pharmacology* 1994; 37(4):321-324.

- 1 Taneja, C., Papakostas, G. I., Jing, Y., Baker, R. A., Forbes, R. A., & Oster, G. (2012). Cost-effectiveness of adjunctive therapy with atypical antipsychotics for acute treatment of major depressive disorder. *Annals of Pharmacotherapy*, 46, 642-649.
- 4 Taylor, D., Sparshatt, A., Varma, S., & Olofinjana, O. (2014). Antidepressant efficacy of agomelatine: meta-analysis of published and unpublished studies. *BMJ*, 348, g1888.
- 6 Taylor, D., Stewart, S., & Connolly, A. (2006). Antidepressant withdrawal symptoms—telephone calls to a national medication helpline. *J Affect Disord*, 95(1-3), 129-133.
- 8 Taylor, R., Brown, J. S., & Weinman, J. (2013). A comparison of the illness perceptions of North Indian and white British women. *J Ment Health*, 22(1), 22-32.
- 10 Teasdale, J. D., Segal, Z. V., Williams, J. M. G., Ridgeway, V. A., Soulsby, J. M., & Lau, M. A. (2000). Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of Consulting and Clinical Psychology*, 68(4), 615.
- 13 Teismann, T., von Brachel, R., Hanning, S., Grillenberger, M., Hebermehl, L., Hornstein, I., & Willutzki, U. (2014). A randomized controlled trial on the effectiveness of a rumination-focused group treatment for residual depression. *Psychother Res*, 24(1), 80-90.
- 16 Teo, A. R., Choi, H., & Valenstein, M. (2013). Social relationships and depression: Ten-year follow-up from a nationally representative study. *PLoS ONE*, 8(4), e62396.
- 18 Teodorczuk, A., Firbank, M. J., Pantoni, L., Poggesi, A., Erkinjuntti, T., Wallin, A., ... O'Brien, J. T. (2009). Relationship between baseline white-matter changes and development of late-life depressive symptoms: 3-year results from the LADIS study. *Psychological Medicine*, 40(04), 603.
- 22 Terman, M., & Terman, J. S. (1995). Treatment of seasonal affective disorder with a high-output negative ionizer. *The Journal of Alternative and Complementary Medicine*, 1(1), 87-92.
- 25 Terman, M., & Terman, J. S. (2006). Controlled trial of naturalistic dawn simulation and negative air ionization for seasonal affective disorder. *American Journal of Psychiatry*, 163(12), 2126-2133.
- 28 Terman, M., Terman, J. S., & Ross, D. C. (1998). A controlled trial of timed bright light and negative air ionization for treatment of winter depression. *Archives of General Psychiatry*, 55(10), 875-882.
- 31 Terra, J., & Montgomery, S. (1998). Fluvoxamine prevents recurrence of depression: results of a long-term, double-blind, placebo-controlled study. *International Clinical Psychopharmacology*, 13(2), 55-62.
- 34 Thase, M. E., Fava, M., Halbreich, U., Kocsis, J. H., Koran, L., Davidson, J., ... Harrison, W. (1996). A placebo-controlled, randomized clinical trial comparing sertraline and imipramine for the treatment of dysthymia. *Archives of General Psychiatry*, 53(9), 777-784.
- 37 Thase, M. E., Friedman, E. S., Biggs, M. M., Wisniewski, S. R., Trivedi, M. H., Luther, J. F., ... Rush, A. J. (2007). Cognitive therapy versus medication in augmentation and switch strategies as second-step treatments: a STAR*D report. *Am J Psychiatry*, 164(5), 739-752.
- 40 Thase, M. E., Nierenberg, A. A., Keller, M. B., & Panagides, J. (2001). Efficacy of mirtazapine for prevention of depressive relapse: a placebo-controlled double-blind trial of recently remitted high-risk patients. *The Journal of Clinical Psychiatry*, 62(10), 782-788.
- 43 Thase, M. E., & Rush, A. J. (1997). When at first you don't succeed: sequential strategies for antidepressant nonresponders. *J Clin Psychiatry*, 58 Suppl 13, 23-29.

- 1 Thompson C. Management of depression in real-life settings: knowledge gained from large-scale clinical trials. International Clinical Psychopharmacology Vol 9 Suppl 3, pp 21-5, 1994.
- 2 Thomas, C. M., & Morris, S. (2003). Cost of depression among adults in England in 2000.
- 3 The British Journal of Psychiatry, 183(6), 514-519.
- 4 Thompson, L. W., Coon, D. W., Gallagher-Thompson, D., Sommer, B. R. & Koin, D. (2001).
- 5 Comparison of desipramine and cognitive/behavioral therapy in the treatment of elderly
- 6 outpatients with mild-to-moderate depression. Am J Geriatr Psychiatry 9, 225-40.
- 7
- 8 Thompson, C., Kinmonth, A. L., Stevens, L., Pevele, R. C., Stevens, A., Ostler, K. J., ...
- 9 Campbell, M. J. (2000). Effects of a clinical-practice guideline and practice-based education
- 10 on detection and outcome of depression in primary care: Hampshire Depression Project
- 11 randomised controlled trial. The Lancet, 355(9199), 185-191.
- 12 Thompson, C., Ostler, K., Peveler, R. C., Baker, N., & Kinmonth, A. L. (2001). Dimensional
- 13 perspective on the recognition of depressive symptoms in primary care: The Hampshire
- 14 Depression Project 3. Br J Psychiatry, 179, 317-323.
- 15 Thompson, C., Thompson, S., & Smith, R. (2004). Prevalence of seasonal affective disorder
- 16 in primary care; a comparison of the seasonal health questionnaire and the seasonal pattern
- 17 assessment questionnaire. Journal of Affective Disorders, 78(3), 219-226.
- 18 Thomson, J., Rankin, H., Ashcroft, G., Yates, C. M., McQueen, J. K., & Cummings, S.
- 19 (1982). The treatment of depression in general practice: a comparison of L-tryptophan,
- 20 amitriptyline, and a combination of L-tryptophan and amitriptyline with placebo. Psychological
- 21 Medicine, 12(4), 741-751.
- 22 Thoren, P., Floras, J. S., Hoffmann, P., & Seals, D. R. (1990). Endorphins and exercise:
- 23 physiological mechanisms and clinical implications. Med Sci Sports Exerc, 22(4), 417-428.
- 24 Tiemens, B. G., Ormel, J., Jenner, J. A., van der Meer, K., Van Os, T. W., van den Brink, R.
- 25 H., . . . van den Brink, W. (1999). Training primary-care physicians to recognize, diagnose
- 26 and manage depression: does it improve patient outcomes? Psychol Med, 29(4), 833-845.
- 27 Titov, N., Dear, B. F., Ali, S., Zou, J. B., Lorian, C. N., Johnston, L., ... Fogliati, V. J. (2015).
- 28 Clinical and cost-effectiveness of therapist-guided internet-delivered cognitive behavior
- 29 therapy for older adults with symptoms of depression: a randomized controlled trial. Behavior
- 30 therapy, 46(2), 193-205.
- 31 Titov N, Dear BF, Staples LG, Terides MD, Karin E, Sheehan J, Johnston L, Gandy M,
- 32 Fogliati VJ, Wootton BM, McEvoy PM. Disorder-specific versus transdiagnostic and clinician-
- 33 guided versus self-guided treatment for major depressive disorder and comorbid anxiety
- 34 disorders: a randomized controlled trial. Journal of Anxiety Disorders. 2015 Oct 31; 35:88-102.
- 35
- 36 Tollefson GD HS. Analysis of the Hamilton Depression Rating Scale factors from a double-
- 37 blind, placebo-controlled trial of fluoxetine in geriatric major depression. International Clinical
- 38 Psychopharmacology 8(4):253-9, 1993.
- 39 Tollefson, G. D., Bosomworth, J. C., Heiligenstein, J. H., Potvin, J. H., & Holman, S. (1995).
- 40 A double-blind, placebo-controlled clinical trial of fluoxetine in geriatric patients with major
- 41 depression. International Psychogeriatrics, 7(1), 89-104.
- 42 Tollefson GD, Greist JH, Jefferson JW, Heiligenstein JH, Sayler ME, Tollefson SL, Koback K.
- 43 Is baseline agitation a relative contraindication for a selective serotonin reuptake inhibitor: a
- 44 comparative trial of fluoxetine versus imipramine. Journal of Clinical Psychopharmacology
- 45 1994 Dec; 14(6):385-91.

- 1 Tor, P.-C., Bautovich, A., Wang, M.-J., Martin, D., Harvey, S. B., & Loo, C. (2015). A systematic review and meta-analysis of brief versus ultrabrief right unilateral electroconvulsive therapy for depression. *The Journal of Clinical Psychiatry*, 8(2), e1092-e1098.
- 5 Torkan H, Blackwell SE, Holmes EA, Kalantari M, Neshat-Doost HT, Maroufi M, Talebi H. Positive imagery cognitive bias modification in treatment-seeking patients with major depression in Iran: A pilot study. *Cognitive therapy and research*. 2014 Apr 1; 38(2):132-45.
- 8 Town, J. M., Abbass, A., & Hardy, G. (2011). Short-term psychodynamic psychotherapy for personality disorders: a critical review of randomized controlled trials. *J Pers Disord*, 25(6), 723-740.
- 11 Traffanstedt, M. K., Mehta, S., & LoBello, S. G. (2016). Major depression with seasonal variation. *Clinical Psychological Science*, 4(5), 825-834.
- 13 Trinquet, L., Abbé, A., & Ravaud, P. (2012). Impact of reporting bias in network meta-analysis of antidepressant placebo-controlled trials. *PLoS ONE*, 7(4), e35219.
- 15 Trivedi, M. H., Rush, A. J., Wisniewski, S. R., Nierenberg, A. A., Warden, D., Ritz, L., . . . Team, S. D. S. (2006). Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry*, 163(1), 28-40.
- 19 Truax, C. B., & Carkhuff, R. R. (1967). Toward effective counseling and psychotherapy; training and practice, by Charles B. Truax and Robert R. Carkhuff. Chicago: Aldine.
- 21 Turner, E. H., Matthews, A. M., Linardatos, E., Tell, R. A., & Rosenthal, R. (2008). Selective publication of antidepressant trials and its influence on apparent efficacy. *New England Journal of Medicine*, 358(3), 252-260.
- 24 Tylee, A., Freeling, P., Kerry, S., & Burns, T. (1995). How does the content of consultations affect the recognition by general practitioners of major depression in women? *Br J Gen Pract*, 45(400), 575-578.
- 27 Tyrer, P. (1984). Psychiatric clinics in general practice. An extension of community care. *The British Journal of Psychiatry*, 145(1), 9-14.
- 29 Uebelacker, L., Marootian, B., Tigue, P., Haggarty, R., Primack, J., & Miller, I. (2011). Telephone Depression Care Management for Latino Medicaid Health Plan Members: A Pilot Randomized Controlled Trial. *Journal of Nervous and Mental Disease*, 199, 678-683.
- 32 Unutzer, J., Katon, W., Callahan, C. M., Williams, J. W., Jr., Hunkeler, E., Harpole, L., . . . Treatment, I. I. I. M.-P. A. t. C. (2002). Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA*, 288(22), 2836-2845.
- 35 Unutzer, J., Katon, W., Callahan, C. M., Williams, J. W., Jr., Hunkeler, E., Harpole, L., . . . Treatment, I. I. I. M.-P. A. t. C. (2002). Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA*, 288(22), 2836-2845.
- 38 Usaf S, Kavanagh DJ. Mechanisms of improvement in treatment for depression: test of a self-efficacy and performance model. *Journal of Cognitive Psychotherapy*. 1990; 4(1):51-70.
- 40 Ustun, T. B., & Sartorius, N. (1995). Mental illness in general health care: an international study: JSTOR.
- 42 Valenstein, M., Pfeiffer, P. N., Brandfon, S., Walters, H., Ganoczy, D., Kim, H. M., . . . Heisler, M. (2016). Augmenting ongoing depression care with a mutual peer support intervention versus self-help materials alone: A randomized trial. *Psychiatric services (Washington, D.C.)*, 67(2), 236-239.

- 1 Valkanova, V., & Ebmeier, K. P. (2013). Vascular risk factors and depression in later life: A systematic review and meta-analysis. *Biological Psychiatry*, 73(5), 406-413.
- 3 Vallejo, J., Gasto, C., Catalan, R., & Salamero, M. (1987). Double-blind study of imipramine versus phenelzine in Melancholias and Dysthymic Disorders. *The British Journal of Psychiatry*, 151(5), 639-642.
- 6 van den Broek, W. W., Birkenhager, T., Mulder, P. G., Bruijn, J. A., & Moleman, P. (2006). Imipramine is effective in preventing relapse in electroconvulsive therapy-responsive depressed inpatients with prior pharmacotherapy treatment failure: a randomized, placebo-controlled trial. *Journal of Clinical Psychiatry*, 67(2), 263.
- 10 van den Broek, W. W., Birkenhager, T. K., Mulder, P. G., Bruijn, J. A., & Moleman, P. (2004). A double-blind randomized study comparing imipramine with fluvoxamine in depressed inpatients. *Psychopharmacology (Berl)*, 175(4), 481-486.
- 13 Van Gurp, G., Meterissian, G. B., Haiek, L. N., McCusker, J., & Bellavance, F. (2002). St John's wort or sertraline? Randomized controlled trial in primary care. *Canadian Family Physician*, 48(5), 905-912.
- 16 van Os, T. W. D. P., van den Brink, R. H. S., Tiemens, B. G., Jenner, J. A., van der Meer, K., & Ormel, J. (2004). Are effects of depression management training for General Practitioners on patient outcomes mediated by improvements in the process of care? *Journal of Affective Disorders*, 80(2-3), 173-179.
- 20 van Schaik, A., van Marwijk, H., Adèr, H., van Dyck, R., de Haan, M., Penninx, B., ... Beekman, A. (2006). Interpersonal psychotherapy for elderly patients in primary care. *The American Journal of Geriatric Psychiatry*, 14(9), 777-786.
- 23 van Schaik, D. J. F., Klijn, A. F. J., van Hout, H. P. J., van Marwijk, H. W. J., Beekman, A. T. F., de Haan, M., & van Dyck, R. (2004). Patients' preferences in the treatment of depressive disorder in primary care. *General Hospital Psychiatry*, 26(3), 184-189.
- 26 van Straten, A., Hill, J., Richards, D., & Cuijpers, P. (2015). Stepped care treatment delivery for depression: a systematic review and meta-analysis. *Psychological Medicine*, 45(02), 231-246.
- 29 van Straten, A., Tiemens, B., Hakkaart, L., Nolen, W. A., & Donker, M. C. (2006). Stepped care vs. matched care for mood and anxiety disorders: a randomized trial in routine practice. *Acta Psychiatr Scand*, 113(6), 468-476.
- 32 van Weel-Baumgarten, E., van den Bosch, W., van den Hoogen, H., & Zitman, F. G. (1998). Ten year follow-up of depression after diagnosis in general practice. *Br J Gen Pract*, 48(435).
- 34 Vanelle, J., Attar-Levy, D., Poirier, M., Bouhassira, M., Blin, P., & Oli, J. (1997). Controlled efficacy study of fluoxetine in dysthymia. *The British Journal of Psychiatry*, 170(4), 345-350.
- 36 Van't Veer-Tazelaar, P., Smit, F., Hout, H., Oppen, P., Horst, H., Beekman, A., & Marwijk, H. (2010). Cost-effectiveness of a stepped care intervention to prevent depression and anxiety in late life: randomised trial. *British Journal of Psychiatry*, 196(4), 319-325.
- 39 van't Veer-Tazelaar, P., van Marwijk, H., van Oppen, P., van Hout, H., van der Horst H., Cuijpers, P., Smit, F. & Beekman, A. (2009) Stepped-Care Prevention of Anxiety and Depression in Late Life: A Randomized Controlled Trial, *Archives of General Psychiatry*, 66, 297-304
- 43 Vartiainen H. Double-blind study of mirtazapine and placebo in hospitalized patients with major depression. *European Neuropsychopharmacology* 4(2):145-50, 1994; 4(2):145-150.

- 1 Vasiliadis HM, Latimer E, Dionne PA, Préville M. The costs associated with antidepressant use in depression and anxiety in community-living older adults. *Can J Psychiatry*. 2013 Apr; 58(4):201-9.
- 4 Veale DM, Le Fevre K, Pantelis C, De Souza V, Mann A, Sargeant A. Aerobic exercise in the adjunctive treatment of depression: a randomized controlled trial. *Journal of the Royal Society of Medicine*. 1992 Sep 1; 85(9):541-4.
- 7 Ventura, D., Armstrong, E.P., Skrepnek, G.H. & Erder, M.H. (2007) Escitalopram versus sertraline in the treatment of major depressive disorder: a randomized clinical trial. *Current Medical Research and Opinion*, 23, 245-250.Juan, Puerto Rico.
- 10 Verduyn C, Barrowclough C, Roberts J, Tarrier N, Harrington R. Maternal depression and child behaviour problems. *The British Journal of Psychiatry*. 2003 Oct 1; 183(4):342-8.
- 12 Vernmark, K., et al. (2010) Internet administered guided self-help versus individualized e-mail therapy: A randomized trial of two versions of CBT for major depression. *Behaviour research and therapy* 48, 368-376
- 15 Versiani, M., Amrein, R., Stabl, M., & Group, I. C. S. (1997). Moclobemide and imipramine in chronic depression (dysthymia): an international double-blind, placebo-controlled trial. *International Clinical Psychopharmacology*, 12(4), 183-194.
- 18 Versiani, M., Mehilane, L., Gaszner, P., & Arnaud-Castiglioni, R. (1999). Reboxetine, a unique selective NRI, prevents relapse and recurrence in long-term treatment of major depressive disorder. *The Journal of Clinical Psychiatry*, 60(6):400-6.
- 21 Versiani M, Moreno R, Ramakers-van Moorsel CJ, Schutte AJ. Comparison of the effects of mirtazapine and fluoxetine in severely depressed patients. *CNS Drugs* 2005; 19(2):137-46.
- 23 Versiani M, Ontiveros A, Mazzotti GG. Fluoxetine versus amitriptyline in the treatment of major depression with associated anxiety (anxious depression): a double-blind comparison. *Int Clin Psychopharmacology* 1999; 14:321-327.
- 26 Vitriol VG, Ballesteros ST, Florenzano RU, Weil KP, Benadof DF. Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. *Psychiatric services*. 2009 Jul; 60(7):936-42.
- 29 Vlasveld, M. C., Feltz-Cornelis, C. M., Adèr, H. J., Anema, J. R., Hoedeman, R., Mechelen, W., & Beekman, A. T. (2012). Collaborative care for major depressive disorder in an occupational healthcare setting. *British Journal of Psychiatry*, 200(6), 510-511.
- 32 Volz, H.-P., Eberhard, R., & Grill, G. (2000). Wirksamkeit und Verträglichkeit des Johanniskrautextraktes D-0496 bei leichten bis mittelschweren depressiven Episoden - eine placebokontrollierte Doppelblindstudie über 6 Wochen. *Nervenheilkunde*, 19(7), 401-405.
- 35 von Knorring, L., Åkerblad, A.-C., Bengtsson, F., Carlsson, Å., & Ekselius, L. (2006). Cost of depression: effect of adherence and treatment response. *European Psychiatry*, 21(6), 349-354.
- 38 Von Korff, M., & Goldberg, D. (2001). Improving outcomes in depression. *BMJ*, 323(7319), 948-949.
- 40 Von Korff, M., Gruman, J., Schaefer, J., Curry, S. J., & Wagner, E. H. (1997). Collaborative management of chronic illness. *Ann Intern Med*, 127(12), 1097-1102.
- 42 Vos, T., Flaxman, A. D., Naghavi, M., Lozano, R., Michaud, C., Ezzati, M., ... Aboyans, V. (2013). Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*, 380(9859), 2163-2196.

- 1 Vythilingam, M., Chen, J., Bremner, J. D., Mazure, C. M., Maciejewski, P. K., & Nelson, J. C. (2003). Psychotic depression and mortality. *Am J Psychiatry*, 160(3), 574-576.
- 2 Waddell, G., & Burton, A. K. (2006). Is work good for your health and well-being? The Stationery Office (TSO), London.
- 3 Wade A, Lemming O, & Hedegaard KB. Escitalopram 10 mg/day is effective and well tolerated in a placebo-controlled study in depression in primary care. *International Clinical Psychopharmacology* 2002; 17(3):95-102.
- 4 Wade, A. G., Toumi, I., & Hemels, M. E. H. (2005). A pharmacoeconomic evaluation of escitalopram versus citalopram in the treatment of severe depression in the United Kingdom. *Clinical Therapeutics*, 27, 486-496.
- 5 Wade, A. G., Toumi, I., & Hemels, M. E. H. (2005). A probabilistic cost-effectiveness analysis of escitalopram, generic citalopram and venlafaxine as a first-line treatment of major depressive disorder in the UK. *Current Medical Research and Opinion*, 21, 631-641.
- 6 Wagner, E. H. (1997). Managed care and chronic illness: health services research needs. *Health Serv Res*, 32(5), 702-714.
- 7 Wagner, E. H., & Groves, T. (2002). Care for chronic diseases. *BMJ*, 325(7370), 913-914.
- 8 Wagner, H., & Bladt, S. (1994). Pharmaceutical quality of hypericum extracts. *J Geriatr Psychiatry Neurol*, 7 Suppl 1, S65-68.
- 9 Wagner B, Horn AB, Maercker A. Internet-based versus face-to-face cognitive-behavioral intervention for depression: a randomized controlled non-inferiority trial. *Journal of affective disorders*. 2014 Jan 31; 152:113-21.
- 10 Wakelin JS. Fluvoxamine in the treatment of the older depressed patient; double-blind, placebo-controlled data. *International Clinical Psychopharmacology* 1(3):221-30, 1986.
- 11 Waller, R., & Gilbody, S. (2009). Barriers to the uptake of computerized cognitive behavioural therapy: a systematic review of the quantitative and qualitative evidence. *Psychol Med*, 39(5), 705-712.
- 12 Walsh, A. E. L., & Harmer, C. J. (2015). The cognitive neuropsychological model of antidepressant response. *Current Opinion in Psychology*, 4, 124-130.
- 13 Wang, E. J., Barecki-Roach, M., & Johnson, W. W. (2004). Quantitative characterization of direct P-glycoprotein inhibition by St John's wort constituents hypericin and hyperforin. *J Pharm Pharmacol*, 56(1), 123-128.
- 14 Wang T, Wang L, Tao W, Chen L. Acupuncture combined with an antidepressant for patients with depression in hospital: a pragmatic randomised controlled trial. *Acupuncture in Medicine*. 2014 Apr 29:acupmed-2013.
- 15 Wang WD, Lu XY, Ng SM, Hong L, Zhao Y, Lin YN, Wang F. Effects of electro-acupuncture on personality traits in depression: a randomized controlled study. *Chinese journal of integrative medicine*. 2013 Oct 1; 19(10):777-82.
- 16 Waraich, P., Goldner, E. M., Somers, J. M., & Hsu, L. (2004). Prevalence and incidence studies of mood disorders: A systematic review of the literature. *The Canadian Journal of Psychiatry*, 49(2), 124-138.
- 17 Ward, E., King, M., Lloyd, M., Bower, P., Sibbald, B., Farrelly, S., ... Addington-Hall, J. (2000). Randomised controlled trial of non-directive counselling, cognitive-behaviour therapy, and usual general practitioner care for patients with depression. I: clinical effectiveness. *BMJ*, 321(7273), 1383-1388.

- 1 Warmerdam L, van Straten A, Twisk J, Riper H, Cuijpers P. Internet-based treatment for adults with depressive symptoms: randomized controlled trial. *Journal of medical Internet research*. 2008 Oct; 10(4).
- 4 Watanabe, N., Omori, I. M., Nakagawa, A., Cipriani, A., Barbui, C., Churchill, R., & Furukawa, T. A. (2011). Mirtazapine versus other antidepressive agents for depression. *Cochrane Database Syst Rev* (12), CD006528.
- 7 Watkins, E. (2016). Rumination-focused cognitive-behavioral therapy for depression. New York: The Guilford Press.
- 9 Watkins, E. R. (2008). Constructive and unconstructive repetitive thought. *Psychol Bull*, 134(2), 163-206.
- 11 Watkins, E. R., Mullan, E., Wingrove, J., Rimes, K., Steiner, H., Bathurst, N., ... Scott, J. (2011). Rumination-focused cognitive-behavioural therapy for residual depression: phase II randomised controlled trial. *British Journal of Psychiatry*, 199(4), 317-322.
- 14 Watkins, E. R., & Nolen-Hoeksema, S. (2014). A habit-goal framework of depressive rumination. *J Abnorm Psychol*, 123(1), 24-34.
- 16 Watkins, E. R., Taylor, R. S., Byng, R., Baeyens, C., Read, R., Pearson, K., & Watson, L. (2012). Guided self-help concreteness training as an intervention for major depression in primary care: a Phase II randomized controlled trial. *Psychological Medicine*, 42(7), 1359-1371.
- 20 Watson, J.C., Gordon, L.B., Stermac, L., Kalogerakos, F., & Steckley, P. (2003) Comparing the effectiveness of process-experiential with cognitive-behavioural psychotherapy in the treatment of depression. *Journal of Consulting and Clinical Psychology*, 71 (4), 773-781.
- 23 Weich, S., Nazroo, J., Sproston, K., McManus, S., Blanchard, M., Erens, B., Tyre, P. (2004). Common mental disorders and ethnicity in England: the EMPIRIC study. *Psychol Med*, 34(8), 1543-1551.
- 26 Weimer, K., Colloca, L., & Enck, P. (2015). Placebo effects in psychiatry: mediators and moderators. *The Lancet Psychiatry*, 2(3), 246-257.
- 28 Weissman, M. M. (1987). Advances in psychiatric epidemiology: rates and risks for major depression. *Am J Public Health*, 77(4), 445-451.
- 30 Weissman, M. M., Markowitz, J. C., & Klerman, G. L. (2000). Comprehensive guide to interpersonal psychotherapy. New York: Basic Books.
- 32 Weissman MM, Prusoff B, Sholomskas AJ, Greenwald S. A double-blind clinical trial of alprazolam, imipramine, or placebo in the depressed elderly. *Journal of Clinical Psychopharmacology* 1992; 12(3):175-182.
- 35 Weitz, E., Kleiboer, A., van Straten, A., Hollon, S. D., & Cuijpers, P. (2017). Individual patient data meta-analysis of combined treatments versus psychotherapy (with or without pill placebo), pharmacotherapy or pill placebo for adult depression: a protocol. *BMJ Open*, 7(2), e013478.
- 39 Wells, K., Sherbourne, C., Schoenbaum, N., Duan, N., Meredith, L., Unutzer, J., ... Rubenstein, L. (2000). Impact of disseminating quality improvement programs for depression in managed primary care: a randomized controlled trial. *Journal of the American Medical Association*, 283, 212-220.
- 43 Wernicke JF, Dunlop SR, Dornseif BE, Bosomworth JC, Humbert M. Low-dose fluoxetine therapy for depression. *Psychopharmacology Bulletin* 1988; 24(1):183-188.

- 1 Wernicke JF, Bosomworth JC, Ashbrook E. Fluoxetine at 20 mg per day: The recommended 2 and therapeutic dose in the treatment of depression. International Clinical 3 Psychopharmacology 1989; 4(1): 63-67.
- 4 Westrin, Å. & Lam, R. W. (2007). Seasonal affective disorder: A clinical update. Annals of 5 Clinical Psychiatry, 19(4), 239-246.
- 6 Wheatley, D. (1997). LI 160, an extract of St. John's wort, versus amitriptyline in mildly to 7 moderately depressed outpatients--a controlled 6-week clinical trial. Pharmacopsychiatry, 30 8 Suppl 2, 77-80.
- 9 Wheatley DP, Van Moffaert M, Timmerman L, Kremer CME. Mirtazapine: Efficacy and 10 tolerability in comparison with fluoxetine in patients with moderate to severe major 11 depressive disorder. Journal of Clinical Psychiatry 1998; 59(6):306-312.
- 12 Whisman, M. A. (2001). The association between depression and marital dissatisfaction 13 Marital and family processes in depression: A scientific foundation for clinical practice. (pp. 3- 14 24): American Psychological Association (APA).
- 15 Whisman, M. A., Sheldon, C. T., & Goering, P. (2000). Psychiatric disorders and 16 dissatisfaction with social relationships: Does type of relationship matter? Journal of 17 Abnormal Psychology, 109(4), 803-808.
- 18 Whisman, M. A., & Uebelacker, L. A. (2003). Comorbidity of relationship distress and mental 19 and physical health problems. In D. K. Snyder & M. A. Whisman (Eds.), Treating difficult 20 couples: helping clients with coexisting mental and relationship disorders. New York: Guilford 21 Press.
- 22 White, A. (2004). A cumulative review of the range and incidence of significant adverse 23 events associated with acupuncture. Acupunct Med, 22(3), 122-133.
- 24 White, K., Razani, J., Cadow, B., Gelfand, R., Palmer, R., Simpson, G., & Sloane, R. B. 25 (1984). Tranylcypromine vs nortriptyline vs placebo in depressed outpatients: a controlled 26 trial. Psychopharmacology, 82(3), 258-262.
- 27 Whiteford, H. A., Degenhardt, L., Rehm, J., Baxter, A. J., Ferrari, A. J., Erskine, H. E., . . . 28 Johns, N. (2013). Global burden of disease attributable to mental and substance use 29 disorders: findings from the Global Burden of Disease Study 2010. The Lancet, 382(9904), 30 1575-1586.
- 31 WHO. (1992). ICD-10, the ICD-10 classification of mental and behavioural disorders: clinical 32 descriptions and diagnostic guidelines. Geneva: World Health Organization.
- 33 Whooley, M. A., Avins, A. L., Miranda, J., & Browner, W. S. (1997). Case-finding instruments 34 for depression. Two questions are as good as many. J Gen Intern Med, 12(7), 439-445.
- 35 Wiersma, J. E., Van Schaik, D. J., Hoogendorn, A. W., Dekker, J. J., Van, H. L., Schoevers, 36 R. A., . . . Van Oppen, P. (2014). The effectiveness of the cognitive behavioral analysis 37 system of psychotherapy for chronic depression: a randomized controlled trial. Psychother 38 Psychosom, 83(5), 263-269.
- 39 Wijkstra, J., Burger, H., van den Broek, W. W., Birkenhager, T. K., Janzing, J. G. E., Boks, 40 M. P. M., . . . Nolen, W. A. (2010). Long-term response to successful acute pharmacological 41 treatment of psychotic depression. Journal of Affective Disorders, 123(1-3), 238-242.
- 42 Wijkstra J, Burger H, van den Broek WW, Birkenhäger TK, Janzing JG, Boks MP, et al. 43 Treatment of unipolar psychotic depression: a randomized, double-blind study comparing 44 imipramine, venlafaxine, and venlafaxine plus quetiapine. Acta Psychiatrica Scandinavica 45 2010; 121(3):190–200."

- 1 Wijkstra, J., Lijmer, J., Burger, H., Cipriani, A., Geddes, J., & Nolen, W. A. (2015).
- 2 Pharmacological treatment for psychotic depression. Cochrane Database Syst Rev (7), CD004044.
- 3
- 4 Wilcox, C., Cohn, J., Katz, B., Mijares, C., Guarino, J., Panagides, J., & DeFrancisco, D.
- 5 (1994). A double-blind, placebo-controlled study comparing mianserin and amitriptyline in
- 6 moderately depressed outpatients. International Clinical Psychopharmacology, 9(4), 271-
- 7 280.
- 8 Wileman, S. M., ANDREW, J. E., HOWIE, F. L., CAMERON, I. M., McCORMACK, K., NAJI,
- 9 S. A., & EAGLES, J. M. (2001). Light therapy for seasonal affective disorder in primary care.
- 10 The British Journal of Psychiatry, 178(4), 311-316.
- 11 Wiles, N., Thomas, L., Abel, A., Ridgway, N., Turner, N., Campbell, J., ... Lewis, G. (2013).
- 12 Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based
- 13 patients with treatment resistant depression: results of the CoBalT randomised controlled
- 14 trial. Lancet, 381(9864), 375-384.
- 15 Wiles, N. J., Thomas, L., Turner, N., Garfield, K., Kounali, D., Campbell, J., ... Hollinghurst,
- 16 S. (2016). Long-term effectiveness and cost-effectiveness of cognitive behavioural therapy
- 17 as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: Follow-
- 18 up of the CoBalT randomised controlled trial. The Lancet Psychiatry, 3, 137-144.
- 19 Wiley-Exley, E., Domino, M. E., Maxwell, J., & Levkoff, S. E. (2009). Cost-effectiveness of
- 20 integrated care for elderly depressed patients in the PRISM-E study. Journal of Mental
- 21 Health Policy and Economics, 12, 205-213+217.
- 22 Wilfley, D. E. (2000). Interpersonal psychotherapy for group (1st ed.). New York: Basic Books.
- 23
- 24 Wilkinson, P., Alder, N., Juszczak, E., Matthews, H., Merritt, C., Montgomery, H., ... Jacoby,
- 25 R. (2009). A pilot randomised controlled trial of a brief cognitive behavioural group
- 26 intervention to reduce recurrence rates in late life depression. International Journal of
- 27 Geriatric Psychiatry, 24(1), 68-75.
- 28 Williams, A. D., Blackwell, S. E., Mackenzie, A., Holmes, E. A., & Andrews, G. (2013).
- 29 Combining imagination and reason in the treatment of depression: a randomized controlled
- 30 trial of internet-based cognitive-bias modification and internet-CBT for depression. Journal of
- 31 Consulting and Clinical Psychology, 81(5), 793-799.
- 32 Williams JM, Crane C, Barnhofer T, Brennan K, Duggan DS, Fennell MJ, Hackmann A,
- 33 Krusche A, Muse K, Von Rohr IR, Shah D. Mindfulness-based cognitive therapy for
- 34 preventing relapse in recurrent depression: a randomized dismantling trial. Journal of
- 35 consulting and clinical psychology. 2014 Apr; 82(2):275.
- 36 Williams AD, O'Moore K, Blackwell SE, Smith J, Holmes EA, Andrews G. Positive imagery
- 37 cognitive bias modification (CBM) and internet-based cognitive behavioral therapy (iCBT): A
- 38 randomized controlled trial. Journal of affective disorders. 2015 Jun 1; 178:131-41.
- 39 Williams, C., Wilson, P., Morrison, J., McMahon, A., Walker, A., Andrew, W., ... Tansey, L.
- 40 (2013). Guided self-help cognitive behavioural therapy for depression in primary care: a
- 41 randomised controlled trial. PLoS ONE, 8(1), e52735.
- 42 Williams, J. B. (1988). A structured interview guide for the Hamilton Depression Rating
- 43 Scale. Archives of General Psychiatry, 45(8), 742-747.
- 44 Williams Jr, J. W., Barrett, J., Oxman, T., Frank, E., Katon, W., Sullivan, M., ... Sengupta, A.
- 45 (2000). Treatment of dysthymia and minor depression in primary care: a randomized
- 46 controlled trial in older adults. JAMA, 284(12), 1519-1526.

- 1 Williams, J. W., Jr., Kerber, C. A., Mulrow, C. D., Medina, A., & Aguilar, C. (1995).
- 2 Depressive disorders in primary care: prevalence, functional disability, and identification. *J Gen Intern Med*, 10(1), 7-12.
- 3
- 4 Williams, R., & Hunt, K. (1997). Psychological distress among British South Asians: the contribution of stressful situations and subcultural differences in the West of Scotland Twenty-07 Study. *Psychol Med*, 27(5), 1173-1181.
- 5
- 6
- 7 Wilson, K., Mottram, P., Ashworth, L., & Abou-Saleh, M. (2003). Older community residents with depression: long-term treatment with sertraline. *The British Journal of Psychiatry*, 182(6), 492-497.
- 8
- 9
- 10 Wilson, M., & MacCarthy, B. (1994). GP consultation as a factor in the low rate of mental health service use by Asians. *Psychol Med*, 24(1), 113-119.
- 11
- 12 Wittchen, H. U., Höfler, M., & Meister, W. (2001). Prevalence and recognition of depressive syndromes in German primary care settings: poorly recognized and treated? *International Clinical Psychopharmacology*, 16(3), 121-135.
- 13
- 14
- 15 Witte, B., Harrer, G., Kaptan, T., Podzuweit, H., & Schmidt, U. (1995). [Treatment of depressive symptoms with a high concentration hypericum preparation. A multicenter placebo-controlled double-blind study]. *Fortschr Med*, 113(28), 404-408.
- 16
- 17
- 18 Woelk, H. (2000). Comparison of St John's wort and imipramine for treating depression: randomised controlled trial. *BMJ*, 321(7260), 536-539.
- 19
- 20 Wong, S. Y., Mercer, S. W., Woo, J., & Leung, J. (2008). The influence of multi-morbidity and self-reported socio-economic standing on the prevalence of depression in an elderly Hong Kong population. *BMC Public Health*, 8(1), 119.
- 21
- 22
- 23 World Bank. (1993). World development report: Investing in health research development. Geneva, Switzerland:
- 24
- 25 Wright JH, Wright AS, Albano AM, Basco MR, Goldsmith LJ, Raffield T, Otto MW. Computer-assisted cognitive therapy for depression: maintaining efficacy while reducing therapist time. *American Journal of Psychiatry*. 2005 Jun 1; 162(6):1158-64.
- 26
- 27
- 28 Wuthrich VM, Rapee RM. Randomised controlled trial of group cognitive behavioural therapy for comorbid anxiety and depression in older adults. *Behaviour Research and Therapy*. 2013 Dec 31; 51(12):779-86.
- 29
- 30
- 31 Yang W, Ding Z, Dai T, Peng F, Zhang JX. Attention bias modification training in individuals with depressive symptoms: a randomized controlled trial. *Journal of behavior therapy and experimental psychiatry*. 2015 Dec 31; 49:101-11.
- 32
- 33
- 34 Yesavage, J. A., Brink, T. L., Rose, T. L., Lum, O., Huang, V., Adey, M., & Leirer, V. O. (1982). Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res*, 17(1), 37-49.
- 35
- 36
- 37 Yeung A, Lepoutre V, Wayne P, Yeh G, Slipp LE, Fava M, Denninger JW, Benson H, Fricchione GL. Tai chi treatment for depression in Chinese Americans: a pilot study. *American Journal of Physical Medicine & Rehabilitation*. 2012 Oct 1; 91(10):863-70.
- 38
- 39
- 40 Yeung, A., Shyu, I., Fisher, L., Wu, S., Yang, H., & Fava, M. (2010). Culturally sensitive collaborative treatment for depressed Chinese Americans in primary care. *American Journal of Public Health*, 100, 2397-2340.
- 41
- 42
- 43 Yevtunshenko, V.Y., Belous, A.I., Yevtushenko, Y.G., Gusinin, S.E., Buzik, O.J. & Agibalova, T.V. (2007) Efficacy and tolerability of escitalopram versus citalopram in major depressive
- 44

- 1 disorder: a 6-week, multicentre, prospective, randomized, double-blind, active-controlled study in adult outpatients. *Clinical Therapeutics*, 29, 2319-2332
- 3 Yiend, J., Paykel, E., Merritt, R., Lester, K., Doll, H., & Burns, T. (2009). Long term outcome of primary care depression. *J Affect Disord*, 118(1-3), 79-86.
- 5 Yirmiya, R., & Bab, I. (2009). Major depression is a risk factor for low bone mineral density: a meta-analysis. *Biol Psychiatry*, 66(5), 423-432.
- 7 Yoshimura R, Hori H, Umena-Nakano W, Ikenouchi-Sugita A, Katsuki A, Atake K, Nakamura J. Comparison of lithium, aripiprazole and olanzapine as augmentation to paroxetine for inpatients with major depressive disorder. *Therapeutic advances in psychopharmacology*. 2014 Jun; 4(3):123-9.
- 11 Young JP, Coleman A, Lader MH. A controlled comparison of fluoxetine and amitriptyline in depressed out-patients. *British Journal of Psychiatry* 1987 Sep; 151:337-40.
- 13 Zanardi, R., Franchini, L., Gasperini, M., Perez, J., & Smeraldi, E. (1996). Double-blind controlled trial of sertraline versus paroxetine in the treatment of delusional depression. *Am J Psychiatry*, 153(12), 1631-1633.
- 16 Zanardi R, Franchini L, Serretti A, Perez J, Smeraldi E. Venlafaxine versus fluvoxamine in the treatment of delusional depression: a pilot double-blind controlled study. *Journal of Clinical Psychiatry* 2000; 61(1):26-9.
- 19 Zanarini, M. C., Frankenburg, F. R., Dubo, E. D., Sickel, A. E., Trikha, A., Levin, A., & Reynolds, V. (1998). Axis I comorbidity of borderline personality disorder. *Am J Psychiatry*, 155(12), 1733-1739.
- 22 Zhang, Z. J., Ng, R., Man, S. C., Li, J. T., Wong, W., Wong, H. K., ... Sze, S. C. (2013). Use of electroacupuncture to accelerate the antidepressant action of selective serotonin reuptake inhibitors: a single-blind, randomised, controlled study. *Xianggang yi xue za zhi [Hong Kong medical journal]*, 19 Suppl 9, 12-16.
- 26 Zhang B, Ma WY, Zhang BT, Liu SL. A control study of clinical therapeutic effects of laser-acupuncture on depressive neurosis. *World Journal of Acupuncture-Moxibustion*. 1996; 6(2):12-7.
- 29 Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatr Scand*, 67(6), 361-370.
- 31 Zivin, K., Pfeiffer, P. N., Bohnert, A. S., Ganoczy, D., Blow, F. C., Nallamothu, B. K., & Kales, H. C. (2013). Evaluation of the FDA warning against prescribing citalopram at doses exceeding 40 mg. *Am J Psychiatry*, 170(6), 642-650.
- 34 Zivkov M. Org 3770 versus amitriptyline: A 6-week randomized double-blind multicentre trial in hospitalized depressed patients. *Human Psychopharmacology* 1995; 10(3):173-180.
- 36 Zu, S., et al. (2014) A comparison of cognitive-behavioral therapy, antidepressants, their combination and standard treatment for Chinese patients with moderate-severe major depressive disorders. *Journal of Affective Disorders* 152-154, 262-267
- 39 Zung, W. W. (1965). A self-rating depression scale. *Arch Gen Psychiatry*, 12, 63-70.
- 40 Zusky, P. M., Biederman, J., Rosenbaum, J. F., Manschreck, T. C., Gross, C. C., Weilberg, J. B., & Gastfriend, D. R. (1988). Adjunct low dose lithium carbonate in treatment-resistant depression: a placebo-controlled study. *Journal of Clinical Psychopharmacology*, 8(2), 120-124.