

Subarachnoid haemorrhage

[S] Evidence review for patient information

NICE guideline <number>

Evidence review underpinning

February 2021

Draft for consultation

*Developed by the National Guideline Centre,
hosted by the Royal College of Physicians*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and, where appropriate, their carer or guardian.

Local commissioners and providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to Notice of rights.

ISBN

[add for final publication version only, delete this text for consultation version]

Contents

1 Patient information	5
1.1 Review question: What patient information (including lifestyle advice) should be given to adults who have had an aneurysmal subarachnoid haemorrhage?	5
1.2 Introduction	5
1.3 Characteristics table	5
1.4 Qualitative evidence	6
1.4.1 Included studies	6
1.4.2 Excluded studies	6
1.4.3 Summary of studies included in the evidence review	7
1.4.4 Qualitative evidence synthesis	10
1.4.5 Qualitative evidence summary	14
1.5 Economic evidence	19
1.6 The committee's discussion of the evidence.....	19
1.6.1 Interpreting the evidence.....	19
1.6.2 Cost effectiveness and resource use	20
1.6.3 Other factors the committee took into account	21
Appendices	24
Appendix A: Review protocols	24
Appendix B: Literature search strategies	30
B.1 Clinical search literature search strategy	30
B.2 Health Economics literature search strategy.....	34
Appendix C: Qualitative evidence selection.....	35
Appendix D: Qualitative evidence tables	36
Appendix E: Excluded studies.....	46
E.1 Excluded qualitative studies	46

1 Patient information

2 Evidence review underpinning recommendations 1.4.1 and 1.5.3 to 1.5.6 and 1.5.8 to 1.5.10
3 in the NICE guideline.

1.1 Review question: What patient information (including lifestyle advice) should be given to adults who have had an aneurysmal subarachnoid haemorrhage?

1.2 Introduction

8 Aneurysmal subarachnoid haemorrhage is a sudden onset, life-threatening condition with
9 variable outcome. Despite advances in treatment and improved survival and functional
10 recovery rates, many subarachnoid haemorrhage survivors report ongoing difficulties. It is
11 common for them to experience fatigue, and cognitive, emotional and psychosocial issues.
12 These problems are associated with reduced quality of life, social participation and
13 community re-integration, and return to work.

14 People with aneurysmal subarachnoid haemorrhage and their family and/or carers require
15 information that explains their diagnosis, treatment, expectations of recovery and follow up.
16 Alongside information about their condition and treatment, patients and their family and/or
17 carers often seek advice on actions to reduce the risk of recurrence and allow return to
18 previous activities. Currently, the provision of this information varies widely between
19 neurosurgical centres and across wider healthcare settings.

1.3 Characteristics table

21 For full details see the review protocol in Appendix A:.

22 **Table 1: Characteristics of review question**

Objective	To determine what information (such as lifestyle advice) should be given to people who have had a subarachnoid haemorrhage.
Population and setting	Inclusion: Adults (16 and older) who have had a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm. Exclusion: <ul style="list-style-type: none">• Adults with subarachnoid haemorrhage caused by head injury, ischaemic stroke or an arteriovenous malformation.• Children and young people aged 15 years and younger.
Context	aSAH is associated with significant sequelae and can have a notable effect on the person's life post-haemorrhage. Adequate information provision for a person with aSAH and their family and/or carer can aid care planning and management, improve understanding and accuracy of expectations, and can influence quality of life. As such, this information for a person with aSAH and their family and/or carer provided both within the immediate care setting and in the community following discharge can be invaluable.
Review strategy	Synthesis of qualitative and quantitative research. Results will be presented in table format. The quality of the qualitative evidence will be assessed by a GRADE CerQual approach for each review finding and results presented in summary tables. Descriptive quantitative data such as incidence rate or frequencies of information preference from survey questionnaires will be synthesised, quality assessed, and considered alongside qualitative evidence.

1.4 1 Qualitative evidence

1.4.1 2 Included studies

3 Six studies were included in the review,^{3, 6, 8, 11, 21, 25} these are summarised in Table 2 below.
4 Four studies provided qualitative data^{3, 8, 11, 21} and 2 provided quantitative data.^{6, 25} Key
5 findings from these studies are summarised in Section 1.4.2 below. See also the study
6 selection flow chart in Appendix C:, study evidence tables in Appendix D:, and excluded
7 studies lists in Appendix E:.

1.4.2 8 Excluded studies

9 See the excluded studies list in Appendix E:.

10

11

© NIHR 2024 All rights reserved. Guidance for National of rights
7

1.4.3 1 Summary of studies included in the evidence review

2 Table 2: Summary of studies included in the review

Study	Design	Population	Research aim	Comments
Berggren 2010 ³	In-depth semi-structured interviews with thematic qualitative analysis. The interviews were reported to have taken place one year and seven months after the event for 60 – 90 minutes in the participants homes.	People who had experienced and were treated for SAH at a university hospital in Sweden were interviewed. Informants who were contactable and native language speaking at the time of their discharge were selected from the hospitals patient record system. N=9	The aim was to analyse peoples accounts of SAH and to describe how they create meaning for the SAH and the events that surround it. Specific questions were: - What is highlighted in the accounts of SAH? - How is the illness reconstructed? - How is meaning created through communicative interaction with others about SAH?	A study with some methodological limitations based in Sweden. Focused on specific topics so results appear to be partially applicable. The relevant data has been extracted for the review.
Dulhanty 2019 ⁶	A cross sectional postal survey, with quantitative collation of responses. The Self-Reported Needs after Stroke Questionnaire was modified and used to measure the self-reported needs of SAH survivors and the extent to which they were met 1–2 years and 3–5 years post haemorrhage.	People who had experienced a subarachnoid haemorrhage (SAH) in Greater Manchester between September 2010–September 2012 (late post-SAH group) and September 2013–September 2014 (early post-SAH group) were recruited. All adult patients admitted to the Regional Neurosciences Centre with SAH within these dates, who survived and were discharged home from the neurosciences centre or inpatient rehabilitation were identified from the centre's	Explore the type and frequency of self reported needs both early and late following SAH specifically to the extent to which they have been met.	A study with some methodological limitations based in UK. Focused on specific topics aside from patient information so results appear to be partially applicable. The relevant data has been extracted for the review.

Study	Design	Population	Research aim	Comments
		prospectively maintained clinical database. N=203		
Hedlund 2010 ⁸	A semi structured interview guide covering the following question areas: (1) what the participants perceived as important for recovery from aSAH, and (2) perceived consequences of SAH. The interviews were carried out 12 months after onset of aSAH. Each participant was asked to choose a time and setting for the interview (outpatient clinic or home).	Men and women of different ages and from different places in the catchment area who had undergone different treatment for aSAH and with no or minor neurological impairments at discharge from the neurosurgical unit. N=20	The study constitutes the qualitative part of an ongoing study investigating psychiatric morbidity, cognitive strategies, coping and quality of life in patients after aSAH. The aims of this study were to describe what participants with no or minor neurological deficits after aSAH perceived as being important for recovery, and perceived consequences of aSAH.	A study with some methodological limitations based in Sweden. Focused on specific topics in relation to depression or no depression, so results appear to be partially applicable. The relevant data has been extracted for the review.
Jarvis 2002 ¹¹	Through qualitative methodology participants were interviewed using a semi-structured approach in their own homes.	Patients who had suffered a SAH within the last 14 – 18 months. Patients were excluded if they were continuing to receive treatment of any description. N=8	The study aimed to illuminate the experience of recovery from SAH.	A study with some methodological limitations based in UK. Focused on specific topics in relation to symptoms or stressful experiences post SAH, so results appear to be partially applicable. The relevant data has been extracted for the review.
Persson 2017 ²¹	Explorative interview study with a qualitative design. Individual interviews, with open ended questions, using an interview guide were performed with all participants. The participants chose the time and location of the interview (home or University of Gothenburg).	Participants were included from the Extended Stroke Arm Longitudinal study at the University of Gothenburg. The inclusion criteria were > 18 years of age, with an SAH between 4 th February 2009 until the 2 nd of December 2010, receiving care at the	To explore experiences of care and rehabilitation as well as the consequences and strategies used to cope with everyday life six years after SAH.	A well conducted study based in Sweden. Focused on the consequences of SAH and the coping strategies patients use to cope. Results should be generally applicable.

Study	Design	Population	Research aim	Comments
		Sahlgrenska University Hospital in Gothenburg, Sweden, resident in the Gothenburg Urban area (<35km from the hospital), able to speak Swedish and at least some participants should be of working age. The participants were contacted by phone and if they agreed to participate, a time and interview was planned. N=16		
Von Vogelsang 2004 ²⁵	Intervention study with quasi experimental design with qualitative and quantitative components to the study. Patients were divided into two groups while in hospital and either provided verbal and written information or only verbal information from medical professionals.	Participants with ruptured intracranial aneurysm treated consecutively at a Swedish neurosurgical clinic. Participants were recruited over a period of 12 months and recruitment was concluded in 2001. All patients in the sample were acute admissions; and those treated for non-ruptured aneurysms were not included. Patients were included if they were be able to understand, speak and read Swedish, and had a score of 4–5 on the Glasgow Outcome Scale (GOS). N=62	An investigation into the effects of increased information for patients treated for intracranial aneurysm rupture.	A study with some methodological limitations based in Sweden. Focused on specific topics in relation to patients information such as ease of understanding and matching of their needs as well as levels of anxiety, so results appear to be applicable. The relevant data has been extracted for the review.

1 See Appendix D: for full evidence tables.

1.4.4 1 Qualitative evidence synthesis

2 **Table 3: Qualitative Review findings**

Main findings	Statement of finding
Theme 1 - Clarity of information	
SAH event	People have difficulty in remembering what happened up to their SAH and immediately thereafter. There are some who want to know what happened, but others do not want to know what happened and prefer that to remain in the past.
Ongoing symptoms	People want to know what symptoms to expect after SAH treatment to enable their families and themselves prepare
Medical information	People want information on what to expect in the period immediately after surgery, both negative and positive outcomes and what kind of care they might expect.
Theme 2 - Support	
Social relationships	People value social relationships from a range of sources, to provide practical and emotional support through the process of care.
Theme 3 - Future	
Long term implications	In addition to immediate symptoms after surgery, people want to know what symptoms might be experienced long- term and potentially may not improve after SAH treatment
Reoccurrence	People value information about SAH, including information about risk factors for SAH, and risk and prevention of recurrent SAH.

3 **Table 4: Quantitative Review findings**

Main findings	Statement of finding	
Information Needs – people needed information and/or advice about		
	N (%)	
Health	<ul style="list-style-type: none"> SAH 105 (58) General health (diet / alcohol / smoking) 40 (21) Physical relationships 17 (9) 	
Travel	<ul style="list-style-type: none"> Driving 19 (10) Public transport 22 (11) 	
Finance	<ul style="list-style-type: none"> Benefits 34 (17) Money management 7 (4) 	
Home	<ul style="list-style-type: none"> Home aids 20 (10) Adaptations 10 (5) Moving home 12 (6) 	
Information Needs – peoples satisfaction with information given		
	P value	
Meeting information needs Scale: 0-5, low is poor outcome	Written + oral info: 4.4 (2.9) Oral info only: 3.3 (1.6)	0.06
Ease of understanding Scale: 0-5, low is poor outcome	Written + oral info: 3.9 (1.8) Oral info only: 3.3 (1.9)	0.24

1.4.4.1 1 Narrative summary of review findings

2 Clarity of information

3 Qualitative findings:

4 Review finding 1: SAH event

5 People felt that having a SAH was an extremely traumatic experience. However, most people
6 can only clearly recollect their experience after intervention and the events leading up to
7 surgery are not clear or missing. For some this was frustrating as they had been through a
8 life changing event and undergone an intervention but could not recollect what happened.
9 This set of people wanted to be know more and understand what an SAH is and what had
10 happened to them. Other people do not wish to know more about their condition and prefer
11 to focus on their current symptoms and recovery.

12 Some respondents could not recollect any information given to them while they were
13 admitted in hospital, but were told by their families that they were given oral and written
14 information. Overall, people and their families value information on the process of care
15 throughout the interventional journey. Information would help to reduce anxiety after the
16 SAH.

17 This review finding was based on primary research addressing the experiences of people
18 who have had an SAH and their families, mostly in Sweden. There was a judgement of
19 moderate confidence in this review finding, because of concerns about direct relevance and
20 coherence.

21 Review finding 2: Ongoing symptoms

22 The side effects and symptoms after having a SAH varied from patient to patient. Fatigue,
23 inability to concentrate and completing routine tasks were complications that people faced
24 once discharged home. It is unclear whether people and their families were informed of what
25 to expect once discharged. Therefore, it seems relevant to ensure that they are appropriately
26 informed in person and through written information about common symptoms after having a
27 SAH and how to manage those once home, with the support of families or carers.

28 This review finding was based on primary research addressing the difficulties people had
29 once they were home and in their recovery period. There was a judgement of high
30 confidence in this review finding, as there were no significant methodological limitations or
31 concerns.

32 Review finding 3: Medical information

33 People were suffering with the immediate and long-term consequences of intervention for
34 SAH, often not knowing what to expect. It is important that they know and understand the
35 impact of surgery on their quality of life, their ability to return to activities of normal daily life,
36 how to manage pain, and how to identify and respond to complications. Topics such as risks
37 for family members, surgical techniques of treatment, rehabilitation and future risks were also
38 highlighted as important to be informed about.

39 Some people reported the feeling of abandonment once they were discharged from hospital
40 and were unclear about ongoing medical support or follow up, and this had the potential to
41 increase anxiety. Those who did receive rehabilitation support and follow up felt supported.
42 However, some participants did not receive any rehabilitation support and felt unable to
43 request it. This added to the feeling of abandonment.

1 This review finding was based on primary research addressing the experiences of people
2 who have been discharged after SAH surgery. There was a judgement of moderate
3 confidence in this review finding, because of minor concerns about relevance.

4 A point that was consistent through all three of these review findings was that respondents
5 wanted information that was easy to understand for themselves, families or carers.

6 **Support**

7 **Review finding 1: Different sources of support**

8 People emphasized the importance of the support of family, friends, and community
9 members throughout their journey. People also valued the support from the health care
10 professionals involved in their care. Particularly after discharge, support involved practical
11 help with activities of daily living, and in employment and emotional support during distress.
12 Some people felt that they might be a burden for family and friends, but they appreciated the
13 practical and emotional support through the process of care. People wanted their primary
14 care provider to have more information about SAH, so they could be a source of information
15 and support as this was lacking.

16 This review finding was based on primary research addressing the experiences of people
17 after SAH intervention. There was a judgement of high confidence in this review finding, as
18 there were no significant methodological limitations or concerns.

19 **Future**

20 **Review finding 1: Long term implications**

21 People wanted more information about the long-term consequences of having a SAH and
22 neurosurgery. This is linked partially to the symptoms they faced once they were discharged,
23 mainly what people called “invisible” symptoms such as fatigue and cognitive symptoms such
24 as memory impairment, planning and concentration. Some participants were able to find their
25 own adaptations to their day to day lives, especially in their places of work. Some
26 participants struggled as their employers or team members were unaware of what support
27 they may need and the consequences of having a SAH. A few also had to face
28 unemployment as they were unable to continue their jobs, which affected them economically
29 and psychologically.

30 This review finding was based on primary research addressing the experiences of people
31 after SAH surgery. There was a judgement of high confidence in this review finding, as there
32 were no significant methodological limitations or concerns.

33 **Review finding 2: Recurrence**

34 People wanted to know how likely it was that they would have another aneurysm or
35 haemorrhage. At times this was referred to as existential threat or worry about reoccurrence
36 and survival. People wish to understand the risks for reoccurrence and what modifiable risk
37 factors they may also have.

38 This review finding was based on primary research addressing the experiences of people
39 who underwent surgery for SAH. There was a judgement of high confidence in this review
40 finding, as there were no significant methodological limitations or concerns.

41 **Quantitative findings**

42 **Information needs:**

43 People highlighted that they wanted more information on a variety of different topics after
44 their SAH. Linked to the other findings, people wanted more information about their SAH and

1 as well as their general health and well being. This included modifications to their lifestyle but
2 also information about resuming physical relationships. As the SAH may have an affect on a
3 person's cognitive and physical abilities, they wanted to know what aids are available for
4 support and what adaptations could be made to support their recovery.

5 As the recovery period after an SAH may affect their ability to work or continue employment,
6 more information was needed on benefit support and managing money. People also wanted
7 more information on travelling and how the SAH would affect their ability to drive and using
8 public transport.

9 Patients also reported that receiving information both written and orally better met their
10 information needs and was slightly easier for them to understand than receiving only oral
11 information.

12 This review finding was based on primary research addressing the experiences of people
13 who had a SAH within the UK. There was a judgement of high confidence in this review
14 finding, as there were no significant methodological limitations or concerns.

15

1.4.5 1 Qualitative evidence summary

2 Table 5: Summary of Qualitative evidence – CERQual checklist

Study design and sample size		Finding	Quality assessment			
Number of studies contributing to the finding	Design		Criteria	Rating	Overall assessment of confidence	
Clarity of Information						
5	Semi structured qualitative interviews + intervention study with quasi-experimental design	<p>SAH event: people have difficulty in remembering what happened up to their SAH and immediately thereafter. There are some who want to know what happened, but others do not want to know what happened and prefer that to remain in the past.</p> <p><i>Berggren 2010³Hedlund 2010⁸</i> <i>Jarvis 2002¹¹</i> <i>Persson 2017²¹</i> <i>Von Vogelsang 2004²⁵</i></p>	<p>Limitations</p> <p>Minor methodological limitations ^{a, b} (five studies with minor methodological limitations due to responses to a variety of different open questions asked between the studies)</p>	<p>Coherence</p> <p>Minor concerns about coherence ^{a, c} (due to consolidation of multiple concerns into one theme)</p>	<p>Relevance</p> <p>Minor concerns about relevance ^a (due to the difference of responses and importance under one theme)</p>	MODERATE
			<p>Adequacy</p> <p>Minor concerns ^a</p>			

				(five studies that overall offered moderately rich data)	
4	<p>Ongoing Symptoms: people want to know what symptoms to expect after SAH treatment to enable their families and themselves prepare</p> <p><i>Berggren 2010³</i> <i>Hedlund 2010⁸</i> <i>Jarvis 2002¹¹</i> <i>Persson 2017²¹</i></p>	Limitations	Minor concerns about limitations ^a (four studies with minor methodological limitations)	MODERATE	
		Coherence	Minor concerns ^a (data generally consistent within and across all studies)		
		Relevance	Minor concerns ^a (studies of people and families who have experienced SAH in the UK and Sweden)		
		Adequacy	Minor concerns ^a (four studies that together offered moderately rich data)		
5	<p>Medical information: people want information on what to expect in the period immediately after surgery, both negative and positive outcomes and what kind of care they might expect.</p> <p><i>Berggren 2010³</i> <i>Hedlund 2010⁸</i> <i>Jarvis 2002¹¹</i> <i>Persson 2017²¹</i> <i>Von Vogelsang 2004²⁵</i></p>	Limitations	Minor concerns about limitations ^a (five studies with minor methodological limitations)	MODERATE	
		Coherence	Minor concerns ^a (data generally consistent within and across all studies)		
		Relevance	Minor concerns ^a (studies of people and families who have		

				experienced SAH in the UK and Sweden)	
			Adequacy	Minor concerns ^a (five studies that together offered moderately rich data)	
Support					
4	Semi structured qualitative interviews	Social support: social relationships - People value social relationships from a range of sources, to provide practical and emotional support through the process of care. <i>Berggren 2010</i> ³ <i>Hedlund 2010</i> ⁸ <i>Jarvis 2002</i> ¹¹ <i>Persson 2017</i> ²¹	Limitations	Minor concerns about limitations (four studies with minor methodological limitations)	MODERATE
			Coherence	Minor concerns (data generally consistent within and across all studies)	
			Relevance	Minor concerns (studies of people and families who have experienced SAH in the UK and Sweden)	
			Adequacy	No concerns (four studies that together offered moderately rich data)	
Future					
4	Semi structured qualitative interviews	Long term consequences: in addition to immediate symptoms after surgery, people want to know what symptoms might be experienced long- term and potentially may not improve after SAH <i>Berggren 2010</i> ³ <i>Hedlund 2010</i> ⁸	Limitations	Minor concerns about limitations (four studies with minor methodological limitations)	MODERATE
			Coherence	Minor concerns	

		<p><i>Jarvis 2002</i>¹¹ <i>Persson 2017</i>²¹</p>		(data generally consistent within and across all studies)	
			Relevance	Minor concerns (studies of people and families who have experienced SAH in the UK and Sweden)	
			Adequacy	Minor concerns (four studies that together offered moderately rich data)	
4		<p>Reoccurrences: people value information about SAH, including information about risk factors for SAH, and risk and prevention of recurrent SAH. <i>Berggren 2010</i>³ <i>Hedlund 2010</i>⁸ <i>Jarvis 2002</i>¹¹ <i>Persson 2017</i>²¹</p>	Limitations	Minor concerns about limitations (four studies with minor methodological limitations)	MODERATE
			Coherence	Minor concerns (data generally consistent within and across all studies)	
			Relevance	Minor concerns (studies of people and families who have experienced SAH in the UK and Sweden)	
			Adequacy	Minor concerns (four studies that together offered moderately rich data)	

1 **Table 6: Summary of Quantitative evidence**

Study design and sample size		Finding	Risk of Bias
Number of studies contributing to the finding	Design		
1	Cross-sectional survey	<p>Information needs – people needed information and/or advice about</p> <p>Health: SAH – 105 patients, general health (diet / alcohol / smoking) – 40 patients, physical relationships – 17 patients</p> <p>Travel: driving – 19 patients, public transport – 22 patients</p> <p>Finance: benefits – 34 patients, money management – 7 patients</p> <p>Home: home aids – 20 patients, adaptations – 10 patients, moving home – 12 patients.</p> <p><i>Dulhanty 2019⁶</i></p>	Moderate ^a
1	Intervention study with quasi experimental design	<p>Information Needs – people satisfaction with information given</p> <p>Meeting information needs: Written + oral info: 4.4 (2.9) compared to Oral info only: 3.3 (1.6)</p> <p>Ease of understanding: Written + oral info: 3.9 (1.8) compared to Oral info only: 3.3 (1.9)</p> <p><i>Von Vogelsang 2004²⁵</i></p>	High ^b

2 *Moderate risk of bias due to (a) outcome measurement bias, 203 (51%) participants responded to the questionnaire (JBI checklist for cross sectional study), and (b)*
 3 *potential selection bias with intervention start and follow-up start differing between intervention and control groups, and potential confounding bias with no adjusting for*
 4 *possible confounding factors between control and intervention groups (ROBINS-I).*

5

1.5 1 Economic evidence

- 2 The committee agreed that health economic studies would not be relevant to this review
3 question, and so were not sought.

1.6 4 The committee's discussion of the evidence

1.6.1 5 Interpreting the evidence

1.6.1.1 6 The quality of the evidence

7 Qualitative Findings

8 The quality of evidence was of moderate quality. Evidence was downgraded for minor
9 concerns about methodological limitations with responses to a variety of different open
10 questions asked between the included studies, minor concerns about contextual relevance
11 with inclusion of studies of people and families who have experienced SAH in the UK and
12 Sweden, and minor concerns about adequacy of data in the quantity and richness of data
13 available. The committee highlighted that each of these concerns were minor and considered
14 the evidence to be reliable.

15 Quantitative Findings

16 The quantitative evidence was considered to be at moderate to high risk of bias. This was
17 mostly due to low rates of participants questionnaire response and potential selection bias.

18 Overall, the committee agreed that the evidence provided a rich understanding of the
19 different themes and topics people would like information about after having a SAH. In
20 addition to this evidence, the committee used their experience to agree topics that should be
21 included, as a minimum, in the information given. Therefore, the committee were able to
22 make strong recommendations for patient information.

1.6.1.2 3 Findings identified in the evidence synthesis

24 The committee discussed the findings of the review and felt the themes identified reflected
25 their experience of the information needs of patients and was from well-conducted qualitative
26 and quantitative studies. The committee considered the evidence helpful in highlighting
27 people's specific information needs at a challenging time due to the effects of aneurysmal
28 SAH on memory and recall of information in the peri-operative and early recovery period.

29 The qualitative evidence showed that patients described some themes that were important
30 for the content of information considered useful after an SAH. One of the themes highlighted
31 was the clarity of information people were given. Patients described that recall of information
32 was a challenge with regards to the SAH event, but any information given after the event too.
33 This contributed to people's understanding of what ongoing symptoms they could expect and
34 their understanding of the long-term medical implications after having an SAH. A specific
35 point raised was that patients wanted to know what their medical follow up after the event
36 would entail, as some felt abandoned once discharged from hospital.

37 Another theme identified from the evidence was the support that people with an SAH
38 received. They were grateful for all the support received from formal medical service
39 providers and informal carers and family, which all contributed to activities of daily living.
40 People wanted their carers and families to also be well informed about SAH and what to
41 expect after an SAH in order to support their family member with the SAH effectively.

1 An important theme drawn from the evidence was that people wanted to know the long term
2 medical and social implications of having a SAH. This ranged from what symptoms they
3 could face because of the SAH, but also the implications on daily functioning including
4 employment. Recurrence of SAH was also a topic which was important to people to be well
5 informed about.

6 The quantitative evidence supplemented what the qualitative evidence showed in that it also
7 highlighted similar areas of information preference from the participant responses. The
8 quantitative data also identified that people wanted more information on specific topics such
9 as future health, travel, finance and home life. Notably, ~60% of participants from one study
10 reported that they wished for more information on SAH and how it could impact on their
11 health. The evidence conveyed that patients wanted information that was easy to
12 understand, so they wanted information in both written and verbal formats. The written format
13 would allow patients and their families to refer to information after they had left hospital and
14 aid ongoing understanding through the recovery phase. One study showed that satisfaction
15 with information received was higher in patients who received written and oral information
16 compared to patients who received only oral information.

17 The committee noted that information provision may help to inform people's expectation of
18 recovery after a SAH and this may help the recovery process and reduce potential stress or
19 anxiety. The evidence also found that some people do not want in-depth information about
20 their SAH as this may contribute to anxiety or stress during their recovery period. A person's
21 choice about the level of information wanted should also be taken into account after an SAH.

22 The recommendations made by the committee outlined the areas important to include in
23 patient information: providing information to the person on what an aneurysmal subarachnoid
24 haemorrhage is and what caused it, treatment options and the possible side effects, ongoing
25 symptoms or complications from having had an aSAH, ongoing support available from local
26 and national groups, and advice on resuming usual activities such as returning to work, and
27 driving. The importance of written information was highlighted, in order that people were able
28 to review the information at a time suitable for themselves.

29 The committee also drew from their experience to suggest some additional and more specific
30 areas of information provision, which might be useful for patients and their families or carers.
31 These included explaining how aSAH has been treated and how effective the treatment has
32 been, and any follow-up plan for the patient including a named contact within the specialist
33 centre who could be contacted for future advice. The committee also agreed that information
34 should be provided about other possible symptoms that in their experience were common
35 such as sleep disturbance, headache, low mood, changes to taste and smell and other
36 advice such as medications the person may be taking, and wound care.

37 There is variation in information provision for people who have had a SAH. The committee
38 highlighted within the recommendations that information provision and support should be
39 given to people on admission with a SAH, after treatment, and post discharge.

40 The committee noted that verbal information and support needs to be tailored to the
41 individual. The guideline committee were aware that this would be especially applicable for
42 people with learning disabilities, concerns with capacity and those with language difficulties.

1.6.23 Cost effectiveness and resource use

44 Cost effectiveness evidence was not sought as this is a qualitative review. The
45 recommendations provide guidance regarding what topics should be covered within a patient
46 review, but the committee agreed that this will not impact consultation time.

47 The committee considered that increased provision of information might be required where it
48 does not already exist in a suitable format. However, this is in line with the general principles
49 of provision of information already established in the existing NICE Patient experience

1 guideline and so the recommendations based on this review do not represent an additional
2 resource impact. The committee noted that good quality information was accessible from
3 sources such as The British Stroke Association and Headway.

1.6.3 4 Other factors the committee took into account

5 When a person has a SAH, patient information should be communicated and delivered
6 clearly, openly and in a manner that enables the person and their family to be involved in
7 shared decision making about their treatment and recovery pathway. The committee
8 considered that improved information provision may enhance understanding of SAH, and
9 could aid recovery and influence expectations of the possible positive and negative
10 outcomes. The committee highlighted that some people may not want information about their
11 SAH or would prefer their family or carers receive the information instead of them. The
12 committee considered these general principles in providing information and support to be
13 covered by the Patient experience in adult NHS services and a cross reference to this
14 guideline was added to the recommendations.
15

1 References

- 2 1. Berggren E, Sidenvall B, Larsson D. Memory ability after subarachnoid haemorrhage:
3 relatives' and patients' statements in relation to test results. *British Journal of*
4 *Neuroscience Nursing*. 2010; 6(8):383-388
- 5 2. Berggren E, Sidenvall B, Larsson D. Subarachnoid haemorrhage has long-term
6 effects on social life. *British Journal of Neuroscience Nursing*. 2011; 7(1):429-435
- 7 3. Berggren E, Sidenvall B, Muhli UH. Identity construction and meaning-making after
8 subarachnoid haemorrhage. *British Journal of Neuroscience Nursing*. 2010; 6(2):86-
9 93
- 10 4. Cedzich C, Roth A. Neurological and psychosocial outcome after subarachnoid
11 haemorrhage, and the hunt and hess scale as a predictor of clinical outcome.
12 *Zentralblatt für Neurochirurgie*. 2005; 66(3):112-118
- 13 5. Covey J, Noble AJ, Schenk T. Family and friends' fears of recurrence: impact on the
14 patient's recovery after subarachnoid hemorrhage. *Journal of Neurosurgery*. 2013;
15 119(4):948-954
- 16 6. Dulhanty LH, Hulme S, Vail A, Patel HC, Tyson SF. The self-reported needs of
17 patients following subarachnoid hemorrhage (SAH). *Disability and Rehabilitation*.
18 2019; <https://dx.doi.org/10.1080/09638288.2019.1595748>
- 19 7. He S, Mack WJ. Health-related quality of life after aneurysmal subarachnoid
20 hemorrhage: interplay between physical, cognitive, and emotional factors. *World*
21 *Neurosurgery*. 2014; 81(1):37-39
- 22 8. Hedlund M, Zetterling M, Ronne-Engstrom E, Ekselius L, Carlsson M. Perceived
23 recovery after aneurysmal subarachnoid haemorrhage in individuals with or without
24 depression. *Journal of Clinical Nursing*. 2010; 19(11-12):1578-1587
- 25 9. Hellawell DJ, Pentland B. Relatives' reports of long term problems following traumatic
26 brain injury or subarachnoid haemorrhage. *Disability and Rehabilitation*. 2001;
27 23(7):300-305
- 28 10. Hutter BO, Kreitschmann-Andermahr I. Subarachnoid hemorrhage as a psychological
29 trauma. *Journal of Neurosurgery*. 2014; 120(4):923-930
- 30 11. Jarvis A. Recovering from subarachnoid haemorrhage: patients' perspective. *British*
31 *Journal of Nursing*. 2002; 11(22):1430-1437
- 32 12. Karic T, Roe C, Haug Nordenmark T, Becker F, Sorteberg A. Impact of early
33 mobilization and rehabilitation on global functional outcome one year after
34 aneurysmal sub arachnoid haemorrhage. *Journal of Rehabilitation Medicine*. 2016;
35 48(8):676-682
- 36 13. King JT, Jr., Horowitz MB, Bissonette DJ, Tsevat J, Roberts MS. What do patients
37 with cerebral aneurysms know about their condition? *Neurosurgery*. 2006; 58(5):824-
38 830; discussion 824-830
- 39 14. Morris PG, Wilson JT, Dunn L. Anxiety and depression after spontaneous
40 subarachnoid hemorrhage. *Neurosurgery*. 2004; 54(1):47-52; discussion 52-44
- 41 15. Murgo M, Cavanagh K, Latham S. Health Related Quality of Life and support needs
42 for sub-arachnoid haemorrhage survivors in New South Wales Australia. *Australian*
43 *Critical Care*. 2016; 29(3):146-150

- 1 16. National Institute for Health and Care Excellence. Developing NICE guidelines: the
2 manual [updated October 2018]. London. National Institute for Health and Care
3 Excellence, 2014. Available from:
4 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20overview>
- 5 17. Nishino A, Sakurai Y, Tsuji I, Arai H, Uenohara H, Suzuki S et al. Resumption of work
6 after aneurysmal subarachnoid hemorrhage in middle-aged Japanese patients.
7 *Journal of Neurosurgery*. 1999; 90(1):59-64
- 8 18. Nordenmark TH, Karic T, Roe C, Sorteberg W, Sorteberg A. The post-aSAH
9 syndrome: a self-reported cluster of symptoms in patients with aneurysmal
10 subarachnoid hemorrhage. *Journal of Neurosurgery*. 2020; 132(5):1556-1565
- 11 19. Ogden JA, Utley T, Mee EW. Neurological and psychosocial outcome 4 to 7 years
12 after subarachnoid hemorrhage. *Neurosurgery*. 1997; 41(1):25-34
- 13 20. Passier PECA, Visser-Meily JMA, Rinkel GJE, Lindeman E, Post MWM.
14 Determinants of health-related quality of life after aneurysmal subarachnoid
15 hemorrhage: a systematic review. *Quality of Life Research*. 2013; 22(5):1027-1043
- 16 21. Persson HC, Tornbom K, Sunnerhagen KS, Tornbom M. Consequences and coping
17 strategies six years after a subarachnoid hemorrhage - A qualitative study. *PloS One*.
18 2017; 12(8):e0181006
- 19 22. Preiss M, Netuka D, Koblihova J, Bernardova L, Charvat F, Kratochvilova D et al.
20 Quality of life 1 year after aneurysmal subarachnoid hemorrhage in good outcome
21 patients treated by clipping or coiling. *Journal of Neurological Surgery*. 2012;
22 73(4):217-223
- 23 23. Ross S, Bhargava D, Al-Tamimi Y, Goddard T, Tennant A, Quinn A. Identifying
24 patient report outcomes relevant to aneurysmal subarachnoid hemorrhage follow-up.
25 'In:' Zuccarello M, Clark J, Pyne-Geithman G, Andaluz N, Hartings J, Adeoye O,
26 editors. *Cerebral Vasospasm: Neurovascular Events After Subarachnoid Hemorrhage*
27 *Acta Neurochirurgica Supplement*. 115. Vienna: Springer. 2013. p. 13-16.
- 28 24. Stegen G, Freckmann N. Outcome and rehabilitation after aneurysmal subarachnoid
29 haemorrhage. *Zentralblatt für Neurochirurgie*. 1991; 52(1):37-39
- 30 25. von Vogelsang AC, Wengstrom Y, Forsberg C. Patient information after ruptured
31 intracranial aneurysm. *Journal of Advanced Nursing*. 2004; 48(6):551-559

32
33

1 Appendices

2 Appendix A: Review protocols

3 Table 7: Review protocol: Patient information

ID	Field	Content
0.	PROSPERO registration number	CRD42019160099
1.	Review title	What patient information (including lifestyle advice) should be given to adults who have had an aneurysmal subarachnoid haemorrhage?
2.	Review question	What patient information (including lifestyle advice) should be given to adults who have had an aneurysmal subarachnoid haemorrhage?
3.	Objective	To determine what information (such as lifestyle advice) should be given to people who have had a subarachnoid haemorrhage.
4.	Searches	<p>The following databases will be searched:</p> <ul style="list-style-type: none"> • Embase • MEDLINE • CINAHL • PsycINFO <p>Searches will be restricted by:</p> <ul style="list-style-type: none"> • English language only <p>The searches may be re-run 6 weeks before the final committee meeting and further studies retrieved for inclusion if relevant.</p> <p>The full search strategies will be published in the final review.</p>
5.	Condition or domain being studied	Aneurysmal subarachnoid haemorrhage
6.	Population	<p>Inclusion: Adults (16 and older) who have had a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm.</p> <p>Exclusion:</p> <ul style="list-style-type: none"> • Adults with subarachnoid haemorrhage caused by head injury, ischaemic stroke or an arteriovenous malformation. • Children and young people aged 15 years and younger.
7.	Intervention/Exposure/Test	Views, opinions and experiences relating to any information, education or support specified in studies
8.	Comparator/Reference standard/Confounding factors	n/a
9.	Types of study to be included	Qualitative studies such as interview and focus group studies (including studies using grounded theory, phenomenology or other appropriate qualitative approaches); quantitative data such as incidence rate or

		frequencies of information preference from questionnaires will also be considered alongside qualitative evidence.
10.	Other exclusion criteria	<p>Exclusions:</p> <ul style="list-style-type: none"> • Adults with subarachnoid haemorrhage caused by head injury, ischaemic stroke or an arteriovenous malformation. • Children and young people aged 15 years and younger. • Non-English language studies.
11.	Context	aSAH is associated with significant sequelae and can have a notable effect on the person's life post-haemorrhage. Adequate information provision for a person with aSAH and their family and/or carer can aid care planning and management, improve understanding and accuracy of expectations, and can influence quality of life. As such, this information for a person with aSAH and their family and/or carer provided both within the immediate care setting and in the community following discharge can be invaluable.
12.	Primary outcomes (critical outcomes)	<p>Themes will be derived from the evidence identified for this review and not pre-specified.</p> <p>Quantitative data such as incidence rate or frequencies of reported information preference will be extracted and presented alongside the themes identified from qualitative analysis.</p>
13.	Secondary outcomes (important outcomes)	n/a
14.	Data extraction (selection and coding)	<p>EndNote will be used for reference management, sifting, citations and bibliographies. All references identified by the searches and from other sources will be screened for inclusion. 10% of the abstracts will be reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer. The full text of potentially eligible studies will be retrieved and will be assessed in line with the criteria outlined above.</p> <p>A standardised form will be used to extract data from studies including the study aim, population setting, design, methodology, findings and limitations (see Developing NICE guidelines: the manual section 6.4).</p>
15.	Risk of bias (quality) assessment	<p>Risk of bias will be assessed using the appropriate checklist as described in Developing NICE guidelines: the manual</p> <p>Critical Appraisal Skills Programme (CASP) qualitative checklist will be used to assess included qualitative studies.</p> <p>Risk of bias for quantitative data will be employed depending on the design of the study:</p> <ul style="list-style-type: none"> • Randomised Controlled Trial: Cochrane RoB (2.0) • Non randomised study, including cohort studies: Cochrane ROBINS-I • Case control study: CASP case control checklist • Controlled before-and-after study or Interrupted time series: Effective Practice and Organisation of Care (EPOC) RoB Tool • Cross sectional study: JBI checklist for cross sectional study

		<ul style="list-style-type: none"> Case series: Institute of Health Economics (IHE) checklist for case series <p>10% of all evidence reviews are quality assured by a senior research fellow. This includes checking:</p> <ul style="list-style-type: none"> papers were included /excluded appropriately a sample of the data extractions correct methods are used to synthesise data a sample of the risk of bias assessments <p>Disagreements between the review authors over the risk of bias in particular studies will be resolved by discussion, with involvement of a third review author where necessary.</p>		
16.	Strategy for data synthesis	<p>The synthesis of qualitative data will follow a thematic analysis approach. Information will be synthesised into main review findings. Results will be presented in a detailed narrative and in table format with summary statements of main review findings.</p> <p>GRADE CERQual will be used to synthesise the qualitative data and assess the certainty of evidence for each review finding.</p> <p>Quantitative data from surveys reporting patient information preferences will be reported narratively and presented alongside thematic analysis. Risk of bias will be assessed to ascertain outcome quality.</p> <p>Endnote will be used for bibliography, citations, sifting and reference management.</p>		
17.	Analysis of sub-groups	n/a		
18.	Type and method of review	<input type="checkbox"/>	Intervention	
		<input type="checkbox"/>	Diagnostic	
		<input type="checkbox"/>	Prognostic	
		<input checked="" type="checkbox"/>	Qualitative	
		<input type="checkbox"/>	Epidemiologic	
		<input type="checkbox"/>	Service Delivery	
		<input checked="" type="checkbox"/>	Other (mixed methods)	
19.	Language	English		
20.	Country	England		
21.	Anticipated or actual start date			
22.	Anticipated completion date	3 February 2021		
23.	Stage of review at time of this submission	Review stage	Started	Completed
		Preliminary searches	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Piloting of the study selection process	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

		Formal screening of search results against eligibility criteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Data extraction	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Risk of bias (quality) assessment	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
		Data analysis	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
24.	Named contact	<p>5a. Named contact National Guideline Centre</p> <p>5b Named contact e-mail SAH@nice.org.uk</p> <p>5e Organisational affiliation of the review National Institute for Health and Care Excellence (NICE) and the National Guideline Centre</p>		
25.	Review team members	<p>From the National Guideline Centre:</p> <ul style="list-style-type: none"> • Ms Gill Ritchie • Mr Ben Mayer • Mr Audrius Stonkus • Mr Vimal Bedia • Ms Emma Cowles • Ms Jill Cobb • Ms Liz Pearton • Ms Amelia Unsworth 		
26.	Funding sources/sponsor	This systematic review is being completed by the National Guideline Centre which receives funding from NICE.		
27.	Conflicts of interest	<p>All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline.</p>		
28.	Collaborators	<p>Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of Developing NICE guidelines: the manual. Members of the guideline committee are available on the NICE website.</p>		

29.	Other registration details	
30.	Reference/URL for published protocol	
31.	Dissemination plans	<p>NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as:</p> <ul style="list-style-type: none"> • notifying registered stakeholders of publication • publicising the guideline through NICE's newsletter and alerts • issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE.
32.	Keywords	Subarachnoid haemorrhage; patient information; lifestyle advice
33.	Details of existing review of same topic by same authors	None
34.	Current review status	<input type="checkbox"/> Ongoing
		<input type="checkbox"/> Completed but not published
		<input type="checkbox"/> Completed and published
		<input type="checkbox"/> Completed, published and being updated
		<input type="checkbox"/> Discontinued
35..	Additional information	
36.	Details of final publication	www.nice.org.uk

1

2 **Table 8: Health economic review protocol**

Review question	All questions where health economic evidence applicable
Objectives	To identify health economic studies relevant to any of the review questions.
Search criteria	<ul style="list-style-type: none"> • Populations, interventions and comparators must be as specified in the clinical review protocol above. • Studies must be of a relevant health economic study design (cost–utility analysis, cost–effectiveness analysis, cost–benefit analysis, cost–consequences analysis, comparative cost analysis). • Studies must not be a letter, editorial or commentary, or a review of health economic evaluations. (Recent reviews will be ordered although not reviewed. The bibliographies will be checked for relevant studies, which will then be ordered.) • Unpublished reports will not be considered unless submitted as part of a call for evidence. • Studies must be in English.
Search strategy	A health economic study search will be undertaken using population-specific terms and a health economic study filter.
Review strategy	Studies not meeting any of the search criteria above will be excluded. Studies published before 2003, abstract-only studies and studies from non-OECD countries or the USA will also be excluded.

Each remaining study will be assessed for applicability and methodological limitations using the NICE economic evaluation checklist which can be found in appendix H of Developing NICE guidelines: the manual.¹⁶

Inclusion and exclusion criteria

- If a study is rated as both 'Directly applicable' and with 'Minor limitations' then it will be included in the guideline. A health economic evidence table will be completed and it will be included in the health economic evidence profile.
- If a study is rated as either 'Not applicable' or with 'Very serious limitations' then it will usually be excluded from the guideline. If it is excluded then a health economic evidence table will not be completed and it will not be included in the health economic evidence profile.
- If a study is rated as 'Partially applicable', with 'Potentially serious limitations' or both then there is discretion over whether it should be included.

Where there is discretion

The health economist will decide based on the relative applicability and quality of the available evidence for that question, in discussion with the guideline committee if required. The ultimate aim is to include health economic studies that are helpful for decision-making in the context of the guideline and the current NHS setting. If several studies are considered of sufficiently high applicability and methodological quality that they could all be included, then the health economist, in discussion with the committee if required, may decide to include only the most applicable studies and to selectively exclude the remaining studies. All studies excluded based on applicability or methodological limitations will be listed with explanation in the excluded health economic studies appendix below.

The health economist will be guided by the following hierarchies.

Setting:

- UK NHS (most applicable).
- OECD countries with predominantly public health insurance systems (for example, France, Germany, Sweden).
- OECD countries with predominantly private health insurance systems (for example, Switzerland).
- Studies set in non-OECD countries or in the USA will be excluded before being assessed for applicability and methodological limitations.

Health economic study type:

- Cost–utility analysis (most applicable).
- Other type of full economic evaluation (cost–benefit analysis, cost-effectiveness analysis, cost–consequences analysis).
- Comparative cost analysis.
- Non-comparative cost analyses including cost-of-illness studies will be excluded before being assessed for applicability and methodological limitations.

Year of analysis:

- The more recent the study, the more applicable it will be.
- Studies published in 2003 or later but that depend on unit costs and resource data entirely or predominantly from before 2003 will be rated as 'Not applicable'.
- Studies published before 2003 will be excluded before being assessed for applicability and methodological limitations.

Quality and relevance of effectiveness data used in the health economic analysis:

- The more closely the clinical effectiveness data used in the health economic analysis match with the outcomes of the studies included in the clinical review the more useful the analysis will be for decision-making in the guideline.

1 Appendix B: Literature search strategies

2 This literature search strategy was used for the following review;

- 3 • What patient information (including lifestyle advice) should be given to adults who have
4 had an aneurysmal subarachnoid haemorrhage?

5 The literature searches for this review are detailed below and complied with the methodology
6 outlined in Developing NICE guidelines: the manual¹⁶

7 For more information, please see the Methods Report published as part of the accompanying
8 documents for this guideline.

B.1.9 Clinical search literature search strategy

10 Searches for patient views were run in Medline (OVID), Embase (OVID), CINAHL and PsycINFO
11 (ProQuest). Search filters were applied to the search where appropriate.

12 **Table 9: Database date parameters and filters used**

Database	Dates searched	Search filter used
Medline (OVID)	1946 – 26 June 2020	Exclusions Qualitative studies
Embase (OVID)	1974 – 26 June 2020	Exclusions Qualitative studies
CINAHL, Current Nursing and Allied Health Literature (EBSCO)	Inception – 26 June 2020	Qualitative studies
PsycINFO (ProQuest)	Inception – 26 June 2020	Qualitative studies

13 Medline (Ovid) search terms

1.	exp Subarachnoid Hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	Intracranial Aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial or brain) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.
15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16

18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23
25.	6 not 24
26.	(exp child/ or exp pediatrics/ or exp infant/) not (exp adolescent/ or exp adult/ or exp middle age/ or exp aged/)
27.	25 not 26
28.	limit 27 to English language
29.	Information centers/
30.	information services/ or library services/
31.	Information Dissemination/
32.	Libraries/
33.	health education/
34.	publications/ or books/ or pamphlets/
35.	Patient Education Handout/
36.	patient education as topic/
37.	consumer health information/
38.	counseling/ or directive counseling/ or distance counseling/
39.	psychosocial support systems/
40.	Needs Assessment/
41.	life style/
42.	Social Support/
43.	Adaptation, Psychological/
44.	Financial Support/
45.	or/29-44
46.	((patient* or user* or consumer*) adj4 (educat* or literature or leaflet* or book* or pamphlet* or fact sheet* or factsheet* or publication* or librar* or inform* or advice or need* or requirement* or support* or service* or communication* or involv)).ti,ab.
47.	((patient* or user* or consumer*) adj4 (attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact*)).ti,ab.
48.	((patient* or user* or consumer) adj4 (financial* or income or earning* or psych* or work* or job or employ* or social)).ti,ab.
49.	(lifestyle* or life style*).ti,ab.
50.	((inform* or advice or educat* or learn* or support* or financial* or counsel* or psych* or mental health or work* or job* or employ* or social) adj4 (service* or literature or leaflet* or booklet* or pamphlet* or manual* or brochure* or publication* or handout* or fact sheet* or factsheet* or material* or program* or service* or centre* or center* or hub* or need* or requirement* or support* or seek* or access* or disseminat* or barrier*)).ti,ab.
51.	((financial* or income or earning* or psych* or counsel* or mental health or work* or job* or employ* or social or coping) adj4 (inform* or educat* or learn* or help or service* or need* or requirement* or support* or communication* or involv* or attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact*)).ti,ab.
52.	((return* or back) adj2 (work* or job or employ*)).ti,ab.

53.	or/46-52
54.	28 and (45 or 53)
55.	Qualitative research/ or Narration/ or exp Interviews as Topic/ or exp "Surveys and Questionnaires"/ or Health care surveys/
56.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*).ti,ab.
57.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* adj3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*).ti,ab.
58.	or/55-57
59.	54 and 58

1 Embase (Ovid) search terms

1.	*subarachnoid hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp intracranial aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	Case report/ or Case study/
11.	(letter or comment*).ti.
12.	or/7-11
13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	Nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental animal/
19.	Animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	(exp child/ or exp pediatrics/) not (exp adult/ or exp adolescent/)
25.	23 not 24
26.	limit 25 to English language
27.	information service/
28.	documentation/
29.	publication/
30.	book/

31.	library/
32.	information dissemination/
33.	health education/
34.	publication/
35.	book/
36.	patient education/
37.	counseling/ or anticipatory guidance/ or directive counseling/ or e-counseling/ or patient counseling/ or patient guidance/ or peer counseling/
38.	social support/
39.	psychosocial care/
40.	needs assessment/
41.	lifestyle/ or healthy lifestyle/ or lifestyle modification/
42.	coping behavior/
43.	consumer health information/
44.	or/27-43
45.	((patient* or user* or consumer*) adj4 (educat* or literature or leaflet* or book* or pamphlet* or fact sheet* or factsheet* or publication* or librar* or inform* or advice or need* or requirement* or support* or service* or communication* or involv)).ti,ab.
46.	((patient* or user* or consumer*) adj4 (attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact*)).ti,ab.
47.	((patient* or user* or consumer) adj4 (financial* or income or earning* or psych* or work* or job or employ* or social)).ti,ab.
48.	(lifestyle* or life style*).ti,ab.
49.	((inform* or advice or educat* or learn* or support* or financial* or counsel* or psych* or mental health or work* or job* or employ* or social) adj4 (service* or literature or leaflet* or booklet* or pamphlet* or manual* or brochure* or publication* or handout* or fact sheet* or factsheet* or material* or program* or service* or centre* or center* or hub* or need* or requirement* or support* or seek* or access* or disseminat* or barrier*)).ti,ab.
50.	((financial* or income or earning* or psych* or counsel* or mental health or work* or job* or employ* or social or coping) adj4 (inform* or educat* or learn* or help or service* or need* or requirement* or support* or communication* or involv* or attitud* or priorit* or perception* or preferen* or expectation* or choice* or perspective* or view* or satisfact*)).ti,ab.
51.	((return* or back) adj2 (work* or job or employ*)).ti,ab.
52.	or/45-51
53.	26 and (44 or 52)
54.	health survey/ or exp questionnaire/ or exp interview/ or qualitative research/ or narrative/
55.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*).ti,ab.
56.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* adj3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*).ti,ab.
57.	or/54-56
58.	53 and 57

1 CINAHL (EBSCO) search terms

51.	SU Subarachnoid Hemorrhage
-----	----------------------------

S2.	TI (((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial) n3 (hemorrhag* or haemorrhag* or bleed* or blood*))) OR AB (((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial) n3 (hemorrhag* or haemorrhag* or bleed* or blood*)))
S3.	TI ((SAH or aSAH)) OR AB ((SAH or aSAH))
S4.	TI (((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial or brain) n3 (aneurysm* or aneurism* or hematoma* or haematoma*))) OR AB (((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial or brain) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.)
S5.	S1 OR S2 OR S3 OR S4
S6.	(MH "Qualitative Studies+")
S7.	(MH "Qualitative Validity+")
S8.	(MH "Interviews+") OR (MH "Focus Groups") OR (MH "Surveys") OR (MH "Questionnaires+")
S9.	(qualitative or interview* or focus group* or theme* or questionnaire* or survey*)
S10.	(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* adj3 analys*) or theoretical sampl* or purposive sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*)
S11.	S6 OR S7 OR S8 OR S9 OR S10
S12.	S5 AND S11

1 PsycINFO (ProQuest) search terms

1.	(ti(((subarachnoid* OR arachnoid* OR cerebral OR intracerebral OR intra-cerebral OR intracranial OR intra-cranial) NEAR/3 (hemorrhag* OR haemorrhag* OR bleed* OR blood*))) OR ab(((subarachnoid* OR arachnoid* OR cerebral OR intracerebral OR intra-cerebral OR intracranial OR intra-cranial) NEAR/3 (hemorrhag* OR haemorrhag* OR bleed* OR blood*))) OR ti(((subarachnoid* OR arachnoid* OR cerebral OR intracerebral OR intra-cerebral OR intracranial OR intra-cranial OR brain) NEAR/3 (aneurysm* OR aneurism* OR hematoma* OR haematoma*))) OR ab(((subarachnoid* OR arachnoid* OR cerebral OR intracerebral OR intra-cerebral OR intracranial OR intra-cranial OR brain) NEAR/3 (aneurysm* OR aneurism* OR hematoma* OR haematoma*))) OR MAINSUBJECT.EXACT.EXPLODE("Subarachnoid Hemorrhage") OR ti(SAH OR aSAH) OR ab(SAH OR aSAH) AND ((su.exact.explode("qualitative methods") or su.exact("narratives") or su.exact.explode("questionnaires") or su.exact.explode("interviews") or su.exact.explode("health care services") or ti,ab(qualitative or interview* or focus group* or theme* or questionnaire* or survey*) or ti,ab(metasynthes* or meta-synthes* or metasummar* or meta-summar* or metastud* or meta-stud* or metathem* or meta-them* or ethno* or emic or etic or phenomenolog* or grounded theory or constant compar* or (thematic* near/3 analys*) or theoretical-sampl* or purposive-sampl* or hermeneutic* or heidegger* or husserl* or colaizzi* or van kaam* or van manen* or giorgi* or glaser* or strauss* or ricoeur* or spiegelberg* or merleau*)))
----	---


B.2.2 Health Economics literature search strategy

3 Health economic evidence was not required for this review.

1 Appendix C: Qualitative evidence

2 selection

Figure 1: Flow chart of qualitative study selection for the review of Patient information



3

4

1 Appendix D: Qualitative evidence tables

2

Study	Berggren 2010 ³
Aim	<p>The aim was to analyse peoples accounts of SAH and to describe how the initiate and create meaning for onset and events surrounding the SAH. Specific questions were:</p> <ul style="list-style-type: none"> - What is highlighted in the accounts of SAH? - How is the illness reconstructed? - How is meaning created through communicative interaction with others about SAH?
Population	<p>Nine people who had experienced and were treated for SAH at a university hospital in Sweden were interviewed. Informants who were contactable and native language speaking at the time of their discharge were selected from the hospitals patient record system.</p> <p>Characteristics: n=9; 3 male/ 6 female; mean age (range) 33 - 67; interviews took place 1 year and 7 months after the SAH.]</p>
Setting	Participants homes
Study design	Qualitative
Methods and analysis	<p>In-depth semi-structured interviews with thematic qualitative analysis.</p> <p>Discourse analysis was used to describe meaning – making in the accounts of the experience onset of a SAH. The text was read several times to acquire a full appreciation of the data and to elucidate patterns. Both the information (what), and the interaction (how) were studied according to the model of Linell and Thunkvist (2003). Interesting content and statements which occurred together were grouped together in content and topics and then subjected to discourse analysis. Sequences and episodes were extracted from the accounts and critical events were identified. The sequences and episodes were then analysed with the support of data extracts. The accounts were rich in fact and interest constructions. The extracts were translated from Swedish to English by a native English speaker so that the subtleties and meaning from the original language were not lost.</p>
Findings	<p>SAH event</p> <p>a) Memory – informants report memory and memory gaps for the event of SAH and the period following the SAH. They may suddenly remember what bits of information they were missing but heavily relied on their support systems to inform them of what had happened. There are differences in understanding related to the event, where some informants remember the symptoms related to the SAH clearly which was linked their own survival and an existential crisis.</p>
	<p>Support</p> <p>a) Different sources of support – extracts included in the review identify that informants relied on information provided by relatives, friends or carer’s to understand what actually happened to them medically and to find some sense of meaning.</p>

Study	Berggren 2010³
Limitations and applicability of evidence	A study based in Sweden with some methodological limitations, due to their line questioning to understand how people initiate and create meaning. The study focused on specific topics but results appear to be generally applicable. The authors have focused on very specific outcomes and areas of interest such as finding meaning after an SAH through communication. The relevant data has been extracted for the review.

1

Study	Dulhanty 2019⁶		
Aim	Explore the type and frequency of self reported needs both early and late following SAH specifically to the extent to which they have been met.		
Population	A census sample of people who had experienced a subarachnoid haemorrhage (SAH) in Greater Manchester between September 2010–September 2012 (late post-SAH group) and September 2013–September 2014 (early post-SAH group) were recruited. All adult patients admitted to the Regional Neurosciences Centre with SAH within these dates, who survived and were discharged home from the neurosciences centre or inpatient rehabilitation were identified from the centre’s prospectively maintained clinical database. People requiring nursing home or residential care were excluded as it is assumed that their needs are being met and were unlikely to be able to consent.		
	Variable	1 – 2 years post SAH (n=122)	3 – 5 years post SAH (n=81)
	Age, mean (SD)	55.0 (11.6)	55.4 (10.5)
	Male / female	45/77	31/50
	WFNS Grade	Grade 1 – 83	Grade 1 – 54
		Grade 2 – 21	Grade 2 – 18
		Grade 3 – 3	Grade 3 – 3
		Grade 4 – 9	Grade 4 – 5
		Grade 5 – 6	Grade 5 – 1
	Fisher Grade	Grade 1 – 19	Grade 1 – 7
		Grade 2 – 18	Grade 2 – 1
		Grade 3 – 22	Grade 3 – 33
		Grade 4 – 63	Grade 4 – 40
	Treatment	Coiled – 78	Coiled – 47
		Clipped – 16	Clipped – 10
		None – 28	None – 24
	Length of stay (IQR)	13.0 (8 to 20.25)	13.0 (9 to 21.5)

Study	Dulhanty 2019⁶		
Setting	Regional Neurosciences centres within Greater Manchester		
Study design	Cross sectional postal survey		
Methods and analysis	Potential participants were sent an invitation letter followed by a study pack containing the patient information sheet, data collection tool, a request for researchers to access the clinical data stored on the centre's database and a stamped-return envelope. They were also given details of how to access the same questionnaire online to submit their data electronically if they preferred, or they could arrange to complete the questionnaires by telephone interview.		
Findings	Patient information needs:		People reporting info need: N (%)
	Health	<ul style="list-style-type: none"> • SAH • General health (diet / alcohol / smoking) 	105 (58) 40 (21)
	Travel	<ul style="list-style-type: none"> • Driving • Public Transport • Benefits • Money management 	19 (10) 22 (11) 34 (17) 7 (4)
	Home	<ul style="list-style-type: none"> • Home aids • Home adaptations • Moving home • Physical relationships 	20 (10) 10 (5) 12 (6) 17 (9)
Limitations and applicability of evidence	<p>This paper had reported some quantitative results directly relevant to patient information needs by respondents.</p> <p>A study based in the UK. The quantitative results were used to understand requirements for patient information and are generally applicable. Moderate risk of bias – outcome measurement bias 203 (51%) participants responded to the questionnaire. The relevant data has been extracted for the review.</p>		

1

Study	Hedlund 2010⁸
Aim	The study constitutes the qualitative part of an ongoing study investigating psychiatric morbidity, cognitive strategies, coping and quality of life in patients after SAH. The aims of this study were to describe what participants with no or minor neurological deficits after SAH perceived as being important for recovery, and perceived consequences of SAH.
Population	Men and women of different ages and from different places in the catchment area who had undergone different treatment for SAH and with no or minor neurological impairments at discharge from the neurosurgical unit.

Study	Hedlund 2010 ⁸
	<p>N=20</p> <p>The median age of the participants was 51 years (range 30 – 64 years). On arrival to the neurosurgical clinic, 18 patients were fully alert, one patient scored 2 and one patient scored 3 (Reaction Level scale RL 85). The most common localization of the aneurysm was arteria communicans anterior. At the time of the qualitative interview, 18 participants lived with spouses, 10 had children at home and 2 participants lived alone. Eleven participants had returned to work, one with an adjusted workplace situation. Five participants were still on sick leave, one participant retired due to age and one retired due to disability prior to the onset of SAH. Only one of the 20 participants had continuous contact with a neurological rehabilitation clinic at the time of the interview.</p>
Setting	<p>The interviews were conducted at the neurosurgical outpatient clinic or in the participants' home on average 12 months after the onset of SAH</p>
Study design	<p>Semi-structured qualitative review</p>
Methods and analysis	<p>A semi structured interview guide covering the following question areas: (1) what the participants perceived as important for recovery from SAH, and (2) perceived consequences of SAH.</p> <p>The data analysis was conducted in two steps:</p> <ol style="list-style-type: none"> 1) To explore the research questions, a qualitative manifest content analysis inspired by Graneheim and Lundman was used as a first step. All interviews were transcribed verbatim. The interview text was read through several times to become familiar with the data and acquire an overall understanding of the text. Discussions on the content of the text in relation to the aim of each research question were held to reach consensus. Meaning units (related by their context and content) were identified, condensed (shortened with preserved core), abstracted and labelled with a code. All codes were compared and sorted into categories based on shared similarities. 2) After categorisation, the authors went backwards through the codes and meaning units to identify the person behind each statement to see if the person had been previously diagnosed with depression. When going forwards again the authors revealed that some categories were synthesized exclusively of codes and meaning units from depressed participants while some of the other categories were exclusively synthesized of codes and meaning units from participants without depression.
Findings	<p>Clarity of information</p>
	<p>a) SAH event – those who continually suffered from memory loss wanted to get more information about SAH and the course of events that led to their SAH from their physician</p>
	<p>b) Symptoms – informants suffered from multiple consequences from their SAH which affected their cognitive, physical, psychosocial and emotional functioning. The findings of these consequences was linked to how much information and support they felt they received after the SAH from families or informal care givers.</p>

Study	Hedlund 2010⁸
	c) Medical information – Participants generally felt that there was insufficient follow up from their local medical providers once they were discharged from the hospital. Some reported not receiving any rehabilitation support and felt unable to ask for support.
	Support
	a) Different types of support – participants were grateful for the care that they had received in hospital. Majority of participants were grateful to their families or care givers for constant support after their SAH, especially once discharged. Some, were worried about the responsibilities at home that were now upon their families and felt worried about being burdensome. These participants were also worried about expressing the ongoing symptoms they were facing, weakening relationships with significant others or families for not fully understanding what they were experiencing.
	Future
	a) Long term implications – most participants were eager to return to their “normal” lives and were slowly adjusting to their new reality. Despite the physical consequences they were making efforts to try and return to their day to day lives. But there were some participants who longed for their old lives, wanting to know when they will be able to go back to “normal” so they don’t feel dependant or have constant uncertainty.
	b) Reoccurrence – Some participants were worried about headaches, follow up scans and procedures as they related these experiences to SAH or potentially to having another SAH
Limitations and applicability of evidence	A study based in Sweden with some methodological limitations due to the consolidation of multiple concerns under one themes and the focus on specific topics in relation to depression or no depression, but results appear to be generally applicable. The relevant data has been extracted for the review.

1

Study	Jarvis 2002¹¹
Aim	The study aimed to illuminate the experience of recovery from SAH.
Population	Patients who had suffered a SAH within the last 14 – 18 months. Patients were excluded if they were continuing to receive treatment of any description. Characteristics: n=8 ; 6 male/ 2 female; age (range) 30 – 80;
Setting	Interviews took place in the participants homes
Study design	Qualitative interviews with a semi – structured approach
Methods and analysis	The researcher opted for an inductive data analysis, however accepted that specific themes in relation to the objective of the review already existed. These were: illness understanding or knowledge; problem experience; experiences of recovery; questions and queries and feelings or emotions. Therefore, the researcher opted for deductive method of analysis. This is where previously identified themes are examined for their presences within data, sometimes referred to as thematic analysis.

Study	Jarvis 2002 ¹¹
	While carrying out the deductive analysis it was vital for the researcher to note any new individual issues that arose in order to keep the link with the process of discovery or inductive analysis. Secondary analysis of the data, inductive analysis, ensured that important content would not be missed or left undiscovered. The third stage of analysis was a validation process and took the form of a brief interpretative account of the patient's story.
Findings	Clarity of information
	<p>a) SAH event – some participants were extremely scared about the potentially fatal incident they had undergone but wanted more information and know exactly what had happened to them so they could piece things together and fill the gaps in their memory. Whereas other participants instead felt a sense of anxiety and dread having such large memory gaps and not feeling in control of their health.</p>
	<p>b) Ongoing symptoms – participants felt uninformed about the persistent symptoms that they would face once discharged home after treatment for the SAH. The main physical consequence was fatigue / tiredness, which was considered to be an invisible complication because it couldn't be seen by others.</p>
	Future
	<p>a) Long term implications – Informants felt that some consequences of SAH persisted beyond the initial recovery period and impacted their lives at a greater scale. This was related to ongoing tiredness, lack of concentration and inability to handle multiple tasks (e.g. at a place of work), which lead to dependence and unemployment.</p>
	<p>b) Reoccurrence – Participants were fearful of reoccurrences of SAH. This was closely linked to the memory of having the initial event - the fear of not knowing what was happening or happened.</p>
Limitations and applicability of evidence	A study with some methodological limitations based in UK. Some themes were put forth to participants and others were offered by participants themselves through questioning. The study focused on specific topics in relation to symptoms or stressful experiences post SAH, but results appear to be generally applicable. The relevant data has been extracted for the review.

1

Study	Persson 2017 ²¹
Aim	To explore experiences of care and rehabilitation as well as the consequences and strategies used to cope with everyday life six years after SAH.
Population	Participants were included from the Extended Stroke Arm Longitudinal study at the University of Gothenburg. The inclusion criteria were > 18 years of age, with an SAH between 4th February 2009 until the 2nd of December 2010, receiving care at the Sahlgrenska University Hospital in Gothenburg, Sweden, Resident in the Gothenburg Urban area (<35km from the hospital), able to speak Swedish and at least some participants should be of working age. The participants were contacted by phone and if they agreed to participate, a time and interview was planned.

Study	Persson 2017²¹
	[Characteristics: n=16; 8 male/ 8 female; mean age (range) 57 (45-76);
Setting	Participant chose location
Study design	Qualitative descriptive design
Methods and analysis	<p>Explorative interview with a qualitative descriptive design, using an inductive driven thematic analysis.</p> <p>All interviews were performed face-to-face using an interview guide with open ended questions: 1) What type of treatment (acute and rehabilitation) did you receive? 2) What type of consequences (such as cognitive and physical) do/did you have post SAH? 3) What strategies have you used to cope with the consequences of SAH in daily life? 4) What impact do the consequences post SAH have on your social function and in daily life? The interview guide was discussed among the authors prior to the interviews, and some questions were changed/added.</p> <p>The first interview was a pilot for the interview guide, and the questions seemed to cover the research areas of interest. The interviews took place on average six years (SD 0.5) post SAH.</p> <p>The data was analysed according to thematic analysis described by Braun and Clarke as a flexible method. The thematic analysis used in this study was inductive and can be referred to a realist/essentialist paradigm.</p>
Findings	Clarity of information
	a) SAH event – while some participants felt grateful for the care they have received, others felt that they were not cared for properly and were not adequately informed about the SAH event and their treatment pathway. Patients also worried about the implications of the event on their families and this wasn't helped by having gaps in their memory and not having a proper understanding of what had happened to them.
	b) Medical information – many patients felt like they were abandoned after discharge from the acute neurosurgical unit and were not given enough information about cognitive symptoms they could face which became evident only once they were discharged. They felt they did not have enough knowledge about the consequences of SAH and wanted more information about the course of the illness prior to discharge. Experiences of medical support after discharge also varied, where some had follow-ups arranged. Others did not have any pre-arranged follow-ups and had to contact their medical providers for support themselves, which added to the feeling of abandonment.
	Conversely, those patients who had been referred to a rehabilitation centre felt quite supported and they were given the right level of information that they required. But this group of patients also felt unclear about the medical follow ups after the discharge from these facilities and would have liked to be followed up.
	Support
	a) Different sources of support – Patients received various levels of support from families and care givers to rehabilitation clinics as well. Patients wished that their primary care provider had more information about SAH, so they could be a source of information and support as this was lacking. They were grateful for the support that they had received but were worried about the feeling of being a burden and having to make major adaptations to their lives to accommodate the consequences of SAH.

Study	Persson 2017²¹
	Future
	a) Long term implications – participants would have liked to know about the long-term implications of SAH, to better understand how their lives would be going forward. On top of the cognitive symptoms, some participants had to change their homes, lost their licenses, and became unemployed
	b) Coping – patients wanted to get back to their daily lives and routines and most were able to do so. Some constantly knew the SAH had changed their life, so tried not to talk about too much and hid symptoms to avoid any extra attention with their friends or even in their workplace. Coping was linked to the level of information they knew about the long-term consequences of SAH. From a practical and physical level, many employers were unaware of what their employees would need to deal with the SAH consequences in the workplace. Coping at the workplace varied for participants from some feeling really well supported, to others feeling neglected due to no knowledge of SAH.
Limitations and applicability of evidence	A well conducted study based in Sweden. Focused on the consequences of SAH and the coping strategies patients use to cope. Results are generally applicable.

1
2

Study	von Vogelsang 2004²⁵	
Aim	An investigation into the effects of increased information for patients treated for intracranial aneurysm rupture.	
Population	Participants were 62 patients with ruptured intracranial aneurysm treated consecutively at a Swedish neurosurgical clinic. Participants were recruited over a period of 12 months and recruitment was concluded in 2001. All patients in the sample were acute admissions; and those treated for non-ruptured aneurysms were not included. Patients were included if they were be able to understand, speak and read Swedish, and had a score of 4–5 on the Glasgow Outcome Scale (GOS).	
	Variable	Comparison group (n=34)
	Age in years, mean (range)	53.7 (32–78)
	Male	14
	Female	20
	Days spent in institutional care, mean (range)	20.6 (9–54)
	Glasgow Outcome Scale, (range 1–5), mean (SD)	4.4 (0.6)
		Intervention group (n=28)
		56.2 (35–76)
		6
		22
		20.9 (10–35)
		4.5 (0.6)

Study	von Vogelsang 2004 ²⁵			
	Surgical treatment	30	25	
	Endovascular treatment	3	3	
	Both surgical and endovascular treatment	1	0	
Setting	Swedish neurosurgical clinic			
Study design	Non randomized comparative study			
Methods and analysis	<p>The intervention group (n = 28) received oral as well as written information, while the comparison group (n = 34) received only oral information.</p> <p>Existing written information, produced by neurosurgical clinics and patient associations, was not considered suitable, hence a new information leaflet was written. The content included explanations of the medical condition and treatment, diagnostic procedures, risk factors, physical and psychological symptoms after treatment, explanation of key terms, and addresses of patient associations. The content was determined by suggestions from a neurosurgeon specializing in intracranial vascular diseases, and by a nurse with 4 years' experience of follow-up interviews with patients who had experienced an intracranial aneurysm. Care was taken to make the written information as easy to read and understand as possible.</p> <p>When the comparison group was complete, the information leaflet was distributed consecutively to patients in the intervention group, who received the same questionnaire and informative letter, and were subject to the same time interval and reminder call as the comparison group. When the information leaflet was distributed, a delivery note was included and the patient's data were recorded on this by the person who distributed the leaflet. The delivery note was handed over to the nurse who conducted the study as a receipt. The delivery note was included to ensure that every patient in the intervention group got the information leaflet.</p>			
Findings	Question/instrument	Comparison group	Intervention group	P value
	Question 1: Information difficult/easy to understand? (range 1–5: the lower the score the more difficult to understand), mean (SD)	3.3 (1.9)	3.9 (1.8)	0.24
	Question 2: Information corresponding/not corresponding to patients' needs for information? (range 1–5: the lower the score the less correspondence with patients' needs for information), mean (SD)	3.3 (1.6)	4.4 (2.9)	0.06
	Clarity of information			

Study	von Vogelsang 2004 ²⁵
	<p>a) SAH event – participants preferred written information to refer back to as most of them did not remember any oral information due to difficulties with memory in the acute phase after an SAH. Families were the initial source of information and some participants felt they couldn't assimilate any information due to the shock.</p> <p>b) Medical information – respondents generally wanted a better quality of information from all medical professionals. Topics such as SAH, heredity, surgical techniques, rehabilitation, and future risks were highlighted as important topics to be covered better in information given to patients, families and carers.</p>
Limitations and applicability of evidence	<p>Paper reported both qualitative and quantitative data which has been extracted for the review.</p> <p>A study based in Sweden. This study was comparing the use of written and oral information compared to oral information only. This was investigated using self report questionnaires. The quantitative information used from this study was elicited using a Likert style response format. However, open ended questions were also used to get more a richer understanding of the participants experiences. Potential selection bias with intervention start and follow-up start differing between intervention and control groups, and potential confounding bias with no adjusting for possible confounding factors between control and intervention groups (ROBINS-I).</p> <p>Overall, the results are generally applicable.</p>

1 Appendix E: Excluded studies

E.1.2 Excluded qualitative studies

3 Table 10: Studies excluded from the qualitative review

Reference	Reason for exclusion
Berggren 2010 ¹	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Berggren 2011 ²	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Cedzich 2005 ⁴	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Covey 2013 ⁵	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
He 2014 ⁷	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Hellawell 2001 ⁹	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Hutter 2014 ¹⁰	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Karic 2016 ¹²	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
King 2006 ¹³	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Morris 2004 ¹⁴	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Murgo 2016 ¹⁵	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Nishino 1999 ¹⁷	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Nordenmark 2020 ¹⁸	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Ogden 1997 ¹⁹	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Passier 2013 ²⁰	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Preiss 2012 ²²	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Ross 2013 ²³	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice
Stegen 1991 ²⁴	No relevant outcome - patient outcome data, no reference to information needs or lifestyle advice

4

5