

Otitis media with effusion in under 12s

Review questions

NICE guideline tbc

Review questions

March 2023

Early submission

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2023. All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

Review questions	5
-------------------------------	----------

Review questions

Review topic	Review question number in the scope	Review question
[1] Risk factors for OME	1.1	What are the modifiable risk factors for developing OME in children under 12 years?
[2] Recognition of OME (to help identify when to refer for further investigation)	2.1	What presenting features are associated with OME in children under 12 years?
[3] Natural history of OME (to help identify when intervention and follow-up is needed)	3.1	What is the progression, resolution and recurrence (natural history) of OME without hearing loss at presentation in children under 12 years?
[3] Natural history of OME (to help identify when intervention and follow-up is needed)	3.2	What is the progression, resolution and recurrence (natural history) of OME-related hearing loss at presentation in children under 12 years?
[4] Interventions for children with OME	4.1	What is the effectiveness of ventilation tubes for managing OME with associated hearing loss in children under 12 years?
[4] Interventions for children with OME	4.2	What is the effectiveness of adenoidectomy (with or without hearing loss) for managing OME with associated hearing loss in children under 12 years?
[4] Interventions for children with OME	4.3	What is the effectiveness of antibiotics for managing OME with associated hearing loss in children under 12 years?
[4] Interventions	4.4	What is the effectiveness of non-antimicrobial pharmacological interventions (such as steroids, antihistamines, leukotriene

for children with OME		receptor antagonists, mucolytics and decongestants) for managing OME in children under 12 years?
[4] Interventions for children with OME	4.5	What is the effectiveness of auto-inflation for managing OME with associated hearing loss in children under 12 years?
[4] Interventions for children with OME	4.6	What is the effectiveness of air conduction and bone conduction hearing aids/devices for hearing loss associated with OME in children under 12 years?
[5] Care during and after surgery	5.1	What intraoperative or postoperative interventions are effective at preventing otorrhoea (ear discharge) after surgery for OME-related hearing loss in children under 12 years?
[5] Care during and after surgery	5.2	What interventions are effective for treating otorrhoea (ear discharge) after surgery for OME-related hearing loss in children under 12 years?
[5] Care during and after surgery	5.3	What should the follow-up strategy be after surgical treatment for OME-related hearing loss in children under 12 years?
[6] Information for children, parents and carers	6.1	What information is valued by children under 12 years with suspected or confirmed OME and their parents and carers?