

## **Economics, Evidence review, Appendix C1**

### **– Selecting areas for economic analysis**

#### **Guideline**

Transition between inpatient mental health settings and community and care home settings

This report was produced by the Personal Social Services Research Unit at the London School of Economics and Political Science. PSSRU (LSE) is an independent research unit and is contracted as a partner of the NICE Collaborating Centre for Social Care (NCCSC) to carry out the economic reviews of evidence and analyses.

Authors of the report: Marija Trachtenberg and Martin Knapp.

## **1 Introduction – Economic work as part of guideline development**

The economic work is comprised of 2 main components. The first is the critical appraisal and review of existing cost-effectiveness literature and interpreting the results to make recommendations for the UK context. These can be found in Appendices C1 and C2 and are not the focus of this report.

The second component is undertaking new economic analyses. This report shows the decision-making process leading up to the decision to conduct new economic analyses. Those new analyses are presented in economic Appendices C3.2 and C3.3 and are not covered in this report.

## **2 Methods to select areas for new economic analysis**

New economic analyses are useful where the existing cost-effectiveness literature is not sufficient to make recommendations for the UK context. However, decisions must also take into account data availability and whether it is feasible to conduct new analyses. Also, as there are potentially many areas on which to do new analyses, prioritisation is necessary. Prioritising one area over another is based on the expectation that the new analysis will help reduce uncertainty associated with an intervention's cost-effectiveness and that the recommendations coming from the analysis has a significant impact on social care outcomes and costs. These criteria are also considered alongside Guideline Committee preferences.

## **3 Results – Economic analysis for this guideline**

The two economic analyses carried out for this guideline relate to review questions 4 and 5. They look at interventions that improve the discharge process and reduce readmissions to hospital. The rationale is that recommendations would have significant impact on social care costs and outcomes. Evidence was of good methodological quality, data was available for the analysis and the Guideline Committee expressed agreement and preference for these areas.

The other review questions were not prioritised for economic analysis because:

- There was a lack of economic evidence and the single effectiveness paper identified (Goldberg et al. 2013) planned to publish an economic analysis (email communication with study authors) and the Guideline Committee did not express preference for this area.<sup>1</sup>
- There was a lack of both effectiveness and economic evidence.<sup>2</sup>

---

<sup>1</sup> Review Question 4: Approaches to care planning and assessment for admission to hospital.

<sup>2</sup> Review Questions 7, 8, 9, 10: Interventions to support people living with dementia during transition/Interventions to support children and young people/Interventions to support carers/Learning and development for mental health and social care staff.

- The review questions did not involve a decision between interventions and therefore are not relevant for economic analysis.<sup>3</sup>

### **3.1 Review Questions 4 and 5 – selecting interventions for economic analysis**

Due to the timing of the evidence review, the first economic analysis was agreed with the chair in advance of the Guideline Committee meeting. If a decision had been left until the seventh and eighth Guideline Committee meetings (29<sup>th</sup>/30<sup>th</sup> September), there would not have been enough time to complete the analysis. However, if one analysis were to be agreed beforehand (based on the studies already available to the economist) there would be enough time to complete two analyses. Such an approach was agreed with the chair.

#### 1. Rationale and method for selecting the first economic analysis

Based on the available evidence, the first economic analysis was based on Kessing et al. (2013), a moderately sized RCT from Denmark (n=158) rated as having moderate quality internal validity (+) and good external validity (++) . It focuses on bipolar patients in the early stages of the disorder. The study was selected on the basis that the intervention was associated with significant reductions in the use of hospitalisation and the time horizon was sufficiently long (2.5 years). It was thought that an economic analysis would be useful because there was some evidence (but not definitive) that patients in the earlier stages of bipolar disorder have greater potential to benefit (Reinares 2014 citing Colom et al. 2010; Scott et al. 2006).<sup>4</sup> Furthermore, there was uncertainty as to the trade-offs between the costs of the intervention (which is relatively resource intensive, as it is delivered over a 2-year period) and whether the benefits of such an intervention would be cost-effective in the English context. For in-depth rationale, please refer to the economics Appendix C3.2, under the section, 'Background – why the analysis is important'.

#### 2. Rationale and method for selecting the second economic analysis

The second analysis was decided with the Guideline Committee on the seventh and eighth meetings. At the time, the Guideline Committee had reviewed the available evidence and had decided to make a general recommendation on peer support workers; there was some discussion as to the type of peer support, as some members of the Guideline Committee had various experiences and ideas of types of peer support, which could be provided in particular contexts. Some of this discussion touched on the one moderate quality RCT from United States on the use of peer support on discharge (Sledge et al. 2011). There was discussion on the role of peer-delivered programmes to which the Guideline Committee had a strong preference that the economic analysis focus on peer support because this

---

<sup>3</sup> Review Questions 1, 2, 3: Views and experiences from service users, families and carers, and practitioners.

<sup>4</sup> Specifically, interventions become less effective in reducing time until next relapse or the time spent ill as the number of bipolar episodes increases.

topic, which potentially benefits the person giving support as well as the recipient, is consistent with recovery and recovery-oriented interventions. To aid the Guideline Committee, three recent systematic reviews that looked at peer support were searched. Unfortunately, none of the reviews were specifically about transition, but interventions were still relevant. Of the three systematic reviews, one review synthesised the evidence more narrowly, meaning that similar interventions were grouped together, and results were reported using meta-analytic techniques (Fuhr et al. 2014). The other two systematic reviews grouped interventions that had fewer similarities and so these were considered unsuitable for use in an economic analysis (Lloyd-Evans et al. 2014; Pitt et al. 2013).

Meta-analytic results were available for a peer-delivered self-management intervention (as identified in Fuhr et al. 2014). The results were based on three high-quality non-UK studies where samples and interventions were of acceptable similarity. The studies in the meta-analysis measured impact on quality of life, hope and clinical symptoms, but did not measure readmissions to hospital. The intervention was delivered among a sample of community-dwelling individuals (and not necessarily those recently discharged from hospital) but the Guideline Committee did not think that was a problem, as they believed that discharged patients could still benefit from this intervention as a part of improving the discharge process. Economic analysis was considered useful due to the lack of cost-effectiveness evidence for this type of intervention. It was thought that the economic analysis could clarify the trade-offs between the additional costs of the intervention alongside reported benefits.

### **3.2 Conclusion**

Both economic analyses were used to support 'consider' recommendations in the guideline. This was due to the nature of the evidence base, primarily that studies were non-UK.

Economic analyses of both interventions are found in Appendix C3.2 (Kessing et al. 2013) and Appendix C3.3 (Fuhr et al. 2014).

#### 4 Bibliography

Colom F, Reinares M, Pacchiarotti I, Popovic D, Mazzarini L., Martinez AA, Torren, C, Rosa AR, Palomino-Otiniano R, Franco C, Bonnín CM, Vieta E (2010) Has number of previous episodes any effect on responses to group psychoeducation in bipolar patients? *Acta Neuropsychiatr.* 22: 50–3

Fuhr et al. (2014) Effectiveness of peer-delivered interventions for severe mental illness and depression on clinical and psychosocial outcomes: a systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol.* 49: 1691–702

Kessing L, Hansen H, Hvenegaard A, Christensen E, Dam H, Glud C, Wetterslev J (2013) Treatment in a specialised out-patient mood disorder clinic V. standard out-patient treatment in the early course of bipolar disorder: randomised clinical trial. *The British Journal of Psychiatry* 202: 212–9

Lloyd-Evans B, Mayo-Wilson E, Harrison B, Istead H, Brown E, Pilling S, Johnson S, Kendall T (2014) A systematic review and meta-analysis of randomised controlled trials of peer support for people with severe mental illness. *BMC Psychiatry* 14: 39

Pitt V, Lowe D, Hill S, Pictor M, Hetrick SE, Ryan R, Berends L (2013). Consumer-providers of care for adult clients of statutory mental health services (review). *Cochrane Database of Systematic Reviews* 3

Reinares M, Sanchezz-Moreno J, Fountoulakis KN (2014) Psychosocial interventions in bipolar disorder: what, for whom, and when. *Journal of Affective Disorders* 156: 46–55

Scott J, Paykel E, Morriss R, Bentall R, Kinderman P, Johnson T, Abbott R, Hayhurst H (2006) Cognitive-behavioural therapy for severe and recurrent bipolar disorders: randomised controlled trial. *Br. J. Psychiatry* 188: 313–20