

Chapter 15 Advance care planning

Emergency and acute medical care in over 16s: service delivery and organisation

NICE guideline 94

March 2018

*Developed by the National Guideline Centre,
hosted by the Royal College of Physicians*

Disclaimer

Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and, where appropriate, their guardian or carer.

Copyright

© NICE 2018. All rights reserved. Subject to [Notice of rights](#).

ISBN: 978-1-4731-2741-8

Chapter 15 Advance care planning

Contents

15 Advance care planning	5
15.1 Introduction	5
15.2 Review question: Does advance care planning improve outcomes compared with usual care?	5
15.3 Clinical evidence.....	6
15.4 Economic evidence	10
15.5 Evidence statements.....	10
15.6 Recommendations and link to evidence.....	11
References	14
Appendices	21
Appendix A: Review protocol	21
Appendix B: Clinical article selection	23
Appendix C: Forest plots	24
Appendix D: Clinical evidence tables.....	26
Appendix E: Economic evidence tables	34
Appendix F: GRADE tables	35
Appendix G: Excluded clinical studies	37
Appendix H: Excluded economic studies.....	39

15 Advance care planning

15.1 Introduction

Advance care planning is established as part of best practice as an integral part of the management of people with multiple chronic conditions, particularly as they approach the limits of their treatment and end of life.

Advance care planning has been defined as a process of formal decision making that aims to help people establish decisions about future care that take effect when they lose the capacity⁴¹ to make informed decisions. In many cases that will lead to decisions about the extent of treatment, location of treatment and cardiopulmonary resuscitation. Some people may also instruct their families and friends and/or delegate power of attorney for such decisions.

This is incorporated in government strategy “End of life care strategy: promoting high quality care for adults at the end of their life” (2008) and supported by GM guidance “treatment and care towards the end of life: good practice in decision making” (2010). Patient and care advice is also available.⁷³

These people are likely to also present with acute medical emergencies and advance care plans have the potential to improve their care, including the treatment in the environment of their choice. Therefore, we asked the question “Does advance care planning improve outcomes compared with usual care?”

15.2 Review question: Does advance care planning improve outcomes compared with usual care?

For full details see review protocol in Appendix A.

Table 1: PICO characteristics of review question

Population	Adults and young people (16 years and over) with a suspected or confirmed or at risk of an AME
Intervention	Advance care planning <ul style="list-style-type: none"> • Patients who receive advance care planning and go on to make an advance care directive (ACP/AD+) • Patients who receive advance care planning and do not go on to make an advance care directive (ACP/AD-) • Patients who receive advance care planning but it is not reported whether they make an advance care directive (ACP)
Comparison	Usual care (no advance care planning) <ul style="list-style-type: none"> • Patients who do not receive advance care planning but make an advance care directive (No ACP/AD+) • Patients who do not receive advance care planning and do not make an advance care directive (No ACP/AD-) • Patients who do not receive advance care planning and it is not reported whether they make an advance care directive (No ACP)
Outcomes	Avoidable adverse events (CRITICAL) Quality of life (CRITICAL) Patient and/or carer satisfaction (CRITICAL) Length of hospital stay/Hospital-free days/Super spell days (IMPORTANT) Number of presentations to Emergency Department (CRITICAL)

	Number of admissions to hospital (CRITICAL) Number of GP presentations (IMPORTANT) Readmission up to 30 days (IMPORTANT) Place of death (CRITICAL)
Study design	Systematic reviews (SRs) of RCTs, RCTs, observational studies only to be included if no relevant SRs or RCTs are identified.

15.3 Clinical evidence

Three studies were included in the review.^{30,34,35} These are summarised in Table 2 below. Evidence from these studies is summarised in the GRADE clinical evidence summary below (Table 3). See also the study selection flow chart in Appendix B, study evidence tables in Appendix D, forest plots in Appendix C, GRADE tables in Appendix F and excluded studies list in Appendix G.

We searched for randomised trials comparing the impact of advance care planning or usual care (no advance care planning) on the outcomes outlined in Table 1.

Three randomised controlled trials were identified in which patients were randomised to receive facilitated advance care planning or usual care.

Table 2: Summary of studies included in the review

Study	Intervention and comparison	Population	Outcomes	comments
Detering, 2010 ³⁰ RCT	Facilitated advance care planning versus usual care	Legally competent medical inpatients aged 80 or above. n=309 Setting: A University hospital in Melbourne, Australia	Patient and/or carer satisfaction; Family satisfaction	Single centre study
Engelhardt 2006 ³⁴ RCT	Advanced illness co-ordinated care program including end of life care versus usual care (69.4% created advance directives in intervention group versus 48.4% in the usual care group)	Patients from Veterans Affairs Medical Centres, a home care organisation and 2 managed care organisations. n=275 Setting: Three Department of Veterans Affairs Medical centres, a home care organisation and 2 managed care organisations in New York, USA	Patient and/or carer satisfaction, family satisfaction	Limited generalisability of the findings to populations with different demographic characteristics (for example, female, non-white), to those with less serious medical problems (for example, outpatients), and to those with other diagnoses is limited.
Engelhardt 2009 ³⁵ RCT	Advanced illness co-ordinated care programme including end of life care versus usual care (47.0% created advance directives in intervention group versus 21.1% in the usual care group)	Patients with advanced illnesses. n=532 Setting: New York, USA	Inpatient admissions, ED visits, quality of life	Limited generalisability-study included a homogeneous population (87.9% white)

Table 3: Clinical evidence profile: advance care planning versus usual care, RCT evidence

Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with Control	Risk difference with Advance care planning (95% CI)
Patient and/or carer satisfaction	272 (1 study)	⊕⊕⊕⊖ MODERATE ^a due to risk of bias	RR 1.05 (1 to 1.09)	Moderate 942 per 1000	47 more per 1000 (from 0 more to 85 more)
Family satisfaction	56 (1 study)	⊕⊕⊖⊖ LOW ^{a,b} due to risk of bias, imprecision	RR 1.15 (0.91 to 1.46)	Moderate 778 per 1000	117 more per 1000 (from 70 fewer to 358 more)
Patient and/or carer satisfaction	186 (1 study)	⊕⊕⊖⊖ LOW ^a due to risk of bias	-	-	The mean patient and/or carer satisfaction in the intervention groups was 0.09 higher (0.1 lower to 0.28 higher)
Family satisfaction (problems reported)	143 (1 study)	⊕⊕⊖⊖ LOW ^a due to risk of bias	-	-	The mean family satisfaction (problems reported) in the intervention groups was 0.12 lower (0.22 to 0.02 lower)
QOL: SF-12 Physical standardised score	403 (1 study)	⊕⊕⊖⊖ LOW ^{a,b} due to risk of bias, imprecision	-	-	The mean QOL: SF-12 physical standardised score in the intervention groups was 1.66 lower (3.71 lower to 0.39 higher)
QOL: SF-12 Mental standardised score	403 (1 study)	⊕⊖⊖⊖ VERY LOW ^{a,b} due to risk of bias, imprecision	-	-	The mean QOL: SF-12 mental standardised score in the intervention groups was 1.1 higher (1.08 lower to 3.28 higher)
McGill QOL questionnaire	403 (1 study)	⊕⊕⊖⊖ LOW ^a due to risk of bias	-	-	The mean McGill QOL questionnaire in the intervention groups was 0.14 higher (0.06 lower to 0.34 higher)

Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with Control	Risk difference with Advance care planning (95% CI)
ED visits	403 (1 study)	⊕⊕⊖⊖ LOW ^a due to risk of bias, imprecision	-	-	The mean number of ED visits in the intervention groups was 1.66 lower (3.62 lower to 0.3 higher)
Inpatient admissions	403 (1 study)	⊕⊕⊖⊖ LOW ^a due to risk of bias, imprecision	-	-	The mean number of inpatient admissions in the intervention groups was 1.89 lower (4.23 lower to 0.45 higher)

(a) Downgraded by 1 increment if the majority of the evidence was at high risk of bias, and downgraded by 2 increments if the majority of the evidence was at very high risk of bias.

(b) Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

Outcomes which could not be analysed in Revman:

In 1 study³⁰ 133/154 patients (or family members if patients died before discharge) in the intervention group and 139/155 patients (or family members if patients died before discharge) in the usual care group completed a discharge questionnaire. Overall patient level of satisfaction with hospital stay was 93% very satisfied, 5% satisfied and 2% not satisfied in the intervention group and 65% very satisfied, 29% satisfied and 6% not satisfied in the usual care group. Family satisfaction with the quality of death was 83% very satisfied, 7% satisfied and 10% not satisfied in the intervention group and 48% very satisfied, 30% satisfied and 22% not satisfied in the usual care group.

15.4 Economic evidence

Published literature

No relevant economic evaluations were identified.

The economic article selection protocol and flow chart for the whole guideline can be found in the guideline's Appendix 41A and Appendix 41B.

Unit costs

Table 4: Unit costs of end of life admissions

	Mean cost
Hospital admission ending in death	£2,506 to £3,587 ^(a)

(a) Source: NHS Improving Quality 2013.⁹⁷ The lower value is based on NICE QIPP calculations while the higher value is based on the NHS Improving Quality study

The Centre for Reviews and Dissemination produced a briefing for Bristol CCG to support decisions regarding the delivery of advance care planning.²³ This briefing highlighted the lack of evidence on the cost-effectiveness of advance care planning. However, it cited an independent data analysis from South West England that was used in an economic evaluation of transferring people to an Electronic Palliative Care Co-ordination system (EPaCCS) that supports the co-ordination of care, including support for conversations about end of life care wishes.⁹⁷ This data analysis showed that deaths in hospital in the area relating to people transferred to EPaCCs were below 10% (compared to a national average of 54.5%). They estimated the support required for death in usual place of residence was £2107, and therefore there would be a saving of £399 to £1480 per death in usual place of residence compared with the lower and upper values of the cost of a hospital admission ending in death (see Table 4).

15.5 Evidence statements

Clinical

Three studies comprising 1116 patients evaluated advance care planning for improving outcomes, in adults and young people at risk of an AME, or with a suspected or confirmed AME. The evidence suggested that compared with usual care, advance care planning may provide a benefit for reduced number of ED visits and inpatient admissions (1 study, low quality), and family satisfaction (dichotomous data) (1 study, low quality). However, the evidence suggested there was no difference to patient and/or carer satisfaction (1 study, moderate quality), family satisfaction problems reported (continuous data) (1 study, low quality), quality of life SF-12 physical standardised score (1 study, low quality) and quality of life SF-12 mental standardised score (1 study, very low quality).

Economic

No relevant economic evaluations were identified.

15.6 Recommendations and link to evidence

Recommendations	9. Offer advance care planning to people in the community and in hospital who are approaching the end of life and are at risk of a medical emergency^a. Ensure that there is close collaboration between the person, their families and carers, and the professionals involved in their care.
Research recommendation	-
Relative values of different outcomes	The guideline committee considered quality of life, place of death, total number of admissions to hospital, avoidable adverse events, and presentations to ED and patient and/or carer satisfaction as critical outcomes for their decision-making. The outcomes considered to be important were: readmission, length of hospital stay and number of presentations to the GP.
Trade-off between benefits and harms	<p>There was evidence from 3 randomised controlled trials comparing advance care planning (ACP) with usual care.</p> <p>The evidence suggested that compared to usual care, advance care planning may provide a benefit for reduced number of ED visits and inpatient admissions and family satisfaction (dichotomous data). However, the evidence suggested there were no differences to patient and/or carer satisfaction, family satisfaction (continuous data), problems reported and quality of life (using several scales), but the estimates of effect generally favoured advance care planning.</p> <p>No evidence was found for the following outcomes: length of hospital stay, avoidable adverse events, number of GP presentations, place of death and readmission.</p> <p>The committee felt that given the benefit in terms of reduced ED visits and inpatient admission and the trend towards a positive benefit for other outcomes, a recommendation could be developed to support the use of advance care plans.</p> <p>The committee made a strong recommendation for ACP, even though the evidence was not of high quality. This was because providing ACP did not seem to have any obvious negative effects. In addition, it is in line with public surveys in which the majority express a wish to die at home. It was felt that ACP would help to preserve dignity; enable patient choice and does not involve a significant burden in terms of costs.</p> <p>While recognising that the prediction of life expectancy was not an exact science, the committee members were content to use the General Medical Council's definition³⁸ of 'approaching the end of life' as meaning that death was likely within 12 months. The NICE quality standard guidance on End of life care for adults⁷² refers to end of life care as being provided by the NHS in England for people who are likely to die within the next 12 months. Choosing the best time for such sensitive discussions needed to take into account individual's particular circumstances. There are useful tools that are available to help identify and care for patients in this phase of their life (for example, The Gold Standards Framework¹).</p>
Trade-off between net effects and costs	<p>No economic evidence was identified and therefore, unit costs were presented to the committee.</p> <p>The wider implementation of advanced care planning would require more input from community based palliative care services for patients and their families. However, the committee felt that for many patients it would be less resource intensive to die at home. ACP reduces hospital admission and emergency</p>

a NICE is developing a guideline on end of life care for adults in the last year of life.

<p>Recommendations</p>	<p>9. Offer advance care planning to people in the community and in hospital who are approaching the end of life and are at risk of a medical emergency^a. Ensure that there is close collaboration between the person, their families and carers, and the professionals involved in their care.</p>
<p>Research recommendation</p>	<p>-</p>
	<p>department visits which might be translated into cost savings. The committee also noted that none of the studies assessed hospital length of stay as an outcome. Additionally, evidence considered in the review of community palliative care (Chapter 14) suggests that this service was cost saving. Based on their collective experience, the committee believed that caring for terminally ill people at home can release hospital beds for other patients and this would allow the hospital to use its available resources more efficiently.</p>
<p>Quality of evidence</p>	<p>The evidence for the outcomes: family satisfaction, ED attendance and inpatient admissions were of low GRADE quality due to risk of bias and imprecision. The evidence for patient and/or carer satisfaction was of moderate to low GRADE quality, due to risk of bias. The evidence for quality of life (physical and mental components) was low to very low GRADE quality respectively, due to imprecision and risk of bias. Evidence reporting quality of life through the McGill pain questionnaire was of low GRADE quality due to risk of bias.</p>
<p>Other considerations</p>	<p>The committee noted that in order to implement ACP (also known as advance statement of wishes) it is crucial that information can be accessed by all relevant parties, such as the GP, hospital staff, paramedics and nursing home staff. Without this information staff may attempt to resuscitate patients who had DNAR orders as part of their advance decision or paramedics may transport patients to hospital against the person’s wishes, particularly if they are unable to express this. The committee noted that in London, senior paramedics have access to the “Coordinate my care” database which alerts them if a person has an advance care directive. This helps the paramedics to make an informed decision about how to manage the patient. It is also important the next of kin and relevant family members are aware of advanced directives to ensure that they do not countermand the wishes of an informed and capacitous person.</p> <p>The committee also considered by whom, how and when advance care planning conversations should be held. It was felt that it was probably best done by the healthcare professional with the closest relationship with the patient, while involving others with specific expertise (for example, in terms of the benefits and burdens of particular medical interventions and technologies, such as intensive care medicine). The GP, care home staff or hospital doctors all had a role in initiating the discussion, which should include the family and carers, and spiritual leaders where appropriate.</p> <p>There was an agreement that an acute medical emergency was not an ideal context for these discussions, but that if the patient’s wishes had not been determined beforehand then it was still better to attempt to do so in the acute situation rather than embarking on a potentially burdensome treatment pathway. The opportunity to broach these sensitive discussions shortly after resolution of an AME should also not be neglected.</p> <p>Advance care directives should be reviewed at regular intervals to permit changes in patients’ wishes to be respected. Mental capacity when discussing ACP must be considered and the appropriate measures should be put in place if needed for example, enduring power of attorney.</p>

<p>Recommendations</p>	<p>9. Offer advance care planning to people in the community and in hospital who are approaching the end of life and are at risk of a medical emergency^a. Ensure that there is close collaboration between the person, their families and carers, and the professionals involved in their care.</p>
<p>Research recommendation</p>	<p>-</p>
	<p>The current use of ACP within the NHS is variable. Greater emphasis on ACP is required, especially with an increasing elderly population with multiple co-morbidities which limit the efficacy of life-sustaining therapies.</p> <p>The committee noted that for ACP to be implemented it is crucial that staff are appropriately trained, with regular updates and readily accessible supporting information. Training should include when and how to implement discussions, and should include background information on the natural history and management of chronic diseases in patients in the community. Identifying that people are progressing towards the end of their lives requires knowledge of the person as well as an understanding of disease and the exclusion of reversible but undiagnosed acute, chronic, or acute-on-chronic illness. Public education is a key component but currently not well-addressed. Initiatives to promote advance care planning should focus on public understanding of the limits of medical technology, and optimising informed decision-making between communities, primary and secondary care.</p> <p>Improvements are required in IT infrastructure to bridge the communication gaps between primary, secondary and social care. An example is provided by the NHS programme 'Coordinate my care' (http://coordinatemycare.co.uk/) which permits information-sharing about treatment preferences and limits, and links to NHS 111 and some ambulance services.</p> <p>NICE has developed guidance on Care of dying adults in the last days of life.⁷¹</p> <p>NICE is developing a guideline on End of life Care for adults in last year of life' due to publish 2018 (see NICE website for further details).</p>

References

- 1 The Gold Standards Framework. 2017. Available from: <http://www.goldstandardsframework.org.uk/> [Last accessed: 8 March 2017]
- 2 Alberta Heritage Foundation for Medical Research. Advance directives and health care costs and the end of life. Canada. Edmonton: Alberta Heritage Foundation for Medical Research (AHFMR), 2005. Available from: <http://www.assembly.ab.ca/lao/library/egovdocs/2005/alhfm/150236.pdf>
- 3 Alifrangis C, Koizia L, Rozario A, Rodney S, Harrington M, Somerville C et al. The experiences of cancer patients. *QJM*. 2011; 104(12):1075-1081
- 4 Allen R, Ventura N. Advance directives use in acute care hospitals. *JONA's Healthcare Law, Ethics, and Regulation*. 2005; 7(3):86-91
- 5 Amering M, Stastny P, Hopper K. Psychiatric advance directives: qualitative study of informed deliberations by mental health service users. *British Journal of Psychiatry*. 2005; 186:247-252
- 6 Amjad H, Towle V, Fried T. Association of experience with illness and end-of-life care with advance care planning in older adults. *Journal of the American Geriatrics Society*. 2014; 62(7):1304-1309
- 7 Anderson JP, Kaplan RM, Schneiderman LJ. Effects of offering advance directives on quality adjusted life expectancy and psychological well-being among ill adults. *Journal of Clinical Epidemiology*. 1994; 47(7):761-772
- 8 Arenson CA, Novielli KD, Chambers CV, Perkel RL. The importance of advance directives in primary care. *Primary Care - Clinics in Office Practice*. 1996; 23(1):67-82
- 9 Atkinson JM, Garner HC, Patrick H, Stuart S. Issues in the development of advance directives in mental health care. *Journal of Mental Health*. 2003; 12(5):463-474
- 10 Attwood S, Anderson K, Mitchell T. Discussing cardiopulmonary resuscitation with patients. *British Journal of Nursing*. 2001; 10:1201-1207
- 11 Au DH, Udris EM, Engelberg RA, Diehr PH, Bryson CL, Reinke LF et al. A randomized trial to improve communication about end-of-life care among patients with COPD. *Chest*. 2012; 141(3):726-735
- 12 Braun KL, Karel H, Zir A. Family response to end-of-life education: differences by ethnicity and stage of caregiving. *American Journal of Hospice and Palliative Medicine*. 2006; 23(4):269-276
- 13 Bravo G, Arcand M, Blanchette D, Boire-Lavigne AM, Dubois MF, Guay M et al. Promoting advance planning for health care and research among older adults: a randomized controlled trial. *BMC Medical Ethics*. 2012; 13:1
- 14 Bravo G, Dubois MF, Wagneur B. Assessing the effectiveness of interventions to promote advance directives among older adults: a systematic review and multi-level analysis. *Social Science and Medicine*. 2008; 67(7):1122-1132
- 15 Briggs LA, Kirchhoff KT, Hammes BJ, Song MK, Colvin ER. Patient-centered advance care planning in special patient populations: a pilot study. *Journal of Professional Nursing*. 2004; 20(1):47-58

- 16 Brink P, Smith TF, Kitson M. Determinants of do-not-resuscitate orders in palliative home care. *Journal of Palliative Medicine*. 2008; 11(2):226-232
- 17 Broadwell AW, Boisauvin EV, Dunn JK, Engelhardt HT, Jr. Advance directives on hospital admission: a survey of patient attitudes. *Southern Medical Journal*. 1993; 86(2):165-168
- 18 Brody KK, Perrin NA, Dellapenna R. Advanced illness index: predictive modeling to stratify elders using self-report data. *Journal of Palliative Medicine*. 2006; 9(6):1310-1319
- 19 Campbell LA, Kisely SR. Advance treatment directives for people with severe mental illness. *Cochrane Database of Systematic Reviews*. 2009; Issue 1:CD005963. DOI:10.1002/14651858.CD005963.pub2
- 20 Capel M, Gazi T, Vout L, Wilson N, Finlay I. Where do patients known to a community palliative care service die? *BMJ Supportive and Palliative Care*. 2012; 2:43-47
- 21 Caplan GA, Meller A, Squires B, Chan S, Willett W. Advance care planning and hospital in the nursing home. *Age and Ageing*. 2006; 35(6):581-585
- 22 Catic AG, Berg AI, Moran JA, Knopp JR, Givens JL, Kiely DK et al. Preliminary data from an advanced dementia consult service: Integrating research, education, and clinical expertise. *Journal of the American Geriatrics Society*. 2013; 61(11):2008-2012
- 23 Centre for Reviews and Dissemination. Advance care planning. England. York: Centre for Reviews and Dissemination (CRD), 2013. Available from: <https://www.york.ac.uk/inst/crd/pdf/Advance%20care%20planning.pdf>
- 24 Chen CY, Thorsteinsdottir B, Cha SS, Hanson GJ, Peterson SM, Rahman PA et al. Health care outcomes and advance care planning in older adults who receive home-based palliative care: a pilot cohort study. *Journal of Palliative Medicine*. 2014; 18(1):38-44
- 25 Clayton JM, Butow PN, Tattersall MH, Devine RJ, Simpson JM, Aggarwal G et al. Randomized controlled trial of a prompt list to help advanced cancer patients and their caregivers to ask questions about prognosis and end-of-life care. *Journal of Clinical Oncology*. 2007; 25(6):715-723
- 26 Connors J, Dawson NV, Desbiens NA, Fulkerson J, Goldman L, Knaus WA et al. A controlled trial to improve care for seriously ill hospitalized patients: the study to understand prognoses and preferences for outcomes and risks of treatments (SUPPORT). *JAMA - Journal of the American Medical Association*. 1995; 274(20):1591-1598
- 27 De Korte-Verhoef MC, Pasman HRW, Schweitzer BPM, Francke AL, Onwuteaka-Philipsen BD, Deliens L. General practitioners' perspectives on the avoidability of hospitalizations at the end of life: a mixed-method study. *Palliative Medicine*. 2014; 28(7):949-958
- 28 De Vleminck A, Houttekier D, Deliens L, Vander Stichele R, Pardon K. Development of a complex intervention to support the initiation of advance care planning by general practitioners in patients at risk of deteriorating or dying: a phase 0-1 study. *BMC Palliative Care*. 2016; 15(1):17
- 29 Dening KH, Jones L, Sampson EL. Advance care planning for people with dementia: a review. *International Psychogeriatrics*. 2011; 23(10):1535-1551
- 30 Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ*. 2010; 340:c1345

- 31 Dev R, Coulson L, Del Fabbro E, Palla SL, Yennurajalingam S, Rhondali W et al. A prospective study of family conferences: effects of patient presence on emotional expression and end-of-life discussions. *Journal of Pain and Symptom Management*. 2013; 46(4):536-545
- 32 Donze J, Lipsitz S, Schnipper JL. Risk factors for potentially avoidable readmissions due to end-of-life care issues. *Journal of Hospital Medicine*. 2014; 9(5):310-314
- 33 Effiong A, Shinn L, Pope TM, Raho JA. Advance care planning for end-stage kidney disease. *Cochrane Database of Systematic Reviews*. 2013; Issue 7:CD010687. DOI:10.1002/14651858.CD010687
- 34 Engelhardt JB, McClive-Reed KP, Toseland RW, Smith TL, Larson DG, Tobin DR. Effects of a program for coordinated care of advanced illness on patients, surrogates, and healthcare costs: a randomized trial. *American Journal of Managed Care*. 2006; 12(2):93-100
- 35 Engelhardt JB, Rizzo VM, Della Penna RD, Feigenbaum PA, Kirkland KA, Nicholson JS et al. Effectiveness of care coordination and health counseling in advancing illness. *American Journal of Managed Care*. 2009; 15(11):817-825
- 36 Evans N, Bausewein C, Menaca A, Andrew EVW, Higginson IJ, Harding R et al. A critical review of advance directives in Germany: attitudes, use and healthcare professionals' compliance. *Patient Education and Counseling*. 2012; 87(3):277-288
- 37 Gade G, Venohr I, Conner D, McGrady K, Beane J, Richardson RH et al. Impact of an inpatient palliative care team: a randomized control trial. *Journal of Palliative Medicine*. United States 2008; 11(2):180-190
- 38 General Medical Council. Treatment and care towards the end of life: good practice in decision making, 2010. Available from: http://www.gmc-uk.org/guidance/ethical_guidance/end_of_life_care.asp
- 39 Glaudemans JJ, Moll van Charante EP, Willems DL. Advance care planning in primary care, only for severely ill patients? A structured review. *Family Practice*. 2015; 32(1):16-26
- 40 Hajizadeh N, Crothers K, Braithwaite RS. Using modeling to inform patient-centered care choices at the end of life. *Journal of Comparative Effectiveness Research*. 2013; 2(5):497-508
- 41 Hayhoe B, Howe A. Advance care planning under the Mental Capacity Act 2005 in primary care. *British Journal of General Practice*. 2011; 61(589):e537-e541
- 42 Henderson ML. Advance directives for patients with cancer. *Cancer Practice*. 1997; 5:186-188
- 43 Hesse KA. Terminal care of the very old. Changes in the way we die. *Archives of Internal Medicine*. 1995; 155(14):1513-1518
- 44 Hirschman KB, Abbott KM, Hanlon AL, Prvu BJ, Naylor MD. What factors are associated with having an advance directive among older adults who are new to long term care services? *Journal of the American Medical Directors Association*. 2012; 13(1):82
- 45 Ho VW, Thiel EC, Rubin HR, Singer PA. The effect of advance care planning on completion of advance directives and patient satisfaction in people with HIV/AIDS. *AIDS Care*. 2000; 12(1):97-108

- 46 Hofmann PB. Decisions near the end of life: resource allocation implications for hospitals. *Cambridge Quarterly of Healthcare Ethics*. 1992; 1(3):229-237
- 47 Houben CH, Spruit MA, Groenen MT, Wouters EF, Janssen DJ. Efficacy of advance care planning: a systematic review and meta-analysis. *Journal of the American Medical Directors Association*. 2014;(4):477-489
- 48 Houben CHM, Spruit MA, Wouters EFM, Janssen DJA. A randomised controlled trial on the efficacy of advance care planning on the quality of end-of-life care and communication in patients with COPD: The research protocol. *BMJ Open*. 2014; 4(1):e004465
- 49 Houttekier D, Cohen J, Cools F, Deliens L. Advance care planning for end-of-life care. *Cochrane Database of Systematic Reviews*. 2012; Issue 2:CD009618. DOI:10.1002/14651858.CD009618
- 50 In der Schmitt J, Rotharmel S, Mellert C, Rixen S, Hammes BJ, Briggs L et al. A complex regional intervention to implement advance care planning in one town's nursing homes: protocol of a controlled inter-regional study. *BMC Health Services Research*. 2011; 11:14
- 51 Ishihara KK, Wrenn K, Wright SW, Socha CM, Cross M. Advance directives in the emergency department: too few, too late. *Academic Emergency Medicine*. 1996; 3(1):50-53
- 52 Jain A, Corriveau S, Quinn K, Gardhouse A, Vegas DB, You JJ. Video decision aids to assist with advance care planning: a systematic review and meta-analysis. *BMJ Open*. 2015; 5(6):e007491
- 53 Jethwa KD, Onalaja O. Advance care planning and palliative medicine in advanced dementia: a literature review. *BJPsych Bulletin*. 2015; 39(2):74-78
- 54 Johnson J, Baranowski-Birkmeier T, O'Donnell JB. Advance directives in the medical intensive care unit of a community teaching hospital. *Chest*. 1995; 107(3):752-756
- 55 Jones L, Harrington J, Barlow CA, Tookman A, Drake R, Barnes K et al. Advance care planning in advanced cancer: can it be achieved? An exploratory randomized patient preference trial of a care planning discussion. *Palliative and Supportive Care*. 2011; 9(1):3-13
- 56 Kass-Bartelmes BL, Hughes R. Advance care planning: preferences for care at the end of life. *Journal of Pain and Palliative Care Pharmacotherapy*. 2004; 18(1):87-109
- 57 Khandelwal N, Kross EK, Engelberg RA, Coe NB, Long AC, Curtis JR. Estimating the effect of palliative care interventions and advance care planning on ICU utilization: a systematic review. *Critical Care Medicine*. 2015; 43(5):1102-1111
- 58 Khazaal Y, Chatton A, Pasandin N, Zullino D, Preisig M. Advance directives based on cognitive therapy: a way to overcome coercion related problems. *Patient Education and Counseling*. 2009; 74(1):35-38
- 59 La Puma J, Orentlicher D, Moss RJ. Advance directives on admission. Clinical implications and analysis of the Patient Self-Determination Act of 1990. *JAMA - Journal of the American Medical Association*. 1991; 266(3):402-405
- 60 Levy C, Morris M, Kramer A. Improving end-of-life outcomes in nursing homes by targeting residents at high-risk of mortality for palliative care: program description and evaluation. *Journal of Palliative Medicine*. 2008; 11(2):217-225

- 61 Lukas L, Foltz C, Paxton H. Hospital outcomes for a home-based palliative medicine consulting service. *Journal of Palliative Medicine*. 2013; 16(2):179-184
- 62 Mei CL, Hinderer KA, Kehl KA. A systematic review of advance directives and advance care planning in Chinese people from Eastern and Western cultures. *Journal of Hospice and Palliative Nursing*. 2014; 16(2):75-85
- 63 Mentz RJ, Tulskey JA, Granger BB, Anstrom KJ, Adams PA, Dodson GC et al. The palliative care in heart failure trial: rationale and design. *American Heart Journal*. 2014; 168(5):645
- 64 Mezey M, Kluger M, Maislin G, Mittelman M. Life-sustaining treatment decisions by spouses of patients with Alzheimer's disease. *Journal of the American Geriatrics Society*. 1996; 44(2):144-150
- 65 Mitchell SL, Morris JN, Park PS, Fries BE. Terminal care for persons with advanced dementia in the nursing home and home care settings. *Journal of Palliative Medicine*. 2004; 7(6):808-816
- 66 Molloy DW, Guyatt GH. A comprehensive health care directive in a home for the aged. *CMAJ Canadian Medical Association Journal*. 1991; 145:307-311
- 67 Molloy DW, Guyatt GH, Russo R, Goeree R, O'Brien BJ, Bédard M et al. Systematic implementation of an advance directive program in nursing homes: a randomized controlled trial. *JAMA - Journal of the American Medical Association*. 2000; 283(11):1437-1444
- 68 Morrison RS, Chichin E, Carter J, Burack O, Lantz M, Meier DE. The effect of a social work intervention to enhance advance care planning documentation in the nursing home. *Journal of the American Geriatrics Society*. 2005; 53(2):290-294
- 69 Motley M. Improving patient-centered care through advance care planning. *JAAPA : Official Journal of the American Academy of Physician Assistants*. 2013; 26(6):38-43
- 70 Mularski RA, Dy SM, Shugarman LR, Wilkinson AM, Lynn J, Shekelle PG et al. A systematic review of measures of end-of-life care and its outcomes. *Health Services Research*. 2007; 42(5):1848-1870
- 71 National Clinical Guideline Centre. Care of dying adults in the last days of life. NICE guideline 31. London. National Clinical Guideline Centre, 2015. Available from: <https://www.nice.org.uk/guidance/ng31>
- 72 National Institute for Health and Clinical Excellence. End of life care for adults. NICE quality standard 13. London. National Institute for Health and Clinical Excellence, 2011. Available from: <http://guidance.nice.org.uk/QS13>
- 73 NHS England and National Council for Palliative Care. Planning for your future care: a guide, 2009. Available from: <http://www.nhs.uk/planners/end-of-life-care/documents/planning-for-your-future-care.pdf>
- 74 Nicholas LH, Bynum JPW, Iwashyna T, Weir DR, Langa KM. Advance directives and nursing home stays associated with less aggressive end-of-life care for patients with severe dementia. *Health Affairs*. 2014; 33(4):667-674
- 75 Nicholas LH, Langa KM, Iwashyna TJ, Weir DR. Regional variation in the association between advance directives and end-of-life Medicare expenditures. *JAMA - Journal of the American Medical Association*. 2011; 306(13):1447-1453

- 76 Norris K, Merriman MP, Curtis JR, Asp C, Tuholske L, Byock IR. Next of kin perspectives on the experience of end-of-life care in a community setting. *Journal of Palliative Medicine*. 2007; 10(5):1101-1115
- 77 Oulton J, Rhodes SM, Howe C, Fain MJ, Mohler MJ. Advance directives for older adults in the emergency department: a systematic review. *Journal of Palliative Medicine*. 2015; 18(6):500-505
- 78 Papageorgiou A, Janmohamed A, King M, Davidson O, Dawson J. Advance directives for patients compulsorily admitted to hospital with serious mental disorders: directive content and feedback from patients and professionals. *Journal of Mental Health*. 2004; 13(4):379
- 79 Papageorgiou A, King M, Janmohamed A, Davidson O, Dawson J. Advance directives for patients compulsorily admitted to hospital with serious mental illness. Randomised controlled trial. *British Journal of Psychiatry*. 2002; 181:513-519
- 80 Piamjariyakul U, Myers S, Werkowitch M, Smith CE. End-of-life preferences and presence of advance directives among ethnic populations with severe chronic cardiovascular illnesses. *European Journal of Cardiovascular Nursing*. 2014; 13:185-189
- 81 Rabow MW, Dibble SL, Pantilat SZ, McPhee SJ. The comprehensive care team: a controlled trial of outpatient palliative medicine consultation. *Archives of Internal Medicine*. 2004; 164(1):83-91
- 82 Raymond M, Warner A, Davies N, Nicholas N, Manthorpe J, Iliffe S. Palliative and end of life care for people with dementia: lessons for clinical commissioners. *Primary Health Care Research and Development*. 2014; 15(4):406-417
- 83 Schellinger S, Sidebottom A, Briggs L. Disease specific advance care planning for heart failure patients: implementation in a large health system. *Journal of Palliative Medicine*. 2011; 14(11):1224-1230
- 84 Schmidt RJ, Weaner BB, Long D. The power of advance care planning in promoting hospice and out-of-hospital death in a dialysis unit. *Journal of Palliative Medicine*. 2014; 18(1):62-66
- 85 Schneiderman LJ, Pearlman RA, Kaplan RM, Anderson JP, Rosenberg EM. Relationship of general advance directive instructions to specific life-sustaining treatment preferences in patients with serious illness. *Archives of Internal Medicine*. 1992; 152(10):2114-2122
- 86 Sherwen E. Advance care planning ensures patient choice on place of death. *Nursing Times*. 2010; 106:8
- 87 Solloway M, LaFrance S, Bakitas M, Gerken M. A chart review of seven hundred eighty-two deaths in hospitals, nursing homes, and hospice/home care. *Journal of Palliative Medicine*. 2005; 8(4):789-796
- 88 Song K, Amatya B, Voutier C, Khan F. Advance care planning in patients with primary malignant brain tumors: a systematic review. *Frontiers in Oncology*. 2016; 6:223
- 89 Song MK. Effects of end-of-life discussions on patients' affective outcomes. *Nursing Outlook*. 2004; 52(3):118-125
- 90 Song MK, Ward SE, Fine JP, Hanson LC, Lin FC, Hladik GA et al. Advance care planning and end-of-life decision making in dialysis: a randomized controlled trial targeting patients and their surrogates. *American Journal of Kidney Diseases*. 2015; 66(5):813-822

- 91 Sudore R, Le Gem M, McMahon R, Feuz M, Katen M, Barnes DE. The advance care planning PREPARE study among older veterans with serious and chronic illness: study protocol for a randomized controlled trial. *Trials*. 2015; 16:570
- 92 Teno J, Lynn J, Connors AF, Jr., Wenger N, Phillips RS, Alzola C et al. The illusion of end-of-life resource savings with advance directives. SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. *Journal of the American Geriatrics Society*. 1997; 45(4):513-518
- 93 Teno JM, Gruneir A, Schwartz Z, Nanda A, Wetle T. Association between advance directives and quality of end-of-life care: a national study. *Journal of the American Geriatrics Society*. 2007; 55(2):189-194
- 94 Teno JM, Lynn J, Phillips RS, Murphy D, Youngner SJ, Bellamy P et al. Do formal advance directives affect resuscitation decisions and the use of resources for seriously ill patients? SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. *Journal of Clinical Ethics*. 1994; 5(1):23-30
- 95 Thomas RE, Wilson D, Sheps S. A literature review of randomized controlled trials of the organization of care at the end of life. *Canadian Journal on Aging*. 2006; 25(3):271-293
- 96 Wenger NS, Oye RK, Desbiens NA, Phillips RS, Teno JM, Connors AF, Jr. et al. The stability of DNR orders on hospital readmission. The SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. *Journal of Clinical Ethics*. 1996; 7(1):48-54
- 97 Whole Systems Partnership. Economic evaluation of the Electronic Palliative Care Coordination System (EPaCCS) early implementer sites. NHS, 2013. Available from: <http://www.england.nhs.uk/wp-content/uploads/2013/05/economic-eval-epaccs.pdf>
- 98 Wu P, Lorenz KA, Chodosh J. Advance care planning among the oldest old. *Journal of Palliative Medicine*. 2008; 11(2):152-157
- 99 Yoo JW, Nakagawa S, Kim S. Relationships among advance directives, principal diagnoses, and discharge outcomes in critically ill older adults. *Palliative and Supportive Care*. 2013; 11(4):315-322

Appendices

Appendix A: Review protocol


Table 5: Review protocol: Advance care planning

Review question	Does advance care planning improve outcomes compared with usual care?
Guideline condition and its definition	Acute medical emergencies. Definition: A medical emergency can arise in anyone, for example, in people: without a previously diagnosed medical condition, with an acute exacerbation of underlying chronic illness, after surgery, after trauma
Objectives	To determine if the provision of advance care planning, with or without advance directives improves outcomes.
Review population	Adults and young people (16 years and over) with a suspected or confirmed AME or at risk of an AME
	Adults
	Line of therapy not an inclusion criterion
Interventions and comparators: generic/class; specific/drug (All interventions will be compared with each other, unless otherwise stated)	Advance care planning; Patients who receive advance care planning and go on to make an advance care directive (ACP/AD+) Advance care planning; Patients who receive advance care planning and do not go on to make an advance care directive (ACP/AD-) Advance care planning; Patients who receive advance care planning but it is not reported whether they make an advance care directive (ACP) No advance care planning; Patients who do not receive advance care planning but make an advance care directive (No ACP/AD+) No advance care planning; Patients who do not receive advance care planning and do not make an advance care directive (No ACP/AD-) No advance care planning; Patients who do not receive advance care planning and it is not reported whether they make an advance care directive (No ACP)
Outcomes	- Quality of life at end of follow-up (Continuous) CRITICAL - Place of death- hospital/home at end of follow-up (Dichotomous) CRITICAL - Number of admissions to hospital at end of follow-up (Dichotomous) CRITICAL - Readmission up to 30 days at end of follow-up (Dichotomous) IMPORTANT - Number of presentations to Emergency Department during study period at end of follow-up (Dichotomous) CRITICAL - Patient and/or carer satisfaction at end of follow-up (Dichotomous) CRITICAL - Length of hospital stay at end of follow-up (Continuous) IMPORTANT - Avoidable adverse events at end of follow-up (Dichotomous) CRITICAL - Number of GP presentations at end of follow-up (Dichotomous) IMPORTANT - Length of stay in programme at end of follow-up (Continuous) IMPORTANT
Study design	Systematic Review RCT Quasi-RCT Non-randomised comparative study Prospective cohort study Retrospective cohort study

	Case control study Historical controlled study Before and after study Controlled before and after study Interrupted Time series
Unit of randomisation	Patient
Crossover study	Permitted
Minimum duration of study	Not defined
Other exclusions	Major trauma
Subgroup analyses if there is heterogeneity	- Frail elderly (Frail elderly; Not frail elderly); Different from younger population
Search criteria	Databases: Medline, Embase, the Cochrane Library Date limits for search: none Language: English only

Appendix B: Clinical article selection

Figure 1: Flow chart of clinical article selection for the review of advance care planning



Appendix C: Forest plots

C.1 Advance care planning versus standard care

Figure 2: Patient and/or carer satisfaction with hospital stay




Figure 3: Family satisfaction with quality of death




Figure 4: Patient and/or carer satisfaction




Figure 5: Family satisfaction (number of problems reported)




Figure 6: Quality of life: SF-12 Physical standardised score




Figure 7: Quality of life: SF-12 Mental standardised score




Figure 8: Quality of life: McGill questionnaire




Figure 9: Number of ED visits





Figure 10: Number of inpatient admissions



Appendix D: Clinical evidence tables

Study	Detering 2010 ³⁰
Study type	RCT (Patient randomised; Parallel)
Number of studies (number of participants)	1 (n=309)
Countries and setting	Conducted in Australia; Setting: Internal medicine, cardiology or respiratory medicine at a large university hospital in Melbourne, Australia
Line of therapy	Not applicable
Duration of study	Intervention + follow up: 6 months
Method of assessment of guideline condition	Adequate method of assessment/diagnosis
Stratum	Overall
Subgroup analysis within study	Not applicable
Inclusion criteria	Elderly patients aged over 80 admitted under internal medicine, cardiology or respiratory medicine in a large university hospital in Melbourne, Australia. Able to complete advance care planning during current hospital admission.
Exclusion criteria	Incompetent, cannot speak English, expected to die or be discharged within 24 hours, had previous formal advance care planning, and had no family.
Age, gender and ethnicity	Age - Median (IQR): Intervention group: 85 (82-88), control group: 84 (81-87). Gender (M:F): Intervention group: 54% men, Control group: 41% men. Ethnicity: nr
Further population details	1. Frail elderly
Extra comments	86% of the intervention group expressed wishes on end of life care by completing an advance care plan or verbally.
Indirectness of population	No indirectness
Interventions	(n=154) Intervention 1: Advance care directives. Formal advance care planning from a trained facilitator (nurse or allied health worker) using the Respecting Patient Choices model. The programme involved a co-ordinated approach to advance care planning whereby trained non-medical facilitators, in collaboration with treating doctors, assist patients and their families to reflect on the patient's goals, values and beliefs and to discuss and document their future choices about healthcare. Patients are encouraged to appoint a surrogate and to document their wishes about end of life care, including the wish for life prolonging treatments and cardiopulmonary resuscitation recorded on an advance care plan. As needed, treating doctors participated in this discussion to ensure that the patients understood their illness, treatment options and likely prognosis. This

Study	Detering 2010 ³⁰
	<p>programme utilises relevant legislation by enabling appointment of legal surrogates and ensures a systematic approach to filing of completed documents in hospital medical records so that they are readily available. Patients were encouraged to include their families, particularly their nominated surrogates, in discussions. The aim was to complete advance care planning before hospital discharge. 81% of the intervention group received advance care planning and of these 84% completed an advance care directive. A 5 question survey was used to assess patient and/or carer satisfaction on their hospital stay. Additionally the quality of end of life care questionnaire, an 8 item locally developed tool to assess a family member's satisfaction with the quality of the patient's death, was used. Duration Median discussion length 60 minutes (range 10-200 minutes) over 1-3 meetings. Concurrent medication/care: None.</p> <p>(n=155) Intervention 2: Usual Care. Consistent with usual practice, control patients received usual medical care but no advance care planning unless it was specifically requested. Duration nr. Concurrent medication/care: N/A</p>
Funding	Academic or government funding
<p>RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ADVANCE CARE DIRECTIVES versus USUAL CARE</p> <p>Of the 108 patients who expressed wishes on end of life care, 82% expressed wishes about cardiopulmonary resuscitation and 75% about life prolonging treatment.</p> <p>Protocol outcome 1: Mortality at during study period - Actual outcome: Mortality at Six months; Group 1: 29/154, Group 2: 27/155 Risk of bias: All domain - Low, Selection - Low, Blinding - Low, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness</p> <p>Protocol outcome 2: Patient and/ or carer satisfaction at during study period - Actual outcome: Patient (or family if patient died before discharge) satisfied with hospital stay at Before discharge; Group 1: 131/133, Group 2: 131/139; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - High, Crossover - Low, Comments - Unvalidated questionnaire; Indirectness of outcome: No indirectness, Comments: No indirectness- Actual outcome: Family satisfied with quality of death at Before discharge; Group 1: 26/29, Group 2: 21/27; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - High, Crossover - Low, Comments - Unvalidated questionnaire; Indirectness of outcome: No indirectness, Comments: No indirectnessindirectness</p>	
Protocol outcomes not reported by the study	Quality of life at during study period; Avoidable adverse events at during study period; Number of presentations to Emergency Department at during study period; Number of admissions to hospital at After 28 days of first admission; Number of GP presentations at during study period; Readmission; Length of stay in programme at during study period; Length of hospital stay at during study period

Study	Engelhardt 2006 ³⁴
Study type	RCT (Patient randomised; Parallel)
Number of studies (number of participants)	1 (n=Intervention group: 133, usual care group: 142)
Countries and setting	Conducted in USA; Setting: Three Department of Veterans Affairs Medical centres, a home care organisation and 2 managed care organisations.
Line of therapy	Not applicable
Duration of study	Intervention + follow up: 3 months
Method of assessment of guideline condition	Adequate method of assessment/diagnosis: Advanced illness defined as: COPD, chronic heart failure or cancer.
Stratum	Overall
Subgroup analysis within study	Not applicable
Inclusion criteria	Patients with chronic obstructive pulmonary disease, chronic heart failure or cancer diagnoses including those of the oesophagus, trachea, colon, liver, pancreas, lung or uterus, cancers of the prostate or breast with metastasis, melanoma, leukaemia, lymphosarcoma, Hodgkin's disease or multiple myeloma. Patients with COPD or CHF were eligible if they had experienced 1 or more admissions to an intensive care unit or 2 or more acute-care admissions in the last 6 months.
Exclusion criteria	Nr
Recruitment/selection of patients	Nr
Age, gender and ethnicity	Age - Mean (SD): Intervention group: 70.72, usual care group: 70.8. Gender (M:F): Intervention group: 18.8% female, 81.2% male; usual care group: 23.9% female, 76.1% male. Ethnicity: Intervention group: 88% white, 11.3% black, 0.8% other; usual care group: 85.7% white, 11.4% black, 2.9% other
Further population details	1. Frail elderly
Indirectness of population	No indirectness
Interventions	(n=133) Intervention 1: Advance care directives. The AICCP delivers care coordination and support through 6 functions. The first is physician support, which consists of helping patients develop well-organised questions to make economical use of provider time and ensuring that physicians have complete information about patients. The second is health literacy, which is the capacity to understand basic health information. The AICCP addresses literacy concerns in each session (for example, by helping patients comprehend specialised medical terminology, which both increases their understanding and reduces their embarrassment). The third function is care coordination, which is locating and arranging linkages to medical services. The fourth is prevention, which refers to a focus on those aspects of EOL planning that often are avoided and emotionally charged. In this study, prevention referred to efforts to reduce or eliminate common psychosocial

Study	Engelhardt 2006 ³⁴
	<p>concerns related to advanced illness such as (1) coping with the loss of ability to perform valued activities; (2) identifying and addressing family conflict around difficult advanced illness and EOL decisions (for example, patient relocation, financial burdens of illness); (3) avoiding caregiver burnout (for example, by dividing care among family members);(4) anticipating emotional reactions (for example, anticipatory grief, fear of death); (5) enhancing self-management skills by preparing patients and families to cope with health system delivery shortfalls (for example, fragmentation of care delivery, gaps in care); and (6) promoting advance planning, because timely planning may avert decision making in crisis situations. Care coordinators help clarify patient preferences for care under different health scenarios, using worksheets designed for this purpose. If patients engage in advance planning, care coordinators assist them in formulating and documenting ADs and discussing them with providers. Family misunderstandings about care issues frequently can be resolved during meetings with care coordinators, reducing physician time spent mediating between family members. Care coordinators also provide emotional and social support. Emotional support consists of attending to affective components of illness, identifying specific motions, helping patients cope with suffering, and providing referrals for on-going counselling. Social support includes guidance and information, as well as tangible support. In the AICCP, structured guidance support helps patients and caregivers complete tasks needed to maintain health and function. The AICCP provides information support in the form of guiding patients through the immense amount of medical information available to sources that are (1)adjusted for health literacy,(2) endorsed by their physicians, and (3) relevant to their situations. It provides tangible support by locating and arranging social support services. These functions are performed by nurses, nurse practitioners or social workers. The AICCP was implemented in a 6-session format and delivered by existing personnel who were familiar with institutional policies and who had on-going relationships with providers. These personnel were chosen because a reported barrier to effective implementation of an EOL program was using staff without an institutional identity and credibility. Care coordinators' salaries were contributed by study sites. Sites replaced care coordinators if their resources allowed; if not, care coordinators' duties were reconfigured to focus on patients with advanced illness. Each care coordinator attended training and reviewed assigned readings, including the AICCP training manual 15, 17. Program delivery was standardised across sites through conference calls and followed a structured-visit format. Care coordinators were taught to individualise the program to meet specific needs; for example, patients could schedule extra meetings. The mean number of visits was 4.92(SD = 2.94). Treatment implementation checklists, covering AICCP-recommended interventions, were examined for a randomly selected subset of patients. Intervention elements were completed in a mean of 83% of patients. The most common reason for not completing an element was that it did not apply to the patient's circumstances. Duration 6 sessions. Concurrent medication/care: 2 AICCP participants crossed over to usual care.</p> <p>(n=142) Intervention 2: Usual Care nr. Duration 3 months including follow up. Concurrent medication/care: 18 usual care participants crossed over to AICCP.</p>

Study	Engelhardt 2006 ³⁴
Funding	Academic or government funding (Robert Wood Johnson Foundation, Fan Fox/Leslie R Samuels Foundation and the Nathan Cummings Foundation)
<p>RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ADVANCE CARE PLANNING versus USUAL CARE</p> <p>Protocol outcome 1: Patient and/ or carer satisfaction at during study period - Actual outcome: Patient and/or carer satisfaction with healthcare measured on a five-point Likert-type scale at Enrolment, 3 and 6 months post enrolment; Group 1: mean 4.07 (SD 0.68); n=86, Group 2: mean 3.98 (SD 0.67); n=100; Risk of bias: All domain - Very high, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - High, Crossover - Low, Subgroups - Low, Other 1 - High, Other 2 - Low, Other 3 - Low, Comments - 18 usual care participants crossed over to AICCP and 2 AICCP participants crossed over onto usual care however ITT analysis was performed.; Indirectness of outcome: No indirectness ; Group 1 Number missing: 47, Reason: nr; Group 2 Number missing: 42, Reason: nr</p> <p>- Actual outcome: Surrogate (family) satisfaction with healthcare using a modified EOL Family Interview. Number of problems in 7 domains (shared decision making, physical comfort and emotional support, advance care planning, co-ordination of care, personal care and respect, family self-efficacy, family emotional and spiritual support. at Three months post-enrolment; Group 1: mean 0.41 (SD 0.3); n=67, Group 2: mean 0.53 (SD 0.32); n=76; Risk of bias: All domain - Very high, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - High, Crossover - Low, Subgroups - Low, Other 1 - High, Other 2 - Low, Other 3 - Low, Comments - 18 usual care participants crossed over to AICCP and 2 AICCP participants crossed over onto usual care however ITT analysis was performed.; Indirectness of outcome: No indirectness ; Group 1 Number missing: 47, Reason: nr; Group 2 Number missing: 42, Reason: nr</p>	
Protocol outcomes not reported by the study	Quality of life at during study period; Avoidable adverse events at during study period; Number of presentations to Emergency Department at during study period; Number of admissions to hospital at After 28 days of first admission; Number of GP presentations at during study period; Readmission; Length of stay in programme at during study period; Number of radiology tests at during study period; Number of outpatient visits at during study period; Number of laboratory tests ordered at during study period; Number of home health visits at during study period; Number of pharmacy prescriptions at during study period; Length of hospital stay at during study period

Study	Engelhardt 2009 ³⁵
Study type	RCT (Patient randomised; Parallel)
Number of studies (number of participants)	1 (n=532)
Countries and setting	Conducted in USA
Line of therapy	Not applicable
Duration of study	Intervention + follow up: 6 session intervention plus a 4 month follow up

Study	Engelhardt 2009 ³⁵
Method of assessment of guideline condition	Adequate method of assessment/diagnosis: Advanced illness defined as: COPD, chronic heart failure or cancer.
Stratum	Overall
Subgroup analysis within study	Not applicable
Inclusion criteria	Patients with chronic obstructive pulmonary disease, chronic heart failure or cancer diagnoses including those of the oesophagus, trachea, colon, liver, pancreas, lung or uterus, cancers of the prostate or breast with metastasis, melanoma, leukaemia, lymphosarcoma, Hodgkin's disease or multiple myeloma. Patients with COPD or CHF were eligible if they had experienced 1 or more admissions to an intensive care unit or 2 or more acute-care admissions in the last 6 months.
Exclusion criteria	NR
Age, gender and ethnicity	Age - -: Gender (M:F): Define. Ethnicity: Intervention: 84.8% white, 15.2% non-white; usual care: 91% white, 9% non-white
Further population details	1. Frail elderly:
Indirectness of population	--
Interventions	<p>(n=267) Intervention 1: Advance care directives.</p> <p>The 6-session model has the following 3 components: (1) nondirective health counselling, (2) education and (3) care coordination. AICCP was delivered by social workers and a health educator with 16 hours of initial training and with 20 hours of follow-up. AICCP meetings were face to face, lasting a mean of 59.0 (SD = 22.1) minutes, including brief follow-up telephone contacts. The mean number of sessions was 4.9 (SD = 2.1) (range 0-10 [mode, 6]), with 81.9% of patients completing 3 to 7 sessions. On average, caregivers attended 50% of sessions based on patient preference, caregiver availability, and need. The topics covered across sessions were structured in a biopsychosocial 3-domain format, including the following: (1)health-related topics included but were not limited to understanding illness, treatment expectations, emerging symptoms, adherence to treatment recommendations, communication with health professionals, and advance planning; (2) coping with loss of role, functional capacity or health status; evaluating whether situations are amenable to change or, if not, whether reactions to unchangeable situations are modifiable; and monitoring for anxiety or depression, interpersonal conflict and existential concerns; and (3) caregiving concerns, maximising health system benefits, home environmental modifications, home care, and long-term care planning. This structure was delivered using a nondirective health counselling format, patient education and care coordination. It facilitated recognition and normalisation of the consequences of living with on-going health problems in domains of function beyond physical health per se. It promoted identification of psychosocial needs and supports and facilitated initiation of discussions about ways to adapt to and compensate for losses induced by reduced health status. An electronic web tool operationalised each session of AICCP by providing a checklist of health education topics and tasks to be completed in interviews. For example, at these meetings, coordinators introduced advance planning. If an</p>

Study	Engelhardt 2009 ³⁵
	<p>expected task was not addressed at a specified meeting, coordinators were given pop-up reminders to complete them at subsequent meetings. Health education also included, as needed, information about health-related benefits within their health system and their community. Duration 6 sessions. Concurrent medication/care: n/a.</p> <p>(n=265) Intervention 2: Usual Care. Usual care. Duration n/a. Concurrent medication/care: n/a</p>
Funding	Academic or government funding (The Garfield Foundation)
RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ADVANCE CARE PLANNING versus USUAL CARE	
<p>Protocol outcome 1: Quality of life at during study period - Actual outcome: SF-12 physical standardised score at Post -test; Group 1: mean 32.99 (SD 10.31); n=198, Group 2: mean 34.65 (SD 10.66); n=205; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - Low, Crossover - Low, Subgroups - Low, Other 1 - Low, Other 2 - Low, Other 3 - Low; Indirectness of outcome: No indirectness ; Baseline details: Age differed between the two groups but statistically controlled for; Group 1 Number missing: 69, Reason: nr; Group 2 Number missing: 60, Reason: nr</p> <p>Protocol outcome 2: Number of presentations to Emergency Department at during study period - Actual outcome: ED visits at Post- test; Group 1: mean 3.69 (SD 6.14); n=198, Group 2: mean 5.35 (SD 12.87); n=205; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - Low, Crossover - Low, Subgroups - Low, Other 1 - Low, Other 2 - Low, Other 3 - Low; Indirectness of outcome: No indirectness ; Baseline details: Age differed between the two groups but statistically controlled for; Group 1 Number missing: 69, Reason: nr; Group 2 Number missing: 60, Reason: nr</p> <p>- Actual outcome: SF-12 mental standardised score at Post-test; Group 1: mean 49.09 (SD 10.77); n=198, Group 2: mean 47.99 (SD 11.55); n=205; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - Low, Crossover - Low, Subgroups - Low, Other 1 - Low, Other 2 - Low, Other 3 - Low; Indirectness of outcome: No indirectness ; Baseline details: Age differed between the two groups but statistically controlled for; Group 1 Number missing: 69, Reason: nr; Group 2 Number missing: 60, Reason: nr</p> <p>- Actual outcome: McGill Quality of Life Questionnaire at Post- test; Group 1: mean 5.03 (SD 0.87); n=198, Group 2: mean 4.89 (SD 1.14); n=205; Risk of bias: All domain - Very high, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - High, Crossover - Low, Subgroups - Low, Other 1 - Low, Other 2 - Low, Other 3 - Low; Indirectness of outcome: No indirectness ; Baseline details: Age differed between the two groups but statistically controlled for; Group 1 Number missing: 69, Reason: nr; Group 2 Number missing: 60, Reason: nr</p> <p>Protocol outcome 3: Number of admissions to hospital at After 28 days of first admission - Actual outcome: Inpatient admissions at Post- test; Group 1: mean 2.44 (SD 5.11); n=198, Group 2: mean 4.33 (SD 16.26); n=205; Risk of bias: All domain - High, Selection - Low, Blinding - Low, Incomplete outcome data - High, Outcome reporting - Low, Measurement - Low, Crossover - Low, Subgroups - Low, Other 1 - Low, Other 2 - Low, Other 3 - Low; Indirectness of outcome: No indirectness ; Baseline details: Age differed between the two groups but statistically controlled for; Group 1 Number missing: 69, Reason: nr; Group 2 Number missing: 60, Reason: nr</p>	
Protocol outcomes not reported by the study	Avoidable adverse events at during study period; Patient and/ or carer satisfaction at during study period; Number of

Study	Engelhardt 2009 ³⁵
	GP presentations at during study period; Readmission; Length of stay in programme at during study period; Number of radiology tests at during study period; Number of outpatient visits at during study period; Number of laboratory tests ordered at during study period; Number of home health visits at during study period; Number of pharmacy prescriptions at during study period; Length of hospital stay at during study period

Appendix E: Economic evidence tables

No studies were included.

Appendix F: GRADE tables

Table 6: Clinical evidence profile: Advance care planning versus usual care, RCT evidence

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Advance care planning	Control	Relative (95% CI)	Absolute		
Patient satisfaction												
1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	no serious imprecision	none	131/133 (98.5%)	94.2%	RR 1.05 (1 to 1.09)	47 more per 1000 (from 0 more to 85 more)	⊕⊕⊕O MODERATE	CRITICAL
Family satisfaction												
1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	Serious ²	none	26/29 (89.7%)	77.8%	RR 1.15 (0.91 to 1.46)	117 more per 1000 (from 70 fewer to 358 more)	⊕⊕OO LOW	CRITICAL
Patient satisfaction (Better indicated by higher values)												
1	randomised trials	very serious ¹	no serious inconsistency	no serious indirectness	no serious imprecision	none	86	100	-	MD 0.09 higher (0.1 lower to 0.28 higher)	⊕⊕OO LOW	CRITICAL
Family satisfaction (problems reported) (Better indicated by lower values)												
1	randomised trials	very serious ¹	no serious inconsistency	no serious indirectness	no serious imprecision	none	67	76	-	MD 0.12 lower (0.22 to 0.02 lower)	⊕⊕OO LOW	CRITICAL
QOL: SF-12 Physical standardised score (Better indicated by higher values)												
1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	Serious ²	none	198	205	-	MD 1.66 lower (3.71 lower to 0.39 higher)	⊕⊕OO LOW	CRITICAL
QOL: SF-12 Mental standardised score (Better indicated by higher values)												

1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	very serious ²	none	198	205	-	MD 1.1 higher (1.08 lower to 3.28 higher)	⊕○○○ VERY LOW	CRITICAL
McGill QOL questionnaire (Better indicated by lower values)												
1	randomised trials	very serious ¹	no serious inconsistency	no serious indirectness	no serious imprecision	none	198	205	-	MD 0.14 higher (0.06 lower to 0.34 higher)	⊕⊕○○ LOW	CRITICAL
Number of ED visits (Better indicated by lower values)												
1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	Serious ²	none	198	205	-	MD 1.66 lower (3.62 lower to 0.3 higher)	⊕⊕○○ LOW	CRITICAL
Number of inpatient admissions (Better indicated by lower values)												
1	randomised trials	serious ¹	no serious inconsistency	no serious indirectness	Serious ²	None	198	205	-	MD 1.89 lower (4.23 lower to 0.45 higher)	⊕⊕○○ LOW	CRITICAL

¹ Downgraded by 1 increment if the majority of the evidence was at high risk of bias, and downgraded by 2 increments if the majority of the evidence was at very high risk of bias

² Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

Appendix G: Excluded clinical studies

Table 7: Studies excluded from the clinical review

Study	Exclusion reason
ALBERTA 2005 ²	Review paper checked for references.
ALIFRANGIS 2011 ³	Qualitative study. No outcomes of interest
ALLEN 2005 ⁴	Observational study (RCT evidence available).
AMERING 2005 ⁵	Incorrect population. Qualitative study.
AMJAD 2014 ⁶	No outcomes of interest
ANDERSON 1994 ⁷	No outcomes of interest
ARENSON 1996 ⁸	Narrative paper checked for references
ATKINSON 2003 ⁹	Qualitative study
ATTWOOD 2001 ¹⁰	Review does not match protocol
AU 2012 ¹¹	No outcomes of interest
BRAUN 2006 ¹²	No outcomes of interest
BRAVO 2008 ¹⁴	Systematic review does not match protocol.
BRAVO 2012 ¹³	Methodological paper (no results reported)
BRIGGS 2004 ¹⁵	No outcomes of interest
BRINK 2008 ¹⁶	No outcomes of interest.
BROADWELL 1993 ¹⁷	Qualitative study
BRODY 2006 ¹⁸	Incorrect topic
CAMPBELL 2009 ¹⁹	Incorrect population
CAPEL 2012 ²⁰	No outcomes of interest.
CAPLAN 2006 ²¹	Observational study (RCT evidence available).
CATIC 2013 ²²	Incorrect intervention
CHEN 2014 ²⁴	Observational study (RCT evidence available).
CLAYTON 2007 ²⁵	Incorrect intervention
CONNORS 1995 ²⁶	Multiple interventions without a clear focus on advance care planning, therefore ACP not tested in trial.
DEKORTE-VERHOEF 2014 ²⁷	No outcomes of interest
DENING 2011A ²⁹	Systematic review checked for references
DEV 2013 ³¹	No outcomes of interest
DEVLEMINCK2016 ²⁸	Incorrect intervention. The objective of the study was to develop an intervention to support the initiation of advance care planning in general practice.
DONZE 2014 ³²	Incorrect study design (case-control)
EFFIONG 2013 ³³	Cochrane protocol does not sufficiently match protocol
EVANS 2012A ³⁶	Review checked for references
GADE 2008 ³⁷	RCT of palliative care team provision; not advanced care planning.
GLAUDEMANS 2015 ³⁹	Structured review. Checked references

HAJIZADEH 2013 ⁴⁰	Theoretical modeling
HENDERSON 1997 ⁴²	Narrative paper
HESSE 1995 ⁴³	Observational study (RCT evidence available).
HIRSCHMAN 2012 ⁴⁴	No outcomes of interest
HO 2000 ⁴⁵	Observational study (RCT evidence available).
HOFMANN 1992 ⁴⁶	Narrative paper
HOUBEN 2014 ⁴⁸	Methodological paper (no results reported)
HOUBEN 2014A ⁴⁷	Systematic review checked for references
HOUTTEKIER 2012 ⁴⁹	Cochrane protocol (review not completed)
INDERSCHMITTEN 2011 ⁵⁰	Methodological paper (no results reported)
ISHIHARA 1996 ⁵¹	No outcomes of interest
JAIN 2015 ⁵²	Systematic review. Checked references
JETHWA 2015 ⁵³	Literature review. Checked references
JOHNSON 1995 ⁵⁴	Observational study (RCT evidence available).
JONES 2011 ⁵⁵	Not an RCT of advance care planning; a trial of whether patients wish to discuss ACP
KASSBARTELMES 2004 ⁵⁶	Review paper checked for references
KHANDELWAL 2015 ⁵⁷	Systematic review. Checked references
KHAZAAL 2009 ⁵⁸	Incorrect population
LA PUMA 1991 ⁵⁹	Narrative checked for references
LEVY 2008 ⁶⁰	Observational study (RCT evidence available).
LUKAS 2013 ⁶¹	Observational study (RCT evidence available).
MEI 2014 ⁶²	Systematic review does not match protocol.
MENTZ 2014 ⁶³	Methodological paper (no results reported)
MEZEY 1996A ⁶⁴	No outcomes of interest
MITCHELL 2004 ⁶⁵	Incorrect comparison (terminal care in nursing homes versus community)
MOLLOY 1991A ⁶⁶	Observational study (RCT evidence available).
MOLLOY 2000A ⁶⁷	Care planning not tested; education about the existence of directives
MORRISON 2005 ⁶⁸	Social workers were randomized to ACP education, not patients randomized to receive or not receive ACP
MOTLEY 2013 ⁶⁹	Review checked for references
MULARSKI 2007 ⁷⁰	Systematic review does not match protocol
NICHOLAS 2011 ⁷⁵	Observational study (RCT evidence available).
NICHOLAS 2014 ⁷⁴	Observational study (RCT evidence available).
NORRIS 2007 ⁷⁶	Qualitative study
OULTON 2015 ⁷⁷	Systematic review. Checked references. Studies included in the review were of methodology not considered in our protocol (surveys, interviews)
PAPAGEORGIOU 2002 ⁷⁹	Incorrect population
PAPAGEORGIOU 2004 ⁷⁸	Incorrect population
PIAMJARIYAKUL 2014 ⁸⁰	Qualitative study
RABOW 2004 ⁸¹	Intervention group also received psychosocial support and family caregiver training.

RAYMOND 2014 ⁸²	Review paper, incorrect population.
SAMPSON 2011 ²⁹	Incorrect intervention. The study aimed to assess the feasibility of implementing a 2 component intervention (palliative assessment and advance care planning) to improve end of life care for people with advance dementia.
SHELLINGER 2011 ⁸³	Observational study (RCT evidence available).
SCHMIDT 2014 ⁸⁴	No outcomes of interest
SCHNEIDERMAN 1992 ⁸⁵	No outcomes of interest
SHERWEN 2010 ⁸⁶	Narrative
SOLLOWAY 2005 ⁸⁷	Observational study (RCT evidence available).
SONG 2004 ⁸⁹	Systematic review checked for references
SONG 2015 ⁹⁰	No protocol outcomes reported
Song2016A ⁸⁸	Systematic review- not AME patients. Advance care planning in patients with primary malignant brain tumours
SUDORE 2015 ⁹¹	Study protocol for a RCT
TENO 1994 ⁹⁴	Observational study (RCT evidence available).
TENO 1997 ⁹²	Incorrect comparison (advance directives that were documented on medical charts versus those that were not).
TENO 2007 ⁹³	Observational study (RCT evidence available).
THOMAS 2006B ⁹⁵	Review paper checked for references
WENGERS 1996 ⁹⁶	Incorrect comparison
WU 2008 ⁹⁸	Observational study (RCT evidence available).
YOO 2013 ⁹⁹	Observational study (RCT evidence available).

Appendix H: Excluded economic studies

No studies were excluded.