

# Appendix I: References

## I.1 Dementia diagnosis

### I.1.1 Introduction

Mercy L, Hodges JR, Dawson K, Barker RA, Brayne C. Incidence of early-onset dementias in Cambridgeshire, United Kingdom. *Neurology*. 2008 Nov 4;71(19):1496-9.

James BD, Bennett DA, Boyle PA, Leurgans S, Schneider JA. Dementia from Alzheimer disease and mixed pathologies in the oldest old. *JAMA*. 2012 May 2;307(17):1798-800.

### I.1.2 Dementia diagnosis

#### I.1.2.1 Systematic Reviews: sources of primary studies

Bloudek Lisa M, Spackman D Eldon, Blankenburg Michael, and Sullivan Sean D (2011) Review and meta-analysis of biomarkers and diagnostic imaging in Alzheimer's disease. *Journal of Alzheimer's disease: JAD* 26(4), 627-45

Brigo Francesco, Turri Giulia, and Tinazzi Michele (2015) 123I-FP-CIT SPECT in the differential diagnosis between dementia with Lewy bodies and other dementias. *Journal of the neurological sciences* 359(1-2), 161-71

Caobelli Federico, Cobelli Milena, Pizzocaro Claudio, Pavia Marco, Magnaldi Silvia, and Guerra Ugo Paolo (2015) The role of neuroimaging in evaluating patients affected by Creutzfeldt-Jakob disease: a systematic review of the literature. *Journal of neuroimaging: official journal of the American Society of Neuroimaging* 25(1), 2-13

Carnero-Pardo Cristobal, Lopez-Alcalde Samuel, Allegri Ricardo Francisco, and Russo Maria Julieta (2014) A systematic review and meta-analysis of the diagnostic accuracy of the Phototest for cognitive impairment and dementia. *Dementia & Neuropsychologia* 8(2), 141-147

Cromarty Ruth A, Elder Greg J, Graziadio Sara, Baker Mark, Bonanni Laura, Onofri Marco, O'Brien John T, and Taylor John-Paul (2016) Neurophysiological biomarkers for Lewy body dementias. *Clinical neurophysiology: official journal of the International Federation of Clinical Neurophysiology* 127(1), 349-59

Davis Daniel H. J, Creavin Sam T, Yip Jennifer L. Y, Noel-Storr Anna H, Brayne Carol, and Cullum Sarah (2015) Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. *The Cochrane database of systematic reviews* 10, CD010775

Dougall Nadine J, Bruggink Sjoerd, and Ebmeier Klaus P (2004) Systematic review of the diagnostic accuracy of 99mTc-HMPAO-SPECT in dementia. *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry* 12(6), 554-70

Durand-Martel Pascali, Tremblay Dominic, Brodeur Catherine, and Paquet Nancy (2010) Autopsy as gold standard in FDG-PET studies in dementia. *The Canadian journal of neurological sciences. Le journal canadien des sciences neurologiques* 37(3), 336-42

Ehreke L, Luppia M, Konig H H, and Riedel-Heller S G (2010) Is the clock drawing test a screening tool for the diagnosis of mild cognitive impairment? A systematic review. *International Psychogeriatrics* 22(1), 56-63

© NICE 2018. All rights reserved. See [Notice of rights](#).

Ferreira Daniel, Perestelo-Perez Lilisbeth, Westman Eric, Wahlund Lars-Olof, Sarria Antonio, and Serrano-Aguilar Pedro (2014) Meta-Review of CSF Core Biomarkers in Alzheimer's Disease: The State-of-the-Art after the New Revised Diagnostic Criteria. *Frontiers in aging neuroscience* 6, 47

Haddow Lewis John, Floyd Sian, Copas Andrew, and Gilson Richard John Cary (2013) A systematic review of the screening accuracy of the HIV Dementia Scale and International HIV Dementia Scale. *PloS one* 8(4), e61826

Health Quality, and Ontario (2006) Functional brain imaging: an evidence-based analysis. *Ontario health technology assessment series* 6(22), 1-79

Health Quality, and Ontario (2014) The appropriate use of neuroimaging in the diagnostic work-up of dementia: an evidence-based analysis. *Ontario health technology assessment series* 14(1), 1-64

Larner Andrew J, and Mitchell Alex J (2014) A meta-analysis of the accuracy of the Addenbrooke's Cognitive Examination (ACE) and the Addenbrooke's Cognitive Examination-Revised (ACE-R) in the detection of dementia. *International psychogeriatrics / IPA* 26(4), 555-63

Lin Jennifer S, O'Connor Elizabeth, Rossom Rebecca C, Perdue Leslie A, and Eckstrom Elizabeth (2013) Screening for cognitive impairment in older adults: A systematic review for the U.S. Preventive Services Task Force. *Annals of internal medicine* 159(9), 601-12

Masterman D L, Mendez M F, Fairbanks L A, and Cummings J L (1997) Sensitivity, specificity, and positive predictive value of technetium 99-HMPAO SPECT in discriminating Alzheimer's disease from other dementias. *Journal of geriatric psychiatry and neurology* 10(1), 15-21

McKeith I G, Ballard C G, Perry R H, Ince P G, O'Brien J T, Neill D, Lowery K, Jaros E, Barber R, Thompson P, Swann A, Fairbairn A F, and Perry E K (2000) Prospective validation of consensus criteria for the diagnosis of dementia with Lewy bodies. *Neurology* 54(5), 1050-8

Mitchell Alex J (2013) The Mini-Mental State Examination (MMSE): An update on its diagnostic validity for cognitive disorders. *Cognitive screening instruments: A practical approach.* , 15-46

Muayqil Taim, Gronseth Gary, and Camicioli Richard (2012) Evidence-based guideline: diagnostic accuracy of CSF 14-3-3 protein in sporadic Creutzfeldt-Jakob disease: report of the guideline development subcommittee of the American Academy of Neurology. *Neurology* 79(14), 1499-506

Mulder Cees, Verwey Nicolaas A, van der Flier , Wiesje M, Bouwman Femke H, Kok Astrid, van Elk , Evert J, Scheltens Philip, and Blankenstein Marinus A (2010) Amyloid-beta(1-42), total tau, and phosphorylated tau as cerebrospinal fluid biomarkers for the diagnosis of Alzheimer disease. *Clinical chemistry* 56(2), 248-53

Naqvi Raza M, Haider Sehrish, Tomlinson George, and Alibhai Shabbir (2015) Cognitive assessments in multicultural populations using the Rowland Universal Dementia Assessment Scale: a systematic review and meta-analysis. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 187(5), E169-75

O'Sullivan Dawn, O'Regan Niamh A, and Timmons Suzanne (2016) Validity and Reliability of the 6-Item Cognitive Impairment Test for Screening Cognitive Impairment: A Review. *Dementia and geriatric cognitive disorders* 42(1-2), 42-9

- Papathanasiou Nikolaos D, Boutsiadis Anastasios, Dickson John, and Bomanji Jamshed B (2012) Diagnostic accuracy of 123I-FP-CIT (DaTSCAN) in dementia with Lewy bodies: a meta-analysis of published studies. *Parkinsonism & related disorders* 18(3), 225-9
- Tang W, Huang Q, Wang Y, Wang Z Y, and Yao Y Y (2014) Assessment of CSF Aβ<sub>42</sub> as an aid to discriminating Alzheimer's disease from other dementias and mild cognitive impairment: A meta-analysis of 50 studies. *Journal of the Neurological Sciences* 345(1), 26-36
- Treglia Giorgio, and Cason Ernesto (2012) Diagnostic performance of myocardial innervation imaging using MIBG scintigraphy in differential diagnosis between dementia with lewy bodies and other dementias: a systematic review and a meta-analysis. *Journal of neuroimaging : official journal of the American Society of Neuroimaging* 22(2), 111-7
- Tsoi Kelvin K. F, Chan Joyce Y. C, Hirai Hoyee W, Wong Samuel Y. S, and Kwok Timothy C. Y (2015) Cognitive Tests to Detect Dementia: A Systematic Review and Meta-analysis. *JAMA internal medicine* 175(9), 1450-8
- van Harten , Argonde C, Kester Maartje I, Visser Pieter-Jelle, Blankenstein Marinus A, Pijnenburg Yolande A. L, van der Flier , Wiesje M, and Scheltens Philip (2011) Tau and p-tau as CSF biomarkers in dementia: a meta-analysis. *Clinical chemistry and laboratory medicine* 49(3), 353-66
- Wahlster Philip, Niederlander Charlotte, Kriza Christine, Schaller Sandra, and Kolominsky-Rabas Peter L (2013) Clinical assessment of amyloid imaging in Alzheimer's disease: a systematic review of the literature. *Dementia and geriatric cognitive disorders* 36(5-6), 263-78
- Yeo Jing Ming, Lim Xuxin, Khan Zubair, and Pal Suvankar (2013) Systematic review of the diagnostic utility of SPECT imaging in dementia. *European archives of psychiatry and clinical neuroscience* 263(7), 539-52
- Zerr I, Pocchiari M, Collins S, Brandel J P, de Pedro Cuesta, J , Knight R S, Bernheimer H, Cardone F, Delasnerie-Laupretre N, Cuadrado Corrales, N , Ladogana A, Bodemer M, Fletcher A, Awan T, Ruiz Bremon, A , Budka H, Laplanche J L, Will R G, and Poser S (2000) Analysis of EEG and CSF 14-3-3 proteins as aids to the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 55(6), 811-5
- Zhang Jian, Zhang Chun-Hua, Li Rong-Jie, Lin Xiao-Li, Chen Ying-Dao, Gao Huai-Qing, and Shi Sheng-Liang (2014) Accuracy of urinary AD7c-NTP for diagnosing Alzheimer's disease: a systematic review and meta-analysis. *Journal of Alzheimer's disease: JAD* 40(1), 153-9

#### **I.1.2.2 Systematic Reviews: sources of data**

- Archer Hilary A, Smailagic Nadja, John Christeena, Holmes Robin B, Takwoingi Yemisi, Coulthard Elizabeth J, and Cullum Sarah (2015) Regional cerebral blood flow single photon emission computed tomography for detection of Frontotemporal dementia in people with suspected dementia. *The Cochrane database of systematic reviews* 6, CD010896
- Chan CCH, Fage BA, Smailagic N, Gill SS, Herrmann N, Nikolaou V, Seitz DP. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a secondary care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 12. Art. No.: CD011414.
- Creavin Sam T, Wisniewski Susanna, Noel-Storr Anna H, Trevelyan Clare M, Hampton Thomas, Rayment Dane, Thom Victoria M, Nash Kirsty J. E, Elhamoui Hosam, Milligan Rowena, Patel Anish S, Tsivos Demitra V, Wing Tracey, Phillips Emma, Kellman Sophie M, Shackleton Hannah L, Singleton Georgina F, Neale Bethany E, Watton Martha E, and Cullum Sarah (2016) Mini-Mental State Examination (MMSE) for the detection of dementia in

clinically unevaluated people aged 65 and over in community and primary care populations. The Cochrane database of systematic reviews 1, CD011145

Davis Daniel H. J, Creavin Sam T, Yip Jennifer L. Y, Noel-Storr Anna H, Brayne Carol, and Cullum Sarah (2015) Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. The Cochrane database of systematic reviews 10, CD010775

Harrison Jennifer K, Fearon Patricia, Noel-Storr Anna H, McShane Rupert, Stott David J, and Quinn Terry J (2015) Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the diagnosis of dementia within a secondary care setting. The Cochrane database of systematic reviews 3, CD010772

McCleery Jenny, Morgan Shirlony, Bradley Kevin M, Noel-Storr Anna H, Ansorge Olaf, and Hyde Chris (2015) Dopamine transporter imaging for the diagnosis of dementia with Lewy bodies. The Cochrane database of systematic reviews 1, CD010633

Seitz DP, Fage BA, Chan CCH, Gill SS, Herrmann N, Smailagic N, Nikolaou V.

Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a primary care setting. Cochrane Database of Systematic Reviews 2014, Issue 12. Art. No.: CD011415.

### **I.1.2.3 Primary studies picked up by the SR search, from within the SRs and additional references from within primary studies**

Abdel-Aziz K, Larner AJ: Six-Item Cognitive Impairment Test (6CIT): pragmatic diagnostic accuracy study for dementia and MCI. *Int Psychogeriatr* 2015; 27: 991–997.

Andreasen N, Minthon L, Davidsson P, Vanmechelen E, Van-derstichele H, Winblad B, et al. Evaluation of CSF-tau and CSF-Abeta 42 as diagnostic markers for Alzheimer disease in clinical practice. *ArchNeurol* 2001; 58: 373–9.

Bahl JM, Heegaard NH, Falkenhorst G, Laursen H, Hogenhaven H, Molbak K, Jespersgaard C, Hougs L, Waldemar G, Johannsen P, Christiansen M. The diagnostic efficiency of biomarkers in sporadic Creutzfeldt-Jakob disease compared to Alzheimer's disease. *Neurobiol Aging* 2009; 30:1834–1841

Beinhoff U, Hilbert V, Bittner D, Gron G and Riepe MW. Screening for cognitive impairment: a triage for out-patient care. *Dementia and Geriatric Cognitive Disorders* 2005; 20: 278–285.

Bergman H, Chertkow H, Wolfson C, Stern J, Rush C, Whitehead V, Dixon R. HM-PAO (CERETEC) SPECT brain scanning in the diagnosis of Alzheimer's disease. *J Am Geriatr Soc* 1997; 45: 15–20

Beaudry P, Cohen P, Brandel JP, et al. 14-3-3 protein, neuron-specific enolase, and S-100 protein in cerebrospinal fluid of patients with Creutzfeldt-Jakob disease. *Dement Geriatr Cogn Disord* 1999; 10: 40–46.

Boutoleau-Bretonniere C, Lebouvier T, Delaroche O, Lamy E, Evrard C, Charriau T, et al. Value of neuropsychological testing, imaging, and CSF biomarkers for the differential diagnosis and prognosis of clinically ambiguous dementia. *Journal of Alzheimer's Disease* 2012; 28(2):323-36.

Brandt C, Bahl JC, Heegaard NH, Waldemar G, Johannsen P. Usability of cerebrospinal fluid biomarkers in a tertiary memory clinic. *Dement Geriatr Cogn Disord* 2008; 25: 553-558.

Burkhard PR; Sanchez JC; Landis T; Hochstrasser DF. CSF detection of the 14-3-3 protein in unselected patients with dementia. *Neurology*. 2001; 56: 1528-33

Callahan CM, Unverzagt FW, Hui SL, et al. Six-item screener to identify cognitive impairment among potential subjects for clinical research. *Med Care*. 2002; 40(9):771-81. PMID: 12218768

Carnero-Pardo 2013 {published and unpublished data} Carnero-Pardo C, Espejo-Martinez B, Lopez-Alcalde S, Espinosa-Garcia M, Saez-Zea C, Vilchez-Carrillo R, et al. Effectiveness and costs of phototest in dementia and cognitive impairment screening. *BMC Neurology* 2011; 11: 92. [Personal communication to Creavin et al. 2016 Cochrane Review. Email from: C Carnero-Pardo 31 October 2014.]

Chen C, Dong YH, Merchant R, Collinson S, Ting E, Quah SL et al. The Montreal Cognitive Assessment (MoCA) is superior to the Mini-Mental State Examination (MMSE) in detecting patients with moderate cognitive impairment, no dementia (CIND) and at high risk of dementia. Conference: Alzheimer's Association International Conference, Paris France. Conference Start: 20110716 Conference End: 20110721. 2011. [Data taken from by Davis et al. 2015 Cochrane Review]

Chohan G, Pennington C, Mackenzie JM, Andrews M, Everington D, Will RG, Knight RS, Green AJ. The role of cerebrospinal fluid 14-3-3 and other proteins in the diagnosis of sporadic Creutzfeldt-Jakob disease in the UK: a 10-year review. *J Neurol Neurosurg Psychiatry*. 2010; 81: 1243-8.

Cruz-Orduna I, Bellon JM, Torrero P, Aparicio E, Sanz A, Mula N, et al. Detecting MCI and dementia in primary care: effectiveness of the MMS, the FAQ and the IQCODE. *Family Practice* 2012; 29: 401-6.

Cuadrado-Corrales N, Jiménez-Huete A, Albo C, Hortigüela R, Vega L, Cerrato L, Sierra-Moros M, Rábano A, de Pedro-Cuesta J, Calero M. Impact of the clinical context on the 14-3-3 test for the diagnosis of sporadic CJD. *BMC Neurology* 2006; 6: 25

Davis HF, Skolasky RL, Selnes OA, Burgess DM, McArthur JC. Assessing HIV-associated dementia: modified HIV dementia scale versus the grooved pegboard. *The AIDS reader* 2002; 12: 29-31.

Dobert N, Pantel J, Frolich L, Hamscho N, Menzel C, Grunwald F. Diagnostic value of FDG-PET and HMPAO-SPET in patients with mild dementia and mild cognitive impairment: Metabolic index and perfusion index. *Dement Geriatr Cogn Disord* 2005; 20: 63-70.

Dumurgier J, Schraen S, Gabelle A, Vercausse O, Bombois S, Laplanche J-L, Peoc'h K et al. Cerebrospinal fluid amyloid- $\beta$  42/40 ratio in clinical setting of memory centers: a multicentric study. *Alzheimer's Research & Therapy* 2015; 7:30-38.

Engedal K, Snaedal J, Hoegh P, Jelic V, Bo Andersen B, Naik M, Wahlund LO, Oeksengaard AR. Quantitative EEG applying the statistical recognition pattern method: a useful tool in dementia diagnostic workup. *Dement Geriatr Cogn Disord* 2015;40: 1–12.

Estorch M, Camacho V, Paredes P, et al. Cardiac (123)I-metaiodobenzylguanidine imaging allows early identification of dementia with Lewy bodies during life. *Eur J Nucl Med Mol Imaging* 2008; 35: 1636-1641.

Flicker L, Logiudice D, Carlin JB, Ames D. The predictive value of dementia screening instruments in clinical populations. *International Journal of Geriatric Psychiatry* 1997; 12: 203–9.

Frisoni GB, Prestia A, Zanetti O, Galluzzi S, Romano M, Cotelli M, Gennarelli M, Binetti G, Bocchio L, Paghera B, Amicucci G, Bonetti M, Benussi L, Ghidoni R, Geroldi C. Markers of Alzheimer's disease in a population attending a memory clinic. *Alzheimers Dement* 2009; 5: 307- 317.

- Gabelle A, Dumurgier J, Vercruysse O, Paquet C, Bombois S, Laplanche J-L., et al. Impact of the 2008-2012 French Alzheimer plan on the use of cerebrospinal fluid biomarkers in research memory center: the PLM Study. *J. Alzheimers Dis.* 2013; 34: 7–305.
- Garcia M, Perlado Ortiz de Pinedo F. Cognitive deterioration: use of the short version of the Informant Test (IQCODE) in the geriatrics consultations. *Revista Española de Geriatria y Gerontología* 2002; 37: 81–5. [In Spanish: data source Harrison et al. 2015 Cochrane Review]
- Gold G, Bouras C, Canuto A, Bergallo M, Herrmann FR, Hof PR, Mayor P-A, Michel J-P, Giannakopoulos P. Clinicopathological Validation Study of Four Sets of Clinical Criteria for Vascular Dementia. *Am J Psychiatry* 2002; 159: 82–87
- Goncalves DC, Arnold E, Appadurai K, Byrne GJ. Case finding in dementia: comparative utility of three brief instruments in the memory clinic setting. *International Psychogeriatrics* 2011; 23: 788–96.
- Goodman I, Golden G, Flitman S, Xie K, McConville M, Levy S, Zimmerman E, Lebedeva Z, Richter R, Minagar A, Averback P. A multi-center blinded prospective study of urine neural thread protein measurements in patients with suspected Alzheimer's disease. *J Am Med Dir Assoc* 2007; 8: 21-30.
- Hancock P, Lerner AJ. Diagnostic utility of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) and its combination with the Addenbrooke's Cognitive Examination-Revised (ACE-R) in a memory clinic-based population. *International Psychogeriatrics* 2009; 21: 526–30.
- Hentschel F, Kreis M, Damian M, Krumm B, Frolich L. The clinical utility of structural neuroimaging with MRI for diagnosis and differential diagnosis of dementia: a memory clinic study. *Int J Geriatr Psychiatry.* 2005; 20: 645-50.
- Hoffman JM, Welsh-Bohmer KA, Hanson M, Krain B, Hulette C, Earl N et al. FDG PET imaging in patients with pathologically verified dementia. *J Nucl Med.* 2000; 41: 1920-8.
- Holman BL, Johnson KA, Gerada B, Carvalho PA, Satlin A. The scintigraphic appearance of Alzheimer's disease: a prospective study using technetium-99m-HMPAO SPECT. *J Nucl Med* 1992; 33: 181–185.
- Ibach B, Binder H, Dragon M, Poljansky S, Haen E, Schmitz E, et al. Cerebrospinal fluid tau and beta-amyloid in Alzheimer patients, disease controls and an age-matched random sample. *Neurobiol Aging* 2006; 27: 1202–11.
- Jagust W, Reed B, Mungas D, Ellis W, De Carli C. What does fluorodeoxyglucose PET imaging add to a clinical diagnosis of dementia? *Neurology.* 2007; 69: 871-7.
- Kemp PM, Clyde K, Holmes C. Impact of 123I-FP-CIT (DaTSCAN) SPECT on the diagnosis and management of patients with dementia with Lewy bodies: a retrospective study. *Nucl Med Commun* 2011; 32: 298-302.
- Kenney K, Brechtel C, Takahashi H, Kurohara K, Anderson P, Gibbs CJ Jr. An enzyme-linked immunosorbent assay to quantify 14-3-3 proteins in the cerebrospinal fluid of suspected Creutzfeldt-Jakob disease patients. *Ann Neurol* 2000; 48: 395–398.
- Knafelc R, Lo Giudice D, Harrigan S, Cook R, Flicker L, Mackinnon A, et al. The combination of cognitive testing and an informant questionnaire in screening for dementia. *Age and Ageing* 2003; 32: 541–7.
- Kukull WA, Larson EB, Teri L, Bowen J, McCormick W, Pfanschmidt ML. The mini mental state examination score and the clinical diagnosis of dementia. *J Clin Epidemiol* 1994; 47: 1061-1067.

- Larner AJ. Addenbrooke's Cognitive Examination (ACE) for the diagnosis and differential diagnosis of dementia. *Clinical Neurology and Neurosurgery* 2007; 109 : 491–494
- Launes J, Sulkava R, Erkinjuntti T, Nikkinen P, Lindroth L, Liewendahl K, et al. 99Tcm-HMPAO SPECT in suspected dementia. *Nuclear Medicine Communications* 1991;12: 757–65.
- Lemstra AW, van Meegen MT, Vreyling JP, Meijerink PH, Jansen GH, Bulk S, Baas F, van Gool WA. 14-3-3 testing in diagnosing Creutzfeldt-Jakob disease: A prospective study in 112 patients. *Neurology* 2000; 55: 514-6
- Maddalena A, Papassotiropoulos A, Muller-Tillmanns B, et al. Biochemical diagnosis of Alzheimer disease by measuring the cerebrospinal fluid ratio of phosphorylated tau protein to beta-amyloid peptide (42). *Arch Neurol* 2003; 60: 1202–6.
- Masterman DL, Mendez MF, Fairbanks LA, Cummings JL. Sensitivity, specificity, and positive predictive value of technetium 99-HMPAO SPECT in discriminating Alzheimer's disease from other dementias. *J Geriatr Psychiatry Neurol* 1997; 10: 15–21
- Mathuranath PS, Nestor PJ, Berrios G.E, Rakowicz W and Hodges JR. A brief cognitive test battery to differentiate Alzheimer's disease and frontotemporal dementia. *Neurology* 2000; 55: 1613– 1620
- Mayeux R, Saunders A, Shea S, Mirra S, Evans D, Roses AD, Hyman BT et al. Utility of the Apolipoprotein E genotype in the diagnosis of Alzheimer's disease. *NEJM* 1998; 338: 506-511.
- McMurdo ME, Grant DJ, Kennedy NS, Gilchrist J, Findlay D, McLennan JM. The value of HMPAO SPECT scanning in the diagnosis of early Alzheimer's disease in patients attending a memory clinic. *Nucl Med Commun* 1994; 15: 405-409
- Milian M, Leiherr AM, Straten G, Muller S, Leyhe T, Eschweiler GW. The Mini-Cog versus the Mini-Mental State Examination and the Clock Drawing Test in daily clinical practice: screening value in a German Memory Clinic. *International Psychogeriatrics* 2012; 24: 766-74
- Nielsen TR, Andersen BB, Gottrup H, et al. Validation of the RUDAS for multicultural screening in Danish memory clinics. *Dement Geriatr Cogn Disord* 2013; 36: 354-62.
- O'Brien JT, Metcalfe S, Swann A, et al. Medial temporal lobe width on CT scanning in Alzheimer's disease: comparison with vascular dementia, depression and dementia with Lewy bodies. *Dement Geriatr Cogn Disord*. 2000;11: 114-118
- O'Brien JT, McKeith IG, Walker Z, Tatsch K, Booij J et al. Diagnostic accuracy of 123I-FP-CIT SPECT in possible dementia with Lewy bodies. *The British Journal of Psychiatry* 2009; 194: 34-39.
- Ossenkoppele R, Prins ND, Pijnenburg YA, Lemstra AW, van der Flier WM, Adriaanse SF, Windhorst AD, Handels RL, Wolfs CA, Aalten P, Verhey FR, Verbeek MM, van Buchem MA, Hoekstra OS, Lammertsma AA, Scheltens P, van Berckel BN: Impact of molecular imaging on the diagnostic process in a memory clinic. *Alzheimers Dement* 2013; 9: 414–421.
- Read SL, Miller BL, Mena I, Kim R, Itabashi H, Darby A. SPECT in dementia: clinical and pathological correlation. *Journal of the American Geriatrics Society* 1995; 43: 1243–7
- Rollin-Silliare A, Bombois S, Deramecourt V, Steinert- Emptaz A, Salleron J, Morvan J, et al. Contribution of single photon emission computed tomography to the differential diagnosis of dementia in a memory clinic. *Journal of Alzheimer's Disease* 2012; 30: 833–45.

- Sager MA, Hermann BP, LaRue A and Woodard JL. Screening for dementia in community-based memory clinics. *Wisconsin Medical Journal*, 2006; 105: 25–29.
- Schroter A, Zerr I, Henkel K et al. Magnetic resonance imaging in the clinical diagnosis of Creutzfeldt–Jakob disease. *Arch Neurol* 2000; 57: 1751-1757.
- Sikkes SA, Van den Berg MT, Knol DL, De-Lange-de Klerk ES, Scheltens P, Uitdehaag BM, et al. How useful is IQCODE for discriminating between Alzheimer’s disease, mild cognitive impairment and subjective memory complaints? *Dementia and Geriatric Cognitive Disorders* 2010; 30: 411–6.
- Silverman DHS, Small GW, Chang CY, Lu CS, Kung De Abarto MA, Chen W, et al. Positron emission tomography in evaluation of dementia; regional brain metabolism and long-term outcome. *JAMA*. 2001; 286: 2120-7.
- Skinner S, Adewale AJ, DeBlock L, Gill MJ, Power C. Neurocognitive screening tools in HIV/AIDS: comparative performance among patients exposed to antiretroviral therapy. *HIV Medicine*, 2009; 10: 246–252.
- Talbot PR, Lloyd JJ, Snowden JS, Neary D, Testa HJ. A clinical role for 99mTc-HMPAO SPECT in the investigation of dementia? *Journal of Neurology, Neurosurgery, and Psychiatry* 1998; 63:306-13.
- Terpening Z, Cordato NJ, Hepner IJ, Lucas SK, Lindley RI. Utility of the Addenbrooke’s Cognitive Examination- Revised for the diagnosis of dementia syndromes. *Australas J Ageing*. 2011; 30: 113-8.
- Toledo JB, Brettschneider J, Grossman M, Arnold SE et al. CSF biomarkers cutoffs: the importance of coincident neuropathological diseases. *Act Neuropathol* 2012; 124: 23-35.
- G.Treglia, E.Cason, P.Cortelli, A.Gabellini, R.Liguori, A.Bagnato, A.Giordano, G. Fagioli, Iodine-123metaiodobenzylguanidine scintigraphy and iodine-123 ioflupane single photon emission computed tomography in Lewy body diseases: complementary or alternative techniques? *J. Neuroimaging* 24 (2012) 149–154.
- Tschampa HJ, Kallenberg K, Urbach H, Meissner B, Nicolay C, Kretschmar HA, et al. MRI in the diagnosis of sporadic Creutzfeldt-Jakob disease: a study on inter-observer agreement. *Brain*. 2005; 128: 9-33.
- Van Everbroeck B, Quoilin S, Boons J, Martin JJ, Cras P. A prospective study of CSF markers in 250 patients with possible Creutzfeldt-Jakob disease. *J neurol Neurosurg Psychiatry* 2003; 74: 1210–4.
- Van Everbroeck B, Dobbeleir I, De Waele M, De Deyn P, Martin JJ, Cras P. Differential diagnosis of 201 possible Creutzfeldt-Jakob disease patients. *J Neurol*. 2004; 251:298-304.
- Velakoulis D, Lloyd JH. The role of SPECT scanning in a neuropsychiatry unit. *Aust N Z J Psychiatry* 1998; 32: 511-22.
- Walker RWH, Walker Z. Dopamine transporter single photon emission computerized tomography in the diagnosis of dementia with Lewy bodies. *Movement Disorders* 2009; 24: S754–9.
- Zerr I, Bodemer M, Gefeller O, et al. Detection of 14-3-3 protein in the cerebrospinal fluid supports the diagnosis of Creutzfeldt-Jakob disease. *Ann Neurol* 1998; 43: 32– 40.
- Zerr I, Pocchiari M, Collins S, et al. Analysis of EEG and CSF 14-3-3 proteins as aids to the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2000; 55: 811– 815.

#### I.1.2.4 Primary studies from the additional primary study search

Alexander S K, Rittman T, Xuereb J H, Bak T H, Hodges J R, and Rowe J B (2014) Validation of the new consensus criteria for the diagnosis of corticobasal degeneration. *Journal of Neurology, and Neurosurgery & Psychiatry* 85(8), 925-9

Ampuero Israel, Alegre-Abarrategui Javier, Rodal Izaskun, Espana Antonio, Ros Raquel, Sendon Jose Luis Lopez, Galloway Eva Garcia, Cervello Angeles, Caminero Ana Belen, Zabala Antxon, Erro Elena, Jarauta Fernando, Morlan Lorenzo, Lopez-Valdes Eva, Aladro Yolanda, Seijo Manuel, Rivas Guillermo Garcia, Munoz David G, de Yebenes , and Justo Garcia (2009) On the diagnosis of CADASIL. *Journal of Alzheimer's disease: JAD* 17(4), 787-94

Apolinario Daniel, Lichtenthaler Daniel Gomes, Magaldi Regina Miksian, Soares Aline Thomaz, Busse Alexandre Leopold, Amaral Jose Renato das Gracias, Jacob-Filho Wilson, and Brucki Sonia Maria Dozzi (2016) Using temporal orientation, category fluency, and word recall for detecting cognitive impairment: the 10-point cognitive screener (10-CS). *International journal of geriatric psychiatry* 31(1), 4-12

Arslan Esra, Ekmekcioglu Ozgul, Gortan Fatma Arzu, Engin Akcan, Zeynep Funda, Erkan Melih Engin, Emul Haci Murat, Halac Metin, Cermik Tevfik Fikret, and Sonmezoglu Kerim (2015) The value of FDG-PET/CT by using 3-dimensional stereotactic surface projection software analysis in the differential diagnosis of dementia. *Turkish journal of medical sciences* 45(5), 1149-58

Bacchetta J P, Kovari E, Merlo M, Canuto A, Herrmann F R, Bouras C, Gold G, Hof P R, and Giannakopoulos P (2007) Validation of clinical criteria for possible vascular dementia in the oldest-old. *Neurobiology of Aging* 28(4), 579-85

Bastide L, De Breucker, S , Van den Berge , M , Fery P, Pepersack T, and Bier J C (2012) The Addenbrooke's Cognitive Examination Revised is as effective as the original to detect dementia in a French-speaking population. *Dementia and geriatric cognitive disorders* 34(5-6), 337-43

Berger Gabriele, Frolich Lutz, Weber Bernhard, and Pantel Johannes (2008) Diagnostic accuracy of the clock drawing test: the relevance of "time setting" in screening for dementia. *Journal of geriatric psychiatry and neurology* 21(4), 250-60

Bonello M, and Lerner A J (2016) Applause sign: Screening utility for dementia and cognitive impairment. *Postgraduate Medicine* 128(2), 250-253

Chan Yung, Yeung Kwok-Hung, Ho Hing-Fung, Ho King-Man, Tin-Keung Lam, Edman , Leung Wai-Lin, and Kam Kai-Man (2014) Use of cerebrospinal fluid enzyme immunoassay for diagnosis of neurosyphilis. *International journal of STD & AIDS* 25(8), 571-8

Christensen IB Thrane, Larsson Elna-Marie, Holm Ida E, Nielsen Ole B. F, and Andersen Stig (2017) Olfactory testing in consecutive patients referred with suspected dementia. *BMC geriatrics* 17(1), 129

Coulthart Michael B, Jansen Gerard H, Olsen Elina, Godal Deborah L, Connolly Tim, Choi Bernard C. K, Wang Zheng, and Cashman Neil R (2011) Diagnostic accuracy of cerebrospinal fluid protein markers for sporadic Creutzfeldt-Jakob disease in Canada: a 6-year prospective study. *BMC neurology* 11, 133

Coutinho G, De Oliveira-Souza , R , Moll J, Tovar-Moll F, and Mattos P (2013) Is it possible to identify individuals with mild cognitive impairment and Alzheimer's disease using a 30-minute neuropsychological battery?. *Revista de Psiquiatria Clinica* 40(4), 139-143

Duits Flora H, Teunissen Charlotte E, Bouwman Femke H, Visser Pieter-Jelle, Mattsson Niklas, Zetterberg Henrik, Blennow Kaj, Hansson Oskar, Minthon Lennart, Andreasen Niels, Marcusson Jan, Wallin Anders, Rikkert Marcel Olde, Tsolaki Magda, Parnetti Lucilla, Herukka Sanna-Kaisa, Hampel Harald, De Leon , Mony J, Schroder Johannes, Aarsland Dag, Blankenstein Marinus A, Scheltens Philip, van der Flier , and Wiesje M (2014) The cerebrospinal fluid "Alzheimer profile": easily said, but what does it mean?. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 10(6), 713-723.e2

Dumaresq Jeannot, Langevin Stephanie, Gagnon Simon, Serhir Bouchra, Deligne Benoit, Tremblay Cecile, Tsang Raymond S. W, Fortin Claude, Coutlee Francois, and Roger Michel (2013) Clinical prediction and diagnosis of neurosyphilis in HIV-infected patients with early Syphilis. *Journal of clinical microbiology* 51(12), 4060-6

Ferman T J, Boeve B F, Smith G E, Lin S C, Silber M H, Pedraza O, Wszolek Z, Graff-Radford N R, Uitti R, Van Gerpen , J , Pao W, Knopman D, Pankratz V S, Kantarci K, Boot B, Parisi J E, Dugger B N, Fujishiro H, Petersen R C, and Dickson D W (2011) Inclusion of RBD improves the diagnostic classification of dementia with Lewy bodies. *Neurology* 77(9), 875-82

Fourier A, Dorey A, Perret-Liaudet A, and Quadrio I (2017) Detection of CSF 14-3-3 Protein in Sporadic Creutzfeldt-Jakob Disease Patients Using a New Automated Capillary Western Assay. *Molecular neurobiology* [Epub ahead of print]

Foutz A, Appleby B S, Hamlin C, Liu X, Yang S, Cohen Y, Chen W, Blevins J, Fausett C, Wang H, Gambetti P, Zhang S, Hughson A, Tatsuoka C, Schonberger L B, Cohen M L, Caughey B, and Safar J G (2017) Diagnostic and prognostic value of human prion detection in cerebrospinal fluid. *Annals of Neurology* 81(1), 79-92

Galvin J, Tolea M, and Sidhu J (2015) Improving the clinical detection and diagnosis of lewy body dementia using a composite risk score. *Neurodegenerative Diseases* 15, 1457

Goldstein Felicia C, Ashley Angela V, Miller Eric, Alexeeva Olga, Zanders Lavezza, and King Veronique (2014) Validity of the montreal cognitive assessment as a screen for mild cognitive impairment and dementia in African Americans. *Journal of geriatric psychiatry and neurology* 27(3), 199-203

Gustafson Lars, Englund Elisabet, Brunnstrom Hans, Brun Arne, Erikson Catarina, Warkentin Siegbert, and Passant Ulla (2010) The accuracy of short clinical rating scales in neuropathologically diagnosed dementia. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry* 18(9), 810-20

Hamlin Clive, Puoti Gianfranco, Berri Sally, Sting Elliott, Harris Carrie, Cohen Mark, Spear Charles, Bizzi Alberto, Debanne Sara M, and Rowland Douglas Y (2012) A comparison of tau and 14-3-3 protein in the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 79(6), 547-52

Hancock P, and Larner A J (2011) Test Your Memory test: diagnostic utility in a memory clinic population. *International journal of geriatric psychiatry* 26(9), 976-80

Hanyu Haruo, Shimizu Soichiro, Hirao Kentaro, Sakurai Hirofumi, Iwamoto Toshihiko, Chikamori Taishiro, Hida Satoshi, Yamashina Akira, Koizumi Kiyoshi, and Abe Kimihiko (2006) The role of 123I-metaiodobenzylguanidine myocardial scintigraphy in the diagnosis of Lewy body disease in patients with dementia in a memory clinic. *Dementia and geriatric cognitive disorders* 22(5-6), 379-84

Harris J M, Gall C, Thompson J C, Richardson A M, Neary D, du Plessis, D , Pal P, Mann D M, Snowden J S, and Jones M (2013) Sensitivity and specificity of FTDC criteria for behavioral variant frontotemporal dementia. *Neurology* 80(20), 1881-7

Heath C A, Cooper S A, Murray K, Lowman A, Henry C, MacLeod M A, Stewart G E, Zeidler M, MacKenzie J M, Ironside J W, Summers D M, Knight R S, and Will R G (2010) Validation of diagnostic criteria for variant Creutzfeldt-Jakob disease. *Annals of Neurology* 67(6), 761-70

Jahn Holger, Wittke Stefan, Zurbig Petra, Raedler Thomas J, Arlt Sonke, Kellmann Markus, Mullen William, Eichenlaub Martin, Mischak Harald, and Wiedemann Klaus (2011) Peptide fingerprinting of Alzheimer's disease in cerebrospinal fluid: identification and prospective evaluation of new synaptic biomarkers. *PloS one* 6(10), e26540

Jubb Michael T, and Evans Jonathan J (2015) An Investigation of the Utility of the Addenbrooke's Cognitive Examination III in the Early Detection of Dementia in Memory Clinic Patients Aged over 75 Years. *Dementia and geriatric cognitive disorders* 40(3-4), 222-32

Kaneta Tomohiro, Nakatsuka Masahiro, Nakamura Kei, Seki Takashi, Yamaguchi Satoshi, Tsuboi Masahiro, and Meguro Kenichi (2016) Improved Diagnostic Accuracy of SPECT Through Statistical Analysis and the Detection of Hot Spots at the Primary Sensorimotor Area for the Diagnosis of Alzheimer Disease in a Community-Based Study: "The Osaka-Tajiri Project". *Clinical nuclear medicine* 41(1), e1-6

Kerklaan B J, Van Berckel , B N M, Herholz K, Dols A, Van Der Flier , W M, Scheltens P, and Pijnenburg Y A. L (2014) The added value of 18-fluorodeoxyglucose-positron emission tomography in the diagnosis of the behavioral variant of frontotemporal dementia. *American Journal of Alzheimer's Disease and other Dementias* 29(7), 607-613

Kiesmann Michele, Chanson Jean-Baptiste, Godet Julien, Vogel Thomas, Schweiger Laetitia, Chayer Said, and Kaltenbach Georges (2013) The Movement Disorders Society criteria for the diagnosis of Parkinson's disease dementia: their usefulness and limitations in elderly patients. *Journal of neurology* 260(10), 2569-79

Knapskog A B, Engedal K, and Braekhus A (2016) Performance of cerebrospinal fluid biomarkers of Alzheimer disease in a memory clinic in Norway. *Alzheimer Disease and Associated Disorders* 30(1), 8-14

Koikkalainen Juha, Rhodius-Meester Hanneke, Tolonen Antti, Barkhof Frederik, Tijms Betty, Lemstra Afina W, Tong Tong, Guerrero Ricardo, Schuh Andreas, Ledig Christian, Rueckert Daniel, Soininen Hilka, Remes Anne M, Waldemar Gunhild, Hasselbalch Steen, Mecocci Patrizia, van der Flier , Wiesje , and Lotjonen Jyrki (2016) Differential diagnosis of neurodegenerative diseases using structural MRI data. *NeuroImage. Clinical* 11, 435-49

Larner A J (2015) AD8 Informant Questionnaire for Cognitive Impairment: Pragmatic Diagnostic Test Accuracy Study. *Journal of geriatric psychiatry and neurology* 28(3), 198-202

Lattanzio Francesca, Abu-Rumeileh Samir, Franceschini Alessia, Kai Hideaki, Amore Giulia, Poggiolini Ilaria, Rossi Marcello, Baiardi Simone, McGuire Lynne, Ladogana Anna, Pocchiari Maurizio, Green Alison, Capellari Sabina, and Parchi Piero (2017) Prion-specific and surrogate CSF biomarkers in Creutzfeldt-Jakob disease: diagnostic accuracy in relation to molecular subtypes and analysis of neuropathological correlates of p-tau and Abeta42 levels. *Acta neuropathologica* 133(4), 559-578

Leitao M J, Baldeiras I, Almeida M R, Ribeiro M H, Santos A C, Ribeiro M, Tomas J, Rocha S, Santana I, and Oliveira C R (2016) Sporadic Creutzfeldt-Jakob disease diagnostic accuracy is improved by a new CSF ELISA 14-3-3gamma assay. *Neuroscience* 322, 398-407

Malhotra Chetna, Chan Angelique, Matchar David, Seow Dennis, Chuo Adeline, and Do Young Kyung (2013) Diagnostic performance of short portable mental status questionnaire

for screening dementia among patients attending cognitive assessment clinics in Singapore. *Annals of the Academy of Medicine, and Singapore* 42(7), 315-9

Manabe Y, Inui Y, Toyama H, and Kosaka K (2017) 123I-metaiodobenzylguanidine myocardial scintigraphy with early images alone is useful for the differential diagnosis of dementia with Lewy bodies. *Psychiatry Research - Neuroimaging* 261, 75-79

Mendez M F, Shapira J S, McMurtray A, Licht E, and Miller B L (2007) Accuracy of the clinical evaluation for frontotemporal dementia. *Archives of Neurology* 64(6), 830-5

Mormont E, Jamart J, and Robaye L (2012) Validity of the five-word test for the evaluation of verbal episodic memory and dementia in a memory clinic setting. *Journal of Geriatric Psychiatry and Neurology* 25(2), 78-84

Motara H, Olusoga T, Russell G, Jamieson S, Ahmed S, Brindle N, Pillai A, Scarsbrook A F, Patel C N, and Chowdhury F U (2017) Clinical impact and diagnostic accuracy of 2-[18F]-fluoro-2-deoxy-D-glucose positron-emission tomography/computed tomography (PET/CT) brain imaging in patients with cognitive impairment: a tertiary centre experience in the UK. *Clinical Radiology* 72(1), 63-73

Panegyres P K, Rogers J M, McCarthy M, Campbell A, and Wu J S (2009) Fluorodeoxyglucose-positron emission tomography in the differential diagnosis of early-onset dementia: a prospective, community-based study. *BMC Neurology* 9, 41

Postel-Vinay Nicolas, Hanon Olivier, Clerson Pierre, Brown Jeremy M, Menard Joel, Paillaud Elena, Alonso Eliana, Pasquier Florence, Pariel Sylvie, Belliard Serge, Pere Jean-Jacques, and Belmin Joel (2014) Validation of the Test Your Memory (F-TYM Test) in a French memory clinic population. *The Clinical neuropsychologist* 28(6), 994-1007

Rohan Zdenek, Smetakova Magdalena, Kukal Jaromir, Rusina Robert, and Matej Radoslav (2015) Proteinase-activated receptor 2 and disease biomarkers in cerebrospinal fluid in cases with autopsy-confirmed prion diseases and other neurodegenerative diseases. *BMC neurology* 15, 50

Sakamoto Fumi, Shiraishi Shinya, Yoshida Morikatsu, Tomiguchi Seiji, Hirai Toshinori, Namimoto Tomohiro, Hashimoto Mamoru, Ikeda Manabu, Uetani Hiroyuki, and Yamashita Yasuyuki (2014) Diagnosis of dementia with Lewy bodies: diagnostic performance of combined 123I-IMP brain perfusion SPECT and 123I-MIBG myocardial scintigraphy. *Annals of nuclear medicine* 28(3), 203-11

Sakamoto Fumi, Shiraishi Shinya, Tsuda Noriko, Hashimoto Mamoru, Tomiguchi Seiji, Ikeda Manabu, and Yamashita Yasuyuki (2017) Diagnosis of dementia with Lewy bodies: can 123I-IMP and 123I-MIBG scintigraphy yield new core features?. *The British journal of radiology* 90(1070), 20160156

Siritho Sasitorn, Senanarong Vorapun, Nako Arpas, Chotinaiwattarakul Wattananchai, Jamjumrus Piyanuj, Udompunthuruk Suthipol, and Pongvarin Niphon (2006) Use of Hachinski Ischemic Score in the memory clinic: Thai experience. *Journal of the Medical Association of Thailand = Chotmaihet thangphaet* 89(11), 1822-8

Skjerve Arvid, Nordhus Inger Hilde, Engedal Knut, Braekhus Anne, Nygaard Harald A, Pallesen Stale, and Haugen Per Kristian (2008) Validation of the Seven Minute Screen and Syndrom Kurztest among elderly Norwegian outpatients. *International psychogeriatrics* 20(4), 807-14

Skogseth Ragnhild, Hortobagyi Tibor, Soennesyn Hogne, Chwiszczuk Luiza, Ffytche Dominic, Rongve Arvid, Ballard Clive, and Aarsland Dag (2017) Accuracy of Clinical Diagnosis of Dementia with Lewy Bodies versus Neuropathology. *Journal of Alzheimer's disease : JAD* 59(4), 1139-1152

- Slaets Sylvie, Van Acker , Frank , Versijpt Jan, Hauth Lothar, Goeman Johan, Martin Jean-Jacques, De Deyn , Peter Paul, and Engelborghs Sebastiaan (2015) Diagnostic value of MIBG cardiac scintigraphy for differential dementia diagnosis. *International journal of geriatric psychiatry* 30(8), 864-9
- Streit Sven, Limacher Andreas, Zeller Andreas, and Burge Markus (2015) Detecting dementia in patients with normal neuropsychological screening by Short Smell Test and Palmo-Mental Reflex Test: an observational study. *BMC geriatrics* 15, 90
- Suppa Per, Anker Ulrich, Spies Lothar, Bopp Irene, Ruegger-Frey Brigitte, Klaghofer Richard, Gocke Carola, Hampel Harald, Beck Sacha, and Buchert Ralph (2015) Fully automated atlas-based hippocampal volumetry for detection of Alzheimer's disease in a memory clinic setting. *Journal of Alzheimer's disease : JAD* 44(1), 183-93
- Tagliapietra M, Zanusso G, Fiorini M, Bonetto N, Zarantonello G, Zambon A, Ermani M, Monaco S, Manara R, and Cagnin A (2013) Accuracy of diagnostic criteria for sporadic creutzfeldt-jakob disease among rapidly progressive dementia. *Journal of Alzheimer's Disease* 34(1), 231-8
- Thomas AJ, Attems J, Colloby Sj, O'Brien Jt, McKeith I, Walker R, Lee L, Burn D, Lett Dj, and Walker Z (2017) Autopsy validation of 123 I-FP-CIT dopaminergic neuroimaging for the diagnosis of DLB. *Neurology* 88(3), 276-283
- Tripathi Madhavi, Tripathi Manjari, Vibha Deepti, Gowda Naveen, Bal Chandrashekhar, and Malhotra Arun (2010) Tc-99m ethylcysteinate dimer SPECT in the differential diagnosis of dementias. *Neurology India* 58(6), 857-62
- Vijverberg Everard G. B, Dols Annemiek, Krudop Welmoed A, Peters Anne, Kerssens Cora J, van Berckel , Bart N M, Wattjes Mike P, Barkhof Frederik, Gossink Flora, Prins Niels D, Stek Max L, Scheltens Philip, and Pijnenburg Yolande A. L (2016a) Diagnostic Accuracy of the Frontotemporal Dementia Consensus Criteria in the Late-Onset Frontal Lobe Syndrome. *Dementia and geriatric cognitive disorders* 41(3-4), 210-9
- Vijverberg Everard G. B, Wattjes Mike P, Dols Annemiek, Krudop Welmoed A, Moller Christiane, Peters Anne, Kerssens Cora J, Gossink Flora, Prins Niels D, Stek Max L, Scheltens Philip, van Berckel, Bart N M, Barkhof Frederik, and Pijnenburg Yolande A. L (2016b) Diagnostic Accuracy of MRI and Additional [18F]FDG-PET for Behavioral Variant Frontotemporal Dementia in Patients with Late Onset Behavioral Changes. *Journal of Alzheimer's disease: JAD* 53(4), 1287-97
- Walker Z, Jaros E, Walker R W, Lee L, Costa D C, Livingston G, Ince P G, Perry R, McKeith I, and Katona C L (2007) Dementia with Lewy bodies: a comparison of clinical diagnosis, FP-CIT single photon emission computed tomography imaging and autopsy. *Journal of Neurology, and Neurosurgery & Psychiatry* 78(11), 1176-81
- Yakushev Igor, Bartenstein Peter, Siessmeier Thomas, Hiemke Christoph, Scheurich Armin, Lotz Johannes, Fellgiebel Andreas, and Muller Matthias J (2010) Cerebrospinal fluid tau protein levels and 18F-fluorodeoxyglucose positron emission tomography in the differential diagnosis of Alzheimer's disease. *Dementia and geriatric cognitive disorders* 30(3), 245-53
- Yeung P Y, Wong L L, Chan C C, Leung Jess L. M, and Yung C Y (2014) A validation study of the Hong Kong version of Montreal Cognitive Assessment (HK-MoCA) in Chinese older adults in Hong Kong. *Hong Kong medical journal = Xianggang yi xue za zhi* 20(6), 504-10
- Zerr I, Kallenberg K, Summers D M, Romero C, Taratuto A, Heinemann U, Breithaupt M, Varges D, Meissner B, Ladogana A, Schuur M, Haik S, Collins S J, Jansen G H, Stokin G B, Pimentel J, Hewer E, Collie D, Smith P, Roberts H, Brandel J P, van Duijn , C , Pocchiari M, Bogue C, Cras P, Will R G, and Sanchez-Juan P (2009) Updated clinical diagnostic criteria

for sporadic Creutzfeldt-Jakob disease.[Erratum appears in Brain. 2012 Apr;135(Pt 4):1335].  
Brain 132(Pt 10), 2659-68

Zwan MD, Bouwman Fh, Konijnenberg E, Flier Wm, Lammertsma Aa, Verhey Frj, Aalten P, Berckel Bnm, and Scheltens P (2017) Diagnostic impact of flutemetamol PET in early-onset dementia. *Alzheimer's research & therapy* 9(1) (no pagination),

### **I.1.3 Distinguishing dementia from delirium or delirium with dementia**

Cole Martin G, McCusker Jane, Dendukuri Nandini, and Han Ling (2002) Symptoms of delirium among elderly medical inpatients with or without dementia. *The Journal of neuropsychiatry and clinical neurosciences* 14(2), 167-75

Erkinjuntti T, Sulkava R, Wikstrom J, and Autio L (1987) Short Portable Mental Status Questionnaire as a screening test for dementia and delirium among the elderly. *Journal of the American Geriatrics Society* 35(5), 412-6

Leonard Maeve, McInerney Shane, McFarland John, Condon Candice, Awan Fahad, O'Connor Margaret, Reynolds Paul, Meaney Anna Maria, Adamis Dimitrios, Dunne Colum, Cullen Walter, Trzepacz Paula T, and Meagher David J (2016) Comparison of cognitive and neuropsychiatric profiles in hospitalised elderly medical patients with delirium, dementia and comorbid delirium-dementia. *BMJ open* 6(3), e009212

Meagher David J, Leonard Maeve, Donnelly Sinead, Conroy Marion, Saunders Jean, and Trzepacz Paula T (2010) A comparison of neuropsychiatric and cognitive profiles in delirium, dementia, comorbid delirium-dementia and cognitively intact controls. *Journal of neurology, neurosurgery, and and psychiatry* 81(8), 876-81

Richardson SJ, Davis DHJ, Bellelli G, Hasemann W, Meagher D, Kreisel SH, MacLulich AMJ, Cerejeira J, and Morandi A (2017) Detecting delirium superimposed on dementia: diagnostic accuracy of a simple combined arousal and attention testing procedure.. *International psychogeriatrics* 29(10), 1585-1593

Trzepacz P T, Mittal D, Torres R, Canary K, Norton J, and Jimerson N (2001) Validation of the Delirium Rating Scale-revised-98: comparison with the delirium rating scale and the cognitive test for delirium. *The Journal of neuropsychiatry and clinical neurosciences* 13(2), 229-42

### **I.1.4 Case finding for people at high risk of dementia**

van den Dungen , P , Moll van Charante, E P, van de Ven , P M, van Marwijk , H W, van der Horst , H E, van Hout , and H P (2016) Case Finding of Mild Cognitive Impairment and Dementia and Subsequent Care; Results of a Cluster RCT in Primary Care. *PLoS ONE* 11(6), e0156958

## **I.2 Involving people living with dementia in decision about care**

### **I.2.1 Barriers and facilitators to involvement in decision making for people living with dementia**

Bisson J I, Hampton V, Rosser A, and Holm S (2009) Developing a care pathway for advance decisions and powers of attorney: qualitative study. *British Journal of Psychiatry* 194(1), 55-61

Dening K H, Greenish W, Jones L, Mandal U, and Sampson E L (2012) Barriers to providing end-of-life care for people with dementia: a whole-system qualitative study. *BMJ supportive & palliative care* 2(2), 103-7

Goodman C, Amador S, Elmore N, Machen I, and Mathie E (2013) Preferences and priorities for ongoing and end-of-life care: a qualitative study of older people with dementia resident in care homes. *International Journal of Nursing Studies* 50(12), 1639-47

Livingston G, Leavey G, Manela M, Livingston D, Rait G, Sampson E, Bavishi S, Shahriyarmolki K, and Cooper C (2010) Making decisions for people with dementia who lack capacity: qualitative study of family carers in UK. *BMJ* 341, c4184

Lord K, Livingston G, and Cooper C (2015) A systematic review of barriers and facilitators to and interventions for proxy decision-making by family carers of people with dementia. *International Psychogeriatrics* 27(8), 1301-12

Mackenzie Jenny (2006) Stigma and dementia: East European and South Asian family carers negotiating stigma in the UK. *Dementia: The International Journal of Social Research and Practice* 5(2), 233-247

Murphy J, and Oliver T (2013) The use of Talking Mats to support people with dementia and their carers to make decisions together. *Health & Social Care in the Community* 21(2), 171-80

Parveen S, Blakey H, and Oyebode J R (2017) Evaluation of a carers' information programme culturally adapted for South Asian families. *International Journal of Geriatric Psychiatry* 02, 02

Poppe M, Burleigh S, and Banerjee S (2013) Qualitative evaluation of advanced care planning in early dementia (ACP-ED). *PLoS ONE [Electronic Resource]* 8(4), e60412

Samsi K, and Manthorpe J (2013) Everyday decision-making in dementia: findings from a longitudinal interview study of people with dementia and family carers. *International Psychogeriatrics* 25(6), 949-61

## **I.3 Care planning, review and co-ordination**

### **I.3.1 Introduction**

Dawn Brooker and Isabelle Latham (2016) *Person-Centred Dementia Care, Second Edition: Making Services Better with VIPS Framework* Jessica Kingsley Publishers.

### **I.3.2 Health and social care co-ordination**

#### **I.3.2.1 Qualitative evidence**

Bunn F, Burn A M, Robinson L, Poole M, Rait G, Brayne C, Schoeman J, Norton S, and Goodman C (2017) Healthcare organisation and delivery for people with dementia and comorbidity: a qualitative study exploring the views of patients, carers and professionals. *BMJ Open* 7, e013067

Brooker Dawn, Dröes Rose-Marie, and Evans Shirley (2017) Framing outcomes of post-diagnostic psychosocial interventions in dementia: the Adaptation-Coping Model and adjusting to change. *Working with Older People: Community Care Policy & Practice* 21, 13-21

Faith Martin, and et al (2015) Qualitative evaluation of a self-management intervention for people in the early stage of dementia. *Dementia: the International Journal of Social Research and Practice* 14(4), 418-435

Gethin-Jones Stephen (2014) Familial perceptions of the impact of outcome-focused homecare with older people experiencing dementia and living alone. *Working with Older People: Community Care Policy & Practice* 18(2), 90-96

Gibson G, Timlin A, Curran S, and Wattis J (2007) The impact of location on satisfaction with dementia services amongst people with dementia and their informal carers: a comparative evaluation of a community-based and a clinic-based memory service. *International Psychogeriatrics* 19(2), 267-77

Gladman J R, Jones R G, Radford K, Walker E, and Rothera I (2007) Person-centred dementia services are feasible, but can they be sustained?. *Age & Ageing* 36(2), 171-6

Górska S, Forsyth K, Irvine L, Maciver D, Prior S, Whitehead J, Flockhart J, Fairnie J, and Reid J (2013) Service-related needs of older people with dementia: perspectives of service users and their unpaid carers. *International Psychogeriatrics* 25(7), 1107-14

Hean S, Nojeed N, and Warr J (2011) Developing an integrated Memory Assessment and Support Service for people with dementia. *Journal of Psychiatric & Mental Health Nursing* 18(1), 81-8

Iliffe Steve, Waugh Amy, Poole Marie, Bamford Claire, Brittain Katie, Chew-Graham Carolyn, Fox Chris, Katona Cornelius, Livingston Gill, Manthorpe Jill, Steen Nick, Stephens Barbara, Hogan Vanessa, and Robinson Louise (2014) The effectiveness of collaborative care for people with memory problems in primary care: results of the CAREDEM case management modelling and feasibility study. *Health Technology Assessment* 18(8), 1-148

Innes A, Szymczynska P, and Stark C (2014) Dementia diagnosis and post-diagnostic support in Scottish rural communities: experiences of people with dementia and their families. *Dementia: the International Journal of Social Research and Practice* 13(2), 233-247

- Kelly F, and Innes A (2016) Facilitating independence: the benefits of a post-diagnostic support project for people with dementia. *Dementia: the International Journal of Social Research and Practice* 15(2), 162-180
- Moore D, and Jones K (2011) Promoting self-directed support for people living with dementia: overcoming the challenges. *Social Care and Neurodisability* 2(2), 66-70
- Popham C, and Orrell M (2012) What matters for people with dementia in care homes?. *Aging and Mental Health* 16(1-2), 181-188
- Rothera I, Jones R, Harwood R, Avery A J, Fisher K, James V, Shaw I, and Waite J (2008) An evaluation of a specialist multiagency home support service for older people with dementia using qualitative methods. *International Journal of Geriatric Psychiatry* 23(1), 65-72
- Sonola, Thiel, Goodwin, Kodner (2013) Oxleas Advanced Dementia Service: supporting carers and building resilience. (URBA)
- Toms Gill R, Quinn Catherine, Anderson Daniel E, and Clare Linda (2015) Help yourself: Perspectives on self-management from people with dementia and their caregivers. *Qualitative Health Research* 25(1), 87-98
- Willis Rosalind, Chan Jenifer, Scriven Issy, Lawrence Vanessa, Matthews David, Murray Joanna, and Banerjee Sube (2011) The Croydon Memory Service: Using generic working to create efficiency, job satisfaction and satisfied customers. In: *Mental health and later life: Delivering an holistic model for practice*. New York, NY: Routledge/Taylor & Francis Group; US, p125-136

### **1.3.2.2 Quantitative evidence**

- Bass D M, Judge K S, Maslow K, Wilson N L, Morgan R O, McCarthy C A, Looman W J, Snow A L, and Kunik M E (2015) Impact of the care coordination program "partners in Dementia Care" on veterans' hospital admissions and emergency department visits. *Alzheimer's and Dementia: Translational Research and Clinical Interventions* 1(1), 13-22
- Bass D M, Judge K S, Snow A L, Wilson N L, Morgan R O, Maslow K, Randazzo R, Moyer J A, Odenheimer G L, Archambault E, Elbein R, Pirraglia P, Teasdale T A, McCarthy C A, Looman W J, and Kunik M E (2014) A controlled trial of Partners in Dementia Care: veteran outcomes after six and twelve months. *Alzheimer's Research & Therapy* 6(1), 9
- Bass D M, Judge K S, Snow A L, Wilson N L, Morgan R, Looman W J, McCarthy C A, Maslow K, Moyer J A, Randazzo R, Garcia-Maldonado M, Elbein R, Odenheimer G, and Kunik M E (2013) Caregiver outcomes of partners in dementia care: effect of a care coordination program for veterans with dementia and their family members and friends. *Journal of the American Geriatrics Society* 61(8), 1377-86
- Bass D M, Clark P A, Looman W J, McCarthy C A, and Eckert S (2003) The Cleveland Alzheimer's managed care demonstration: outcomes after 12 months of implementation. *Gerontologist* 43(1), 73-85
- Callahan C M, Boustani M A, Unverzagt F W, Austrom M G, Damush T M, Perkins A J, Fultz B A, Hui S L, Counsell S R, and Hendrie H C (2006) Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA* 295(18), 2148-57
- Chien W T, and Lee Y M (2008) A disease management program for families of persons in Hong Kong with dementia. *Psychiatric Services* 59(4), 433-6

Chien W T, and Lee I Y (2011) Randomized controlled trial of a dementia care programme for families of home-resided older people with dementia. *Journal of Advanced Nursing* 67(4), 774-87

Chodosh J, Colaiaco B A, Connor K I, Cope D W, Liu H, Ganz D A, Richman M J, Cherry D L, Blank J M, Carbone Rdel, P , Wolf S M, and Vickrey B G (2015) Dementia Care Management in an Underserved Community: The Comparative Effectiveness of Two Different Approaches. *Journal of Aging & Health* 27(5), 864-93

Chu P, Edwards J, Levin R, and Thomson J (2000) The use of clinical case management for early stage Alzheimer's patients and their families. *American-Journal-of-Alzheimer's-Disease* 15(5), 284-90

Dias A, Dewey M E, D'Souza J, Dhume R, Motghare D D, Shaji K S, Menon R, Prince M, and Patel V (2008) The effectiveness of a home care program for supporting caregivers of persons with dementia in developing countries: a randomised controlled trial from Goa, India. *PLoS ONE* 3(6), e2333

Eloniemi-Sulkava U, Notkola I L, Hentinen M, Kivela S L, Sivenius J, and Sulkava R (2001) Effects of supporting community-living demented patients and their caregivers: a randomized trial. *Journal of the American Geriatrics Society* 49(10), 1282-7

Eloniemi-Sulkava U, Saarenheimo M, Laakkonen M L, Pietila M, Savikko N, Kautiainen H, Tilvis R S, and Pitkala K H (2009) Family care as collaboration: effectiveness of a multicomponent support program for elderly couples with dementia. Randomized controlled intervention study. *Journal of the American Geriatrics Society* 57(12), 2200-8

Fortinsky R H, Kulldorff M, Kleppinger A, and Kenyon-Pesce L (2009) Dementia care consultation for family caregivers: collaborative model linking an Alzheimer's association chapter with primary care physicians. *Aging & Mental Health* 13(2), 162-70

Jansen A P, van Hout , H P, Nijpels G, Rijmen F, Droes R M, Pot A M, Schellevis F G, Stalman W A, van Marwijk , and H W (2011) Effectiveness of case management among older adults with early symptoms of dementia and their primary informal caregivers: a randomized clinical trial. *International Journal of Nursing Studies* 48(8), 933-43

Kwak J, Montgomery R J, Kosloski K, and Lang J (2011) The impact of TCARE on service recommendation, use, and caregiver well-being. *Gerontologist* 51(5), 704-13

Lam L C, Lee J S, Chung J C, Lau A, Woo J, and Kwok T C (2010) A randomized controlled trial to examine the effectiveness of case management model for community dwelling older persons with mild dementia in Hong Kong. *International Journal of Geriatric Psychiatry* 25(4), 395-402

Meeuwssen E J, Melis R J, Van Der Aa , G C, Goluke-Willemsse G A, De Leest , B J, Van Raak , F H, Scholzel-Dorenbos C J, Verheijen D C, Verhey F R, Visser M C, Wolfs C A, Adang E M, Olde Rikkert, and M G (2012) Effectiveness of dementia follow-up care by memory clinics or general practitioners: randomised controlled trial. *BMJ* 344, e3086

Miller R, Newcomer R, and Fox P (1999) Effects of the Medicare Alzheimer's Disease Demonstration on nursing home entry. *Health Services Research* 34(3), 691-714

Newcomer R, Yordi C, DuNah R, Fox P, and Wilkinson A (1999) Effects of the Medicare Alzheimer's Disease Demonstration on caregiver burden and depression. *Health Services Research* 34(3), 669-89

Samus Q M, Johnston D, Black B S, Hess E, Lyman C, Vavilikolanu A, Pollutra J, Leoutsakos J M, Gitlin L N, Rabins P V, and Lyketsos C G (2014) A multidimensional home-based care coordination intervention for elders with memory disorders: the maximizing

independence at home (MIND) pilot randomized trial. *American Journal of Geriatric Psychiatry* 22(4), 398-414

Schoenmakers Birgitte, Buntinx Frank, and DeLepeleire Jan (2010) Supporting the dementia family caregiver. *Aging & Mental Health* 14(1),

Shelton P, Schraeder C, Dworak D, Fraser C, and Sager M A (2001) Caregivers' utilization of health services: results from the Medicare Alzheimer's Disease Demonstration, Illinois site. *Journal of the American Geriatrics Society* 49(12), 1600-5

Tanner J A, Black B S, Johnston D, Hess E, Leoutsakos J M, Gitlin L N, Rabins P V, Lyketsos C G, and Samus Q M (2015) A randomized controlled trial of a community-based dementia care coordination intervention: effects of MIND at Home on caregiver outcomes. *American Journal of Geriatric Psychiatry* 23(4), 391-402

Van Mierlo , L D, Meiland F J, Van de Ven , P M, Van Hout , H P, and Droes R M (2015) Evaluation of DEM-DISC, customized e-advice on health and social support services for informal carers and case managers of people with dementia; a cluster randomized trial. *International Psychogeriatrics* 27(8), 1365-78

Vickrey B G, Mittman B S, Connor K I, Pearson M L, Della Penna, R D, Ganiats T G, Demonte R W, Jr , Chodosh J, Cui X, Vassar S, Duan N, and Lee M (2006) The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial.[Summary for patients in *Ann Intern Med.* 2006 Nov 21;145(10):131; PMID: 17116913]. *Annals of Internal Medicine* 145(10), 713-26

Xiao L D, De Bellis A, Kyriazopoulos H, Draper B, Ullah S (2016) The effect of a personalized dementia care intervention for caregivers from Australian minority groups. *Current Topics in Research.* 31(1): 57-67

### **I.3.3 Post diagnosis review for people living with dementia**

Bass D M, Clark P A, Looman W J, McCarthy C A, and Eckert S (2003) The Cleveland Alzheimer's managed care demonstration: Outcomes after 12 months of implementation. *Gerontologist* 43(1), 73-85

Crotty M, Halbert J, Rowett D, Giles L, Birks R, Williams H, and Whitehead C (2004) An outreach geriatric medication advisory service in residential aged care: A randomised controlled trial of case conferencing. *Age and Ageing* 33(6), 612-617

Fati Nourhashemi, and et al (2010) Effectiveness of a specific care plan in patients with Alzheimer's disease: cluster randomised trial (PLASA study). *British Medical Journal* 19.6.10, 1346

J Meeuwse Els, and et al (2012) Effectiveness of dementia follow-up care by memory clinics or general practitioners: randomised controlled trial. *British Medical Journal* 2.6.12, 15

Kohler L, Meinke-Franze C, Hein J, Fendrich K, Heymann R, Thyrian J R, and Hoffmann W (2014) Does an interdisciplinary network improve dementia care? Results from the IDemUck-Study. *Current Alzheimer Research* 11(6), 538-548

## **I.4 Inpatient care**

### **I.4.1 Caring for people living with dementia who are admitted to hospital**

Baldwin R, Pratt H, Goring H, Marriott A, and Roberts C (2004) Does a nurse-led mental health liaison service for older people reduce psychiatric morbidity in acute general medical wards? A randomised controlled trial. *Age & Ageing* 33(5), 472-8

Boltz M, Chippendale T, Resnick B, and Galvin J E (2015) Testing family-centered, function-focused care in hospitalized persons with dementia. *Neurodegenerative Disease Management* 5(3), 203-15

Campbell M L, and Guzman J A (2004) A proactive approach to improve end-of-life care in a medical intensive care unit for patients with terminal dementia. *Critical Care Medicine* 32(9), 1839-1843

Goldberg Se, Bradshaw Le, Kearney Fc, Russell C, Whittamore Kh, Foster Pe, Mamza J, Gladman Jr, Jones Rg, Lewis Sa, Porock D, and Harwood Rh (2013) Care in specialist medical and mental health unit compared with standard care for older people with cognitive impairment admitted to general hospital: randomised controlled trial (NIHR TEAM trial). *BMJ (Clinical research ed.)* 347, f4132

Villars H, Dupuy C, Soler P, Gardette V, Soto M E, Gillette S, Nourhashemi F, and Vellas B (2013) A follow-up intervention in severely demented patients after discharge from a special Alzheimer acute care unit: impact on early emergency room re-hospitalization rate. *International Journal of Geriatric Psychiatry* 28(11), 1131-40

## **I.5 Care setting transitions**

### **I.5.1 Introduction**

Naylor M, Keating SA. (2008). Transitional Care: moving patients from one care setting to another. *The American journal of nursing*. 108(9):58-63.

### **I.5.2 Managing the transition between different settings for people living with dementia**

Davies JD, Tremont G, Bishop DS, Fortinsky RH. (2010). A telephone-delivered psychosocial intervention improves dementia caregiver adjustment following nursing home placement. *Int J Geriatr Psychiatry*, 26: 380-387

Gaugler J, Roth DL, Haley WE, Mittelman MS. (2011). Modeling trajectories and transitions: Results from the New York University Caregiver Intervention. *Nurs Res*. 60(3 Suppl): S28-S37

Gaugler JE, Reese M, Sauld J. (2015). A pilot evaluation of psychosocial support for family caregivers of relatives with dementia in long-term care: The Residential Care Transition Module. *Research in Gerontological Nursing*. 8(4): 161-172.

McGilton KS, Rivera TM, Dawson P. (2003). Can we help persons with dementia find their way in a new environment. *Aging & Mental Health*, 7(5): 363-371.

## **I.6 Modifying risk factors for dementia progression**

### **I.6.1 Risk factors for dementia progression**

#### **I.6.1.1 Studies evaluating antidiabetic medicines**

Gold M, Alderton C, Zvartau-Hind M, et al. (2010) Rosiglitazone monotherapy in mild-to-moderate Alzheimer's disease: results from a randomized, double-blind, placebo-controlled phase III study. *Dement Geriatr Cogn Disord*. 30(2): 131-46.

Risner ME, Saunders AM, Altman JF et al. (2006) Efficacy of rosiglitazone in a genetically defined population with mild-to-moderate Alzheimer's disease. *Pharmacogenomics J*. 6(4): 246-54.

#### **I.6.1.2 Studies evaluating NSAIDs**

Aisen PS, Schafer KA, Grundman M et al. (2003) Effects of rofecoxib or naproxen vs placebo on Alzheimer disease progression: a randomized controlled trial. *JAMA*. 289(21):2819-26.

Bentham P, Gray R, Sellwood E et al. (2008) Aspirin in Alzheimer's disease (AD2000): a randomised open-label trial. *Lancet Neurol*. 7(1): 41-9.

de Jong D, Jansen R, Hoefnagels W et al. (2008) No effect of one-year treatment with indomethacin on Alzheimer's disease progression: a randomized controlled trial. *PLoS One* 3(1): e1475.

Green RC, Schneider LS, Amato DA et al. (2009) Effect of tarenflurbil on cognitive decline and activities of daily living in patients with mild Alzheimer disease: a randomized controlled trial. *JAMA*. 302(23): 2557-64

Pasqualetti P, Bonomini C, Dal Forno G et al. (2009) A randomized controlled study on effects of ibuprofen on cognitive progression of Alzheimer's disease. *Aging Clin Exp Res*. 21(2): 102-10.

Reines SA, Block GA, Morris JC et al. (2004) Rofecoxib: no effect on Alzheimer's disease in a 1-year, randomized, blinded, controlled study. *Neurology*. 62(1): 66-71

Rogers J, Kirby LC, Hempelman SR et al (1993) Clinical trial of indomethacin in Alzheimer's disease. *Neurology*. 43(8): 1609-1611

Scharf S, Mander A, Ugoni A (1999) A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology*. 53(1): 197-201.

Soininen H, West C, Robbins J, Niculescu L (2007) Long-term efficacy and safety of celecoxib in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 23(1): 8-21.

Wilcock GK, Black SE, Hendrix SB et al. (2008) Efficacy and safety of tarenflurbil in mild to moderate Alzheimer's disease: a randomised phase II trial. *Lancet Neurol*. 7(6): 483-93.

#### **I.6.1.3 Studies evaluating statins**

Feldman HH, Doody RS, Kivipelto M et al (2010) Randomized controlled trial of atorvastatin in mild to moderate Alzheimer disease: LEADe. *Neurology*. 74(12): 956-64.

Sano M, Bell KL, Galasko D et al. (2011) A randomized, double-blind, placebo-controlled trial of simvastatin to treat Alzheimer disease. *Neurology*. 77(6): 556-63.

Simons M, Schwärzler F, Lütjohann D et al. (2002) Treatment with simvastatin in normocholesterolemic patients with Alzheimer's disease: A 26-week randomized, placebo-controlled, double-blind trial. *Ann Neurol.* 52(3): 346-50.

Sparks DL, Sabbagh MN, Connor DJ et al. (2005) Atorvastatin for the treatment of mild to moderate Alzheimer disease: preliminary results. *Arch Neurol.* 62(5): 753-7.

Sparks DL, Connor DJ, Sabbagh MN et al (2006) Circulating cholesterol levels, apolipoprotein E genotype and dementia severity influence the benefit of atorvastatin treatment in Alzheimer's disease: results of the Alzheimer's Disease Cholesterol-Lowering Treatment (ADCLT) trial. *Acta Neurol Scand Suppl.* 185:3-7.

#### **I.6.1.4 Studies evaluating antihypertensive medicines**

Kume K, Hanyu H, Sakurai H, et al. (2012) Effects of telmisartan on cognition and regional cerebral blood flow in hypertensive patients with Alzheimer's disease. *Geriatr Gerontol Int.* 12(2): 207-14.

Ohrai T, Tomita N, Sato-Nakagawa T, et al. (2004) Effects of brain-penetrating ACE inhibitors on Alzheimer disease progression. *Neurology.* 63(7): 1324-1325.

Morich FJ, Bieber F, Lewis JM et al. (2012) Nimodipine in the Treatment of Probable Alzheimer's Disease. *Neurology.* 11(4): 185-195.

Pantoni L, del Ser T, Sogliani AG, et al. (2005) Efficacy and safety of nimodipine in subcortical vascular dementia: a randomized placebo-controlled trial. *Stroke.* 36(3): 619-24.

## **I.7 Cholinesterase inhibitors and memantine for dementia**

### **I.7.1 Introduction**

O'Brien JT and Thomas A (2015) Vascular dementia. *The Lancet*, 386(10004), 1698-1706.

O'Brien JT, Holmes C, Jones M, Jones R, Livingston G, McKeith I, Mittler P, Passmore P, Robinson L, Sampson EL, Taylor JP, Thomas A, Burns A (2017). Clinical practice with anti-dementia drugs: A revised (third) consensus statement from the British Association for Psychopharmacology. *Journal of Psychopharmacology*; 31(2): 147-168.

Stinton C, McKeith I, Taylor JP, Lafortune L, Mioshi E, Mak E, Cambridge V, Mason J, Thomas A, O'Brien JT (2015) Pharmacological management of lewy body dementia: a systematic review and meta-analysis. *American Journal of Psychiatry*, 172(8), 731-42.

### **I.7.2 Cholinesterase inhibitors and memantine for people living with Alzheimer's disease**

Aupperle,P.M., Coyne,A.C., 20000717, Primary vs subspecialty care: a structured follow-up of dementia patients and their caregivers, *American Journal of Geriatric PsychiatryAm.J.Geriatri.Psychiatry*, 8, 167-170, 2000

Aupperle,P.M., MacPhee,E.R., Coyne,A.C., Blume,J., Sanchez,B., 20030716, Health service utilization by Alzheimer's disease patients: a 2-year follow-up of primary versus subspecialty care, *Journal of Geriatric Psychiatry & Neurology*, 16, 15-17, 2003

Watanabe,N., Yamamura,K., Suzuki,Y., Umegaki,H., Shigeno,K., Matsushita,R., Sai,Y., Miyamoto,K., Yamada,K., 20121002, Pharmacist-based Donepezil Outpatient Consultation Service to improve medication persistence, *Patient preference & adherence*, 6, 605-611, 2012

### **I.7.3 Cholinesterase inhibitors and memantine in Alzheimer's disease**

Araki T, Wake R, Miyaoka T, Kawakami K et al (2014) The effects of combine treatment of memantine and donepezil on Alzheimer's disease patients and its relationship with cerebral blood flow in the prefrontal area (2014) *International Journal of Geriatric Psychiatry*, 29, 881-889

Choi SH, Park KW, Na DL, Han JH, Kim E-J, Shim YS, Lee J-H (2011) Tolerability and efficacy of memantine add-on therapy to rivastigmine transdermal patches in mild to moderate Alzheimer's disease; a multicentre randomised, open label, parallel-group study, *Current Medical Research and Opinion*, 27 (7), 1375-1383

Dysken MW, Sano M, Asthana S, Vertrees JE, Pallaki M et al (2014) Effect of vitamin E and memantine on functional decline in Alzheimer's disease The TEAM AD VA cooperative randomized trial, *JAMA*, 311, (1), 33-44

Grossberg GT, Manes F, Allegri RF, Gutierrez-Robledo LM, Gloger S, Xie L, Jia D, Pejovic V, Miller MT, Perhach JL, Graham SM (2013) The safety, tolerability and efficacy of once daily memantine (28mg): A multinational randomised double blind , placebo controlled trial in patients with moderate to severe Alzheimer's disease taking cholinesterase inhibitors , *CNS Drugs*, 27, 469-478

Herrmann, N., O'Regan, J., Ruthirakuhan, M., Kiss, A., Eryavec, G., Williams, E., Lanctot, K.L. A Randomized Placebo-Controlled Discontinuation Study of Cholinesterase Inhibitors in Institutionalized Patients With Moderate to Severe Alzheimer Disease. *Journal of the American Medical Directors Association*, 17:142-147, 2016.

Howard R, McShane R, Lindesay J, Ritchie C, Baldwin A, Barber R, Burns A, Dening T et al (2012) Donepezil and memantine for moderate to severe Alzheimer's disease, *New England Journal of Medicine*, 2012, 366, (10), 893-903

Porsteinsson AP, Grossberg GT, Mintzer J, Olin JT (2008) Memantine treatment in patients with mild to moderate Alzheimer's disease already receiving a cholinesterase inhibitor: A randomized double-blind placebo-controlled trial, *Current Alzheimer Research* 5, 83-89

Shao Z-Q, (2015) Comparison of the efficacy of four cholinesterase inhibitors in combination with memantine for the treatment of Alzheimer's disease

Tariot PN, Farlow MR, Grossberg GT, Graham SM, McDonald S, Gergel I (2004) Memantine treatment in patients with moderate to severe Alzheimer's disease already receiving donepezil, *JAMA*, 291 (3), 317-324

#### **I.7.4 Pharmacological management of Parkinson's disease dementia**

Aarsland, D., Laake, K., Larsen, J.P., Janvin, C., Donepezil for cognitive impairment in Parkinson's disease: a randomised controlled study, *J Neurol Neurosurg Psychiatry*, 72, 708-712, 2002

Aarsland, D., Ballard, C., Walker, Z., Bostrom, F., Alves, G., Kossakowski, K., Leroi, I., Pozo-Rodriguez, F., Minthon, L., Londos, E., 2009 08 14, Memantine in patients with Parkinson's disease dementia or dementia with Lewy bodies: a double-blind, placebo-controlled, multicentre trial, *Lancet Neurology*, 8, 613-618, 2009

Dubois B, Tolosa E, Katzenschlager R, Emre M, Lees A J, Schumann G, Pourcher E, Gray J, Thomas G, Swartz J, Hsu T, and Moline M L (2012) Donepezil in Parkinson's disease dementia: a randomized, double-blind efficacy and safety study. *Movement Disorders*, 1230-1238

Emre, M., Aarsland, D., Albanese, A., Byrne, E., Deuschl, G., De Deyn, P., Durif, F., Kulisevsky, J., van Laar, T., Lees, A., Poewe, W., Robillard, A., Rosa, M., Wolters, E., Quarg, P., Tekin, S., Lane, S., Rivastigmine for dementia associated with Parkinson's disease, *N Engl J Med*, 351, 2509-2518, 2004

Emre M, Tsolaki M, Bonuccelli U, Destee A, Tolosa E, Kutzelnigg A, Ceballos-Baumann A, Zdravkovic S, Bladstrom A, Jones R, and Study Investigators (2010) Memantine for patients with Parkinson's disease dementia or dementia with Lewy bodies: a randomised, double-blind, placebo-controlled trial. [Review]. *Lancet Neurology*, 969-977

Emre M, Poewe W, De Deyn P P, Barone P, Kulisevsky J, Pourcher E, van Laar T, Storch A, Micheli F, Burn D, Durif F, Pahwa R, Callegari F, Tenenbaum N, and Strohmaier C (2014) Long-term safety of rivastigmine in parkinson disease dementia: an open-label, randomized study. *Clinical Neuropharmacology*, 9-16

Ikeda M, Mori E, Matsuo K, Nakagawa M, and Kosaka K (2015) Donepezil for dementia with Lewy bodies: a randomized, placebo-controlled, confirmatory phase III trial. *Alzheimer's Research & Therapy*, 4-

Leroi I, Overshott R, Byrne E J, Daniel E, and Burns A (2009) Randomized controlled trial of memantine in dementia associated with Parkinson's disease. *Movement Disorders*, 1217-1221

McKeith I, Del Ser T, Spano P, Emre M, Wesnes K, Anand R, Cicin-Sain A, Ferrara R, and Spiegel R (2000) Efficacy of rivastigmine in dementia with Lewy bodies: A randomised, double-blind, placebo-controlled international study. *Lancet*. 356 (9247) (pp 2031-2036), and 2000. Date of Publication: 16 Dec 2000. 2031-2036

Mori E, Ikeda M, Kosaka K, Donepezil-DLB Study, and I 20121024 (2012) Donepezil for dementia with Lewy bodies: a randomized, placebo-controlled trial. *Annals of Neurology*, 41-52

Ravina B, Putt M, Siderowf A, Farrar J T, Gillespie M, Crawley A, Fernandez H H, Trieschmann M M, Reichwein S, and Simuni T (2005) Donepezil for dementia in Parkinson's disease: a randomised, double blind, placebo controlled, crossover study. *Journal of Neurology, Neurosurgery &, and Psychiatry*, 934-939

### **I.7.5 Cholinesterase inhibitors and memantine for types of dementia other than typical Alzheimer's disease**

Auchus A, Brashear H, Salloway S, Korczyn A, De Deyn , P , and Gassmann-Mayer C (2007) Galantamine treatment of vascular dementia: A randomized trial. *Neurology* 69(5), 448-458

Ballard C, Sauter M, Scheltens P, He Y, Barkhof F, Straaten E C, Flier W M, Hsu C, Wu S, and Lane R (2008) Efficacy, safety and tolerability of rivastigmine capsules in patients with probable vascular dementia: the VantagE study. *Current medical research and opinion* 24(9), 2561-74

Black S, Roman G C, Geldmacher D S, Salloway S, Hecker J, Burns A, Perdomo C, Kumar D, and Pratt R (2003) Efficacy and tolerability of donepezil in vascular dementia: Positive results of a 24-week, multicenter, international, randomized, placebo-controlled clinical trial. *Stroke* 34(10), 2323-2330

Boxer A L, Knopman D S, Kaufer D I, Grossman M, Onyike C, Graf-Radford N, Mendez M, Kerwin D, Lerner A, Wu C K, Koestler M, Shapira J, Sullivan K, Klepac K, Lipowski K, Ullah J, Fields S, Kramer J H, Merrilees J, Neuhaus J, Mesulam M M, and Miller B L (2013) Memantine in patients with frontotemporal lobar degeneration: a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet Neurology* 12(2), 149-56

Kertesz A, Morlog D, Light M, Blair M, Davidson W, Jesso S, and Brashear R (2008) Galantamine in frontotemporal dementia and primary progressive aphasia. *Dementia & Geriatric Cognitive Disorders* 25(2), 178-85

Krupp LB, Christodoulou C, Melville P, Scherl WF, Pai LY et al (2011) Multicenter randomized clinical trial of donepezil for memory impairment in multiple sclerosis , *Neurology*, 76, 1500-1507

Maurer M, Ortler S, Baier M, Meergans M, Scherer P (2012) Randomised multicentre trial on safety and efficacy of rivastigmine in cognitively impaired multiple sclerosis patients, *Multiple Sclerosis journal*, 19, 631-638

Mok V, Wong A, Ho S, Leung T, Lam W W, and Wong K S (2007) Rivastigmine in Chinese patients with subcortical vascular dementia. *Neuropsychiatric Disease & Treatment* 3(6), 943-8

Orgogozo J, Rigaud A S, Stoeffler A, Moebius H J, and Forette F (2002) Efficacy and safety of memantine in patients with mild to moderate vascular dementia. *Stroke, and a journal of cerebral circulation* 33, 1834-9

Roman G C, Salloway S, Black S E, Royall D R, Decarli C, Weiner M W, Moline M, Kumar D, Schindler R, and Posner H (2010) Randomized, placebo-controlled, clinical trial of donepezil in vascular dementia: differential effects by hippocampal size. *Stroke* 41(6), 1213-21

Saint Paul LP, Creveuil C, Heinzele O, De Seze J, Vermesch P et al (2016) Efficacy and safety of memantine in cognitive impairment in multiple sclerosis: randomized placebo-controlled study, *Journal of the neurological sciences*, 363, 69-76

Sesok S, Bolle N, Kobal J, Bucik V, and Vodusek D B (2014) Cognitive function in early clinical phase huntington disease after rivastigmine treatment. *Psychiatria Danubina* 26(3), 239-48

Small G, Erkinjuntti T, Kurz A, and Lilienfeld S (2003) Galantamine in the treatment of cognitive decline in patients with vascular dementia or Alzheimer's disease with cerebrovascular disease. *CNS Drugs* 17(12), 905-14

Vercelletto M, Boutoleau-Bretonnière C, Volteau C, Puel M, Auriacombe S, Sarazin M, Michel B F, Couratier P, Thomas-Antérion C, Verpillat P, Gabelle A, Golfier V, Cerato E, and Lacomblez L (2011) Memantine in behavioral variant frontotemporal dementia: negative results. *Journal of Alzheimer's disease : JAD* 23(4), 749-59

Wilcock G, Mobius H J, and Stoffler A (2002) A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (MMM500). *International Clinical Psychopharmacology* 17(6), 297-305

Wilkinson D, Doody R, Helme R, Taubman K, Mintzer J, Kertesz A, and Pratt R (2003) Donepezil in vascular dementia: A randomized, placebo-controlled study. *Neurology* 61(4), 479-486

## **I.8 Drugs that may worsen cognitive decline**

### **I.8.1 Drugs that may cause cognitive decline**

Ancelin ML, Aretero S, Portet F, Dupuy AM, Touchon J, Ritchie K (2006) Non-degenerative mild cognitive impairment in elderly people and use of anticholinergic drugs: longitudinal cohort study, *BMJ*, 332 (7539) 455-459

Boustani M, Campbell N, Munger S, Maidment I, Fox C (2008) Impact of anticholinergics on the aging brain: a review and practical application, *Aging Health*, 4 (3) 311-320

Carnahan RM, Lund BC, Perry PJ, Pollock BG, Culp KR (2006) The Anticholinergic Drug Scale as a measure of drug-related anticholinergic burden: Associations with Serum Anticholinergic Activity, *Journal of Clinical Pharmacology*, 46: 1481-1486

Ehrt U, Broich K, Larssen JP, Ballard C, Aarsland D (2010) Use of drugs with anticholinergic effect and impact on cognition in Parkinson's disease: a cohort study, *BMJ*, 81, 160-65

Han L, Agostini JV, Allore HG (2008) Cumulative Anticholinergic Exposure Is Associated with Poor Memory and Executive Function in Older Men, *Journal of the American Geriatric Society*, 56 (12) 2203-2210

Rudolph JL, Salow MJ, Angelini MC, McGlinchy RE (2008) The anticholinergic risk scale and anticholinergic adverse effects in older persons, *American Medical Association*, 168 (5) 518-513

Salahudeen MS, Duffull SB, Nishtala PS (2015) Anticholinergic burden quantified by anticholinergic risk scales and adverse outcomes in older people: a systematic review, *BMC Geriatrics*, 15(31)

Sitirronarit G, Ames D, Bush AL, Faux N, et al (2010) Effects of anticholinergic drugs on cognitive function in older Australians: Results from the AIBL study, *Dementia and Geriatric Cognitive Disorders*, 31 173-178

## **I.9 Non-pharmacological interventions for people living with dementia**

### **I.9.1 Non-pharmacological interventions for people living with dementia**

#### **I.9.1.1 Cognitive stimulation therapy**

Alves J, Alves-Costa F, Magalhaes R, et al. Cognitive stimulation for Portuguese older adults with cognitive impairment: a randomized controlled trial of efficacy, comparative duration, feasibility, and experiential relevance. *American Journal of Alzheimer's Disease & Other Dementias* 2014; 29:503-12.

Baines S, Saxby P, Ehler K. Reality orientation and reminiscence therapy: a controlled cross-over study of elderly confused people. *British Journal of Psychiatry* 1987; 151:222–31.

Baldelli MV, Pirani A, Motta M, Abati E, Mariani E, Manzi V. Effects of reality orientation therapy on elderly patients in the community. *Archives of Gerontology and Geriatrics* 1993a; 17(3):211–8.

Baldelli MV, Boiardi R, Fabbo A, Pradelli JM, Neri M. The role of reality orientation therapy in restorative care of elderly patients with dementia plus stroke in the subacute nursing home setting. *Archives of Gerontology and Geriatrics* 2002; 35 Suppl 8:15–22.

Bottino CMC, Carvalho IAM, Alvarez AM, Avila R, Zukauskas PR, Bustamante SEZ, et al. Cognitive rehabilitation combined with drug treatment in Alzheimer's disease patients: a pilot study. *Clinical Rehabilitation* 2005; 19:861–9.

Breuil V, De Rotrou J, Forette F, Tortrat D, Ganansia Ganem A, Frambourt A, et al. Cognitive stimulation of patients with dementia: Preliminary results. *International Journal of Geriatric Psychiatry* 1994; 9(3):211–7.

Buschert VC, Friese U, Teipel SJ, Schneider P, Merensky W, Rujescu D, et al. Effects of a newly developed cognitive intervention in amnesic mild cognitive impairment and mild Alzheimer's disease: a pilot study. *Journal of Alzheimer's Disease* 2011; 25:679–94.

Camargo CHF, Justus FF, Retzlaff G. The effectiveness of reality orientation in the treatment of Alzheimer's disease. *American Journal of Alzheimer's Disease & Other Dementias* 2015; 30:527-32.

Capotosto E, Belacchi C, Gardini S, Faggian S, Piras F, Mantoan V, Salvalaio E, Pradelli S, and Borella E. Cognitive stimulation therapy in the Italian context: its efficacy in cognitive and non-cognitive measures in older adults with dementia. *International journal of geriatric psychiatry* 2017; 32(3):331-340.

Chapman SB, Weiner MF, Rackley A, Hynan LS, Zientz J. Effects of cognitive-communication stimulation for Alzheimer's disease patients treated with donepezil. *Journal of Speech, Language, and Hearing Research* 2004; 47(5): 1149–63.

Coen RF, Flynn B, Rigney, E, O'Connor E, Fitzgerald L, Murray C, et al. Efficacy of a cognitive stimulation therapy programme for people with dementia. *Irish Journal of Psychological Medicine* 2011; 28(3):145–7.

Cove J, Jacobi N, Donovan H, et al. Effectiveness of weekly cognitive stimulation therapy for people with dementia and the additional impact of enhancing cognitive stimulation therapy with a carer training program. *Clinical Interventions in Aging* 2014; 9:2143-50.

- Ferrario E, Cappa G, Molaschi M, Rocco M, Fabris F. Reality orientation therapy in institutionalized elderly patients: Preliminary results. *Archives of Gerontology and Geriatrics* 1991;12 Suppl 2:139–42.
- Kim HJ, Yang Y, Oh, JG, et al. Effectiveness of a community-based multidomain cognitive intervention program in patients with Alzheimer’s disease. *Geriatrics Gerontology International* 2016; 16:191-9.
- Mapelli D, Di Rosa E, Nocita R, et al. Cognitive stimulation in patients with dementia: randomized controlled trial. *Dementia and Geriatric Cognitive Disorders* 2013; 3:263-71.
- Onder G, Zanetti, O, Giacobini E, Frisoni GB, Bartorelli L, et al. Reality orientation therapy combined with cholinesterase inhibitors in Alzheimer’s disease: randomised controlled trial. *British Journal of Psychiatry* 2005; 187:450–5.
- Orgeta V, Leung P, Yates L, et al. Individual cognitive stimulation therapy for dementia: a clinical effectiveness and cost-effectiveness pragmatic, multicentre, randomised controlled trial. *Health Technology Assessment* 2015; 19(64).
- Orrell M, Aguirre E, Spector A, et al. Maintenance cognitive stimulation therapy for dementia: single-blind, multicentre, pragmatic randomised controlled trial. *The British Journal of Psychiatry* 2014; 204:454-61.
- Paddick SM, Mkenda S, Mbowe G, Kisoli A, Gray WK, Dotchin CL, Ternent L, Ogunniyi A, Kissima J, Olakehinde O, Mushi D, and Walker RW. Cognitive stimulation therapy as a sustainable intervention for dementia in sub-Saharan Africa: feasibility and clinical efficacy using a stepped-wedge design. *International psychogeriatrics* 2017; 29(6):979-989.
- Requena C, Maestu F, Fernandez A, Ortiz T. Effects of cholinergic drugs and cognitive training on dementia: 2-year follow-up. *Dementia and Geriatric Cognitive Disorders* 2006;22:339–45.
- Spector A, Orrell M, Davies S, Woods B. Can reality orientation be rehabilitated? Development and piloting of an evidence-based programme of cognition-based therapies for people with dementia. *Neuropsychological Rehabilitation* 2001; 11(3-4):377–97.
- Spector A, Thorgrimsen L, Woods B, Royan L, Davies S, Butterworth M, et al. Efficacy of an evidence-based cognitive stimulation therapy programme for people with dementia: randomised controlled trial. *British Journal of Psychiatry* 2003; 183:248–54.
- Wallis GG, Baldwin M, Higginbotham P. Reality orientation therapy-a controlled trial. *British Journal of Medical Psychology* 1983; 56(3):271–7.
- Woods RT. Reality Orientation and Staff attention: A Controlled Study. *British Journal of Psychiatry* 1979; 134:502–7.
- Yamanaka K, Kawano Y, Noguchi D, et al. Effects of cognitive stimulation therapy Japanese version (CST-J) for people with dementia: a single-blind, controlled clinical trial. *Aging & Mental Health* 2013; 17(5):579-86.

### **1.9.1.2 Cognitive training**

- Amieva H, Robert PH, Grandoulier AS, et al. Group and individual cognitive therapies in Alzheimer’s disease: the ETNA3 randomized trial. *International Psychogeriatrics* 2015.
- Bergamaschi S, Arcara G, Calza A, et al. One-year repeated cycles of cognitive training (CT) for Alzheimer’s disease. *Aging Clin Exp Res* 2013; 25:421-6.

Cahn-Weiner DA, Malloy PF, Rebok GW, Ott BR. Results of a placebo-controlled study of memory training for mildly-impaired Alzheimer's disease patients. *Applied Neuropsychology* 2003;10:215–23.

Davis RN, Massman PJ, Doody RS. Cognitive intervention in Alzheimer disease: a randomized placebo controlled study. *Alzheimer Disease and Associated Disorders* 2001; 15(1):1–9.

de Luca R, Bramanti A, de Cola MC, et al. Cognitive training for patients with dementia living in a Sicilian nursing home: a novel web-based approach. *Neurological Sciences* 2016; 37:1685-91.

de Vreese LP, Neri M. Ecological impact of combined cognitive training programs (CTP) and drug treatment (ChE-I) in AD. *International Psychogeriatrics* 1999; 11 Suppl:S91.

Galante E, Venturini G, Fiaccadori C. Computer-based cognitive intervention for dementia: preliminary results of a randomized clinical trial. *Giornale Italiano di Medicina del Lavoro ed Ergonomia* 2007; 29(3):26–32.

Heiss W-D, Kessler J, Mielke R, Szelies B, Herholz K. Long-term effects of phosphatidylserine, pyritinol and cognitive training in Alzheimer's disease: a neuropsychological, EEG, and PET investigation. *Dementia* 1994; 5(2):88–98.

Koltai DC, Welsh-Bohmer KA, Smechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychological Rehabilitation* 2001; 11(3/4): 455–75.

Lee GY, Yip CCK, Yu ECS, et al. Evaluation of a computer-assisted errorless learning-based memory training program for patients with early Alzheimer's disease in Hong Kong: a pilot study. *Clinical Interventions in Aging* 2013; 13(8):623-33.

Loewenstein DA, Acevedo A, Czaja SJ, Duara R. Cognitive rehabilitation of mildly impaired Alzheimer's disease patients on cholinesterase inhibitors. *American Journal of Geriatric Psychiatry* 2004; 12:395–402.

Quayhagen MP, Quayhagen M, Corbeil RR, Roth PA, Rodgers JA. A dyadic remediation program for care recipients with dementia. *Nursing Research* 1995; 44(3): 153–9.

Quayhagen MP, Quayhagen M, Corbeil RR, Hendrix RC, Jackson JE, Snyder L, et al. Coping with dementia: evaluation of four nonpharmacologic interventions. *International Psychogeriatrics* 2000; 12(2):249–65.

### **1.9.1.3 Cognitive rehabilitation**

Amieva H, Robert PH, Grandoulier AS, et al. Group and individual cognitive therapies in Alzheimer's disease: the ETNA3 randomized trial. *International Psychogeriatrics* 2015.

Brunelle-Hamann L, Thivierge S, Simard M. Impact of a cognitive rehabilitation intervention on neuropsychiatric symptoms in mild to moderate Alzheimer's disease. *Neuropsychological Rehabilitation* 2015; 25(5):677-707.

Clare L, Linden DEJ, Woods R, Whitaker R, Evans SJ, Parkinson CH, et al. Goal-oriented cognitive rehabilitation for people with early-stage Alzheimer disease: a single-blind randomized controlled trial of clinical efficacy. *American Journal of Geriatric Psychiatry* 2010; 18:928–39.

Clare L, Kudlicka A, Oyebode JR, et al. Goal-oriented cognitive rehabilitation in early-stage Alzheimer's and related dementias: A multi-centre single-blind randomised controlled trial (GREAT). *Health Technology Assessment* (in press).

Seyun K. Cognitive rehabilitation for elderly people with early-stage Alzheimer's disease. *Journal of Physical Therapy Science* 2015; 27(2):543-6.

Thivierge S, Jean L, Simard M. A randomized cross-over controlled study on cognitive rehabilitation of instrumental activities of daily living in Alzheimer disease. *American Association for Geriatric Psychiatry* 2014; 22(11): 1188-99.

#### **1.9.1.4 Self-management groups**

Laakkonen M, Kautiainen H, Hölttä E, Savikko N, Tilvis R, Strandberg T, et al. Effects of Self-Management Groups for People with Dementia and Their Spouses--Randomized Controlled Trial. *J Am Geriatr Soc.* 2016;64(4):752-60.

Logsdon RG, Pike KC, McCurry SM, Hunter P, Maher J, Snyder L, et al. Early-stage memory loss support groups: outcomes from a randomized controlled clinical trial. *J Gerontol B Psychol Sci Soc Sci.* 2010;65(6):691-7.

Quinn C, Toms G, Jones C, Brand A, Edwards RT, Sanders F, et al. A pilot randomized controlled trial of a self-management group intervention for people with early-stage dementia (The SMART study). *Int Psychogeriatr.* 2016;28(5):787-800.

#### **1.9.1.5 Reminiscence therapy**

Amieva H, Robert PH, Grandoulier AS, et al. Group and individual cognitive therapies in Alzheimer's disease: the ETNA3 randomized trial. *International Psychogeriatrics* 2015.

Charlesworth G, Burnell K, Crellin N, et al. Peer support and reminiscence therapy for people with dementia and their family carers: A factorial pragmatic randomised trial. *J Neurol Neurosurg Psychiatry.* 2016;August 12:Article In Press.

Eritz H, Hadjistavropoulos T, Williams J, Kroeker K, R Martin R, M Lix L, et al. A life history intervention for individuals with dementia: a randomised controlled trial examining nursing staff empathy, perceived patient personhood and aggressive behaviours. *Ageing Soc.* 2016;36(10):2061-89.

Hsieh C, Chang C, Su S, Hsiao Y, Shih Y, Han W, et al. Reminiscence group therapy on depression and apathy in nursing home residents with mild-to-moderate dementia. *J Exp Clin Med* 2010;2(2):72-78.

Ito T, Meguro K, Akanuma K. A randomized controlled trial of the group reminiscence approach in patients with Vascular Dementia. *Dementia Geriatric Cognitive Disorders.* 2007;24:48-54.

Lopes TS, Afonso RM, Ribeiro OM. A quasi-experimental study of a reminiscence program focused on autobiographical memory in institutionalized older adults with cognitive impairment. *Arch Gerontol Geriatr.* 2016;66:183-92.

Subramaniam P, Woods B, Whitaker C. Life review and life story books for people with mild to moderate dementia: a randomised controlled trial. *Aging Ment Health.* 2014;18(3):363-75.

Tadaka E, Kanagawa K. Effects of reminiscence group in elderly people with Alzheimer's disease and vascular dementia in a community setting. *Geriatric Gerontology International.* 2007;7:167-73.

Tanaka S., Honda S., Nakano H., et al. Comparison between group and personal rehabilitation for dementia in a geriatric health service facility: single-blinded randomized controlled study. *Psychogeriatrics,* 2017;17(3), pp.177-185.

Van Bogaert P, Van Grinsven R Tolsen D. Effects of SOICos model-based individual reminiscence on older adults with mild to moderate dementia due to Alzheimer disease: A pilot study. *Journal of the American Medical Directors Association*. 2013;14:528.e9-528.e13.

Wang J. Group reminiscence therapy for cognitive and affective function of demented elderly in Taiwan. *International Journal of Geriatric Psychiatry*. 2007;22:1235-40.

Woods RT, Orrell M, Bruce E, Edwards RT, Hoare Z, Hounsome B, et al. REMCARE: Pragmatic multi-centre randomised trial of reminiscence groups for people with dementia and their family carers: Effectiveness and economic analysis. *PLOS ONE*. 2016;11(4):e0152843.

Wu LF, Koo M. Randomized controlled trial of a six-week spiritual reminiscence intervention on hope, life satisfaction, and spiritual well-being in elderly with mild and moderate dementia. *Int J Geriatr Psych*. 2015;May 11:Epub.

#### **I.9.1.6 Occupational therapy**

Graff MJL, Vernooij-Dassen MJM, Thijssen M, Dekker J, Hoefnagels WHL, OldeRikkert MGM. Effects of community occupational therapy on quality of life mood, and health status in dementia patients and their caregivers: a randomized control trial. *Journal of Gerontology*. 2007;62:1002-9.

Gitlin LN, Winter L, Burke J, Chernett N, Dennis MP, Hauck WW. Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: a randomized pilot study. *Am J Geriatr Psychiatry*. 2008;16(3):229–39.

Gitlin LN, Winter L, Dennis MP, Hodgson N, Hauck WW. A biobehavioral home-based intervention and the well-being of patients with dementia and their caregivers: the COPE randomized trial. *JAMA*. 2010;304(9):983-91.

Voigt-Radloff S, Graff M, Leonhart R, Schornstein K, Jessen F, Bohlken J. A multicentre RCT on community occupational therapy in Alzheimer's disease: 10 sessions are not better than one consultation. *BMJ Open*. 2011;doi:10.1136/bmjopen-2011-000096.

#### **I.9.1.7 Psychotherapy**

Burns A, Guthrie E, Marino-Francis F, Busby C, Morris J, Russell E, et al. Brief psychotherapy in Alzheimer's disease: randomized controlled trial. *British Journal of Psychiatry*. 2005;187:143-7.

Marshall A, Spreadbury J, Cheston R. A pilot randomised controlled trial to compare changes in quality of life for participants with early diagnosis dementia who attend a 'Living well with Dementia' group compared to waiting list control. *Aging Mental Health*. 2014;19:526-35.

Tappen RM, Williams CL. Therapeutic conversation to improve mood in nursing home residents with Alzheimer's disease. *Research in Gerontological Nursing*. 2009;4:267-75.

#### **I.9.1.8 Exercise**

Arcoverde C, Deslandes A, Moraes H, Almeida C, Araujo NB, Vasques PE, Silveira H, Laks J. Treadmill training as an augmentation treatment for Alzheimer's disease: a pilot randomized controlled study. *Arq Neuropsiquiatr*. 2014 Mar;72(3):190-6.

Bossers WJ, van der Woude LH, Boersma F, Hortobágyi T, Scherder EJ, van Heuvelen MJ. Comparison of Effect of Two Exercise Programs on Activities of Daily Living in Individuals with Dementia: A 9-Week Randomized, Controlled Trial. *J Am Geriatr Soc*. 2016 Jun;64(6):1258-66.

- Burgener SC, Yang Y, Gilbert R, Marsh-Yant S. The effects of a multimodal intervention on outcomes of persons with early-stage dementia. *Am J Alzheimers Dis Other Demen*. 2008 Aug-Sep;23(4):382-94.
- Cancela JM, Ayan C, Varela S, Seijo M. Effects of a long-term aerobic exercise intervention on institutionalized patients with dementia. *J Sci Med Sport*. 2016 Apr;19(4):293-8.
- Cheng ST, Chow PK, Song YQ, Yu EC, Lam JH. Can leisure activities slow dementia progression in nursing home residents? A cluster-randomized controlled trial. *Int Psychogeriatr*. 2014 Apr;26(4):637-43.
- Christofoletti G, Oliani MM, Gobbi S, Stella F, Bucken-Gobbi LT, Renato Canineu P. A controlled clinical trial on the effects of motor intervention on balance and cognition in institutionalized elderly patients with dementia. *Clinical Rehabilitation* 2008;22(7):618–26.
- Hwang HH, Choi YJ. The effects of the dance therapy program through rhythmic exercise on cognitive memory performance of the elderly with dementia. *Proceedings of the 21st Pan-Asian Congress of Sports and Physical Education* 2010;4:12–7.
- Kemoun G, Thibaud M, Roumagne N, Carette P, Albinet C, Toussaint L, et al. Effects of a physical training programme on cognitive function and walking efficiency in elderly persons with dementia. *Dementia and Geriatric Cognitive Disorders* 2010;29(2):109–14.
- Littbrand H, Lundin-Olsson L, Gustafson Y, Rosendahl E. The effect of a high-intensity functional exercise program on activities of daily living: a randomized controlled trial in residential care facilities. *J Am Geriatr Soc*. 2009 Oct;57(10):1741-9.
- Luttenberger K, Donath C, Uter W, Graessel E. Effects of multimodal nondrug therapy on dementia symptoms and need for care in nursing home residents with degenerative dementia: a randomized-controlled study with 6-month follow-up. *J Am Geriatr Soc*. 2012 May;60(5):830-40.
- Öhman H, Savikko N, Strandberg TE, Kautiainen H, Raivio MM, Laakkonen ML, Tilvis R, Pitkälä KH. Effects of Exercise on Cognition: The Finnish Alzheimer Disease Exercise Trial: A Randomized, Controlled Trial. *J Am Geriatr Soc*. 2016 Apr;64(4):731-8.
- Pitkälä KH, Pöysti MM, Laakkonen ML, Tilvis RS, Savikko N, Kautiainen H, Strandberg TE. Effects of the Finnish Alzheimer disease exercise trial (FINALEX): a randomized controlled trial. *JAMA Intern Med*. 2013 May 27;173(10):894-901.
- Rolland Y, Pillard F, Klapouszczak A, Reynish E, Thomas D, Andrieu S, et al. Exercise program for nursing home residents with Alzheimer's Disease: a one-year randomized, controlled trial. *Journal of the American Geriatric Society* 2007; Vol. 55, issue 2:158–65.
- Steinberg M, Leoutsakos JM, Podewils LJ, Lyketsos CG. Evaluation of a home-based exercise program in the treatment of Alzheimer's disease: the Maximizing Independence in Dementia (MIND) study. *International Journal of Geriatric Psychiatry* 2009;24(7):680–5.
- Suttanon P, Hill KD, Said CM, Williams SB, Byrne KN, LoGiudice D, Lautenschlager NT, Dodd KJ. Feasibility, safety and preliminary evidence of the effectiveness of a home-based exercise programme for older people with Alzheimer's disease: a pilot randomized controlled trial. *Clin Rehabil*. 2013 May;27(5):427-38.
- Telenius EW, Engedal K, Bergland A. Effect of a high-intensity exercise program on physical function and mental health in nursing home residents with dementia: an assessor blinded randomized controlled trial. *PLoS One*. 2015 May 14;10(5):e0126102.
- Toots A, Littbrand H, Lindelöf N, Wiklund R, Holmberg H, Nordström P, Lundin-Olsson L, Gustafson Y, Rosendahl E. Effects of a High-Intensity Functional Exercise Program on

Dependence in Activities of Daily Living and Balance in Older Adults with Dementia. *J Am Geriatr Soc.* 2016 Jan;64(1):55-64.

Van de Winckel A, Feys H, De Weerd W, Dom R. Cognitive and behavioural effects of music-based exercises in patients with dementia. *Clinical Rehabilitation* 2004;18 (3):253–60.

Venturelli M, Scarsini R, Schena F. Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *American Journal of Alzheimer's Disease & Other Dementias* 2011;26(5):381–8.

Vreugdenhil A, Cannell J, Davies A, Razay G. A community based exercise programme to improve functional ability in people with Alzheimer's disease: a randomized controlled trial. *Scandinavian Journal of Caring Sciences* 2012;26:12–9.

Yang SY, Shan CL, Qing H, Wang W, Zhu Y, Yin MM, Machado S, Yuan TF, Wu T. The Effects of Aerobic Exercise on Cognitive Function of Alzheimer's Disease Patients. *CNS Neurol Disord Drug Targets.* 2015;14(10):1292-7.

### **I.9.1.9 Nutrition**

Aisen PS, Schneider LS, Sano M. High-dose vitamin B supplementation and cognitive decline in Alzheimer disease: a randomized controlled trial. *JAMA.* 300(15), 2008.

Chen H, Liu S, Ji L, et al. Folic acid supplementation mitigates Alzheimer's disease by reducing inflammation: a randomized controlled trial. *Mediators of Inflammation.*

Connelly PJ, Prentice NP, Cousland G, et al. A randomised double-blind placebo controlled trial of folic acid supplementation of cholinesterase inhibitors in Alzheimer's disease. *International Journal of Geriatric Psychiatry.* 23:155-160, 2008.

De Sousa OLV, Amaral TF. Three-week nutritional supplementation effect on long-term nutritional status of patients with mild Alzheimer's disease. *Alzheimer Dis Assoc Disord.* 26(2), 2012.

Dong HT, Jin YG, Bai Y. 11 cases of Alzheimer's disease treated by acupuncture. *Acta Universita Tis Traditionis Medical Is Sinensis Pharmacologiaeque Shanghai* 2002;16(3): 26–8.

Dysken MW, Sano M, Asthana S, et al. Effect of vitamin E and memantine on functional decline in Alzheimer disease: The TEAM-AD VA cooperative randomized trial. *JAMA.* 2014;311(1):33-44.

Freund-Levi Y, Eriksdotter-Jonhagen M, Cederholm T, Basun H, Faxen-Irving G, Garlind A, Vedin I, Vessby B, Wahlund LO, Palmblad J. Omega-3 fatty acid treatment in 174 patients with mild to moderate Alzheimer disease: OmegaAD study: a randomized double-blind trial. (2006) *Arch Neurol* 63(10): 1402-8.

Furukawa K, Tomita N, Uematsu D, et al. Randomized double-blind placebo-controlled multicentre trial of Yokukansan for neuropsychiatric symptoms in Alzheimer's disease. *Geriatr Gerontol Int* 2017. 17:211-18.

Gu C, Shen T, An H, et al. Combined therapy of Di-Huang-Yi-Zhi with donepezil in patients with Parkinson's disease dementia. *Neroscience Letters* 606:13-17 (2015).

Heo JH, Lee ST, Chu K, et al. An open-label trial of Korean red ginseng as an adjuvant treatment for cognitive impairment in patients with Alzheimer's disease. *European Journal of Neurology* 2008, 15:365-8.

- Herrschaft H, Nacu A, Likhachev S, Sholomov I, Hoerr R, Schlaefke S. 2012. Ginkgo biloba extract EGb 761 ® in dementia with neuropsychiatric features: a randomised, placebo-controlled trial to confirm the efficacy and safety of a daily dose of 240 mg. *J Psychiatr Res.* 46: 716 – 723.
- Ihl R, Bachinskaya N, Korczyn A, Vakhapova V, Tribanek M, Hoerr R, et al . 2011a. Efficacy and safety of a once-daily formulation of Ginkgo biloba extract EGb 761 ® in dementia with neuropsychiatric features. A randomized controlled trial. *Int J Geriatr Psychiatry* 26: 1186 – 1194
- Kanowski S, Hoerr R: Ginkgo biloba extract EGb 761 in dementia: intent-to-treat analyses of a 24-week, multi-center, double-blind, placebo-controlled, randomized trial. *Pharmacopsychiatry* 2003, 36:297-303.
- Lauque S, Arnaud-Battandier F, Mansourian R, et al. Protein-energy oral supplementation in malnourished nursing-home residents. A controlled trial. *Age and Aging* 2000, 29:51-6.
- Le Bars PL, Katz MM, Berman N, Itil TM, Freedman AM, Schatzberg AF: A placebo-controlled, double-blind, randomized trial of an extract of Ginkgo biloba for dementia. North American EGb Study Group. *JAMA* 1997, 278:1327-1332.
- Liu FG, Fang YS, Gao ZX, Zhou JD, Sou ML. Double-blind controlled treatment of huperzine-A and placebo in 28 patients with Alzheimer disease. *Chinese Journal of Pharmacoepidemiology* 1995;4(4):196–8.
- Maurer K, Ihl R, Dierks T, Frolich L, Maurer K, Ihl R, et al: Clinical efficacy of Ginkgo biloba special extract EGb 761 in dementia of the Alzheimer type. *J Psychiatr Res* 1997, 31:645-655.
- Mazza M, Capuanob A, Briaa P, Mazza S. Ginkgo biloba and donepezil: a comparison in the treatment of Alzheimer's dementia in a randomized placebo-controlled double-blind study. *European Journal of Neurology* 2006, 13: 981–985
- Napryeyenko O, Borzenko I, GINDEM-NP Study Group. 2007. Ginkgo biloba special extract in dementia with neuropsychiatric features. A randomised, placebo-controlled, double-blind clinical trial. *Arzneimittelforschung* 57: 4 – 11.
- Nikolova G, Yancheva S, Raychev I, Hoerr R, for the PLAGIN Study Group. 2013. Ginkgo biloba extract in dementia: A 22-week randomised, placebo-controlled, double-blind trial. *Bulgarian Neurology* 14: 139 – 143.
- Quinn JF, Raman R, Thomas RG, et al. Docosahexaenoic acid supplementation and cognitive decline in Alzheimer disease: a randomized trial. *JAMA.* 2010 November 3; 304(17): 1903–1911.
- Rafii MS1, Walsh S, Little JT, Behan K, Reynolds B, Ward C, Jin S, Thomas R, Aisen PS; Alzheimer's Disease Cooperative Study. A phase II trial of huperzine A in mild to moderate Alzheimer disease. *Neurology.* 2011 Apr 19;76(16):1389-94.
- Remington R, Bechtel C, Larsen D, et al. A phase II randomized clinical trial of a nutritional formulation for cognition and mood in Alzheimer's disease. *Journal of Alzheimer's Disease* 45 (2015) 395–405.
- Salas-Savado J, Torres M, Planas M, et al. Effect of oral administration of a whole formula diet on nutritional and cognitive status in patients with Alzheimer's disease. *Clinical Nutrition* 2005, 24:390-7.

Scheltens P, Kamphuis PJ, Verhey FR, Olde Rikkert MG, Wurtman RJ, Wilkinson D, Twisk JW, Kurz A. Efficacy of a medical food in mild Alzheimer's disease: A randomized, controlled trial. *Alzheimers Dement*. 2010 Jan;6(1):1-10.

Scheltens P, Twisk JW, Blesa R, Scarpini E, von Arnim CA, Bongers A, Harrison J, Swinkels SH, Stam CJ, de Waal H, Wurtman RJ, Wieggers RL, Vellas B, Kamphuis PJ. Efficacy of Souvenaid in mild Alzheimer's disease: results from a randomized, controlled trial. *J Alzheimers Dis*. 2012;31(1):225-36.

Schneider LS, DeKosky ST, Farlow MR, Tariot PN, Hoerr R, Kieser M: A randomized, double-blind, placebo-controlled trial of two doses of Ginkgo biloba extract in dementia of the Alzheimer's type. *Curr Alzheimer Res* 2005, 2:541-551.

Shah RC, Kamphuis PJ, Leurgans S, Swinkels SH, Sadowsky CH, Bongers A, Rappaport SA, Quinn JF, Wieggers RL, Scheltens P, Bennett DA. The S-Connect study: results from a randomized, controlled trial of Souvenaid in mild-to-moderate Alzheimer's disease. *Alzheimers Res Ther*. 2013 Nov 26;5(6):59.

Shinto L, Quinn J, Montine T, et al. A randomized placebo-controlled pilot trial of omega-3 fatty acids and alpha lipoic acid in Alzheimer's disease. *J Alzheimers Dis*. 2014 ; 38(1): . doi:10.3233/JAD-130722

Sun Y, Lu CJ, Chien KL, et al. Efficacy of multivitamin supplementation containing vitamins B6 and B12 and folic acid as adjunctive treatment with a cholinesterase inhibitor in Alzheimer's disease: a 26-week, randomized, double-blind, placebo-controlled study in Taiwanese patients. *Clinical Therapeutics*. 29(10), 2007.

Suominen MH, Puranen TM, Jyvakrpi SK, et al. Nutritional guidance improves nutrient intake and quality of life, and may prevent falls in aged persons with Alzheimer disease living with a spouse (NuAD trial). *J Nutr Health Agin*. 19(9), 2015.

van Dongen MC, van Rossum E, Kessels AG, Sielhorst HJ, Knipschild PG: The efficacy of ginkgo for elderly people with dementia and age-associated memory impairment: new results of a randomized clinical trial. *J Am Geriatr Soc* 2000, 48:1183-1194.

Xu SS, Gao ZX, Weng Z, Du ZM, Xu WA, Tong ZH, et al. Efficacy of tablet huperzine-A on memory, cognition and behavior in Alzheimer's disease. *International Medical Journal* 1997;4(2):127-31.

Yang CY, Lu ZP, Zheng CG. Efficacy and reliability of huperzine a in mild and moderate Alzheimer's disease. *Chinese Journal of Clinical Rehabilitation* 2003;7(31): 4258-9.

Zhang Y, Lin C, Zhang L, et al. Cognitive improvement during treatment for mild Alzheimer's disease with a Chinese herbal formula: a randomized controlled trial. *PLoS ONE* 10(6): e0130353.

Zhang ZX, Wang XD, Chen QT, Shu L, Wang JZ, Shan GL. Clinical efficacy and safety of huperzine A in treatment of mild to moderate Alzheimer disease, a placebo-controlled, double-blind, randomized trial. *Zhonghua yi xue za zhi* 2002;82(14):941-4.

Zhou BR, Xu ZQ, Kuang YF, Deng YH. Curative effect of aspirin for postponing the development of Alzheimer's disease. *Chinese Journal of Clinical Rehabilitation* 2004;8 (16):3020-1.

#### **1.9.1.10 Music therapy**

Ceccato E, Vigato G, Bonetto C, Bevilacqua A, Pizziolo P, Crociani S, Zanfretta E, Pollini L, Caneva PA, Baldin L, Frongillo C, Signorini A, Demoro S, and Barchi E (2012) STAM

- protocol in dementia: a multicenter, single-blind, randomized, and controlled trial. *American journal of Alzheimer's disease and other dementias* 27(5), 301-10
- Chu H, Yang CY, Lin Y, Ou KL, Lee TY, O'Brien AP, and Chou KR (2014) The impact of group music therapy on depression and cognition in elderly persons with dementia: a randomized controlled study. *Biological research for nursing* 16(2), 209-17
- Cooke M, Moyle W, Shum D, Harrison S, and Murfield J (2010) A randomized controlled trial exploring the effect of music on quality of life and depression in older people with dementia. *Journal of health psychology* 15(5), 765-76
- Guetin S, Portet F, Picot MC, Pommie C, Messaoudi M, Djabelkir L, Olsen AL, Cano MM, Lecourt E, and Touchon J (2009) Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study. *Dementia and geriatric cognitive disorders* 28(1), 36-46
- Hong IS, and Choi MJ (2011) Songwriting oriented activities improve the cognitive functions of the aged with dementia. *Arts Psychother* 38(4), 221-28
- Lin Y, Chu H, Yang CY, Chen CH, Chen SG, Chang HJ, Hsieh CJ, and Chou KR (2011) Effectiveness of group music intervention against agitated behavior in elderly persons with dementia. *International journal of geriatric psychiatry* 26(7), 670-8
- Narme P, Clement S, Ehrle N, Schiaratura L, Vachez S, Courtaigne B, Munsch F, and Samson S (2014) Efficacy of musical interventions in dementia: evidence from a randomized controlled trial. *Journal of Alzheimer's disease: JAD* 38(2), 359-69
- Raglio A, Bellandi D, Baiardi P, Gianotti M, Ubezio MC, Zancacchi E, Granieri E, Imbriani M, and Stramba-Badiale M (2015) Effect of Active Music Therapy and Individualized Listening to Music on Dementia: A Multicenter Randomized Controlled Trial. *Journal of the American Geriatrics Society* 63(8), 1534-9
- Remington R (2002) Calming music and hand massage with agitated elderly. *Nursing research* 51(5), 317-23
- Ridder HM, Stige B, Qvale LG, and Gold C (2013) Individual music therapy for agitation in dementia: an exploratory randomized controlled trial. *Aging & mental health* 17(6), 667-78
- Sakamoto M, Ando H, and Tsutou A (2013) Comparing the effects of different individualized music interventions for elderly individuals with severe dementia. *International psychogeriatrics* 25(5), 775-84
- Sarkamo T, Laitinen S, Numminen A, Kurki M, Johnson JK, and Rantanen P (2016) Clinical and Demographic Factors Associated with the Cognitive and Emotional Efficacy of Regular Musical Activities in Dementia. *Journal of Alzheimer's disease: JAD* 49(3), 767-81
- Sung HC, Chang AM, and Abbey J (2006). The effects of preferred music on agitation of older people with dementia in Taiwan. *International journal of geriatric psychiatry* 21(10), 999-1000
- Sung HC, Lee WL, Li TL, and Watson R (2012) A group music intervention using percussion instruments with familiar music to reduce anxiety and agitation of institutionalized older adults with dementia. *International journal of geriatric psychiatry* 27(6), 621-7
- Thornley J, Hirjee H, and Vasudev A (2016) Music therapy in patients with dementia and behavioral disturbance on an inpatient psychiatry unit: results from a pilot randomized controlled study. *International psychogeriatrics* 28(5), 869-71

Van de Winckel A, Feys H, De Weerd W, and Dom R (2004) Cognitive and behavioural effects of music-based exercises in patients with dementia. *Clinical rehabilitation* 18(3), 253-60

#### **I.9.1.11 Aromatherapy**

Ballard CG, O'Brien J, Reichelt K, et al. Aromatherapy as a safe and effective treatment for the management of agitation in severe dementia: the results of a double-blind, placebo-controlled trial with Melissa. *J Clin Psychiatry* 2002;53:553-8.

Burns A, Perry E, Holmes C, Francis P, Morris J, Howes MJ, Chazot P, Lees G, and Ballard C (2011) A double-blind placebo-controlled randomized trial of Melissa officinalis oil and donepezil for the treatment of agitation in Alzheimer's disease. *Dementia and geriatric cognitive disorders* 31(2), 158-64.

Yang YP, Wang CJ, and Wang JJ (2016) Effect of aromatherapy massage on agitation and depressive mood in individuals with dementia. *Journal of gerontological nursing* 42(9), 38-46.

#### **I.9.1.12 Light therapy**

Burns A, Allen H, Tomenson B, Duignan D, and Byrne J (2009) Bright light therapy for agitation in dementia: a randomized controlled trial. *International psychogeriatrics* 21(4), 711-21.

Graf A, Wallner C, Schubert V, Willeit M, Wik W, Fischer P, Kasper S, and Neumeister A (2001) The effects of light therapy on mini-mental state examination scores in demented patients. *Biological psychiatry* 50(9), 725-7.

Onega LL, Pierce TW, and Epperly L (2016) Effect of Bright Light Exposure on Depression and Agitation in Older Adults with Dementia. *Issues in mental health nursing* 37(9), 660-667.

#### **I.9.1.13 Non-invasive brain stimulation**

Ahmed MA, Darwish ES, Khedr EM, El Serogy YM, and Ali AM (2012) Effects of low versus high frequencies of repetitive transcranial magnetic stimulation on cognitive function and cortical excitability in Alzheimer's dementia. *Journal of neurology* 259(1), 83-92

Andre S, Heinrich S, Kayser F, Menzler K, Kesselring J, Khader PH, Lefaucheur JP, and Mylius V (2016) At-home tDCS of the left dorsolateral prefrontal cortex improves visual short-term memory in mild vascular dementia. *Journal of the neurological sciences* 369, 185-90

Cotelli M, Calabria M, Manenti R, Rosini S, Zanetti O, Cappa SF, and Miniussi C (2011) Improved language performance in Alzheimer disease following brain stimulation. *Journal of neurology, neurosurgery, and and psychiatry* 82(7), 794-7

Cotelli M, Manenti R, Brambilla M, Petesi M, Rosini S, Ferrari C, Zanetti O, and Miniussi C (2014) Anodal tDCS during face-name associations memory training in Alzheimer's patients. *Frontiers in aging neuroscience* 6, 38

Lee J, Choi BH, Oh E, Sohn EH, and Lee AY (2016) Treatment of Alzheimer's Disease with Repetitive Transcranial Magnetic Stimulation Combined with Cognitive Training: A Prospective, Randomized, Double-Blind, Placebo-Controlled Study. *Journal of clinical neurology (Seoul, and Korea)* 12(1), 57-64

Rabey JM, Dobronevsky E, Aichenbaum S, Gonen O, Marton RG, and Khaigrekht M (2013) Repetitive transcranial magnetic stimulation combined with cognitive training is a safe and effective modality for the treatment of Alzheimer's disease: a randomized, double-blind study. *Journal of neural transmission (Vienna, and Austria: 1996)* 120(5), 813-9

#### **I.9.1.14 Acupuncture**

Liu Q, Wang XJ, Zhang ZC, Xue R, Li P, and Li B (2016) Neuroprotection against vascular dementia after acupuncture combined with donepezil hydrochloride: P300 event related potential. *Neural regeneration research* 11(3), 460-4.

Wang Y, Qin WG, and Yu CD (2014) Clinical observation on effect of cranial suture acupuncture combined with donepezil hydrochloride tablets for Alzheimer's disease. *WJAM* 24(2), 19-24.

#### **I.9.1.15 Assistive technology**

Van der Roest HG, Wenborn J, Pastink C, et al (2017) Assistive technology for memory support in dementia. *Cochrane Database of Systematic Reviews*. DOI: 10.1002/14651858.CD009627.pub2

#### **I.9.1.16 Animal assisted therapy**

Olsen, C., Pedersen, I., Bergland, A., et al. Effect of animal-assisted interventions on depression, agitation and quality of life in nursing home residents suffering from cognitive impairment or dementia: a cluster randomized controlled trial. *International journal of geriatric psychiatry* 2016; 31(12), pp.1312-1321.

#### **I.9.1.17 Robotic pet therapy**

Petersen, S., Houston, S., Qin, H., et al. The Utilization of Robotic Pets in Dementia Care. *Journal of Alzheimer's Disease* 2017; 55(2), pp.569-574.

#### **I.9.1.18 Adapted mindfulness program**

Churcher Clarke, A., Chan, JMY., Stott, J., et al. An adapted mindfulness intervention for people with dementia in care homes: feasibility pilot study. *Int J Geriatr Psychiatry* 2017; DOI: 10.1002/gps.4669

### **I.9.2 Pre, peri and post-diagnostic counselling and support for people living with dementia and their families**

Koivisto Am, Hallikainen I, Välimäki T et al. (2016) Early psychosocial intervention does not delay institutionalization in persons with mild Alzheimer disease and has impact on neither disease progression nor caregivers' well-being: ALSOVA 3-year follow-up. *International Journal of Geriatric Psychiatry* 31(3):273-283

Laakkonen M L, Kautiainen H, Holttä E, et al. (2016) Effects of Self-Management Groups for People with Dementia and Their Spouses--Randomized Controlled Trial. *Journal of the American Geriatrics Society* 64(4):752-60

Phung Kieu T. T, Waldorff F B, Buss D V, et al. (2013) A three-year follow-up on the efficacy of psychosocial interventions for patients with mild dementia and their caregivers: the multicentre, rater-blinded, randomised Danish Alzheimer Intervention Study (DAISY). *BMJ Open* 3(11):e003584

Waldorff FB, Buss DV, Eckermann A, et al. (2012) Efficacy of psychosocial intervention in patients with mild Alzheimer's disease: the multicentre, rater blinded, randomised Danish Alzheimer Intervention Study (DAISY). *BMJ*;345:e4693.

## **I.10 Managing non-cognitive symptoms**

### **I.10.1 Interventions for treating illness emergent non-cognitive symptoms in people living with dementia**

#### **I.10.1.1 Anxiety, depression, antidepressants and antipsychotics**

##### **Systematic reviews**

Ing-Randolph AR, Phillips LR, Williams AB (2014) Group music interventions for dementia-associated anxiety: a -systematic review, *International Journal of Nursing Studies*, 37:1775-84

Leong C (2014) Antidepressants for depression in patients with dementia: a review of the literature, *American Society of Consultant Pharmacists*, 29(4):254-63

Ma H, Huang Y, Cong Z et al (2014) The efficacy and safety of atypical antipsychotics for the treatment of dementia: a meta-analysis of randomized placebo-controlled trials, *Journal of Alzheimer's Disease*, 42(3):915-37

Moulton CD, Hopkins CWP, Bevan-Jones WR (2014) Systematic review of pharmacological treatments for depressive symptoms in Huntington's disease, *Movement disorders*, 29(12):1556-61

Orgeta V, Qazi A, Spector A et al (2015) Psychological treatments for depression and anxiety in dementia and mild cognitive impairment: systematic review and meta-analysis, *British Journal of Psychiatry*, 207:293-8

Pan YJ, Wu CS, Gau SSF et al (2014) Antipsychotic discontinuation in patients with dementia: a systematic review and meta-analysis of published randomized controlled studies, *Dementia and Geriatric Cognitive Disorders*, 37:125-40

Petrovsky D, Cacchione PZ, George M (2015) Review of the effect of music interventions on symptoms of anxiety and depression in older adults with mild dementia, *American Society of Consultant Pharmacists*, 29(4):254-63

Richter T, Meyer G, Möhler R et al (2012) Psychosocial interventions for reducing antipsychotic medication in care home residents, *Cochrane Database of Systematic Reviews*, 12:CD008634

Schneider LS, Dagerman KS, Higgins JPT, et al (2011) Lack of evidence for the efficacy of memantine in mild Alzheimer's disease, *Archives of Neurology*, 68(8):991-8

Seitz DP, Adunuri N, Gill SS et al (2011) Antidepressants for agitation and psychosis in dementia, *Cochrane Database of Systematic Reviews*, 2:CD008191

##### **Randomised controlled trials**

Ballard C, Lana MM, Theodoulou M et al (2008) A randomised, blinded, placebo-controlled trial in dementia patients continuing or stopping neuroleptics, *Plos Medicine*, 5(4):587-99

Ballard C, Hanney ML, Theodoulou M et al (2009) The dementia antipsychotic withdrawal trial (DART-AD): long-term follow-up of a randomised placebo-controlled trial, *lancet Neurology*, 8:151-57

Ballard C, Thomas A, Gerry S et al (2015) A double-blind randomized placebo-controlled withdrawal trial comparing memantine and antipsychotics for the long-term treatment of

function and neuropsychiatric symptoms in people with Alzheimer's disease (MAIN-AD), *JAMDA*, 16(4):316-22

Banerjee S, Hellier J, Dewey M et al (2011) Sertraline or mirtazapine for depression in dementia (HTA-SADD): a randomised, multicentre double-blind placebo-controlled trial, *Lancet*, 378:403-11

Boström G, Conradsson M, Hörnsten C et al (2015) Effects of a high-intensity functional exercise program on depressive symptoms among people with dementia in residential care: a randomised controlled trial, *International Journal of Geriatric Psychiatry*

Brodaty H, Draper BM, Millar J et al (2003) Randomised controlled trial of different models of care for nursing home residents with dementia complicated by depression or psychosis, *Journal of Health Psychology*, 15(5):765-76

Cooke M, Moyle W, Shum D et al (2010) A randomised controlled trial exploring the effect of music on quality of life and depression in older people with dementia, *Journal of Health Psychology*, 15(5):765-76

Fossey J, Ballard C, Juszczak E et al (2006) Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial, *BMJ Online*

Hickman SE, Barrick AL, Williams CS et al (2007) The effect of ambient bright light therapy on depressive symptoms in persons with dementia, *Journal of the American Geriatrics Society*, 55:1817-24

Holmes C, Wilkinson D, Dean C et al (2007) Risperidone and rivastigmine and agitated behaviour in severe Alzheimer's disease: a randomised double blind placebo controlled trial, *International Journal of Geriatric Psychiatry*, 22:380-1

Kiosses DN, Ravdin LD, Gross JJ et al (2015) Problem adaptation therapy (PATH) for older adults with major depression and cognitive impairment: a randomised clinical trial, *JAMA Psychiatry*, 72(1):22-30

Leontjevas R, Gerritsen DL, Smalbrugge M et al (2013) A structural multidisciplinary approach to depression management in nursing-home residents: a multicentre, stepped-wedge cluster-randomised trial, *Lancet*, 381:2255-64

Lyketos CG, Del Campo LM, Steinberg M et al (2003) Treating depression in Alzheimer's disease. Efficacy and safety of sertraline therapy and the benefits of depression reduction: the DIADS, *Archives of General Psychiatry*, 60:737-46

Porsteinsson AP, Drye LT, Pollock BG et al (2014) Effect of citalopram on agitation in Alzheimer's disease, *JAMA*, 311(7):682-91

Sung HC, Chang AM, Lee WL (2010) A preferred music listening intervention to reduce anxiety in older adults with dementia in nursing homes, *Journal of Clinical Nursing*, 19:1056-64

Verhey FRJ, Verkaaik M, Lousberg R, et al (2006) Olanzapine versus haloperidol in the treatment of agitation in elderly patients with dementia: results of a randomized controlled double-blind trial, *Dementia and Geriatric Cognitive Disorders*, 21:1-8

Weintraub D, Rosenberg PB, Drye LT et al (2010) Sertraline for the treatment of depression in Alzheimer's disease: week-24 outcomes, *American Journal of Geriatric Psychiatry*, 18(4):332-40

### I.10.1.2 Sleep problems

#### Systematic reviews

Forbes D, Culum I, Lischka AR, Morgan DG, Peacock S, Forbes J, Forbes S. (2009) Light-therapy for managing cognitive, sleep, functional behavioural or psychiatric disturbances in dementia. *Cochrane Database of Systematic Reviews*, 4: CD003946.

McCleery J, Cohen DA, Sharpley AL (2016) Pharmacotherapies for sleep disturbances in Alzheimer's disease (Review), *Cochrane Database of Systematic Reviews*, 3:CD009178

#### Randomised controlled trials

Alessi CA, Martin JL, Webber AP, et al (2005) Randomized, controlled trial of a nonpharmacological intervention to improve abnormal sleep/wake patterns in nursing home residents. *JAGS*, 53(8):803-10

Chong MS, Ayalon L, Marler M, Loreda JS, Corey-Bloom J, Palmer BW, Liu L, Anconi-Israel S. (2006) Continuous positive airway pressure reduces subjective daytime sleepiness in patients with mild to moderate Alzheimer's disease with sleep disordered breathing. *Journal of American Geriatric Society*, 54: 777-781.

Harris M, Richards KC, Grando VT. (2012) The effects of slow-stroke back massage on minutes of night-time sleep in persons with dementia and sleep disturbances in the nursing home: A pilot study. *Journal of Holistic Nursing*, 30: 255-263.

Larsson V, Aarsland D, Ballard C, Mintho L, Londos E (2010) The effect of memantine on sleep behaviour in dementia with Lewy bodies and Parkinson's disease dementia. *Int. J. Geriatr. Psychiatry*, 25:1030-1038

McCurry SM, Gibbons LE, Logsdon RG, et al (2005) Night-time insomnia treatment and education for Alzheimer's disease: a randomized, controlled trial. *JAGS*, 53:609-18

McCurry SM, Gibbons LE, Logsdon RG, et al (2011) Increasing walking and bright light exposure to improve sleep in community-dwelling persons with Alzheimer's disease: results of a randomized, controlled trial. *JAGS*, 59(8):1393-1402

Richards KC, Beck C, O'Sullivan PS, Shue VM. (2005) Effect of Individualised social activity on sleep in nursing home residents with dementia. *Journal American Geriatric Society*, 53: 1510-1517.

### I.10.1.3 Agitation, aggression and apathy

#### Systematic reviews

Brown R, Howards R, Candy B, Sampson EL (2015) Opioids for agitation in dementia (review), *Cochrane database of Systematic Reviews*, 5: Art. No, CD009705

Forrester LT, Maayan N, Orrell M, Spector AE, Buchan LD, Soares-Weiser K (2014) Aromatherapy for dementia, *Cochrane database of systematic reviews*, 2: CD003150

Jutkowitz E, Brasure M, Fuchs E, Shippee T et al (2016) Care delivery interventions to manage agitation and aggression in dementia nursing home and assisted living residents: A systematic review and meta analysis, *JAGS*, 64: 477-488

Kong, EH, Evans, L, Guevara, J (2009) Nonpharmacological intervention for agitation in dementia: a systematic review and meta-analysis, *Aging and Mental Health*, 13: 512-520

Von Gunten A, Schlaefe S, Uberla K (2015) Efficacy of Gingko Biloba extract EGb 761 in dementia with behavioural and psychological symptoms: A systematic review, *The World Journal of Biological Psychiatry*, 2: 1-12

Xiao H, Su Y, Cao X, Sun S, Liang Z (2010) A meta analysis of mood stabilisers for Alzheimer's disease, *Journal of Science and Technology*, 5, 652-658

### **Randomised controlled trials**

Burns A, Allen H, Tomensen B, Duigan D, Byrne J (2009) Bright light therapy for agitation in dementia: a randomized controlled trial, *International Psychogeriatrics*, 21, 711-721

Cohen Mansfield J, Libin A, Marx MS (2007) Nonpharmacological treatment of agitation: A controlled trial of systematic individualised Intervention *Journal of Gerontology*, 8, 908-916

Cohen-Mansfield J, Thein K, Marx MS et al (2012), Efficacy of nonpharmacological interventions for agitation in advanced dementia: A randomised placebo controlled trial, *Focus on Alzheimers's disease and Related Disorders*, 25, 564-575

Cummings JL, Lykestos CG, Peskind ER, Porsteinsson AP et al (2015) Effect of dextromethorphan quinodine on agitation in people with alzheimer's disease dementia: a randomised controlled trial, *JAMA*, 314:1242-1254

Deudon A, Maubourguet N, Gervais X, Leone E, Brocker P et al (2009) Non pharmacological management of behavioural symptoms in nursing homes, *Journal of Geriatric Psychiatry*, 24, 1386-1395

Fox C, Crugel M, Maidment I, Austad H et al (2012) Efficacy of memantine for agitation in Alzheimer's disease: a randomised double-blind placebo controlled trial : *Plos One*, 7: e35185

Frakey LL, Dalloway S Buelow M, Mallo A (2012) A randomized, double-blind, placebo controlled trial of modafinil for the treatment of apathy in individuals for mild-to- moderate Alzheimer's disease patients with agitation and aggression, *Journal of Clinical Psychiatry*, 73: 796-801

Holmes C, Wilkinson D, Dean C, (2004) The efficacy of donepezil in the treatment of neuropsychiatric symptoms in Alzheimer's disease, *Neurology*, 2, 214-219

Howard RJ, Juszcak E, Ballard CG, Bentham P, Brown RG et al (2007) Donepezil for the treatment of agitation in Alzheimer's disease, *New England Journal of Medicine*, 357: 1382-1392

Lin Y, Chu H, Yang CY, Chen CH et al (2011) Effectiveness of group music intervention against agitated behaviour in elderly people with dementia, *International Journal of Geriatric Psychiatry* 7: 670-678

Mahlberg R, Walther S, Eichman U, Tracik F, Kunz D (2007) Effects of rivastigmine on actigraphically monitored motor activity in severe agitation related to Alzheimer's disease, *Archives of gerontology and geriatrics*, 1: 19-26

McCabe MP, Bird M, Davison TE, Mellor D et al (2015) An RCT to evaluate the utility of a clinical protocol for staff in the management of behavioural and psychological symptoms of dementia in residential aged care settings, *Aging and Mental Health*, 9, 799-807

Porsteinsson AP, Tariot PN, Erb R, Cox, C Smith E et al (2001) Placebo-controlled study of divalproex sodium for agitation in dementia, *Journal of Geriatric Psychiatry*, 9, 58-66

- Rapp MA, Mell T, Majic T, Treusch Y et al (2013) Agitation in nursing home residents with dementia (VIDEANT trial): effects of a cluster randomised controlled, guideline intervention trial, *International Journal of the American Medical Directors*, 9: 690-695
- Rea R, Carotenuto A, Traini E, Fasanaro AM, Manzo V, Amenta F (2015) Apathy treatment in Alzheimer's disease: Interim results of ASVCOMALVA trial, *Journal of Alzheimer's disease*, 48: 377-383
- Ridder, HMO, Stige B, Qvale LG, Gold C (2013) Individual music therapy for agitation in dementia: an exploratory randomised controlled trial, *Aging and Mental Health*, 17, 667-678
- Sung H, Chang SM, Lee W, Lee M (2006) The effects of group music with movement intervention on agitated behaviours of institutionalised elders with dementia in Taiwan, *Complementary Therapies in Medicine* 2: 113-119
- Van der Ploeg E, Eppingstall B, Camp CJ, Runci SJ et al (2013) A Randomised crossover trial to study the effect of personalised one to one interaction using Montessori-based activities on agitation, affect and engagement in nursing home residents with Dementia, *International Psychogeriatrics*, 25, 564-575
- van den Elsen GAH, Amir IA, Verkes R-J, Kramers C et al (2015) Tetrahydrocannabinol for neuropsychiatric symptoms in dementia: a randomised controlled trial, *American Academy of Neurology*, 84:2338-2344
- Wang LY, Shofer JB, Rohde K, Hart KL, Hoff DJ et al (2008) Prazosin for the treatment of behavioral symptoms in Alzheimer's disease patients with agitation and aggression, *American Journal of Geriatric Psychiatry*, 17: 744-751
- Yang MH, Lin LC, Wu, SC Chiu JH (2015) Comparison of the efficacy of aroma-acu-pressure and aromatherapy for the treatment of dementia associated agitation, *Bio Med Central* 15: 377-383
- Zwijssen SA, Smallbrugge M, Eefsting J et al (2014) Coming to Grips with Challenging Behaviour: A cluster randomised controlled trial on the effects of a multidisciplinary care program on challenging behaviour in dementia, *JAMDA*, 15, 531.e1 – 531.e10

## **I.11 Supporting informal carers**

### **I.11.1 Supporting informal carers of people living with dementia**

#### **I.11.1.1 Systematic reviews**

Jensen M, Agbata I N, Canavan M, and McCarthy G (2015) Effectiveness of educational interventions for informal caregivers of individuals with dementia residing in the community: systematic review and meta-analysis of randomised controlled trials. *International Journal of Geriatric Psychiatry* 30(2), 130-43

Laver K, Milte R, Dyer S, Crotty M (2016) A systematic review and meta analysis comparing carer focused and dyadic multicomponent interventions for carers of people with dementia, *Journal of Aging and Health*, 1-42

Lins S, Hayder-Beichel D, Rucker G, Motschall E, Antes G, Meyer G, and Langer G (2014) Efficacy and experiences of telephone counselling for informal carers of people with dementia. *Cochrane Database of Systematic Reviews* 9, CD009126

Maayan N, Soares-Weiser K, and Lee H (2014) Respite care for people with dementia and their carers. *Cochrane Database of Systematic Reviews* 1, CD004396

Parker D, Mills S, and Abbey J (2008) Effectiveness of interventions that assist caregivers to support people with dementia living in the community: a systematic review. *International Journal of Evidence-Based Healthcare* 6(2), 137-72

Scott Jennifer L, Dawkins Sarah, Quinn Michael G, Sanderson Kristy, Elliott Kate-Ellen J, Stirling Christine, Schuz Ben, and Robinson Andrew (2016) Caring for the carer: A systematic review of pure technology-based cognitive behavioral therapy (TB-CBT) interventions for dementia carers. *Aging & Mental Health* 20(8), 793-803

Thompson C A, Spilsbury K, Hall J, Birks Y, Barnes C, and Adamson J (2007) Systematic review of information and support interventions for caregivers of people with dementia. *BMC Geriatrics* 7, 18

Vernooij-Dassen M, Draskovic I, McCleery J, and Downs M (2011) Cognitive reframing for carers of people with dementia. *Cochrane Database of Systematic Reviews* (11), CD005318

#### **I.11.1.2 Randomised controlled trials**

Arango-Lasprilla J C, Panyavin I, Merchan E J. H, Perrin P B, Arroyo-Anllo E M, and Snipes D J (2014) Evaluation of a group cognitive-behavioral dementia caregiver intervention in Latin America. *American Journal of Alzheimer's Disease and Other Dementias* 29(6), 548-55

Au Alma, Li Sapphire, Lee Kristy, Leung Patrick, Pan Pey-Chyou, Thompson Larry, and Gallagher-Thompson Dolores (2010) Educational/Counseling Model Health Care: The Coping with Caregiving Group Program for Chinese caregivers of patients with Alzheimer's disease in Hong Kong. *Patient Education and Counseling* 78(2), 256-260

Blom M M, Zarit S H, Groot Zwaafink, R B, Cuijpers P, and Pot A M (2015) Effectiveness of an Internet intervention for family caregivers of people with dementia: results of a randomized controlled trial. *PLoS ONE [Electronic Resource]* 10(2), e0116622

Bruvik F K, Allore H G, Ranhoff A H, and Engedal K (2013) The effect of psychosocial support intervention on depression in patients with dementia and their family caregivers: an assessor-blinded randomized controlled trial. *Dementia and Geriatric Cognitive Disorders Extra* 3(1), 386-97

- Burgio L, Stevens A, Guy D, Roth DL, Haley WE (2003) Impact of two psychosocial interventions on white and African American family caregivers of individuals with Dementia, *Gerontologist*, 43, 568-579
- Charlesworth Georgina, and et al (2008) Befriending carers of people with dementia: randomised controlled trial. *British Medical Journal* (7.6.08), 1295-1297
- Cheng S T, Lau R W, Mak E P, Ng N S, and Lam L C (2014) Benefit-finding intervention for Alzheimer caregivers: conceptual framework, implementation issues, and preliminary efficacy. *Gerontologist* 54(6), 1049-58
- Chu H, Yang C Y, Liao Y H, Chang L I, Chen C H, and Lin C C (2011) The effects of a support group on dementia caregivers' burden and depression. *Journal of aging and health* 23(2), 228-41
- Connell Cathleen M, and Janevic Mary R (2009) Effects of a Telephone-Based Exercise Intervention for Dementia Caregiving Wives: A Randomized Controlled Trial. *The Journal of Applied Gerontology* 28(2), 171-194
- Cristancho-Lacroix V, Wrobel J, Cantegreil-Kallen I, Dub T, Rouquette A, and Rigaud A S (2015) A web-based psychoeducational program for informal caregivers of patients with Alzheimer's disease: a pilot randomized controlled trial. *Journal of Medical Internet Research* 17(5), e117
- Danucalov M Ad, Kozasa E H, Ribas K T, Galduroz J Cf, Garcia M C, and Verreschi I Tn (2013) A yoga and compassion meditation program reduces stress in familial caregivers of Alzheimer's disease patients. *Evidence-based complementary and alternative medicine : eCAM* 2013, 513149
- Dias A, Dewey M E, D'Souza J, Dhume R, Motghare D D, Shaji K S, Menon R, Prince M, and Patel V (2008) The effectiveness of a home care program for supporting caregivers of persons with dementia in developing countries: a randomised controlled trial from Goa, India. *PLoS ONE [Electronic Resource]* 3(6), e2333
- Ducharme F C, Levesque L L, Lachance L M, Kergoat M J, Legault A J, Beaudet L M, and Zarit S H (2011) "Learning to become a family caregiver" efficacy of an intervention program for caregivers following diagnosis of dementia in a relative. *Gerontologist* 51(4), 484-94
- Fortinsky R H, Kulldorff M, Kleppinger A, and Kenyon-Pesce L (2009) Dementia care consultation for family caregivers: collaborative model linking an Alzheimer's association chapter with primary care physicians. *Aging & Mental Health* 13(2), 162-70
- Gallagher-Thompson Dolores, Gray Heather L, Dupart Tamarra, Jimenez Daniel, and Thompson Larry W (2008) Effectiveness of Cognitive/Behavioral Small Group Intervention for Reduction of Depression and Stress in Non-Hispanic White and Hispanic/Latino Women Dementia Family Caregivers: Outcomes and Mediators of Change. *Journal of Rational-Emotive Cognitive-Behavior Therapy* 26(4), 286-303
- Gaugler J E, Reese M, and Mittelman M S (2015) Effects of the Minnesota Adaptation of the NYU Caregiver Intervention on Depressive Symptoms and Quality of Life for Adult Child Caregivers of Persons with Dementia. *American Journal of Geriatric Psychiatry* 23(11), 1179-1192
- Gaugler J E, Reese M, and Sauld J (2015) A Pilot Evaluation of Psychosocial Support for Family Caregivers of Relatives with Dementia in Long-Term Care: The Residential Care Transition Module. *Research in Gerontological Nursing* 8(4), 161-72

- Gonyea Judith G, López Luz M, and Velásquez Esther H (2016) The Effectiveness of a Culturally Sensitive Cognitive Behavioral Group Intervention for Latino Alzheimer's Caregivers. *Gerontologist* 56(2), 292-302 11p
- Hattink B, Meiland F, van der Roest H, Kevern P, Abiuso F, Bengtsson J, Giuliano A, Duca A, Sanders J, Basnett F, Nugent C, Kingston P, and Droes R M (2015) Web-Based STAR E-Learning Course Increases Empathy and Understanding in Dementia Caregivers: Results from a Randomized Controlled Trial in the Netherlands and the United Kingdom. *Journal of Medical Internet Research* 17(10), e241
- Hirano A, Suzuki Y, Kuzuya M, Onishi J, Ban N, and Umegaki H (2011) Influence of regular exercise on subjective sense of burden and physical symptoms in community-dwelling caregivers of dementia patients: a randomized controlled trial. *Archives of Gerontology & Geriatrics* 53(2), e158-63
- Horvath K J, Trudeau S A, Rudolph J L, Trudeau P A, Duffy M E, and Berlowitz D (2013) Clinical trial of a home safety toolkit for Alzheimer's disease. *International journal of Alzheimer's disease* 2013, 913606
- Jansen A P, Hout H P, Nijpels G, Rijmen F, Dröes R M, Pot A M, Schellevis F G, Stalman W A, and Marwijk H W (2011) Effectiveness of case management among older adults with early symptoms of dementia and their primary informal caregivers: a randomized clinical trial. *International journal of nursing studies* 48(8), 933-43
- Kwok T, Wong B, Ip I, Chui K, Young D, and Ho F (2013) Telephone-delivered psychoeducational intervention for Hong Kong Chinese dementia caregivers: a single-blinded randomized controlled trial. *Clinical Interventions In Aging* 8, 1191-7
- Lavretsky H, Epel E S, Siddarth P, Nazarian N, Cyr N S, Khalsa D S, Lin J, Blackburn E, and Irwin M R (2013) A pilot study of yogic meditation for family dementia caregivers with depressive symptoms: effects on mental health, cognition, and telomerase activity. *International Journal of Geriatric Psychiatry* 28(1), 57-65
- Leach M J, Francis A, and Ziaian T (2015) Transcendental Meditation for the improvement of health and wellbeing in community-dwelling dementia caregivers [TRANSCENDENT]: a randomised wait-list controlled trial. *BMC Complementary & Alternative Medicine* 15, 145
- Liddle J, Smith-Conway E R, Baker R, Angwin A J, Gallois C, Copland D A, Pachana N A, Humphreys M S, Byrn G J, and Chener H J (2012) Memory and communication support strategies in dementia: Effect of a training program for informal caregivers. *International Psychogeriatrics* 24(12), 1927-1942
- Logiudice D, Waltrowicz W, Brown K, Burrows (1999) Do memory clinics improve the quality of life of carers? A randomised pilot trial, *International Journal of Geriatric Psychiatry*, 14, 626-632
- Losada A, Marquez-Gonzalez M, and Romero-Moreno R (2011) Mechanisms of action of a psychological intervention for dementia caregivers: effects of behavioral activation and modification of dysfunctional thoughts. *International Journal of Geriatric Psychiatry* 26(11), 1119-27
- Losada A, Marquez-Gonzalez M, Romero-Moreno R, Mausbach B T, Lopez J, Fernandez-Fernandez V, and Nogales-Gonzalez C (2015) Cognitive-behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for dementia family caregivers with significant depressive symptoms: Results of a randomized clinical trial. *Journal of Consulting & Clinical Psychology* 83(4), 760-72

Mittelman MS, Ferris SH, Steinberg G, Shulman E, Mackell JA, Ambinder A, Cohen J (1993) An intervention that delays institutionalization of Alzheimer's disease patients: treatment of spouse-caregivers, *The Gerontologist*, 33, 730-740

Mittelman MS, Brodaty H, Wallen AS, Burns A (2008) A 3 country randomised controlled trial of a psychosocial intervention for patients with Alzheimer's disease: Effects on carer depression, *American Journal of Geriatric Psychiatry*, 16, 893-904

Oken B S, Fonareva I, Haas M, Wahbeh H, Lane J B, Zajdel D, and Amen A (2010) Pilot controlled trial of mindfulness meditation and education for dementia caregivers. *Journal of Alternative & Complementary Medicine* 16(10), 1031-8

Prick A E, De Lange J, Twisk J, and Pot A M (2015) The effects of a multi-component dyadic intervention on the psychological distress of family caregivers providing care to people with dementia: A randomized controlled trial. *International Psychogeriatrics* 27(12), 2031-2044

Rose K M (2006) The effects of cranial electrical stimulation on sleep disturbances, depressive symptoms, and caregiving appraisal in elderly caregivers of persons with Alzheimer's disease or related dementia. , University of Virginia.

Spijker A, Wollersheim H, Teerenstra S, Graff M, Adang E, Verhey F, and Vernooij-Dassen M (2011) Systematic care for caregivers of patients with dementia: a multicenter, cluster-randomized, controlled trial. *American Journal of Geriatric Psychiatry* 19(6), 521-31

Torkamani M, McDonald L, Saez Aguayo, I, Kanios C, Katsanou M N, Madeley L, Limousin P D, Lees A J, Haritou M, and Jahanshahi M (2014) A randomized controlled pilot study to evaluate a technology platform for the assisted living of people with dementia and their carers. *Journal of Alzheimer's disease : JAD* 41(2), 515-23

Tremont G, Davis J D, Papandonatos G D, Ott B R, Fortinsky R H, and Gozalo P (2014) Psychosocial telephone intervention for dementia caregivers: a randomized, controlled trial. *Alzheimer's & Dementia* , No Pagination Specified

Ulstein I D, Sandvik L, Bruun Wyller, T, and Engedal K (2007) A one-year randomized controlled psychosocial intervention study among family carers of dementia patients - Effects on patients and carers. *Dementia and Geriatric Cognitive Disorders* 24(6), 469-475

Whitebird R R, Kreitzer M, Crain A L, Lewis B A, Hanson L R, and Enstad C J (2013) Mindfulness-based stress reduction for family caregivers: a randomized controlled trial. *Gerontologist* 53(4), 676-86

Xiao L D, Bellis A, Kyriazopoulos H, Draper B, and Ullah S (2016) The effect of a personalized dementia care intervention for caregivers from Australian minority groups. *American journal of Alzheimer's disease and other dementias* 31(1 // (UNSW) \*University of New South Wales\*), 57-67

## I.12 Staff training

### I.12.1 Staff training

Beer, C. et al. (2011). A cluster-randomised trial of staff education to improve the quality of life of people with dementia living in residential care: the DIRECT study. *PLoS ONE*, 6, e28155.

Beer, C. et al. (2010). Dementia in residential care: education intervention trial (DIRECT); protocol for a randomised controlled trial. *Trials*, 11, 63

Burgio, L. D., Stevens, A., Burgio, K. L., Roth, D. L., Paul, P. and Gerstle, M. S. (2002). Teaching and maintaining behavior management skills in the nursing home. *The Gerontologist*, 32, 487–496.

Chang, C. C. and Lin, L.-C. (2005). Effects of a feeding skills training program on nursing assistants and dementia patients. *Journal of Clinical Nursing*, 14, 1185–1192.

Chenoweth, L. et al. (2009). Caring for aged dementia care resident study (CADRES) of person-centred care, dementia-care mapping, and usual care in dementia: a cluster-randomised trial. *The Lancet Neurology*, 8, 317–325.

Chenoweth, L. et al. (2014). PerCEN: a cluster randomized controlled trial of person-centered residential care and environment for people with dementia. *International Psychogeriatrics*, 26, 1147–1160.

Clare, L. et al. (2013). AwareCare: a pilot randomized controlled trial of an awareness-based staff training intervention to improve quality of life for residents with severe dementia in long-term care settings. *International Psychogeriatrics*, 25, 128–139.

Davison, T. E., McCabe, M. P., Visser, S., Hudgson, C., Buchanan, G. and George, K. (2007). Controlled trial of dementia training with a peer support group for aged care staff. *International Journal of Geriatric Psychiatry*, 22, 868–873.

Deudon, A. et al. (2009). Non-pharmacological management of behavioural symptoms in nursing homes. *International Journal of Geriatric Psychiatry*, 24, 1386–1395.

Döpp C.M.E., Graff M.J.L., Teerenstra S., Olde Rikkert M.G.M., Nijhuis–van der Sanden M.W.G., Vernooij-Dassen M.J.F.J., (2015) Effectiveness of a training package for implementing a community-based occupational therapy program in dementia: a cluster randomized controlled trial. *Clinical Rehabilitation*, 29, 974–986.

Finnema, E. et al. (2005). The effect of integrated emotion-oriented care versus usual care on elderly persons with dementia in the nursing home and on nursing assistants: a randomized clinical trial. *International Journal of Geriatric Psychiatry*, 20, 330–343.

Fossey, J. et al. (2006). Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial. *British Medical Journal*, 332, 756–758.

Huizing, A., Hamers, J., Gulpers, M. and Berger, M. (2006). Short-term effects of an educational intervention on physical restraint use: a cluster randomized trial. *BMC Geriatrics*, 6, 17.

Leone, E. et al. (2013). Management of apathy in nursing homes using a teaching program for care staff: the STIM-EHPAD study. *International Journal of Geriatric Psychiatry*, 28, 383–392.

- Magai, C., Cohen, C. I. and Gomberg, D. (2002). Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *International Psychogeriatrics*, 14, 25–38.
- McCallion, P., Tosel, R. W., Lacey, D. and Banks, S. (1999). Educating nursing assistants to communicate more effectively with nursing home residents with dementia. *The Gerontologist*, 39, 546–558.
- Pellfolk, T. J., Gustafson, Y., Bucht, G. and Karlsson, S. (2010). Effects of a restraint minimization program on staff knowledge, attitudes, and practice: a cluster randomized trial. *Journal of the American Geriatrics Society*, 58, 62–69.
- Robison, J., Curry, L., Gruman, C., Porter, M., Henderson, C. R. J. and Pillemer, K. (2007). Partners in caregiving in a special care environment: cooperative communication between staff and families on dementia units. *The Gerontologist*, 47, 504–515.
- Sloane, P. D. et al. (2004). Effect of person-centered showering and the towel bath on bathing-associated aggression, agitation, and discomfort in nursing home residents with dementia: a randomized, controlled trial. *Journal of American Geriatric Society*, 52, 1795–1894.
- Testad I., Aasland A.M., Aarsland D. (2005) The effect of staff training on the use of restraint in dementia: a single-blind randomised controlled trial. *International Journal of Geriatric Psychiatry*, 20, 587-590.
- Testad, I., Ballard, C., Brønnick, K. and Aarsl, D. (2010). The effect of staff training on agitation and use of restraint in nursing home residents with dementia: a single-blind, randomized controlled trial. *The Journal of Clinical Psychiatry*, 71, 80–86.
- Van De Ven, G. et al. (2013). Effects of dementia-care mapping on residents and staff of care homes: a pragmatic cluster-randomised controlled trial. *PLoS One*, 8, e67325.
- van Weert, J.C.M., van Dulmen, A.M., Spreeuwenberg, P.M.M., Ribbe, M.W., Bensing, J.M. Effects of snoezelen, integrated in 24 h dementia care, on nurse-patient communication during morning care. *Patient Educ. Couns.* 2005;58:312–326.
- Verkaik, R., Francke, A. L., Van Meijel, B., Spreeuwenberg, P. M., Ribbe, M. W. and Bensing, J. M. (2011). The effects of a nursing guideline on depression in psychogeriatric nursing home residents with dementia. *International Journal of Geriatric Psychiatry*, 26, 723–732.
- Visser, S. M., McCabe, M. P., Hudgson, C., Buchanan, G., Davison, T. E. and George, K. (2008). Managing behavioural symptoms of dementia: effectiveness of staff education and peer support. *Aging and Mental Health*, 12, 47–55.
- Wenborn, J. et al. (2013). Providing activity for people with dementia in care homes: a cluster randomised controlled trial. *International Journal of Geriatric Psychiatry*, 28, 1296–1304.

## **I.13 The specific needs of younger people living with dementia**

### **I.13.1 Needs of younger people living with dementia**

Chaplin R, and Davidson I (2016) What are the experiences of people with dementia in employment? *Dementia* 15, 147-61

Clayton-Turner, and et al (2015) *Approaching an unthinkable future: understanding the support needs of people living with young onset dementia*. Wolverhampton: Dementia Pathfinders

Clemerson Gemma, Walsh Sue, and Isaac Claire (2014) Towards living well with young onset dementia: An exploration of coping from the perspective of those diagnosed. *Dementia: The International Journal of Social Research and Practice* 13(4), 451-466

Hegarty, Ackermann, and Evans (2014) Walking side by side. *Journal of Dementia Care* 22(2), 18-19

Higgins Alan (2010) The ACE approach: promoting well-being and peer support for younger people with dementia. *Journal of Mental Health Training Education and Practice* 5(3), 41-50

Johnson Ann, and Walker Denise (2008) The ladies who lunch. *Journal of Dementia Care* 16(2), 14-15

Pipon-Young Fiona, and et al (2012) I'm not all gone, I can still speak: the experiences of younger people with dementia. An action research study. *Dementia: the International Journal of Social Research and Practice* 11(5), 597-616

## **I.14 Assessing and managing comorbidities**

### **I.14.1 Assessing and treating intercurrent illness in people living with dementia**

#### **I.14.1.1 Assessing intercurrent illness**

DeWaters T, Faut-Callahan M, McCann J J, Paice J A, Fogg L, Hollinger-Smith L, Sikorski K, and Stanaitis H (2008) Comparison of self-reported pain and the PAINAD Scale in hospitalized cognitively impaired and intact older adults after hip fracture surgery. *Orthopaedic Nursing* 27, 21-28 8p

Horgas A L, Nichols A L, Schapson C A, and Vietes K (2007) Assessing pain in persons with dementia: relationships among the non-communicative patient's pain assessment instrument, self-report, and behavioral observations. *Pain Management Nursing* 8, 77-85 9p

Kato-Narita E M, Nitrini R, and Radanovic M (2011) Assessment of balance in mild and moderate stages of Alzheimer's disease: implications on falls and functional capacity. *Arquivos de Neuro-Psiquiatria* 69, 202-7

Lukas A, Barber J, Johnson P, and Gibson S (2013) Observer-rated pain assessment instruments improve both the detection of pain and the evaluation of pain intensity in people with dementia. *European Journal of Pain* 17, 1558-1568

Mosele M, Inelmen E, Toffanello E, Girardi A, Coin A, Sergi G, and Manzato E (2012) Psychometric properties of the Pain Assessment in Advanced Dementia Scale compared to self assessment of pain in elderly patients. *Dementia and Geriatric Cognitive Disorders* 34, 38-43

Sepulveda E, Franco J G, Trzepacz P T, Gaviria A M, Vinuelas E, Palma J, Ferre G, Grau I, and Vilella E (2015) Performance of the Delirium Rating Scale-Revised-98 Against Different Delirium Diagnostic Criteria in a Population With a High Prevalence of Dementia. *Psychosomatics* 56, 530-41

Van Herk , R , Van Dijk , M , Tibboel D, Baar F P. M, De Wit , R , and Duivenvoorden H J (2009) The Rotterdam Elderly Pain Observation Scale (REPOS): A new behavioral pain scale for non-communicative adults and cognitively impaired elderly persons. *Journal of Pain Management* 1, 367-378

#### **I.14.1.2 Managing intercurrent illness**

Allen J, Koziak A, Buddingh S, Liang J, Buckingham J, and Beaupre L A (2012) Rehabilitation in patients with dementia following hip fracture: a systematic review. *Physiotherapy Canada* 64, 190-201

Booth Vicky, Logan Pip, Harwood Rowan, and Hood Victoria (2015) Falls prevention interventions in older adults with cognitive impairment: A systematic review of reviews. *International Journal of Therapy & Rehabilitation* 22, 289-296 8p

Chan W C, Fai Yeung, J W, Man Wong, C S, Wa Lam, L C, Chung K F, Hay Luk, J K, Wah Lee, J S, Kin Law, and A C (2015) Efficacy of physical exercise in preventing falls in older adults with cognitive impairment: A systematic review and meta-analysis. *Journal of the American Medical Directors Association* 16, 149-154

Husebo B S, Ballard C, Fritze F, Sandvik R K, and Aarsland D (2014) Efficacy of pain treatment on mood syndrome in patients with dementia: a randomized clinical trial. *International Journal of Geriatric Psychiatry* 29, 828-36

Kolanowski A M, Fick D M, Clare L, Steis M, Boustani M, and Litaker M (2011) Pilot study of a nonpharmacological intervention for delirium superimposed on dementia. *Research in Gerontological Nursing* 4, 161-7

Oliver D, Connelly J B, Victor C R, Shaw F E, Whitehead A, Genc Y, Vanoli A, Martin F C, and Gosney M A (2007) Strategies to prevent falls and fractures in hospitals and care homes and effect of cognitive impairment: systematic review and meta-analyses. *BMJ* 334, 82

Sandvik R K, Selbaek G, Seifert R, Aarsland D, Ballard C, Corbett A, and Husebo B S (2014) Impact of a stepwise protocol for treating pain on pain intensity in nursing home patients with dementia: a cluster randomized trial. *European Journal of Pain* 18, 1490-500

Shaw F E, Bond J, Richardson D A, Dawson P, Nicholas Steen, I , McKeith I G, Anne Kenny, and R (2003) Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: Randomised controlled trial. *British Medical Journal* 326, 73-75

Stenvall M, Olofsson B, Lundstrom M, Englund U, Borssen B, Svensson O, Nyberg L, and Gustafson Y (2007) A multidisciplinary, multifactorial intervention program reduces postoperative falls and injuries after femoral neck fracture. *Osteoporosis International* 18, 167-75

Suttanon P, Hill K D, Said C M, Williams S B, Byrne K N, LoGiudice D, Lautenschlager N T, and Dodd K J (2013) Feasibility, safety and preliminary evidence of the effectiveness of a home-based exercise programme for older people with Alzheimer's disease: a pilot randomized controlled trial. *Clinical Rehabilitation* 27, 427-38

Tchalla Achille, E , Lachal Florent, Cardinaud Noelle, Saulnier Isabelle, Rialle Vincent, and Preux Pierre-Marie (2013) Preventing and managing indoor falls with home-based technologies in mild and moderate Alzheimer's disease patients: Pilot study in a community dwelling. *Dementia and Geriatric Cognitive Disorders* 36, 251-61

Toulotte C, Fabre C, Dangremont B, Lensel G, and Thevenon A (2003) Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a randomised controlled trial. *Age & Ageing* 32, 67-73

Wesson J, Clemson L, Brodaty H, Lord S, Taylor M, Gitlin L, and Close J (2013) A feasibility study and pilot randomised trial of a tailored prevention program to reduce falls in older people with mild dementia. *BMC Geriatrics* 13, 89

Fuchs-Lacelle S, Hadjistavropoulos T, and Lix L (2008) Pain assessment as intervention: a study of older adults with severe dementia. *Clinical Journal of Pain* 24, 697-707

#### **I.14.2 Management strategies for people living with dementia and co-existing physical long term conditions**

Engberg S, Sereika SM, McDowell B, Weber E, Brodak I (2002) Effectiveness of prompted voiding in treating urinary incontinence in cognitively impaired homebound adults, *Journal of wound ostomy & continence*, 5, pp 252-265

Jirovec M M, Templin T (2001) Predicting success using individualised scheduled toileting for memory-impaired elders at home, *Research in Nursing and Health*, 24, pp1-8

Kume K, Hanyu H, Sakurai, H, Takada Y (2012) Effects of telmisartan on cognition and regional cerebral blood flow in hypertensive patients with Alzheimer's disease, *Geriatrics & gerontology international* 12 1444-1586

Ostaskiewicz J, Johnston L, Roe B (2010), Timed oiding for the management of urinary incontinence in adults, *Cochrane Database of Systematic Reviews*

Plichart M, Seux ML, Caillard L et al (2013) Home based blood pressure measurement in elderly patients with cognitive impairment: comparison of agreement, *Blood pressure monitoring*, 18, 208-214

Richard E, Kuiper R, Dijkgraaf MG, van Gool MD (2009) Vascular care in patients with Alzheimer's disease with cerebrovascular lesions – a randomised clinical trial, *Journal of the American Geriatrics Society*, 5 pp 797-805

Sato T, Hanyu H, Hirao K, Kanetaka H, Sakurai H, Iwamoto T (2011) Efficacy of PPAR- $\gamma$  agonist pioglitazone in Alzheimer disease, *Neurobiology of Aging*, 32, 1626 -1633

### **I.14.3 Managing mental health conditions alongside dementia**

No relevant evidence was identified for inclusion

## **I.15 Palliative care**

### **I.15.1 Palliative care**

#### **I.15.1.1 Qualitative evidence**

Crowther J, Wilson K C, Horton S, and Lloyd-Williams M. (2013). Compassion in healthcare - lessons from a qualitative study of the end of life care of people with dementia. *Journal of the Royal Society of Medicine*, 106(12), pp.492-7.

Davies N, Maio L, Vedavanam K, Manthorpe J, Vernooij-Dassen M, Iliffe S, and Team Impact Research. (2014). Barriers to the provision of high-quality palliative care for people with dementia in England: a qualitative study of professionals' experiences. *Health & Social Care in the Community*, 22(4), pp.386-94.

Dening KH, Greenish W, Jones L, et al. Barriers to providing end-of-life care for people with dementia: a whole-system qualitative study. *BMJ Supportive & Palliative Care* (2012). doi:10.1136/bmjspcare-2011-000178

Grisaffi K, Robinson L. (2010). Timing of end-of-life care in dementia: difficulties and dilemmas for GPs. *Journal of Dementia Care*, 18(3), pp.36-39.

Lamahewa K, Mathew R, Iliffe S, Wilcock J, Manthorpe J, Sampson E L, and Davies N (2017) A qualitative study exploring the difficulties influencing decision making at the end of life for people with dementia. *Health Expectations* 22, 22

Lawrence V, Samsi K, Murray J, Harari D, Banerjee S. (2011). Dying well with dementia: qualitative examination of end-of-life care. *British Journal of Psychiatry*, 199(5), 417-22.

Moore K J, Davis S, Gola A, Harrington J, Kupeli N, Vickerstaff V, King M, Leavey G, Nazareth I, Jones L, and Sampson E L (2017) Experiences of end of life amongst family carers of people with advanced dementia: longitudinal cohort study with mixed methods. *BMC Geriatrics* 17, 135

Treloar A, Crugel M, and Adamis D. (2009). Palliative and end of life care of dementia at home is feasible and rewarding: results from the 'Hope for Home' study. *Dementia* (14713012), 8(3), pp.335-347.

#### **I.15.1.2 Quantitative evidence**

Ahronheim JC, Morrison RS, Morris J, Baskin S, Meier DE. Palliative care in advanced dementia: a randomized controlled trial and descriptive analysis. *Journal of Palliative Medicine* 2000;3(3):265-73

Hanson LC, Carey TS, Caprio AJ, Lee TJ, Ersek M, Garrett J, et al. Improving decision making for feeding options in advanced dementia: a randomized, controlled trial. *Journal of the American Geriatric Society* 2011;59(11):2009-16

Hanson LC, Zimmerman S, Song, MH, et al. Effect of the Goals of Care intervention for advanced dementia: a randomized clinical trial. *JAMA Internal Medicine* 2017;177(1):24-31

Sampson EL, Candy B, Jones L. Enteral tube feeding for older people with advanced dementia. *Cochrane Database of Systematic Reviews* 2009, Issue 2. Art. No.: CD007209