

Chronic obstructive pulmonary disease in adults

Quality standard

Published: 28 July 2011

[nice.org.uk/guidance/qs10](https://www.nice.org.uk/guidance/qs10)

Contents

Introduction	5
Why this quality standard is needed	5
How this quality standard supports delivery of outcome frameworks	6
Patient experience and safety issues	10
Coordinated services.....	10
List of quality statements.....	12
Quality statement 1: Diagnosis with spirometry	14
Quality statement.....	14
Rationale	14
Quality measures	14
What the quality statement means for service providers, healthcare professionals and commissioners...	15
What the quality statement means for patients, service users and carers	15
Source guidance.....	16
Definitions of terms used in this quality statement	16
Quality statement 2: Inhaler technique	18
Quality statement.....	18
Rationale	18
Quality measures	18
What the quality statement means for service providers, healthcare professionals and commissioners...	20
What the quality statement means for patients, service users and carers.....	20
Source guidance.....	21
Equality and diversity considerations.....	21
Quality statement 3: Assessment for long-term oxygen therapy.....	22
Quality statement.....	22
Rationale	22
Quality measures	22
What the quality statement means for service providers, healthcare professionals and commissioners ..	23

What the quality statement means for patients, service users and carers.....	23
Source guidance.....	23
Definitions.....	24
Quality statement 4: Pulmonary rehabilitation for stable COPD and exercise limitation	25
Quality statement.....	25
Rationale	25
Quality measures	25
What the quality statement means for service providers, healthcare professionals and commissioners...	27
What the quality statement means for patients, service users and carers.....	27
Source guidance.....	27
Definitions of terms used in this quality statement	27
Equality and diversity considerations.....	28
Quality statement 5: Pulmonary rehabilitation after an acute exacerbation.....	29
Quality statement.....	29
Rationale	29
Quality measures	29
What the quality statement means for service providers, healthcare professionals and commissioners...	30
What the quality statement means for patients, service users and carers.....	30
Source guidance.....	31
Definitions of terms used in this quality statement	31
Equality and diversity considerations.....	32
Quality statement 6: Emergency oxygen during an exacerbation	33
Quality statement.....	33
Rationale.....	33
Quality measures	33
What the quality statement means for service providers, healthcare professionals and commissioners...	34
What the quality statement means for patients, service users and carers	34
Source guidance.....	34

Definitions of terms used in this quality statement	35
Quality statement 7: Non-invasive ventilation	36
Quality statement.....	36
Rationale	36
Quality measures	36
What the quality statement means for service providers, healthcare professionals and commissioners ..	37
What the quality statement means for patients, service users and carers.....	37
Source guidance.....	37
Definitions of terms used in this quality statement	38
Quality statement 8 (placeholder): Hospital discharge care bundle	40
What is a placeholder statement?	40
Rationale	40
Using the quality standard.....	41
Quality measures	41
Levels of achievement	41
Using other national guidance and policy documents.....	41
Diversity, equality and language	42
Development sources.....	43
Evidence sources.....	43
Policy context	43
Definitions and data sources for the quality measures	43
Related NICE quality standards	44
Quality Standards Advisory Committee and NICE project team	45
Quality Standards Advisory Committee.....	45
NICE project team	47
Update information	48
About this quality standard.....	49

This standard is based on NG115.

This standard should be read in conjunction with QS13, QS15, QS25, QS43, QS117, QS121, QS122, QS110 and QS82.

Introduction

This quality standard covers the assessment, diagnosis and management of chronic obstructive pulmonary disease (COPD). It does not cover prevention, screening or case finding. For more information see the [COPD topic overview](#).

This quality standard has been updated. The topic was identified for update following the annual review of quality standards in 2014. The review identified that there had been changes in the areas for improvement for COPD. For further information about the update, see [update information](#). Statements from the 2011 quality standard that are no longer national priorities for improvement, but are still underpinned by current accredited guidance, are included after the updated statements in the [list of quality statements](#).

This quality standard should be considered alongside [quality statement 2](#) in the NICE quality standard on smoking: supporting people to stop, which sets out the high-quality requirements for referring people who smoke to an evidence-based smoking cessation service. Smoking is one of the main causes of COPD and encouraging people with COPD to stop smoking is one of the most important components in managing COPD.

This quality standard should also be considered alongside the NICE quality standard on [end of life care for adults](#), which sets out the high-quality requirements for adults approaching the end of their life.

Why this quality standard is needed

COPD is a long-term respiratory condition characterised by airflow obstruction that is not fully reversible. The airflow obstruction does not change markedly over several months and is usually progressive. COPD is predominantly caused by smoking. Other factors, particularly occupational exposures, such as harmful dust and chemicals, may also contribute to developing COPD. People with COPD often have exacerbations, when there is rapid and sustained worsening of symptoms beyond their usual day-to-day variation.

In the UK, it is estimated that 3 million people have COPD, of whom 2 million are undiagnosed. Prevalence increases with age and most people are not diagnosed until they are in their 50s. There are significant geographic variations in the prevalence of COPD, and it is closely associated with levels of deprivation. Unlike many other common chronic diseases, the prevalence of COPD has not declined in recent years.

There is no single diagnostic test for COPD. Making a diagnosis relies on clinical judgement based on a combination of history, physical examination and confirmation of the presence of airflow obstruction using spirometry.

COPD is treatable but not curable, and early diagnosis and treatment can help to slow the decline in lung function and increase the amount of time that people with COPD have to enjoy an active life. Pharmacological and other therapies can help to manage symptoms and disability caused by COPD, and improve the person's quality of life, despite having only limited or no impact on the airflow obstruction.

The quality standard is expected to contribute to improvements in the following outcomes:

- COPD diagnosis
- morbidity
- mortality
- acute exacerbations
- hospital admissions
- A&E attendance
- quality of life
- change in breathlessness
- exercise capacity
- inappropriate non-invasive ventilation.

How this quality standard supports delivery of outcome frameworks

NICE quality standards are a concise set of prioritised statements designed to drive measurable improvements in the 3 dimensions of quality – patient safety, patient experience and clinical

effectiveness – for a particular area of health or care. They are derived from high-quality guidance, such as that from NICE or other sources accredited by NICE. This quality standard, in conjunction with the guidance on which it is based, should contribute to the improvements outlined in the following 3 outcomes frameworks published by the Department of Health:

- [NHS Outcomes Framework 2015–16](#)
- [Public Health Outcomes Framework 2013–16](#)
- [Adult Social Care Outcomes Framework 2015–16](#)

Tables 1–3 show the outcomes, overarching indicators and improvement areas from the frameworks that the quality standard could contribute to achieving.

Table 1 [NHS Outcomes Framework 2015–16](#)

Domain	Overarching indicators and improvement areas
1 Preventing people from dying prematurely	<p><i>Improvement areas</i></p> <p>Reducing premature mortality from the major causes of death</p> <p>1.2 Under 75 mortality rate from respiratory disease*</p>
2 Enhancing quality of life for people with long-term conditions	<p><i>Overarching indicator</i></p> <p>2 Health-related quality of life for people with long-term conditions**</p> <p><i>Improvement areas</i></p> <p>Ensuring people feel supported to manage their condition</p> <p>2.1 Proportion of people feeling supported to manage their condition**</p> <p>Reducing time spent in hospital by people with long-term conditions</p> <p>2.3 i Unplanned hospitalisation for chronic ambulatory care sensitive conditions (adults)</p> <p>Improving quality of life for people with multiple long-term conditions</p> <p>2.7 Health-related quality of life for people with three or more long-term conditions**</p>

<p>3 Helping people to recover from episodes of ill health or following injury</p>	<p>Improvement areas</p> <p>Helping older people to recover their independence after illness or injury</p> <p>3.6 i Proportion of older people (65 and over) who were still at home 91 days after discharge from hospital into reablement/ rehabilitation service*</p> <p>3.6 ii Proportion offered rehabilitation following discharge from acute or community hospital*</p>
<p>4 Ensuring that people have a positive experience of care</p>	<p>Overarching indicators</p> <p>4a Patient experience of primary care</p> <p>i. GP services</p> <p>ii. GP out-of-hours services</p> <p>4b Patient experience of hospital care</p> <p>4c Friends and family test</p> <p>4d Patient experience characterised as poor or worse</p> <p>i. Primary care</p> <p>ii. Hospital care</p> <p>Improvement areas</p> <p>Improving people's experience of outpatient care</p> <p>4.1 Patient experience of outpatient services</p> <p>Improving hospitals' responsiveness to personal needs</p> <p>4.2 Responsiveness to inpatients' personal needs</p> <p>Improving the experience of care for people at the end of their lives</p> <p>4.6 Bereaved carers' views on the quality of care in the last 3 months of life</p>
<p>Alignment across the health and social care system</p> <p>* Indicator is shared</p> <p>** Indicator is complementary</p>	

Table 2 Public health outcomes framework for England 2013– 16

Domain	Objectives and indicators
--------	---------------------------

2 Health improvement	<p>Objective</p> <p>People are helped to live healthy lifestyles, make healthy choices and reduce health inequalities</p> <p>Indicators</p> <p>2.14 Smoking prevalence – adults (over 18s)</p>
4 Healthcare public health and preventing premature mortality	<p>Objective</p> <p>Reduced numbers of people living with preventable ill health and people dying prematurely, whilst reducing the gap between communities</p> <p>Indicators</p> <p>4.7 Mortality rate from respiratory diseases*</p> <p>4.11 Emergency readmissions within 30 days of discharge from hospital*</p> <p>4.13 Health-related quality of life for older people</p>
<p>Alignment across the health and social care system</p> <p>* Indicator shared with the NHS Outcomes Framework</p>	

Table 3 The Adult Social Care Outcomes Framework 2015–16

Domain	Overarching and outcome measures
2 Delaying and reducing the need for care and support	<p>Overarching measure</p> <p>2A Permanent admissions to residential and nursing care homes, per 100,000 population</p> <p>Outcome measures</p> <p>Everybody has the opportunity to have the best health and wellbeing throughout their life, and can access support and information to help them manage their care needs</p> <p>Earlier diagnosis, intervention and reablement means that people and their carers are less dependent on intensive services</p> <p>2B Proportion of older people (65 and over) who were still at home 91 days after discharge from hospital into reablement/rehabilitation services*</p>

Aligning across the health and care system

* Indicator shared

** Indicator complementary

Patient experience and safety issues

Ensuring that care is safe and that people have a positive experience of care is vital in a high-quality service. It is important to consider these factors when planning and delivering services relevant to COPD.

A patient safety alert on [Risk of severe harm and death from unintentional interruption of non-invasive ventilation](#) has been issued by NHS England to raise awareness of patient safety incidents. Non-invasive ventilation is used for some people with COPD who have acute exacerbations that do not respond to medical therapy.

NICE has developed guidance and an associated quality standard on patient experience in adult NHS services (see the NICE Pathway on [patient experience in adult NHS services](#)), which should be considered alongside this quality standard. They specify that people receiving care should be treated with dignity, have opportunities to discuss their preferences, and are supported to understand their options and make fully informed decisions. They also cover the provision of information to patients and service users. Quality statements on these aspects of patient experience are not usually included in topic-specific quality standards. However, recommendations in the development sources for quality standards that affect patient experience and are specific to the topic are considered during quality statement development.

Coordinated services

The quality standard for COPD specifies that services should be commissioned from and coordinated across all relevant agencies encompassing the whole COPD care pathway. A person-centred, integrated approach to providing services is fundamental to delivering high-quality care to adults with COPD in secondary, primary and community services.

The Health and Social Care Act 2012 sets out a clear expectation that the care system should consider NICE quality standards in planning and delivering services, as part of a general duty to secure continuous improvement in quality. Commissioners and providers of health and social care should refer to the library of NICE quality standards when designing high-quality services. Other quality standards that should also be considered when choosing, commissioning or providing a

high-quality COPD service are listed in [related quality standards](#).

Training and competencies

The quality standard should be read in the context of national and local guidelines on training and competencies. All healthcare professionals involved in assessing, caring for and treating people with COPD should have sufficient and appropriate training and competencies to deliver the actions and interventions described in the quality standard. Quality statements on staff training and competency are not usually included in quality standards. However, recommendations in the development sources on specific types of training for the topic that exceed standard professional training are considered during quality statement development.

Role of families and carers

Quality standards recognise the important role families and carers have in supporting people with COPD. If appropriate, healthcare professionals should ensure that family members and carers are involved in the decision-making process about investigations, treatment and care.

List of quality statements

Statement 1 People aged over 35 years who present with a risk factor and one or more symptoms of chronic obstructive pulmonary disease (COPD) have post-bronchodilator spirometry. [2011, updated 2016]

Statement 2 People with COPD who are prescribed an inhaler have their inhaler technique assessed when starting treatment and then regularly during treatment. [2011, updated 2016]

Statement 3 People with stable COPD and a persistent resting stable oxygen saturation level of 92% or less have their arterial blood gases measured to assess whether they need long-term oxygen therapy. [2011, updated 2016]

Statement 4 People with stable COPD and exercise limitation due to breathlessness are referred to a pulmonary rehabilitation programme. [2011, updated 2016]

Statement 5 People admitted to hospital for an acute exacerbation of COPD start a pulmonary rehabilitation programme within 4 weeks of discharge. [2011, updated 2016]

Statement 6 People receiving emergency oxygen for an acute exacerbation of COPD have oxygen saturation levels maintained between 88% and 92%. [new 2016]

Statement 7 People with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical therapy have non-invasive ventilation. [2011, updated 2016]

Statement 8 (Placeholder) Hospital discharge care bundle. [new 2016]

In 2016 this quality standard was updated and statements prioritised in 2011 were updated (2011, updated 2016) or replaced (new 2016). For more information, see [update information](#).

Statements from the 2011 quality standard for COPD that may still be useful at a local level, but are no longer considered national priorities for improvement:

- People with COPD have a current individualised comprehensive management plan, which includes high-quality information and educational material about the condition and its management, relevant to the stage of disease.

- People with COPD have a comprehensive clinical and psychosocial assessment, at least once a year or more frequently if indicated, which includes degree of breathlessness, frequency of exacerbations, validated measures of health status and prognosis, presence of hypoxaemia and comorbidities.
- People with COPD who smoke are regularly encouraged to stop and are offered the full range of evidence-based smoking cessation support.
- People who have had an exacerbation of COPD are provided with individualised written advice on early recognition of future exacerbations, management strategies (including appropriate provision of antibiotics and corticosteroids for self-treatment at home) and a named contact.
- People with COPD receiving long-term oxygen therapy are reviewed in accordance with NICE guidance, at least annually, by a specialist oxygen service as part of the integrated clinical management of their COPD.
- People admitted to hospital with an exacerbation of COPD are cared for by a respiratory team, and have access to a specialist early supported-discharge scheme with appropriate community support.
- People admitted to hospital with an exacerbation of COPD are reviewed within 2 weeks of discharge.
- People with advanced COPD, and their carers, are identified and offered palliative care that addresses physical, social and emotional needs.

The [2011 quality standard for COPD](#) is available as a pdf.

Quality statement 1: Diagnosis with spirometry

Quality statement

People aged over 35 years who present with a risk factor and one or more symptoms of chronic obstructive pulmonary disease (COPD) have post-bronchodilator spirometry. [2011, updated 2016]

Rationale

A diagnosis of COPD is confirmed by post-bronchodilator spirometry. To ensure early diagnosis, spirometry should be done in primary care when a person presents with a risk factor for COPD (which is usually smoking) and one or more symptoms of COPD.

Quality measures

Structure

a) Evidence of local arrangements and written clinical protocols to ensure that people aged over 35 years presenting with a risk factor and one or more symptoms of COPD have post-bronchodilator spirometry.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Evidence of local arrangements and written clinical protocols to ensure that healthcare professionals in primary care using post-bronchodilator spirometry are trained and competent in its use.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

c) Evidence of local arrangements to ensure that primary care services providing post-bronchodilator spirometry are supported by quality control processes.

Data source: Local data collection.

Process

Proportion of people aged over 35 years presenting with a risk factor and one or more symptoms of COPD who have post-bronchodilator spirometry.

Numerator – the number in the denominator who have post-bronchodilator spirometry.

Denominator – the number of people aged over 35 years presenting with a risk factor and one or more symptoms of COPD.

Data source: Local data collection. Quality and Outcomes Framework indicator COPD002: The percentage of patients with COPD in whom the diagnosis has been confirmed by post-bronchodilator spirometry between 3 months before and 12 months after entering on to the register.

Outcome

COPD incidence.

Data source: Local data collection. Royal College of Physicians' National COPD Audit Programme.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (primary care services) ensure that quality-assured post-bronchodilator spirometry is carried out in people aged 35 years and over who have a risk factor and one or more symptoms of COPD, to confirm diagnosis of COPD. Service providers ensure that healthcare professionals are trained and competent in performing and interpreting post-bronchodilator spirometry.

Healthcare professionals (in primary care services) ensure that they perform quality-assured post-bronchodilator spirometry in people aged 35 years and over who have a risk factor and one or more symptoms of COPD, to confirm diagnosis of COPD. Healthcare professionals ensure they remain up to date with training and competencies in performing and interpreting post-bronchodilator spirometry.

Commissioners (clinical commissioning groups) ensure that they commission services in which people aged 35 years and over who present with a risk factor and one or more symptoms of COPD receive quality-assured post-bronchodilator spirometry to confirm a diagnosis of COPD.

What the quality statement means for patients, service users and carers

People aged 35 or older who have an increased risk of COPD and who have one or more symptoms

of COPD are offered a test to check how well their lungs work (called post-bronchodilator spirometry). This test is used to diagnose COPD. People are at an increased risk of COPD if they smoke or have smoked in the past, or if they have been exposed to harmful fumes, dust or chemicals, often at work. Symptoms of COPD include breathlessness, long-lasting cough, coughing up phlegm, frequent winter 'bronchitis' and wheezing.

Source guidance

Chronic obstructive pulmonary disease in over 16s: diagnosis and management (2018) NICE guideline NG115, recommendations 1.1.1, 1.1.5, 1.1.8 and 1.1.10

Definitions of terms used in this quality statement

Risk factors

Risk factors for COPD include:

- smoking history
- occupational exposure to harmful fumes, dust or chemicals
- exposure to fumes, such as biomass fuels.

[NICE's guideline on [chronic obstructive pulmonary disease](#) and expert opinion]

Symptoms of COPD

Symptoms of COPD are:

- exertional breathlessness
- chronic cough
- regular sputum production
- frequent winter 'bronchitis'
- wheeze.

[NICE's guideline on [chronic obstructive pulmonary disease](#), recommendation 1.1.1]

Post-bronchodilator spirometry

Post-bronchodilator spirometry is used to identify abnormalities in lung volumes and air flow. Spirometry should be performed by a healthcare professional who has had appropriate training and who has up-to-date skills. The use of post-bronchodilator spirometry should be supported by quality control processes.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#), recommendations 1.1.9 and 1.1.10]

Quality statement 2: Inhaler technique

Quality statement

People with chronic obstructive pulmonary disease (COPD) who are prescribed an inhaler have their inhaler technique assessed when starting treatment and then regularly during treatment. [2011, updated 2016]

Rationale

Bronchodilator therapy is usually delivered using a hand-held inhaler device. People with COPD need to use their inhaler correctly to receive the optimal treatment dose. Assessing inhaler technique should happen at the first prescription once a person has been taught the correct technique, and then be reassessed regularly (for example, at their annual review, if their treatment changes or after an acute exacerbation) throughout the duration of a person's treatment in primary, community and secondary care services.

Quality measures

Structure

a) Evidence of local arrangements and written clinical protocols to ensure that people with COPD who are prescribed an inhaler have their technique assessed at the start of treatment and then regularly during their treatment.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Evidence of local arrangements and written clinical protocols to ensure that healthcare professionals in primary, community and secondary care services are trained and competent in teaching inhaler technique.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

Process

a) Proportion of people with COPD prescribed an inhaler who have their inhaler technique assessed at the start of treatment.

Numerator – the number in the denominator who have their inhaler technique assessed at the start

of treatment.

Denominator – the number of people with COPD prescribed an inhaler.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Proportion of people with COPD prescribed an inhaler who have their inhaler technique assessed at their annual review.

Numerator – the number in the denominator whose last inhaler annual review was no longer than 12 months since the previous one or since inhaler initiation.

Denominator – the number of people with COPD prescribed an inhaler for more than 12 months.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

c) Proportion of people with COPD prescribed an inhaler who have their inhaler technique assessed after a change in treatment.

Numerator – the number in the denominator who had their inhaler technique assessed after a change in treatment.

Denominator – the number of people with COPD prescribed an inhaler who have had their inhaler changed.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

d) Proportion of people with COPD prescribed an inhaler who have their inhaler technique assessed after an acute exacerbation.

Numerator – the number in the denominator who had their inhaler technique assessed after an acute exacerbation.

Denominator – the number of people with COPD prescribed an inhaler who have had an acute exacerbation.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

Outcome

a) Exacerbation rates.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Hospital admissions.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (primary care services, community services and secondary care services) ensure that systems are in place and healthcare professionals are trained and competent to teach people with COPD who are prescribed an inhaler the correct inhaler technique and to assess their inhaler technique when starting treatment and regularly during their treatment.

Healthcare professionals (nurses, GPs, secondary care doctors, physiotherapists, occupational therapists and pharmacists) ensure that they provide training in the correct inhaler technique to people with COPD when they have been prescribed an inhaler. Healthcare professionals ensure that they assess the person's inhaler technique when starting treatment and regularly during their treatment.

Commissioners (clinical commissioning groups) ensure that they commission services in which people with COPD who are prescribed an inhaler are trained and assessed in the correct inhaler technique when they start treatment, and have their technique reassessed regularly during their treatment.

What the quality statement means for patients, service users and carers

People with COPD who are given an inhaler have a check to make sure that they can use it correctly when they start treatment and at least once a year at their annual review. They should also have a check if their treatment changes or after a sudden flare up of their symptoms (called an acute exacerbation).

Source guidance

Chronic obstructive pulmonary disease in over 16s: diagnosis and management (2018) NICE guideline NG115, recommendations 1.2.20, 1.2.21 and 1.3.46

Equality and diversity considerations

Elderly people, or people with learning disabilities, physical disabilities or cognitive impairment may experience difficulties learning and retaining the adequate inhaler technique to ensure that they get the optimal treatment dose. An individual patient assessment should be carried out before choosing the most appropriate device for delivery of inhaled therapy.

Quality statement 3: Assessment for long-term oxygen therapy

Quality statement

People with stable chronic obstructive pulmonary disease (COPD) and a persistent resting stable oxygen saturation level of 92% or less have their arterial blood gases measured to assess whether they need long-term oxygen therapy (LTOT). [2011, updated 2016]

Rationale

LTOT is used to treat people with stable COPD who have developed daytime hypoxaemia. People with COPD and a persistent resting stable oxygen saturation of 92% or less should be assessed for their suitability for LTOT, which can improve survival, pulmonary haemodynamics, polycythaemia and neuropsychological health.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that people with stable COPD and a persistent resting stable oxygen saturation level of 92% or less have their arterial blood gases measured to assess whether they need LTOT.

Data source: Local data collection Royal College of Physicians' [National COPD Audit Programme](#).

Process

Proportion of people with stable COPD and a persistent resting stable oxygen saturation level of 92% or less who have their arterial blood gases measured to assess whether they need LTOT.

Numerator – the number in the denominator who have their arterial blood gases measured to assess whether they need LTOT.

Denominator – the number of people with stable COPD and a persistent resting stable oxygen saturation level of 92% or less.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

Outcome

a) Hospital admission for acute exacerbation.

Data source: Local data collection.

b) Quality of life.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (primary and secondary care services) ensure that systems are in place for people with stable COPD and a persistent resting oxygen saturation level of 92% or less to have their arterial blood gases measured to assess whether they need LTOT.

Healthcare professionals ensure that they measure the arterial blood gases of people with stable COPD and a persisting resting oxygen saturation level of 92% or less to assess whether they need LTOT.

Commissioners (clinical commissioning groups) ensure that they commission services in which people with stable COPD and a persisting resting oxygen saturation level of 92% or less have their arterial blood gases measured to assess whether they need LTOT.

What the quality statement means for patients, service users and carers

People with COPD that is stable and who have low levels of oxygen in their blood (when checked using a device that clips to their finger) have this confirmed by a blood test, to assess whether they need long-term oxygen therapy. Long-term oxygen therapy is treatment with oxygen breathed in through a tube (placed just inside the nose) or a mask connected to an oxygen supply. It is usually given for at least 15 hours during the day or night.

Source guidance

- [Chronic obstructive pulmonary disease in over 16s: diagnosis and management \(2018\) NICE guideline NG115, recommendation 1.2.53](#)
- British Thoracic Society (2015) [Guidelines for home oxygen use in adults, Referral and](#)

- assessment of patients for LTOT, page i11, bullet point 5

Definitions

Long-term oxygen therapy (LTOT)

The provision of oxygen therapy for continuous use at home, usually given for at least 15 hours during the day or night.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#)]

Assessment for LTOT

Assessing people for LTOT should comprise measuring arterial blood gases on 2 occasions at least 3 weeks apart in people who have a confident diagnosis of COPD, who are receiving optimum medical management and whose COPD is stable.

[NICE's guideline on [chronic obstructive pulmonary disease](#), recommendation 1.2.54]

Stable COPD

The absence of any of the features of a recent acute exacerbation, such as worsening breathlessness, cough, increased sputum production and change in colour of sputum.

[NICE's guideline on [chronic obstructive pulmonary disease](#), section 1.3]

Persistent resting stable oxygen saturation

An oxygen saturation (measured with a pulse oximeter) that is persistently 92% or less when the person is in a chronic stable state and is at rest (is not, and has not recently, been exercising).

[Expert opinion]

Quality statement 4: Pulmonary rehabilitation for stable COPD and exercise limitation

Quality statement

People with stable chronic obstructive pulmonary disease (COPD) and exercise limitation due to breathlessness are referred to a pulmonary rehabilitation programme. [2011, updated 2016]

Rationale

Pulmonary rehabilitation programmes improve a person's exercise capacity, quality of life, symptoms and levels of anxiety and depression.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that people with stable COPD and exercise limitation due to breathlessness are referred to a pulmonary rehabilitation programme.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit and organisational audit](#).

Process

a) Proportion of people with stable COPD and exercise limitation due to breathlessness who are referred to a pulmonary rehabilitation programme.

Numerator – the number in the denominator who are referred to a pulmonary rehabilitation programme.

Denominator – the number of people with stable COPD and exercise limitation due to breathlessness.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit](#).

b) Proportion of referrals of people with stable COPD and exercise limitation due to breathlessness that result in the person attending a pulmonary rehabilitation programme.

Numerator – the number in the denominator that result in the person attending a pulmonary rehabilitation programme.

Denominator – the number of referrals of people with stable COPD and exercise limitation due to breathlessness to pulmonary rehabilitation programmes.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit](#).

c) Proportion of attendances of people with stable COPD and exercise limitation due to breathlessness that result in the person completing a pulmonary rehabilitation programme.

Numerator – the number in the denominator that result in the person completing a pulmonary rehabilitation programme.

Denominator – the number of attendances of people with stable COPD and exercise limitation due to breathlessness at pulmonary rehabilitation programmes.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit](#).

Outcome

a) Hospital admissions for acute exacerbation.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Quality of life.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

c) Exercise capacity.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: Pulmonary rehabilitation clinical audit](#).

d) GP attendances.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care and community services) ensure that systems are in place for people with stable COPD and exercise limitation due to breathlessness to be referred to a pulmonary rehabilitation programme.

Healthcare professionals refer people with stable COPD and exercise limitation due to breathlessness are referred to a pulmonary rehabilitation programme.

Commissioners (clinical commissioning groups) ensure that they commission services in which people with stable COPD and exercise limitation due to breathlessness are referred to a pulmonary rehabilitation programme.

What the quality statement means for patients, service users and carers

People with COPD that is stable and who have difficulty walking and have to walk slowly and stop often or soon become breathless, are referred to a pulmonary rehabilitation programme. This includes exercises, information about COPD, diet advice and support depending on the person's needs.

Source guidance

[Chronic obstructive pulmonary disease in over 16s: diagnosis and management \(2018\) NICE guideline NG115](#), recommendation 1.2.79

Definitions of terms used in this quality statement

Exercise limitation

Medical Research Council dyspnoea scale of breathlessness grade 3 and above. A breathlessness of grade 3 is defined as 'walks slower than contemporaries on level ground because of breathlessness, or has to stop for breath when walking at own pace'.

[NICE's guideline on [chronic obstructive pulmonary disease](#)]

Pulmonary rehabilitation programme

A multidisciplinary programme of care for people with chronic respiratory impairment that is individually tailored and designed to optimise each person's physical and social performance and autonomy.

[NICE's guideline on [chronic obstructive pulmonary disease](#) and British Thoracic Society's [guideline on pulmonary rehabilitation in adults](#)]

Pulmonary rehabilitation programmes should be held at times that suit people with COPD and in locations that are easy for people with COPD to get to, and have good access for people with disabilities. Programmes should be available within a reasonable time from referral.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#), recommendation 1.2.80]

Programmes comprise individualised exercise programmes and education, and:

- are at least 6 weeks in duration and include a minimum of twice-weekly supervised sessions
- include supervised, individually tailored and prescribed, progressive exercise training including both aerobic and resistance training
- include a defined, structured education programme.

[British Thoracic Society's [guideline on pulmonary rehabilitation in adults](#)]

Equality and diversity considerations

Pulmonary rehabilitation is not suitable for people with unstable cardiac disease, locomotor or neurological difficulties precluding exercise such as severe arthritis or peripheral vascular disease, and people in a terminal phase of an illness or with significant cognitive or psychiatric impairment.

Quality statement 5: Pulmonary rehabilitation after an acute exacerbation

Quality statement

People admitted to hospital for an acute exacerbation of chronic obstructive pulmonary disease (COPD) start a pulmonary rehabilitation programme within 4 weeks of discharge. [2011, updated 2016]

Rationale

Starting a pulmonary rehabilitation programme within 4 weeks of hospital discharge after an acute exacerbation reduces the short-term risk of hospital readmission, and improves the quality of life and the short-term exercise capacity of people with COPD.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that people with COPD admitted to hospital for an acute exacerbation start a pulmonary rehabilitation programme within 4 weeks of discharge.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit and organisational audit](#).

Process

Proportion of people discharged from hospital after an acute exacerbation of COPD who start a pulmonary rehabilitation programme within 4 weeks of discharge.

Numerator – the number in the denominator who start a pulmonary rehabilitation programme within 4 weeks of discharge.

Denominator – the number of people discharged from hospital after an acute exacerbation of COPD.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit](#).

Outcome

a) Hospital admissions for acute exacerbations.

Data source:Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Quality of life.

Data source:Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

c) Exercise capacity.

Data source:Local data collection. Royal College of Physicians' [National COPD Audit Programme: pulmonary rehabilitation clinical audit](#).

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care and community services) ensure that systems are in place for people admitted to hospital for an acute exacerbation of COPD to start a pulmonary rehabilitation programme within 4 weeks of discharge.

Healthcare professionals ensure that people admitted to hospital for an acute exacerbation of COPD are referred for and receive a pulmonary rehabilitation programme within 4 weeks of discharge.

Commissioners (clinical commissioning groups) ensure that they commission services in which people who are admitted to hospital for an acute exacerbation of COPD are referred for and receive a pulmonary rehabilitation programme within 4 weeks of discharge.

What the quality statement means for patients, service users and carers

People with COPD who have had a hospital stay because of a sudden flare up of their symptoms (called an acute exacerbation) start a pulmonary rehabilitation programme within 4 weeks of leaving hospital. This includes exercises, information about COPD, diet advice and support depending on the person's needs.

Source guidance

- [Chronic obstructive pulmonary disease in over 16s: diagnosis and management \(2018\) NICE guideline NG115](#), recommendation 1.2.78
- British Thoracic Society (2013) [Guideline on pulmonary rehabilitation in adults](#), Post-exacerbation pulmonary rehabilitation page ii15, paragraph 6

Definitions of terms used in this quality statement

Acute exacerbation

An exacerbation is a sustained worsening of a person's symptoms from their usual stable state and which is beyond usual day-to-day variations and acute in onset. Commonly reported symptoms are: worsening breathlessness, cough, increased sputum production and change in sputum colour.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#)]

Exercise capacity and physical activity levels are impaired during and after an exacerbation, contributing to skeletal muscle dysfunction, particularly of the lower limbs.

[Adapted from British Thoracic Society's [guideline on pulmonary rehabilitation in adults](#)]

Pulmonary rehabilitation programme

A multidisciplinary programme of care for people with chronic respiratory impairment that is individually tailored and designed to optimise each person's physical and social performance and autonomy.

[NICE's guideline on [chronic obstructive pulmonary disease](#) and British Thoracic Society's [guideline for pulmonary rehabilitation in adults](#)]

Pulmonary rehabilitation programmes should be held at times that suit people with COPD and in locations that are easy for people with COPD to get to, and have good access for people with disabilities. Programmes should be available within a reasonable time from referral.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#), recommendation 1.2.80]

Programmes comprise individualised exercise programmes and education, and:

- are at least 6 weeks in duration and include a minimum of twice-weekly supervised sessions
- include supervised, individually tailored and prescribed, progressive exercise training including both aerobic and resistance training
- include a defined, structured education programme.

[British Thoracic Society's [guideline on pulmonary rehabilitation in adults](#)]

Equality and diversity considerations

Pulmonary rehabilitation is not suitable for people with unstable cardiac disease, locomotor or neurological difficulties precluding exercise such as severe arthritis or peripheral vascular disease, and people in a terminal phase of an illness or with significant cognitive or psychiatric impairment.

Some people with COPD may not be well enough to attend a pulmonary rehabilitation programme within 4 weeks of an acute exacerbation, may not have attended hospital after an acute exacerbation of COPD or may not have been admitted to hospital after their exacerbation of COPD.

Quality statement 6: Emergency oxygen during an exacerbation

Quality statement

People receiving emergency oxygen for an acute exacerbation of chronic obstructive pulmonary disease (COPD) have their oxygen saturation levels maintained between 88% and 92%. [new 2016]

Rationale

During an exacerbation, people with COPD may experience a worsening of gas exchange in the lungs, which can lead to low blood oxygen levels. Emergency oxygen is often given during the treatment of an exacerbation, either in the community, during transfer to hospital in an ambulance or while being assessed at hospital.

In some people, uncontrolled oxygen therapy may reduce the depth and frequency of breathing, leading to a rise in blood carbon dioxide levels and a fall in the blood pH (acidosis). Controlled oxygen therapy must therefore be administered by a delivery device and at a flow rate that helps the oxygen saturation to be maintained between 88% and 92%.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that people receiving emergency oxygen for an acute exacerbation of COPD have their oxygen saturation levels maintained between 88% and 92%.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

Process

Proportion of people receiving emergency oxygen for an acute exacerbation of COPD who have their oxygen saturation levels maintained between 88% and 92%.

Numerator – the number in the denominator whose oxygen saturation levels are maintained between 88% and 92%.

Denominator – the number of people with an acute exacerbation of COPD receiving emergency oxygen.

Outcome

a) Frequency of non-invasive ventilation due to oxygen toxicity.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

b) Morbidity rates.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (community and secondary care services, ambulance trusts, A&E departments) ensure that devices and flow rates are used to enable oxygen saturation levels to be maintained between 88% and 92% in people receiving emergency oxygen for an acute exacerbation of COPD.

Healthcare professionals ensure that devices and flow rates are used to enable oxygen saturation levels to be maintained between 88% and 92% in people receiving emergency oxygen for an acute exacerbation of COPD.

Commissioners ensure that they commission services that use devices and flow rates to enable oxygen saturation levels to be maintained between 88% and 92% in people receiving emergency oxygen for an acute exacerbation of COPD.

What the quality statement means for patients, service users and carers

People with COPD who need emergency oxygen because of a sudden flare up of their symptoms (called an acute exacerbation) receive the correct amount of oxygen to keep the oxygen levels in their blood at a safe level.

Source guidance

[Chronic obstructive pulmonary disease in over 16s: diagnosis and management \(2018\) NICE guideline NG115, recommendation 1.3.31](#)

Definitions of terms used in this quality statement

Acute exacerbation

An exacerbation is a sustained worsening of a person's symptoms from their stable state beyond usual day-to-day variations and is acute in onset. Commonly reported symptoms are: worsening breathlessness, cough, increased sputum production and change in the colour of the sputum.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#)]

Quality statement 7: Non-invasive ventilation

Quality statement

People with an acute exacerbation of chronic obstructive pulmonary disease (COPD) and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical therapy have non-invasive ventilation. [2011, updated 2016]

Rationale

Non-invasive ventilation is used to treat persistent hypercapnic ventilatory failure and acidosis during an exacerbation of COPD, when a person's arterial blood gases (especially the pH and carbon dioxide levels) are not responding (or worsening) despite optimal medical management. Non-invasive ventilation should be delivered in a dedicated setting by staff trained and experienced in its use because of safety concerns with using the equipment.

Quality measures

Structure

Evidence of local arrangements to ensure that people with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical treatment have non-invasive ventilation.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

Process

Proportion of people with an exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical treatment who have non-invasive ventilation.

Numerator – the number in the denominator who have non-invasive ventilation.

Denominator – the number of people with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical therapy.

Outcome

Mortality rates.

Data source: Local data collection. Royal College of Physicians' [National COPD Audit Programme](#).

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care services and A&E departments) ensure that people with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical treatment have non-invasive ventilation.

Healthcare professionals ensure that people with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical treatment have non-invasive ventilation. Healthcare professionals are trained and experienced in using non-invasive ventilation.

Commissioners (clinical commissioning groups) ensure that they commission services in which people with an acute exacerbation of COPD and persistent acidotic hypercapnic ventilatory failure that is not improving after 1 hour of optimal medical treatment have non-invasive ventilation.

What the quality statement means for patients, service users and carers

People with COPD who have 'ventilatory failure' during a sudden flare up of their symptoms (called an acute exacerbation) are given an emergency treatment called non-invasive ventilation if they do not improve after 1 hour of treatment with medicine and oxygen. Ventilatory failure happens when a person can't breathe deeply enough and waste carbon dioxide builds up in the blood causing acid to form. Non-invasive ventilation involves wearing a mask connected to a machine that pumps oxygen into the lungs.

Source guidance

[Chronic obstructive pulmonary disease in over 16s: diagnosis and management \(2018\) NICE guideline NG115, recommendation 1.3.34](#)

Definitions of terms used in this quality statement

Acute exacerbation

An acute exacerbation is a sustained worsening of a person's symptoms from their stable state, and which is beyond usual day-to-day variations and acute in onset. Commonly reported symptoms are worsening breathlessness, cough, increased sputum production and change in sputum colour.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#)]

Persistent acidotic hypercapnic ventilatory failure

Acute acidotic hypercapnic respiratory failure results from an inability of the respiratory system to provide sufficient alveolar ventilation to maintain a normal arterial PCO₂ and blood pH level. Co-existent hypoxaemia is usually mild and easily corrected. Conventionally, a pH <7.35 and a PCO₂ >6.5 kPa, persisting after initial medical therapy, define acute respiratory acidosis and have been used as threshold values for considering the use of non-invasive ventilation. More severe degrees of acidosis, such as pH <7.25, have been used as a threshold for considering provision of invasive mechanical ventilation.

[Adapted from NICE's guideline on [chronic obstructive pulmonary disease](#) and expert consensus]

Non-invasive ventilation

Non-invasive ventilation is a method of providing ventilatory support that does not require an endotracheal tube. It is usually delivered through a mask that covers the nose or a mask covering the nose and the mouth.

[NICE's guideline on [chronic obstructive pulmonary disease](#)]

Non-invasive ventilation should be given once it is recognised that a person is not responding to 1 hour of optimal medical therapy.

[Expert consensus]

Optimal medical treatment

Controlled oxygen therapy, nebulised bronchodilator therapy, systemic corticosteroids and antibiotics if indicated, in line with the NICE guideline.

[NICE's guideline on [chronic obstructive pulmonary disease](#)]

Quality statement 8 (placeholder): Hospital discharge care bundle

What is a placeholder statement?

A placeholder statement is an area of care that has been prioritised by the Quality Standards Advisory Committee but for which no source guidance is currently available. A placeholder statement indicates the need for evidence-based guidance to be developed in this area.

Rationale

Hospital discharge care bundles are designed to ensure that every person leaving hospital receives the best care. They emphasise the key interventions in the management pathway, including details of settings for care and treatment. There are several elements of ongoing care that an adult with COPD should start before discharge from hospital, which can improve their outcome. There is currently a lack of evidence-based guidance about the details that should be included in these care bundles. [new 2016]

Using the quality standard

Quality measures

The quality measures accompanying the quality statements aim to improve the structure, process and outcomes of care in areas identified as needing quality improvement. They are not a new set of targets or mandatory indicators for performance management.

We have indicated if current national indicators exist that could be used to measure the quality statements. If there is no national indicator that could be used to measure a quality statement, the quality measure should form the basis for audit criteria developed and used locally.

See NICE's [how to use quality standards](#) for further information, including advice on using quality measures.

Levels of achievement

Expected levels of achievement for quality measures are not specified. Quality standards are intended to drive up the quality of care, and so achievement levels of 100% should be aspired to (or 0% if the quality statement states that something should not be done). However, NICE recognises that this may not always be appropriate in practice, taking account of safety, choice and professional judgement, and therefore desired levels of achievement should be defined locally.

Using other national guidance and policy documents

Other national guidance and current policy documents have been referenced during the development of this quality standard. It is important that the quality standard is considered alongside the documents listed in [development sources](#).

Diversity, equality and language

During the development of this quality standard, equality issues have been considered and [equality assessments](#) are available.

Good communication between health, public health and social care practitioners and people with COPD in primary, community and secondary care is essential. Treatment, care and support, and the information given about it, should be culturally appropriate. It should also be accessible to people with additional needs such as physical, sensory or learning disabilities, and to people who do not speak or read English. People with COPD in primary, community or secondary care should have access to an interpreter or advocate if needed.

Commissioners and providers should aim to achieve the quality standard in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this quality standard should be interpreted in a way that would be inconsistent with compliance with those duties.

Development sources

Further explanation of the methodology used can be found in the quality standards [process guide](#).

Evidence sources

The documents below contain recommendations from NICE guidance or other NICE-accredited recommendations that were used by the Quality Standards Advisory Committee to develop the quality standard statements and measures.

- [BTS guideline for home oxygen use in adults](#) (2015) British Thoracic Society
- [BTS guideline on pulmonary rehabilitation in adults](#) (2013) British Thoracic Society
- [Chronic obstructive pulmonary disease in over 16s: diagnosis and management](#) (2018) NICE guideline NG115

Policy context

It is important that the quality standard is considered alongside current policy documents, including:

- Department of Health (2011) [An outcomes strategy for people with chronic obstructive pulmonary disease \(COPD\) and asthma in England](#)
- Department of Health (2012) [An outcomes strategy for COPD and asthma: NHS companion document](#)

Definitions and data sources for the quality measures

- Royal College of Physicians (2015) [National chronic obstructive pulmonary disease \(COPD\) Audit Programme](#)

Related NICE quality standards

- [Bronchiolitis in children](#) (2016) NICE quality standard 122
- [Pneumonia in adults](#) (2016) NICE quality standard 110
- [Smoking: supporting people to stop](#) (2013) NICE quality standard 43
- [Asthma](#) (2013) NICE quality standard 25
- [Patient experience in adult NHS services](#) (2012) NICE quality standard 15
- [End of life care for adults](#) (2011) NICE quality standard 13

The full list of quality standard topics referred to NICE is available from the [quality standards topic library](#) on the NICE website.

Quality Standards Advisory Committee and NICE project team

Quality Standards Advisory Committee

This quality standard has been developed by Quality Standards Advisory Committee 3. Membership of this committee is as follows:

Ms Deryn Bishop

Public Health Behaviour Change Specialist, Solihull Public Health Department

Jan Dawson

Registered Dietitian

Dr Matthew Fay

GP, Westcliffe Medical Practice, Shipley, West Yorkshire

Dr Malcolm Fisk

Co-Director, Ageing Society Grand Challenge Initiative, Coventry University

Margaret Goose

Lay member

Dr Madhavan Krishnaswamy

Consultant Clinical Oncologist, Southend University Hospital NHS Trust

Mrs Geeta Kumar

Clinical Director, Women's Services (East) Betsi Cadwaladr University Health Board

Dr Hugh McIntyre (Chair)

Consultant Physician, East Sussex Healthcare Trust

Ms Ann Nevinson

Lay member

Professor Gillian Parker

Professor of Social Policy Research, Social Policy Research Unit, University of York

Mr David Pugh

Independent Consultant, Gloucestershire County Council

Dr Eve Scott

Head of Safety and Risk, The Christie NHS Foundation Trust, Manchester

Dr Jim Stephenson

Consultant Medical Microbiologist, Epsom and St Helier NHS Trust

Mr Darryl Thompson

Registered Nurse (Mental Health), South West Yorkshire Partnership NHS Foundation Trust

Mrs Julia Thompson

Health Improvement Principal, Sheffield City Council

Mrs Sarah Williamson

Clinical Quality Assurance and Performance Manager, NHS Stockport Clinical Commissioning Group

The following specialist members joined the committee to develop this quality standard:

Dr Michael Rudolf

Honorary Consultant Physician, London North West Healthcare NHS Trust

Chris Loveridge

Nurse, Education for Health

Dr Kevin Gruffyard-Jones

GP, Box Surgery, Wiltshire

Mr Alex Woodward

Physiotherapist, Leicestershire Partnership NHS Trust

Professor Mike Roberts

Consultant Physician, Barts Health

Michael McKevitt

Lay member

NICE project team

Nick Baillie

Associate Director

Craig Grime

Technical Adviser

Alison Tariq

Lead Technical Analyst

Esther Clifford

Programme Manager

Liane Marsh

Coordinator

Update information

February 2016: This quality standard was updated and statements prioritised in 2011 were replaced.

Statements are marked as [new 2016] or [2011, updated 2016]:

- [new 2016] if the statement covers a new area for quality improvement
- [2011, updated 2016] if the statement covers an area for quality improvement included in the 2011 quality standard and has been updated.

Statements numbered 1, 3, 6, 8 and 11 in the 2011 version have been updated and included in the updated quality standard, marked as [2011, updated 2016].

Statements from the 2011 version (numbered 2, 4, 5, 7, 9, 10, 12 and 13) that are no longer considered national priorities for improvement, but may still be useful at a local level, are included after the updated statements in the [list of quality statements](#) section.

- The [2011 quality standard for COPD](#) is available as a pdf.

Minor changes since publication

December 2018: Source guidance references have been updated to align this quality standard with the NICE guideline on [chronic obstructive pulmonary disease](#).

About this quality standard

NICE quality standards describe high-priority areas for quality improvement in a defined care or service area. Each standard consists of a prioritised set of specific, concise and measurable statements. NICE quality standards draw on existing NICE or NICE-accredited guidance that provides an underpinning, comprehensive set of recommendations, and are designed to support the measurement of improvement.

The methods and processes for developing NICE quality standards are described in the quality standards [process guide](#).

This quality standard has been incorporated into the NICE Pathway on [chronic obstructive pulmonary disease](#).

NICE produces guidance, standards and information on commissioning and providing high-quality healthcare, social care, and public health services. We have agreements to provide certain NICE services to Wales, Scotland and Northern Ireland. Decisions on how NICE guidance and other products apply in those countries are made by ministers in the Welsh government, Scottish government, and Northern Ireland Executive. NICE guidance or other products may include references to organisations or people responsible for commissioning or providing care that may be relevant only to England.

ISBN: 978-1-4731-1652-8

Endorsing organisation

This quality standard has been endorsed by NHS England, as required by the Health and Social Care Act (2012)

Supporting organisations

Many organisations share NICE's commitment to quality improvement using evidence-based guidance. The following supporting organisations have recognised the benefit of the quality standard in improving care for patients, carers, service users and members of the public. They have agreed to work with NICE to ensure that those commissioning or providing services are made aware of and encouraged to use the quality standard.

- [Society for Acute Medicine \(SAM\)](#)

- [Chartered Society of Physiotherapy](#)
- [Primary Care Respiratory Society UK](#)
- [British Thoracic Society](#)
- [Royal College of General Practitioners](#)