

Preterm labour and birth

Quality standard

Published: 19 October 2016

[nice.org.uk/guidance/qs135](https://www.nice.org.uk/guidance/qs135)

Contents

Introduction	5
Why this quality standard is needed	5
How this quality standard supports delivery of outcome frameworks	6
Safety and people's experience of care	8
Coordinated services.....	8
List of quality statements.....	10
Quality statement 1: Providing information about potential signs and symptoms of preterm labour	11
Quality statement.....	11
Rationale	11
Quality measures	11
What the quality statement means for service providers, healthcare professionals and commissioners ..	12
What the quality statement means for women and their companions	12
Source guidance.....	12
Definitions of terms used in this quality statement	12
Equality and diversity considerations.....	14
Quality statement 2: Prophylactic vaginal progesterone and prophylactic cervical cerclage	15
Quality statement.....	15
Rationale	15
Quality measures	15
What the quality statement means for service providers, healthcare professionals and commissioners ..	16
What the quality statement means for women and their companions	16
Source guidance.....	17
Definitions of terms used in this quality statement	17
Quality statement 3: Information for women having a planned preterm birth	18
Quality statement.....	18
Rationale	18

Quality measures	18
What the quality statement means for service providers, healthcare professionals and commissioners ..	19
What the quality statement means for women and their companions	19
Source guidance.....	19
Definitions of terms used in this quality statement	20
Equality and diversity considerations.....	20
Quality statement 4: Tocolysis and corticosteroids for women in suspected preterm labour between 26+0 and 29+6 weeks of pregnancy.....	21
Quality statement.....	21
Rationale	21
Quality measures	21
What the quality statement means for service providers, healthcare professionals and commissioners ..	23
What the quality statement means for women and their companions	23
Source guidance.....	23
Definitions of terms used in this quality statement	23
Quality statement 5: Corticosteroids for women between 30+0 and 33+6 weeks of pregnancy	25
Quality statement.....	25
Rationale	25
Quality measures	25
What the quality statement means for service providers, healthcare professionals and commissioners ..	26
What the quality statement means for women and their companions	27
Source guidance.....	27
Definitions of terms used in this quality statement	27
Quality statement 6: Magnesium sulfate for women between 24+0 and 29+6 weeks of pregnancy.....	29
Quality statement.....	29
Rationale	29
Quality measures	29
What the quality statement means for service providers, healthcare professionals and commissioners ..	30

What the quality statement means for women and their companions	30
Source guidance.....	31
Definitions of terms used in this quality statement	31
Using the quality standard.....	32
Quality measures	32
Levels of achievement	32
Using other national guidance and policy documents.....	32
Diversity, equality and language	33
Development sources.....	34
Evidence sources.....	34
Policy context	34
Related NICE quality standards	35
Published	35
Future quality standards.....	35
Quality standards advisory committee and NICE project team	36
Quality standards advisory committee	36
NICE project team	38
About this quality standard.....	39

This standard is based on NG25.

This standard should be read in conjunction with QS121, QS120, QS109, QS75, QS60, QS35, QS32, QS15, QS4, QS105, QS162 and QS169.

Introduction

This quality standard covers care for pregnant women who are considered to be at risk of, or with symptoms and signs of, preterm labour and birth. It does not cover women with a multiple pregnancy. For more information see the [preterm labour and birth topic overview](#).

Why this quality standard is needed

Preterm birth (that is, before 37 weeks of pregnancy) is the single biggest cause of neonatal mortality and morbidity in the UK. Babies born preterm have high rates of early, late and post neonatal mortality, and the risk of mortality increases as gestational age at birth decreases. Babies who survive have increased rates of disability.

The most important long-term consequence of prematurity is neurodevelopmental disability. This can range from severe motor abnormalities, such as cerebral palsy, to less severe cognitive disabilities. Although the risk for the individual child is greatest for those born at the earliest gestational ages, the global burden of neurodevelopmental disabilities depends on the number of babies born at each of these gestations, and so is very significant for babies born between 32 and 36 weeks, less for those born between 28 and 31 weeks, and least for those born at less than 28 weeks of gestation.

Around 75% of women delivering preterm do so after preterm labour, which may or may not be preceded by preterm prelabour rupture of membranes (P-PROM). The remaining 25% have a planned preterm birth because of medical complications (for example, extreme growth retardation in the baby or maternal conditions such as pre-eclampsia).

The quality standard is expected to contribute to improvements in the following outcomes:

- incidence of preterm births
- fetal morbidity and mortality

- neonatal morbidity and mortality
- maternal morbidity
- neurodevelopmental disabilities
- women's experience of childbirth.

How this quality standard supports delivery of outcome frameworks

NICE quality standards are a concise set of prioritised statements designed to drive measurable improvements in the 3 dimensions of quality – safety, experience and effectiveness of care – for a particular area of health or care. They are derived from high-quality guidance, such as that from NICE or other sources accredited by NICE. This quality standard, in conjunction with the guidance on which it is based, should contribute to the improvements outlined in the following 2 outcomes frameworks published by the Department of Health:

- [NHS outcomes framework 2016–17](#)
- [Public health outcomes framework for England, 2016–19](#)

Tables 1 and 2 show the outcomes, overarching indicators and improvement areas from the frameworks that the quality standard could contribute to achieving.

Table 1 [NHS outcomes framework 2016–17](#)

Domain	Overarching indicators and improvement areas
1 Preventing people from dying prematurely	<p><i>Overarching indicators</i></p> <p>1a Potential Years of Life Lost (PYLL) from causes considered amenable to healthcare</p> <p>i Adults ii Children and young people</p> <p>1c Neonatal mortality and stillbirths</p> <p><i>Improvement areas</i></p> <p>Reducing mortality in children</p> <p>1.6 i Infant mortality*</p>

<p>4 Ensuring that people have a positive experience of care</p>	<p>Overarching indicators</p> <p><i>4c Friends and family test</i></p> <p><i>4d Patient experience characterised as poor or worse</i></p> <p><i>i Primary care</i></p> <p><i>ii Hospital care</i></p> <p>Improvement areas</p> <p>Improving women and their families' experience of maternity services</p> <p>4.5 Women's experience of maternity services</p>
<p>5 Treating and caring for people in a safe environment and protecting them from avoidable harm</p>	<p>Overarching indicators</p> <p><i>5a Deaths attributable to problems in healthcare</i></p> <p><i>5b Severe harm attributable to problems in healthcare</i></p> <p>Improvement areas</p> <p>Improving the culture of safety reporting</p> <p>5.6 Patient safety incidents reported</p>
<p>Alignment with Public health outcomes framework</p> <p>* Indicator is shared</p> <p>Indicators in italics in development</p>	

Table 2 Public health outcomes framework for England, 2016–19

Domain	Objectives and indicators
<p>4 Healthcare public health and preventing premature mortality</p>	<p>Objective</p> <p>Reduced numbers of people living with preventable ill health and people dying prematurely, whilst reducing the gap between communities</p> <p>Indicators</p> <p>4.01 Infant mortality*</p> <p>4.03 Mortality rate from causes considered preventable**</p>

Alignment with NHS outcomes framework

* Indicator is shared

** Indicator is complementary

Safety and people's experience of care

Ensuring that care is safe and that people have a positive experience of care is vital in a high-quality service. It is important to consider these factors when planning and delivering services relevant to pregnant women who are considered to be at risk of preterm labour and birth.

NICE has developed guidance and an associated quality standard on [patient experience in adult NHS services](#) (see the NICE pathway on [patient experience in adult NHS services](#)), which should be considered alongside this quality standard. This specifies that people receiving care should be treated with dignity, have opportunities to discuss their preferences, and be supported to understand their options and make fully informed decisions. It also covers the provision of information to patients and service users. Quality statements on these aspects of patient experience are not usually included in topic-specific quality standards. However, recommendations in the development sources for quality standards that affect people's experience of using services are specific to the topic are considered during quality statement development.

Coordinated services

The quality standard for preterm labour and birth specifies that services should be commissioned from and coordinated across all relevant agencies encompassing the whole preterm labour and birth care pathway. A person-centred, integrated approach to providing services is fundamental to delivering high-quality care to pregnant women who are considered to be at risk of preterm labour and birth.

The Health and Social Care Act 2012 sets out a clear expectation that the care system should consider NICE quality standards in planning and delivering services, as part of a general duty to secure continuous improvement in quality. Commissioners and providers of health and social care should refer to the library of NICE quality standards when designing high-quality services. Other quality standards that should also be considered when choosing, commissioning or providing a high-quality preterm labour and birth service are listed in [related quality standards](#).

Resource impact considerations

NICE quality standards should be achievable by local services. The potential resource impact is

considered by the quality standards advisory committee, drawing on resource impact work for the source guidance. Organisations are encouraged to use the [costing statement](#) for the NICE guideline on preterm labour and birth to help estimate local costs.

Training and competencies

The quality standard should be read in the context of national and local guidelines on training and competencies. All healthcare professionals involved in assessing, caring for and treating pregnant women who are considered to be at risk of preterm labour and birth should have sufficient and appropriate training and competencies to deliver the actions and interventions described in the quality standard. Quality statements on staff training and competency are not usually included in quality standards. However, recommendations in the development source on specific types of training for the topic that exceed standard professional training are considered during quality statement development.

Role of families and carers

Quality standards recognise the important role families and carers have in supporting pregnant women who are considered to be at risk of preterm labour and birth. If appropriate, healthcare professionals should ensure that family members and carers are involved in the decision-making process about investigations, treatment and care.

List of quality statements

Statement 1. Pregnant women at increased risk of preterm labour are given information about the potential signs and symptoms.

Statement 2. Women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage.

Statement 3. Women having a planned preterm birth are given information about the risks and potential outcomes.

Statement 4. Women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are offered tocolysis and maternal corticosteroids.

Statement 5. Women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have preterm prelabour rupture of membranes (P-PROM) are offered maternal corticosteroids.

Statement 6. Women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours are offered magnesium sulfate.

Quality statement 1: Providing information about potential signs and symptoms of preterm labour

Quality statement

Pregnant women at increased risk of preterm labour are given information about the potential signs and symptoms.

Rationale

Not all pregnant women at increased risk of preterm labour know what symptoms and signs to look out for. At such a vulnerable time in their lives, women and their families need information about the potential signs and symptoms of preterm labour and the care available that meets their needs and supports shared decision-making, without causing unnecessary anxiety.

Quality measures

Structure

Evidence of local arrangements and clinical protocols to ensure that pregnant women at increased risk of preterm labour are given information about the potential signs and symptoms.

Data source: Local data collection.

Process

Proportion of pregnant women at increased risk of preterm labour who are given information about the potential signs and symptoms.

Numerator – the number in the denominator who are given information about the potential signs and symptoms.

Denominator – the number of pregnant women at increased risk of preterm labour.

Data source: Local data collection.

Outcome

Pregnant women's satisfaction with the information provided.

Data source:Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care services) ensure that pregnant women at increased risk of preterm labour are given written information about the potential signs and symptoms by healthcare professionals, who also discuss this with them.

Healthcare professionals (such as midwives and obstetricians) give written information to pregnant women at increased risk of preterm labour about the potential signs and symptoms, and discuss this with them.

Commissioners (clinical commissioning groups) commission services that ensure that pregnant women at increased risk of preterm labour are given written information about the potential signs and symptoms by healthcare professionals, who also discuss this with them.

What the quality statement means for women and their companions

Pregnant women at increased risk of preterm labour(that is, going into labour before the 37th week of pregnancy) are given written information about the signs and symptoms that might suggest preterm labour. A healthcare professional also talks to them about this and the care available.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, recommendation 1.1.1

Definitions of terms used in this quality statement

Pregnant women at increased risk of preterm labour

Pregnant women who:

- have a history of:
 - spontaneous preterm birth
 - preterm prelabour rupture of membranes

- – mid-trimester loss
 - cervical trauma (including surgery – for example, previous cone biopsy [cold knife or laser], large loop excision of the transformation zone [LLETZ – any number] and radical diathermy)
- are considered to be at risk of preterm labour and birth because they have a short cervix (less than 25 mm) that has been identified on a transvaginal ultrasound scan and/or bulging membranes in the current pregnancy
- have preterm prelabour rupture of membranes (P-PROM).

[Adapted from the NICE full guideline on [preterm labour and birth](#)]

Information

Pregnant women should be given oral and written information, and be directed to organisations that can provide further support.

[Adapted from the NICE full guideline on [preterm labour and birth](#)]

Potential signs of preterm labour

Potential signs could include:

- watery, mucosal or bloody vaginal discharge
- regular or frequent (often painless) contractions or uterine tightening
- P-PROM.

[Expert opinion]

Potential symptoms of preterm labour

Potential symptoms could include:

- pelvic or lower abdominal pressure
- constant low, dull backache
- mild abdominal cramps, with or without diarrhoea.

[Expert opinion]

Equality and diversity considerations

Pregnant women at increased risk of preterm labour should have access to information that is understandable if they:

- have additional needs, such as physical, sensory or learning disabilities
- do not speak or read English
- have religious, ethnic or cultural needs.

Interpreters and advocates should be provided if needed.

Quality statement 2: Prophylactic vaginal progesterone and prophylactic cervical cerclage

Quality statement

Women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage.

Rationale

Preterm birth causes significant neonatal morbidity and mortality, as well as long-term disability. Therefore strategies for preventing preterm birth are important. Both prophylactic cervical cerclage and prophylactic vaginal progesterone are effective in preventing or delaying preterm birth in women with a short cervix and a history of spontaneous preterm birth or mid-trimester loss between 16⁺⁰ and 34⁺⁰ weeks of pregnancy. Which treatment is best for each woman will depend on her individual preferences and circumstances, and women should be given information as part of shared decision-making.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage.

Data source: Local data collection.

Process

Proportion of women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy who are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage.

Numerator – the number in the denominator who are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage.

Denominator – the number of women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy.

Data source: Local data collection.

Outcome

Timing of labour and birth.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care services) ensure that women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage, with the choice of treatment depending on the woman's preferences and circumstances.

Healthcare professionals (such as obstetricians caring for women with high-risk pregnancies) offer women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage, with the choice of treatment depending on the woman's preferences and circumstances.

Commissioners (clinical commissioning groups) commission services that ensure that women who have had a previous preterm birth or mid-trimester loss and have a cervical length of less than 25 mm measured between 16⁺⁰ and 24⁺⁰ weeks of pregnancy are offered a choice of either prophylactic vaginal progesterone or prophylactic cervical cerclage, with the choice of treatment depending on the woman's preferences and circumstances.

What the quality statement means for women and their companions

Women at increased risk of preterm labour in whom an ultrasound scan has shown that they have a short cervix are offered a treatment to stop the cervix (neck of the womb) opening early and so delay labour and birth. This could be either progesterone (a natural female sex hormone) inserted into the vagina or a stitch in the cervix, with the choice of treatment depending on the woman's

preferences and circumstances.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, recommendation 1.2.1

Definitions of terms used in this quality statement

Prophylactic vaginal progesterone

Progesterone by vaginal suppository to reduce the incidence of spontaneous preterm birth.

[NICE full guideline on [preterm labour and birth](#)]

Prophylactic cervical cerclage

A treatment for cervical weakness (also termed cervical incompetence or insufficiency) to prevent preterm birth and miscarriage.

[NICE full guideline on [preterm labour and birth](#)]

Quality statement 3: Information for women having a planned preterm birth

Quality statement

Women having a planned preterm birth are given information about the risks and potential outcomes.

Rationale

Women who are having a planned preterm birth need information about the level and nature of the risks, including how likely it is that their baby will survive. This should be given as early as possible in the antenatal period. It can help the woman and her family to understand what neonatal care their baby might need and inform their discussions with their neonatologist or paediatrician. The woman and her family should also be offered a tour of the neonatal unit.

Quality measures

Structure

Evidence of local arrangements and clinical protocols to ensure that women having a planned preterm birth are given information about the risks and potential outcomes.

Data source: Local data collection.

Process

Proportion of women having a planned preterm birth who are given information about the risks and potential outcomes.

Numerator – the number in the denominator who are given information about the risks and potential outcomes.

Denominator – the number of women having a planned preterm birth.

Data source: Local data collection.

Outcome

Women's awareness of the risks and potential outcomes of having a planned preterm birth.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care services) ensure that women having a planned preterm birth are given information about the risks and potential outcomes and the care available.

Healthcare professionals (such as obstetricians, neonatologists and paediatricians) give information to women having a planned preterm birth about the risks and potential outcomes and the care available.

Commissioners (clinical commissioning groups) commission services that ensure that women having a planned preterm birth are given information about the risks and potential outcomes and the care available.

What the quality statement means for women and their companions

Women who are having a planned preterm birth (before the 37th week of pregnancy) for medical reasons are told about what may happen. This should include:

- information about the care that the woman and her baby might need, and whether this might include being transferred to another hospital for specialist care
- information about the types of problems that a preterm baby might have, both at birth and as they grow up, including how likely these are
- offering to show the woman round the neonatal unit, and an opportunity for her to talk with a neonatologist or paediatrician.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, recommendation 1.1.2

Definitions of terms used in this quality statement

Planned preterm birth

A planned birth before 37⁺⁰ weeks of pregnancy because of medical complications.

[Adapted from the NICE full guideline on [preterm labour and birth](#)]

Information

Women should be given oral and written information, and directed to organisations that can provide further support.

[Adapted from the NICE full guideline on [preterm labour and birth](#)]

Potential outcomes

These include the likelihood of the baby surviving and long-term neurodevelopmental outcomes.

[Adapted from the NICE full guideline on [preterm labour and birth](#)]

Equality and diversity considerations

Women having a planned preterm labour should have access to information that is understandable if they:

- have additional needs, such as physical, sensory or learning disabilities
- do not speak or read English
- have religious, ethnic or cultural needs.

Interpreters and advocates should be provided if needed.

Quality statement 4: Tocolysis and corticosteroids for women in suspected preterm labour between 26⁺⁰ and 29⁺⁶ weeks of pregnancy

Quality statement

Women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are offered tocolysis and maternal corticosteroids.

Rationale

For women in suspected preterm labour, tocolysis and maternal corticosteroids may delay the birth and reduce the risk of problems such as cerebral palsy and of neonatal death. Not all women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are currently offered these treatments. It is important that the potential benefits and risks of both of these treatments are discussed with the woman and her family members. Tocolysis is appropriate only under particular circumstances, and a range of factors need to be taken into account. Giving corticosteroids to a woman before a preterm birth reduces the severity of lung disease of prematurity and of other associated complications for her baby. Maternal corticosteroids also have the potential to reduce the number of days that the baby needs to be on a ventilator.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are offered tocolysis and maternal corticosteroids.

Data source: Local data collection.

Process

a) Proportion of women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy in suspected preterm labour who receive tocolysis.

Numerator – the number in the denominator who receive tocolysis.

Denominator – the number of women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy in suspected

preterm labour.

Data source: Local data collection.

b) Proportion of women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy in suspected preterm labour who receive maternal corticosteroids.

Numerator – the number in the denominator who receive maternal corticosteroids.

Denominator – the number of women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy in suspected preterm labour.

Data source: Local data collection.

Outcome

a) Neonatal death.

Data source: Local data collection.

b) Intraventricular haemorrhage.

Data source: Local data collection.

c) Incidence of neonatal sepsis.

Data source: Local data collection.

d) Use of antibiotics.

Data source: Local data collection.

e) Ventilation.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (such as secondary or tertiary care services) ensure that women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are offered tocolysis and maternal corticosteroids as appropriate.

Healthcare professionals (such as midwives and obstetricians) offer tocolysis and maternal corticosteroids as appropriate to women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour.

Commissioners (clinical commissioning groups) commission services that ensure that women between 26⁺⁰ and 29⁺⁶ weeks of pregnancy who are in suspected preterm labour are offered tocolysis and maternal corticosteroids as appropriate.

What the quality statement means for women and their companions

Women who are more than 26 weeks but less than 30 weeks pregnant and in suspected preterm labour are offered tocolytics (medicines that slow down or stop labour) and steroid injections (to help the baby's lungs develop before the birth) if these medicines are likely to help their baby. The benefits and risks of each of these treatments are explained to them.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, sections 1.8 and 1.9

Definitions of terms used in this quality statement

Suspected preterm labour

A woman is in suspected preterm labour if she has reported symptoms of preterm labour and has had a clinical assessment (including a speculum or digital vaginal examination) that confirms the possibility of preterm labour but rules out established labour.

[NICE guideline on [preterm labour and birth](#)]

Tocolysis

Drugs used to stop or delay the progress of labour. The NICE guideline recommends nifedipine as

the first choice, or oxytocin receptor antagonists if nifedipine is contraindicated.

[NICE full guideline on [preterm labour and birth](#)]

Maternal corticosteroids

Corticosteroids (glucocorticosteroids) are anti-inflammatory medicines given to the woman (usually by intramuscular injection) which cross the placenta and accelerate fetal lung maturation.

[NICE full guideline on [preterm labour and birth](#)]

Quality statement 5: Corticosteroids for women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy

Quality statement

Women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have preterm prelabour rupture of membranes (P-PROM) are offered maternal corticosteroids.

Rationale

Giving corticosteroids to a woman before a preterm birth reduces the severity of lung disease of prematurity and of other associated complications for her baby. Maternal corticosteroids also have the potential to reduce the number of days that the baby needs to be on a ventilator.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have P-PROM are offered maternal corticosteroids.

Data source: Local data collection.

Process

a) Proportion of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy in diagnosed preterm labour who receive maternal corticosteroids.

Numerator – the number in the denominator who receive maternal corticosteroids.

Denominator – the number of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy in diagnosed preterm labour.

Data source: Local data collection.

b) Proportion of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy having a planned preterm birth

who receive maternal corticosteroids.

Numerator – the number in the denominator who receive maternal corticosteroids.

Denominator – the number of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are having a planned preterm birth.

Data source: Local data collection.

c) Proportion of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy with P-PROM who receive maternal corticosteroids.

Numerator – the number in the denominator who receive maternal corticosteroids.

Denominator – the number of women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy with P-PROM.

Data source: Local data collection.

Outcome

a) Ventilation.

Data source: Local data collection.

b) Incidence of neonatal sepsis.

Data source: Local data collection.

c) Use of antibiotics.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (secondary care services) ensure that women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have P-PROM are offered maternal corticosteroids by healthcare professionals.

Healthcare professionals (such as obstetricians and midwives) offer maternal corticosteroids to women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have P-PROM.

Commissioners (clinical commissioning groups) commission services that ensure that women between 30⁺⁰ and 33⁺⁶ weeks of pregnancy who are in diagnosed preterm labour, are having a planned preterm birth or have P-PROM are offered maternal corticosteroids.

What the quality statement means for women and their companions

Women over 30 weeks and under 34 weeks of pregnancy are offered corticosteroid injections to help their baby's lungs develop if:

- they are in diagnosed preterm labour (they have had a test that shows they are in labour) or
- they are having a planned preterm birth or
- they have preterm prelabour rupture of membranes (also called P-PROM) – this is when a woman's waters break early but labour hasn't started.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, recommendation 1.9.3

Definitions of terms used in this quality statement

Diagnosed preterm labour

A woman is in diagnosed preterm labour if she is in suspected preterm labour and has had a positive diagnostic test for preterm labour.

[NICE guideline on [preterm labour and birth](#)]

Planned preterm birth

A planned birth before 37⁺⁰ weeks of pregnancy because of medical complications.

[Adapted from NICE full guideline on [preterm labour and birth](#)]

Preterm prelabour rupture of membranes (P-PROM)

A woman is described as having P-PROM if she has ruptured membranes before 37⁺⁰ weeks of pregnancy but is not in established labour.

[NICE full guideline on [preterm labour and birth](#)]

Maternal corticosteroids

Corticosteroids (glucocorticosteroids) are anti-inflammatory medicines given to the woman (usually by intramuscular injection) which cross the placenta and accelerate fetal lung maturation.

[NICE full guideline on [preterm labour and birth](#)]

Quality statement 6: Magnesium sulfate for women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy

Quality statement

Women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours are offered magnesium sulfate.

Rationale

With advances in neonatal care in recent years, more babies born preterm are surviving. These children frequently have long-term complications associated with preterm birth. Neurological effects are common and may cause severe disability. Magnesium sulfate can protect the developing fetal brain and so has significant potential to reduce disability.

Quality measures

Structure

Evidence of local arrangements and written clinical protocols to ensure that women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours are offered magnesium sulfate.

Data source: Local data collection.

Process

a) Proportion of women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy in established preterm labour who receive magnesium sulfate.

Numerator – the number in the denominator who receive magnesium sulfate.

Denominator – the number of women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy in established preterm labour.

Data source: Local data collection.

b) Proportion of women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are having a planned

preterm birth within 24 hours who receive magnesium sulfate.

Numerator – the number in the denominator who receive magnesium sulfate in the 24 hours before the birth.

Denominator – the number of women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who have a planned preterm birth.

Data source: Local data collection.

Outcome

Incidence of cerebral palsy.

Data source: Local data collection.

What the quality statement means for service providers, healthcare professionals and commissioners

Service providers (such as secondary or tertiary care services) ensure that women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours are offered magnesium sulfate.

Healthcare professionals (such as midwives, obstetricians and neonatologists) offer magnesium sulfate to women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours.

Commissioners (clinical commissioning groups) commission services that ensure that women between 24⁺⁰ and 29⁺⁶ weeks of pregnancy who are in established preterm labour or having a planned preterm birth within 24 hours are offered magnesium sulfate.

What the quality statement means for women and their companions

Women who are more than 24 weeks but less than 30 weeks pregnant and in established preterm labour, or having a planned preterm birth within 24 hours, are offered magnesium sulfate, as this medicine is likely to help their baby. The benefits and risks of this treatment are explained to them.

Source guidance

- [Preterm labour and birth](#) (2015) NICE guideline NG25, recommendation 1.10.1

Definitions of terms used in this quality statement

Established preterm labour

A woman is in established preterm labour if she has progressive cervical dilatation from 4 cm with regular contractions.

[NICE guideline on [preterm labour and birth](#)]

Planned preterm birth

A planned birth before 37⁺⁰ weeks of pregnancy because of medical complications.

[Adapted from NICE full guideline on [preterm labour and birth](#)]

Using the quality standard

Quality measures

The quality measures accompanying the quality statements aim to improve the structure, process and outcomes of care in areas identified as needing quality improvement. They are not a new set of targets or mandatory indicators for performance management.

We have indicated if current national indicators exist that could be used to measure the quality statements. If there is no national indicator that could be used to measure a quality statement, the quality measure should form the basis for audit criteria developed and used locally.

See [how to use quality standards](#) for further information, including advice on using quality measures.

Levels of achievement

Expected levels of achievement for quality measures are not specified. Quality standards are intended to drive up the quality of care, and so achievement levels of 100% should be aspired to (or 0% if the quality statement states that something should not be done). However, NICE recognises that this may not always be appropriate in practice, taking account of safety, choice and professional judgement, and therefore desired levels of achievement should be defined locally.

NICE's [quality standard service improvement template](#) helps providers to make an initial assessment of their service compared with a selection of quality statements. It includes assessing current practice, recording an action plan and monitoring quality improvement. This tool is updated monthly to include new quality standards.

Using other national guidance and policy documents

Other national guidance and current policy documents have been referenced during the development of this quality standard. It is important that the quality standard is considered alongside the documents listed in [development sources](#).

Diversity, equality and language

During the development of this quality standard, equality issues have been considered and [equality assessments](#) are available.

Good communication between healthcare professionals and pregnant women who are considered to be at risk of preterm labour and birth is essential. Treatment, care and support, and the information given about it, should be culturally appropriate. It should also be accessible to people with additional needs such as physical, sensory or learning disabilities, and to people who do not speak or read English. Pregnant women who are considered to be at risk of preterm labour and birth should have access to an interpreter or advocate if needed.

Commissioners and providers should aim to achieve the quality standard in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this quality standard should be interpreted in a way that would be inconsistent with compliance with those duties.

Development sources

Further explanation of the methodology used can be found in the [quality standards process guide](#).

Evidence sources

The documents below contain recommendations from NICE guidance or other NICE-accredited recommendations that were used by the quality standards advisory committee to develop the quality standard statements and measures.

- [Preterm labour and birth](#) (2015) NICE guideline NG25

Policy context

It is important that the quality standard is considered alongside current policy documents, including:

- Department of Health (2013) [Maternity care facilities: planning and design \(HBN 09-02\)](#)
- NHS England (2013) [The national friends and family test \(FFT\): guidance for maternity services](#)
- Care Quality Commission (2008) [Towards better births: a review of maternity services in England](#)

Related NICE quality standards

Published

- [Antimicrobial stewardship](#) (2016) NICE quality standard 121
- [Medicines optimisation](#) (2016) NICE quality standard 120
- [Diabetes in pregnancy](#) (2016) NICE quality standard 109
- [Neonatal infection](#) (2014) NICE quality standard 75
- [Inducing labour](#) (2014) NICE quality standard 60
- [Hypertension in pregnancy](#) (2013) NICE quality standard 35
- [Caesarean section](#) (2013) NICE quality standard 32
- [Patient experience in adult NHS services](#) (2012) NICE quality standard 15
- [Neonatal specialist care](#) (2010) NICE quality standard 4

Future quality standards

This quality standard has been developed in the context of all quality standards referred to NICE, including the following topics scheduled for future development:

- Cerebral palsy
- Developmental follow-up of preterm babies

The full list of quality standard topics referred to NICE is available from the [quality standards topic library](#) on the NICE website.

Quality standards advisory committee and NICE project team

Quality standards advisory committee

This quality standard has been developed by quality standards advisory committee 1. Membership of this committee is as follows:

Dr Ivan Benett

Clinical Director, Central Manchester Clinical Commissioning Group

Dr Gita Bhutani

Associate Director for Psychological Professions, Lancashire Care NHS Foundation Trust

Mrs Jennifer Bostock

Lay member

Dr Helen Bromley

Consultant in Public Health, Cheshire West and Chester Council

Ms Amanda de la Motte

Deputy Chief Nurse, South Lincolnshire Clinical Commissioning Group

Mr Phillip Dick

Psychiatric Liaison Team Manager, West London Mental Health Trust

Ms Phyllis Dunn

Clinical Lead Nurse, University Hospital of North Staffordshire

Dr Steve Hajioff

Director of Public Health, London Borough of Hillingdon

Dr Ian Manifold

Head of Measures Development, National Peer Review Programme, NHS England

Mr Gavin Maxwell

Lay member

Ms Teresa Middleton

Deputy Director of Quality, NHS Gloucestershire Clinical Commissioning Group

Mrs Juliette Millard

UK Nursing and Health Professions Adviser, Leonard Cheshire Disability

Hazel Trender

Senior Vascular Nurse Specialist, Sheffield Teaching Hospital Trust

Dr Hugo van Woerden

Director of Public Health, NHS Highland

Dr Bee Wee (Chair)

Consultant and Senior Clinical Lecturer in Palliative Medicine, Oxford University Hospitals NHS Trust and Oxford University

Ms Alyson Whitmarsh

Programme Head for Clinical Audit, Health and Social Care Information Centre

Ms Jane Worsley

Chief Operating Officer, Options Group, Alcester Heath, Warwickshire

Dr Arnold Zermansky

GP, Leeds

The following specialist members joined the committee to develop this quality standard:

Dr Sam Oddie

Consultant Neonatologist, Bradford Royal infirmary

Mrs Jane Plumb

Lay member

Dr Meekai To

Consultant in Obstetrics and Fetal medicine, King's College Hospital

Louise Weaver-Lowe

Lead Nurse, Central Manchester

NICE project team

Nick Baillie

Associate Director

Karen Slade

Consultant Clinical Adviser

Alaster Rutherford

Clinical Adviser

Stephanie Birtles

Technical Adviser

Sabina Keane

Lead Technical Analyst

Esther Clifford

Programme Manager

Jenny Mills

Project Manager

Julia Sus

Coordinator

About this quality standard

NICE quality standards describe high-priority areas for quality improvement in a defined care or service area. Each standard consists of a prioritised set of specific, concise and measurable statements. NICE quality standards draw on existing NICE or NICE-accredited guidance that provides an underpinning, comprehensive set of recommendations, and are designed to support the measurement of improvement.

The methods and processes for developing NICE quality standards are described in the [quality standards process guide](#).

This quality standard has been incorporated into the NICE pathway on [preterm labour and birth](#).

NICE produces guidance, standards and information on commissioning and providing high-quality healthcare, social care and public health services. We have agreements to provide certain NICE services to Wales, Scotland and Northern Ireland. Decisions on how NICE guidance and other products apply in those countries are made by ministers in the Welsh government, Scottish government, and Northern Ireland Executive. NICE guidance or other products may include references to organisations or people responsible for commissioning or providing care that may be relevant only to England.

ISBN: 978-1-4731-2118-8

Endorsing organisation

This quality standard has been endorsed by NHS England, as required by the Health and Social Care Act (2012)

Supporting organisations

Many organisations share NICE's commitment to quality improvement using evidence-based guidance. The following supporting organisations have recognised the benefit of the quality standard in improving care for patients, carers, service users and members of the public. They have agreed to work with NICE to ensure that those commissioning or providing services are made aware of and encouraged to use the quality standard.

- [Royal College of General Practitioners](#)
- [Bliss](#)
- [Royal College of Midwives](#)

- Royal College of Obstetricians and Gynaecologists