

Hip fracture in adults

Quality standard

Published: 27 March 2012

[nice.org.uk/guidance/qs16](https://www.nice.org.uk/guidance/qs16)

Contents

Quality statements.....	4
Quality statement 1: Multidisciplinary management	6
Quality statement.....	6
Rationale	6
Quality measures	6
What the quality statement means for different audiences.....	7
Source guidance.....	7
Definitions of terms used in this quality statement	8
Quality statement 2: Timing and expertise for surgery.....	9
Quality statement.....	9
Rationale	9
Quality measures	9
What the quality statement means for different audiences	10
Source guidance.....	11
Definitions of terms used in this quality statement	11
Quality statement 3: Intracapsular fracture	12
Quality statement.....	12
Rationale	12
Quality measures	12
What the quality statement means for different audiences	13
Source guidance.....	13
Definitions of terms used in this quality statement	13
Quality statement 4: Trochanteric fracture	15
Quality statement.....	15
Rationale	15
Quality measures	15
What the quality statement means for different audiences.....	16

Source guidance.....	16
Definitions of terms used in this quality statement	16
Quality statement 5: Subtrochanteric fracture	18
Quality statement.....	18
Rationale	18
Quality measures	18
What the quality statement means for different audiences.....	19
Source guidance.....	19
Definitions of terms used in this quality statement	19
Quality statement 6: Rehabilitation after surgery	20
Quality statement.....	20
Rationale	20
Quality measures	20
What the quality statement means for different audiences.....	21
Source guidance.....	22
Definitions of terms used in this quality statement	22
Update information	23
About this quality standard.....	24
Improving outcomes	24
Resource impact.....	25
Diversity, equality and language	25

This standard is based on CG124.

This standard should be read in conjunction with QS3, QS15, QS86, QS63, QS149 and QS166.

Quality statements

Statement 1 Adults with hip fracture are cared for within a Hip Fracture Programme at every stage of the care pathway. [2012, updated 2016]

Statement 2 Adults with hip fracture have surgery on a planned trauma list on the day of, or the day after, admission. [2012, updated 2016]

Statement 3 Adults with displaced intracapsular hip fracture have a total hip replacement if clinically eligible, rather than hemiarthroplasty. [2012, updated 2017]

Statement 4 Adults with trochanteric fractures above and including the lesser trochanter receive extramedullary implants. [2012, updated 2016]

Statement 5 Adults with subtrochanteric fracture are treated with an intramedullary nail. [new 2016]

Statement 6 Adults with hip fracture start rehabilitation at least once a day, no later than the day after surgery. [2012, updated 2016]

In 2016 this quality standard was reviewed and statements prioritised in 2012 were updated (2012, updated 2016) or replaced (new 2016). Statement 3 was updated again in 2017 (2012, updated 2017). For more information, see [update information](#).

Statements from the 2012 quality standard for hip fracture that may still be useful at a local level, but are no longer considered national priorities for improvement:

- People with hip fracture have their cognitive status assessed, measured and recorded from admission.
- People with hip fracture receive prompt and effective pain management, in a manner that takes into account the hierarchy of pain management drugs, throughout their hospital stay.

- People with hip fracture are offered early supported discharge (if they are eligible), led by the Hip Fracture Programme team.
- People with hip fracture are offered a multifactorial risk assessment to identify and address future falls risk, and are offered individualised intervention if appropriate.
- People with hip fracture are offered a bone health assessment to identify future fracture risk and offered pharmacological intervention as needed before discharge from hospital.

The [2012 quality standard for hip fracture](#) is available as a pdf.

NICE has developed guidance and a quality standard on patient experience in adult NHS services (see the NICE pathway on [patient experience in adult NHS services](#)), which should be considered alongside these quality statements.

Other quality standards that should be considered when commissioning or providing services for people with hip fracture include:

- [Osteoporosis](#) (2017) NICE quality standard 149
- [Falls in older people](#) (2015, updated 2017) NICE quality standard 86
- [Delirium in adults](#) (2014) NICE quality standard 63

A full list of NICE quality standards is available from the [quality standards topic library](#).

Quality statement 1: Multidisciplinary management

Quality statement

Adults with hip fracture are cared for within a Hip Fracture Programme at every stage of the care pathway. [2012, updated 2016]

Rationale

People with hip fracture, including those cared for in the community, often have comorbidities and complex care needs. The multidisciplinary approach of a Hip Fracture Programme, with regular assessment and continuous rehabilitation, has been found to better meet those needs, and lead to improved functional outcomes and reduced mortality.

Quality measures

Structure

Evidence of local arrangements to ensure that people with hip fracture are cared for within a Hip Fracture Programme at every stage of the care pathway.

Data source: Local data collection.

Process

a) Proportion of presentations of hip fracture in which the person receives an orthogeriatric assessment prior to surgery.

Numerator – the number in the denominator in which the person receives an orthogeriatric assessment prior to surgery.

Denominator – the number of presentations of hip fracture.

Data source: Local data collection. The [National Hip Fracture Database](#) records access to orthogeriatric assessment.

b) Proportion of presentations of hip fracture in which the person has their goals for multidisciplinary rehabilitation identified.

Numerator – the number in the denominator in which the person has their goals for multidisciplinary rehabilitation identified.

Denominator – the number of people having surgery for hip fracture.

Data source: Local data collection.

Outcome

a) Mortality for people with hip fracture at discharge.

Data source: Local data collection.

b) Functional outcome at 1 year.

Data source: Local data collection.

What the quality statement means for different audiences

Service providers (such as hospitals) have systems in place to ensure that people with hip fracture are cared for within a Hip Fracture Programme at every stage of the care pathway.

Commissioners (such as clinical commissioning groups) ensure that they commission hip fracture services that provide care within a Hip Fracture Programme at every stage of the care pathway.

People with hip fracture are looked after within a programme of care, called a Hip Fracture Programme. This involves a team of healthcare professionals with different skills working together to provide care. Hip Fracture Programmes provide care at every stage, in hospital and at home, which includes regular assessment, and coordination of care and rehabilitation.

Source guidance

[Hip fracture: management](#) (2011) NICE guideline CG124, recommendation 1.8.1 (key priority for implementation)

Definitions of terms used in this quality statement

Hip Fracture Programme

A coordinated multidisciplinary approach ensuring continuity of care and responsibility across the clinical pathway. It covers care in all settings, including ambulances, A&E departments, radiology, operating theatres, wards and in the community and primary care, and at all stages, including diagnosis, treatment, recovery, discharge planning, rehabilitation, long-term after care and secondary prevention.

It involves formal 'orthogeriatric' care, with the geriatric medical team contributing to joint preoperative patient assessment, and increasingly taking the lead in postoperative medical care, multidisciplinary rehabilitation and discharge planning.

It includes all of the following:

- orthogeriatric assessment
- rapid optimisation of fitness for surgery
- early identification of individual goals for multidisciplinary rehabilitation to recover mobility and independence, and to facilitate return to pre-fracture residence and long-term wellbeing
- continued, coordinated, orthogeriatric and multidisciplinary review
- liaison or integration with related services, particularly mental health, falls prevention, bone health, primary care and social services
- clinical and service governance responsibility for all stages of the pathway of care and rehabilitation, including those delivered in the community.

[Adapted from NICE's guideline on [hip fracture](#), recommendation 1.8.1 and expert consensus]

Quality statement 2: Timing and expertise for surgery

Quality statement

Adults with hip fracture have surgery on a planned trauma list on the day of, or the day after, admission. [2012, updated 2016]

Rationale

People with hip fracture can experience pain and anxiety while waiting for an operation. Delays in surgery are associated with negative outcomes for mortality and return to mobility. Therefore, it is important to avoid any unnecessary delays for people who are assessed as fit for surgery. A planned trauma list includes specific healthcare professionals with the expertise required for hip surgery. Senior staff supervision can help to reduce the risk of complications during the surgery.

Quality measures

Structure

a) Evidence of local arrangements to ensure that people with hip fracture have surgery on a planned trauma list.

Data source: Local data collection.

b) Evidence of local arrangements to ensure that people with hip fracture have surgery on the day of, or the day after, admission.

Data source: Local data collection.

Process

a) Proportion of operations for hip fracture that are performed on a planned trauma list.

Numerator – the number in the denominator that are performed on a planned trauma list.

Denominator – the number of operations for hip fracture.

Data source: Local data collection. The [National Hip Fracture Database](#) records the time of the operation in relation to the admission.

b) Proportion of operations for hip fracture that are performed on the day of, or the day after, admission.

Numerator – the number in the denominator that are performed on the day of, or the day after, admission.

Denominator – the number of operations for hip fracture.

Data source: Local data collection. The NHS Digital [Compendium of Clinical and Health Indicators](#) records emergency hospital admissions and timely surgery: fractured proximal femur. The [National Hip Fracture Database](#) records the time of the operation in relation to the admission.

Outcome

a) Postoperative complications for people with hip fracture.

Data source: Local data collection.

b) Length of hospital stay for people with hip fracture.

Data source: Local data collection.

c) Mortality for people having hip fracture surgery.

Data source: Local data collection.

What the quality statement means for different audiences

Service providers (hospitals) ensure that systems are in place for people with hip fracture to have surgery on a planned trauma list on the day of, or the day after, admission.

Healthcare professionals (such as specialists, orthogeriatricians and anaesthetists) perform hip fracture surgery on a planned trauma list on the day of, or the day after, admission.

Commissioners (such as clinical commissioning groups) ensure that they commission services that have sufficient capacity for people with hip fracture to have surgery on a planned trauma list on the day of, or the day after, admission.

People with hip fracture have an operation carried out by a team of senior specialists on the day they are admitted to hospital or the next day.

Source guidance

Hip fracture: management (2011) NICE guideline CG124, recommendations 1.2.1 (key priority for implementation) and 1.5.1

Definitions of terms used in this quality statement

Planned trauma list

A planned trauma list is one with a rostered senior anaesthetist, senior surgeon and dedicated theatre time.

[NICE's guideline on hip fracture, full guideline]

Quality statement 3: Intracapsular fracture

Quality statement

Adults with displaced intracapsular hip fracture have a total hip replacement if clinically eligible, rather than hemiarthroplasty. [2012, updated 2017]

Rationale

Many people with hip fracture are concerned with recovery and getting back to usual activity. Total hip replacement is associated with improved functional status compared with hemiarthroplasty, if the patient was active before the fracture, is not cognitively impaired and is medically fit for the procedure.

Quality measures

Structure

Evidence of local arrangements to perform total hip replacement if people with displaced intracapsular fracture are clinically eligible, rather than hemiarthroplasty.

Data source: Local data collection.

Process

Proportion of people with displaced intracapsular fracture and eligible for total hip replacement who receive total hip replacement.

Numerator – the number in the denominator who receive total hip replacement.

Denominator – the number of people with displaced intracapsular fracture and eligible for total hip replacement.

Data source: Local data collection. The [National Hip Fracture Database](#) records procedure type for intracapsular displaced fracture and cementing of arthroplasties.

Outcome

Functional status at 1 and 5 years after total hip replacement for displaced intracapsular fracture.

Data source:Local data collection.

What the quality statement means for different audiences

Service providers (hospitals) ensure that total hip replacement is performed for displaced intracapsular fracture, rather than hemiarthroplasty, if the person is assessed as clinically eligible.

Healthcare professionals (orthopaedic surgeons) treat displaced intracapsular fracture using total hip replacement, rather than hemiarthroplasty, if the person is assessed as clinically eligible.

Commissioners (clinical commissioning groups) ensure that orthopaedic services perform total hip replacement, rather than hemiarthroplasty, to treat displaced intracapsular fracture in people who are clinically eligible.

People with a fracture inside the socket of their hip joint and where the bones have moved out of position (called a displaced intracapsular fracture) who were fit and active before the fracture have both parts of the hip joint (the ball and socket) replaced with artificial parts (called a total hip replacement).

Source guidance

[Hip fracture: management](#) (2011) NICE guideline CG124, recommendations 1.6.2 and 1.6.3 (key priorities for implementation)

Definitions of terms used in this quality statement

Intracapsular fractures

Fractures above the insertion of the capsular attachment of the hip joint are called intracapsular.

[NICE's guideline on [hip fracture](#), full guideline]

Clinical eligibility for total hip replacement

Patients with a displaced intracapsular fracture who:

- were able to walk independently out of doors with no more than the use of a stick and
- are not cognitively impaired and

- are medically fit for anaesthesia and the procedure.

[NICE's guideline on [hip fracture](#), recommendation 1.6.3]

Quality statement 4: Trochanteric fracture

Quality statement

Adults with trochanteric fractures above and including the lesser trochanter receive extramedullary implants. [2012, updated 2016]

Rationale

Extramedullary implants, such as sliding hip screws, have similar clinical outcomes to intramedullary devices. However, some studies have shown that intramedullary implants have a higher reoperation rate because of periprosthetic fracture. In addition, extramedullary implants are less expensive than intramedullary implants. Therefore, extramedullary implants are recommended in preference to intramedullary nails for the treatment of trochanteric fractures.

Quality measures

Structure

Evidence of local arrangements to ensure that people with trochanteric fractures above and including the lesser trochanter receive extramedullary implants.

Data source: Local data collection.

Process

Proportion of presentations of trochanteric fractures above and including the lesser trochanter in which the person receives extramedullary implants.

Numerator – the number in the denominator in which the person receives extramedullary implants.

Denominator – the number of presentations of trochanteric fractures above and including the lesser trochanter.

Data source: Local data collection. The [National Hip Fracture Database](#) records procedure type for trochanteric fracture.

Outcome

Reoperation rates for people with trochanteric fractures.

Data source: Local data collection.

What the quality statement means for different audiences

Service providers (hospitals) ensure that systems are in place for people with trochanteric fractures above and including the lesser trochanter to receive extramedullary implants in preference to intramedullary nails.

Healthcare professionals (orthopaedic surgeons) ensure that people with trochanteric fractures above and including the lesser trochanter receive extramedullary implants in preference to intramedullary nails.

Commissioners (such as clinical commissioning groups) ensure that they commission services where people with trochanteric fractures above and including the lesser trochanter receive extramedullary implants in preference to intramedullary nails.

People with a fracture outside the socket of their hip joint and near the top of the thigh bone (called a trochanteric fracture) have an operation to reposition the broken bone and hold it in place while it heals. This is done using one or more special screws inserted into the bone and attached to a metal plate (called an extramedullary implant).

Source guidance

[Hip fracture: management](#) (2011) NICE guideline CG124, recommendation 1.6.7 (key priorities for implementation)

Definitions of terms used in this quality statement

Trochanteric fractures

Fractures that occur outside or distal to the hip joint capsule, which can be two-part fractures (stable) or multi-fragmentary (unstable).

[NICE's guideline on [hip fracture](#), full guideline]

Extramedullary implants

A screw that is attached to a plate on the outside of the femoral head and neck.

[NICE's guideline on [hip fracture](#), full guideline]

Quality statement 5: Subtrochanteric fracture

Quality statement

Adults with subtrochanteric fracture are treated with an intramedullary nail. [new 2016]

Rationale

Using an intramedullary device can provide mechanical protection to a potentially diseased bone. Intramedullary fixation is the treatment of choice for subtrochanteric fractures because it allows splinting of the whole of the femoral shaft. Although intramedullary nails are more expensive than extramedullary implants, they lead to fewer patients with non-union of fracture needing reoperation.

Quality measures

Structure

Evidence of local arrangements to ensure that people with subtrochanteric fracture are treated with an intramedullary nail.

Data source: Local data collection.

Process

Proportion of presentations of subtrochanteric fractures treated with an intramedullary nail.

Numerator – the number in the denominator that are treated with an intramedullary nail.

Denominator – the number of presentations of subtrochanteric fractures.

Data source: Local data collection.

Outcome

a) Number of people with non-union of fracture.

Data source: Local data collection.

What the quality statement means for different audiences

Service providers (hospitals) ensure that systems are in place for people with subtrochanteric fractures to be treated with an intramedullary nail.

Healthcare professionals (orthopaedic surgeons) perform surgery on people with subtrochanteric fractures using an intramedullary nail.

Commissioners (such as clinical commissioning groups) ensure that they commission services where people with subtrochanteric fractures are treated with an intramedullary nail.

People with a fracture outside the socket of their hip joint and a small way down the thigh bone (called a subtrochanteric fracture) have an operation to reposition the broken bone and hold it in place while it heals. This is done using a metal rod, called an intramedullary nail, which is inserted into the bone.

Source guidance

[Hip fracture: management](#) (2011) NICE guideline CG124, recommendation 1.6.8

Definitions of terms used in this quality statement

Subtrochanteric fracture

The fracture is predominantly in the 5 cm of bone immediately distal to the lesser trochanter.

[NICE's guideline on [hip fracture](#), full guideline]

Intramedullary nail

A metal rod, which is inserted down the middle of the femoral shaft.

[NICE's guideline on [hip fracture](#), full guideline]

Quality statement 6: Rehabilitation after surgery

Quality statement

Adults with hip fracture start rehabilitation at least once a day, no later than the day after surgery.
[2012, updated 2016]

Rationale

Early restoration of mobility after hip fracture surgery can be beneficial for the person because it can reduce the length of hospital stay and avoid the complications of prolonged bed confinement. Rehabilitation at least once a day has potential benefits of improved mobility, increased independence, and reduced need for institutional care. A physiotherapist assessment is needed before the rehabilitation starts. People should be offered support with rehabilitation every day while in hospital, which can be given by members of the multidisciplinary team when the physiotherapist is not present. This support should continue after discharge from hospital.

Quality measures

Structure

Evidence of local arrangements to ensure that people with hip fracture start rehabilitation at least once a day, no later than the day after surgery.

Data source: Local data collection.

Process

a) Proportion of hip fracture operations after which the person starts rehabilitation no later than the day after surgery.

Numerator – the number in the denominator after which the person starts rehabilitation no later than the day after surgery.

Denominator – the number of hip fracture operations.

Data source: Local data collection. The [National Hip Fracture Database](#) records if the patient was mobilised on the day after surgery.

b) Proportion of hip fracture operations after which the person has rehabilitation at least once a day.

Numerator – the number in the denominator after which the person has rehabilitation at least once a day.

Denominator – the number of hip fracture operations.

Data source: Local data collection.

Outcome

a) Length of hospital stay for people with hip fracture.

Data source: Local data collection.

b) Return to the pre-hip fracture place of residence.

Data source: Local data collection.

c) Return to the pre-hip fracture level of mobility.

Data source: Local data collection. The [National Hip Fracture Database](#) records the routine follow-up of hip fracture patients.

What the quality statement means for different audiences

Service providers (such as hospitals) ensure that systems are in place for people with hip fracture to start rehabilitation at least once a day, no later than the day after surgery.

Healthcare professionals (such as physiotherapists and nurses) offer rehabilitation at least once a day to people with hip fracture, starting no later than the day after surgery.

Commissioners (such as clinical commissioning groups) ensure that they commission services in which people with hip fracture start rehabilitation at least once a day, no later than the day after surgery.

People who have had an operation for hip fracture are offered rehabilitation at least once a day to

help them recover. Rehabilitation should be started by the day after their operation (unless there is a medical or surgical reason not to). Rehabilitation after a hip fracture operation includes support with sitting and standing and keeping an upright posture to improve movement and strength, and help with their recovery.

Source guidance

Hip fracture: management (2011) NICE guideline CG124, recommendations 1.7.1 and 1.7.2

Definitions of terms used in this quality statement

Rehabilitation

Rehabilitation is the process of re-establishing the ability to move between postures (for example, from sitting to standing), maintain an upright posture and to ambulate with increasing levels of complexity (speed, changes of direction, dual and multi-tasking).

[Adapted from NICE's guideline on hip fracture, full guideline and expert input]

Update information

May 2017: Statement 3 on intracapsular fracture has been updated to reflect changes to recommendations 1.6.2 and 1.6.3 in the NICE guideline on [hip fracture](#). This is marked as [2012, updated 2017].

November 2016: This quality standard was updated and statements prioritised in 2012 were replaced.

Statements are marked as:

- [new 2016] if the statement covers a new area for quality improvement
- [2012, updated 2016] if the statement covers an area for quality improvement included in the 2012 quality standard and has been updated.

Statements numbered 1, 2, 5, 6, 7, 8 and 9 in the 2012 version have been updated and are marked as [2012, updated 2016].

Statements from the 2012 version (numbered 3, 4, 10, 11 and 12) that are no longer considered national priorities for improvement, but may still be useful at a local level, are listed after the updated statements in [quality statements](#).

[The 2012 quality standard for hip fracture](#) is available as a pdf.

About this quality standard

NICE quality standards describe high-priority areas for quality improvement in a defined care or service area. Each standard consists of a prioritised set of specific, concise and measurable statements. NICE quality standards draw on existing NICE or NICE-accredited guidance that provides an underpinning, comprehensive set of recommendations, and are designed to support the measurement of improvement.

Information about how NICE quality standards are developed is available from [developing NICE quality standards](#) on the website.

See [quality standard advisory committees](#) on the website for details of standing committee 4 members who advised on this quality standard. Information about the topic experts invited to join the standing members is available on the [quality standard's webpage](#).

This quality standard has been incorporated into the NICE pathway on [hip fracture](#).

NICE produces guidance, standards and information on commissioning and providing high-quality healthcare, social care, and public health services. We have agreements to provide certain NICE services to Wales, Scotland and Northern Ireland. Decisions on how NICE guidance and other products apply in those countries are made by ministers in the Welsh government, Scottish government, and Northern Ireland Executive. NICE guidance or other products may include references to organisations or people responsible for commissioning or providing care that may be relevant only to England.

Improving outcomes

The quality standard is expected to contribute to improvements in the following outcomes:

- length of hospital stay
- readmission to hospital
- reoperation rates
- mortality rates.

It is also expected to support delivery of the Department of Health's outcome frameworks:

- [NHS outcomes framework 2016–17](#)

- [Adult social care outcomes framework 2015–16](#)
- [Public health outcomes framework for England 2016–19](#).

Resource impact

NICE quality standards should be achievable by local services. The potential resource impact is considered by the quality standards advisory committee, drawing on resource impact work for the source guidance. Organisations are encouraged to use the [costing report and template](#) for the NICE guideline on hip fracture to help estimate local costs.

Diversity, equality and language

During the development of this quality standard, equality issues have been considered and [equality assessments](#) are available.

Good communication between health, public health and social care practitioners and people with hip fracture is essential. Treatment, care and support, and the information given about it, should be culturally appropriate. It should also be accessible to people with additional needs such as physical, sensory or learning disabilities, and to people who do not speak or read English. People with hip fracture in hospital should have access to an interpreter or advocate if needed.

Commissioners and providers should aim to achieve the quality standard in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this quality standard should be interpreted in a way that would be inconsistent with compliance with those duties.

ISBN: 978-1-4731-2016-7

Endorsing organisation

This quality standard has been endorsed by NHS England, as required by the Health and Social Care Act (2012)

Supporting organisations

Many organisations share NICE's commitment to quality improvement using evidence-based guidance. The following supporting organisations have recognised the benefit of the quality standard in improving care for patients, carers, service users and members of the public. They have

agreed to work with NICE to ensure that those commissioning or providing services are made aware of and encouraged to use the quality standard.

- [British Orthopaedic Association](#)
- [Chartered Society of Physiotherapy](#)
- [Royal College of General Practitioners](#)
- [Royal College of Physicians](#)