

# Familial hypercholesterolaemia

Quality standard

Published: 13 August 2013

[www.nice.org.uk/guidance/qs41](http://www.nice.org.uk/guidance/qs41)

## Contents

Introduction .....	6
Why this quality standard is needed .....	6
How this quality standard supports delivery of outcome frameworks .....	7
Coordinated services.....	8
Training and competencies.....	8
List of quality statements.....	9
Quality statement 1: Diagnosis .....	10
Quality statement.....	10
Rationale.....	10
Quality measures .....	10
What the quality statement means for service providers, general practitioners, and commissioners .....	11
What the quality statement means for patients, service users and carers.....	11
Source guidance.....	11
Definitions of terms used in this quality statement .....	11
Equality and diversity considerations.....	12
Quality statement 2: Specialist referral .....	13
Quality statement.....	13
Rationale.....	13
Quality measures .....	13
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	14
What the quality statement means for patients, service users and carers.....	14
Source guidance.....	14
Definitions of terms used in this quality statement .....	14
Quality statement 3: DNA testing.....	16
Quality statement.....	16
Rationale.....	16
Quality measures .....	16

What the quality statement means for service providers, healthcare practitioners, and commissioners ..	17
What the quality statement means for patients, service users and carers.....	17
Source guidance.....	17
Definitions of terms used in this quality statement .....	17
<b>Quality statement 4: Diagnosis in children under 10 years .....</b>	<b>19</b>
Quality statement.....	19
Rationale.....	19
Quality measures .....	19
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	20
What the quality statement means for patients, service users and carers.....	20
Source guidance.....	20
Definitions of terms used in this quality statement .....	20
<b>Quality statement 5: Cascade testing.....</b>	<b>22</b>
Quality statement.....	22
Rationale.....	22
Quality measures .....	22
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	23
What the quality statement means for patients, service users and carers.....	24
Source guidance.....	24
Definitions of terms used in this quality statement .....	24
<b>Quality statement 6: Drug treatment in adults.....</b>	<b>26</b>
Quality statement.....	26
Rationale.....	26
Quality measures .....	26
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	27
What the quality statement means for patients, service users and carers.....	27
Source guidance.....	27
Definitions of terms used in this quality statement .....	27

---

Equality and diversity considerations.....	28
Quality statement 7: Drug treatment in children .....	29
Quality statement.....	29
Rationale.....	29
Quality measures .....	29
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	29
What the quality statement means for patients, service users and carers.....	30
Source guidance.....	30
Definitions of terms used in this quality statement .....	30
Equality and diversity considerations.....	31
Quality statement 8: Annual review.....	32
Quality statement.....	32
Rationale.....	32
Quality measures .....	32
What the quality statement means for service providers, healthcare practitioners, and commissioners ..	32
What the quality statement means for patients, service users and carers.....	33
Source guidance.....	33
Definitions of terms used in this quality statement .....	33
Using the quality standard.....	35
Quality measures .....	35
Levels of achievement .....	35
Using other national guidance and policy documents.....	35
Diversity, equality and language .....	36
Development sources.....	37
Evidence sources.....	37
Policy context .....	37
Related NICE quality standards .....	38
Topic Expert Group and NICE project team .....	39

Topic Expert Group.....	39
NICE project team .....	40
Update information.....	41
About this quality standard.....	42

This standard is based on CG71.

This standard should be read in conjunction with QS15, QS43, QS100, QS99, QS140, QS120, QS111, QS94 and QS84.

## Introduction

This quality standard covers the identification and management of heterozygous familial hypercholesterolaemia (FH) in adults, young people and children. Homozygous FH has been excluded from this quality standard because it has a low incidence and people with homozygous FH need specialist care. Please see the NICE guideline on [familial hypercholesterolaemia](#) for recommendations on the diagnosis and treatment of homozygous FH. For more information see the [scope](#) of the quality standard.

### *Why this quality standard is needed*

FH is an inherited condition caused by an alteration in a gene, which results in a high cholesterol concentration in the blood. Raised cholesterol concentrations are present from birth and lead to early development of atherosclerosis and coronary heart disease. The condition is transmitted from generation to generation in such a way that siblings and children of a person with FH have a 1 in 2 chance (50:50 risk) of also having FH.

Most people with FH have inherited an altered gene for FH in an autosomal dominant pattern from only 1 parent and are therefore 'heterozygous'. Occasionally, a person will inherit an altered gene from both parents and will have 'homozygous' FH or 'compound heterozygous' FH. Homozygous FH is rare, with an incidence of approximately one in a million.

The prevalence of heterozygous FH in the UK population is estimated to be 1 in 500, which means that approximately 120,000 people are expected to be affected<sup>[1]</sup>. However, more than 80% of these are currently undiagnosed and untreated. If left untreated, more than 50% of men with heterozygous FH will develop coronary heart disease by the age of 50 years and more than 50% of women by the age of 60 years<sup>[2]</sup>. Life expectancy is restored to near normal with early preventive treatment, particularly statin treatment and smoking cessation.

The importance of better identification of families/individuals at very high risk of cardiovascular disease, including those with FH, is recognised in the Department of Health's [Cardiovascular Disease Outcomes Strategy](#).

## How this quality standard supports delivery of outcome frameworks

NICE quality standards are a concise set of prioritised statements designed to drive measureable quality improvements within a particular area of health or care. They are derived from high-quality guidance, such as that from NICE or other sources accredited by NICE. This quality standard, in conjunction with the guidance on which it is based, should contribute to the improvements outlined in the following outcomes framework published by the Department of Health:

- [NHS Outcomes Framework 2013/14](#) (Department of Health, November 2012)

Table 1 shows the outcomes, overarching indicators and improvement areas from the framework that the quality standard could contribute to achieving.

**Table 1** [NHS Outcomes Framework 2013/14](#)

Domain	Overarching indicators and improvement areas
1 Preventing people from dying prematurely	<p><b>Overarching indicator</b></p> <p>1a Potential years of life lost (PYLL) from causes considered amenable to healthcare</p> <p>i Adults ii Children and young people</p> <p>1b Life expectancy at 75</p> <p>i Males ii Females</p>
2 Enhancing quality of life for people with long-term conditions	<p><b>Overarching indicator</b></p> <p>2 Health-related quality of life for people with long-term conditions*</p> <p><b>Improvement areas</b></p> <p>Ensuring people feel supported to manage their condition</p> <p>2.1 Proportion of people feeling supported to manage their condition*</p>

<p>4 Ensuring people have a positive experience of care</p>	<p><b>Overarching indicator</b></p> <p>4a Patient experience of primary care i) GP services</p> <p><b>Improvement areas</b></p> <p>Improving people's experience of outpatient care</p> <p>4.1 Patient experience of outpatient care</p>
<p><b>Alignment across the health and social care system</b></p> <p>*Indicator complementary with Adult Social Care Outcomes Framework (ASCOF).</p>	

### Coordinated services

The quality standard for FH specifies that services should be commissioned from and coordinated across all relevant agencies encompassing the whole FH care pathway. A person-centred, integrated approach to providing services is fundamental to delivering high-quality care to people with FH.

The Health and Social Care Act 2012 sets out a clear expectation that the care system should consider NICE quality standards in planning and delivering services, as part of a general duty to secure continuous improvement in quality. Commissioners and providers of health and social care should refer to the library of NICE quality standards when designing high-quality services. Other quality standards that should also be considered when choosing, commissioning or providing a high-quality FH service are listed in [related quality standards](#).

### Training and competencies

The quality standard should be read in the context of national and local guidelines on training and competencies. All healthcare practitioners involved in assessing, caring for and treating people with FH should have sufficient and appropriate training and competencies to deliver the actions and interventions described in the quality standard.

<sup>[1]</sup> Seed M, Roughton M, Pedersen K et al. (2012) [Current statin treatment, DNA testing and cascade testing of UK patients with familial hypercholesterolaemia](#). Primary Care Cardiovascular Journal 5: 181–5.

<sup>[2]</sup> Slack J (1969) Risks of ischaemic heart-disease in familial hyperlipoproteinaemia states. Lancet 2: 1380–2.



## List of quality statements

Statement 1 Adults with a baseline total cholesterol above 7.5 mmol/l are assessed for a clinical diagnosis of familial hypercholesterolaemia (FH).

Statement 2 People with a clinical diagnosis of FH are referred for specialist assessment.

Statement 3 People with a clinical diagnosis of FH are offered DNA testing as part of a specialist assessment.

Statement 4 Children at risk of FH are offered diagnostic tests by the age of 10 years.

Statement 5 Relatives of people with a confirmed diagnosis of monogenic FH are offered DNA testing through a nationwide, systematic cascade process.

Statement 6 Adults with FH receive lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

Statement 7 Children with FH are assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting by the age of 10 years.

Statement 8 People with FH are offered a structured review at least annually.

## Quality statement 1: Diagnosis

### *Quality statement*

Adults with a baseline total cholesterol above 7.5 mmol/l are assessed for a clinical diagnosis of familial hypercholesterolaemia (FH).

### *Rationale*

Most of the 120,000 people estimated to have FH are undiagnosed and untreated. Because untreated FH carries a very high risk of cardiovascular disease, it is important that every opportunity is taken to identify people with FH and offer them treatment. Considering a clinical diagnosis of FH in people with high cholesterol will result in greater identification of FH and support cascade testing of their relatives. This will lead to more treatment to reduce cholesterol levels and prevention of coronary events among people with FH.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that adults with a baseline total cholesterol above 7.5 mmol/l are assessed for a clinical diagnosis of FH.

**Data source:** Local data collection.

#### **Process**

Proportion of adults with a baseline total cholesterol above 7.5 mmol/l who are assessed for a clinical diagnosis of FH.

Numerator – The number of people in the denominator assessed for a clinical diagnosis of FH.

Denominator – The number of adults with a baseline total cholesterol above 7.5 mmol/l.

**Data source:** Local data collection.

## *What the quality statement means for service providers, general practitioners, and commissioners*

**Service providers** ensure that systems are in place for adults with a baseline total cholesterol above 7.5 mmol/l to be assessed for a clinical diagnosis of FH.

**General practitioners** assess adults with a baseline total cholesterol above 7.5 mmol/l for a clinical diagnosis of FH.

**Commissioners** ensure that they commission services that identify and assess adults with a baseline total cholesterol above 7.5 mmol/l for a clinical diagnosis of FH.

## *What the quality statement means for patients, service users and carers*

Adults with a total cholesterol above 7.5 mmol/l before treatment have an assessment for FH.

## *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendations 1.1.1, 1.1.2, 1.1.4, 1.1.5 and 1.1.9

## *Definitions of terms used in this quality statement*

### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

### **Adults**

People aged 16 and older.

### **Baseline total cholesterol**

The total cholesterol concentration before treatment.

### **Assessment for a clinical diagnosis of FH**

NICE's guideline on familial hypercholesterolaemia recommends assessment for a clinical diagnosis of FH using all 3 criteria below:

- exclusion of secondary causes of hypercholesterolaemia
- 2 measurements of LDL-C concentration
- assessment against Simon Broome criteria or Dutch Lipid Clinic Network (DLCN) criteria to make a clinical diagnosis of FH in primary care settings.

### *Equality and diversity considerations*

The statement has been restricted to adults because the criteria for assessment for a clinical diagnosis of FH are not appropriate for children and young people under 16 years.

## Quality statement 2: Specialist referral

### *Quality statement*

People with a clinical diagnosis of familial hypercholesterolaemia (FH) are referred for specialist assessment.

### *Rationale*

Diagnosing and managing FH in an individual and their relatives can be complex, and is best achieved when there is access to specialist services. Specialist assessments, which include DNA testing, can confirm a diagnosis. Once an accurate diagnosis has been made, people with FH can receive appropriate treatment, and cascade testing can be started to identify affected family members.

### *Quality measures*

#### **Structure**

a) Evidence of local arrangements to ensure that people with a clinical diagnosis of FH are referred for specialist assessment.

**Data source:** Local data collection.

b) Evidence of local arrangements to ensure that a protocol for referral for a specialist assessment is agreed between primary and secondary care.

**Data source:** Local data collection.

#### **Process**

Proportion of people with a clinical diagnosis of FH referred for specialist assessment.

Numerator – The number of people in the denominator referred for specialist assessment.

Denominator – The number of people with a clinical diagnosis of FH.

**Data source:** Local data collection.

## Outcome

Ratio of observed to estimated numbers of people with FH, using an estimate based on the area's estimated prevalence of FH (based on 1 in 500) and population size.

*Data source:* Local data collection using a dedicated database.

### *What the quality statement means for service providers, healthcare practitioners, and commissioners*

Service providers ensure that systems are in place for people with a clinical diagnosis of FH to be referred for specialist assessment.

Healthcare practitioners refer people with a clinical diagnosis of FH for specialist assessment.

Commissioners ensure that they commission services that can offer specialist assessment for people with a clinical diagnosis of FH.

### *What the quality statement means for patients, service users and carers*

People who are given a clinical diagnosis of FH because they have high cholesterol and family history or other signs are referred for specialist assessment.

## Source guidance

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendations 1.2.2 and 1.3.1.17

## Definitions of terms used in this quality statement

### Familial hypercholesterolaemia (FH)

FH relates to heterozygous FH only.

### Assessment for a clinical diagnosis of FH

NICE's guideline on familial hypercholesterolaemia recommends assessment for a clinical diagnosis of FH using all 3 criteria below:

- exclusion of secondary causes of hypercholesterolaemia
- 2 measurements of LDL-C concentration
- assessment against Simon Broome criteria or Dutch Lipid Clinic Network (DLCN) criteria to make a clinical diagnosis of FH in primary care settings.

## Specialist assessment

This should include:

- confirmation of the clinical diagnosis of FH made by GP or other healthcare professional
- an offer of DNA testing to increase the certainty of the diagnosis
- initiation of cascade testing if a diagnosis is confirmed.

A specialist assessment should be performed by a healthcare professional with expertise in FH who has access to the wider skills of a multidisciplinary team. This team should include a dietitian, cardiologist and paediatrician, and a clinical genetic specialist to take a family history and obtain informed consent for a DNA test. For children and young people, this should be a specialist with expertise in FH in children and young people.

Children refers to people younger than 10, young people refers to those aged 10 up to and including age 15, and adults refers to people aged 16 and older.

## Quality statement 3: DNA testing

### *Quality statement*

People with a clinical diagnosis of familial hypercholesterolaemia (FH) are offered DNA testing as part of a specialist assessment.

### *Rationale*

DNA testing is important because it increases the certainty of a diagnosis of FH and allows the identification of affected and unaffected relatives through cascade testing.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that people with a clinical diagnosis of FH are offered DNA testing as part of a specialist assessment.

*Data source:* Local data collection.

#### **Process**

a) Proportion of people with a clinical diagnosis of FH who receive DNA testing as part of a specialist assessment.

Numerator – The number of people in the denominator receiving DNA testing as part of a specialist assessment.

Denominator – The number of people with a clinical diagnosis of FH.

*Data source:* Local data collection using a dedicated database.

b) Proportion of people with a clinical diagnosis of FH receiving DNA testing as part of a specialist assessment who give informed consent for the test.

Numerator – The number of people in the denominator who give informed consent for the test.

Denominator – The number of people with a clinical diagnosis of FH receiving DNA testing as part



of a specialist assessment.

*Data source:* Local data collection using a dedicated database.

## Outcome

Patient satisfaction with process of informed consent.

*Data source:* Local data collection.

### *What the quality statement means for service providers, healthcare practitioners, and commissioners*

Service providers ensure that systems are in place for people with a clinical diagnosis of FH to be offered DNA testing as part of a specialist assessment.

Specialists with expertise in FH offer DNA testing to people with a clinical diagnosis of FH as part of a specialist assessment.

Commissioners ensure that they commission services that offer DNA testing to people with a clinical diagnosis of FH as part of a specialist assessment.

### *What the quality statement means for patients, service users and carers*

People who are given a clinical diagnosis of FH because they have high cholesterol and family history or other signs are offered DNA testing as part of a specialist assessment.

## Source guidance

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendation 1.1.6

## Definitions of terms used in this quality statement

### Familial hypercholesterolaemia (FH)

FH relates to heterozygous FH only.

## Assessment for a clinical diagnosis of FH

NICE's guideline on [familial hypercholesterolaemia](#) recommends assessment for a clinical diagnosis of FH using all 3 criteria below:

- exclusion of secondary causes of hypercholesterolaemia
- 2 measurements of LDL-C concentration
- assessment against Simon Broome or Dutch Lipid Clinic Network (DLCN) criteria to make a clinical diagnosis of FH in primary care settings.

## DNA testing

DNA testing should test for all gene mutations known to cause FH. Methods should meet the standards set out by the [UK Genetic Testing Network](#). Informed consent should be given for DNA testing.

## Quality statement 4: Diagnosis in children under 10 years

### *Quality statement*

Children at risk of familial hypercholesterolaemia (FH) are offered diagnostic tests by the age of 10 years.

### *Rationale*

Children with FH begin to develop cardiovascular disease before clinical signs appear, with thickening of the carotid artery wall identifiable by the age of 10 years. Diagnosis by the age of 10 years allows lifestyle changes and tailored therapy if indicated, which will reduce long-term problems associated with high cholesterol and improve long-term health.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that children at risk of FH are offered diagnostic tests by the age of 10 years.

*Data source:* Local data collection.

#### **Process**

Proportion of children at risk of FH who receive a specified diagnostic test by the age of 10 years.

Numerator – The number of people in the denominator who had received a specified diagnostic test.

Denominator – The number of children aged 10 years at risk of FH.

*Data source:* Local data collection.

#### **Outcome**

Ratio of observed to estimated numbers of children at risk of FH, using an estimate based on the area's estimated prevalence of FH (based on 1 in 500) and population size.

*Data source:* Local data collection.

### *What the quality statement means for service providers, healthcare practitioners, and commissioners*

Service providers ensure that systems are in place for children at risk of FH to be offered diagnostic tests by the age of 10 years.

Specialist with expertise in FH in children and young people offer children at risk of FH diagnostic tests by the age of 10 years.

Commissioners ensure that they commission services that offer children at risk of FH diagnostic tests by the age of 10 years.

### *What the quality statement means for patients, service users and carers*

Children at risk of FH because they have 1 parent with the condition are offered diagnostic tests by the age of 10 years.

### *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendation 1.1.15

### *Definitions of terms used in this quality statement*

#### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

#### **Children at risk of FH**

Children (under 10 years) with 1 affected parent.

#### **Specified diagnostic tests**

NICE's guideline on familial hypercholesterolaemia recommends that children at risk of FH because of 1 affected parent receive either of the following diagnostic tests:

- A DNA test if the family mutation is known.
- LDL-C concentration measurement if the family mutation is not known. When excluding a diagnosis of FH a further LDL-C measurement should be repeated after puberty because LDL-C concentrations change during puberty.

### **Specialist with expertise in FH in children and young people**

A healthcare professional with expertise in FH in children and young people who has access to the wider skills of a multidisciplinary team. This team should include a dietitian, cardiologist and paediatrician, and a clinical genetic specialist to take a family history and obtain informed consent for a DNA test. All children and young people being investigated for a diagnosis of FH should be referred to a specialist with expertise in FH in children and young people in a child-focused setting.

### **Child-focused setting**

NICE's guideline on [familial hypercholesterolaemia](#) defines a child-focused setting as valuing the child's view and validating their voice in making decisions impacting their lives. A child-focused facility or space is one designed with the children's needs in mind.

## Quality statement 5: Cascade testing

### *Quality statement*

Relatives of people with a confirmed diagnosis of monogenic familial hypercholesterolaemia (FH) are offered DNA testing through a nationwide, systematic cascade process.

### *Rationale*

Most people in the UK with FH are undiagnosed. Cascade testing has been shown to be effective for identifying people with FH, especially when provided nationwide. Nationwide cascade testing ensures that all family members can access DNA testing wherever they live.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that relatives of people with a confirmed diagnosis of monogenic FH are offered DNA testing through a nationwide systematic cascade processes.

*Data source:* Local data collection.

#### **Process**

a) Proportion of untested first-degree relatives of people with a confirmed diagnosis of monogenic FH who are offered cascade testing.

Numerator – The number of people in the denominator offered cascade testing.

Denominator – The number of untested first-degree relatives of people with a confirmed diagnosis of monogenic FH.

*Data source:* Local data collection using a dedicated database.

b) Proportion of at-risk, untested, second- and third-degree relatives of people with a confirmed diagnosis of monogenic FH who are offered cascade testing.

Numerator – The number of people in the denominator offered cascade testing.

Denominator – The number of at-risk, untested second- and third-degree relatives of people with a confirmed diagnosis of monogenic FH.

*Data source:* Local data collection using a dedicated database.

c) Proportion of untested first-degree relatives of people with a confirmed diagnosis of monogenic FH who receive cascade testing.

Numerator – The number of people in the denominator receiving cascade testing.

Denominator – The number of untested first-degree relatives of people with a confirmed diagnosis of monogenic FH.

*Data source:* Local data collection using a dedicated database.

d) Proportion of at-risk, untested, second- and third-degree relatives of people with a confirmed diagnosis of monogenic FH who receive cascade testing.

Numerator – The number of people in the denominator receiving cascade testing.

Denominator – The number of at-risk, untested second- and third-degree relatives of people with a confirmed diagnosis of monogenic FH.

*Data source:* Local data collection using a dedicated database.

## Outcome

Prevalence of FH.

*Data source:* Local data collection using a dedicated database.

## *What the quality statement means for service providers, healthcare practitioners, and commissioners*

Service providers ensure that systems are in place for relatives of people with a confirmed diagnosis of monogenic FH to be offered DNA testing through a nationwide, systematic cascade process.

Specialists with expertise in FH offer DNA testing to relatives of people with a confirmed diagnosis

of monogenic FH through a nationwide, systematic cascade process.

Commissioners ensure that they commission services that offer DNA testing to relatives of people with a confirmed diagnosis of monogenic FH, through a nationwide, systematic cascade process.

### *What the quality statement means for patients, service users and carers*

Relatives of people with a confirmed diagnosis of FH and a known DNA mutation are offered DNA testing themselves as part of a national scheme.

### *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendation 1.2.1

### *Definitions of terms used in this quality statement*

#### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

#### **Monogenic FH**

Present when an autosomal dominant pattern of inheritance of elevated LDL-C levels is seen in the extended family of the proband (for example, on average 50% of first-degree relatives have elevated levels). In relatives, the age- and gender-specific diagnostic cut-offs in NICE's guideline on familial hypercholesterolaemia should be used because the Simon Broome diagnostic cut-offs are not appropriate for relatives. The diagnosis of monogenic FH can also be given when the index case carries a documented FH-causing mutation in the *LDLR*, *APOB* or *PCSK9* genes.

#### **Confirmed diagnosis of monogenic FH**

This requires evidence from DNA testing of an FH-causing mutation in the *LDLR*, *APOB* or *PCSK9* genes. Before cascade testing is initiated in relatives, a diagnosis of monogenic FH in the index person should be confirmed by a specialist with expertise in FH.

#### **Relatives**

At least first- and second-degree biological relatives and third-degree biological relatives if



possible, as defined in the glossary for NICE's guideline on [familial hypercholesterolaemia](#).

## Cascade testing

A mechanism for identifying people at risk of a genetic condition by a process of family tracing. For FH the test employed is a DNA test where a disease-causing mutation has been identified in the index individual/proband.

## Quality statement 6: Drug treatment in adults

### *Quality statement*

Adults with familial hypercholesterolaemia (FH) receive lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

### *Rationale*

Lipid-modifying drug treatment reduces LDL-C levels and prevents the development of cardiovascular disease. Studies indicate that treatment that lowers LDL-C levels by more than 50% from baseline offers greater benefit for plaque stabilisation than treatment that is less effective at reducing LDL-C.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that adults with FH receive lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

*Data source:* Local data collection.

#### **Process**

Proportion of adults with FH who receive appropriate lipid-modifying drug treatment.

Numerator – The number of people in the denominator receiving appropriate lipid-modifying drug treatment.

Denominator – The number of adults with FH.

*Data source:* Local data collection.

#### **Outcome**

Number of adults with FH whose LDL-C concentration is reduced by more than 50% from baseline within 1 year.

**Data source:** Local data collection using a dedicated database.

### *What the quality statement means for service providers, healthcare practitioners, and commissioners*

**Service providers** ensure that systems are in place for adults with FH to receive lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

**Healthcare practitioners** offer adults with FH lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

**Commissioners** ensure that they commission services that offer adults with FH lipid-modifying drug treatment to reduce LDL-C concentration by more than 50% from baseline.

### *What the quality statement means for patients, service users and carers*

**Adults with FH** are offered drugs to reduce the low-density cholesterol (bad cholesterol) in their blood to less than a half of the level before treatment.

### *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendations 1.3.1.2, 1.3.1.3, 1.3.1.4, 1.3.1.6, 1.3.1.7, 1.3.1.8 and 1.3.1.14

### *Definitions of terms used in this quality statement*

#### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

#### **Adults with FH**

Adults (aged 16 and older) should have a diagnosis of FH made by a specialist with expertise in FH.

#### **Baseline LDL-C**

The concentration before treatment.

## Lipid-modifying drug treatment

Lipid-modifying drug treatment should be given in accordance with the recommendations in NICE's guideline on [familial hypercholesterolaemia](#), a summary of which is given below:

- to achieve the recommended reduction:
  - offer a high-intensity statin with the lowest acquisition cost as the initial treatment
  - increase dose of statin to maximum licensed or tolerated dose
- ezetimibe monotherapy is recommended if the person is intolerant to statin therapy or there are contraindications to initial statin therapy
- co-administer ezetimibe with initial statin therapy when serum total or LDL-C concentration is not appropriately controlled and a change from initial statin therapy to an alternative statin is being considered.

Statin are classified as high intensity if they produce greater LDL-C reductions than simvastatin 40 mg (for example, simvastatin 80 mg and appropriate doses of atorvastatin and rosuvastatin).

Treatment for FH is usually provided by either a specialist with expertise in FH or a GP through a shared care arrangement.

### *Equality and diversity considerations*

The statement has been restricted to adults only because there is currently no evidence on which to base any specific target for lowering LDL-C in children and young people under 16 years. However, lipid-modifying drug treatment should be considered by the age of 10 years in line with NICE's guideline on [familial hypercholesterolaemia](#).

Women with FH should be advised that lipid-modifying drug treatment should not be taken if they are planning to conceive or during pregnancy because of the risk of fetal abnormality. Women should be advised that lipid-modifying drug treatment should be stopped 3 months before they attempt to conceive. Women with FH should be advised about the potential risks and benefits of re-starting lipid-modifying drug treatment for the mother and breastfed infant. Resins are the only lipid-modifying drug treatment that should be considered during breastfeeding.

## Quality statement 7: Drug treatment in children

### *Quality statement*

Children with familial hypercholesterolaemia (FH) are assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting by the age of 10 years.

### *Rationale*

Children with FH begin to develop cardiovascular disease before clinical signs appear, with thickening of the carotid artery wall identifiable by the age of 10 years. Once a child is diagnosed as having FH, it is important they should be assessed for lipid-modifying drug treatment as soon as possible. The assessment should include a discussion of the harms and benefits of different treatments. This allows children to start treatment as soon as it is appropriate and before significant atherosclerosis has developed.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that children with FH are assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting by the age of 10 years.

**Data source:** Local data collection.

#### **Process**

Proportion of children with FH who are assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting by the age of 10 years.

**Numerator** – The number of people in the denominator assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting.

**Denominator** – The number of children with FH aged 10 years.

**Data source:** Local data collection.

### *What the quality statement means for service providers, healthcare*

## *practitioners, and commissioners*

**Service providers** ensure that systems are in place for children with FH to be assessed for lipid-modifying drug treatment by a specialist with expertise in FH in a child-focused setting by the age of 10 years.

**Specialists with expertise in FH in children and young people** assess children with FH for lipid-modifying drug treatment in a child-focused setting by the age of 10 years.

**Commissioners** ensure that they commission services in which specialists with expertise in FH assess children with FH for lipid-modifying drug treatment, in a child-focused setting, by the age of 10 years.

## *What the quality statement means for patients, service users and carers*

**Children with FH** have an assessment for possible drug treatment to reduce the low-density cholesterol (bad cholesterol) in their blood by a specialist in a children's department, by the age of 10 years.

## *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendation 1.3.1.20

## *Definitions of terms used in this quality statement*

### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

### **Specialist with expertise in FH in children and young people**

A healthcare professional with expertise in FH in children and young people who has access to the wider skills of a multidisciplinary team. This team should include a dietitian, cardiologist and paediatrician, and a clinical genetic specialist to take a family history and obtain informed consent for a DNA test.

## Assessment for lipid-modifying drug treatment

NICE's guideline on [familial hypercholesterolaemia](#) recommends that the decision to offer or defer lipid-modifying drug treatment for a child or young person should take into account their age, age at onset of coronary heart disease within the family and the presence of other cardiovascular risk factors, including their LDL-C concentration.

## Child-focused setting

NICE's guideline on [familial hypercholesterolaemia](#) defines a child-focused setting as valuing the child's view and validating their voice in making decisions impacting their lives. A child-focused facility or space is one designed with the children's needs in mind.

## *Equality and diversity considerations*

Boys and girls should have equal access to lipid-modifying drug treatment for FH. There is anecdotal evidence that some clinicians are less likely to recommend statins in a girl than a boy at the same level of individual risk, arguing that because the age at onset of coronary heart disease in women is later than in men, the start of drug treatment can be 'safely' delayed until adulthood. However, many young women will need to stop statins for several years when they are trying to conceive, and during pregnancy and breastfeeding. Because the accumulation of LDL-C burden is similar in boys and girls, this will result in the exposure to high LDL-C being greater in early adulthood in young women than in their male siblings. Gender should not influence a clinician's decision to offer treatment; the decision should be made in accordance with the recommendations in NICE's guideline on [familial hypercholesterolaemia](#), which indicate that lipid lowering with statins should be considered by the age of 10 years.

## Quality statement 8: Annual review

### *Quality statement*

People with familial hypercholesterolaemia (FH) are offered a structured review at least annually.

### *Rationale*

Regular structured review enables treatment to be monitored and adjusted to achieve the recommended LDL-C concentration. It also enables monitoring for the possible development of symptoms and signs of coronary heart disease and optimising management. Records can be maintained of affected family members and information can be tailored to individual circumstances. Progress with cascade testing of at-risk relatives can be monitored.

### *Quality measures*

#### **Structure**

Evidence of local arrangements to ensure that people with FH are offered a structured review at least annually.

*Data source:* Local data collection.

#### **Process**

Proportion of people with FH who receive a structured review at least annually.

Numerator – The number of people in the denominator who had a structured review within 12 months of the last review or diagnosis.

Denominator – The number of people with FH.

*Data source:* Local data collection using a dedicated database.

### *What the quality statement means for service providers, healthcare practitioners, and commissioners*

Service providers ensure that systems are in place for people with FH to be offered a structured



review at least annually.

**Healthcare practitioners** offer people with FH a structured review at least annually.

**Commissioners** ensure that they commission services that offer a structured review at least annually to people with FH.

### *What the quality statement means for patients, service users and carers*

People with FH are offered a detailed review of their condition at least once a year.

### *Source guidance*

Familial hypercholesterolaemia: identification and management (NICE guideline CG71), recommendations 1.3.1.27, 1.4.2.1, 1.4.3.2, 1.5.1.1, 1.5.1.3, 1.5.1.4 and 1.5.1.5

### *Definitions of terms used in this quality statement*

#### **Familial hypercholesterolaemia (FH)**

FH relates to heterozygous FH only.

#### **People with FH**

People should have a diagnosis of FH made by a specialist with expertise in FH.

#### **Structured review**

Structured review should include all the following components if appropriate:

- Recording progress of cascade testing among relatives.
- Updating the family pedigree and noting changes in the coronary heart disease status of relatives.
- Assessing any symptoms of coronary heart disease.
- Assessing smoking status, and offering advice and information on smoking cessation services.
- Measuring fasting lipid profile.

- Discussing adherence to treatment, and possible side effects of treatment the person may be experiencing.
- Discussing any changes in lifestyle or lipid-modifying drug treatment that may be needed to achieve the recommended LDL-C concentration.
- Giving advice on contraception and pregnancy (to women and girls only).
- Monitoring growth and pubertal development (in children and young people only).

## Using the quality standard

### *Quality measures*

The quality measures accompanying the quality statements aim to improve the structure, process and outcomes of care in areas identified as needing quality improvement. They are not a new set of targets or mandatory indicators for performance management.

For further information, including guidance on using quality measures, please see [how to use quality standards](#).

### *Levels of achievement*

Expected levels of achievement for quality measures are not specified. Quality standards are intended to drive up the quality of care, and so achievement levels of 100% should be aspired to (or 0% if the quality statement states that something should not be done). However, NICE recognises that this may not always be appropriate in practice, taking account of safety, choice and professional judgement, and therefore desired levels of achievement should be defined locally.

### *Using other national guidance and policy documents*

Other national guidance and current policy documents have been referenced during the development of this quality standard. It is important that the quality standard is considered by commissioners, providers, health and social care practitioners, patients, service users and carers alongside the documents listed in [development sources](#).

## Diversity, equality and language

During the development of this quality standard, equality issues have been considered and [equality assessments](#) are available.

Good communication between healthcare practitioners and people with familial hypercholesterolaemia (FH) is essential. Treatment, care and support, and the information given about it, should be culturally appropriate. It should also be accessible to people with additional needs such as physical, sensory or learning disabilities, and to people who do not speak or read English. People with FH should have access to an interpreter or advocate if needed.

Commissioners and providers should aim to achieve the quality standard in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this quality standard should be interpreted in a way that would be inconsistent with compliance with those duties.

## Development sources

Further explanation of the methodology used can be found in the [quality standards process guide](#) on the NICE website.

## Evidence sources

The documents below contain recommendations from NICE guidance or other NICE-accredited recommendations that were used by the Topic Expert Group to develop the quality standard statements and measures.

- [Familial hypercholesterolaemia: identification and management](#) (2008, updated 2017) NICE guideline CG71

## Policy context

It is important that the quality standard is considered alongside current policy documents, including:

- Department of Health (2013) [Cardiovascular Disease Outcomes Strategy: Improving outcomes for people with or at risk of cardiovascular disease](#)
- Royal College of Physicians (2010) [National audit of the management of familial hypercholesterolaemia 2010](#)
- Royal College of Physicians (2010) [National clinical audit of the management of familial hypercholesterolaemia pilot 2009](#)
- NHS Wales (2009) [Cardiac disease national service framework for Wales](#)
- Department of Health (2004) [National service framework for children, young people and maternity services: core standards](#)

## Related NICE quality standards

- [Transition from children's to adults' services](#) (2016) NICE quality standard 140
- [Medicines optimisation](#) (2016) NICE quality standard 120
- [Obesity in adults: prevention and lifestyle weight management programmes](#) (2016) NICE quality standard 111
- [Cardiovascular risk assessment and lipid modification](#) (2015) NICE quality standard 100
- [Secondary prevention after a myocardial infarction](#) (2015) NICE quality standard 99
- [Obesity in children and young people: prevention and lifestyle weight management programmes](#) (2015) NICE quality standard 94
- [Physical activity: for NHS staff, patients and carers](#) (2015) NICE quality standard 84
- [Smoking: supporting people to stop](#) (2013) NICE quality standard 43
- [Patient experience in adult NHS services](#) (2012) NICE quality standard 15

## Topic Expert Group and NICE project team

### *Topic Expert Group*

**Dr Rubin Minhas (Chair)**

Group Medical and Scientific Director, Nuffield Health

**Dr Daphne Austin**

Public Health Physician, Right Care, QIPP, Department of Health

**Mrs Yvonne Dumsday**

Patient/carer member, Heart UK

**Mr Steve Humphries**

Head of Division, Centre for Cardiovascular Genetics, University College London

**Mrs Katharine Kear**

Patient/carer member, Heart UK

**Dr Ian McDowell**

Senior Lecturer and Consultant, University Hospital of Wales

**Mrs Clare Neuwirth**

Lipid Specialist Nurse/Research Nurse, Imperial College Healthcare NHS Trust/MRC Clinical Sciences Centre

**Dr Nadeem Qureshi**

Professor of Primary Care, University of Nottingham

**Dr Uma Ramaswami**

Consultant Metabolic Paediatrician, Willink Biochemical Genetics Unit, Central Manchester University Hospitals NHS Foundation Trust

**Mrs Helen Stracey**

Dietetic Services Manager, Chelsea and Westminster Hospital Foundation Trust

**Dr Adie Viljoen**

Consultant Chemical Pathologist, East and North Herts NHS Trust

**Ms Melanie Watson**

Principal Genetic Counsellor, Wessex Clinical Genetic Service, University Hospital Southampton

**Dr Catherine Woodward**

Director of Public Health and Director of Infection Prevention and Control, NHS Telford and Wrekin

*NICE project team*

**Nick Baillie**

Associate Director

**Rachel Neary**

Programme Manager

**Beth Shaw**

Technical Advisor

**Nicola Greenway**

Lead Technical Analyst

**Esther Clifford**

Project Manager

**Lee Berry**

Coordinator



## Update information

**November 2017:** Changes were made to definitions and source guidance sections throughout to ensure alignment with the updated NICE guideline on [familial hypercholesterolaemia: identification and management](#).

## About this quality standard

NICE quality standards describe high-priority areas for quality improvement in a defined care or service area. Each standard consists of a prioritised set of specific, concise and measurable statements. NICE quality standards draw on existing NICE or NICE-accredited guidance that provides an underpinning, comprehensive set of recommendations, and are designed to support the measurement of improvement.

Information about [how NICE quality standards are developed](#) is available from the NICE website.

This quality standard has been incorporated into the NICE Pathway on [familial hypercholesterolaemia](#).

ISBN: 978-1-4731-0260-6

## *Endorsing organisation*

This quality standard has been endorsed by NHS England, as required by the Health and Social Care Act (2012)

## *Supporting organisations*

Many organisations share NICE's commitment to quality improvement using evidence-based guidance. The following supporting organisations have recognised the benefit of the quality standard in improving care for patients, carers, service users and members of the public. They have agreed to work with NICE to ensure that those commissioning or providing services are made aware of and encouraged to use the quality standard.

- [HEART UK](#)
- [Royal College of General Practitioners](#)
- [Royal College of Nursing](#)