



# Pregnancy and neonates: smokers at delivery

NICE indicator

Published: 1 August 2012

Last updated: 30 November 2021

[www.nice.org.uk/indicators/ind19](https://www.nice.org.uk/indicators/ind19)

## Indicator

Proportion of pregnant women who were smokers at the time of delivery.

## Indicator type

Network / system level indicator.

The indicator would be appropriate to understand and report on the performance of networks or systems of providers.

This document does not represent formal NICE guidance. For a full list of NICE indicators, see our [menu of indicators](#).

To find out how to use indicators and how we develop them, see our [NICE indicator process guide](#).

## Rationale

Smoking during pregnancy is detrimental to the health of both the mother and unborn baby. Pregnant women who smoke should be referred to an evidence-based stop smoking service at the booking appointment to help woman to stop smoking during pregnancy and beyond.

## Source guidance

Antenatal care. NICE guideline NG201 (2021), recommendations 1.2.4, 1.3.9

Tobacco: preventing uptake, promoting quitting and treating dependence. NICE guideline NG209 (2021, updated 2023), recommendations 1.11.1, 1.18.2, 1.19.1, 1.19.2, 1.19.3, 1.19.4, 1.19.5, 1.19.6, 1.19.7

## Specification

**Numerator:** The number of pregnant women who were known to be smokers at the time of delivery.

**Denominator:** The number of pregnant women.

**Calculation:** Numerator divided by the denominator, multiplied by 100.

**Exclusions:** Mothers with an unknown smoking status.

**Data source:** NHS Digital statistics on women's smoking status at time of delivery: England.

Minimum population: The indicator would be appropriate to understand and report on the performance of networks or systems of providers.

ISBN: 978-1-4731-5881-8