



Diabetes: annual HbA1c (children T2DM)

NICE indicator

Published: 16 April 2025

www.nice.org.uk/indicators/ind299

This indicator replaces IND49.

Indicator

Proportion of children and young people aged under 18 years with type 2 diabetes who have had their glycated haemoglobin A1c (HbA1c) monitored in the previous 12 months.

Indicator type

Network / system level indicator. The indicator would be appropriate to understand and report on the performance of networks or systems of providers.

This document does not represent formal NICE guidance. For a full list of NICE indicators, see our [menu of indicators](#).

To find out how to use indicators and how we develop them, see our [NICE indicator process guide](#).

Rationale

This indicator aims to reduce the risk of complications associated with diabetes, and support early identification of any complications, in children and young people by monitoring glycated haemoglobin A1C (HbA1c). Good glycaemic control, measured using HbA1c, is important for the proactive management of diabetes to prevent complications, such as diabetic ketoacidosis.

Source guidance

[Diabetes \(type 1 and type 2\) in children and young people: diagnosis and management. NICE guideline NG18](#) (2015, updated 2023), recommendation 1.3.36

Specification

Numerator: The number of people in the denominator who had their glycated haemoglobin A1c (HbA1c) monitored in the previous 12 months.

Denominator: The number of children and young people aged under 18 years with type 2 diabetes.

Calculation: Numerator divided by the denominator, multiplied by 100.

Exclusions: None.

Personalised care adjustments or exception reporting should be considered to account for situations where the patient declines, does not attend or if HbA1c monitoring is not appropriate.

Data source: National Paediatrics Diabetes Audit and National Diabetes Audit Young People with Type 2 Diabetes Dashboard.

Expected population size: The National Diabetes Audit Young People with Type 2 Diabetes Dashboard 2022 to 2023 shows that 0.004% of people in England are aged under 18 years with type 2 diabetes: 0.4 per 10,000 patients served by a network. There is no minimum number of patients required for network level indicators. However, consideration should be given to whether the majority of results would require suppression because of small numbers.

ISBN: 978-1-4731-6950-0